Title: Environmental legacy effects impact maize growth and microbiome assembly under drought stress

Authors

Joel F. Swift[1,2*], Matthew R. Kolp[1,2,4], Amanda Carmichael[1,4], Natalie E. Ford[1], Paige M. Hansen[2,4], Benjamin A. Sikes[1,2], Manuel Kleiner[3], Maggie R. Wagner[1,2*]

Affiliations

[1] Department of Ecology and Evolutionary Biology, University of Kansas, Lawrence, KS 66045, USA

[2] Kansas Biological Survey & Center for Ecological Research, University of Kansas, Lawrence, KS 66045, USA

[3] Department of Plant and Microbial Biology, North Carolina State University, Raleigh, NC, 27695, USA

[4] Current Address: Center for Animal and Human Health in Appalachia, College of Veterinary Medicine, Lincoln Memorial University, Harrogate, TN 37752, USA (M.R.K.); Department of Conservation and Research, Southeastern Center for Conservation, Atlanta Botanical Garden, Atlanta, GA, 30309, USA (A.C.); Department of Soil and Crop Science, Colorado State University, Fort Collins, CO 80521, USA (P.M.H.)

*Correspondence: joel.swift@ku.edu and maggie.r.wagner@ku.edu

Abstract

Background and Aims: As the climate changes, plants and their associated microbiomes face greater water limitation and increased frequency of drought. Historical environmental patterns can leave a legacy effect on soil and root-associated microbiomes, but the impact of this conditioning on future drought performance is poorly understood. Precipitation gradients provide a means to assess these legacy effects.

Methods: We collected soil microbiomes from four native prairies across a steep precipitation gradient in Kansas, USA. Seedlings of two *Zea mays* genotypes were inoculated with each soil microbiome in a factorial drought experiment. We investigated plant phenotypic and root

microbiome responses to drought and modeled relationships between plant growth metrics and

climatic conditions from the soil microbiome origin sites.

Results: Drought caused plants to accumulate shoot mass more slowly and achieve greater

root/shoot mass ratios. Drought restructured the bacterial root-associated microbiome via

depletion of Pseudomonadota and enrichment of Actinomycetota, whereas the fungal microbiome

was largely unaffected. An environmental legacy effect on prairie soil microbiomes influenced

plants' drought responses: counterintuitively, prairie soil inocula from historically wetter locations

increased shoot biomass under drought more than inocula from historically drier prairie soils.

Conclusion: We demonstrated links between soil microbiome legacy effects and plant

performance under drought, suggesting that future drying climates may condition soils to

negatively impact plant performance.

Keywords: Root microbiome, soil microbiome, maize, drought, precipitation gradient

Introduction

Increasing drought frequency due to global climate change will impact both soil and plant-

associated microbiomes (Chiang et al. 2021; Balting et al. 2021). Drought directly limits microbial

growth, mobility, and access to resources, but also alters plant host traits (e.g., root physiology,

morphology, and exudate composition) that comprise the habitat available to root-associated

microorganisms (Naylor and Coleman-Derr 2018). Historical environmental patterns and drought

can leave lasting impacts, or legacy effects, on soil and root-associated microbiomes. Naturally-

occurring precipitation gradients—which shape the biogeography of both plants (Field et al. 2009;

Korell et al. 2021) and soil microorganisms (Bachar et al. 2010; Hawkes et al. 2011)—provide

opportunities to study legacy effects of the environment on plant-microbe interactions. Historical

patterns of water availability across gradients constrain the functional potential of soil

microbiomes (Hawkes and Keitt 2015; Averill et al. 2016; Canarini et al. 2021); however, the

implications for host plants, including crops, are not well understood.

Because plants recruit root microbes primarily from the soil (Zarraonaindia et al. 2015),

we would expect drought-induced changes in the soil microbiome to be reflected in root-associated

microbiomes. However, because many root-associated microbes are also sensitive to plant root

traits, their responses to drought may also reflect drought-induced changes in the host phenotype.

Both soil- and root-associated microbiomes under drought exhibit consistent enrichment of Gram-

positive bacteria and depletion of Gram-negative bacteria (Acosta-Martínez et al. 2014;

Fuchslueger et al. 2016; Naylor et al. 2017). Within those categories, however, some bacterial

phyla respond to drought differently in soil versus roots (Naylor and Coleman-Derr 2018). For

example, enrichment of Bacillota (formerly Firmicutes) and Actinomycetota (formerly

Actinobacteria) during drought are less pronounced in soil than in root-associated microbiomes,

suggesting that these phyla respond to drought-induced changes in root physiology (Naylor et al.

2017; Fitzpatrick et al. 2018). Furthermore, the presence of a host plant can influence the drought

responses of the taxa that comprise each phylum (Santos-Medellín et al. 2017). Overall, drought

impacts both soil- and root-associated microbiomes, but in different ways, owing in part to the

influence of the plant phenotype.

Changes in soil- and root-associated microbiomes caused by historical precipitation may

affect plant fitness under drought. Drought-induced restructuring of root-associated microbiomes

is hypothesized to be a conserved mechanism that supports survival of both the plant host and its

symbionts (Naylor et al. 2017; Xu et al. 2018; Xu and Coleman-Derr 2019). For example, studies

have shown that plant fitness is increased when a soil microbiome's historic moisture conditions

match contemporary experimental conditions (Lau and Lennon 2012; Munoz-Ucros et al. 2022),

with this effect observed more commonly with co-adapted plants and soils that historically share

a habitat (Lau et al. 2017; Remke et al. 2021; de Vries et al. 2023). Additional research suggests

that exposure to short-term drought creates a microbial legacy effect that moderates drought stress

in plants (Kaisermann et al. 2017; De Long et al. 2019) and strengthens positive plant-soil

feedbacks, although many such patterns were species-specific (Buchenau et al. 2022). Thus, how

historical precipitation patterns condition soil- and root-associated microbiomes and their

relevance to plant fitness under drought stress is an open question.

90   To address this open question about legacy effects on microbiome function, we designed

an experiment to assess plant fitness under a contemporary drought when inoculated with soil

microbiomes from different historic precipitation regimes. Specifically, we collected soil

microbiomes from native prairie remnants spanning an east-west precipitation gradient in Kansas

(Figure 1). This precipitation gradient is coupled with multiple climatic metrics, with the

environment becoming increasingly arid from east to west resulting in lower soil moisture and

increased reference evapotranspiration (Figure 1C-E). We inoculated *Zea mays* L. (maize)

seedlings with these soil microbiomes and measured plant growth during our experimental

drought. Our goals were to: **1**) characterize plant phenotype and microbiome responses to drought,

2) test how soil microbiome histories affected maize growth, and **3**) identify microbial taxa that

were associated with plant growth. Our results suggest that both historical and experimental

drought shape soil microbiomes' interactions with maize.

Material and Methods

Soil inoculum sources and preparation

Soil samples were collected in May 2019 from four prairie sites across a steep precipitation gradient in Kansas, USA (Figure 1). Prairies were selected at the same latitude and relative elevation such that they have very similar temperature profiles throughout the year. Western prairie sites (SVR and HAY) are predominantly short- to mixed-grass prairies (e.g., *Bouteloua* spp.), while eastern sites (TLI and KNZ) are dominated by tallgrass prairie species (e.g., *Andropogon gerardi*, *Schizachyrium scoparium*, *Panicum virgatum*). All collection sites were high-quality remnant prairies that have never experienced tillage or irrigation, which are strong perturbations that may obscure environmental legacy effects. Soil was collected from five evenly-spaced sub-samples per site, sampling to an approximate depth of 60 cm. Soil sub-samples were homogenized and sieved to remove rocks and plant material and then stored in plastic bags at 4ºC until use, approximately four months after collection.

Figure 1. Kansas precipitation gradient. A) Soil samples were collected from four prairies in Kansas along an increasing precipitation gradient. Locations from east to west, Smoky Valley Ranch prairie (SVR; latitude 38.8665, longitude -100.9951), Hayes prairie (HAY; 38.8355, -99.3033), The Land Institute prairie (TLI; 38.9698, -97.4690), and Konza native prairie (KNZ;

121 39.1056, -96.6099). Boxplots to the right of the map depict annual climatic metrics extracted from
122 the TerraClimate database for each collection site from 1990-2021: **B)** precipitation, **C)** reference
123 evapotranspiration, **D)** aridity index, and **E)** soil moisture. Boxplot hinges represent the 1st and 3rd
124 quartiles; whiskers represent 1.5 times the interquartile range.
125

Inocula were prepared by mixing 20 g of each soil into 100 mL 1x phosphate-buffered

saline (PBS) and 0.0001% Triton X-100. After settling, the soil slurries were filtered through

Miracloth (22-25 µm pore size; Calbiochem, San Diego, CA, USA) and then centrifuged for 30

minutes (3000 x g). After discarding the supernatant, we resuspended the microbial pellets in 20

mL 1x PBS and kept them shaking for aeration until use, later that day. Each final inoculum

consisted of 10 mL of the resuspended pellet in 1 L 0.5X Murashige and Skoog basal salt mixture

(MS; Sigma-Aldrich, Darmstadt, Germany). A control treatment was made by adding 10 mL

sterile 1x PBS into 1 L 0.5X MS.

Planting, inoculation, and greenhouse setup

To increase the generalizability of our results, two genotypes of *Zea mays* L. (maize) were

used: B73 and Mo17. These genotypes have been shown to vary in their growth response to

beneficial endophytes (Schultz et al. 2022) and so may differ in response based on microbiome

origin. In a laminar flow hood, seeds were surface sterilized by submerging in 70% ethanol, then

5% NaClO, followed by three rinses with sterile water. To avoid confounding effects of soil

physicochemical properties, autoclaved calcined clay ("Pro's Choice Rapid Dry"; Oil-Dri

Corporation, Chicago, IL) was used as the growth medium. Seeds were planted one inch deep in

107mL "cone-tainers" (Stuewe & Sons Inc., Tangent, OR, USA) with a sterilized cotton ball

pushed to the bottom to prevent clay from passing through drainage holes. Approximately 100mL

of sterilized calcined clay was used per cone-tainer. The day after planting, cone-tainers were

assigned randomly to treatments, inoculated with 25 mL of one of the four soil inocula (or the

sterile control), and placed two inches apart in randomized complete blocks. Overall, each

combination of treatments (drought treatment × soil inocula × genotype) was replicated 29 to 31

times across 27 randomized blocks (Figure S1A). Plants were grown in a temperature-controlled

greenhouse (23/20 ℃ day/night, respectively) at the University of Kansas (Lawrence, KS, USA)

arranged under growth lights (250 W Sun Systems Compact Cool fluorescent lamps) to

supplement natural light for 13 hours/day during the experiment (January-March 2020).

Monitoring and harvesting plants

All plants received 30 mL of reverse osmosis water twice weekly until emergence. After

emergence, "well-watered" plants continued twice-weekly watering while "droughted" plants

were watered every 8-12 days. Plant height was measured weekly to the nearest 0.25 cm on the

tallest fully expanded leaf held parallel to the stem. Plants were sequentially harvested to measure

biomass at one of three time points during the experiment: 68 plants on day 24 and day 39, and the

remainder (n=224) on day 50 (Figure S1B). Calcined clay was removed from roots by hand. Shoot

and root fractions were separated and oven-dried at 65 ℃ to a constant weight. Before drying, one

nodal root (from the uppermost soil-borne root whorl) per plant was collected and stored at -80 ℃

until DNA extraction.

Phenotypic analysis

All statistical analysis for phenotypic and microbiome data (see below) was conducted in

R v3.6 (R Core Team 2021). Differences among group means were assessed using analysis of

variance (ANOVA) with Type III sums of squares in the car v3.0.13 (Fox and Weisberg 2019) or

lmertest v3.1.3 packages (Kuznetsova et al. 2017) with post-hoc comparisons made using the

emmeans package v1.7.4.1 (Lenth et al. 2020). Assumptions of all models were checked by

inspecting model residuals (Harrison et al. 2018). Dry shoot and root mass were converted to

accumulation rates (g/day) to account for the variable number of growing days plants experienced

post-emergence. The lme4 package v1.1.29 (Bates et al. 2015) was used to fit a generalized linear

mixed model for emergence (binary) and linear mixed models for other growth responses

(continuous). Models used the following formula (*growth rate ~ Inoculum × Treatment ×*

Genotype) with all possible interactions included. Experimental block was treated as a random

effect with time point included as an additional random effect in the root/shoot ratio model.

Root/shoot ratio was natural log-transformed, and both shoot and root mass accumulation rates

were square-root-transformed to meet the assumptions of ANOVA.

Models were also fit using climate metrics to test for linear relationships between plant

growth and climatic conditions from the soil microbiome origin sites. Monthly precipitation,

reference evapotranspiration (grass reference crop), Palmer's Drought Severity Index (PDSI), and

soil moisture values were extracted from TerraClimate database (Abatzoglou et al. 2018), selecting

the closest 4x4 km grid cell to each site. Annual normals (30-year averages; Figures 1B-D and S2)

were calculated for each metric, PDSI was calculated as a monthly average to reduce

autocorrelation, and an aridity index was calculated by dividing normal annual precipitation by

evapotranspiration.

DNA extraction and library preparation

Nodal roots were freeze-dried for 48 hours (FreeZone lyophilizer; Labconco, Kansas City,

MO, USA) and ground to powder with a HT Lysing Homogenizer (OHAUS, Parsippany, NJ,

USA). Each ground root sample was mixed into 800 µL lysis buffer (1M Tris pH=8.0; 0.5 mM

NaCl; 10 mM EDTA) and transferred to 96-well plates containing 500 µL of 2 mm garnet rocks

per well (Biospec Products, Bartlesville, OK, USA), after which DNA purification proceeded via

bead-beating and chemical lysis as previously described (Wagner et al. 2020).

Amplicon sequencing libraries were prepared using established protocols and PCR

conditions (Wagner et al. 2020). Briefly, we targeted the V4 region of the 16S SSU rRNA gene

for bacteria/archaea using 515F-806R primers (Apprill et al. 2015; Parada et al. 2016) and the

ITS1 region for fungi using ITS1f and ITS2 primers (White et al. 1990; Gardes and Bruns 1993).

Both amplicons were amplified separately using an initial PCR and then barcoded via a second

index PCR. Amplicons were sequenced (2x250 bp) on a NovaSeq 6000 SP flow cell (Illumina,

San Diego, CA, USA). Each "batch" of 96 samples (n=6) remained together throughout DNA

extraction, amplification, and sequencing; individual samples were randomized across batches.

To confirm that we inoculated the experimental plants with realistic microbial

communities, we also sequenced the original soils and freshly derived inocula. From homogenized

soil collected from each site, we created eight technical replicates which we used to test whether

our protocol for creating microbial slurries preserved the original community composition. DNA

from each soil replicate and from freshly generated inocula was extracted in December 2022 using

the same protocol that we used for roots, except omitting the lyophilization step and placing 250

mg soil or 800 µL inoculum directly into the garnet plate. From there, library preparation and

sequencing proceeded as described above for roots.

Bioinformatic processing

Sequence processing and quality filtering were conducted using an established workflow

(Wagner et al. 2020). Briefly, Cutadapt v2.3 (Martin 2011) was used to remove primers and

demultiplex reads into 16S and ITS sequence files. DADA2 v1.14.1 (Callahan et al. 2016) was

used to denoise reads, merge paired-end sequences, dereplicate, and remove chimeric sequences.

The forward and reverse 16S reads were truncated to 220 and 170bp, respectively and filtered

based on the maximum number of expected errors of eight and nine bp per read, respectively. ITS

reads were processed using DADA2 in an identical manner, except that reads were not truncated

to preserve biologically relevant length variation. Forward and reverse 16S and ITS reads were

then used separately to estimate sequencing error rates for denoising prior to dereplication of reads

into Amplicon Sequence Variants (ASVs) and merging of pair-end reads. Taxonomy was assigned

to the resulting ASVs by a naïve Bayesian classifier (Wang et al. 2007) trained on the 11.5 release

of the Ribosomal Database Project (Callahan 2017) for bacteria and the 01.12.2017 release of the

UNITE database (UNITE Community 2017) for fungi. Phyloseq v1.40.0 (McMurdie and Holmes

2013) was used to remove host sequences (chloroplast and mitochondria; 0.35% of 16S reads) and

ASVs unassigned to a kingdom. Chao1 richness and diversity indices (Shannon and Inverse

Simpson) were calculated before the removal of ASVs not observed ≥ 25 times in ≥ 5 samples and

samples with <400 reads (200 bacterial and 133 fungal samples); 98.0% and 94.5% of reads

remained after host filtering and thresholding (bacterial and fungal reads, respectively). Finally,

ASV counts were normalized via a centered-log-ratio (CLR) transformation using the ALDEx2

package v1.28.1 (Fernandes et al. 2014; Gloor et al. 2017). Due to low replication and low

sequencing depth at the earlier timepoints, we analyzed only microbiome samples from the last

time point. This retained 131 bacterial (mean read depth 123,554 ± SE 18,276) and 92 fungal

samples (41,081 ± 10,129).

Microbiome Analysis

To test the similarity of inocula to their soils of origin, a linear model was fit for each alpha

diversity metric with collection site, substrate (soil *vs.* inoculum), and sequencing depth (Z-

transformed) as main effects. Community composition was assessed by permutational multivariate

analysis of variance (PERMANOVA) and constrained partial redundancy analysis (RDA). For

PERMANOVA, the *adonis2()* function from the vegan package v2.6.2 (Oksanen et al. 2019) was

used to model the matrix of between-sample Aitchison distances (Gloor et al. 2017) with collection

site and substrate as main effects. The RDA was fit with the same model, removing variance

attributed to ln(Sequencing Depth), and visualized using an ordination plot.

To investigate root microbiome responses, a model was fit for each alpha diversity metric

(*Metric ~ Soil Inoculum × Drought Treatment × Genotype*), with *Block* and *Batch* as random

effects. The CLR-transformed abundance of each family was modeled to test responses to

experimental factors. Across all models and where appropriate, p-values were adjusted to correct

for multiple comparisons using a Benjamini-Hochberg correction (Benjamini and Hochberg

1995). To explore changes in community composition, a PERMANOVA model was fit (*Aitchison*

distance matrix ~ Soil Inoculum × Drought Treatment × Genotype) with permutations constrained

by *Batch.* Main effects were tested for homogeneity of dispersions using the vegan package.

Pairwise dispersions among factor levels were assessed using a permutation test with 999

permutations constrained by *Batch.* Constrained partial redundancy analysis of the Aitchison

distance matrix was conducted to remove variance attributed to *Batch* and *ln(Sequencing Depth)*

and to visualize the variance in microbial composition explained by experimental factors.

The machine-learning package caret v6.0.92 (Kuhn 2008) with ranger's v0.14.1 (Wright

and Ziegler 2017) implementation of the random forest algorithm was used to assess how

accurately roots could be classified as droughted or well-watered based on microbiome community

data (bacterial and fungal, separately). Samples were randomly split into a training set (80%) and

a test set (20%), with the test set withheld from the model, and used to determine the final model's

performance. Hyperparameters were optimized using a grid search over minimum node size (1, 5,

10) and the number of features available at each node (10-100% of the ASVs). For each

combination in the grid, classifier accuracy was used to assess performance on out-of-bag samples

with 10-fold cross validation. Final predictions were made with the trained model on the withheld

test set and visualized using confusion matrices. The contribution of each ASV to classification

accuracy was assessed using permuted importance: mean decrease in accuracy (MDA) values were

calculated for each ASV across the 10-fold cross validation. The benefits of this strategy are

twofold, in that we can assess whether drought induced a predictable shift in the microbiome as a

whole and identify individual ASVs with the greatest differential abundance as a result of drought.

Connecting plant growth to microbial abundances

Linear models were used to test associations between taxon abundances and plant growth.

To focus on the phenotypic variation that could not be explained by Genotype, Soil Inoculum,

Block and Batch effects, we extracted residuals from a model that contained these terms and used

them as the response variable for a second model (*Growth Response$_{residual}$ ~ Taxon abundance $\times$*

Drought Treatment). The *Taxon abundance* coefficient described the association of the taxon

(CLR-transformed counts of bacterial and fungal families) with plant growth rate; the interaction

term described whether the taxon's effect on plant growth differed under droughted vs. well-

watered conditions. We chose family and order as the taxonomic levels for testing bacteria and

fungi, respectively, as a conservative measure to limit the number of tests. P-values were adjusted

by term to account for multiple testing.

Results

Soil and inocula community composition differs across collection sites

We assessed the microbial community composition of soils and their derived inocula from

the four collection sites (n=8 samples per site). Bacterial, but not fungal, alpha diversity of soils

differed across collection sites (Tables S1 and S2); bacterial diversity was highest in the

westernmost prairie soil (Figure S3). Community composition of both bacterial and fungal

microbiomes differed among sites (Table S3; R^2=0.22-0.24; *p*-value<0.01). Relative abundances

of bacterial phyla and fungal classes further showed site-specific patterns (Figures S5 and S6).

Relative to the original soils, we observed that inocula generally had lower richness and

diversity, but this difference was non-significant for most metrics (Figure S4; Tables S1 and S2).

The composition of both bacterial and fungal communities also shifted between soils and their

derived inocula (Figure S5-6; Table S3); however, both soil and inocula samples clustered in

ordination space by collection site (Figure 2). Thus, the derived inocula largely retained the

differences among the original soil microbial communities.

Figure 2. Soil community composition is strongly differentiated across the soil sources and

substrates. Redundancy analysis for **A**) bacterial and **B**) fungal communities from soil sources

used to derive soil inocula. Soil source and substrate are represented by differing colors or

shapes, respectively.

Plant growth rates vary across soil microbiomes, genotype, and drought treatment

We found no evidence that soil inocula impacted emergence rates (p-value = 0.19,

Pearson's χ^2 test; Table S4), however, maize genotype B73 (81.3%) emerged more successfully

than Mo17 (62.0%; p-value=0.05, Pearson's χ^2 test). Post-emergence, both drought treatment and

maize genotype affected plant growth (Table 1). Drought decreased both shoot and root mass

accumulation rates (Figure 3A-B; p-value=<0.01) and increased root/shoot ratio (Figure 3C; p-

value<0.01). These traits also differed between genotypes, with Mo17 investing more into root

growth than B73 (Figure S7; p-value=<0.01). Shoot mass accumulation rate and root/shoot ratio

were both influenced by soil inoculum (Table 1; p-value=<0.01-0.02); this effect was driven by

the 'TLI' inoculum which caused the slowest accumulation of shoot mass and the lowest root/shoot

ratio. We observed no evidence of interactions between drought treatment, genotype, and soil

inoculum on plant growth.

Table 1. Type three analysis of variance (ANOVA) for plant growth.

Factor	DF	sqrt(Shoot mass rate)		sqrt(Root Mass Rate)		log(Root/shoot Ratio)	
		F-value	p-value	F-value	p-value	F-value	p-value
Drought Treatment (DT)	1	335.67	**<0.01**	16.96	**<0.01**	161.21	**<0.01**
Genotype (G)	1	33.56	**<0.01**	22.74	**<0.01**	110.11	**<0.01**
Soil Inoculum (SI)	3	5.08	**<0.01**	0.67	0.57	3.30	**0.02**
DT x G	1	0.08	0.77	0.93	0.34	0.12	0.73
DT x SI	3	2.25	0.08	1.73	0.16	0.63	0.59
G x SI	3	2.13	0.10	0.61	0.61	0.44	0.73
DT x G x SI	3	0.25	0.86	0.09	0.96	0.13	0.94

311 Bold values denote significant effects, p-value<0.05

Figure 3. Shoot and root growth vary across drought treatment. The effect of drought treatment on **A)** shoot mass accumulation rate, **B)** root mass accumulation rate, and **C)** root/shoot ratio. Boxplot hinges represent the 1st and 3rd quartiles; whiskers represent 1.5 times the interquartile range; letters above box plots denote significant differences in estimated marginal means (*p*-value=<0.01; see table 1 for full ANOVA table).

Root microbiomes and community responses to drought differed between bacteria and fungi

We observed different patterns of richness and diversity for bacterial and fungal root microbiomes exposed to prairie inoculum and drought (Tables S5 and S6). Bacterial Chao1 richness varied across soil inocula (Table S5; *p*-value=0.03), with roots receiving the 'KNZ' inoculum exhibiting the highest richness. Drought increased Shannon and Inverse Simpson's diversity of fungal communities (Table S6). We observed no difference in bacterial or fungal richness or diversity between the maize genotypes (Tables S5 and S6).

Root bacterial community composition was also structured by soil inoculum (Table 2; R^2=0.05, *p*-value=<0.01, PERMANOVA) and drought treatment (Figure 4A; R^2=0.01, *p*-value=0.01, PERMANOVA). We found no evidence that bacterial community dispersion differed among treatments, genotypes, or inocula (*p*-value>0.05, Multivariate Levene's test). Redundancy analysis aligned with PERMANOVA results: soil inoculum and drought treatment explained the

largest proportions of variance (6.09% and 1.57%; Figure S8 and 4B, respectively), while genotype

explained <1% of variance (Figure S9). Among bacterial families, Sphingomonadaceae was

responsive to drought treatment (q=7.1e^{-4}), with this family decreasing from 1.11% relative

abundance to 0.46% under drought conditions.

Table 2. Permutational Multivariate ANOVA (PERMANOVA) table bacterial community
composition of maize nodal roots.

Factor	DF	SS	R^2	F-value	p-value
Genotype (G)	1	2687	0.01	0.88	0.47
Soil Inoculum (SI)	3	19091	0.05	2.10	**<0.01**
Drought Treatment (DT)	1	5484	0.01	1.81	**0.01**
G x SI	3	9849	0.02	1.08	0.47
G x DT	1	2289	0.01	0.75	0.72
SI x DT	3	10026	0.02	1.10	0.43
G x SI x DT	3	6352	0.02	0.70	0.95
Residual	115	349180	0.86		

339   Bold values denote significant effects, p-value<0.05
340

Figure 4. Bacterial community composition across drought treatments. A) Bacterial
taxonomic barplots for drought-stressed and well-watered plants. Each site is represented by a
facet with samples denoted by individual lines and the relative abundance (non CLR transformed
data) of each phylum displayed by color. **B)** Redundancy analysis on bacterial communities
constrained by drought treatment, conditioned on sequencing batch and ln-transformed usable
reads per sample. **C)** Confusion matrix depicting the classification accuracy of a random forest
classifier predicting drought treatment trained on bacterial communities. Above the confusion
matrix balanced accuracy is shown; for additional classifier statistics, see Table S8.
350
In contrast to bacteria, drought treatment did not alter relative abundances of any fungal

families, nor overall root fungal community profiles (Figure 5A). Variance in fungal community

composition was mostly due to initial soil inoculum source (Figure S10; Table 3; R^2=0.09, p-

value=<0.01). As observed for bacterial communities, neither drought treatment, maize genotype,

nor soil inoculum source affected fungal community dispersion (p-value=>0.05, Multivariate

Levene's test). Drought and maize genotype each explained only a small portion of the variance

in fungal community composition (Figure 5B and S11, respectively). Thus, while root-associated

bacterial community composition responded to drought treatment, we observed no such trend for

fungal community composition.

Table 3. PERMANOVA table for fungal community composition of maize nodal roots.

Factor	DF	SS	R^2	F-value	p-value
Genotype (G)	1	586	0.01	0.89	0.49
Soil Inoculum (SI)	3	5896	0.09	3.00	**<0.01**
Drought Treatment (DT)	1	541	0.01	0.83	0.63
G x SI	3	2181	0.03	1.11	0.60
G x DT	1	740	0.01	1.13	0.18
SI x DT	3	2335	0.04	1.19	0.17
G x SI x DT	3	1969	0.03	1.00	0.61
Residual	76	49822	0.78		

Bold values denote significant effects, p-value<0.05

Figure 5. Fungal community composition across drought treatment. A) Fungal taxonomic

barplots for drought-stressed and well-watered plants. Each site is represented by a facet with

samples denoted by individual lines and the relative abundance (non CLR transformed data) of

each phylum displayed by color. **B)** Redundancy analysis on fungal communities constrained by

drought treatment, conditioned on sequencing batch and ln-transformed usable reads per sample.

C) Confusion matrix depicting the classification accuracy of random forest a classifier predicting

drought treatment trained on fungal communities. Above the confusion matrix balanced accuracy

is shown; for additional classifier statistics, see Table S9.

Identification of bacterial and fungal ASVs responding to drought

Our random forest models showed greater accuracy in predicting the drought treatment of

a given sample when trained on bacterial communities than on fungal communities (Figures 4C

and 5C). Three bacterial ASVs greatly aided the prediction of treatment (Figures 4C and S12-13):

two from the family Sphingomonadaceae (Pseudomonadota) and one from the family

Sphingobacteriaceae (Bacteroidota). These ASVs showed mean decrease in accuracy (MDA)

values of 1.6%, 1.2%, and 1.2% when withheld from the model, respectively. In other words, these

three bacterial ASVs contributed to ~4% increase of classifier accuracy in predicting. All these

bacterial ASVs were depleted in drought-stressed plants relative to well-watered plants: the

Sphingomonadaceae ASVs decreased in relative abundance by an order of magnitude (0.37% and

0.31% to 0.03% and 0.04% relative abundance, respectively) and the Sphingobacteriaceae ASV

responded similarly (0.9% to 0.1% relative abundance). We also detected two fungal ASVs that

aided the prediction of treatment (Figure 5C and S14-15), from the families Aspergillaceae (1.1%

MDA; Eurotiomycetes) and Sarocladiaceae (0.76% MDA; Sordariomycetes). These fungal ASVs

showed different patterns in response to drought treatment: the Aspergillaceae ASV was enriched

while the Sarocladiaceae ASV was depleted in drought-stressed plants.

Connecting plant growth to bacterial and fungal communities

Soil inocula derived from collection sites across Kansas had only a small impact on plant

growth (Table 1; Figure 6A and S15), with no single inoculum causing significantly improved

growth over other inocula or the control. Examining the prairie sites spanning the east-west

precipitation gradient, we detected an interaction between mean annual precipitation and drought

treatment on shoot mass accumulation rate (Figure 6B; *p*-value=0.04). The shoot mass

accumulation rate of well-watered plants was negatively associated with mean annual precipitation

of the soil microbiome collection site (slope = -2.88e^{-6} g/day/mm precipitation) whereas droughted

plants displayed a weak positive association (slope = 4.53e^{-7} g/day/mm precipitation). Put another

way, counterintuitive to expectations, plants that experienced drought benefited from microbiomes

from historically wetter soils, while the opposite was true for well-watered plants in the experiment

(we discuss this striking pattern below). We observed that this pattern was consistent when

substituting other climatic metrics (PDSI, aridity index, evapotranspiration, and soil moisture) for

mean annual precipitation (Figure S17; *p*-values = 0.03, 0.06, 0.08, and 0.18, respectively).

Notably, B73 and Mo17 exhibited contrasting associations with mean annual precipitation across

the Kansas precipitation gradient (Figure 6C; *p*-value=0.02; Slope= -2.94e^{-6} and 5.10e^{-7},

respectively); the same pattern was observed for other climatic metrics except for PDSI (Figure

S17).

407
Figure 6. Shoot growth rate by soil inoculum. Boxplots are colored by soil inoculum for **A)**
shoot mass accumulation rate with separate facets representing drought-stressed and well-
watered plants. Boxplot hinges represent the 1st and 3rd quartiles; whiskers represent 1.5 times
the interquartile range. **B-C)** shoot mass accumulation rate trend lines depicting significant
interactions of mean annual precipitation (last climate normal 1991-2020) with drought treatment
and genotype, respectively, for plants inoculated with prairie soils. For each regression line, grey
shading represents the 95% confidence interval.
415

We identified bacterial and fungal taxa that were correlated with promotion or suppression

of plant growth. Overall, 11 bacterial families showed a significant positive (3/11) or negative

(8/11) association with shoot mass accumulation rate (Table 4). These represented only a few phyla

(5 Actinomycetota, 4 Pseudomonadota, and 2 Bacillota) and were a mix of Gram-positive (7) and

Gram-negative (5). One of the identified families, Streptococcaceae, interacted with drought

treatment, indicating that its relationship with shoot mass accumulation was modulated by the

amount of water the plant received. The abundance of Streptococcaceae was negatively correlated

with shoot mass rate in well-watered plants but not in droughted plants (Figure S18). Relative to

shoot mass accumulation rate, fewer bacterial families were associated with root mass

accumulation rate (only Acetobacteraceae and Corynebacteriaceae, Table 4). This result may be

due to variance in this trait decreasing over the course of the experiment as roots expanded to fill

the entire container regardless of treatment or inoculum (Figure S19). Notably, Acetobacteraceae

abundance was negatively associated with shoot mass accumulation rate but positively associated

with root mass accumulation rate. We found that the families associated with root/shoot ratio,

Sinobacteraceae and Sphingomonadaceae (Table 4), were a subset of families associated with

shoot mass accumulation rate, suggesting that a change in shoot mass was responsible for this

relationship. For fungi, only a single family, Sporidiobolaceae, was correlated with shoot mass

accumulation rate (Table 4). Similar results were found at lower taxonomic levels for both bacteria

and fungi (Table S10-11).

Table 4. Bacterial and fungal families associated with plant traits; square root transformed shoot
mass accumulation rate (SMR), root mass accumulation rate (RMR), and root/shoot ratio (RSR).

Taxon	Trait	Model Term	R	X	PVE$_{Res}$	Rel. Abund.	q
Acetobacteraceae	SMR	Abundance	-1.25E-04	-1.33E-02	6.67	0.02	2.76E-03
Corynebacteriaceae	SMR	Abundance	-5.46E-04	-1.29E-02	8.60	0.01	1.90E-04
Dietziaceae	SMR	Abundance	-1.07E-03	-1.31E-02	4.94	0.00	6.48E-03
Lactobacillaceae	SMR	Abundance	5.64E-04	-1.46E-02	4.26	0.09	1.55E-02
Microbacteriaceae	SMR	Abundance	-1.07E-05	-1.32E-02	5.27	1.23	6.48E-03
Micrococcaceae	SMR	Abundance	-8.43E-06	-1.31E-02	3.84	8.47	2.46E-02
Nocardiaceae	SMR	Abundance	-3.71E-04	-1.33E-02	3.58	0.02	2.75E-02
Rhodocyclaceae	SMR	Abundance	-4.92E-04	-1.14E-02	3.51	0.06	2.75E-02
Sinobacteraceae	SMR	Abundance	2.12E-04	-1.38E-02	3.70	0.05	2.07E-02
Sphingomonadaceae	SMR	Abundance	-1.03E-04	-1.24E-02	5.31	1.13	6.48E-03
Streptococcaceae*	SMR	Abundance	1.38E-04	-1.34E-02	5.50	0.01	4.10E-03
Acetobacteraceae	RMR	Abundance	2.32E-05	-5.94E-03	8.15	0.02	1.41E-02
Corynebacteriaceae	RMR	Abundance	-2.65E-04	-5.74E-03	6.29	0.01	3.39E-02
Polyangiaceae	RSR	Abundance	2.19E-02	1.37E-01	8.01	0.08	1.81E-02

Sinobacteraceae	RSR	Abundance	-1.48E-03	1.67E-01	5.8	0.05	2.96E-02
Sphingomonadaceae	RSR	Abundance	6.96E-04	1.58E-01	7.44	1.13	1.81E-02
Sporidiobolaceae	SMR	Abundance	3.33E-04	-1.24E-02	6.17	0.48	2.95E-02

*Streptococcaceae shows a significant interaction between the family abundance and drought treatment,
presented in Figure S11.

Discussion

Plant and microbiome responses to drought

Understanding plant-microbiome interactions under drought stress is critical to safeguard

agricultural systems in the face of global climate change. In our experiment, maize displayed

phenotypic tradeoffs in response to drought. Drought-treated plants increased root growth relative

to shoot growth, resulting in shorter plants over the course of the experiment. This supports our

understanding of plant responses to drought, as plants prioritize root over shoot growth (Sharp and

Davies 1979; Lynch et al. 2014), favoring deeper and often denser root systems in order to obtain

sufficient water (Zhan et al. 2015; Gao and Lynch 2016; Sebastian et al. 2016). The maize

genotypes, B73 and Mo17, showed significant differences between root/shoot allocation patterns,

but in general both genotypes responded to drought in a similar manner. These phenotypic changes

in root growth, along with direct effects of water limitation in the soil, contributed to the responses

of the root microbiomes in our study.

Drought altered the bacterial communities of maize roots, although the magnitude of

compositional change within our study was small (R^2=0.01) relative to other studies (R^2=0.04-

0.10) (Santos-Medellín et al. 2017; Naylor et al. 2017; Fitzpatrick et al. 2018; Xu et al. 2018).

Similar to previous studies (Xu and Coleman-Derr 2019), we observed a depletion of Gram-

negative and an enrichment of Gram-positive bacteria in response to drought, likely relating to the

desiccation resistance provided by the thick peptidoglycan cell wall of Gram-positive bacteria

(Manzoni et al. 2012). Enrichment of Actinomycetota during drought tends to be more pronounced

in root-associated microbiomes than in soil, suggesting that shifts in plant root physiology also

play a role (Naylor and Coleman-Derr 2018; Xu and Coleman-Derr 2019); however, the specific

root traits driving this effect are unknown. During drought, root exudate production is generally

maintained or increased (Henry et al. 2007; Preece and Peñuelas 2016; Karlowsky et al. 2018),

while the exudate composition is typically altered (Svenningsson et al. 1990; Canarini et al. 2016;

Calvo et al. 2017) which can contribute to the root microbiome drought response (Zhalnina et al.

2018). In *Sorghum*, for instance, drought-induced changes in root exudate composition and iron

homeostasis contribute to Actinomycetota enrichment in the rhizosphere (Xu et al. 2018, 2021).

Fungal communities of maize roots were less sensitive to drought relative to their bacterial

neighbors. While drought increased root fungal diversity in the plants inoculated with the TLI

prairie soil inoculum, it did not significantly affect overall fungal community composition.

Machine learning classification by drought treatment further illustrated this difference: when

trained on bacteria, we achieved 70% balanced accuracy, while for fungi, we achieved only 47%

accuracy. This discrepancy indicates that bacterial communities responded in a predictable manner

to drought, whereas fungal communities were less predictable between well-watered and drought

plants. This cross-kingdom difference agrees with previous reports, as both free living and root-

associated fungal communities tend to tolerate desiccation (de Vries et al. 2012; Furze et al. 2017).

The observed difference in survival between bacteria and fungi under drought may relate to both

the vegetative and dormant growth forms of fungi: compared to bacterial cells, fungal hyphae

comprise most vegetative growth and can cross air-gaps within dry soil in search of resources

(Griffin 1985; Manzoni et al. 2012), and dormant spores are generally desiccation-resistant (Wyatt

et al. 2013).

Historical precipitation, soil community legacies, and their impact on plant drought tolerance

Our hypothesis that historical precipitation patterns would impart legacy effects on soil

microbiomes that influence plant growth under drought stress was supported. Bacterial and fungal

compositional differences were apparent among the soils and inocula from prairie sites across a

steep precipitation gradient in Kansas (Figures 1&2). We found that the interaction between the

drought treatment and the mean annual precipitation of the soil inoculum source influenced shoot

mass accumulation rate. This pattern held for alternative climate metrics (PDSI, aridity index,

evapotranspiration, and soil moisture).

Following the stress gradient hypothesis (Bertness and Callaway 1994), we expected

drought-stressed maize seedlings to perform better in the presence of microorganisms from

historically dry soils, where chronic drought stress would have fostered positive interactions with

neighboring plants. However, we found that plants performed better when the historic conditions

of inocula were opposite from the contemporary conditions. The reason for this result is not

immediately clear, but there are several possible explanations. Contemporary well-watered

conditions may have relaxed the fitness consequences of expression of microbial traits that are

critical under drier environments. For instance, microorganisms counteract the physiologic stress

of drought by accumulating solutes, which reduces their internal water potential as a mechanism

to avoid desiccation (Harris 1981). The accumulation of solutes requires considerable energy

expenditure and constitutes a large sink of C and N resources (Killham and Firestone 1984;

Schimel et al. 1989, 2007). The contemporary well-watered conditions relax these costly solute

needs, perhaps allowing resources to be reallocated to functions that are beneficial to plants.

Another possibility is that soils from historically wetter sites generally possess greater microbial

biomass and diversity (Neilson et al. 2017; Araya et al. 2020; Wan et al. 2021), which is predicted

to promote positive interactions and improved community function (Hooper et al. 2005). We found

this to be the case for fungi, where our wettest prairie site showed the highest diversity, but this

was less consistent for bacterial communities (Figure S3-4). A third potential explanation involves

the balance of beneficial and pathogenic microbes in the soil community. Pathogen richness in

soil, root, and shoot microbiomes is often positively associated with mean annual temperature and

precipitation (Větrovský et al. 2019; Delgado-Baquerizo et al. 2020; Chen et al. 2021). Thus, soil

communities from historically dry sites may have relatively fewer plant pathogens, resulting in

greater plant performance under contemporary well-watered conditions. We cannot distinguish

between these possible explanations using the data in hand; however, a future experiment that

leverages rain exclusion structures to manipulate precipitation for a given site could effectively

monitor changes in diversity and composition over time. In addition, this experimental

manipulation strategy paired with a technique to quantify microbial biomass (e.g. quantitative

stable-isotope probing) (Greenlon et al. 2022) could enable the exploration of the balance between

beneficial, neutral, and pathogenic microbes.

Microbial taxa associated with plant growth

Multiple bacteria and fungi had abundances correlated with plant responses to drought,

with more taxa associated with shoot growth than root growth. This is likely the result of space

limitation in the pots, as root mass accumulation rates tended to decrease across the experiment

regardless of the drought treatment (Figure S19). Even so, we identified two families that were

associated with both shoot and root mass, Acetobacteraceae and Corynebacteriaceae.

Acetobacteraceae was associated with suppressed shoot growth but increased root growth.

Although in our dataset Acetobacteraceae was represented mainly by the poorly-studied genus

Rhodopila, this family contains many known diazotrophs, which may improve mineral nutrition

of the plant (Reis and Teixeira 2015). Members of Acetobacteraceae, including the particularly

well-studied *Gluconacetobacter diazotrophicus*, have been observed to colonize internal root

tissues under both natural conditions and inoculation experiments (Luna et al. 2010; Rouws et al.

2010). This root colonization proficiency may help to explain the increased root mass we observed

from this association; however, given the decreased shoot mass we also observed,

Acetobacteraceae may have compartment-specific dynamics that warrant further investigation. In

contrast, Corynebacteriaceae was associated with suppressed growth of both roots and shoots.

Members of this family have been identified as root endophytes of multiple plant species (Stafford

et al. 2005; Vik et al. 2013), with some showing phytopathogenic properties and specificity to

maize (Vidaver and Mandel 1974; Dye and Kemp 1977). Additionally, during a multi-generation

experimental evolution study, Corynebacteriaceae (among other bacterial families) was

consistently associated with delayed flowering in multiple plant species (Panke-Buisse et al. 2015).

Streptococcaceae was the only family that showed an association with plant growth that was

modulated by the drought treatment, with this family's relative abundance negatively correlated

with shoot mass accumulation rate in well-watered plants but not in droughted plants. Less is

known about the relevance of Streptococcaceae to plant growth. In *Zingiber officinale*, this family

was significantly enriched in rhizomes with advanced stages of soft rot (Huang et al. 2022).

However, Streptococcaceae is prevalent in soil and plants (Chuah et al. 2016; Wang et al. 2021),

typically at higher abundance in aerial plant compartments than in the rhizosphere (Minervini et

al. 2015; Yu et al. 2020). These families represent targets for direct manipulation and *in-planta*

testing to better understand the mechanism responsible for the growth suppression we observed

under well-watered conditions.

Conclusions

Water is a critical and limiting resource for plant growth as well as a beneficial microbiome.

As maize is cultivated widely across the United States, production will have to contend with more

frequent episodes of drought as the global climate warms. Engineering plant microbiomes to assist

in plant resilience under drought remains an ambitious, but worthwhile goal. Our work illustrates

the differential responses of root and shoot growth allocation to drought in two well-studied maize

genotypes and shows that bacterial, but not fungal, microbiota undergo restructuring with drought,

with decreases in the relative abundance of drought-intolerant taxa. In contrast to predictions based

on the stress gradient hypothesis, soil microbiomes from perennially dry environments along a

natural precipitation gradient did not enhance plant growth under drought. Instead, plants under

drought tended to produce more aboveground biomass with soil microbiomes from historically

wetter sites. The bacterial and fungal groups associated with biomass can serve to direct future

cultivation and *in-planta* testing on maize growth.

Acknowledgements

We thank members of the Wagner lab for helpful comments in preparing this article. In addition,

we thank Zachary Harris and Nichole Ginnan for helpful discussions. This work was funded by

the United States Department of Agriculture National Institute of Food and Agriculture (USDA-

NIFA) grant #2022-67013-36672 to MK and MRW, National Science Foundation grant #IOS-

2016351 to MRW, Novo Nordisk Foundation INTERACT project under award number

NNF19SA0059360 to MK, and seed funding from the NSF under award No. OIA-1656006 to Dr.

Kristin Bowman-James and matching support from the State of Kansas through the Kansas Board

of Regents. We thank the University of Kansas Medical Center – Genomics Core for generating

the sequence data sets. The Genomics Core is supported by the Kansas Intellectual and

Developmental Disabilities Research Center (NIH U54 HD 090216), the Molecular Regulation of

Cell Development and Differentiation – COBRE (P30 GM122731), The NIH S10 High End

Instrumentation Grant (NIH S10OD021743) and the Frontiers CTSA Grant (UL1TR002366).

Competing Interests

The authors declare that the research was conducted in the absence of any commercial or financial

relationships that could be construed as a potential conflict of interest. Any opinion, findings, and

conclusions or recommendations expressed in this material are those of the authors and do not

necessarily reflect the views of the United States Department of Agriculture or the National

Science Foundation.

Data Availability

Raw sequencing data is available on NCBI SRA under BioProject ID PRJNA913622 and SRA

accessions SAMN32302538-SAMN32303072 for root-associated samples and SAMN33863891-

SAMN33864082 for soil and inoculum samples. Code to reproduce the analysis and figures are

available on GitHub at https://github.com/Kenizzer/Maize_Drought_Microbiome.

Author Contributions

BAS and PMH collected soil samples across Kansas. MRK, MK, and MRW designed the

experiment and collection methods. MRK performed sample collection and processing. AC and

NEF performed DNA extractions and library preparations. NEF and JFS performed the data

analysis. JFS wrote the first draft of the manuscript. All authors contributed to data interpretation,

the writing and reviewing of the manuscript, and approved the final draft.

References

Abatzoglou JT, Dobrowski SZ, Parks SA, Hegewisch KC (2018) TerraClimate, a high-resolution global dataset of monthly climate and climatic water balance from 1958–2015. Sci Data 5:170191. https://doi.org/10.1038/sdata.2017.191

Acosta-Martínez V, Cotton J, Gardner T, et al (2014) Predominant bacterial and fungal assemblages in agricultural soils during a record drought/heat wave and linkages to enzyme activities of biogeochemical cycling. Appl Soil Ecol 84:69–82. https://doi.org/10.1016/j.apsoil.2014.06.005

Apprill A, McNally S, Parsons R, Weber L (2015) Minor revision to V4 region SSU rRNA 806R gene primer greatly increases detection of SAR11 bacterioplankton. Aquat Microb Ecol 75:129–137. https://doi.org/10.3354/ame01753

Araya JP, González M, Cardinale M, et al (2020) Microbiome Dynamics Associated With the Atacama Flowering Desert. Front Microbiol 10:1–13. https://doi.org/10.3389/fmicb.2019.03160

Averill C, Waring BG, Hawkes C V. (2016) Historical precipitation predictably alters the shape and magnitude of microbial functional response to soil moisture. Glob Chang Biol 22:1957–1964. https://doi.org/10.1111/gcb.13219

Bachar A, Al-Ashhab A, Soares MIM, et al (2010) Soil Microbial Abundance and Diversity Along a Low Precipitation Gradient. Microb Ecol 60:453–461. https://doi.org/10.1007/s00248-010-9727-1

Balting DF, AghaKouchak A, Lohmann G, Ionita M (2021) Northern Hemisphere drought risk in a warming climate. npj Clim Atmos Sci 4:61. https://doi.org/10.1038/s41612-021-00218-2

Bates D, Mächler M, Bolker B, Walker S (2015) Fitting Linear Mixed-Effects Models Using lme4. J Stat Softw 67:. https://doi.org/10.18637/jss.v067.i01

Benjamini Y, Hochberg Y (1995) Controlling the False Discovery Rate : A Practical and Powerful Approach to Multiple Testing. J R Stat Soc 57:289–300. https://doi.org/https://www.jstor.org/stable/2346101

Buchenau N, van Kleunen M, Wilschut RA (2022) Direct and legacy-mediated drought effects on plant performance are species-specific and depend on soil community composition. Oikos 2022:1–14. https://doi.org/10.1111/oik.08959

Callahan BJ (2017) RDP taxonomic training data formatted for DADA2 (RDP trainset 16/release 11.5)

Callahan BJ, McMurdie PJ, Rosen MJ, et al (2016) DADA2: High-resolution sample inference from Illumina amplicon data. Nat Methods 13:581–583. https://doi.org/10.1038/nmeth.3869

Calvo OC, Franzaring J, Schmid I, et al (2017) Atmospheric CO_2 enrichment and drought stress

modify root exudation of barley. Glob Chang Biol 23:1292–1304.
https://doi.org/10.1111/gcb.13503

Canarini A, Merchant A, Dijkstra FA (2016) Drought effects on Helianthus annuus and Glycine
max metabolites: from phloem to root exudates. Rhizosphere 2:85–97.
https://doi.org/10.1016/j.rhisph.2016.06.003

Canarini A, Schmidt H, Fuchslueger L, et al (2021) Ecological memory of recurrent drought
modifies soil processes via changes in soil microbial community. Nat Commun 12:5308.
https://doi.org/10.1038/s41467-021-25675-4

Chen QL, Hu HW, Yan ZZ, et al (2021) Precipitation increases the abundance of fungal plant
pathogens in Eucalyptus phyllosphere. Environ Microbiol 23:7688–7700.
https://doi.org/10.1111/1462-2920.15728

Chiang F, Mazdiyasni O, AghaKouchak A (2021) Evidence of anthropogenic impacts on global
drought frequency, duration, and intensity. Nat Commun 12:2754.
https://doi.org/10.1038/s41467-021-22314-w

Chuah L-O, Yap K-P, Thong KL, et al (2016) Genome Sequence of "Anthococcus," a Novel
Genus of the Family Streptococcaceae Isolated from Flowers. Genome Announc 4:.
https://doi.org/10.1128/genomeA.01410-16

De Long JR, Semchenko M, Pritchard WJ, et al (2019) Drought soil legacy overrides maternal
effects on plant growth. Funct Ecol 33:1400–1410. https://doi.org/10.1111/1365-
2435.13341

de Vries F, Lau J, Hawkes C, Semchenko M (2023) Plant–soil feedback under drought: does
history shape the future? Trends Ecol Evol 38:708–718.
https://doi.org/10.1016/j.tree.2023.03.001

de Vries FT, Liiri ME, Bjørnlund L, et al (2012) Land use alters the resistance and resilience of
soil food webs to drought. Nat Clim Chang 2:276–280.
https://doi.org/10.1038/nclimate1368

Delgado-Baquerizo M, Guerra CA, Cano-Díaz C, et al (2020) The proportion of soil-borne
pathogens increases with warming at the global scale. Nat Clim Chang 10:550–554.
https://doi.org/10.1038/s41558-020-0759-3

Dye DW, Kemp WJ (1977) A taxonomic study of plant pathogenic Corynebacterium species.
New Zeal J Agric Res 20:563–582. https://doi.org/10.1080/00288233.1977.10427375

Fernandes AD, Reid JNS, Macklaim JM, et al (2014) Unifying the analysis of high-throughput
sequencing datasets: characterizing RNA-seq, 16S rRNA gene sequencing and selective
growth experiments by compositional data analysis. Microbiome 2:15.
https://doi.org/10.1186/2049-2618-2-15

Field R, Hawkins BA, Cornell H V., et al (2009) Spatial species-richness gradients across scales:

a meta-analysis. J Biogeogr 36:132–147. https://doi.org/10.1111/j.1365-2699.2008.01963.x

Fitzpatrick CR, Copeland J, Wang PW, et al (2018) Assembly and ecological function of the root
microbiome across angiosperm plant species. Proc Natl Acad Sci 115:E1157–E1165.
https://doi.org/10.1073/pnas.1717617115

Fox J, Weisberg S (2019) An R Companion to Applied Regression, 3rd edn. Sage, Thousands
Oaks, CA

Fuchslueger L, Bahn M, Hasibeder R, et al (2016) Drought history affects grassland plant and
microbial carbon turnover during and after a subsequent drought event. J Ecol 104:1453–
1465. https://doi.org/10.1111/1365-2745.12593

Furze JR, Martin AR, Nasielski J, et al (2017) Resistance and resilience of root fungal
communities to water limitation in a temperate agroecosystem. Ecol Evol 7:3443–3454.
https://doi.org/10.1002/ece3.2900

Gao Y, Lynch JP (2016) Reduced crown root number improves water acquisition under water
deficit stress in maize (Zea mays L.). J Exp Bot 67:4545–4557.
https://doi.org/10.1093/jxb/erw243

Gardes M, Bruns TD (1993) ITS primers with enhanced specificity for basidiomycetes -
application to the identification of mycorrhizae and rusts. Mol Ecol 2:113–118.
https://doi.org/10.1111/j.1365-294X.1993.tb00005.x

Gloor GB, Macklaim JM, Pawlowsky-Glahn V, Egozcue JJ (2017) Microbiome Datasets Are
Compositional: And This Is Not Optional. Front Microbiol 8:1–6.
https://doi.org/10.3389/fmicb.2017.02224

Greenlon A, Sieradzki E, Zablocki O, et al (2022) Quantitative Stable-Isotope Probing (qSIP)
with Metagenomics Links Microbial Physiology and Activity to Soil Moisture in
Mediterranean-Climate Grassland Ecosystems. mSystems 7:.
https://doi.org/10.1128/msystems.00417-22

Griffin DM (1985) A comparison of the roles of bacteria and fungi, Volume 1: Springer US,
Boston, MA

Harris RF (1981) Effect of Water Potential on Microbial Growth and Activity. In: Parr JF,
Gardner WR, Elliott LF (eds) Water Potential Relations in Soil Microbiology, 9th edn. Soil
Science Society of America, Madison W.I., pp 23–95

Harrison XA, Donaldson L, Correa-Cano ME, et al (2018) A brief introduction to mixed effects
modelling and multi-model inference in ecology. PeerJ 6:e4794.
https://doi.org/10.7717/peerj.4794

Hawkes C V., Keitt TH (2015) Resilience vs. historical contingency in microbial responses to
environmental change. Ecol Lett 18:612–625. https://doi.org/10.1111/ele.12451

Hawkes C V., Kivlin SN, Rocca JD, et al (2011) Fungal community responses to precipitation.
Glob Chang Biol 17:1637–1645. https://doi.org/10.1111/j.1365-2486.2010.02327.x

Henry A, Doucette W, Norton J, Bugbee B (2007) Changes in Crested Wheatgrass Root
Exudation Caused by Flood, Drought, and Nutrient Stress. J Environ Qual 36:904–912.
https://doi.org/10.2134/jeq2006.0425sc

Hooper DU, Chapin FS, Ewel JJ, et al (2005) Effects of biodiversity on ecosystem functioning:
A consensus of current knowledge. Ecol Monogr 75:3–35. https://doi.org/10.1890/04-0922

Huang K, Sun X, Zou Y, et al (2022) Comparison of the Endophytic Bacterial Microbiota of
Asymptomatic and Symptomatic Ginger Rhizomes During the Activation of Adventitious
Bud Development. Plant Dis 106:2470–2479. https://doi.org/10.1094/PDIS-09-21-2069-RE

Kaisermann A, de Vries FT, Griffiths RI, Bardgett RD (2017) Legacy effects of drought on
plant–soil feedbacks and plant–plant interactions. New Phytol 215:1413–1424.
https://doi.org/10.1111/nph.14661

Karlowsky S, Augusti A, Ingrisch J, et al (2018) Land use in mountain grasslands alters drought
response and recovery of carbon allocation and plant-microbial interactions. J Ecol
106:1230–1243. https://doi.org/10.1111/1365-2745.12910

Killham K, Firestone MK (1984) Salt stress control of intracellular solutes in streptomycetes
indigenous to saline soils. Appl Environ Microbiol 47:301–306.
https://doi.org/10.1128/aem.47.2.301-306.1984

Korell L, Auge H, Chase JM, et al (2021) Responses of plant diversity to precipitation change
are strongest at local spatial scales and in drylands. Nat Commun 12:2489.
https://doi.org/10.1038/s41467-021-22766-0

Kuhn M (2008) Building Predictive Models in R Using the caret Package. J Stat Softw 28:159–
160. https://doi.org/10.18637/jss.v028.i05

Kuznetsova A, Brockhoff PB, Christensen RHB (2017) lmerTest Package: Tests in Linear Mixed
Effects Models. J Stat Softw 82:. https://doi.org/10.18637/jss.v082.i13

Lau JA, Lennon JT (2012) Rapid responses of soil microorganisms improve plant fitness in
novel environments. Proc Natl Acad Sci 109:14058–14062.
https://doi.org/10.1073/pnas.1202319109

Lau JA, Lennon JT, Heath KD (2017) Trees harness the power of microbes to survive climate
change. Proc Natl Acad Sci 114:11009–11011. https://doi.org/10.1073/pnas.1715417114

Lenth R, Love J, Herve M (2020) emmeans: Estimated Marginal Means, aka Least-Squares
Means. https://cran.r-project.org/package=emmeans

Luna MF, Galar ML, Aprea J, et al (2010) Colonization of sorghum and wheat by seed
inoculation with Gluconacetobacter diazotrophicus. Biotechnol Lett 32:1071–1076.

https://doi.org/10.1007/s10529-010-0256-2

Lynch JP, Chimungu JG, Brown KM (2014) Root anatomical phenes associated with water
acquisition from drying soil: Targets for crop improvement. J Exp Bot 65:6155–6166.
https://doi.org/10.1093/jxb/eru162

Manzoni S, Schimel JP, Porporato A (2012) Responses of soil microbial communities to water
stress: results from a meta-analysis. Ecology 93:930–938. https://doi.org/10.1890/11-0026.1

Martin M (2011) Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet.journal 17:10. https://doi.org/10.14806/ej.17.1.200

McMurdie PJ, Holmes S (2013) phyloseq: An R Package for Reproducible Interactive Analysis
and Graphics of Microbiome Census Data. PLoS One 8:e61217.
https://doi.org/10.1371/journal.pone.0061217

Minervini F, Celano G, Lattanzi A, et al (2015) Lactic Acid Bacteria in Durum Wheat Flour Are
Endophytic Components of the Plant during Its Entire Life Cycle. Appl Environ Microbiol
81:6736–6748. https://doi.org/10.1128/AEM.01852-15

Munoz-Ucros J, Wilhelm RC, Buckley DH, Bauerle TL (2022) Drought legacy in rhizosphere
bacterial communities alters subsequent plant performance. Plant Soil 471:443–461.
https://doi.org/10.1007/s11104-021-05227-x

Naylor D, Coleman-Derr D (2018) Drought Stress and Root-Associated Bacterial Communities.
Front Plant Sci 8:1–16. https://doi.org/10.3389/fpls.2017.02223

Naylor D, DeGraaf S, Purdom E, Coleman-Derr D (2017) Drought and host selection influence
bacterial community dynamics in the grass root microbiome. ISME J 11:2691–2704.
https://doi.org/10.1038/ismej.2017.118

Neilson JW, Califf K, Cardona C, et al (2017) Significant Impacts of Increasing Aridity on the
Arid Soil Microbiome. mSystems 2:. https://doi.org/10.1128/mSystems.00195-16

Oksanen J, Guillaume Blanchet F, Friendly M, et al (2019) vegan: Community Ecology Package

Panke-Buisse K, Poole AC, Goodrich JK, et al (2015) Selection on soil microbiomes reveals
reproducible impacts on plant function. ISME J 9:980–989.
https://doi.org/10.1038/ismej.2014.196

Parada AE, Needham DM, Fuhrman JA (2016) Every base matters: assessing small subunit
rRNA primers for marine microbiomes with mock communities, time series and global field
samples. Environ Microbiol 18:1403–1414. https://doi.org/10.1111/1462-2920.13023

Preece C, Peñuelas J (2016) Rhizodeposition under drought and consequences for soil
communities and ecosystem resilience. Plant Soil 409:1–17. https://doi.org/10.1007/s11104-
016-3090-z

R Core Team (2021) R: A Language and Environment for Statistical Computing

Reis VM, Teixeira KR dos S (2015) Nitrogen fixing bacteria in the family Acetobacteraceae and
their role in agriculture. J Basic Microbiol 55:931–949.
https://doi.org/10.1002/jobm.201400898

Remke MJ, Johnson NC, Wright J, et al (2021) Sympatric pairings of dryland grass populations,
mycorrhizal fungi and associated soil biota enhance mutualism and ameliorate drought
stress. J Ecol 109:1210–1223. https://doi.org/10.1111/1365-2745.13546

Rouws LFM, Meneses CHSG, Guedes HV, et al (2010) Monitoring the colonization of
sugarcane and rice plants by the endophytic diazotrophic bacterium Gluconacetobacter
diazotrophicus marked with gfp and gusA reporter genes. Lett Appl Microbiol 51:325–330.
https://doi.org/10.1111/j.1472-765X.2010.02899.x

Santos-Medellín C, Edwards J, Liechty Z, et al (2017) Drought Stress Results in a Compartment-
Specific Restructuring of the Rice Root-Associated Microbiomes. MBio 8:1–15.
https://doi.org/10.1128/mBio.00764-17

Schimel J, Balser TC, Wallenstein M (2007) Microbial stress-response physiology and its
implications for ecosystem function. Ecology 88:1386–1394. https://doi.org/10.1890/06-
0219

Schimel JP, Scott WJ, Killham K (1989) Changes in Cytoplasmic Carbon and Nitrogen Pools in
a Soil Bacterium and a Fungus in Response to Salt Stress. Appl Environ Microbiol
55:1635–1637. https://doi.org/10.1128/aem.55.6.1635-1637.1989

Schultz CR, Brantley KM, Wallace JG (2022) The role of genetic variation in Zea mays response
to beneficial endophytes. Plant Growth Regul 98:167–177. https://doi.org/10.1007/s10725-
022-00842-9

Sebastian J, Yee M-C, Goudinho Viana W, et al (2016) Grasses suppress shoot-borne roots to
conserve water during drought. Proc Natl Acad Sci 113:8861–8866.
https://doi.org/10.1073/pnas.1604021113

Sharp RE, Davies WJ (1979) Solute regulation and growth by roots and shoots of water-stressed
maize plants. Planta 147:43–49. https://doi.org/10.1007/BF00384589

Stafford WHL, Baker GC, Brown SA, et al (2005) Bacterial diversity in the rhizosphere of
Proteaceae species. Environ Microbiol 7:1755–1768. https://doi.org/10.1111/j.1462-
2920.2005.00929.x

Svenningsson H, Sundin P, Liljenberg C (1990) Lipids, carbohydrates and amino acids exuded
from the axenic roots of rape seedlings exposed to water-deficit stress. Plant, Cell Environ
13:155–162. https://doi.org/10.1111/j.1365-3040.1990.tb01287.x

UNITE Community (2017) UNITE general FASTA release

Větrovský T, Kohout P, Kopecký M, et al (2019) A meta-analysis of global fungal distribution
reveals climate-driven patterns. Nat Commun 10:1–9. https://doi.org/10.1038/s41467-019-
13164-8

Vidaver AK, Mandel M (1974) Corynebacterium nebraskense, a New, Orange-Pigmented
Phytopathogenic Species. Int J Syst Bacteriol 24:482–485.
https://doi.org/10.1099/00207713-24-4-482

Vik U, Logares R, Blaalid R, et al (2013) Different bacterial communities in ectomycorrhizae
and surrounding soil. Sci Rep 3:3471. https://doi.org/10.1038/srep03471

Wagner MR, Roberts JH, Balint-Kurti P, Holland JB (2020) Heterosis of leaf and rhizosphere
microbiomes in field-grown maize. New Phytol 228:1055–1069.
https://doi.org/10.1111/nph.16730

Wan X, Chen X, Huang Z, Chen HYH (2021) Global soil microbial biomass decreases with
aridity and land-use intensification. Glob Ecol Biogeogr 30:1056–1069.
https://doi.org/10.1111/geb.13282

Wang Q, Garrity GM, Tiedje JM, Cole JR (2007) Naïve Bayesian Classifier for Rapid
Assignment of rRNA Sequences into the New Bacterial Taxonomy. Appl Environ
Microbiol 73:5261–5267. https://doi.org/10.1128/AEM.00062-07

Wang X, Shi W, Wang J, et al (2021) Control of Southern Root-knot Nematodes on Tomato and
Regulation of Soil Bacterial Community by Biofumigation with Zanthoxylum bungeanum
Seed. Hortic Plant J 7:49–58. https://doi.org/10.1016/j.hpj.2020.12.005

White TJ, Burns T, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal
ribosomal RNA genes for phylogenetics. In: PCR protocols: a guide to methods and
applications. Academic Press, San Diego, CA, pp 315–322

Wright MN, Ziegler A (2017) ranger : A Fast Implementation of Random Forests for High
Dimensional Data in C++ and R. J Stat Softw 77:. https://doi.org/10.18637/jss.v077.i01

Wyatt TT, Wösten HAB, Dijksterhuis J (2013) Fungal Spores for Dispersion in Space and Time.
In: Advances in Applied Microbiology. pp 43–91

Xu L, Coleman-Derr D (2019) Causes and consequences of a conserved bacterial root
microbiome response to drought stress. Curr Opin Microbiol 49:1–6.
https://doi.org/10.1016/j.mib.2019.07.003

Xu L, Dong Z, Chiniquy D, et al (2021) Genome-resolved metagenomics reveals role of iron
metabolism in drought-induced rhizosphere microbiome dynamics. Nat Commun 12:3209.
https://doi.org/10.1038/s41467-021-23553-7

Xu L, Naylor D, Dong Z, et al (2018) Drought delays development of the sorghum root
microbiome and enriches for monoderm bacteria. Proc Natl Acad Sci 115:E4284–E4293.
https://doi.org/10.1073/pnas.1717308115

Yu AO, Leveau JHJ, Marco ML (2020) Abundance, diversity and plant-specific adaptations of
plant-associated lactic acid bacteria. Environ Microbiol Rep 12:16–29.
https://doi.org/10.1111/1758-2229.12794

Zarraonaindia I, Owens SM, Weisenhorn P, et al (2015) The Soil Microbiome Influences
Grapevine-Associated Microbiota. MBio 6:e02527-14. https://doi.org/10.1128/mBio.02527-
14

Zhalnina K, Louie KB, Hao Z, et al (2018) Dynamic root exudate chemistry and microbial
substrate preferences drive patterns in rhizosphere microbial community assembly. Nat
Microbiol 3:470–480. https://doi.org/10.1038/s41564-018-0129-3

Zhan A, Schneider H, Lynch JP (2015) Reduced Lateral Root Branching Density Improves
Drought Tolerance in Maize. Plant Physiol 168:1603–1615.
https://doi.org/10.1104/pp.15.00187

Supplemental Figures

Figure S1. Experimental design. A) A complete randomized block design was used; depicted is
a single representative block (number of blocks=27). Each pot was filled with sterile calcined
clay soil mix and received a seed of either *Zea mays* genotype B73 or Mo17, was subject to
either well-watered or drought conditions, and was inoculated with one of the four inocula. One
uninoculated control (denoted with white stripe) and three additional randomized replicates were
included per randomized block. **B)** Plant biomass was collected in a time series: for time point
one (day 25) and two (day 39), 68 plants each were destructively harvested, then dried for root
and shoot biomass. For the third time point (day 50), all remaining plants (n=379 – non-
germinates) were harvested. Prior to drying for biomass measures, one nodal root from each
plant was collected for amplicon sequencing for bacterial and fungal microbiome composition.

Figure S2. Climatic metrics for each collection site from 1990-2021. A) Annual precipitation, B) annual evapotranspiration, C) aridity index, D) average monthly Palmers Drought Stress Index (PDSI), and E) annual soil moisture. Boxplot hinges represent the 1st and 3rd quartiles; whiskers represent 1.5 times the interquartile range.

868
869 **Figure S3.** Bacterial alpha diversity values for **A)** Chao1 **B)** Shannon Diversity and **C)** Inverse
870 Simpson's Diversity. Black squares represent estimated marginal means (EMM) for each
871 collection site and colored bars show the 95% confidence intervals for the EMMs.

872
873   **Figure S4.** Fungal alpha diversity values for **A/C)** Chao1 and **B/D)** Inverse Simpson's Diversity.
874   Black squares represent estimated marginal means (EMM) for each collection site or substrate
875   and colored bars show the 95% confidence intervals for the EMMs.

876
877 **Figure S5.** Taxonomic bar plots for bacterial phyla for **A)** soil and **B)** inocula samples. Historical
878 annual precipitation for each collection site (SVR, HAY, TLI, KNZ) increases from left to right.
879 Each sample is represented by a single line with the relative abundance (non CLR transformed
880 data) of each phylum displayed by color.

881
Figure S6. Taxonomic bar plots for fungal classes for **A)** soil and **B)** inocula samples. Historical
annual precipitation for each collection site (SVR, HAY, TLI, KNZ) increases from left to right.
Each sample is represented by a single line with the relative abundance (non CLR transformed
data) of each class displayed by color.

Figure S7. Maize genotypes growth differs across **A)** shoot mass rate, **B)** root mass rate, and **C)**

root/shoot ratio. Boxplot hinges represent the 1st and 3rd quartiles; whiskers represent 1.5 times

the interquartile range; letters above box plots denote significant differences in estimated

marginal means (*p*-value=<0.01; see Table 1 for full ANOVA table).

891
892 **Figure S8.** Redundancy analysis for root bacterial community composition. Points are colored
893 by soil inoculum. Normal data ellipses are colored by soil inoculum assuming a multivariate
894 normal distribution.

895
896 **Figure S9.** Redundancy analysis for root bacterial community composition. Points are colored
897 by maize genotype.

898
899 **Figure S10**. Redundancy analysis for root fungal community composition. Points are colored by
900 soil inoculum. Normal data ellipses are colored by soil inoculum assuming a multivariate normal
901 distribution.

Figure S11. Redundancy analysis for fungal community composition. Points are colored by

maize genotype.

905
Figure S12. Confusion matrices depicting the classification accuracy of random forest classifiers
trained on bacterial data predicting **A)** Soil Inoculum and **B)** Genotype. The models were trained
using an 80:20 data split (80% train, 20% test) with 10-fold cross validation, shading represents
the proportion of predictions for a label. Labels on the left represent the predicted labels and the
labels on the bottom represent the actual labels. The model wide accuracy is given above each
confusion matrix, class-wise statistics are provided in table in Table S8.

Figure S13. Bacterial ASV importance plots per random forest classifier. Results for **A)**

Drought Treatment, **B)** Soil Inoculum, and **C)** Genotype classifiers. For each random forest

classifier, 10-fold cross validation was used to assess the mean decrease in accuracy of the

classifier when a particular ASV is withheld. Error bars represent the standard error across the

10-folds. Only ASVs with a mean decrease in accuracy greater than 0.001 are shown.

919
Figure S14. Confusion matrices depicting the classification accuracy of random forest classifiers
trained on fungal data predicting **A)** Soil Inoculum and **B)** Genotype. The models were trained
using an 80:20 data split (80% train, 20% test) with 10-fold cross validation, shading represents
the proportion of predictions for a label. Labels on the left represent the predicted labels and the
labels on the bottom represent the actual labels. The model wide accuracy is given above each
confusion matrix, class-wise statistics are provided in table in Table S9.

Figure S15. Fungal ASV importance plots per random forest classifier. Results for **A)** Drought Treatment, **B)** Soil Inoculum, and **C)** Genotype classifiers. For each random forest classifier, 10-fold cross validation was used to assess the mean decrease in accuracy of the classifier when a particular ASV is withheld. Error bars represent the standard error across the 10-folds. Only ASVs with a mean decrease in accuracy greater than 0.001 are shown.

Figure S16. Root growth by soil inoculum. Boxplots are colored by soil inoculum for **A)** root mass rate and **B)** root/shoot ratio (three R/S data points were removed to reduce the y axis range and aid in visualization). Separate facets represent drought-stressed and well-watered plants. Boxplot hinges represent the 1st and 3rd quartiles; whiskers represent 1.5 times the interquartile range.

Figure S17. Shoot growth rate across climatic metrics. Trend lines for shoot mass accumulation rate against **A-B)**
Evapotranspiration, **C-D)** Aridity Index, **E-F)** Soil Moisture, and **G-H)** Monthly Palmers Drought Stress Index (PDSI), for
interactions with drought treatment and genotype, respectively, for plants inoculated with prairie soils. For PDSI, colors represent the
prairie sites as PDSI values do not follow an east to west linear relationship (see Figure S2). For each regression line, grey shading
represents the 95% confidence interval.

944
945 **Figure S18.** The bacterial family Streptococcaceae showed a significant interaction between
946 CLR transformed abundance and drought treatment for residual shoot mass rate after accounting
947 for other experimental and technical factors . For each regression line, grey shading represents
948 the 95% confidence interval.

Figure S19. Plant growth rates across the collection time points of the study. A) Root mass
rate tended to decrease over the course of the experiment and was modulated by the drought
treatment (ANOVA; *'Timepoint × Treatment' p*-value = 0.02). **B**) Shoot mass rate tended to
increase over the course of the experiment (ANOVA; *'Timepoint' p*-value< 0.01). Boxplot
hinges represent the 1st and 3rd quartiles; whiskers represent 1.5 times the interquartile ran

955 **Supplemental Tables**

957 **Table S1.** ANOVA table for bacterial alpha diversity metrics of soil samples collected from six

958 locations across the Kansas precipitation gradient.

		ln(Chao1)			ln(Shannon's Diversity)			ln(Inverse Simpson's Diversity)		
Factor	DF	SS	F-value	p-value	SS	F-value	p-value	SS	F-value	p-value
Substrate	1	1.12	2.94	0.09	2.17	4.09	**0.05**	2.01	3.83	0.06
Collection Site	3	8.36	7.32	**<0.01**	7.64	4.80	**0.01**	6.50	4.12	**0.01**
Sequence Depth	1	69.73	183.32	**<0.01**	50.99	96.14	**<0.01**	32.09	60.99	**<0.01**
Residual	38	14.46			20.15			19.99		

959 Bold values denote significant effects, p-value<0.05

961 **Table S2.** ANOVA table for fungal alpha diversity metrics of soil samples collected from six

962 locations across the Kansas precipitation gradient.

		Ln(Chao1)			ln(Shannon's Diversity)			ln(Inverse Simpson's Diversity)		
Factor	DF	SS	F-value	p-value	SS	F-value	p-value	SS	F-value	p-value
Substrate	1	8.58	36.27	**<0.01**	0.02	0.06	0.80	0.85	1.87	0.18
Collection Site	3	0.84	1.19	0.32	2.25	2.08	0.11	3.46	2.55	0.06
Sequence Depth	1	10.89	46.01	**<0.01**	0.03	0.08	0.78	2.41	5.32	**0.02**
Residual	55	13.01			19.79			24.88		

963 Bold values denote significant effects, p-value<0.05

965 **Table S3.** PERMANOVA table for bacterial and fungal community composition of soil and

966 inocula samples collected from four locations across the Kansas precipitation gradient.

	Bacteria					Fungi				
Factor	DF	SS	R^2	F-value	p-value	DF	SS	R^2	F-value	p-value
Substrate	1	40923	0.13	7.96	**<0.01**	1	29645	0.10	8.22	**<0.01**
Collection Site	3	68034	0.22	4.41	**<0.01**	3	71904	0.24	6.64	**<0.01**
Residual	39	200477	0.65			56	202016	0.67		

967 Bold values denote significant effects, p-value<0.05

969 **Table S4.** ANOVA table for maize emergence rate.

	DF	ChiSq	p-value
Intercept	1	1035.92	**0.00**
Genotype (G)	1	7.12	**0.01**
Soil Inoculum (SI)	3	4.71	0.19
G x SI	3	2.51	0.47

970 Bold values denote significant effects, p-value<0.05

972 **Table S5.** ANOVA table for bacterial alpha diversity metrics of maize nodal roots.

Factor	DF	Chao1			ln(Shannon's Diversity)			Ln(Inverse Simpson's Diversity)		
		SS	F-value	p-value	SS	F-value	p-value	SS	F-value	p-value
Genotype (G)	1	11476	0.60	0.44	0.06	0.13	0.72	0.04	0.08	0.77
Soil Inoculum (SI)	3	179893	3.11	**0.03**	0.88	0.61	0.61	0.09	0.07	0.98
Drought Treatment (DT)	1	55466	2.88	0.09	0.20	0.42	0.52	0.62	1.42	0.24
G X SI	3	79499	1.37	0.25	0.88	0.61	0.61	0.78	0.59	0.62
G X DT	1	25967	1.35	0.25	0.28	0.57	0.45	0.24	0.55	0.46
SI x DT	3	34773	0.60	0.62	0.66	0.46	0.71	0.43	0.33	0.80
G X SI X DT	3	6233	0.11	0.96	0.07	0.05	0.99	0.23	0.18	0.91

973 Bold values denote significant effects, p-value<0.05
974

975 **Table S6.** ANOVA table for fungal alpha diversity metrics of maize nodal roots.

Factor	DF	Chao1			ln(Shannon's Diversity)			ln(Inverse Simpson's Diversity)		
		SS	F-value	p-value	SS	F-value	p-value	SS	F-value	p-value
Genotype (G)	1	107.32	0.59	0.45	0.56	1.62	0.21	0.71	2.47	0.12
Soil Inoculum (SI)	3	428.46	0.78	0.51	0.30	0.29	0.83	0.17	0.20	0.90
Drought Treatment (DT)	1	10.54	0.06	0.81	1.73	5.00	**0.03**	1.79	6.17	**0.02**
G X SI	3	805.97	1.47	0.23	0.53	0.51	0.68	0.63	0.72	0.54
G X DT	1	41.51	0.23	0.64	0.20	0.57	0.45	0.37	1.27	0.26
SI x DT	3	1044.15	1.90	0.14	2.71	2.60	0.06	2.12	2.44	0.07
G X SI X DT	3	557.96	1.01	0.39	1.05	1.01	0.39	1.06	1.22	0.31
Genotype (G)	1	107.32	0.59	0.45	0.56	1.62	0.21	0.71	2.47	0.12

976 Bold values denote significant effects, p-value<0.05
977

978 **Table S7.** Optimal hyperparameters for bacterial and fungal machine learning classifiers trained
979 to predict treatment, soil inoculum, and genotype.

Factor	Bacteria		Fungi	
	mtry	Minimum Node Side	mtry	Minimum Node Side
Treatment	559	10	45	10
Soil Inoculum	437	5	66	1
Genotype	559	5	10	1

981 **Table S8.** Output statistics for bacterial machine learning models predicting treatment, genotype,
982 and soil inoculum.

Metric	Treatment	Genotype	Soil Inoculum			
			SVR	HAY	TLI	KNZ
Precision	0.700	0.708	0.333	1.000	0.600	0.667
Recall	0.875	0.850	1.000	0.100	0.500	0.857
F1	0.778	0.773	0.500	0.182	0.546	0.750
Prevalence	0.593	0.741	0.148	0.370	0.222	0.259
Balanced Accuracy	0.665	0.425	0.826	0.550	0.702	0.854

984 **Table S9.** Output statistics for fungal machine learning models predicting treatment, genotype,
985 and soil inoculum.

Metric	Treatment	Genotype	Soil Inoculum			
			SVR	HAY	TLI	KNZ
Precision	0.500	0.625	0.667	0.875	0.333	0.400
Recall	0.400	0.882	1.000	1.000	0.200	0.400
F1	0.444	0.732	0.800	0.933	0.250	0.400
Prevalence	0.526	0.654	0.105	0.368	0.263	0.263
Balanced Accuracy	0.478	0.441	0.971	0.958	0.529	0.593

987 **Table S10.** Bacterial taxa across taxonomic levels associated with square root transformed shoot
988 mass rate.

Rank	Taxon	Trait	Model Term	R	X	PVE$_{Res}$	Rel. Abund.	q
Phylum	Actinobacteria	SMR	Abundance	-2.31E-05	-1.20E-02	5.69	13.46	2.74E-03
Class	Actinobacteria	SMR	Abundance	-2.31E-05	-1.20E-02	5.69	13.46	4.56E-03
Order	Actinomycetales	SMR	Abundance	-2.75E-05	-1.18E-02	5.81	13.43	7.49E-03
Order	Rhodocyclales	SMR	Abundance	-4.92E-04	-1.14E-02	3.51	0.06	4.14E-02
Order	Sphingomonadales	SMR	Abundance	-1.03E-04	-1.24E-02	5.31	1.13	8.05E-03
Family	Acetobacteraceae	SMR	Abundance	-1.25E-04	-1.33E-02	6.67	0.02	2.76E-03
Family	Corynebacteriaceae	SMR	Abundance	-5.46E-04	-1.29E-02	8.60	0.01	1.90E-04
Family	Dietziaceae	SMR	Abundance	-1.07E-03	-1.31E-02	4.94	0.00	6.48E-03
Family	Lactobacillaceae	SMR	Abundance	5.64E-04	-1.46E-02	4.26	0.09	1.55E-02
Family	Microbacteriaceae	SMR	Abundance	-1.07E-05	-1.32E-02	5.27	1.23	6.48E-03

Rank	Taxon	Trait	Model Term	R	X	PVE_Res	Rel. Abund.	q
Family	Micrococcaceae	SMR	Abundance	-8.43E-06	-1.31E-02	3.84	8.47	2.46E-02
Family	Nocardiaceae	SMR	Abundance	-3.71E-04	-1.33E-02	3.58	0.02	2.75E-02
Family	Rhodocyclaceae	SMR	Abundance	-4.92E-04	-1.14E-02	3.51	0.06	2.75E-02
Family	Sinobacteraceae	SMR	Abundance	2.12E-04	-1.38E-02	3.70	0.05	2.07E-02
Family	Sphingomonadaceae	SMR	Abundance	-1.03E-04	-1.24E-02	5.31	1.13	6.48E-03
Family	Streptococcaceae	SMR	Abundance	1.38E-04	-1.34E-02	5.50	0.01	4.10E-03
Genus	*Corynebacterium*	SMR	Abundance	-5.46E-04	-1.29E-02	8.60	0.01	3.61E-04
Genus	*Cupriavidus*	SMR	Abundance	-3.53E-04	-1.35E-02	3.95	0.02	3.55E-02
Genus	*Dietzia*	SMR	Abundance	-1.07E-03	-1.31E-02	4.94	0.00	1.45E-02
Genus	*Kribbella*	SMR	Abundance	-3.66E-04	-1.26E-02	3.81	0.03	3.55E-02
Genus	*Lactobacillus*	SMR	Abundance	5.64E-04	-1.46E-02	4.26	0.09	2.92E-02
Genus	*Minicystis*	SMR	Abundance	-7.91E-04	-1.32E-02	5.32	0.00	1.45E-02
Genus	*Modestobacter*	SMR	Abundance	-1.04E-03	-1.35E-02	4.29	0.01	2.92E-02
Genus	*Rhodococcus*	SMR	Abundance	-3.71E-04	-1.33E-02	3.58	0.02	4.79E-02
Genus	*Siphonobacter*	SMR	Abundance	3.24E-04	-1.56E-02	3.97	0.99	3.55E-02
Genus	*Sphingobium*	SMR	Abundance	-1.96E-04	-1.33E-02	4.84	0.01	2.13E-02
Genus	*Sphingomonas*	SMR	Abundance	-1.19E-04	-1.22E-02	4.32	1.15	2.92E-02
Genus	*Streptococcus*	SMR	Abundance	1.38E-04	-1.34E-02	5.50	0.01	1.17E-02
Family	Acetobacteraceae	RMR	Abundance	2.32E-05	-5.94E-03	8.15	0.02	1.41E-02
Family	Corynebacteriaceae	RMR	Abundance	-2.65E-04	-5.74E-03	6.29	0.01	3.39E-02
Phylum	Actinobacteria	RSR	Abundance	-3.31E-04	1.82E-01	6.78	13.46	6.35E-03
Phylum	Nitrospirae	RSR	Abundance	1.18E-02	1.64E-01	4.15	0.01	3.89E-02
Class	Actinobacteria	RSR	Abundance	-3.31E-04	1.82E-01	6.78	13.46	1.06E-02
Class	Deltaproteobacteria	RSR	Abundance	7.37E-03	1.55E-01	4.34	0.11	4.32E-02
Class	Nitrospira	RSR	Abundance	1.18E-02	1.64E-01	4.15	0.01	4.32E-02
Order	Actinomycetales	RSR	Abundance	-3.68E-04	1.84E-01	6.75	13.43	1.01E-02
Order	Sphingomonadales	RSR	Abundance	6.96E-04	1.58E-01	7.44	1.13	1.01E-02
Family	Polyangiaceae	RSR	Abundance	2.19E-02	1.37E-01	8.01	0.08	1.81E-02
Family	Sinobacteraceae	RSR	Abundance	-1.48E-03	1.67E-01	5.8	0.05	2.96E-02
Family	Sphingomonadaceae	RSR	Abundance	6.96E-04	1.58E-01	7.44	1.13	1.81E-02
Genus	*Minicystis*	RSR	Abundance	-7.91E-04	-1.32E-02	8.74	0.00	1.58E-02

990 **Table S11.** Fungal taxa across taxonomic levels associated with square root transformed shoot
991 mass rate.

Rank	Taxon	Trait	Model Term	R	X	PVE_Res	Rel. Abund.	q
Class	Microbotryomycetes	SMR	Abundance	3.33E-04	-1.24E-02	6.17	0.48	2.50E-02
Order	Sporidiobolales	SMR	Abundance	3.33E-04	-1.24E-02	6.17	0.48	2.95E-02
Family	Sporidiobolaceae	SMR	Abundance	3.33E-04	-1.24E-02	6.17	0.48	2.95E-02

992