Genetic sex determination in three closely related hydrothermal vent gastropods, one of which has intersex individuals

Castel J[1], Pradillon F[2] , Cueff V[2], Leger G[2], Daguin-Thiébaut C[1], Ruault S[1], Mary J[1], Hourdez S[3], Jollivet D[1], and Broquet T[1*]

1 UMR 7144 AD2M CNRS-Sorbonne Université, Station Biologique de Roscoff, Place Georges Tessier, 29680 Roscoff, France

2 Univ Brest, CNRS, Ifremer, UMR6197 Biologie et Ecologie des Ecosystèmes marins Profonds , F-29280 Plouzané, France

3 UMR 8222 LECOB CNRS-Sorbonne Université, Observatoire Océanologique de Banyuls, 1 Avenue Pierre Fabre, 66650 Banyuls-sur-Mer, France

*Corresponding author: thomas.broquet@sb-roscoff.fr

Keywords: deep-sea, molluscs, XY sex chromosomes, male heterogamety, ddRAD-seq.

Abstract

Molluscs have a wide variety of sexual systems and have undergone many transitions from separate sexes to hermaphroditism or vice versa, which is of interest for studying the evolution of sex determination and differentiation. Following the serendipitous observation that sex was the primary driver of genetic structure in the hydrothermal vent gastropod *Alviniconcha boucheti*, we investigated sexual systems and sex determination in this species and two others of the same genus. We combined genome-wide multi-locus genotypes obtained from RAD sequencing with anatomical observations of the gonads of the three *Alviniconcha* species occurring in the southwest Pacific Ocean: *A. boucheti* (*n*=207), *A. strummeri* (*n*=41 ind.) and *A. kojimai* (*n*=251). In two of the species (*A. boucheti* and *A. strummeri*), the sexes are separate and genetically determined by a male-heterogametic (XY) system. External observation of the gonads in the third species (*A. kojimai*) also suggested that the sexes were separate, but histological analyses revealed that 76% of the individuals classified as females from external observation of the gonads presented a mosaic of male and female reproductive tissue. Empirical analyses and simulations showed nonetheless that 14 RAD loci were sex-linked with an XY signature in *A. kojimai* (as compared with 64 in *A. strummeri* and 373 in *A. boucheti*). Comparison across species and mapping of RAD loci to a non-contiguous reference genome assembly of the related species *A. marisindica* showed that all sex-linked loci identified in *A. kojimai* are located on five scaffolds that also contain 25 and 73 sex-linked RAD loci in the other two species, respectively. These results suggest that all three *Alviniconcha* species share the same XY sex determination system, but that the gonad of XX *A. kojimai* individuals are invaded by a variable proportion of male reproductive tissue. It remains to be seen whether the male tissue in these intersex individuals is functional or not. This situation of three species showing variation in gonadal development around a probably common sex determination system provides new insights into the reproductive mode of poorly known deep-sea species and questions the evolution of gametogenetic polymorphism in these species.

Introduction

A wide variety of sexual systems are observed in eukaryotes, raising questions about the evolution of this polymorphism (recent reviews in Leonard, 2018; Pannell & Jordan, 2022). For instance, male and female gametes are produced by different individuals in about 95% of animal species (gonochoric species, Jarne & Auld, 2006), while an almost exact opposite figure was estimated for angiosperms (dioecious species, Renner, 2014). On closer examination, the seemingly simple, contrasted situation of animals and plants hides a diversity of sexual systems scattered within each of these broad taxonomic groups (and this diversity of course extends to other regions of the tree of life, e.g. Coelho & Umen, 2021). Considering animals, hermaphroditism (defined as a sexual system where male and female reproductive organs are simultaneously or sequentially found on the same individual) is nearly absent in insects but widespread across many other groups where it occurs in variable frequency (Jarne & Auld, 2006; Leonard, 2018). This means that hermaphroditism and gonochorism have evolved multiple times, in a wide variety of phylogenetic lineages. Some large clades (e.g. insects, tetrapods, chondrychtyes) have a mostly invariant, stable sexual system (Leonard, 2018; Pannell & Jordan, 2022), but others show much polymorphism. For instance, Jarne and Auld (2006) estimated that hermaphroditism evolved from gonochorism at least 40 times within prosobranch gastropods alone, raising questions as to how and why such transitions occur.

Evolutionary transitions between sexual systems are intimately linked with the evolution of sex determination and sex differentiation. Sex determination and differentiation can vary from purely genetic, when variation at some position in the genome triggers the male or female specific developmental program, to purely epigenetic, when sex-specific programs are activated by differential expression (Beukeboom & Perrin, 2014). A variety of determination systems along this continuum may lead to separate sexes (gonochory), but hermaphroditism is normally found towards the epigenetic end of the continuum (Beukeboom & Perrin, 2014). Although the evolution of sex determination is a lively field of research, many questions remain regarding the evolution of sex determination mechanisms involved in transitions between gonochory and hermaphroditism and the directionality

of these changes in animals (Leonard, 2018), and also in the evolution of complex sexual systems involving for instance a mixture of hermaphroditic individuals and males (androdioecy) or females (gynodioecy). Separate sexes may evolve *de novo* from hermaphroditism through a series of mutations that create one sex and then the other one, followed by selection for linkage between these loci (that is, formation and evolution of sex-chromosomes). This process and its molecular bases have been described in angiosperms (reviewed in Charlesworth, 2006) and have recently been investigated in Platyhelminthes and nematodes for instance (Wang *et al.*, 2022). In the reverse direction, transitions from separate sexes to hermaphroditism can also be achieved through a series of mutations affecting the sex determination pathway that will allow individuals of one sex to become hermaphrodites, which may provide them with a fitness advantage when access to mating partners is limited. For instance, self-fertile hermaphrodites have evolved from XX females following different modifications of the core sex determination cascade in *Caenorhabditis* species (therefore resulting in androdioecious systems, e.g. Kiontke *et al.*, 2004; Hill *et al.*, 2006).

Here we report the serendipitous discovery of genetic sex determination in three species of deep-sea hydrothermal vent symbiotic gastropods (genus *Alviniconcha*), as well as the unanticipated observation that one of these species contains individuals with a mixture of male and female reproductive tissue, therefore providing an interesting case of a mixed or transitory sexual system.

Gastropods exhibit a wide range of sexual systems and sex determination systems whose evolution has involved a high frequency of transitions (Jarne & Auld, 2006; Collin, 2013; Leonard, 2018). To our knowledge there is no known case of functional andro- or gynodioecy in this group (Weeks, 2012) but gastropods have been particularly studied in the context of exogeneous, anthropogenic, effects on intersex (defined in molluscs as the simultaneous presence of oocytes or spermatogonia, at varying degrees of development, within the normal gonad of the opposite gender of a gonochoristic species, Grilo & Rosa, 2017). Sex determination and sex differentiation of gastropods as a group seem to be very labile, which is interesting for understanding the evolution of these systems.

Although marine gastropods are primarily gonochoristic (reviewed in Collin, 2013), this is not a universal rule (e.g. simultaneous hermaphroditic Ophistobranchia, sex-changing Patellidae and Calyptreidae). In particular, we know very little about the biology of many deep-sea species because these environments are difficult to access. Hydrothermal vents are found along tectonically active locations such as mid-oceanic ridges and active volcanic arcs where seawater penetrates through faults in the lithosphere and resurfaces as a hot, anoxic, acidic fluid enriched in metals and dissolved gases. The chemical and physical characteristics of these peculiar environments have led to specific adaptations, including the establishment of symbioses with chemoautotrophic bacteria (Cavanaugh, 1983; Childress, 1995), and constraints on immune systems (Papot *et al.*, 2017), respiratory systems (Hourdez & Lallier, 2007), detoxification systems (Powell & Somero, 1986), and dispersal allowing the colonization of fragmented and ephemeral habitats (Lutz, 1988; Vrijenhoek, 2010). Regarding reproductive strategies, the hydrothermal environment seems to have generally favoured continuous gametogenesis, oocyte and sperm storage, internal fertilisation, and lecithotrophic development based on yolk reserves (Tyler & Young, 1999; Jollivet *et al.*, 2000; Faure *et al.*, 2007), although some abundant species do not follow this broad pattern (e.g. high fecundity and planktotrophic larvae in *Bathymodiolus* mussels, Laming *et al.*, 2018, and *Alviniconcha* gastropods). The sexual system of many hydrothermal vent species, not to mention their sex determination system, remains to be elucidated.

Here we studied *Alviniconcha* gastropods, large hairy snails that form dense aggregations in warm (7-42°C), sulphide-rich (up to 250 µmol.l^{-1}), and low-oxygen (< 50 µmol.l^{-1}) habitats associated with hydrothermal vents (Desbruyères *et al.*, 1994; Podowski *et al.*, 2010). There are six species in the genus (Johnson *et al.*, 2015; Breusing *et al.*, 2020; Castel *et al.*, 2022), all forming symbiotic associations with sulfo-oxidizing chemoautotrophic bacteria (Stein *et al.*, 1988). We focus here on *Alviniconcha boucheti*, *A. strummeri*, and *A. kojimai*, three species that inhabit the hydrothermal vents of the back-arc basins in the Western Pacific Ocean.

Following the unexpected observations of i) sex-linked patterns of genetic structure and ii) a proportion of male and female tissue mixed within the gonad of some individuals in *A. kojimai*, the

aims of this study were, firstly, to identify the sexual system and sex determination in the three species
105 and, secondly, to assess whether the genomic position of sex-linked loci is conserved across species to better understand the apparent flexibility of sexual systems in this genus.

Methods

Sampling and sexing

110 We used a subset of a previously published genomic data set of ddRAD-seq sequences from Castel et al. (2022). In short, here we used data from 251 *A. kojimai*, 207 *A. boucheti* and 41 *A. strummeri* individuals (total 499) sampled from 10 vent fields in four back-arc basins and a volcanic arc of the Western Pacific Ocean (Figure 1; Table S1). All samples were collected during the Chubacarc expedition on board of the R/V L'Atalante with the ROV Victor 6000 in Spring 2019 (Hourdez & Jollivet,
115 2019). Details about sampling conditions and the preparation of samples on board are described in (Castel *et al.*, 2022).

As no sexual dimorphism is externally visible in *Alviniconcha* gastropods, we determined the sex of 276 individuals from observations of the gonads using a two-step approach. First, sex was identified from the external colour of gonads for a fraction of individuals that were dissected on-board during
120 the sampling cruise (273 out of the 499 ind. that have been processed, 55%). The presence of oocytes gives a whitish colour to the gonads that indicates female sex while brown gonadal tissue is specific to males, but this method only allowed a rough identification of the sex, with some uncertainty. Second, a sub-sample of 170 individuals (34%) were studied using histological observations of the gonad (including 167 individuals previously sexed on board from gonad colour). A cross section of the central
125 part of the formalin-fixed gonad was taken and embedded in paraffin. Sections 5-μm thick were then cut using a Leica RM2035 microtome and stained with standard histological protocols for Harry's haematoxylin and eosin. The stained sections were then observed using a Leica DM2500 LED binocular microscope. The sex-ratio was not estimated from these internal observations of the gonads because

these individuals were not strictly randomly chosen onboard (we tended to balance the number of males and females selected for dissection to study the gametogenesis in the three species).

Figure 1: Origin of samples. Manus Basin: Susu (Su) and Pacmanus (Pac). Woodlark Basin: La Scala (Sca). North-Fiji Basin: Phoenix (Phoe). Futuna volcanic arc: Fatu-Kapa (FK) and Fati-Ufu (FU), these two sites are indistinguishable at this scale. Lau Basin: Mangatolo (Ma), Tow Cam (TC), ABE (ABE), and Tu'i Malila (TM). Species *A. boucheti* is represented in purple, *A. kojimai* in turquoise, and *A. strummeri* in yellow.

Genotyping

After DNA extraction from foot tissue using NucleoSpin® Tissue kits (Macherey-Nagel) or a CTAB protocol (Doyle & Dickson, 1987; modified in Jolly *et al.*, 2003), individual-based ddRAD libraries were produced following the protocol detailed in Daguin-Thiébaut et al. (2021). The libraries were sequenced in 150-bp paired-end reads using a Novaseq 6000 (library construction and sequencing detailed in Castel *et al.*, 2022).

Raw reads were demultiplexed using the process_radtags module of Stacks software version 2.52 (Catchen *et al.*, 2011, 2013) to remove adapters and reads with low quality scores (Phred score < 30). We used Kraken v.2 (Wood & Salzberg, 2014) to remove potential reads originating from chemoautotrophic bacteria and keep only the eukaryotic reads for the assembly (between 0.9% and 1.5% of the initial reads assigned to prokaryotes were discarded, depending on the species considered). Average sequence quality per read and GC-content were checked using Fastqc and multiQC version 1.7. Barcode sequences were trimmed and the remaining 144 bp reads were then treated independently for each species.

The assembly parameters for each species were chosen on the basis of pilot studies consisting of a systematic exploration of the effect of the main Stacks parameters *m*, *M*, and *n*. Details of the pilot analyses and final conditions of data treatment are presented in the supplementary material. The SNPs identified by Stacks were filtered in R version 4.1.1 (R Core Team, 2021) to keep the SNPs present in at least 90% of the individuals and only keep the individuals which were genotyped at least at 80% of the total number of SNPs (allowing a maximum of missing data of 20% for a given individual), except in *A. strummeri* for which we had less samples and thus permitted slightly more missing data per individual (25%).

Our final data set contained 207 individuals and 163 117 SNPs for *A. boucheti*, 251 individuals and 70 122 SNPs for *A. kojimai,* and 41 individuals and 36 568 SNPs for *A. strummeri*. We visualized the genetic structure of each species using principal components analyses in R (adegenet package, Jombart, 2008).

Because sex appeared to be a key driver of genetic structure (see Results), the distribution of genetic diversity was further investigated using functions in the R package hierfstat (Goudet & Jombart, 2021), using only the sexed individuals (100 *A. boucheti*, 151 *A. kojimai*, and 22 *A. strummeri*, see table 1). We characterised the global genetic diversity within each species by estimating total gene diversity H_T. The diversity within each sex *i* was then described by calculating the gene diversity $H_{S,i}$ and observed heterozygosity $H_{O,i}$, and the discrepancy between these parameters (indicating a

potential departure from random association of gametes) was estimated as F_{IS} within each sex. Finally, the genetic differentiation between sexes was estimated using Weir and Cockerham (1984) estimator of F_{ST} between males and females and tested for statistical significance using 100 bootstraps over loci

175   in StAMPP (Pembleton *et al.*, 2013).

Identification of the genetic sex determination system

        Identifying sex chromosomes can be achieved by looking for loci that are present on only one of the sex chromosomes (Gautier, 2014; Stovall *et al.*, 2018; Feron *et al.*, 2021; Garcia-Raventós *et al.*,

180   2023), or by looking at heterozygosity and differentiation between sexes at homologous loci (e.g. Berset-Brändli *et al.*, 2006; Brelsford *et al.*, 2017; Hearn *et al.*, 2022; see also Pyne *et al.*, 2022 for an association study). Here we followed the second approach. To detect genetic systems of sex determination, we searched for SNP markers with alleles differentially fixed on sex chromosomes. If sex was genetically determined and if some SNPs in a non-recombining region had alleles that were

185   differentially fixed on sex chromosomes, then these SNPs would be strictly homozygous in the homogametic sex (XX females or ZZ males) and heterozygous in the heterogametic sex (XY males or ZW females). We looked for such SNPs by estimating the observed heterozygosity in males $H_{O,m}$ and females $H_{O,f}$ at each SNP using functions from the hierfstat R package (Goudet & Jombart, 2021).

190   *Identification of sex-linked loci*

        Besides sex-linked SNPs with fixed, divergent alleles on sex chromosomes, other polymorphic SNPs located on sex chromosomes may have various levels of heterozygosity and sex-linked differentiation depending on the genomic landscape of recombination along these chromosomes. The distribution of genetic diversity between sexes (F_{ST}) and between individuals within each sex (F_{IS}) is

195   more informative than H_O to detect sex-linked loci that are not differentially fixed, because sex linkage will tend to result in an excess of heterozygotes in the heterogamous sex (e.g. XY males) and a simultaneous increased differentiation between sexes.

However, we expect some variance associated with these statistics measured using a finite number of male and female individuals (Table 1). Therefore, for each species we computed F_{IS} and F_{ST}

200   values from simulated genotypes randomly created from a range of allelic frequencies assuming autosomal segregation for two groups of individuals with sample sizes equal to the actual size of the male and female groups in our real dataset (thereby introducing estimation of error variance in the simulations). The simulations were coded in R and consisted of the following steps:

1- Create a number of theoretical bi-allelic SNPs equal to the number of SNPs genotyped in each

205   species and where each SNP has a reference allele with frequency p and an alternative allele with frequency 1-p, where p was randomly sampled from a uniform distribution between 0 and 1 (such as to cover all possible initial values for allelic frequencies).

2- Create a group of "male" and a group of "female" genotypes (with sample sizes equal to the number of males and females sampled in each species) by sampling two alleles randomly from a

210   binomial distribution parameterized using the expected allelic frequencies simulated in step 1.

3- From these simulated genotypes, estimate F_{ST} between the "male" and "female" groups, and F_{IS} within each of these groups at each simulated SNP. These estimates were calculated following (Nei, 1987) as described in R package hierfstat and simplified for bi-allelic SNPs:

$$F_{ST} = (H_T - H_S)/H_T$$

215   and

$$F_{IS} = (H_{S,i} - H_{O,i})/H_{S,i} \text{ for sex } i$$

where:

$$H_T = 2p(1 - p) + H_s/(n_f + n_m) - H_O/2(n_f + n_m)$$

$$H_S = (n_f H_{S,f} + n_m H_{S,m})/(n_f + n_m)$$

220    $$H_{S,i} = n_i/(n_i - 1)\left[2p_i(1 - p_i) - H_{O,i}/2n_i\right]$$

with p the allelic frequency of the reference allele (recalculated from the simulated genotypes) at the SNP under consideration, n_i is the sample size of sex i. In these equations, $2p(1-p)$ is the gene diversity expected under random mating (Hardy-Weinberg, autosomal segregation at a bi-allelic loci

with allelic frequencies p and (1-p)), and all other terms are bias corrections for small sample sizes as described in Nei (1987).

These statistics were computed for each SNP. After simulation, observed F_{ST} and sex-specific F_{IS} values at each SNP from our dataset can be compared to the simulated values. All SNPs from the real dataset that had combinations of F_{IS} and F_{ST} values outside of a convex envelope encompassing all possible theoretical values obtained from autosomal simulations had properties that were not compatible with autosomal segregation. The convex envelope was built using slightly modified R code from F. Briatte (https://github.com/briatte/moralecon/blob/master/moralecon.r).

Synteny of sex-linked loci across the three species

As we used different SNP catalogues for each species in Stacks, sex-linked RAD loci identified within each species were compared to each other through reciprocal blastn (e-value threshold= 10^{-5}, word size=28) in Geneious (Kearse *et al.*, 2012) in order to identify orthologous RAD loci between two or three species.

In a second step, we search for the position of all sex-linked RAD sequences in the closest reference genome available (*A. marisindica*, assembly ASM1885773v1, Yang *et al.*, 2022). Although the level of divergence expected between *A. marisindica* and the species studied here prevented us from using this genome for processing RAD-sequences (we used a de-novo approach instead, see methods in Castel *et al.*, 2022) and although this reference genome is not contiguous (3926 scaffolds, N50 = 0.73 Mb, L50 = 336 contigs), it is still useful to identify potential RAD sequences that would be obtained in only one or two of the three species but belonging to a single genomic scaffold. To this end, we ran a blastn analysis using the software Geneious to identify the 100 best hits of each sex-linked RAD locus on the *A. marisindica* assembly, with minimum coverage of 75% of the RAD sequence and an e-value threshold of 10^{-5}.

Results

Gender identification and sex ratio

The sex of *A. boucheti* individuals was determined by eye on-board by external observation of the gonads for 101 individuals and later reassessed in the lab by histological observations for 55 of these individuals and two others. Of the 55 possible comparisons between the two observations, only 5 mismatches were observed. That is, sex identification based on external observation of the gonads generated about 9% error. Combining the two datasets (as reported in Table S1), we obtained 49 females and 54 males.

For *A. strummeri*, 22 individuals were sexed on-board, and all of them were also used for gonad cross-sections. We observed 2 discrepancies, that is, again 9% error from external gonad observation. Histological observations indicated that we had 12 females and 10 males.

In *A. kojimai*, 68 females and 82 males were identified on-board. Surprisingly, from the subset of 41 females later used in histological analyses, 31 presented a gonad cross-section that included some proportion of male tissue, as shown in figure 2. Visual inspection of gonad sections from these individuals revealed variable proportions of male tissue. By contrast, histological observations confirmed the sex of 48 out of the 49 males for which we took gonad cross-sections (the remaining individual was a female wrongly identified from external observation of the gonad, which brings the total number of females identified by histology to 11).

Focusing on data from histology only, we identified 49 males, 11 females, and 31 'morphological hermaphrodites' or 'intersex individuals' (which we define as individuals with coexisting male and female gonadal tissue but no information on the functionality of these tissues, see discussion).

Because our histological observations were limited to a single middle cross section per individual, it is possible that some male tissue was missed in the individuals identified as females (see discussion). Therefore, we considered two groups of *A. kojimai* individuals in downstream analyses, depending on their sex identified on-board and/or from gonad sections: a group of males (n = 82) and

a group of 'females or morphological hermaphrodites' (*n* = 69). See the discussion for a detailed
argumentation about this dichotomy.

Figure 2: Gonad of an *A. kojimai* 'morphological hermaphrodite'. A) Gonad removed from fresh tissue
on board. Note its overall white colour used as a diagnostic for the female sex on board, while the
brownish spots scattered over it reflect male tissues. B) Transverse cross-section of a formalin-fixed
gonad (same individual as in A), taken approximately at the middle of the antero-posterior axis.
Asterisks indicate male tissue. C) Hematoxylin-eosin coloured semi-thin section prepared from the
section shown in B where male tissues appear in blue (also marked with asterisks) and female tissues
appear in pink. D) Close-up of gonadal tissues with both male and female gametes developing in close
proximity. dg: digestive gland, go: gonad, ovo: oocytes, spz: spermatozoa. Scale bars are 5 mm in A,
1 mm in B & C, 100 μm in D.

 Distribution of genetic variation

The principal components analysis of SNP genotypes in *A. boucheti* revealed an unexpected pattern (Fig. 3A): the first axis (1% of the total variance) separated two groups of individuals clearly determined by their sex, except for three individuals that segregated with the opposite sex. Two of these three individuals had been processed consecutively on-board and their numbering was most likely inverted by mistake. The sex of the third one was identified from external observation only, and given the rate of error of this method it is possible that sex was wrongly identified. All other sexed individuals (i.e. 48 females and 52 males) clustered into two distinct groups along the axis 1 of the PCA. In addition, the 104 unsexed individuals were evenly distributed between the two clusters (52 with the male cluster and 52 with the female cluster, Fig. 3A). The second axis (0.7% of the total variance) further subdivided the male and female groups according to the main geographical subdivision of sampling locations: Western sites (Manus and Woodlark basins) versus Eastern sites (Lau basin and Futuna arc), which are separated by well over 2000 km (Fig. 1).

Remarkably, sex was also a key driver of genetic variation in *A. strummeri* (Fig. 3B), but with a limited effect since it separated individuals along the second component of the PCA (3% of the total variance), while the geographical distribution of sites explained the distribution of genotypes along the first axis (4.3% of the total variance, despite the smaller geographic range of this species). The clustering of individuals according to axis 2 (sex) was also less clear-cut than in the case of *A. boucheti*, making it more uncertain to estimate the number of unsexed individuals in either group.

Finally, there was no visible effect of sex (that is, considering males versus females combined with morphological hermaphrodites) in the PCA of *A. kojimai* genotypes (Fig. 3C). With this species, the only obvious structuring factor was the main West-East geographic separation of sampling locations (PCA axis 1, 1.2% of the total variance), while axis 2 (0.6% of the total variance) was driven by a single individual (not shown in Fig. 3C, see Fig. S4). Importantly, this pattern was left unchanged when considering three groups (males, females, and morphological hermaphrodites identified from histology) rather than two (Fig. S5). Only one morphological hermaphrodite was identified in the

Western region, but this is due to the fact that histological analyses concentrated on Eastern sites.

Within the Eastern region, no genetic structure was apparent between morphological hermaphrodites, females, and males (Fig. S5).

Figure 3: Principal components analyses of SNP genotypes for the three species of *Alviniconcha* gastropods. Solid symbols correspond to the Eastern basins, while empty symbols correspond to the Western basins (Figure 1). A) *A. boucheti* (207 individuals, 163 117 SNPs). The first axis separates individuals according to their sex, while the second axis is consistent with geography. B) *A. strummeri*

325   (41 individuals, 36 568 SNPs). While the first axis separates individuals according to their geographic

origin, the second axis is driven by sex. C) *A. kojimai* (251 individuals, 70 122 SNPs). The first axis

separates individuals according to their geographic origin. The second axis is driven by a single

individual (not shown here, see Fig S4 in supplementary material). There is no obvious clustering by

sex in this species.

To quantify the role of sex in the distribution of genetic variation in the three *Alviniconcha*

species, we estimated population genetics parameters considering males and females (or females

combined with morphological hermaphrodites in the case of *A. kojimai*) for each species (Table 1). The

genetic differentiation between sexes was low but significantly different from zero in all three species.

335 In agreement with results from PCA, it was strongest in *A. boucheti* (F_{ST} = 0.0105, *p*-value < 0.001),

followed by *A. strummeri* (F_{ST} = 0.0024, *p*-value < 0.001) and *A. kojimai* (F_{ST} = 0.0002, *p*-value = 0.037),

reflecting minute differences in gene diversity within sexes versus overall individuals (Table 1). In

species *A. kojimai* there was no significant differentiation between sexes when calculated with a

reduced dataset composed by individuals identified as males, females, and morphological

340 hermaphrodites from histology exclusively (see supplementary material text and Table S2).

Table 1: Diversity indices for the three species of *Alviniconcha*. H_T : total gene diversity, H_S : sex-specific gene diversity, where *m*

is for male and *f* stands for female in *A. boucheti* and *A. strummeri*, and either female or morphological hermaphrodites in *A.*

kojimai (see main text); F_{ST} : differentiation between sexes; $H_{O,i}$: observed heterozygosity in sex *i* and $F_{IS,i}$: heterozygosity

deficiency in sex *i*. Column "Unsexed" gives the number of individuals with uncertain or unknown phenotypic sex (none of these

individuals were used in diversity calculations).

Species	Males	Females	Unsexed	Total	SNPs	H_T	$H_{s,m}$	$H_{s,f}$	F_{ST}	$H_{o,m}$	$H_{o,f}$	$F_{IS,m}$	$F_{IS,f}$
A. boucheti	52[a]	48[a]	107[a]	207	163 117	0.14	0.14	0.138	0.0105	0.14	0.136	0.001	0.017
A. strummeri	10	12	19	41	36 568	0.267	0.267	0.266	0.0024	0.259	0.258	0.03	0.031
A. kojimai	82	69	100	251	70 122	0.111	0.112	0.111	0.0002	0.11	0.109	0.014	0.016

[a] Numbers of males, females, and unsexed individuals in *A. boucheti* differ slightly from those reported in table S1 since here the

three individuals with inconsistent phenotypic vs genotypic sexes were reclassified as "Unsexed" (see text).

Sex determination system

In *A. boucheti*, 553 SNPs on 237 RAD-loci were strictly homozygous in females (n = 48) and heterozygous in males (n = 52, Figure 4a, top-left corner), pointing towards a XY genetic system of sex determination where each of these SNPs has one allele fixed on the X chromosome and the other fixed on the Y. By contrast, no SNP was found to be indicative of a ZW system. Furthermore, a large number of additional SNPs showed a combination of strong $H_{O,m}$ and low $H_{O,f}$, in agreement with the hypothesis of a XY system where SNPs will be linked to the sex chromosome but located in a still recombining region (see next section below).

For *A. strummeri*, although we had much fewer sexed individuals (12 females and 10 males) and thus the error associated with $H_{O,m}$ and $H_{O,f}$ at each locus was higher, we found that 92 SNPs (on 55 RAD-loci) were strictly homozygous in females and heterozygous in males (Figure 4b, top-left corner), again indicating an XY genetic system of sex determination.

Surprisingly, in the species *A. kojimai*, where at least a third of the individuals have a mixture of male and female gonadal tissues, we found 9 SNPs (on 7 RAD-loci) with strong H_O in males but low in hermaphrodites/females (top-left corner of Figure 4c). This pattern, which remained consistent when separating females and morphological hermaphrodites (Fig. S6), again pointed towards a XY-like system where males are XY and morphological hermaphrodites / females are XX (see Discussion). However, this sex determination was supported by a very small number of SNPs (9 out of 70 122, about 0.13 %) and contrary to the two other species, none of these SNPs were strictly homozygous in females and heterozygous in males.

In all three species, a number of SNPs had simultaneously high $H_{O,m}$ and $H_{O,f}$ values (top-right corner of Figs. 4a-c) possibly indicative of paralogous loci that were wrongly assembled together, or, perhaps less likely, sites under strong balancing selection.

Figure 4: Locus by locus female and male-specific observed heterozygosity $H_{O,f}$ and $H_{O,m}$ in three *Alviniconcha* species. A) *A. boucheti*: 163 117 SNPs (48 females and 52 males). In the top-left corner, 553 SNPs (on 237 RAD) have $H_{O,f} = 0$ and $H_{O,m} = 1$, hence compatible with position of these SNPs on XY sex chromosomes with distinct alleles fixed on X and Y ; B) *A. strummeri*: 36 568 SNPs (12 females and 10 males). In the top-left corner, 92 SNPs (on 55 RAD) have $H_{O,f} = 0$ and $H_{O,m} = 1$. Note that sample size was low in this species (22 sexed individuals), resulting in noisy estimates of $H_{O,f}$ and $H_{O,m}$; C) *A. kojimai*: 70 122 SNPs (69 hermaphrodites/females and 82 males). In the top-left corner, nine SNPs (on 7 RAD) showed strong heterozygosity in males but low heterozygosity in hermaphrodites/females.

Sex-linked loci

In *A. boucheti*, 1 011 SNPs (on 373 RAD-loci) had a combination of low $F_{IS,m}$ and strong F_{ST} compatible with XY features that were never obtained by simulation of autosomal segregation (Figs. 5A-B). This represented ca. 0.7% of the set of SNPs for which $F_{IS,m}$ and F_{ST} could be calculated (blue dots in Fig. 5B, *n* = 143 248).

Similarly, in *A. strummeri*, 111 SNPs (on 64 RAD-loci) out of 32 397 SNPs had $F_{IS,m}$ and F_{ST} that were never obtained by simulation of autosomal SNPs (Figs. 5C-D). These SNPs represented 0.34% of the markers analysed here (blue dots in Fig. 5F, *n* = 32 397), but the power for detecting sex-linked SNPs was very reduced for *A. strummeri* because only 22 individuals were sexed, and this number is often even slightly lower for each specific locus (due to locus-specific missing data).

Figure 5: Simulated and observed F_{ST} values between males and females and sex-specific F_{IS} for three *Alviniconcha* species (top row: *A. boucheti*, middle: *A. strummeri*, bottom: *A. kojimai*). Panels A, C, and E show simulated values obtained for autosomal SNPs with a range of expected allelic frequencies (uniform between 0 and 1) and using real male and female sample sizes from each species (table 1). The grey polygon is a convex envelope that covers all possible values obtained by simulation for an autosomal locus. Panels B and D show observed values for male F_{IS} (in blue) or female F_{IS} (pink) and male/female F_{ST} (i.e., each SNP is represented once in blue and once in pink according to each gender).

395   Panel F is similar but, in this case, pink symbols refer to morphological hermaphrodites and females,

and F_{ST} was calculated between males and morphological hermaphrodites/females. Any SNP outside

of the grey polygon has properties that were never obtained by neutral autosomal simulations.

    Finally, in *A. kojimai*, 16 SNPs (on 14 RAD-loci) had $F_{IS,m}$ and F_{ST} values that were never obtained

400   by simulation of autosomal SNPs (Figs. 5E-F). This represented 0.025% of the set of SNPs for which $F_{IS,m}$

and F_{ST} could be calculated (*n* = 64 709).

    In line with the expectations for XY sex chromosomes for the three species, all SNPs showing

strong differentiation between sexes (F_{ST}) tended to have negative F_{IS} values in males (as stated above)

and null F_{IS} in females (shown in pink in Fig. 5).

Synteny of sex-linked loci in the three Alviniconcha species

    A reciprocal BLASTn of sex-linked loci between pairs of species revealed that 28 of them were

common to at least two species (including 3 RAD loci common to the three species, Fig. 6A). In

particular, 10 of the 14 RAD loci identified as sex-linked in *A. kojimai* were also found to be sex-linked

410   in *A. boucheti* (*n* = 4), *A. strummeri* (*n* = 3), or both (*n* = 3). By contrast, a large proportion of loci were

found to be sex-linked in *A. boucheti* only (n = 348), or *A. strummeri* only (n= 40), which may be due to

differences in sample sizes and power or the length of sex chromosomes (see discussion).

    Thirteen of the 14 sex-linked RAD-loci identified in *A. kojimai* were retrieved by blastn on 5

scaffolds of the *A. marisindica* genome assembly (the remaining locus also mapped on one of these

415   scaffolds, but with a size coverage of less than 75%). The total length of these scaffolds amounted to

2.8 Mb (Fig. 6B). Most importantly, these 5 scaffolds were also found to include sex-linked RAD-loci

for the two other species: *A. boucheti* (73 RAD-loci out of 373) and *A. strummeri* (25 RAD-loci out of

64).

    Out of the 64 sex-linked loci identified in *A. strummeri*, 34 loci were distributed on 11 scaffolds

420   of the *A. marisindica* genome (total length 3.7 Mb which also includes the 5 scaffolds identified with

kojimai loci). The remaining loci did not return any significant hits. Out of these 11 scaffolds, 9 were also found to bear sex-linked RAD-loci in *A. boucheti* (total length 3.6 Mb, including the region that is sex-linked in all three species), while the other two (total length 0.12 Mb) seem to be sex-linked only in *A. strummeri*.

425   Finally, 300 of the 373 sex-linked RAD loci identified in *A. boucheti* segregated on 48 scaffolds, with a total length of 25.2 Mb (including 21.6 Mb bearing RAD-loci that were found to be sex-linked only in *A. boucheti*).

Figure 6: A) Distribution of sex-linked RAD-loci shared by *A. boucheti* (100 individuals, 373 sex-linked
430   RAD loci), *A. strummeri* (22 ind., 64 loci) and *A. kojimai* (151 individuals, 14 loci). B) : total length and interspecies distribution of genomic scaffolds found to be sex-linked in each species, using *A. marisindica* reference genome (Yang *et al.*, 2022). These sizes are indicative and should not be taken as face values since they depend on the contiguity of the genome assembly used as a reference (especially in *A. boucheti*) and the unknown frequency of repeated elements on sex chromosomes.

Discussion

The observation that sex is the primary driver of the genetic structure of *A. boucheti* (Figure 3A) was an unexpected by-product of our analyses of the genetic structure of *Alviniconcha* gastropods (Castel, 2022; Castel *et al.*, 2022). This was surprising because, given the fragmentation of vent habitats and the disconnected ridge system typical of the Western Pacific back-arc basins (Desbruyères *et al.*, 1994), we expected the geographic distribution of these gastropods (spanning over 4 000-kilometres) to have a predominant effect on their genetic differentiation. Although the volcanic arc associated with the Solomon islands (unsampled in this study) and the Woodlark basin may act as stepping stones for some species (Mitarai *et al.*, 2016; Boulart *et al.*, 2022), the large divide between hydrothermal vent fields located in the Western part of the study area (Manus basin) vs the Eastern part (Fiji and Lau basins, and Futuna arc) has acted as a barrier limiting exchanges between populations of *Alviniconcha* species (Breusing *et al.*, 2021) and other hydrothermal vent gastropods such as *Ifremeria nautilei* (Tran Lu Y, 2022; Tran Lu Y *et al.*, 2022), among many other species (Poitrimol *et al.*, 2022).

We discuss below how downstream analyses of sex and genetic variation have provided new insights into the sexual system and sex determination in the three species of hydrothermal *Alviniconcha* snails of the southwest Pacific, and why this genus provides an interesting opportunity to study the evolution of sex determination.

Genome-wide genetic structure provides clues to sexual systems

Anatomical observations combined with the effect of sex on genetic variation provided three new pieces of information for *A. boucheti*: this species is gonochoric, sex is genetically determined, and sex chromosomes are sufficiently differentiated to drive the first principal component of a genome-wide PCA (1% genetic variance explained, Fig. 3A), ahead of the effect of geography (0.7%, which is far less than observed for gastropod *I. nautilei* at the same spatial scale, Tran Lu Y *et al.*, 2022). This was confirmed by the measure of genome-wide genetic differentiation between males and females, regardless of their geographic origin (F_{ST} = 0.0105, $p < 0.001$, Table 1). In agreement with a

genetic sex determination system, the sex ratio appeared to be balanced when inferring the number of males and females using the coordinates of genotypes on PCA axis 1 (excluding the three mislabelled individuals: 104 males and 100 females, r = 0.51, exact binomial test p=0.83). The sex-ratio based on morphological or histological observations was balanced as well, but, as mentioned in the methods, the estimation could have resulted from a non-random selection of individuals for sexing.

We came to the same conclusions for *A. strummeri*: this species also has separate sexes and genetic sex determination. Yet in this case the effect of sex was second to that of geography (despite the distribution range being smaller than in *A. boucheti*, Figs. 1 and 3B), and differentiation between sexes was smaller (F_{ST} = 0.0024, p < 0.001, Table 1). This suggests that sex chromosomes are less differentiated in this species. However, further conclusions regarding the relative effects of sex and geography (i.e., local effective population size and gene flow) are hindered by the small sample size (22 ind. with known sex, 41 ind. overall), which limits the robustness of the results and may reflect not only a lower abundance but perhaps also a sampling bias linked with the selection of snail patches during collection (a possibility already mentioned in Castel *et al.*, 2022).

Most interestingly, the third species, *A. kojimai*, presented a strikingly different pattern. Close examination of histological cross-sections revealed that approximately 75% of the individuals that had been classified on board the ship as female based on external observations of the gonad actually had a mixture of female and male gonadal tissue (Fig. 2). In addition, the sex of individuals did not drive any obvious pattern of genetic differentiation on the multi-locus PCA, whether we considered two broad classes based on external and internal observations (i.e. males vs. females combined with morphological hermaphrodites, Fig. 3C) or three classes using internal observations only (males vs. females vs. morphological hermaphrodites, see supplementary information). Here geography was the main driver of the genome-wide genetic differentiation (1.2% genetic variance explained, Fig. 3C) but sex differentiation, although very low, still appeared to be statistically significant when opposing males to females combined with hermaphrodites (F_{ST} = 0.0002, p = 0.037). The sexual system of this species thus appeared to be different from the two others, potentially with a mixture of males, females, and

'morphological hermaphrodites' or 'intersex individuals' (see discussion below) and a weak signature of sex effects on genome-wide patterns of genetic structure.

490   *Sex is genetically determined through a XY system in the three species*

Although genome-wide estimates of genetic diversity gave some hints about sex-determination (significant differentiation between sexes in all species, H_S and H_0 always higher, and F_{IS} lower, in males than females, Table 1), locus-specific parameters were much more conclusive: the distribution of observed heterozygosity for each SNP in males and females pointed to a male-heterogametic genetic

495   system of sex determination for all three species.

The evidence was strongest for *A. boucheti*, where 553 SNPs on 237 RAD-loci were strictly homozygous in females while heterozygous in males (Fig. 4A), meaning that these SNPs have one allele fixed on the X chromosome and another allele fixed on the Y. The same situation was observed in *A. strummeri* (Fig. 4B), but with a smaller number of loci (92 SNPs on 55 RAD-loci) and much less

500   robustness due to the limited sample size. Most interestingly, while no such SNP (i.e. strictly heterozygous in males) was found in *A. kojimai*, nine markers located on seven RAD-loci showed observed heterozygosity values (strong in males, low in females) that were clearly outside the pattern defined by the other 70 113 SNPs (Fig. 4C) and, again, pointed towards a XY system. On the contrary, there was no evidence for female-heterogametic sex determination in any of the three species (see

505   e.g. Hearn *et al.*, 2022 for an example of ZW signature using a similar approach in Littorina gastropods).

The increase in genetic differentiation between homologous chromosomes relies on the reduction of recombination, or even its complete cessation in the case of loci that appear to be differentially fixed on the X and Y chromosomes (e.g. Berset-Brändli *et al.*, 2006; Brelsford *et al.*, 2017). Therefore, we conclude from the above observations that the loci that are strictly homozygous in

510   females and heterozygous in males are located on a non-recombining segment of XY sex chromosomes, at least in *A. boucheti* and *A. strummeri*, and that recombination is also strongly reduced in *A. kojimai* (albeit over a much smaller genomic region, see below).

In addition, the distribution of heterozygosity values in *A. boucheti* (Fig. 4A) suggested that additional loci were located on a sex-linked genomic region with little or no recombination as they had

515   retained some polymorphism on the Y chromosome (heterozygosity null in females, between 0.5 and 1 in males) or on X and Y (heterozygosity between 0 and 0.5 in females, between 0.5 and 1 in males).

It is also possible that the Y chromosomes of *A. boucheti* and *A. strummeri* contain repeated elements, which could create spurious RAD-loci identified as homologous sequences. Such repeated elements can accumulate mutations that would be wrongly identified as heterozygous SNPs. Tandem

520   replications of non-coding DNA, and in particular transposable elements can accumulate as a direct consequence of recombination arrest and may end up representing a large fraction of Y chromosomes (Beukeboom & Perrin, 2014; Charlesworth, 2021). Therefore, one should keep in mind that some of the SNPs that we identified has having one allele fixed on the X and another one fixed on the Y could in fact have the two alleles on distinct repeated sequences on the Y. Nonetheless, the conclusion

525   remains the same: these loci are a good indicator of XY genetic sex determination, but they should not be automatically seen as simple homologous markers present in a single copy (the results shown in figure 4 do in fact suggest that there are a number of autosomal paralogs, identified as loci with simultaneously high male and female heterozygosities).

While observed heterozygosity is an excellent way to identify sex determination systems, it is

530   not as efficient as among-sexes F_{ST} and within-sex F_{IS} to identify all the SNPs that are sex-linked. Using simulations to take into account the error associated with the estimation of locus-specific genetic parameters (in particular when based on a small number of individuals as in the case of *A. strummeri*), we found 1011 SNPS (373 RAD) sex-linked in *A. boucheti*, 111 (64 RAD) in *A. strummeri*, and 16 (14 RAD) in *A. kojimai* (Fig. 5). These numbers are not easily comparable, as *A. strummeri* had both far

535   fewer individuals for which sex was determined and fewer SNPs than the two other species (meaning much less statistical power). Yet *A. kojimai* did not suffer from these limitations (151 sexed individuals genotyped at 70 122 SNPs), and locus-specific genetic estimates clearly identified outlier patterns associated with sex-linkage (Figs. 4C and 5E-F). Importantly, the analyses comparing males vs. females

+ morphological hermaphrodites were confirmed by the analyses restricted to those individuals for which sex was determined through histological observations: sex-linked outlier loci were detected when comparing males *vs* females or males vs morphological hermaphrodites, but not when comparing females vs hermaphrodites (Figs. S6 and S7). Females and hermaphrodites therefore form a homogeneous group. Altogether, these results indicate that sex is at least partly determined genetically in *A. kojimai*, with XY individuals developing into males and XX individuals developing into females or morphological hermaphrodites. Our observations also suggest the main sex-determining gene is contained in a non-recombining region that is smaller than that of *A. boucheti* and probably *A. strummeri*.

A unique pair of XY chromosomes across the three species?

To optimize the efficiency of the genotyping, RAD-sequences were treated independently for each species, meaning that we used three distinct catalogs of loci. Reciprocal blasts of sex-linked loci for each pair of species revealed that three RAD-loci were found to be sex-linked in all three species, and 25 more were identified as sex-linked in two species (Fig. 6A). In addition, all 14 *A. kojimai* sex-linked loci mapped on five scaffolds of the *A. marisindica* reference genome that were also found to include sex-linked loci in the two other species (Fig. 6B). These results demonstrate that the three species share the same sex-linked genomic region, which contains or is close to the region that triggers sex determination in the three species, on the same pair of XY chromosomes.

The size of the sex-linked region can be roughly estimated by looking at the two species for which we had most data (not considering *A. strummeri* here because we probably had limited power to detect most sex-linked markers). In *A. boucheti*, 0.7% of the SNPs were identified as sex-linked, while this figure dropped to 0.025% in *A. kojimai*. The total genome size was estimated for two *Alviniconcha* species: 0.92 Gb for an unidentified species from our study area (formerly called *A. hessleri* before taxonomic revision) estimated by flow cytometry (Bonnivard *et al.*, 2009), and 0.83 Gb for the genome assembly ASM1885773v1 of *A. marisindica* (Yang *et al.*, 2022). If *A. boucheti* and *A. kojimai* have a

565   genome of about 0.9 Gb then 0.7% and 0.025% of this genome represent sex-linked regions of 6.3 Mb

for *A. boucheti* and 2.25 Mb for *A. kojimai*, respectively. A comparable estimate was obtained for *A.*

kojimai by summing the lengths of the reference genome scaffolds found to harbor sex-linked loci

(2.82 Mb, Fig. 6B), but the value obtained for *A. boucheti* was much larger (24.9 Mb).

These values should be interpreted as rough estimates because they depend on sample size

570   (number of sexed individuals), methodology (conservative threshold set by our simulations to decide

whether a locus is sex-linked), and XY polymorphism. Also, they do not consider the potential genome

sequences present only on the X or only on the Y (not investigated in this study, see Gautier, 2014;

Feron *et al.*, 2021; Garcia-Raventós *et al.*, 2023). Moreover, further analyses are needed to reveal i) if

repeated elements are involved in the length of the Y chromosome, and ii) the position of sex-linked

575   RAD-loci within each scaffold. A precise estimation of the size and synteny of sex-linked genomic

regions in the different *Alviniconcha* species will require additional full genome sequences.

What is the sexual system of Alviniconcha *kojimai, and how is sex determined in this species?*

All our observations point towards a XY system of sex determination in *A. kojimai*, with XX

580   individuals developing a mixture of male and female tissues in variable proportion. We have yet to

determine if the mixed gonadal tissues allow these individuals to reproduce using both sex functions,

and, if so, simultaneously or not, and whether selfing is possible. Here we analysed a single gonad cross

section per individual, and further investigations are therefore needed to quantify these proportions

precisely, and in particular, determine if some females that do not show any male tissue (11 such

585   individuals identified here using a single gonad section) are indeed devoid of male tissue. All this

information is necessary to understand the current sexual system of *A. kojimai*, and to understand

whether it may be stable or represent a transitional stage towards gonochorism or hermaphroditism.

The presence of male tissue within hermaphrodites would be only beneficial if the sperm is used for

selfing or cross-interbreeding via sperm storage when the number of individuals is restricted to found

590   a new population: a situation possibly met when geo-tectonic rearrangements or eruptive events

destroy most of the vent communities in a given region (see the recent eruption of the submarine volcano Hunga Tonga – Hunga Ha'apai in December 2021, Gupta *et al.*, 2022). In a recent study of *Gigantidas haimaensis* mussels from cold seep ecosystems in the South China Sea, Shi *et al.* (2022) found that male gonads contained oocytes (while female gonads did not contain spermatocytes).

595   Combining proteomic and transcriptomic analyses with their histological observations, the authors hypothesized that these males were intersex individuals undergoing sex change, a strategy observed in other Bathymodiolinae (Laming *et al.*, 2018) possibly favoured by the ephemeral and patchy distribution of deep sea vent and cold seep habitats.

        Although the ultimate forces that drive transitions between gonochorism and hermaphroditism

600   in animals are far from fully resolved, selection may arise from the costs and benefits of factors such as reproductive assurance, inbreeding, sexual specialization, and plasticity in sex allocation (Weeks, 2012; Leonard, 2018; Pannell & Jordan, 2022). Mixed sexual systems characterized by the co-existence of females and hermaphrodite individuals (gynodioecy) or males and hermaphrodites (androdioecy) are particularly relevant to understand the evolution of sexual systems, and may in some cases

605   represent intermediate stages between hermaphroditism and gonochorism (Charlesworth, 2006; Pannell & Jordan, 2022). However, there are in fact very few species with such mixed systems: Weeks (2012) listed only 115 androdioecious and up to 9 gynodioecious animals. In these species, fully functional hermaphrodites effectively gain reproductive success through both their male and female functions (functional gyno/androdioecy *sensus* Charlesworth, 1984 and Pannell, 2002). Surprisingly, it

610   seems that many more gonochoristic species appear to contain individuals that have both male and female reproductive traits, but with only one type being functional (i.e. a single type of functional gamete produced and used for reproduction). In the perspective of the evolution of sexual systems, this condition has been discussed mostly in the botanical literature (where it is called cryptic dioecy, e.g. Charlesworth, 1984; Mayer & Charlesworth, 1991). In the animal biology literature, this topic has

615   been more intensively examined in the perspective of the anthropogenic stressors possibly causing the development of reproductive tissue typically seen as abnormal (reviewed in Bahamonde *et al.*,

2013; Grilo & Rosa, 2017). Some gonochoristic animal species appear to contain individuals called intersex that present both male and female traits. This is particularly frequent in fish and molluscs, where intersex refers more specifically to "individuals possessing oocytes or distinct stages of spermatogonia, at varying degrees of development, within the normal gonad of the opposite gender (i.e. spermatocytes in the ovary or oocytes in the testis)" (Grilo & Rosa, 2017). Intersex was shown to be exacerbated by environmental contaminants that can disrupt sex differentiation, and thus should be detrimental for the species, but exogenous factors may not be necessarily required, and the frequency of "naturally" occurring intersex individuals remains an open question. In all cases, these species are in fact functionally gonochoristic, but they are interesting as they may nonetheless represent intermediate steps in transitions between fully functional sexual systems (e.g. in angiosperms, Mayer & Charlesworth, 1991). Here, the evolutionary success of *A. kojimai* in colonizing nearly all hydrothermal vents of four back-arc basins and the volcanic arc of Wallis & Futuna despite a high ratio of morphologically hermaphroditic individuals found in every population should indicate that such a transitional stage has been favoured for quite a long time.

Gonochorism is thought to be the ancestral sexual system of gastropods (Collin, 2013) but molluscs have very diversified sexual systems relatively to other major animal clades, with much flexibility in sex determination and sex differentiation (Collin, 2013; Leonard, 2018). Collin (2013) mentions in particular that intersex individuals were observed in many molluscan taxa, sometimes in non-negligible numbers, and that these species deserve further investigations to ascertain their sexual system. Extending the analyses presented here to sexual anatomy and sex-linked genomic patterns of the six species of *Alviniconcha* gastropods (Johnson *et al.*, 2015) should prove interesting with respect to the evolution of sexual systems and sex determination.

Acknowledgements

We thank the crew of the RV L'Atalante and the team of the ROV Victor6000 who participated in the Chubacarc 2019 expedition. We thank P-A. Gagnaire, who suggested that we look at the sex of

individuals to understand the observed genetic structure, and we are grateful to Mathieu Gautier for insightful discussions. This work benefited from access to the Genomer molecular biology platform, member of BioGenOuest and EMBRC-France, and the bioinformatics platform ABiMS of the Station Biologique de Roscoff. This study was supported by the French Oceanographic Fleet program and the ANR CERBERUS (ANR-17-CE02-0003 to S. H.).

Data availability

ddRAD-sequences and associated metadata are available on NCBI within the BioProject PRJNA768636. Additional data are deposited on zenodo.org (DOI 10.5281/zenodo.7817088) and will be publicly available after peer-review.

Literature

Bahamonde, P.A., Munkittrick, K.R. & Martyniuk, C.J. 2013. Intersex in teleost fish: Are we distinguishing endocrine disruption from natural phenomena? *General and Comparative Endocrinology* **192**: 25–35.

Berset-Brändli, L., Jaquiéry, J., Dubey, S. & Perrin, N. 2006. A sex-specific marker reveals male heterogamety in European tree frogs. *Molecular Biology and Evolution* **23**: 1104–1106.

Beukeboom, L.W. & Perrin, N. 2014. *The Evolution of Sex Determination*. Oxford University Press.

Bonnivard, E., Catrice, O., Ravaux, J., Brown, S.C. & Higuet, D. 2009. Survey of genome size in 28 hydrothermal vent species covering 10 families. *Genome* **52**: 524–536.

Boulart, C., Rouxel, O., Scalabrin, C., Le Meur, P., Pelleter, E., Poitrimol, C., *et al.* 2022. Active hydrothermal vents in the Woodlark Basin may act as dispersing centres for hydrothermal fauna. *Commun Earth Environ* **3**: 1–16. Nature Publishing Group.

Brelsford, A., Lavanchy, G., Sermier, R., Rausch, A. & Perrin, N. 2017. Identifying homomorphic sex chromosomes from wild-caught adults with limited genomic resources. *Molecular Ecology*

Resources **17**: 752–759.

670   Breusing, C., Johnson, S.B., Mitarai, S., Beinart, R.A. & Tunnicliffe, V. 2021. Differential patterns of connectivity in Western Pacific hydrothermal vent metapopulations: A comparison of biophysical and genetic models. *Evolutionary Applications* **n/a**.

Breusing, C., Johnson, S.B., Tunnicliffe, V., Clague, D.A., Vrijenhoek, R.C. & Beinart, R.A. 2020. Allopatric and Sympatric Drivers of Speciation in Alviniconcha Hydrothermal Vent Snails. *Mol Biol Evol*,
675   doi: 10.1093/molbev/msaa177.

Castel, J. 2022. Rôle de l'hétérogénéité de l'environnement hydrothermal profond sur la distribution et les mécanismes de spéciation du complexe de gastéropodes Alviniconcha spp. Sorbonne Université.

Castel, J., Hourdez, S., Pradillon, F., Daguin-Thiébaut, C., Ballenghien, M., Ruault, S., *et al.* 2022.
680   Inter-Specific Genetic Exchange Despite Strong Divergence in Deep-Sea Hydrothermal Vent Gastropods of the Genus Alviniconcha. *Genes* **13**: 985. Multidisciplinary Digital Publishing Institute.

Catchen, J., Hohenlohe, P.A., Bassham, S., Amores, A. & Cresko, W.A. 2013. Stacks: an analysis tool set for population genomics. *Molecular Ecology* **22**: 3124–3140.

Catchen, J.M., Amores, A., Hohenlohe, P., Cresko, W. & Postlethwait, J.H. 2011. Stacks: Building
685   and Genotyping Loci De Novo From Short-Read Sequences. *G3 (Bethesda)* **1**: 171–182.

Cavanaugh, C.M. 1983. Symbiotic chemoautotrophic bacteria in marine invertebrates from sulphide-rich habitats. *Nature* **302**: 58–61. Nature Publishing Group.

Charlesworth, D. 1984. Androdioecy and the evolution of dioecy. *Biological Journal of the Linnean Society* **22**: 333–348.

690   Charlesworth, D. 2006. Evolution of Plant Breeding Systems. *Current Biology* **16**: R726–R735.

Charlesworth, D. 2021. When and how do sex-linked regions become sex chromosomes? *Evolution* **75**: 569–581.

Childress, J.J. 1995. Life in Sulfidic Environments: Historical Perspective and Current Research Trends1. *American Zoologist* **35**: 83–90.

695   Coelho, S.M. & Umen, J. 2021. Switching it up: algal insights into sexual transitions. *Plant Reprod* **34**: 287–296.

Collin, R. 2013. Phylogenetic Patterns and Phenotypic Plasticity of Molluscan Sexual Systems. *Integrative and Comparative Biology* **53**: 723–735.

Daguin-Thiébaut, C., Ruault, S., Roby, C., Broquet, T., Viard, F. & Brelsford, A. 2021. Construction
700   of individual ddRAD libraries. *protocols.io*, doi: 10.17504/protocols.io.bv4tn8wn.

Desbruyères, D., Alayse-Danet, A.-M., Ohta, S., & the Scientific Parties of biolauand starmerCruises. 1994. Deep-sea hydrothermal communities in Southwestern Pacific back-arc basins (the North Fiji and Lau Basins): Composition, microdistribution and food web. *Marine Geology* **116**: 227–242.

705   Doyle, J.J. & Dickson, E.E. 1987. Preservation of Plant Samples for Dna Restriction Endonuclease Analysis. *TAXON* **36**: 715–722.

Faure, B., Chevaldonné, P., Pradillon, F., Thiébaut, E. & Jollivet, D. 2007. Spatial and temporal dynamics of reproduction and settlement in the Pompeii worm Alvinella pompejana (Polychaeta: Alvinellidae). *Marine Ecology Progress Series* **348**: 197–211.

710   Feron, R., Pan, Q., Wen, M., Imarazene, B., Jouanno, E., Anderson, J., *et al.* 2021. RADSex: A computational workflow to study sex determination using restriction site-associated DNA sequencing data. *Molecular Ecology Resources* **21**: 1715–1731.

Garcia-Raventós, A., Muñoz-Merida, A., Lapiedra, O., Unzeta, M., Ferrandiz-Rovira, M. & Sol, D. 2023. *Identification of sex-linked SNP markers in wild populations of monomorphic birds*. Preprints.

715   Gautier, M. 2014. Using genotyping data to assign markers to their chromosome type and to infer the sex of individuals: a Bayesian model-based classifier. *Molecular Ecology Resources* **14**: 1141–1159.

Goudet, J. & Jombart, T. 2021. hierfstat: Estimation and Tests of Hierarchical F-Statistics.

Grilo, T.F. & Rosa, R. 2017. Intersexuality in aquatic invertebrates: Prevalence and causes.
720   *Science of The Total Environment* **592**: 714–728.

Gupta, A.K., Bennartz, R., Fauria, K.E. & Mittal, T. 2022. Eruption chronology of the December 2021 to January 2022 Hunga Tonga-Hunga Ha'apai eruption sequence. *Commun Earth Environ* **3**: 1–10. Nature Publishing Group.

Hearn, K.E., Koch, E.L., Stankowski, S., Butlin, R.K., Faria, R., Johannesson, K., *et al.* 2022. Differing associations between sex determination and sex-linked inversions in two ecotypes of Littorina saxatilis. *Evolution Letters* **6**: 358–374.

Hill, R.C., Egydio de Carvalho, C., Salogiannis, J., Schlager, B., Pilgrim, D. & Haag, E.S. 2006. Genetic Flexibility in the Convergent Evolution of Hermaphroditism in Caenorhabditis Nematodes. *Developmental Cell* **10**: 531–538.

Hourdez, S. & Jollivet, D. 2019. CHUBACARC cruise,L'Atalante R/V. , doi: 10.17600/18001111. Sismer.

Hourdez, S. & Lallier, F.H. 2007. Adaptations to hypoxia in hydrothermal-vent and cold-seep invertebrates. *Rev Environ Sci Biotechnol* **6**: 143–159.

Jarne, P. & Auld, J.R. 2006. Animals Mix It up Too: The Distribution of Self-Fertilization Among Hermaphroditic Animals. *Evolution* **60**: 1816–1824.

Johnson, S.B., Waren, A., Tunnicliffe, V., Van Dover, C., Wheat, C.G., Schultz, T.F., *et al.* 2015. Molecular taxonomy and naming of five cryptic species of Alviniconcha snails (Gastropoda: Abyssochrysoidea) from hydrothermal vents. *Syst Biodivers* **13**: 278–295.

Jollivet, D., Empis, A., Baker, M.C., Hourdez, S., Comtet, T., Jouin-Toulmond, C., *et al.* 2000. Reproductive biology, sexual dimorphism, and population structure of the deep sea hydrothermal vent scale-worm, *Branchipolynoe seepensis* (Polychaeta: Polynoidae). *J. Mar. Biol. Ass.* **80**: 55–68.

Jolly, M., Viard, F., Weinmayr, G., Gentil, F., Thiébaut, E. & Jollivet, D. 2003. Does the genetic structure of Pectinaria koreni (Polychaeta: Pectinariidae) conform to a source–sink metapopulation model at the scale of the Baie de Seine? *Helgol Mar Res* **56**: 238–246.

Jombart, T. 2008. adegenet: a R package for the multivariate analysis of genetic markers. *Bioinformatics* **24**: 1403–1405.

Kearse, M., Moir, R., Wilson, A., Stones-Havas, S., Cheung, M., Sturrock, S., *et al.* 2012. Geneious Basic: An integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* **28**: 1647–1649.

750 Kiontke, K., Gavin, N.P., Raynes, Y., Roehrig, C., Piano, F. & Fitch, D.H.A. 2004. Caenorhabditis phylogeny predicts convergence of hermaphroditism and extensive intron loss. *Proceedings of the National Academy of Sciences* **101**: 9003–9008. Proceedings of the National Academy of Sciences.

Laming, S.R., Gaudron, S.M. & Duperron, S. 2018. Lifecycle Ecology of Deep-Sea Chemosymbiotic Mussels: A Review. *Frontiers in Marine Science* **5**.

755 Leonard, J.L. 2018. The Evolution of Sexual Systems in Animals. In: *Transitions Between Sexual Systems: Understanding the Mechanisms of, and Pathways Between, Dioecy, Hermaphroditism and Other Sexual Systems* (J. L. Leonard, ed), pp. 1–58. Springer International Publishing, Cham.

Lutz, R.A. 1988. Dispersal of organisms at deep-sea hydrothermal vents: A review. *Oceanologica Acta, Special issue*. Gauthier-Villars.

760 Mayer, S.S. & Charlesworth, D. 1991. Cryptic dioecy in flowering plants. *Trends in Ecology & Evolution* **6**: 320–325.

Mitarai, S., Watanabe, H., Nakajima, Y., Shchepetkin, A.F. & McWilliams, J.C. 2016. Quantifying dispersal from hydrothermal vent fields in the western Pacific Ocean. *Proceedings of the National Academy of Sciences* **113**: 2976–2981. Proceedings of the National Academy of Sciences.

765 Nei, M. 1987. *Molecular Evolutionary Genetics*. Columbia University Press.

Pannell, J.R. 2002. What Is Functional Androdioecy? *Functional Ecology* **16**: 862–865. [British Ecological Society, Wiley].

Pannell, J.R. & Jordan, C.Y. 2022. Evolutionary Transitions Between Hermaphroditism and Dioecy in Animals and Plants. *Annu. Rev. Ecol. Evol. Syst.* **53**: annurev-ecolsys-102320-085812.

770 Papot, C., Massol, F., Jollivet, D. & Tasiemski, A. 2017. Antagonistic evolution of an antibiotic and its molecular chaperone: how to maintain a vital ectosymbiosis in a highly fluctuating habitat. *Sci Rep* **7**: 1454.

Pembleton, L.W., Cogan, N.O.I. & Forster, J.W. 2013. StAMPP: an R package for calculation of genetic differentiation and structure of mixed-ploidy level populations. *Molecular Ecology Resources* **13**: 946–952.

Podowski, E.L., Ma, S., Iii, G.W.L., Wardrop, D. & Fisher, C.R. 2010. Biotic and abiotic factors affecting distributions of megafauna in diffuse flow on andesite and basalt along the Eastern Lau Spreading Center, Tonga. *Marine Ecology Progress Series* **418**: 25–45.

Poitrimol, C., Thiébaut, É., Daguin-Thiébaut, C., Port, A.-S.L., Ballenghien, M., Y, A.T.L., *et al.* 2022. Contrasted phylogeographic patterns of hydrothermal vent gastropods along South West Pacific: Woodlark Basin, a possible contact zone and/or stepping-stone. *PLOS ONE* **17**: e0275638. Public Library of Science.

Powell, M.A. & Somero, G.N. 1986. Adaptations to sulfide by hydrothermal vent animals: sites and mechanisms of detoxification and metabolism. *The Biological Bulletin* **171**: 274–290. The University of Chicago Press.

Pyne, C.B., McFarlane, S.E. & Mandeville, E.G. 2022. Identification of sex-determining loci in hybridizing Catostomus fish species. bioRxiv.

R Core Team. 2021. R: A language and environment for statistical computing. R Foundation for Statistical Computing., Vienna, Austria.

Renner, S.S. 2014. The relative and absolute frequencies of angiosperm sexual systems: Dioecy, monoecy, gynodioecy, and an updated online database. *American Journal of Botany* **101**: 1588–1596.

Shi, Y., Yao, G., Zhang, H., Jia, H., Xiong, P. & He, M. 2022. Proteome and Transcriptome Analysis of Gonads Reveals Intersex in Gigantidas haimaensis. *BMC Genomics* **23**: 174.

Stein, J.L., Cary, S.C., Hessler, R.R., Vetter, R.D., Felbeck, H., Ohta, S., *et al.* 1988. Chemoautotrophic Symbiosis in a Hydrothermal Vent Gastropod. *The Biological Bulletin* **174**: 373–378. The University of Chicago Press.

Stovall, W.R., Taylor, H.R., Black, M., Grosser, S., Rutherford, K. & Gemmell, N.J. 2018. Genetic sex assignment in wild populations using genotyping-by-sequencing data: A statistical threshold

approach. *Mol Ecol Resour* **18**: 179–190.

Tran Lu Y, A. 2022. La phylogéographie comparée d'espèces hydrothermales du Pacifique Ouest à l'heure de la génomique des populations. Université Montpellier.

Tran Lu Y, A., Ruault, S., Daguin-Thiébaut, C., Castel, J., Bierne, N., Broquet, T., *et al.* 2022. Subtle limits to connectivity revealed by outlier loci within two divergent metapopulations of the deep-sea hydrothermal gastropod Ifremeria nautilei. *Molecular Ecology* **31**: 2796–2813.

Tyler, P.A. & Young, C.M. 1999. Reproduction and dispersal at vents and cold seeps. *Journal of the Marine Biological Association of the United Kingdom* **79**: 193–208. Cambridge University Press.

Vrijenhoek, R.C. 2010. Genetic diversity and connectivity of deep-sea hydrothermal vent metapopulations. *Molecular Ecology* **19**: 4391–4411.

Wang, Y., Gasser, R.B., Charlesworth, D. & Zhou, Q. 2022. Evolution of sexual systems, sex chromosomes and sex-linked gene transcription in flatworms and roundworms. *Nat Commun* **13**: 3239. Nature Publishing Group.

Weeks, S.C. 2012. The Role of Androdioecy and Gynodioecy in Mediating Evolutionary Transitions Between Dioecy and Hermaphroditism in the Animalia. *Evolution* **66**: 3670–3686.

Weir, B.S. & Cockerham, C.C. 1984. Estimating F-statistics for the analysis of population structure. *Evolution* **38**: 1358–1370.

Wood, D.E. & Salzberg, S.L. 2014. Kraken: ultrafast metagenomic sequence classification using exact alignments. *Genome Biology* **15**: R46.

Yang, Y., Sun, J., Chen, C., Zhou, Y., Van Dover, C.L., Wang, C., *et al.* 2022. Metagenomic and metatranscriptomic analyses reveal minor-yet-crucial roles of gut microbiome in deep-sea hydrothermal vent snail. *Animal Microbiome* **4**: 3.