Accurate prediction of in vivo protein abundances by coupling constraint-based modelling

and machine learning

Maurício Alexander de Moura Ferreira[1], Philipp Wendering[2,3], Marius Arend[2,3], Wendel Batista da

Silveira[1], Zoran Nikoloski[2,3,*]

1 Department of Microbiology, Federal University of Viçosa, Viçosa, Minas Gerais, 36570900,

Brazil

2 Bioinformatics, Institute of Biochemistry and Biology, University of Potsdam, Potsdam, 14476,

Germany

3 Systems Biology and Mathematical Modelling, Max Planck Institute of Molecular Plant

Physiology, Potsdam, 14476, Germany

* Corresponding author**:** zoran.nikoloski@uni-potsdam.de

Abstract

Quantification of how different environmental cues affect protein allocation can provide important

insights for understanding cell physiology. While absolute quantification of proteins can be

obtained by resource-intensive mass-spectrometry-based technologies, prediction of protein

abundances offers another way to obtain insights into protein allocation. Here we present

CAMEL, a framework that couples constraint-based modelling with machine learning to predict

protein abundance for any environmental condition. This is achieved by building machine learning

models that leverage static features, derived from protein sequences, and condition-dependent

features predicted from protein-constrained metabolic models. Our findings demonstrate that

CAMEL results in excellent prediction of protein allocation in *E. coli* (average Pearson correlation

of at least 0.9), and moderate performance in *S. cerevisiae* (average Pearson correlation of at

least 0.5). Therefore, CAMEL outperformed contending approaches without using molecular

read-outs from unseen conditions and provides a valuable tool for using protein allocation in

biotechnological applications.

Keywords: Protein allocation, Environmental effects, Multi-model framework

Introduction

Proteomics technologies facilitate system-wide profiling of the identity as well as changes in

abundance, distribution, modifications, and interactions of proteins that drive different cellular

processes (Schubert et al., 2017). As cells are dynamic systems, these characteristics of the

proteome change in response to different environmental cues to facilitate the system's adaptation

to different physiological states (Liu et al., 2019; Nielsen, 2019). As a result, high-throughput

quantitative proteomics technologies, based on mass spectrometry (MS) techniques, have

provided valuable information for various biotechnological and medical applications (Kültz, 2020;

Lill et al., 2021). However, the measurement of absolute protein abundance still poses great

challenges, due to physicochemical properties that interfere with ionization efficiency in mass

spectrometry (Otto et al., 2014; Pappireddi et al., 2019) and the lack of a standardized approach,

affecting the reproducibility of resulting data (Calderón-Celis et al., 2018).

Another approach for protein quantification relies on predicting protein abundances by

using a variety of machine learning approaches using features that are more facile to measure or

quantify. For instance, with gene expression data as predictors, Torres-García et al. (2009)

trained gradient-boosted trees to predict protein abundances in *Desulfovibrio vulgaris*, resulting in

prediction accuracies, quantified by the coefficient of determination, R^2, ranging from 0.39 to 0.58.

Similarly, Mehdi et al. (2014) used Bayesian networks to predict protein abundance for

Saccharomyces cerevisiae and *Schizosaccharomyces pombe* using gene expression data,

yielding Pearson's correlation coefficients ranging from 0.50 to 0.71. Mergner et al. (2020) used

stepwise and LASSO regression models to predict protein abundances in *Arabidopsis thaliana*

based on gene expression as well as sequence-derived features, achieving Pearson correlation

in the range from 0.61 to 0.79 across different tissues. Codon usage metrics have also been used

to predict protein abundance in *Saccharomyces cerevisiae*, with the best model achieving an R^2

value of 0.74 (Ferreira et al., 2021). In addition, structural features of mRNA molecules have been

used to predict protein abundances in *Escherichia coli*, achieving a Spearman correlation

coefficient of 0.71. However, these machine learning models have been developed using

quantitative proteomics and transcriptomics data from optimal growth conditions and their

performance in sub-optimal growth conditions remains unexplored. In addition, given the dynamic

nature of the proteome, it is expected that the predictions across different conditions from

machine learning models based on environment-invariant features (e.g., codon usage and

structure-related), are poor.

The advent of protein-constrained genome-scale metabolic models (pcGEMs) (Beg et al.,

2007) has facilitated not only the prediction of protein abundance but also the usage of

proteomics data to predict metabolic and physiological traits (Bekiaris and Klamt, 2020;

Domenzain et al., 2022; Sánchez et al., 2017). This is achieved by parameterizing pcGEMs with

data on enzyme turnover numbers, providing the basis to link steady-state fluxes with enzyme

abundances. For instance, a pcGEM of *E. coli* was used by Adadi et al. (2012) to predict protein

abundance, achieving Pearson correlation coefficient of 0.84 with gene expression data of *E. coli*

grown on a glucose minimal medium. However, given that comparisons were made with gene

expression data, it remains unexplored how these predictions fare against measured proteomics

data. Further, Heckmann et al. (2018) trained machine learning approaches to predict enzyme

catalytic efficiency, namely *in vitro* (k_{cat}) and *in vivo* values (k_{max}^{vivo}), that were later used to predict

protein abundances with the approach of Adadi et al. (2012). Comparing the predicted protein

abundances to experimental data from Schmidt et al. (2016), they found that using k_{max}^{vivo} values

resulted in a 43% lower root mean squared error (RMSE) in comparison to k_{cat} values. However,

when comparing the prediction to measured proteomics data, the obtained R^2 values were poor

to modest, ranging from 0.11 to 0.68. As a result, we still lack an understanding of the factors that

affect the quality of the protein abundance predictions. Moreover, like the machine learning

approaches above, the existing predictions of protein abundance based on pcGEMs have only

been assessed under conditions that achieve maximal measured specific growth rate for the wild-

type strain that we refer to as optimal. Therefore, the problem of predicting protein abundance

under sub-optimal, stress-related conditions, that result in smaller specific growth rate than the

optimal, is particularly difficult since many mechanisms affecting protein abundance (e.g., codon

usage (Novoa et al., 2019), RNA levels (Eraslan et al., 2019), protein interactions (Mergner et al.,

2020), 3' or 5' UTR motifs (Terai and Asai, 2020)) are not included in pcGEMs.

To address these issues, we coupled constraint-based modelling with machine learning

to predict *in vivo* protein abundances, leading to a multi-model framework termed CAMEL

(Coupled Approach of MEtabolic modelling and machine Learning) that leverages the strengths of

both approaches. First, CAMEL predicts the protein abundance in different conditions using a

constraint-based approach. Next, these protein abundance predictions are employed together

with experimentally measured protein abundances to calculate protein reserve ratios. CAMEL

then trains machine learning (ML) models to predict the protein reserve ratios using different

static features, derived from the protein sequence, and condition-dependent features, obtained

from the pcGEM. Lastly, the predicted protein reserve ratios along with the predicted protein

abundances from the pcGEM are used to calculate *in vivo* protein abundances. Once trained, the

models of CAMEL can be used to make predictions by using features predicted by the pcGEM

instead of data on molecular read-outs from unseen conditions. Our results showed that the

CAMEL approach outperformed contending methods on data from two model organisms, *E. coli*

and *S. cerevisiae*, highlighting the advantages of a coupling of constraint-based modelling and

machine learning approaches. In addition, these applications of CAMEL point to particular

proteins and metabolic pathways for which protein abundance is difficult to predict, requiring the

consideration of additional features to improve prediction performance in future developments.

Material and Methods

Data set construction

We obtained the experimental enzyme abundances for *E. coli* over 31 growth conditions from

three different studies (Peebo et al., 2015; Schmidt et al., 2016; Valgepea et al., 2013), as

reported in Davidi et al. (2016). The growth conditions in these experiments range from

alternative carbon substrates (e.g., acetate, glucosamine, glucose, glycerol, mannose, pyruvate,

and xylose) to chemostats with increasing dilution rates, with glucose as a carbon source. For *S.*

cerevisiae, we used the enzyme abundance values from the data set of Chen and Nielsen (2021),

which contains quantitative proteomics measurements for 30 growth conditions obtained from

four studies (Chen and Nielsen, 2021; Di Bartolomeo et al., 2020; Lahtvee et al., 2017; Yu et al.,

2020), including: optimal batch growth with glucose as carbon source, nitrogen- and glucose-

limited chemostats with increasing dilution rates, stress states achieved with high ethanol and

osmolarity, and reduced nutrient availability.

Condition-dependent features are predicted from the pcGEMs and include the enzyme

usage and the metabolic fluxes associated with each enzyme i. These features are obtained by

solving a linear problem to minimize excess enzyme usage (Eqs. (1)-(8)), detailed below. To

predict the enzyme usage distributions, we used the pcGEMs eciML1515 (Domenzain et al.,

2022) and ecYeast8 (Lu et al., 2019), from *E. coli* and *S. cerevisiae*, respectively, both without

integrated enzyme measurements (batch model). To this end, we minimized excess enzyme

usage. Maximal enzyme usage is linked to efficient usage of resources, which has been shown to

result in better estimates of k_{cat} values over other approaches that focus on minimization of total

flux (Xu et al., 2021). This objective function also ensures that cellular burden is alleviated, which

can lead to proteotoxic effects for the cell (Kintaka et al., 2020). We solve the following linear

programming (LP) problem, termed LP1:

132
$$\min_{\mathbf{v,E}} E_{s,tot} - \sum_{i=1}^{p} MW_i \cdot E_{s,i} \qquad (1)$$

133
$$s.\ t.$$

134
$$\mathbf{N} \cdot \mathbf{v} = \mathbf{0} \qquad (2)$$

135
$$v_{min,j} \le v_j \le v_{max,j}, 1 \le j \le r \qquad (3)$$

136
$$v_{uptake,s} = v_{uptake,s}^{exp} \quad (4)$$

137
$$v_{secretion,s} = v_{secretion,s}^{exp} \quad (5)$$

138
$$v_{bio} = \mu_s \qquad (6)$$

139
$$v_j \le k_{cat}^{ij} \cdot E_{s,i}, 1 \le j \le r, 1 \le i \le p \quad (7)$$

140
$$E_{s,tot} = P_{s,tot}^{exp} \cdot f \cdot \sigma \quad (8)$$

where $E_{s,i}$ is the enzyme i, ranging from 1 to the number of proteins p in the model ($1 \le i \le p$), in

condition s, MW_i is the molecular weight of an enzyme i, $\mathbf{N}$ is the stoichiometric matrix, $\mathbf{v}$ is the

flux distribution vector, v_j is the metabolic flux through reaction j, ranging from 1 to the number of

reactions r in the model ($1 \le j \le r$), v_{bio} is flux through the biomass reaction, and μ is the specific

growth rate, E_{tot} is the total enzyme content, P_{tot}^{exp} is the total protein content, f is the mass

fraction of all measured proteins included in the model, and σ is the average *in vivo* enzyme

saturation, assumed to be a specific value of 0.5 for conditions in which this parameter is not

available (Domenzain et al., 2022; Sánchez et al., 2017). The constraints applied to the problem

are based on the available data about specific growth rates, secretion, and nutrient uptake rates,

protein usage and total protein content.

For both organisms, we constrained the models given the measurements of nutrient

uptake rates and specific growth rates using data from Davidi et al. (2016) and Chen and Nielsen

(2021). We excluded the experimental conditions that lacked physiological data (e.g., nutrient

uptake rates, specific growth rates, or protein content). In addition, we opted to exclude the

temperature stress conditions from Lahtvee et al. (2017), as temperature stress triggers

responses that entail cellular changes that can impact the function of enzymes (Li et al., 2021)

and require more tailored modelling approaches (Wendering et al., 2023). This filtering resulted in

data from 20 conditions for *E. coli*, and 19 conditions for *S. cerevisiae*. For conditions for which

the models were overconstrained, we relaxed the constraints on the uptake and secretion rates

by increments of 1% until the measured specific growth rate could be achieved.

We were also interested in whether using pcGEMs integrated with *in vivo* catalytic rates

(k_{max}^{vivo}) instead of *in vitro* values (k_{cat}) would impact the performance of the ML models. To this

end, we exchanged the integrated k_{cat} values with the k_{max}^{vivo} values, obtained from the compilation

by (Wendering et al., 2023) of estimated catalytic rates using pFBA from the Davidi et al. (2016)

and Chen and Nielsen (2021) studies. For both *E. coli* and *S. cerevisiae*, we concatenated the

predictions generated from the considered growth conditions into a single data set for each

organism. The resulting data set for *E. coli* included 2256 enzymes for the pcGEM using k_{cat}

values, and a data set with 2246 enzymes for the pcGEM k_{max}^{vivo} values. For *S. cerevisiae*, the

resulting data set included 4596 enzymes for the pcGEM using k_{cat} values, and a data set with

4590 enzymes for the pcGEM k_{max}^{vivo} values.

Training and assessment of machine learning models

To train the machine learning models, we used the Tree-based Pipeline Optimization Tool

(TPOT), an automated machine learning tool that optimizes machine learning pipelines to predict

the target variable using genetic programming (Le et al., 2020). We configured TPOT for

regression problems since we aimed to obtain quantitative predictions of the protein reserve ratio,

ϕ_i, for each enzyme $i, 1 \leq i \leq p$. For all conditions in each species, we iterated TPOT over 100

generations to identify the optimized pipeline. We used a population size of 50, which is the

number of the best pipelines that are predicted by TPOT in one generation that are then carried

to next generation, but with randomly altered parameters in their constituent ML algorithms. We

separated 80% of the constructed data sets for training and 20% for validation. The training data

subset was used through all steps, while the validation data subsets were kept out of the model

during TPOT optimization to prevent data leakage and/or training bias. For each training run, we

further applied 10-fold cross-validation to the training data subset. All condition-dependent

features were $\log_{10}$-transformed before training.

For performance assessment, we employed the adjusted coefficient of determination

score (R^2_{adj}). It is often not possible to retrieve feature importance from TPOT-optimized pipelines,

since many of the scikit-learn functions selected by TPOT do not support ranking of feature

importance. As a result, we assessed how the condition-dependent features impact the

predictions by removing them from the data sets and assessing how the TPOT-optimized

pipelines of each data set perform. To further assess the prediction results, we used ϕ and E_s to

obtain predictions about *in vivo* enzyme abundances E_s^{expl} which were compared with the

experimental value using the Pearson's correlation coefficient. We performed GO enrichment

analysis using the clusterProfiler R package (Wu et al., 2021; Yu et al., 2012). Further, we

performed flux variability analysis (FVA) to assess the variability of central metabolic pathways

and of enzyme usage pseudo-reactions.

Validation of models on genetically modified strains

To assess the validity of CAMEL on unseen conditions and its usefulness for metabolic

engineering, we predicted the *in vivo* enzyme abundances E_s^{expl} of *E. coli* strains subjected to

knock-out mutations followed by adaptive laboratory evolution (ALE) (McCloskey et al., 2018a,

2018b, 2018c, 2018d). For the optimization problem, we replicated the growth conditions by

constraining the pcGEM eciML1515 using the estimated k_{max}^{vivo} values obtained by Heckmann et

al., (2020), which were calculated from quantitative proteomics experiments on the same

conditions as McCloskey et al. (2018a, 2018b, 2018c, 2018d). Next, we used the predicted E_s to

predict ϕ using the machine learning models trained with either the k_{cat}-parameterized dataset or

the k_{max}^{vivo}-parameterized dataset. We used the predicted ϕ and E_s to obtain calculate the *in vivo*

enzyme abundances $E_s^{exp\prime}$ which we then compared with the experimental measurements of

Heckmann et al., (2020) using the Pearson's correlation coefficient.

Results and Discussion

Machine learning accurately predicts protein reserve ratios

Our first observation is that the predicted abundance, $E_{s,i}$, of an enzyme i in condition s from

pcGEMs is usually smaller than the measured *in vivo* abundance $E_{s,i}^{exp}$. This is the case since

predictions on $E_{s,i}$ must match the corresponding flux, as $v_j = k_{cat}^{i,j} \cdot E_{s,i}$, and $E_{s,i}$ cannot exceed

the measured protein abundance. We term this discrepancy between the measured and

predicted protein abundance for an enzyme i in condition s the protein reserve ratio, $\phi_{s,i}$, which is

calculated by $\frac{E_{s,i}}{E_{s,i}^{exp}}$.

Since $\phi_{s,i}$ cannot be predicted from pcGEMs alone, we rely on machine learning to train

models for ϕ_i, given data on measured protein abundance and predicted protein usage from

pcGEMs over multiple conditions (Figure 1). We refer to the collection ϕ of reserve ratios over all

proteins present in a pcGEM and measured by quantitative proteomic techniques as the

distribution of **protein reserve ratios**. Since pcGEMs can be parameterized with *in vitro*

measured or with *in vivo* estimated turnover numbers, denoted by k_{cat} and k_{max}^{vivo}, respectively, we

performed a comparative analysis of protein reserve ratios in the two parameterization scenarios.

Our first observation is that proteins differ with respect to their distributions of reserve ratios,

which are dependent on the parameterization of the pcGEMs (Figure 2). For example, the protein

reserve ratios of phosphoglycerate kinase in *E. coli* exhibit a narrow distribution for the pcGEM

parameterization based on k_{cat}, while the distribution is bimodal for the parameterization with

k_{max}^{vivo}. In contrast, the phosphoglycerate kinase in *S. cerevisiae* has the same narrow distribution

for both parameterizations. We also observed that the distributions of protein reserve ratios for all

investigated enzymes have a heavy tail, due to very large reserve ratios in few conditions.

Next, we used TPOT (Le et al., 2020) to optimize machine learning pipelines to predict ɸ

based on two types of features—static and condition-dependent (Table S1). Static features were

obtained from codon usage metrics, while condition-dependent features included enzyme usage

and fluxes of the catalyzed reactions predicted by the pcGEM in the two parameterization

scenarios (see Material and Methods). Our results indicated that the optimized pipelines for all

considered cases exhibited similar structures, composed of two parts: starting with a stacked

ensemble of linear and tree-based algorithms and then following with an Extreme Gradient

Boosting (XGBoost) algorithm (Chen and Guestrin, 2016), except for the *S. cerevisiae* model

using k_{max}^{vivo} values, whose pipeline included the Extra-trees Regressor (Table S2). Using the

trained and optimized pipelines, we then predicted ɸ for the validation data set, which had not

been used during model training and optimization. For *E. coli*, the machine learning models using

either the k_{cat} or k_{max}^{vivo} values showed excellent performance (R_{adj}^2 > 0.9), while the model for *S.*

cerevisiae showed worse performance (R_{adj}^2≈ 0.7) (Figure 3). By comparing the R_{adj}^2 values, we

found that for *E. coli*, the pipelines with condition-dependent features derived by using k_{max}^{vivo}

values outperformed those based on k_{cat} values by 6.7% (Figure 3). For yeast, however, we did

not observe a difference in the overall performance of the machine learning models between the

two parameterizations.

The reasons for the lower predictive performance for yeast in comparison to *E. coli* could

be due to several factors: First, from a modelling perspective, there is more and better-curated

knowledge about the metabolism of *E. coli* compared to that of *S. cerevisiae* (Bernstein et al.,

2023). In addition, the metabolism of yeast is compartmentalized and includes mechanisms for

enzyme regulation not accounted for in pcGEMs of any organism (e.g., organellar enzyme pools,

post-translational modifications, diffusion effects). Second, from a data perspective, a lack of

variability in the protein measurements, resulting from the consideration of samples from similar

conditions, can hamper the training of machine learning models. Putting the two perspectives

together, the eciML1515 model (1259 enzymes, (Domenzain et al., 2022)) includes a larger

number of enzymes in comparison to the ecYeast8 model (965 enzymes, (Lu et al., 2019)),

allowing to capture enzymes that may vary more across stress conditions. Lastly, the accuracy of

the k_{cat} values in the BRENDA database (Chang et al., 2021) can also affect the quality of the

predictions, given the challenges in measuring them experimentally.

Importance of fluxes for predicting protein reserve ratios

We were also interested in identifying the extent to which fluxes for reactions associated with a

protein affect the predictions of the corresponding protein reserve ratios. To this end, we removed

these condition-dependent features from the training data set and re-trained the previously

obtained TPOT-optimized pipelines using the reduced training data set. We found that the

removal of fluxes resulted in reduced predictive performance in all considered scenarios. For

instance, the obtained R^2_{adj} values for *E. coli* were at least 15.4% lower using k_{cat} values and

4.5% lower using k^{vivo}_{max}. For *S. cerevisiae*, there was a 10.8% decrease for the model using k_{cat}

values and an 8.1% decrease using k^{vivo}_{max} values (Figure 3). These findings demonstrated that the

usage of fluxes as condition-dependent features improved model performance in both organisms,

particularly in the case when pcGEMs were parametrized with k_{cat} values. In addition, this shows

that the usage of fluxes as features may, to a certain extent, mitigate the effects of model

parameterization.

The optimization problem of minimizing excess enzyme usage plays an important role in

defining how the flux distribution and enzyme usage distribution are obtained. The objective

function ensures efficient utilization of cellular resources, as it avoids burdening the cell with

excess enzymes (Bruggeman et al., 2020). We emphasize that, biologically, excess enzymes are

important for robustness and adaptability to changing environments, and its minimization could

limit the flexibility needed by cells to adapt to these conditions, leading to reduced fitness (Alter et

al., 2021). However, excess enzymes result in cellular burden, which can lead to proteotoxic

effects for the cell, also reducing fitness (Kintaka et al., 2020). We emphasize that the protein

reserve ratios we define do not necessarily correspond to biological protein reserves — we use

this terminology to account for differences between predictions from pcGEMs and measured

abundances. In that respect, one can use another objective to predict protein fraction used in

fluxes; however, the key point is that we will still rely on ratio between measured and predicted

values to build ML models — which is the essence of CAMEL.

Proteins with particular prediction error of protein reserve ratio are enriched in different

pathways

Next, we were interested if there are any distinguishing characteristics between the two groups of

proteins -- with low or high relative error in predictions. We considered a protein to show low

prediction error if the relative error between the predicted and measured protein reserve ratio was

lower than 50%, and a high prediction error if the relative error was higher than 100% in each

model (noting that varying the definitions with lower, or respectively higher, values have a

negligible effect on the conclusions).

300   First, we checked whether the two groups of proteins were enriched in any specific GO

terms (Ashburner et al., 2000; Carbon et al., 2021). For *E. coli*, we found that proteins with low

prediction error were enriched in biosynthesis of secondary metabolites, biosynthesis of

cofactors, terpenoid biosynthesis and riboflavin metabolism for models using either k_{cat} and k_{max}^{vivo}

values (Figure S1). For proteins with high prediction error, enriched GO terms shared between

models using either k_{cat} or k_{max}^{vivo} values included biosynthesis of secondary metabolites, carbon

metabolism, fatty acid metabolism and degradation (Figure S1). The five proteins with the highest

relative error (Table S3) for both model parameterizations were part of the same metabolic

pathways, namely: glycolysis, fatty acid biosynthesis, beta-oxidation, and cell wall biogenesis.

309   Similarly, for *S. cerevisiae*, the proteins with low prediction error for the model using k_{cat}

and k_{max}^{vivo} values were enriched in the biosynthesis of secondary metabolites, biosynthesis of

amino acids, and carbon metabolism. Likewise, proteins with high prediction error were enriched

in biosynthesis of secondary metabolites, biosynthesis of amino acids, carbon metabolism and

biosynthesis of cofactors for the models using k_{cat} and k_{max}^{vivo} values (Figure S2). The GO term

biosynthesis of secondary metabolites encompasses enzymes catalyzing a broad set of

reactions, that may explain its significant enrichment in all tested protein sets. The five proteins

with the highest relative error (Table S4) showed overlap between the two model

parameterizations and were part of purine metabolism, hexose metabolism, and porphyrin-

containing compound metabolism. These comparisons demonstrated that the protein reserve

ratios did not exhibit distinct patterns based on which metabolic pathway(s) in which the proteins

are involved.

To further compare the two groups of proteins, we examined whether the differences in

the relative error of predictions were associated with the coefficient of variation (CV) of protein

reserve ratios calculated from the models using either k_{cat} and k_{max}^{vivo} values. For all compared

scenarios, many proteins with low CV also had high relative errors. However, we did not identify a

statistically significant difference in CV between proteins with low and high relative errors in all

scenarios except the *E. coli* model using k_{max}^{vivo} values (Pairwise Wilcoxon rank sum test with

Bonferroni correction, Figure S3). Taken together, these results indicated that proteins with low or

high prediction errors are used in similar metabolic contexts, suggesting that other physiological

factors might contribute to the predictive performance of their abundances.

CAMEL accurately predicts *in vivo* protein abundances

Next, we used the predicted distributions of protein reserve ratios, $\hat{\phi}$, to estimate protein

abundance, E_s', which is given by $E_{s,i}' = \frac{E_{s,i}}{\hat{\phi}_{s,i}}$, where $E_{s,i}$ is the condition-specific protein abundance

predicted from the pcGEM. The obtained protein abundance estimates were compared to

experimental measurements (Chen and Nielsen, 2021; Davidi et al., 2016) by calculating the

Pearson correlation coefficient of $\log_{10}$- transformed protein abundances for each growth

condition separately (shown in Tables S5-S8) as well as the average across all conditions (Figure

4). For *E. coli*, we found a high and significant average value for the Pearson correlation

coefficient of 0.987 (p-values < 2.5e-5) over the considered conditions using the models for $\hat{\phi}$

based on the model parameterized with k_{cat} values, and 0.994 (p-values < 3.1e-11) using the

model parameterized with k_{max}^{vivo} values. In *S. cerevisiae*, using the k_{cat} parameterization, we

found a smaller, yet significant average Pearson correlation of 0.483 (p-values < 0.05 for all

significant correlations) over the considered conditions; using the pcGEM with k_{max}^{vivo} values

resulted in an average Pearson's correlation of 0.513 (p-values < 0.03 for all significant

correlations).

Here, too, we took a closer look at proteins with very high relative errors and how their

CV compare to proteins with very low relative errors. In *E. coli*, the proteins with the highest

relative error showed values larger than 1000% (Table S9). These proteins were part of the

chorismate biosynthesis, fatty acid beta-oxidation, pentose phosphate pathway and

peptidoglycan biosynthesis when predictions were based on the pcGEM with k_{cat} values. In the

case when k_{max}^{vivo} parameterization was employed, these proteins were part of fatty acid

biosynthesis, pentose phosphate pathway and purine metabolism. For *S. cerevisiae*, the proteins

with the highest errors showed more extreme values than in the case of *E. coli* (Table S10). For

both pcGEM parameterizations, the proteins with high error are related to amino acid

biosynthesis. As also seen by Lu et al., (2019) and Xia et al., (2022), this observation could be

attributed to how different strains and growth conditions affect amino acid biosynthesis pathways.

Lu et al. (2019) details that differences in energy pathways used for ATP regeneration directly

impacts the biosynthesis of amino acids, given that central metabolic pathways are the main

supplier of precursor molecules. Indeed, we observed that some of the proteins with the highest

relative error are also related to energy metabolism, such as the pyruvate decarboxylase isozyme

1 and the pyruvate dehydrogenase complex protein X component (Table S10). Xia et al. (2022)

have also made a similar observation, stating that the proteome fraction allocated to amino acid

biosynthesis is metabolism-dependent. As the cell increases protein translation, the proteome

fraction allocated to other sections of metabolism is decreased. However, for the growth

conditions we have chosen, the growth rates did not exceed the threshold to trigger the onset of

the Crabtree effect. Further, the models were not constrained on amino acid exchange reactions,

allowing the model to freely uptake required amino acids. This could explain the high error on

these proteins, since they are not being actively used for amino acid biosynthesis. Regarding the

CV, we observed no significant difference between proteins with very high or very low errors

(Figure S4) (pairwise Wilcoxon rank sum test with Bonferroni correction), except for the yeast

pcGEM with k_{max}^{vivo} values (p-value < 0.027).

The FVA analysis revealed that for *S. cerevisiae*, there is little variation on metabolic flux

through central metabolic pathways (median flux ratio of 16.3 across conditions). For *E. coli*,

there was higher variability (median flux ratio of 64.9 across conditions), likely due to the

constraints used, since less exchange reactions of central metabolites were constrained than for

S. cerevisiae. For the enzyme usage pseudo-reactions, we assessed if enzymes with high

relative error could be related to pseudo-reactions with high variability. For both *E. coli* and *S.*

cerevisiae, using either k_{cat} or k_{max}^{vivo} values, there was no relation between the enzyme usage

variability and the relative error. Proteins with either high or low relative errors exhibited similar

variability. Some proteins with high variability displayed low relative errors, although the difference

in enzyme usage variability between the proteins with low variability and proteins with high

variability was very small (median ratio variability of 1.0) (Table S11).

To further demonstrate the capabilities of CAMEL, we validated its predictive

performance using data from unseen conditions. We predicted the E_s and ϕ of mutant *E. coli*

strains subjected to ALE. We then used the predictions to calculate the E_s^{expr} and compared with

experimental values using the Pearson correlation coefficient. For the ML trained with the k_{cat}-

parameterized pcGEM data, we obtained a value of 0.626 for the Pearson correlation coefficient,

while for the ML trained with the k_{max}^{vivo}-parameterized pcGEM data, we obtained a value of 0.533

for the Pearson correlation coefficient. This result shows that CAMEL is able to obtain good

predictions for genetically modified strains, where the rationale behind metabolic optimality may

not hold.

Previous attempts for predicting protein abundance using constraint-based models

achieved good predictive performances, but they underestimated *in vivo* protein abundance. The

predictions performed by Adadi et al. (2012) using the MOMENT approach achieved a Pearson's

correlation of 0.84 between predicted protein contents and gene expression data of *E. coli* grown

on a glucose minimal medium. We showed that CAMEL with k_{max}^{vivo} parameterized pcGEM

resulted in Pearson correlations of over 0.9 for *E. coli* over different conditions, demonstrating

considerably improved performance by using the protein reserve ratios to predict protein usage

values compatible to the *in vivo* protein usage. In addition, CAMEL does not require the use of

any additional molecular read-outs, thus saving valuable resources for making predictions in

unseen conditions. We note that the performance of CAMEL is comparable to that of Resource

Balance Analysis (RBA) model of *Bacillus subtilis*, achieving an R^2 value of 0.94 (Goelzer et al.,

2015); however, predictions from RBA models consider more information about transcription,

protein translation, translocation, compartmentalization, folding and thermostability, which are

difficult to parameterize even in prokaryotic model organisms. CAMEL, on the other hand,

depends on pcGEMs, quantitative proteomics data and coding sequences for generating its

predictions.

Attempts to predict protein abundance using machine learning have previously achieved

good predictive performances but rely on either transcriptomics data or sequence-derived

features, which have low correlation to protein content, and are invariable to changes in

environmental conditions, respectively (Ferreira et al., 2021; Li et al., 2019). For instance, the

model constructed by Terai and Asai (Terai and Asai, 2020), based on three different algorithms,

was trained to predict protein abundance in *E. coli* using sequence-derived features and mRNA

structure information. The model performance was assessed by Spearman correlation, which

ranged from 0.55 to 0.71. Predictions for *S. cerevisiae* were carried out by the Bayesian network

constructed by Mehdi et al. (Mehdi et al., 2014), which combined transcriptomics measurements

and sequence-derived features, using data from *S. cerevisiae* and *Schizosaccharomyces pombe*.

The predictions were evaluated by Spearman correlation, which ranged from 0.61 to 0.77.

Ferreira et al. (Ferreira et al., 2021) generated a predictive model using only codon usage metrics

as features, which achieved a Spearman correlation of 0.74. Although the resulting correlations

were higher than those from CAMEL, both studies relied only on data obtained from optimal

growth conditions. However, CAMEL can be effectively used to make predictions of protein

abundance in sub-optimal scenarios, which is a notoriously difficult endeavor (Li et al., 2019).

Given that predicted protein abundance based on the protein reserve ratios are in good

agreement with experimental values, these results indicated that the coupling of pcGEMs and

machine learning can accurately predict the protein allocation *in vivo*.

By coupling constraint-based approaches and machine learning approaches, we

demonstrated that our proposed approach, CAMEL, generated accurate predictions of the protein

reserve ratios. In addition, we showed that in combination with constraint-based models, CAMEL

leverages these machine learning models to obtain a prediction of protein abundance that

overcome notable limitations inherent to pcGEMs (e.g., over-accumulation of enzymes,

uncertainty in k_{cat} values). As its most notable merit, CAMEL predicts protein abundance by

relying on physiological data, used in the prediction of fluxes, rather than requiring additional

omics measurements such as gene expression. Therefore, CAMEL is readily applicable to any

condition given that measurements of growth rates and exchange fluxes are available, widening

the possibilities for its biotechnological applications (e.g., design of strains). In addition, as our

findings demonstrated, CAMEL results in good predictions for protein abundance for sub-optimal

conditions, opening the possibility for using these predictions as features to model other complex

physiological traits.

Code and data availability

The code and data can be found in the GitHub repository: https://github.com/mauricioamf/CAMEL

Author Contributions:

Conceptualization, MAMF, PW, MA, WBS and ZN.; Methodology, MAMF, PW, MA, WBS and ZN.; Formal Analysis, MAMF.; Investigation, MAMF.; Writing – Original Draft, MAMF and ZN.; Writing – Review & Editing, MAMF, PW, MA, WBS and ZN.; Supervision, WBS and ZN. Funding Acquisition, WBS and ZN. Project Administration, WBS and ZN.

Acknowledgments

This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brasil (CAPES) – Finance Code 001. P.W. and Z.N. acknowledge the support from the Research Focus Area "Evolutionary Systems Biology" of the University of Potsdam. M.A. and Z.N. acknowledge the support of the Max Planck Society. M.A. and Z.N were supported by the European Union's Horizon 2020 research and innovation programme, project PlantaSYST (SGA-CSA No. 739582 under FPA No. 664620).

Declaration of interests

The authors declare no competing interests.

References

Adadi, R., Volkmer, B., Milo, R., Heinemann, M., Shlomi, T., 2012. Prediction of Microbial Growth
Rate versus Biomass Yield by a Metabolic Network with Kinetic Parameters. PLoS Comput
Biol 8, e1002575. https://doi.org/10.1371/JOURNAL.PCBI.1002575

Alter, T.B., Blank, L.M., Ebert, B.E., 2021. Proteome Regulation Patterns Determine Escherichia
coli Wild-Type and Mutant Phenotypes. mSystems 6.
https://doi.org/10.1128/msystems.00625-20

Ashburner, M., Ball, C.A., Blake, J.A., Botstein, D., Butler, H., Cherry, J.M., Davis, A.P., Dolinski, K.,
Dwight, S.S., Eppig, J.T., Harris, M.A., Hill, D.P., Issel-Tarver, L., Kasarskis, A., Lewis, S.,
Matese, J.C., Richardson, J.E., Ringwald, M., Rubin, G.M., Sherlock, G., 2000. Gene
Ontology: tool for the unification of biology. Nat Genet 25, 25.
https://doi.org/10.1038/75556

Beg, Q.K., Vazquez, A., Ernst, J., De Menezes, M.A., Bar-Joseph, Z., Barabási, A.L., Oltvai, Z.N.,
2007. Intracellular crowding defines the mode and sequence of substrate uptake by
Escherichia coli and constrains its metabolic activity. Proc Natl Acad Sci U S A 104, 12663–
12668. https://doi.org/10.1073/PNAS.0609845104/SUPPL_FILE/09845DATASET5.XLS

Bekiaris, P.S., Klamt, S., 2020. Automatic construction of metabolic models with enzyme
constraints. BMC Bioinformatics 21, 1–13. https://doi.org/10.1186/S12859-019-3329-
9/TABLES/2

Bernstein, D.B., Akkas, B., Price, M.N., Arkin, A.P., 2023. Critical assessment of E. coli genome-
scale metabolic model with high-throughput mutant fitness data. bioRxiv
2023.01.05.522875. https://doi.org/10.1101/2023.01.05.522875

Bruggeman, F.J., Planqué, R., Molenaar, D., Teusink, B., 2020. Searching for principles of
microbial physiology. FEMS Microbiol Rev 44, 821–844.
https://doi.org/10.1093/FEMSRE/FUAA034

Calderón-Celis, F., Encinar, J.R., Sanz-Medel, A., 2018. Standardization approaches in absolute
quantitative proteomics with mass spectrometry. Mass Spectrom Rev 37, 715–737.
https://doi.org/10.1002/mas.21542

Carbon, S., Douglass, E., Good, B.M., Unni, D.R., Harris, N.L., Mungall, C.J., Basu, S., Chisholm,
R.L., Dodson, R.J., Hartline, E., Fey, P., Thomas, P.D., Albou, L.P., Ebert, D., Kesling, M.J., Mi,
H., Muruganujan, A., Huang, X., Mushayahama, T., LaBonte, S.A., Siegele, D.A., Antonazzo,
G., Attrill, H., Brown, N.H., Garapati, P., Marygold, S.J., Trovisco, V., dos Santos, G., Falls, K.,
Tabone, C., Zhou, P., Goodman, J.L., Strelets, V.B., Thurmond, J., Garmiri, P., Ishtiaq, R.,
Rodríguez-López, M., Acencio, M.L., Kuiper, M., Lægreid, A., Logie, C., Lovering, R.C.,
Kramarz, B., Saverimuttu, S.C.C., Pinheiro, S.M., Gunn, H., Su, R., Thurlow, K.E., Chibucos,
M., Giglio, M., Nadendla, S., Munro, J., Jackson, R., Duesbury, M.J., Del-Toro, N., Meldal,
B.H.M., Paneerselvam, K., Perfetto, L., Porras, P., Orchard, S., Shrivastava, A., Chang, H.Y.,
Finn, R.D., Mitchell, A.L., Rawlings, N.D., Richardson, L., Sangrador-Vegas, A., Blake, J.A.,
Christie, K.R., Dolan, M.E., Drabkin, H.J., Hill, D.P., Ni, L., Sitnikov, D.M., Harris, M.A., Oliver,
S.G., Rutherford, K., Wood, V., Hayles, J., Bähler, J., Bolton, E.R., de Pons, J.L., Dwinell,

M.R., Hayman, G.T., Kaldunski, M.L., Kwitek, A.E., Laulederkind, S.J.F., Plasterer, C., Tutaj,
M.A., Vedi, M., Wang, S.J., D'Eustachio, P., Matthews, L., Balhoff, J.P., Aleksander, S.A.,
Alexander, M.J., Cherry, J.M., Engel, S.R., Gondwe, F., Karra, K., Miyasato, S.R., Nash, R.S.,
Simison, M., Skrzypek, M.S., Weng, S., Wong, E.D., Feuermann, M., Gaudet, P., Morgat, A.,
Bakker, E., Berardini, T.Z., Reiser, L., Subramaniam, S., Huala, E., Arighi, C.N., Auchincloss,
A., Axelsen, K., Argoud-Puy, G., Bateman, A., Blatter, M.C., Boutet, E., Bowler, E., Breuza, L.,
Bridge, A., Britto, R., Bye-A-Jee, H., Casas, C.C., Coudert, E., Denny, P., Es-Treicher, A.,
Famiglietti, M.L., Georghiou, G., Gos, A.N., Gruaz-Gumowski, N., Hatton-Ellis, E., Hulo, C.,
Ignatchenko, A., Jungo, F., Laiho, K., Le Mercier, P., Lieberherr, D., Lock, A., Lussi, Y.,
MacDougall, A., Ma-Grane, M., Martin, M.J., Masson, P., Natale, D.A., Hyka-Nouspikel, N.,
Orchard, S., Pedruzzi, I., Pourcel, L., Poux, S., Pundir, S., Rivoire, C., Speretta, E., Sundaram,
S., Tyagi, N., Warner, K., Zaru, R., Wu, C.H., Diehl, A.D., Chan, J.N., Grove, C., Lee, R.Y.N.,
Muller, H.M., Raciti, D., van Auken, K., Sternberg, P.W., Berriman, M., Paulini, M., Howe, K.,
Gao, S., Wright, A., Stein, L., Howe, D.G., Toro, S., Westerfield, M., Jaiswal, P., Cooper, L.,
Elser, J., 2021. The Gene Ontology resource: enriching a GOld mine. Nucleic Acids Res 49,
D325–D334. https://doi.org/10.1093/NAR/GKAA1113

Chang, A., Jeske, L., Ulbrich, S., Hofmann, J., Koblitz, J., Schomburg, I., Neumann-Schaal, M.,
Jahn, D., Schomburg, D., 2021. BRENDA, the ELIXIR core data resource in 2021: New
developments and updates. Nucleic Acids Res 49, D498–D508.
https://doi.org/10.1093/nar/gkaa1025

Chen, T., Guestrin, C., 2016. XGBoost: A scalable tree boosting system. Proceedings of the ACM
SIGKDD International Conference on Knowledge Discovery and Data Mining 13-17-Augu,
785–794. https://doi.org/10.1145/2939672.2939785

Chen, Y., Nielsen, J., 2021. In vitro turnover numbers do not reflect in vivo activities of yeast
enzymes. Proc Natl Acad Sci U S A 118, e2108391118.
https://doi.org/10.1073/PNAS.2108391118/SUPPL_FILE/PNAS.2108391118.SD08.XLSX

Davidi, D., Noor, E., Liebermeister, W., Bar-Even, A., Flamholz, A., Tummler, K., Barenholz, U.,
Goldenfeld, M., Shlomi, T., Milo, R., 2016. Global characterization of in vivo enzyme
catalytic rates and their correspondence to in vitro kcat measurements. Proc Natl Acad Sci
U S A 113, 3401–3406. https://doi.org/10.1073/pnas.1514240113

Di Bartolomeo, F., Malina, C., Campbell, K., Mormino, M., Fuchs, J., Vorontsov, E., Gustafsson,
C.M., Nielsen, J., 2020. Absolute yeast mitochondrial proteome quantification reveals
trade-off between biosynthesis and energy generation during diauxic shift. Proc Natl Acad
Sci U S A 117, 7524–7535.
https://doi.org/10.1073/PNAS.1918216117/SUPPL_FILE/PNAS.1918216117.SD07.XLSX

Domenzain, I., Sánchez, B., Anton, M., Kerkhoven, E.J., Millán-Oropeza, A., Henry, C., Siewers, V.,
Morrissey, J.P., Sonnenschein, N., Nielsen, J., 2022. Reconstruction of a catalogue of
genome-scale metabolic models with enzymatic constraints using GECKO 2.0. Nat Commun
13, 1–13. https://doi.org/10.1038/s41467-022-31421-1

Eraslan, B., Wang, D., Gusic, M., Prokisch, H., Orn, B., Hallstr€ Om, M., Uhl En, M., Asplund, A.,
Pont En, F., Wieland, T., Hopf, T., Hahne, H., Kuster, B., Gagneur, J., 2019. Quantification

and discovery of sequence determinants of protein-per-mRNA amount in 29 human
tissues. Mol Syst Biol 15, e8513. https://doi.org/10.15252/MSB.20188513

Ferreira, M., Ventorim, R., Almeida, E., Silveira, S., Silveira, W., 2021. Protein Abundance
Prediction Through Machine Learning Methods. J Mol Biol 433, 167267.
https://doi.org/10.1016/J.JMB.2021.167267

Goelzer, A., Muntel, J., Chubukov, V., Jules, M., Prestel, E., Nölker, R., Mariadassou, M.,
Aymerich, S., Hecker, M., Noirot, P., Becher, D., Fromion, V., 2015. Quantitative prediction
of genome-wide resource allocation in bacteria. Metab Eng 32, 232–243.
https://doi.org/10.1016/J.YMBEN.2015.10.003

Heckmann, D., Campeau, A., Lloyd, C.J., Phaneuf, P. V., Hefner, Y., Carrillo-Terrazas, M., Feist,
A.M., Gonzalez, D.J., Palsson, B.O., 2020. Kinetic profiling of metabolic specialists
demonstrates stability and consistency of in vivo enzyme turnover numbers. Proc Natl
Acad Sci U S A 117, 23182–23190.
https://doi.org/10.1073/PNAS.2001562117/SUPPL_FILE/PNAS.2001562117.SD01.XLSX

Heckmann, D., Lloyd, C.J., Mih, N., Ha, Y., Zielinski, D.C., Haiman, Z.B., Desouki, A.A., Lercher,
M.J., Palsson, B.O., 2018. Machine learning applied to enzyme turnover numbers reveals
protein structural correlates and improves metabolic models. Nat Commun 9, 1–10.
https://doi.org/10.1038/s41467-018-07652-6

Kintaka, R., Makanae, K., Namba, S., Kato, H., Kito, K., Ohnuki, S., Ohya, Y., Andrews, B.J., Boone,
C., Moriya, H., 2020. Genetic profiling of protein burden and nuclear export overload. Elife
9, 1–22. https://doi.org/10.7554/ELIFE.54080

Kültz, D., 2020. Evolution of cellular stress response mechanisms. J Exp Zool A Ecol Integr Physiol
333, 359–378. https://doi.org/10.1002/JEZ.2347

Lahtvee, P.J., Sánchez, B.J., Smialowska, A., Kasvandik, S., Elsemman, I.E., Gatto, F., Nielsen, J.,
2017. Absolute Quantification of Protein and mRNA Abundances Demonstrate Variability in
Gene-Specific Translation Efficiency in Yeast. Cell Syst 4, 495-504.e5.
https://doi.org/10.1016/J.CELS.2017.03.003

Le, T.T., Fu, W., Moore, J.H., 2020. Scaling tree-based automated machine learning to
biomedical big data with a feature set selector. Bioinformatics 36, 250–256.
https://doi.org/10.1093/BIOINFORMATICS/BTZ470

Li, G., Hu, Y., Jan Zrimec, Luo, H., Wang, H., Zelezniak, A., Ji, B., Nielsen, J., 2021. Bayesian
genome scale modelling identifies thermal determinants of yeast metabolism. Nat
Commun 12, 1–12. https://doi.org/10.1038/s41467-020-20338-2

Li, H., Siddiqui, O., Zhang, H., Guan, Y., 2019. Joint learning improves protein abundance
prediction in cancers. BMC Biol 17, 1–14. https://doi.org/10.1186/S12915-019-0730-
9/FIGURES/6

Lill, J.R., Mathews, W.R., Rose, C.M., Schirle, M., 2021. Proteomics in the pharmaceutical and
biotechnology industry: a look to the next decade.

https://doi.org/10.1080/14789450.2021.1962300 18, 503–526.
https://doi.org/10.1080/14789450.2021.1962300

Liu, Y., Lu, S., Liu, K., Wang, S., Huang, L., Guo, L., 2019. Proteomics: a powerful tool to study
plant responses to biotic stress. Plant Methods 2019 15:1 15, 1–20.
https://doi.org/10.1186/S13007-019-0515-8

Lu, H., Li, F., Sánchez, B.J., Zhu, Z., Li, G., Domenzain, I., Marcišauskas, S., Anton, P.M., Lappa, D.,
Lieven, C., Beber, M.E., Sonnenschein, N., Kerkhoven, E.J., Nielsen, J., 2019. A consensus S.
cerevisiae metabolic model Yeast8 and its ecosystem for comprehensively probing cellular
metabolism. Nat Commun 10. https://doi.org/10.1038/s41467-019-11581-3

McCloskey, D., Xu, S., Sandberg, T.E., Brunk, E., Hefner, Y., Szubin, R., Feist, A.M., Palsson, B.O.,
2018a. Adaptive laboratory evolution resolves energy depletion to maintain high aromatic
metabolite phenotypes in Escherichia coli strains lacking the Phosphotransferase System.
Metab Eng 48, 233–242. https://doi.org/10.1016/J.YMBEN.2018.06.005

McCloskey, D., Xu, S., Sandberg, T.E., Brunk, E., Hefner, Y., Szubin, R., Feist, A.M., Palsson, B.O.,
2018b. Adaptation to the coupling of glycolysis to toxic methylglyoxal production in tpiA
deletion strains of Escherichia coli requires synchronized and counterintuitive genetic
changes. Metab Eng 48, 82–93. https://doi.org/10.1016/J.YMBEN.2018.05.012

McCloskey, D., Xu, S., Sandberg, T.E., Brunk, E., Hefner, Y., Szubin, R., Feist, A.M., Palsson, B.O.,
2018c. Multiple optimal phenotypes overcome redox and glycolytic intermediate
metabolite imbalances in Escherichia coli pgi knockout evolutions. Appl Environ Microbiol
84, 823–841. https://doi.org/10.1128/AEM.00823-18/SUPPL_FILE/ZAM019188775SD8.CSV

McCloskey, D., Xu, S., Sandberg, T.E., Brunk, E., Hefner, Y., Szubin, R., Feist, A.M., Palsson, B.O.,
2018d. Growth adaptation of gnd and sdhCB Escherichia coli deletion strains diverges from
a similar initial perturbation of the transcriptome. Front Microbiol 9, 398699.
https://doi.org/10.3389/FMICB.2018.01793/BIBTEX

Mehdi, A.M., Patrick, R., Bailey, T.L., Boden, M., 2014. Predicting the Dynamics of Protein
Abundance. Molecular & Cellular Proteomics 13, 1330–1340.
https://doi.org/10.1074/MCP.M113.033076

Mergner, J., Frejno, M., List, M., Papacek, M., Chen, X., Chaudhary, A., Samaras, P., Richter, S.,
Shikata, H., Messerer, M., Lang, D., Altmann, S., Cyprys, P., Zolg, D.P., Mathieson, T.,
Bantscheff, M., Hazarika, R.R., Schmidt, T., Dawid, C., Dunkel, A., Hofmann, T., Sprunck, S.,
Falter-Braun, P., Johannes, F., Mayer, K.F.X., Jürgens, G., Wilhelm, M., Baumbach, J., Grill,
E., Schneitz, K., Schwechheimer, C., Kuster, B., 2020. Mass-spectrometry-based draft of the
Arabidopsis proteome. Nature 579, 409–414. https://doi.org/10.1038/s41586-020-2094-2

Nielsen, J., 2019. Yeast Systems Biology: Model Organism and Cell Factory. Biotechnol J 14, 1–9.
https://doi.org/10.1002/biot.201800421

Novoa, E.M., Jungreis, I., Jaillon, O., Kellis, M., Leitner, T., 2019. Elucidation of Codon Usage
Signatures across the Domains of Life. Mol Biol Evol 36, 2328–2339.
https://doi.org/10.1093/molbev/msz124

Otto, A., Becher, D., Schmidt, F., 2014. Quantitative proteomics in the field of microbiology.
Proteomics 14, 547–565. https://doi.org/10.1002/pmic.201300403

Pappireddi, N., Martin, L., Wühr, M., 2019. A Review on Quantitative Multiplexed Proteomics.
ChemBioChem 20, 1210–1224. https://doi.org/10.1002/cbic.201800650

Peebo, K., Valgepea, K., Maser, A., Nahku, R., Adamberg, K., Vilu, R., 2015. Proteome
reallocation in Escherichia coli with increasing specific growth rate. Mol Biosyst 11, 1184–
1193. https://doi.org/10.1039/C4MB00721B

Sánchez, B.J., Zhang, C., Nilsson, A., Lahtvee, P., Kerkhoven, E.J., Nielsen, J., 2017. Improving the
phenotype predictions of a yeast genome-scale metabolic model by incorporating
enzymatic constraints. Mol Syst Biol 13, 935. https://doi.org/10.15252/msb.20167411

Schmidt, A., Kochanowski, K., Vedelaar, S., Ahrné, E., Volkmer, B., Callipo, L., Knoops, K., Bauer,
M., Aebersold, R., Heinemann, M., 2016. The quantitative and condition-dependent
Escherichia coli proteome. Nat Biotechnol 34, 104–110. https://doi.org/10.1038/nbt.3418

Schubert, O.T., Röst, H.L., Collins, B.C., Rosenberger, G., Aebersold, R., 2017. Quantitative
proteomics: challenges and opportunities in basic and applied research. Nature Protocols
2017 12:7 12, 1289–1294. https://doi.org/10.1038/nprot.2017.040

Terai, G., Asai, K., 2020. Improving the prediction accuracy of protein abundance in Escherichia
coli using mRNA accessibility. Nucleic Acids Res 48, e81–e81.
https://doi.org/10.1093/nar/gkaa481

Torres-García, W., Zhang, W., Runger, G.C., Johnson, R.H., Meldrum, D.R., 2009. Integrative
analysis of transcriptomic and proteomic data of Desulfovibrio vulgaris: a non-linear model
to predict abundance of undetected proteins. Bioinformatics 25, 1905–1914.
https://doi.org/10.1093/BIOINFORMATICS/BTP325

Valgepea, K., Adamberg, K., Seiman, A., Vilu, R., 2013. Escherichia coli achieves faster growth by
increasing catalytic and translation rates of proteins. Mol Biosyst 9, 2344–2358.
https://doi.org/10.1039/C3MB70119K

Wendering, P., Arend, M., Razaghi-Moghadam, Z., Nikoloski, Z., 2023. Data integration across
conditions improves turnover number estimates and metabolic predictions. Nature
Communications 2023 14:1 14, 1–12. https://doi.org/10.1038/s41467-023-37151-2

Wu, T., Hu, E., Xu, S., Chen, M., Guo, P., Dai, Z., Feng, T., Zhou, L., Tang, W., Zhan, L., Fu, X., Liu,
S., Bo, X., Yu, G., 2021. clusterProfiler 4.0: A universal enrichment tool for interpreting
omics data. Innovation 2, 100141. https://doi.org/10.1016/j.xinn.2021.100141

Xia, J., Sánchez, B.J., Chen, Y., Campbell, K., Kasvandik, S., Nielsen, J., 2022. Proteome allocations
change linearly with the specific growth rate of Saccharomyces cerevisiae under glucose
limitation. Nature Communications 2022 13:1 13, 1–12. https://doi.org/10.1038/s41467-
022-30513-2

Xu, R., Razaghi-Moghadam, Z., Nikoloski, Z., 2021. Maximization of non-idle enzymes improves
the coverage of the estimated maximal in vivo enzyme catalytic rates in Escherichia coli.
Bioinformatics 37, 3848–3855. https://doi.org/10.1093/BIOINFORMATICS/BTAB575

Yu, G., Wang, L.G., Han, Y., He, Q.Y., 2012. ClusterProfiler: An R package for comparing biological
themes among gene clusters. OMICS 16, 284–287.
https://doi.org/10.1089/OMI.2011.0118/ASSET/IMAGES/LARGE/FIGURE1.JPEG

Yu, R., Campbell, K., Pereira, R., Björkeroth, J., Qi, Q., Vorontsov, E., Sihlbom, C., Nielsen, J.,
2020. Nitrogen limitation reveals large reserves in metabolic and translational capacities of
yeast. Nat Commun 11, 1–12. https://doi.org/10.1038/s41467-020-15749-0

Figures

667
Figure 1. Schematic overview of CAMEL, a constraint-based approach to predict protein

abundance. CAMEL uses two types of features, marked as blue boxes: static, obtained from

codon usage metrics, and condition-dependent, given by metabolic flux and enzyme usage

predicted by constraint-based modelling with pcGEMs. The predicted enzyme usage is employed

together with experimental proteomics data to calculate protein reserve ratios that are then used

as the target variable for machine learning (green box). The predictive model is obtained by

selecting the optimal ensemble of machine learning approaches using TPOT. The performance is

assessed by using cross-validation based on regression metrics. Abbreviations: Tr – training; Val

– validation.

Protein reserve ratios

In vitro k_{cat} values   *In vivo* k_{max}^{vivo} values

678
679   **Figure 2. Distribution of protein reserve ratios.** The selected proteins take part in central

680   metabolic pathways such as glycolysis, pentose phosphate pathway and citric acid cycle, and are

681   thus present in both organisms. Protein reserve ratios are unitless values.

683
684 **Figure 3. Performance of models from optimized pipelines**. The figure shows a comparison of

685 R^2_{adj} scores for the optimized pipelines based on all features (blue) or excluding condition-

686 dependent features (grey) with either *in vitro* k_{cat} or k^{vivo}_{max} values. Negative percentages (red)

687 indicate a decrease in predictive performance compared to scores obtained from the models

688 trained using all features. Error bars indicate R^2_{adj} scores obtained from 10-fold cross-validation.

690
Figure 4. Comparison of measured and predicted protein abundances. The colour scheme

indicates the relative error when comparing the measured (E_s^{exp}) and predicted protein

abundances ($E_s^{exp\prime}$). Relative error was calculated using protein abundances in nominal scale.

Panel (A) shows predictions using the *E. coli* model with *in vitro* k_{cat} values (A), and in vivo k_{max}^{vivo}

values (B). The panels (C) and (D), depict the prediction performance of the *S. cerevisiae* model

with k_{cat} values (C) and k_{max}^{vivo} values (D).