Title:

Structures of Langya virus fusion protein ectodomain in pre and post fusion conformation

Authors:

Aaron J. May[1,2], Karunakar Reddy Pothula[1], Katarzyna Janowska[1], Priyamvada Acharya[1,2,3]

Affiliations:
[1] Duke Human Vaccine Institute, Durham NC 27710, USA
[2] Duke University, Department of Biochemistry, Durham NC 27710, USA
[3] Duke University, Department of Surgery, Durham NC 27710, USA
11
12 *Correspondence: aaron.may@duke.edu (A.J.M), priyamvada.acharya@duke.edu (P.A.)

Abstract:

Langya virus (LayV) is a paramyxovirus in the *Henipavirus* genus, closely related to the deadly Nipah and Hendra viruses, that was identified in August 2022 through disease surveillance following animal exposure in eastern China. Paramyxoviruses present two glycoproteins on their surface, known as attachment and fusion proteins, that mediate entry into cells and constitute the primary antigenic targets for immune response. Here, we determine cryo-EM structures of the uncleaved LayV fusion protein (F) ectodomain in pre- and post-fusion conformations. The LayV-F protein exhibits pre- and post-fusion architectures that, despite being highly conserved across paramyxoviruses, show differences in their surface properties, in particular at the apex of the prefusion trimer, that may contribute to antigenic variability. While dramatic conformational changes were visualized between the pre- and post-fusion forms of the LayV-F protein, several domains remained invariant, held together by highly conserved disulfides. The LayV-F fusion peptide is buried within a highly conserved, hydrophobic, interprotomer pocket in the pre-fusion state and is notably less flexible than the rest of the protein, highlighting its "spring-loaded" state and suggesting that the mechanism of pre-to-post transition must involve perturbations to the pocket and release of the fusion peptide. Together, these results offer a structural basis for how the Langya virus fusion protein compares to its Henipavirus relatives and propose a mechanism for the initial step of pre- to post-fusion conversion that may apply more broadly to paramyxoviruses.

Importance:

The *Henipavirus* genus is quickly expanding into new animal hosts and geographic locations. This study compares the structure and antigenicity of the Langya virus fusion protein to other henipaviruses, which has important vaccine or therapeutic development implications. Furthermore, the study proposes a new mechanism to explain the early steps of the fusion initiation process that can be more broadly applied to the *Paramyxoviridae* family.

 Figure Legends:

Figure 1: Structures of LayV-F in pre- and post-fusion conformation:

A-B. Cryo-EM reconstructions of LayV-F in **A.** pre- and **B.** post-fusion conformation, colored
by protomer. Map regions colored in different shades of blue (pre-fusion) and yellow (post-
fusion) for each protomer. Representative reference-free 2D classes used for the reconstruction
are displayed above each map. **C-D.** Cartoon representation of atomic models of LayV-F in **C.**
pre-fusion and **D.** post-fusion conformation that were built into the maps shown in **A-B**. In each,
one protomer is colored by domain, based on the color scheme in panel **E**, and the other two
protomers colored gray. **E.** Sequence domain key for LayV-F. SS: Secretion signal, cleaved prior
to purification. DI, DII, DIII: Domains I, II, and III. FP: Fusion Peptide. HRA, HRB: Heptad
repeats A and B. Foldon: Trimerization domain. TS: Twin-Strep tag, used for purification with
modified streptavidin resin.

Figure 2: Role of conserved cysteines during conformational conversion.

A. Zoomed-in view of the DIII domain, including the fusion peptide (orange) and HRA (yellow),
shown for the (left) pre- and (right) post-fusion structures. Color scheme continued from Figure
1E. The cleavage site at R104 is identified, as well as the disulfide bond formed by C66 and
C187 (yellow sticks). This disulfide bond is a hinge point around which the HRA straightens into
the central coiled-coil in the post-fusion F structure. **B-C.** Superimposition of pre- and post-
fusion conformation domains DI (**B**) and DII (**C**). Color scheme continued from figure 1E, with
the post-fusion conformation colored in a lighter shade. Disulfide bonds are shown as yellow
sticks and labelled. For DII, the movement of the HRB linker region is indicated by an arrow. **D.**
Sequence alignment of paramyxovirus fusion proteins in the regions surrounding conserved
cysteines, with cysteine residue number and domain labelled. Viruses colored blue are from the
Henipavirus genus and colored tan for other paramyxoviruses. Residues are colored by type, per
MView standard (1). Blue: Alcohol, Green: Hydrophobic, Dark Blue: Negative charge, Red:
Positive charge, Purple: Polar, Yellow: Cysteine.

Figure 3: Fusion peptide burial and conservation.

A. Structure of the pre-fusion LayV-F protein shown in cartoon representation and colored by B-
factors. **B.** Difference Distance Matrix plot comparing relative residue positions between pre-
fusion and post-fusion conformation. Sequence key from figure 1E included along each axis.
Gray areas in the bottom right half of the plot are the result of sequences not present in the post-
fusion model. **C.** Hydrophobic fusion peptide pocket in pre-fusion state, with fusion peptide
depicted as orange cartoon, and surface electrostatics view of the pocket. **D.** Side-chain stick
view of the fusion peptide pocket, with hydrophobic residues involved shown with side chains as
green sticks. An enlarged view of this panel with residues labeled is seen in supplemental figure
S5. **E.** Cartoon representations of a single pre-fusion state protomer, colored by sequence
conservation across *Henipavirus* fusion proteins. Color scale generated by ChimeraX default
AL2CO entropy measure. **F.** Zoom in of the fusion peptide pocket, colored by sequence

conservation as in H. The fusion peptide pocket is among the most strongly conserved region of
the protein.

Figure 4: Structure and antigenicity comparison of Henipavirus fusion proteins.

A. Superimposition of a single pre-fusion protomer of LayV-F (blue) NiV-F (cyan) and HeV-F
(green), in cartoon representation. PDBs: LayV-F: 8FEJ, NiV-F: 5EVM, HeV-F: 5EJB (2, 3). **B.**
Zoomed in view of a loop within the folded HRA, one of the few areas of significant structural
difference between LayV-F and NiV/HeV-F. **C.** Binding site of known paramyxovirus F protein-
neutralizing antibodies, labeled by the species targeted. NiV-F trimer from 12B2-bound structure
shown as cartoon and antibody Fabs shown in surface view. Only one Fab per type shown, all
expect PIA174 bind with 3:1 stoichiometry. PDBs: PIA174: 6MJZ, 12B2: 7KI4, mAb66: 6T3F,
1F5: 7KI6, 5B3: 6TYS (4-6). **D.** Surface view of LayV-F and NiV-F with glycosylation sites
(residues meeting the N-X-S/T glycosylation sequence) colored teal. **E.** Surface representation of
LayV-F colored by sequence conservation across *Henipavirus* fusion proteins. Color scale
generated by ChimeraX default AL2CO entropy measure; magenta indicates high conservation
and teal indicates high variability (7). **F.** Surface representation of paramyxovirus fusion
proteins, showing only main chain atoms, colored by residue type and shown from top-down
view. Tan: hydrophobic, Pink: polar, Red: negative, Blue: positive, teal: Potential glycosylation.
PIV5 PBD: 4GIP (8).

Figure 5: Fusion triggering hypothesis for paramyxoviruses.

A. Existing model of *Henipavirus* (HNV) fusion protein triggering based on Liu *et al.* (2013) (9).
HNV-G, brown, shown in a two-receptor binding domain down "closed" state, bound to HNV-F
in pre-fusion conformation. Exposure to Ephrin B2/B3 receptor induces opening of HNV-G,
triggering conversion of HNV-F from pre-fusion to the pre-hairpin state. The triggering
interaction is labeled with a star. **B.** Hypothesized structural basis for the G-F interaction shown
in **A**, wherein a two-step mechanism involving disruption of the fusion peptide burial pocket
through action of the HNV-G stalk on DII frees the fusion peptide, allowing conformational
conversion.

Supplemental Figure 1: Sequence Alignment of *Paramyxoviridae* Fusion Proteins.

Sequence alignment of select paramyxovirus fusion proteins with LayV-F used as the reference
sequence. Cov: percent sequence coverage. Pid: percent sequence identity. Residues are colored
by type, per MView standard (1). Blue: Alcohol, Green: Hydrophobic, Dark Blue: Negative
charge, Red: Positive charge, Purple: Polar, Yellow: Cysteine. Consensus residue types assigned
per MView standard. Alcohol: (o), Aliphatic: (l), Aromatic: (a), Charged: (c), Hydrophobic: (h),
Negative: (-), Polar: (p), Positive: (+), Small: (s), Tiny: (u), Turnlike: (t).

Supplemental Figure 2: Purification and Antibody Binding of *Henipavirus* Proteins.

A. Size-exclusion chromatogram of *Henipavirus* fusion protein purification. LayV-F, blue, NiV-
F WT, dashed red, pre-fusion stabilized NiVop08 (10), dashed green. The fractions identified as
the ectodomain trimer at 180kDa are indicated with a vertical dashed line **B.** Differential
scanning fluorimetry profiles of the products from A, LayV-F measured in triplicate. **C-E.** SDS-

PAGE of *Henipavirus* constructs from A. Molecular weight markers, MW, non-reducing
conditions, NR, reducing conditions, R., **C.** LayV-F loaded at 1µg per lane. **D.** LayV-F loaded at
8µg per lane. **E.** NiVop08 loaded at 3µg per lane, with left two bands sampling the first peak at
440kDa and the second two bands sampling the trimer peak at 180kDa. **F.** BLI sensorgram of
anti-NiV-F antibodies h5B3.1 and 12B2 binding to immobilized LayV-F or NiVop08 (6, 11).

Supplemental Figure 3: Cryo-EM processing of LayV-F structures.

Cryo-EM processing of pre- and post-fusion LayV-F structures. Steps performed using
CryoSPARC (12, 13). CTF Estimation performed on micrographs followed by default blob
picker algorithm and a trained AI-based algorithm (Topaz) for particle picking. Particles for the
post-fusion state were obtained through the blob picker, while particles contributing to the pre-
fusion state came from both sets. Heterogeneous refinement was used to classify and further sort
particles. After non-uniform refinement, a local refinement was performed on the entire
structure. The supplemental map of the post-fusion state prior to local refinement where the N65
glycan is detected has also been deposited.

Supplemental Figure 4: Fusion Peptide comparison.

A-E: Comparison of fusion peptide location in other viral fusion proteins. In the top portion of
each panel, fusion proteins are shown in cartoon representation in gray, with the fusion peptide
as orange spheres, and in the bottom portion, fusion peptides are shown as cartoon representation
with side chains as sticks, orange, with the surface of the rest of the fusion protein colored based
on protomer. The most proximal protomer to the highlighted fusion peptide is colored cyan, with
the neighboring protomers colored green and tan. In cases where there are are multiple subunits
in the fusion protein, the attachment subunit is colored a darker shade than the corresponding
fusion subunit. **A.** HIV Envelope (Env) closed conformation, with (PDB: 4TVP) and without
fusion peptide-targeting antibody VRC34.01 (PDB: 5I8H), shown in cartoon, red. **B**. HIV Env in
open conformation (PDB: 5VN3). **C.** Influenza HA in closed conformation (PDB: 6HJN). **D.**
SARS-CoV-2 Spike in pre-fusion, receptor-binding domain down conformation (PDB: 6XR8).
E. Ebola GP in pre-fusion conformation (PDB: 3CSY). (14-19)

Supplemental Figure 5: Fusion Peptide Pocket

Enlarged view of figure 3D. Side-chain stick view of the fusion peptide pocket, with
hydrophobic residues involved labelled and shown with side chains as green sticks.

Supplemental Note 1.

Comparison of the LayV-F fusion peptide (FP) arrangement to other viral fusion peptides highlights the unique nature of the paramyxovirus FP. Whereas a large portion of the LayV-F FP is buried within an interprotomer pocket, the HIV envelope protein (Env), the FP remains largely surface exposed in the closed conformation (Supplemental Figure 4A) (19). The most N-terminal residues of the HIV-1 FP are often unresolved in structures, further highlighting the flexibility of the FP in this Env conformation. During receptor-induced Env opening, the FP begins to be pulled into the core of the protein, and though not buried, it has a somewhat lesser degree of surface exposure (Supplemental Figure 4B) (18). In contrast, the Influenza Hemagglutinin protein (HA), in the pre-fusion state, maintains the N-terminal end of the FP in a hydrophobic cavity at the core of the protein (Supplemental Figure 4C) (16). While the Influenza HA FP is not mobile and flexible as with the HIV-1 FP in the closed state of Env, the Influenza HA FP arrangement does not constitute as true burial, as the FP is inserted into a recess in the structure but not occluded by other residues. The exact nature of the SARS-CoV-2 spike FP is uncertain, as several regions in S2 may play a role. Many structures have identified the stretch of residues directly following the S2' cleavage site as the fusion peptide. This region is not buried, and like the FP of HIV Env, is often poorly resolved in spike structures (Supplemental Figure 4D) (15). However, a recent study (20) indicates that it is residues 866-910 that insert into the host membrane. Close examination of the SARS-CoV-2 sequence reveals that the residues that in the middle of this FP region, which form an internal FP, much more closely matching the glycine and alanine-rich sequences of other known FPs than those directly following the S2' site. This FP arrangement has a much greater degree of similarity in pre-fusion arrangement to paramyxoviruses. The FP region is almost completely buried by the surrounding domain and has limited surface accessibility (Supplemental Figure 4D). While this FP burial is more comparable to paramyxoviruses, the usage of an internal FP loop draws a comparison to the Ebola glycoprotein (GP) FP, also an internal FP. This Ebola FP region is not buried in the GP structure, instead forming the exterior of the fusion subunit. Though it is an internal FP, as with SARS-CoV-2, the fusion loop is positioned between two beta strands rather than two alpha helices (Supplemental Figure 4E) (14).

Introduction

Langya virus (LayV) is a newly identified member of the *Henipavirus* genus, detected after surveillance of disease following animal exposure in eastern China (21). Over the course of several years, LayV infections were identified in 35 individuals, 26 of whom were infected solely with LayV. In these 26 individuals, common symptoms included fever, fatigue, cough, nausea, and headaches. The genome organization of LayV is identical to that of other henipaviruses, including the better known and highly virulent Nipah and Hendra viruses. Based on phylogenetic analysis, LayV is most closely related to Mojiang virus (MojV), which was discovered in rats in southern China (22), and Gamak (GAKV) and Daeryong (DARV) viruses, which were detected in shrews in the Republic of Korea (23).

The henipaviruses infect a range of animals with Nipah and Hendra viruses having their natural reservoir in fruit bats (24). However, several additional *Henipavirus* species have been discovered in recent years with differing animal reservoirs. A survey of both domestic and wild animals revealed a low level of seropositivity for LayV in goats and dogs (2% and 5% respectively) and the highest positivity rate (52.1%) in shrews (*Crocidura lasiura* specifically), indicating that shrews may be the natural reservoir for LayV. The *Henipavirus* genus is a member of the *Paramyxoviridae* family (25), which includes both extremely infectious human pathogens, such as measles and mumps, and extremely deadly pathogens, such as Nipah and Hendra. This diversity of infectivity and virulence, as well as diversity of animal reservoirs are risk factors that necessitate paramyxovirus research. Accordingly, Nipah and Hendra virus are listed as priority pathogens by the WHO (26).

Henipaviruses present two surface glycoproteins known as the attachment and fusion proteins. These proteins work together to mediate viral entry into host cells and in Nipah virus both have been shown to be required for viral entry (27). As these are the sole virus surface-expressed proteins, they are also the primary targets of neutralizing antibodies against henipaviruses. The *Henipavirus* attachment protein has receptor-binding capability and, in Nipah and Hendra viruses, binds to Ephrin B2 and B3, which is found mostly in the brain and in endothelial cells in the heart and lungs (28, 29). The *Henipavirus* fusion (F) protein is a class-I fusion protein that has a metastable pre-fusion conformation which is displayed on virion

surfaces prior to receptor engagement and a post-fusion conformation that is adopted after virus-

cell fusion (30, 31). Proteolytic cleavage splits F into two subunits, F_1 and F_2, still connected by

disulfide linkages (30, 32), freeing a series of hydrophobic residues known as the fusion peptide

to be inserted into the host cell membrane during this conformational conversion. This process

anchors the virus to the host cell and allows the formation of a six-helix bundle in the F protein

to bring the viral and host membranes together, facilitating fusion. Although it has many

mechanistic features in common with other class-I fusion proteins, the F protein in

paramyxoviruses does not appear to have any primary receptor-binding functionality. This role is

fulfilled by a separate attachment glycoprotein, which is believed to undergo conformational

changes during receptor binding which pass a triggering signal to F (33, 34). The structural basis

for this process is poorly understood.

Here, we determine cryo-EM structures of the LayV fusion protein (LayV-F) in pre- and

post-fusion conformations, both at 4.64Å resolution. The LayV-F ectodomain sequence as

reported by Zhang *et al*. (2022) was used (21), replacing the transmembrane and cytoplasmic

domains with purification tags and a trimerization domain. No prefusion stabilizing mutations

were made, nor was the fusion protein cleaved or otherwise knowingly induced to convert to the

post-fusion state. We compare it to other known *Henipavirus* fusion protein structures and

elucidate the structural basis for pre- to post-fusion conversion in both LayV and more broadly

for *Henipavirus* fusion proteins. Our study demonstrates that the highly conserved

paramyxovirus fusion protein architecture is utilized by LayV-F, identifies a region of variability

among *Henipavirus* fusion proteins in an important antigenic site, and provides evidence for a

mechanism to describe how *Henipavirus* fusion proteins are triggered to undergo conformational

changes during the fusion process.

Results

Purification and Structural Determination of LayV-F

We purified the LayV-F ectodomain based on protocols previously developed for NiV-F, by expressing in 293F cells and purifying via StrepTactin (IBA lifesciences) affinity chromatography followed by size exclusion chromatography. Like the NiV-F protein, the LayV F ectodomain elutes with a distinct peak at ~180 kDa. Higher molecular weight species, broader than the main peak, were also observed at ~400 kDa and above (Supplemental Figure 2A). These could be the result of low-affinity interactions between individual trimers. SDS-PAGE analysis of these high molecular weight species in both NiV-F and LayV-F purifications has consistently produced identical bands corresponding to the ~180 kDa species, consistent with the uncleaved F protein (Supplemental Figure 2E). Higher order multimers of Henipavirus F proteins with rings described as "hexamers of trimers" have been previously reported (2). While larger molecular weight species were detected during purification of LayV-F (Supplemental Figure 2A), they comprised a far lower proportion of total protein as compared to purifications of wild-type NiV-F and a prefusion-stabilized NiV-F mutant, NiVop08 (Supplemental Figure 2A) (10). The yield of LayV-F at ~1.2mg/L was considerably higher than that of wild-type NiV-F at ~250µg/L. The NiVop08 yield was vastly higher than the unstabilized NiV-F construct at ~3mg/L, consistent with the general trend of prefusion-stabilized fusion protein constructs expressing more efficiently than their wild-type counterparts (35-38). Differential scanning fluorimetry (DSF) suggested reduced stability of LayV-F compared to NiV-F (Supplemental Figure 2B). The LayV-F DSF profile closely resembles that of wild-type NiV-F, but is left-shifted, indicating that the temperature-induced conformational changes or unfolding events are occurring at a lower temperature for LayV-F.

We determined the structure of the LayV-F ectodomain trimer using cryo-EM. Reference-free 2D classes revealed both pre- and post-fusion conformations of LayV-F in the cryo-EM dataset (Figure 1A-B). After several rounds of classification and sorting we obtained 4.64 Å reconstructions (calculated within cryoSparc according to the FSC 0.143 gold-standard criterion) of the pre-fusion and post-fusion forms of the LayV F protein from approximately 200,000 and 80,000 particles, respectively. Assessment of the three-dimensional FSC (3DFSC) plot for these maps indicated resolutions of 5.42 Å and 5.25 Å for the pre- and post-fusion conformations,

respectively (39). The structures determined revealed pre- and post-fusion architectures that were
similar to those determined for other paramyxoviruses (2-4, 8, 40, 41). The pre-fusion LayV F
structure has a compact "clove-like" shape with a large internal cavity and spans ~110 Å from
the bottom of the stalk to the top of the apex. In comparison, the post-fusion structure is
elongated, spanning ~160 Å, with regions that formed the pre-fusion apex straightening into a
central coiled-coil, ending with the trademark six-helix bundle of the post-fusion conformation.
347
The LayV F ectodomain includes a fusion peptide (FP) and two heptad repeat sequences that
both contribute to three domains, previously described by Yin *et al*. for the parainfluenzavirus 5
(PIV5) F protein (Figure 1C-E) (41). Domain I (DI) is comprised of the most N-terminal residues
after cleavage of the signal peptide, 22-45, and residues 278-369. Its central β-sheet connects
Domain II (DII) to the apical Domain III (DIII). DII, from residues 370-446, makes significant
contact with the cleavage site and fusion peptide of the neighboring protomer before leading to
heptad repeat B (HRB), located at the most C-terminal end of the ectodomain. DIII, from
residues 46-277, contains the cleavage site, fusion peptide, and heptad repeat A (HRA) (Figure
1C-E). Residues 93-140 from DIII and 441-450 from DII, which become linker regions leading
to the repositioned heptad repeats now forming the six-helix bundle, are not resolved in the post-
fusion structure (Figure 1E).
359
Pre- to post-fusion Conversion Mechanism of LayV-F
Though the *Henipavirus* genus was established with the discovery of Hendra virus in 1994
(24, 42), the number of members of the genus has dramatically expanded in recent years with the
discovery not only of Langya virus, but also Gamak and Daeryong virus, both identified in 2021
(23). Within the past three years, Henipaviruses have been identified in an expanded range of
animal hosts and geographic locations. No longer contained to southeast Asia and Australia, new
species have been discovered in Africa and South America, specifically Angavokely virus and
Peixe-Boi virus, respectively (43, 44). Although there is very high structural similarity for fusion
proteins across not only the *Henipavirus* genus, but within the larger *Paramyxoviridae* family,
there tends to be low sequence identity between species, with LayV-F having ~40% sequence
identity with NiV-F, and ~30% or less identity to other non-*Henipavirus* paramyxoviruses
(Supplemental Figure 1). However, with more *Henipavirus* members identified, some conserved

sequence features have become more evident. These include 10 highly conserved cysteines

present in the ectodomain of paramyxovirus F proteins (Figure 2). Identifying the locations of

these cysteines in the pre- and post-fusion LayV-F structures reveals their critical role in

maintaining domain fold integrity during the dramatic conformational changes that occur during

fusion.

Despite the drastically different shapes of the pre- and post-fusion conformations of LayV-F

resulting in a 51.6 Å Cα RMSD, superimposition of the DI and DII domains demonstrates that

these domains remain largely invariant through the pre- to post-fusion transition of the F protein

(Figures 2A-C). Both DI and DII domains have two internal pairs of disulfide bonds that help

maintain the rigidity of these domains (Figure 2B-D). DI shows a pre- to post-fusion Cα RMSD

of 2.16 Å, while DII undergoes more total movement, resulting in pre- to post-fusion Cα RMSD

of 16.5 Å with the linker region from S423 to I446 that leads to HRB extending away from the

domain as HRB moves to its post-fusion location (Figure 2C). Excluding these residues from the

RMSD calculations, DII shifts only 2.3 Å. The difference in conformation of DIII from pre- to

post-fusion is more dramatic. Based on sequence alignment, the F_1-F_2 cleavage site for LayV-F is

expected to be immediately after the highly conserved residue R104 (Supplemental Figure 1),

however cleavage did not occur in this construct (Supplemental Figure 2C-D). The largest

positional shifts during the pre- to post-fusion transition occurred in the fusion peptide and HRA,

which extend and continue the along the path of the apical helix, now forming the central coiled-

coil (Figure 1A). These movements occur around a hinge point in DIII where the apical helix is

connected to the rest of DIII by a disulfide bond, again demonstrating the role of the conserved

cysteines in providing an anchor for large conformational changes in the F protein (Figure 2A).

<u>Fusion triggering Mechanism of *Henipavirus* Fusion proteins</u>

Insertion of the hydrophobic fusion peptide (FP) into the host membrane is an essential step

that leads to the fusion of the virus and host cell membranes. In the pre-fusion LayV-F, a

substantial portion of the FP is buried within a pocket formed between two LayV-F protomers,

requiring the FP to be released from this pocket during the pre-to-post fusion conformational

transition (Figure 3). Hydrophobic residues lining the interprotomer pocket into which the FP is

buried in the pre-fusion LayV-F include L51, L89, M92, L93, V96, V216, and I262 from DIII,

A108, M110, G112, A114, L115, and G116 from the N-terminal end of the FP upstream of the

helical segment from the same protomer, and residues from the adjacent protomer include M292

from DI and P369, F371, A372, L373, G376, V378, I420, L421, and I422 from DII (Figure 3D,

Supplemental figure 5).

Consistent with the close packing of the FP within an interprotomer pocket in the pre-fusion

F conformation, the modeled coordinates exhibit the lowest B-factors at the FP, while HRB, its

DII linker, and surrounding DI region have the highest B-factors in the structure (Figure 3A).

The HRA, HRB and FP undergo dramatic shifts in their positions during pre- to post-fusion

conformational change of the LayV-F protein (Figure 3B). Although the FP is not resolved in the

post-fusion state, based on its location it is expected to show a position shift similar to that of

HRA. The burial of the pre-fusion FP and its interactions with the residues lining the

interprotomer pocket suggest that the purpose of this pocket is to hold the FP in place to prevent

premature conversion to the post-fusion form and therefore is the source of metastability of the

pre-fusion LayV F structure. Consistent with a key role in pre-to post-fusion conformational

dynamics, the FP and the surrounding pocket are among the most conserved regions in

Henipavirus fusion proteins (Figure 3E-F). Release of the FP from this pocket is a requirement

for the conformation transition of the F protein from the pre-fusion to post-fusion conformation

and disruption of this pocket may be the mechanism by which the FP is released to undergo these

changes.

<u>Structural Conservation and Antigenic Diversity of *Henipavirus* Fusion proteins</u>

Despite low overall sequence conservation, the structural similarity of paramyxovirus fusion

proteins is striking. Structures have previously been determined for the F proteins of two

Henipaviruses in the pre-fusion state, NiV-F and HeV-F (2, 3). Despite having only 38%

sequence identity with these viruses, a superimposition of LayV-F with these structures reveals a

conserved architecture (Figure 4A), with Cα RMSDs of 2.64 Å and 3.00 Å for LayV-F to NiV-F

and HeV-F, respectively, with NiV-F and HeV-F differing between each other by 0.94 Å. One

of the few areas between these species where there is a notable shift is in HRA. In the pre-fusion

conformation, the central helix in HRA is broken by a short, two-strand beta sheet with a loop

that extends to the exterior of the protein. As compared to the homologous HRA loop in

NiV/HeV-F, in LayV-F there is a shift of ~9 Å in the direction of the fusion peptide (Figure 4B).

It is unclear if this shift results in any mechanistic effect on the HRA region, although the loop

resides within a site targeted by antibodies and shift in its position may alter antigenicity of the F

protein. (Figure 4C).

An overlay of all experimentally determined structures of paramyxovirus-targeted antibodies

highlights the antigenicity of the HRA loop site and its adjacent regions (Figure 4C). Several

antibodies target DIII in the pre-fusion F near the HRA region and likely interfere with pre- to

post-fusion conversion. In addition to the HRA loop shift described above, LayV-F and other

paramyxovirus species show differences in their surface-exposed residues in these regions that

could define the unique antigenic properties of the different F proteins. These include a

glycosylation site in LayV-F at N279 (Figure 4D), which is positioned between DI and DIII near

several of the previously identified epitopes, specifically 15F, an HeV-neutralizing antibody and

5B3, a cross-reactive NiV/HeV-neutralizing antibody (6).

The sequences of both NiV-F and HeV-F have two plausible glycosylation sites at N64 and

N67, both at the apex, though previous studies have established that only N67 is glycosylated

(45, 46). The N65 glycosylation position in LayV-F is shifted by 8.2 Å away from the apex

residues of HRA. This glycan was particularly visible in the cryo-EM reconstructions of the

post-fusion conformation before local refinement, where the density at roughly the half-height

point of the map corresponds to N65 (Supplemental Figure 3). Interestingly, sequence alignment

reveals that the LayV-F N65 glycosylation site is not homologous to either N64 or N67 of

NiV/HeV, but rather is equivalent to residue 70 (Supplemental Figure 1). Viewing the fusion

protein colored by sequence conservation further highlights the variability of this region (Figure

4E). While there are highly conserved residues at discrete points of the surface, namely the

highly conserved disulfide bond near the apex, the surrounding region ranges from moderately

conserved to highly variable, especially in surface-accessible regions of DIII and the HRA loop.

This variability includes differing residue types between LayV and other henipaviruses,

particularly at the apex site. The differences in the pre-fusion F protein trimer apex are

highlighted by a negatively charged residue, D188, in both NiV and HeV, while the LayV-F

protein instead has polar residues (N183, Q184) and an overall greater hydrophobic character.

This residue arrangement in the pre-fusion LayV-F trimer apex is more similar to that of the

PIV5 pre-fusion F protein (Figure 4F). This change in residue type is likely a source of antigenic

variability between paramyxovirus F proteins. Consistent with this variability, antibody 12B2,

that binds an epitope in pre-fusion NiV-F that includes the trimer apex (6, 11), did not show

binding to LayV-F (supplemental Figure 2F). Therefore, despite similar F protein architecture,

antigenic variability within paramyxoviruses is driven by variability of surface residues.

Discussion

The continuing emergence of new paramyxovirus species necessitates heightened focus on a family that includes both some of the world's deadliest pathogens in Nipah virus, and most infectious in measles and mumps. Members specifically of the *Henipavirus* genus have spread to humans through zoonotic transfer. While Nipah and Hendra viruses are spread to humans through bats, new henipaviruses have been detected in a range of species, such as shrews and rats. The role of animal reservoirs in the SARS-CoV-2 pandemic highlights the risk factor that a diverse range of animal hosts can be, further necessitating efforts to develop treatments or vaccines against henipaviruses. Structural determinations of *Henipavirus* glycoproteins serve as a foundation for future immunogen or therapeutic design.

Though the structure of LayV-F is strikingly similar to other known *Henipavirus* F structures, key differences in surface characteristics at the apex have implications for pan-*Henipavirus* or pan-paramyxovirus vaccine development. When developing vaccinations for a virus that is quickly mutating, or with many distinct strains, as observed with henipaviruses, it is advisable to target relatively invariable sites for immunogen design. Our pre-fusion LayV-F protein structure helps to demonstrate the particularly variable nature of the apex and DIII in *Henipavirus* F proteins, a trait to be considered when selecting antigenic sites to target for a broad immunogenic response.

In contrast to existing structures of paramyxovirus fusion proteins that were determined in either pre- or post-fusion conformation, our study resolves the LayV-F pre- and post-fusion structures within the same cryo-EM dataset, suggesting that for LayV-F, being in solution at ambient temperatures may allow the proteins to sample enough conformational states such that some will lead to conversion. Though paramyxovirus fusion proteins must be cleaved to enable fusion peptide insertion, prior studies have observed uncleaved fusion proteins assuming a post-fusion conformation (47). Our structures reveal a highly stable configuration of the fusion peptide (FP) in the pre-fusion F conformation, buried within an interprotomer pocket, which must be disrupted to transition to the post-fusion conformation. These observations suggest that FP configuration is a source of metastability in the pre-fusion LayV-F protein, where a shift in the position of the FP or the residues surrounding it in the interprotomer pocket too far from its

pre-fusion configuration may lead to irreversible conversion to the post-fusion form. This,

together with the biochemical properties of the FP burial pocket, suggests tight control of FP

positioning is an important part of pre-fusion metastability.

An elusive question regarding paramyxovirus fusion is how the fusion process is first

initiated. For the many virus families that utilize a fusion protein, the metastable pre-fusion state

is maintained prior to a triggering event that initiates conformational changes. In many cases, this

process involves cleavage and removal of entire domains of the fusion protein, such as with the

removal of the S1 subunit from the SARS-CoV-2 spike protein prior to fusion. The

paramyxovirus fusion proteins, however, have no such attachment subunit to remove. Instead,

paramyxoviruses utilize a separate attachment protein, which is thought to pass on a triggering

signal to the fusion protein after receptor binding (33, 34) (Figure 5A). Within the *Henipavirus*

and *Morbilivirus* genera, the attachment and fusion proteins are believed to form complexes prior

to receptor engagement (33) (Figure 5A). While this might suggest a mechanism where the

attachment protein functions as a clamp on the fusion protein, maintaining the pre-fusion state

before receptor binding, studies assaying fusion competency have demonstrated that the fusion

protein alone is not sufficient for virus and host cell fusion, indicating that fusion triggering

involves a more complex mechanism of interaction between the two proteins (9, 48).

The fusion peptide in the pre-fusion LayV-F protein is in a configuration that is conserved

among paramyxoviruses and distinct from other class-I fusion proteins (Supplemental Figure 5,

Supplemental Note 1). It is not yet known which residues are involved in the interaction

between the attachment and fusion proteins, but the critical role of the FP pocket created at the

DII-DIII interface suggests that DII could be the site of the interaction. The triggering signal may

be transferred by way of the attachment protein repositioning DII such that the fusion peptide is

released from the pocket, irreversibly extending towards the apex (Figure 5B). This mechanism

would be consistent with previous paramyxovirus studies that suggested an attachment protein-

binding role for DII (49, 50). Further supporting this hypothesis is the likelihood that the FP in

the pre-fusion conformation is in a high-energy state due to the kink in the helix comprised of the

FP from T119 through HRA up to N148. Such a helix shape is often brought about by the

presence of a proline at the kink, introducing bond-angle limitations that force this shape into the

helix (51). However, in the helix containing the FP, no proline is present. The bent shape could

be induced by the FP being buried within and locked into the hydrophobic FP pocket. When

disrupted, the position of the FP would therefore move rapidly as the tension in the helix is

resolved. Though it could be hypothesized that cleavage of the FP is the trigger that allows for

rearrangement, our study, in line with others, has established that cleavage is not required for

conformational conversion in paramyxovirus fusion proteins. The existence of a high-energy

state FP helix, held in place by an interprotomer DII, which is then disrupted by the action of an

attachment protein offers a logical explanation for how *Henipavirus* attachment proteins activate

their conjugate fusion proteins, and establishes the basis for how paramyxovirus fusion proteins

maintain their metastability in the pre-fusion state.

Materials and Methods

Data Availability. Cryo-EM reconstructions and atomic models generated during this study are available at wwPDB and EMBD (https://www.rcsb.org; http://emsearch.rutgers.edu) under the accession codes PDB IDs 8FEJ and 8FEL and EMDB IDs EMD-29029 and EMD-29032.

Plasmids. Gene synthesis for all plasmids generated by this study were performed and the sequence confirmed by GeneImmune Biotechnology (Rockville, MD). The fusion protein ectodomain constructs included F protein residues 1 to 438 (GenBank: UUV47205.1), a C-terminal T4 fibritin trimerization motif (FOLDON), a C-terminal HRV3C protease cleavage site, a TwinStrepTag and an 8XHisTag. The ectodomain was cloned into the mammalian expression vector pαH. Synthetic heavy and light chain variable domain genes for Fabs were cloned into a modified pVRC8400 expression vector, as previously described (5, 24, 25). Antibody variable regions for both heavy and light chain replaced the variable regions from the VRC01 antibody plasmid (52), with the IL-2 secretion signal and human kappa constant domain being used for light chain plasmids. All plasmids have been deposited to Addgene (https://www.addgene.org) under the codes XXX.

Cell culture and protein expression. For F ectodomains, GIBCO FreeStyle 293-F cells (embryonal, human kidney) were maintained at 37°C and 9% CO_2 in a 75% humidified atmosphere in FreeStyle 293 Expression Medium (GIBCO). The Plasmid was transiently transfected using Turbo293 (SpeedBiosystems) and incubated at 37°C, 9% CO2, 75% humidity with agitation at 120 rpm for 6 days. On the day following transfection, HyClone CDM4HEK293 media (Cytiva, MA) was added to the cells. For antibodies, GIBCO Expi293F cells (embryonal, human kidney) were maintained at 37°C and 9% CO_2 in a 75% humidified atmosphere in Expi293 Expression Medium (GIBCO). The Plasmid was transiently transfected using Expifectamine 293 (GIBCO) and incubated at 37°C, 9% CO2, 75% humidity with agitation at 120 rpm for 6 days. On the day following transfection, Expifectamine 293 Enhancers (GIBCO) were added to the cells.

Protein purification. On the 6th day post transfection, the fusion protein ectodomains were harvested from the concentrated supernatant, purified using StrepTactin resin (IBA LifeSciences) and size exclusion chromatography (SEC) using a Superose 6 10/300 GL Increase column (Cytiva, MA) equilibrated in PBS buffer (Thermo Scientific 137mM NaCl, 2.7mM KCl, pH 7.4, 10 mM Phosphate Buffer, 1.8mM Potassium Phosphate Monobasic). Antibodies were purified using Protein A affinity (Thermo Scientific) and SEC using a Superose 6 10/300 GL Increase column (Cytiva, MA) equilibrated in PBS buffer (Thermo Scientific 137mM NaCl, 2.7mM KCl, pH 7.4, 10 mM Phosphate Buffer, 1.8mM Potassium Phosphate Monobasic). All steps of the purification were performed at room temperature and in a single day. Protein quality was assessed by SDS-Page using NuPage 4-12% (Invitrogen, CA). The purified proteins were flash frozen and stored at -80 °C in single-use aliquots. Each aliquot was thawed by a 5-minute incubation at 37 °C before use.

Cryo-EM. Purified ectodomain was diluted to a concentration of 1.5 mg/mL in PBS (described above) with 0.0005% DDS and 0.38% glycerol added. A 2.2μL drop of protein was deposited on a Quantifoil R1.2/1.3 grid (Electron Microscopy Sciences, PA) that had been glow discharged for 10 seconds using a PELCO easiGlow™ Glow Discharge Cleaning System. After a 30-second incubation in >95% humidity, excess protein was blotted away for 2.5 seconds before being

plunge frozen into liquid ethane using a Leica EM GP2 plunge freezer (Leica Microsystems).
Frozen grids were imaged using a Titan Krios (Thermo Fisher) equipped with a K3 detector
(Gatan). Movie frame alignment was carried out using UNBLUR (53).The cryoSPARC (12)
software was used for data processing. Phenix (54), PyMOL(55), ChimeraX (56) and Isolde (57)
were used for model building and refinement. Phenix was first used to fit the initial models into
maps. ISOLDE was used to manually adjust residues to address rotamer and Ramachandran
outliers. Phenix was again used to perform real-space refinement, energy-minimization of side-
chain positions, and to set B-factors during an ADP-only real-space refinement.
639
SDS-PAGE. Prepared 1, 3, or 8µg of sample with Laemmli sample buffer (BioRad), PBS buffer,
and with or without 300mM DTT (Reduced/Non-Reduced). Loaded to NuPAGE 4-12% Bis-Tris
gel (Invitrogen) and ran at 175V with MES-SDS running buffer until complete. Stained with
Coomassie blue (Novex) for 30 minutes before water destain and imaging.

Differential scanning fluorimetry. DSF assay was performed using Tycho NT. 6 (NanoTemper
Technologies). Spike ectodomains were diluted to approximately 0.15 mg/mL. Intrinsic
fluorescence was measured at 330 nm and 350 nm while the sample was heated from 35 to 95 °C
at a rate of 30°C/min. The ratio of fluorescence (350/330 nm) and inflection temperatures (T_i)
were calculated by the Tycho NT. 6 apparatus.

Difference distance matrices (DDM). DDM plots were generated using the Bio3D package
(Grant *et al*., 2021) implemented in R (R Core Team, 2018. R: A language and environment for
statistical computing. R Foundation for Statistical Computing, Vienna, Austria. http://www.R-
project.org/).

Bio-layer Interferometry (BLI). Antibody binding to *Henipavirus* F proteins was assessed using
BLI on an Octet RED 384 (Sartorius, formerly ForteBio) with Forte Bio Kinetics Buffer. All
binding assays were performed at 30°C. Antibodies were immobilized on Anti-Human Fc Capture
tips (ForteBio), loaded at 20µg/mL for 300s. F proteins were used as analytes at 100nM, with an
association time of 480s and dissociation time of 600s. Sensorgram data were reference-subtracted
and analyzed using the Octet Analysis Studio software (Sartorius), with a reference tip for each
immobilized antibody and reference sample for each analyte.

Sequence Alignment. Paramyxovirus F sequences aligned with Clustal Omega tool (58). Output
alignment was colored and assigned consensus residue types by MVeiw (1). Sequences obtained
from either Uniprot or Genbank. Uniprot accession numbers: NiV-F: Q9IH63, HeV-F: O89342,
hPIV3-F: P06828, PIV5-F: P04849, MeV-F (ICB Strain): Q786F3, MojV-F: W8SKT3, CedV-F:
J7GX38, hPIV1-F: P12605, NDV-F (Texas Strain): P26628, SeV-F (Z Strain): P04855. Genbank
accession numbers: LayV-F: UUV47205.1, GakV-F: QYO90524.1, DarV-F: QYO90531.1.

References

672   1.   Brown NP, Leroy C, Sander C. 1998. MView: a web-compatible database search or
673   multiple alignment viewer. Bioinformatics 14:380-1.
674   2.   Xu K, Chan Y-P, Bradel-Tretheway B, Akyol-Ataman Z, Zhu Y, Dutta S, Yan L, Feng
675   Y, Wang L-F, Skiniotis G, Lee B, Zhou ZH, Broder CC, Aguilar HC, Nikolov DB. 2015.
676   Crystal Structure of the Pre-fusion Nipah Virus Fusion Glycoprotein Reveals a Novel
677   Hexamer-of-Trimers Assembly. PLOS Pathogens 11:e1005322.
678   3.   Wong JJW, Paterson RG, Lamb RA, Jardetzky TS. 2016. Structure and stabilization of
679   the Hendra virus F glycoprotein in its prefusion form. Proceedings of the National
680   Academy of Sciences 113:1056-1061.
681   4.   Stewart-Jones GBE, Chuang G-Y, Xu K, Zhou T, Acharya P, Tsybovsky Y, Ou L, Zhang
682   B, Fernandez-Rodriguez B, Gilardi V, Silacci-Fregni C, Beltramello M, Baxa U, Druz A,
683   Kong W-P, Thomas PV, Yang Y, Foulds KE, Todd J-P, Wei H, Salazar AM, Scorpio
684   DG, Carragher B, Potter CS, Corti D, Mascola JR, Lanzavecchia A, Kwong PD. 2018.
685   Structure-based design of a quadrivalent fusion glycoprotein vaccine for human
686   parainfluenza virus types 1–4. Proceedings of the National Academy of Sciences
687   115:12265-12270.
688   5.   Avanzato VA, Oguntuyo KY, Escalera-Zamudio M, Gutierrez B, Golden M, Kosakovsky
689   Pond SL, Pryce R, Walter TS, Seow J, Doores KJ, Pybus OG, Munster VJ, Lee B,
690   Bowden TA. 2019. A structural basis for antibody-mediated neutralization of Nipah virus
691   reveals a site of vulnerability at the fusion glycoprotein apex. Proceedings of the National
692   Academy of Sciences 116:25057-25067.
693   6.   Dang HV, Cross RW, Borisevich V, Bornholdt ZA, West BR, Chan Y-P, Mire CE, Da
694   Silva SC, Dimitrov AS, Yan L, Amaya M, Navaratnarajah CK, Zeitlin L, Geisbert TW,
695   Broder CC, Veesler D. 2021. Broadly neutralizing antibody cocktails targeting Nipah
696   virus and Hendra virus fusion glycoproteins. Nature Structural & Molecular Biology
697   28:426-434.
698   7.   Pei J, Grishin NV. 2001. AL2CO: calculation of positional conservation in a protein
699   sequence alignment. Bioinformatics 17:700-12.
700   8.   Welch BD, Liu Y, Kors CA, Leser GP, Jardetzky TS, Lamb RA. 2012. Structure of the
701   cleavage-activated prefusion form of the parainfluenza virus 5 fusion protein.
702   Proceedings of the National Academy of Sciences 109:16672-16677.
703   9.   Liu Q, Stone JA, Bradel-Tretheway B, Dabundo J, Benavides Montano JA, Santos-
704   Montanez J, Biering SB, Nicola AV, Iorio RM, Lu X, Aguilar HC. 2013. Unraveling a
705   Three-Step Spatiotemporal Mechanism of Triggering of Receptor-Induced Nipah Virus
706   Fusion and Cell Entry. PLoS Pathogens 9:e1003770.
707   10.   Loomis RJ, Stewart-Jones GBE, Tsybovsky Y, Caringal RT, Morabito KM, McLellan JS,
708   Chamberlain AL, Nugent ST, Hutchinson GB, Kueltzo LA, Mascola JR, Graham BS.
709   2020. Structure-Based Design of Nipah Virus Vaccines: A Generalizable Approach to
710   Paramyxovirus Immunogen Development. Frontiers in Immunology 11.
711   11.   Dang HV, Chan Y-P, Park Y-J, Snijder J, Da Silva SC, Vu B, Yan L, Feng Y-R, Rockx
712   B, Geisbert TW, Mire CE, Broder CC, Veesler D. 2019. An antibody against the F
713   glycoprotein inhibits Nipah and Hendra virus infections. Nature Structural &
714   Molecular Biology 26:980-987.
715   12.   Punjani A, Rubinstein JL, Fleet DJ, Brubaker MA. 2017. cryoSPARC: algorithms for
716   rapid unsupervised cryo-EM structure determination. Nature Methods 14:290-296.

717   13.   Punjani A, Zhang H, Fleet DJ. 2020. Non-uniform refinement: adaptive regularization
718   improves single-particle cryo-EM reconstruction. Nature Methods 17:1214-1221.
719   14.   Lee JE, Fusco ML, Hessell AJ, Oswald WB, Burton DR, Saphire EO. 2008. Structure of
720   the Ebola virus glycoprotein bound to an antibody from a human survivor. Nature
721   454:177-182.
722   15.   Cai Y, Zhang J, Xiao T, Peng H, Sterling SM, Walsh RM, Rawson S, Rits-Volloch S,
723   Chen B. 2020. Distinct conformational states of SARS-CoV-2 spike protein. Science
724   369:1586-1592.
725   16.   Benton DJ, Nans A, Calder LJ, Turner J, Neu U, Lin YP, Ketelaars E, Kallewaard NL,
726   Corti D, Lanzavecchia A, Gamblin SJ, Rosenthal PB, Skehel JJ. 2018. Influenza
727   hemagglutinin membrane anchor. Proceedings of the National Academy of Sciences
728   115:10112-10117.
729   17.   Ozorowski G, Pallesen J, de Val N, Lyumkis D, Cottrell CA, Torres JL, Copps J,
730   Stanfield RL, Cupo A, Pugach P, Moore JP, Wilson IA, Ward AB. 2017. Open and
731   closed structures reveal allostery and pliability in the HIV-1 envelope spike. Nature
732   547:360-363.
733   18.   Kong R, Xu K, Zhou T, Acharya P, Lemmin T, Liu K, Ozorowski G, Soto C, Taft JD,
734   Bailer RT, Cale EM, Chen L, Choi CW, Chuang G-Y, Doria-Rose NA, Druz A, Georgiev
735   IS, Gorman J, Huang J, Joyce MG, Louder MK, Ma X, McKee K, O'Dell S, Pancera M,
736   Yang Y, Blanchard SC, Mothes W, Burton DR, Koff WC, Connors M, Ward AB, Kwong
737   PD, Mascola JR. 2016. Fusion peptide of HIV-1 as a site of vulnerability to neutralizing
738   antibody. Science 352:828-833.
739   19.   Pancera M, Zhou T, Druz A, Georgiev IS, Soto C, Gorman J, Huang J, Acharya P,
740   Chuang G-Y, Ofek G, Stewart-Jones GBE, Stuckey J, Bailer RT, Joyce MG, Louder MK,
741   Tumba N, Yang Y, Zhang B, Cohen MS, Haynes BF, Mascola JR, Morris L, Munro JB,
742   Blanchard SC, Mothes W, Connors M, Kwong PD. 2014. Structure and immune
743   recognition of trimeric pre-fusion HIV-1 Env. Nature 514:455-461.
744   20.   Shi W, Cai Y, Zhu H, Peng H, Voyer J, Rits-Volloch S, Cao H, Mayer ML, Song K, Xu
745   C, Lu J, Zhang J, Chen B. 2022. Cryo-EM structure of SARS-CoV-2 postfusion spike in
746   membrane. bioRxiv doi:10.1101/2022.12.05.519151:2022.12.05.519151.
747   21.   Zhang X-A, Li H, Jiang F-C, Zhu F, Zhang Y-F, Chen J-J, Tan C-W, Anderson DE, Fan
748   H, Dong L-Y, Li C, Zhang P-H, Li Y, Ding H, Fang L-Q, Wang L-F, Liu W. 2022. A
749   Zoonotic Henipavirus in Febrile Patients in China. New England Journal of Medicine
750   387:470-472.
751   22.   Wu Z, Yang L, Yang F, Ren X, Jiang J, Dong J, Sun L, Zhu Y, Zhou H, Jin Q. 2014.
752   Novel Henipa-like virus, Mojiang Paramyxovirus, in rats, China, 2012. Emerg Infect Dis
753   20:1064-6.
754   23.   Lee S-H, Kim K, Kim J, No JS, Park K, Budhathoki S, Lee SH, Lee J, Cho SH, Cho S,
755   Lee G-Y, Hwang J, Kim H-C, Klein TA, Uhm C-S, Kim W-K, Song J-W. 2021.
756   Discovery and Genetic Characterization of Novel Paramyxoviruses Related to the Genus
757   Henipavirus in Crocidura Species in the Republic of Korea. Viruses 13:2020.
758   24.   Field H, Young P, Yob JM, Mills J, Hall L, Mackenzie J. 2001. The natural history of
759   Hendra and Nipah viruses. Microbes Infect 3:307-14.
760   25.   Rima B, Balkema-Buschmann A, Dundon WG, Duprex P, Easton A, Fouchier R, Kurath
761   G, Lamb R, Lee B, Rota P, Wang L, Ictv Report C. 2019. ICTV Virus Taxonomy Profile:
762   Paramyxoviridae. J Gen Virol 100:1593-1594.

763   26.   Anonymous. Prioritizing diseases for research and development in emergency contexts,
764   *on* WHO. https://www.who.int/activities/prioritizing-diseases-for-research-and-
765   development-in-emergency-contexts. Accessed 2/28/23.

766   27.   Aguilar HC, Iorio RM. 2012. Henipavirus Membrane Fusion and Viral Entry, p 79-94. *In*
767   Lee B, Rota PA (ed), Henipavirus: Ecology, Molecular Virology, and Pathogenesis
768   doi:10.1007/82_2012_200. Springer Berlin Heidelberg, Berlin, Heidelberg.

769   28.   Negrete OA, Levroney EL, Aguilar HC, Bertolotti-Ciarlet A, Nazarian R, Tajyar S, Lee
770   B. 2005. EphrinB2 is the entry receptor for Nipah virus, an emergent deadly
771   paramyxovirus. Nature 436:401-5.

772   29.   Xu K, Rajashankar KR, Chan Y-P, Himanen JP, Broder CC, Nikolov DB. 2008. Host cell
773   recognition by the henipaviruses: Crystal structures of the Nipah G attachment
774   glycoprotein and its complex with ephrin-B3. Proceedings of the National Academy of
775   Sciences 105:9953-9958.

776   30.   White JM, Delos SE, Brecher M, Schornberg K. 2008. Structures and mechanisms of
777   viral membrane fusion proteins: multiple variations on a common theme. Crit Rev
778   Biochem Mol Biol 43:189-219.

779   31.   Lamb RA, Jardetzky TS. 2007. Structural basis of viral invasion: lessons from
780   paramyxovirus F. Current Opinion in Structural Biology 17:427-436.

781   32.   Pager CT, Dutch RE. 2005. Cathepsin L is involved in proteolytic processing of the
782   Hendra virus fusion protein. J Virol 79:12714-20.

783   33.   Bose S, Jardetzky TS, Lamb RA. 2015. Timing is everything: Fine-tuned molecular
784   machines orchestrate paramyxovirus entry. Virology 479-480:518-31.

785   34.   Iorio RM, Melanson VR, Mahon PJ. 2009. Glycoprotein interactions in paramyxovirus
786   fusion. Future Virol 4:335-351.

787   35.   Hsieh C-L, Goldsmith JA, Schaub JM, DiVenere AM, Kuo H-C, Javanmardi K, Le KC,
788   Wrapp D, Lee AG, Liu Y, Chou C-W, Byrne PO, Hjorth CK, Johnson NV, Ludes-
789   Meyers J, Nguyen AW, Park J, Wang N, Amengor D, Lavinder JJ, Ippolito GC, Maynard
790   JA, Finkelstein IJ, McLellan JS. 2020. Structure-based design of prefusion-stabilized
791   SARS-CoV-2 spikes. Science 369:1501-1505.

792   36.   Wrapp D, Wang N, Corbett KS, Goldsmith JA, Hsieh C-L, Abiona O, Graham BS,
793   McLellan JS. 2020. Cryo-EM structure of the 2019-nCoV spike in the prefusion
794   conformation. Science 367:1260-1263.

795   37.   Do Kwon Y, Pancera M, Acharya P, Georgiev IS, Crooks ET, Gorman J, Joyce MG,
796   Guttman M, Ma X, Narpala S, Soto C, Terry DS, Yang Y, Zhou T, Ahlsen G, Bailer RT,
797   Chambers M, Chuang G-Y, Doria-Rose NA, Druz A, Hallen MA, Harned A, Kirys T,
798   Louder MK, O'Dell S, Ofek G, Osawa K, Prabhakaran M, Sastry M, Stewart-Jones GBE,
799   Stuckey J, Thomas PV, Tittley T, Williams C, Zhang B, Zhao H, Zhou Z, Donald BR,
800   Lee LK, Zolla-Pazner S, Baxa U, Schön A, Freire E, Shapiro L, Lee KK, Arthos J,
801   Munro JB, Blanchard SC, Mothes W, Binley JM, et al. 2015. Crystal structure,
802   conformational fixation and entry-related interactions of mature ligand-free HIV-1 Env.
803   Nature Structural & Molecular Biology 22:522-531.

804   38.   Rawi R, Rutten L, Lai Y-T, Olia AS, Blokland S, Juraszek J, Shen C-H, Tsybovsky Y,
805   Verardi R, Yang Y, Zhang B, Zhou T, Chuang G-Y, Kwong PD, Langedijk JPM. 2020.
806   Automated Design by Structure-Based Stabilization and Consensus Repair to Achieve
807   Prefusion-Closed Envelope Trimers in a Wide Variety of HIV Strains. Cell Reports
808   33:108432.

39. Tan YZ, Baldwin PR, Davis JH, Williamson JR, Potter CS, Carragher B, Lyumkis D.
2017. Addressing preferred specimen orientation in single-particle cryo-EM through
tilting. Nature Methods 14:793-796.
40. Hashiguchi T, Fukuda Y, Matsuoka R, Kuroda D, Kubota M, Shirogane Y, Watanabe S,
Tsumoto K, Kohda D, Plemper RK, Yanagi Y. 2018. Structures of the prefusion form of
measles virus fusion protein in complex with inhibitors. Proceedings of the National
Academy of Sciences 115:2496-2501.
41. Yin HS, Wen X, Paterson RG, Lamb RA, Jardetzky TS. 2006. Structure of the
parainfluenza virus 5 F protein in its metastable, prefusion conformation. Nature 439:38-
44.
42. Murray K, Selleck P, Hooper P, Hyatt A, Gould A, Gleeson L, Westbury H, Hiley L,
Selvey L, Rodwell B, Ketterer P. 1995. A Morbillivirus that Caused Fatal Fisease in
Horses and Humans. Science 268:94-97.
43. Madera S, Kistler A, Ranaivoson HC, Ahyong V, Andrianiaina A, Andry S, Raharinosy
V, Randriambolamanantsoa TH, Ravelomanantsoa NAF, Tato CM, DeRisi JL, Aguilar
HC, Lacoste V, Dussart P, Heraud J-M, Brook CE. 2022. Discovery and Genomic
Characterization of a Novel Henipavirus, Angavokely Virus, from Fruit Bats in
Madagascar. Journal of Virology 96:e00921-22.
44. Hernández LHA, da Paz TYB, Silva SPd, Silva FSd, Barros BCVd, Nunes BTD, Casseb
LMN, Medeiros DBA, Vasconcelos PFdC, Cruz ACR. 2022. First Genomic Evidence of
a Henipa-like Virus in Brazil. Viruses 14:2167.
45. Aguilar HC, Matreyek KA, Filone CM, Hashimi ST, Levroney EL, Negrete OA,
Bertolotti-Ciarlet A, Choi DY, McHardy I, Fulcher JA, Su SV, Wolf MC, Kohatsu L,
Baum LG, Lee B. 2006. N-glycans on Nipah virus fusion protein protect against
neutralization but reduce membrane fusion and viral entry. J Virol 80:4878-89.
46. Carter JR, Pager CT, Fowler SD, Dutch RE. 2005. Role of N-linked glycosylation of the
Hendra virus fusion protein. J Virol 79:7922-5.
47. Yin H-S, Paterson RG, Wen X, Lamb RA, Jardetzky TS. 2005. Structure of the
uncleaved ectodomain of the paramyxovirus (hPIV3) fusion protein. Proceedings of the
National Academy of Sciences 102:9288-9293.
48. Yeo YY, Buchholz DW, Gamble A, Jager M, Aguilar HC. 2021. Headless Henipaviral
Receptor Binding Glycoproteins Reveal Fusion Modulation by the Head/Stalk Interface
and Post-receptor Binding Contributions of the Head Domain. Journal of Virology
95:e00666-21.
49. Bose S, Heath CM, Shah PA, Alayyoubi M, Jardetzky TS, Lamb RA. 2013. Mutations in
the parainfluenza virus 5 fusion protein reveal domains important for fusion triggering
and metastability. J Virol 87:13520-31.
50. Poor TA, Jones LM, Sood A, Leser GP, Plasencia MD, Rempel DL, Jardetzky TS,
Woods RJ, Gross ML, Lamb RA. 2014. Probing the paramyxovirus fusion (F) protein-
refolding event from pre- to postfusion by oxidative footprinting. Proceedings of the
National Academy of Sciences 111:E2596-E2605.
51. Wilman HR, Shi J, Deane CM. 2014. Helix kinks are equally prevalent in soluble and
membrane proteins. Proteins 82:1960-70.
52. McCarthy KR, Lee J, Watanabe A, Kuraoka M, Robinson-McCarthy LR, Georgiou G,
Kelsoe G, Harrison SC. 2021. A Prevalent Focused Human Antibody Response to the
Influenza Virus Hemagglutinin Head Interface. mBio 12:e01144-21.

53.   Grant T, Grigorieff N. 2015. Measuring the optimal exposure for single particle cryo-EM
856   using a 2.6 Å reconstruction of rotavirus VP6. Elife 4:e06980.
857   54.   Liebschner D, Afonine PV, Baker ML, Bunkoczi G, Chen VB, Croll TI, Hintze B, Hung
858   L-W, Jain S, McCoy AJ, Moriarty NW, Oeffner RD, Poon BK, Prisant MG, Read RJ,
859   Richardson JS, Richardson DC, Sammito MD, Sobolev OV, Stockwell DH, Terwilliger
860   TC, Urzhumtsev AG, Videau LL, Williams CJ, Adams PD. 2019. Macromolecular
861   structure determination using X-rays, neutrons and electrons: recent developments in
862   Phenix. Acta Crystallographica Section D 75:861-877.
863   55.   Anonymous. The PyMOL Molecular Graphics System, v2.0. Schrödinger, LLC,
864   56.   Pettersen EF, Goddard TD, Huang CC, Meng EC, Couch GS, Croll TI, Morris JH, Ferrin
865   TE. 2021. UCSF ChimeraX: Structure visualization for researchers, educators, and
866   developers. Protein Sci 30:70-82.
867   57.   Croll T. 2018. ISOLDE: a physically realistic environment for model building into low-
868   resolution electron-density maps. Acta Crystallographica Section D 74:519-530.
869   58.   Madeira F, Pearce M, Tivey ARN, Basutkar P, Lee J, Edbali O, Madhusoodanan N,
870   Kolesnikov A, Lopez R. 2022. Search and sequence analysis tools services from EMBL-
871   EBI in 2022. Nucleic acids research doi:10.1093/nar/gkac240:gkac240.

Acknowledgments

Cryo-EM data were collected at the Duke Krios at the Duke University Shared Materials Instrumentation Facility (SMIF), a member of the North Carolina Research Triangle Nanotechnology Network (RTNN), which is supported by the National Science Foundation (award number ECCS-2025064) as part of the National Nanotechnology Coordinated Infrastructure (NNCI). This study utilized the computational resources offered by Duke Research Computing (http://rc.duke.edu; NIH 1S10OD018164-01) at Duke University. This work was supported by funding from the Translating Duke Health Initiative. The comparative analysis of diverse Class I fusion peptides was supported in part by funding from NIAID grant R01 AI145687, R01 AI165947 and U54 AI170752. Figure 5 was made using BioRender.

Figure 1: Structures of LayV-F in pre- and post-fusion conformation:

A-B. Cryo-EM reconstructions of LayV-F in **A.** pre- and **B.** post-fusion conformation, colored by protomer. Map regions colored in different shades of blue (pre-fusion) and yellow (post-fusion) for each protomer. Representative reference-free 2D classes used for the reconstruction are displayed above each map. **C-D.** Cartoon representation of atomic models of LayV-F in **C.** pre-fusion and **D.** post-fusion conformation that were built into the maps shown in **A-B**. In each, one protomer is colored by domain, based on the color scheme in panel **E**, and the other two protomers colored gray. **E.** Sequence domain key for LayV-F. SS: Secretion signal, cleaved prior to purification. DI, DII, DIII: Domains I, II, and III. FP: Fusion Peptide. HRA, HRB: Heptad repeats A and B. Foldon: Trimerization domain. TS: Twin-Strep tag, used for purification with modified streptavidin resin.

Figure 2: Role of conserved cysteines during conformational conversion.

A. Zoomed-in view of the DIII domain, including the fusion peptide (orange) and HRA (yellow), shown for the (left) pre- and (right) post-fusion structures. Color scheme continued from Figure 1E. The cleavage site at R104 is identified, as well as the disulfide bond formed by C66 and C187 (yellow sticks). This disulfide bond is a hinge point around which the HRA straightens into the central coiled-coil in the post-fusion F structure. **B-C.** Superimposition of pre- and post-fusion conformation domains DI (**B**) and DII (**C**). Color scheme continued from figure 1E, with the post-fusion conformation colored in a lighter shade. Disulfide bonds are shown as yellow sticks and labelled. For DII, the movement of the HRB linker region is indicated by an arrow. **D.** Sequence alignment of paramyxovirus fusion proteins in the regions surrounding conserved cysteines, with cysteine residue number and domain labelled. Viruses colored blue are from the *Henipavirus* genus and colored tan for other paramyxoviruses. Residues are colored by type, per MView standard (1). Blue: Alcohol, Green: Hydrophobic, Dark Blue: Negative charge, Red: Positive charge, Purple: Polar, Yellow: Cysteine.

Figure 3: Fusion peptide burial and conservation.

A. Structure of the pre-fusion LayV-F protein shown in cartoon representation and colored by B-factors. **B.** Difference Distance Matrix plot comparing relative residue positions between pre-fusion and post-fusion conformation. Sequence key from figure 1E included along each axis. Gray areas in the bottom right half of the plot are the result of sequences not present in the post-fusion model. **C.** Hydrophobic fusion peptide pocket in pre-fusion state, with fusion peptide depicted as orange cartoon, and surface electrostatics view of the pocket. **D.** Side-chain stick view of the fusion peptide pocket, with hydrophobic residues involved shown with side chains as green sticks. An enlarged view of this panel with residues labeled is seen in supplemental figure S5. **E.** Cartoon representations of a single pre-fusion state protomer, colored by sequence conservation across *Henipavirus* fusion proteins. Color scale generated by ChimeraX default AL2CO entropy measure. **F.** Zoom in of the fusion peptide pocket, colored by sequence conservation as in H. The fusion peptide pocket is among the most strongly conserved region of the protein.

Figure 4: Structure and antigenicity comparison of Henipavirus fusion proteins.

A. Superimposition of a single pre-fusion protomer of LayV-F (blue) NiV-F (cyan) and HeV-F (green), in cartoon representation. PDBs: LayV-F: 8FEJ, NiV-F: 5EVM, HeV-F: 5EJB (2, 3). **B.** Zoomed in view of a loop within the folded HRA, one of the few areas of significant structural difference between LayV-F and NiV/HeV-F. **C.** Binding site of known paramyxovirus F protein-neutralizing antibodies, labeled by the species targeted. NiV-F trimer from 12B2-bound structure shown as cartoon and antibody Fabs shown in surface view. Only one Fab per type shown, all expect PIA174 bind with 3:1 stoichiometry. PDBs: PIA174: 6MJZ, 12B2: 7KI4, mAb66: 6T3F, 1F5: 7KI6, 5B3: 6TYS (4-6). **D.** Surface view of LayV-F and NiV-F with glycosylation sites (residues meeting the N-X-S/T glycosylation sequence) colored teal. **E.** Surface representation of LayV-F colored by sequence conservation across *Henipavirus* fusion proteins. Color scale generated by ChimeraX default AL2CO entropy measure; magenta indicates high conservation and teal indicates high variability (7). **F.** Surface representation of paramyxovirus fusion proteins, showing only main chain atoms, colored by residue type and shown from top-down view. Tan: hydrophobic, Pink: polar, Red: negative, Blue: positive, teal: Potential glycosylation. PIV5 PBD: 4GIP (8).

Figure 5: Fusion triggering hypothesis for paramyxoviruses.

A. Existing model of *Henipavirus* (HNV) fusion protein triggering based on Liu *et al.* (2013) (9). HNV-G, brown, shown in a two-receptor binding domain down "closed" state, bound to HNV-F in pre-fusion conformation. Exposure to Ephrin B2/B3 receptor induces opening of HNV-G, triggering conversion of HNV-F from pre-fusion to the pre-hairpin state. The triggering interaction is labeled with a star. **B.** Hypothesized structural basis for the G-F interaction shown in **A**, wherein a two-step mechanism involving disruption of the fusion peptide burial pocket through action of the HNV-G stalk on DII frees the fusion peptide, allowing conformational conversion.