Spatial autocorrelation dimension as a potential determinant for the temporal persistence of human atrial and ventricular fibrillation.

Dhani Dharmaprani[1,2], Evan V. Jenkins[1,2], Jing X. Quah[2], Kathryn Tiver[2], Lewis Mitchell[3], Matthew Tung[4], Waheed Ahmad[5], Nik Stoyanov[6], Martin Aguilar[7], Martyn P. Nash[8], Richard H. Clayton[9], Stanley Nattel[8,10], Anand N. Ganesan[1,2*].

1. College of Science and Engineering, Flinders University, Adelaide, Australia
2. Department of Cardiovascular Medicine, Flinders Medical Centre, Australia
3. School of Mathematical Sciences, University of Adelaide, Australia
4. Department of Cardiovascular Medicine, Sunshine Coast University Hospital, Birtinya, Queensland, Australia
5. Department of Cardiovascular Medicine, Princess Alexandra Hospital, Queensland, Australia
6. Department of Cardiology, Fiona Stanley Hospital, Perth, Western Australia, Australia
7. The Department of Medicine and Research Center, Montréal Heart Institute
8. Auckland Bioengineering Institute, University of Auckland, New Zealand
9. Insigneo Institute for in-silico Medicine and Department of Computer Science, University of Sheffield, UK
10. Université de Montréal, Montréal, QC, Canada; Pharmacology Institute, University Duisburg-Essen, Germany; CHU Liryc Institute, Bordeaux, France

Correspondence:

Professor Anand Ganesan

Flinders University and Flinders Medical Centre, Adelaide, Australia

anand.ganesan@flinders.edu.au

Word count: 4284 (including introduction, methods, results, discussion, and figure legends)

Supplementary information is provided for this manuscript.

Acknowledgements and funding: We appreciate the comments from Prof. Emeritus Henry Greenside of Duke University, North Carolina, that helped to improve our paper. This work is supported by the National Health and Medical Research Council of Australia Project Grant (1063754) and National Heart Foundation of Australia (101188).

Disclosures: Authors confirm they do not have any disclosures.

ABSTRACT

Background: Despite being central to atrial fibrillation (AF) and ventricular fibrillation (VF) mechanisms and therapy, the factors governing AF and VF termination are poorly understood. It has been noted that system size (L) and the two-point spatial correlation length (ξ_2) link to time until termination in some spatiotemporally chaotic systems, but the relationship between these determinants has not been studied in spontaneously terminating AF/VF.

Objective: We assess whether the time to cardiac fibrillation termination can be predicted using a newly developed metric referred to as the '*spatial autocorrelation dimension*' (D_i), given by the ratio of L and ξ_2, in human AF/VF.

Methods: D_i was computed and compared in a multi-centre, multi-system study with data for sustained versus spontaneously terminating human AF/VF. VF data was collected during coronary-bypass surgery; AF data during clinically indicated AF ablation. We analyzed: i) VF mapped using a 256-electrode epicardial sock (n=12pts); ii) AF mapped using a 64-electrode constellation basket-catheter (n=15pts); iii) AF mapped using a 16-electrode HD-grid catheter (n=42pts). To investigate temporal fibrillation persistence, the response of AF-episodes to flecainide (n=7pts) was also studied.

Results: Spontaneously terminating fibrillation demonstrated a lower D_i (P<0.001 all systems). Lower D_i was also seen in paroxysmal compared to persistent AF (P=0.002). Post-flecainide, D_i decreased over time (P<0.001). Lower D_i was also associated with longer-lasting episodes of AF/VF (R^2>0.90, P<0.05 in all cases). Using k-means clustering, two distinct clusters and their centroids were identified i) a cluster of spontaneously terminating episodes, and ii) a cluster of sustained epochs.

Conclusion: Lower D_i correlates with less temporally persistent cardiac fibrillation. This finding provides potentially important insights into a possible common pathway to termination and therapeutic approaches.

Key words: *Atrial fibrillation, ventricular fibrillation, mechanisms, spiral waves, rotors, chaos*

INTRODUCTION

Cardiac fibrillation is a spatiotemporally disordered electromechanical state observed in either the atria or ventricles.[1] A fundamental property of the arrhythmia is the propensity for some episodes of fibrillation to spontaneously terminate and transition into sinus rhythm, while other episodes may persist. In the case of atrial fibrillation (AF), individual episodes may be as short as a few seconds, but may also last for hours, days or become permanent in individual patients. Ventricular fibrillation (VF) episodes may also have the propensity VF to self-terminate. However, if VF persists for longer than 10 minutes without resuscitation, the arrhythmia can lead to irreversible neurologic harm or death.[2-4]

The general properties of fibrillatory termination and persistence are of particular clinical interest as they have the potential to inform ongoing efforts to achieve personalized prediction of prognosis and therapy in both human fibrillatory disorders. Although extensive efforts have occurred over many decades to empirically identify predictors of termination or persistence in cardiac fibrillation,[5-7] at present there is no universally accepted conceptual paradigm or metric to explain why individual episodes of fibrillation may have the propensity to persist or spontaneously terminate. Such a general paradigm, if it could be developed, would have significant practical relevance, as it could potentially inform unresolved contemporary clinical questions. Examples of such challenges could include the persistence of AF despite pulmonary vein isolation[8,9], the upper limit of effectiveness observed with PVI,[8,9] objective delineation of paroxysmal from persistent forms of AF,[10] or understanding why some episodes of VF are responsive or unresponsive to defibrillation shocks.[11]

Here, we sought to contribute to efforts to develop a conceptual paradigm, to account for the potential for episodes of fibrillation to either persist or self-terminate. The electromechanical complexity that characterizes AF and VF has frequently been compared to the sustained disorder of nonequilibrium chaotic systems in nature that have been observed in physical,[12,13] chemical[14] and biological systems.[15,16]. An important general property observed in these systems is synchronization of behavior between spatially connected neighboring regions, that gradually breaks down with distance, a property that can be measured with the two-point spatial correlation length (ξ_2).

In an early study of spatiotemporal chaotic systems by Egolf and Greenside, it was postulated that

the ratio of system size (L) to correlation length (ξ_2), along with knowledge of spatial dimensionality

(d) could be used to decompose the behavior of chaotic systems into 'independent regions.'[17] Here,

we name this quantity, which was not formally defined in the original paper, as the spatial

autocorrelation dimension (D_i). The spatial autocorrelation dimension is defined with the following

equation:

103
$$D_i = (L/\xi_2)^d \quad (1)$$

104   where L the relative system size, ξ_2 the two-point correlation between pairwise electrodes spatially

105   spaced Δx apart (i.e. inter-electrode spacing), and d the spatial dimensionality of the system. As an

106   example of how the D_i estimator could potentially be computed, Egolf and Greenside suggested the

107   following estimator for a heart with a radius R and $d = 2$ (2):

108
$$D_i \approx 4\pi R^2/\pi\xi_2 \quad (2)$$

To our knowledge, the association of the spatial autocorrelation dimension estimator with the

persistence of termination with human cardiac fibrillation has never formally been assessed. In this

study, we evaluate the hypothesis that differences in D_i may account for the relative time until

termination in AF and VF.[18, 19] We explore this by calculating D_i in spontaneously-terminating versus

sustained episodes of AF and VF, as well as in episodes of AF post pharmacological intervention, with

supporting data from computer simulations. Evidence supporting this hypothesis may offer potentially

important clinical and mechanistic implications including: i) provide a quantitative framework to

understand the termination of cardiac fibrillation; ii) link termination in both atrial and ventricular to a

common mechanism; iii) link the termination of cardiac fibrillation to other nonlinear systems; and iv)

inform future investigations of unresolved clinical challenges in human fibrillatory disorders.

MATERIALS & METHODS

Epicardial sock recordings of human VF

The human VF study uses prospectively collected data as described by Nash et al.[20] The study recruited n=12 patients (n=8 sustained VF, n=4 self-terminating VF) undergoing routine coronary bypass graft procedures for ischemic heart disease with cross-clamp fibrillation. These studies were approved by the Hospital Ethics Committee (REC 01/0130) with written informed consent. Individual patient details are given in **Supplementary Materials S1, Supplemental Table 1**. Recordings were obtained using a 256-electrode epicardial sock (Δx = 10mm inter-electrode spacing) and unipolar electrograms (1kHz sampling) were preprocessed as previously described.[20, 21] In a subset of patients, VF spontaneously terminated before the full 210 seconds without requiring defibrillation, defined as 'self-terminating episodes'. This was compared to sustained VF episodes lasting the full duration. Additional study details are provided in **Supplement S1**.

Basket and HD grid recordings of human AF

The human AF study extends our previously published study.[22] The study was a multi-center observational design analyzing prospectively collected electrophysiologic data acquired prior to ablation (HREC110634). The inclusion criterion was AF undergoing ablation. Patient participation was by informed consent, with recruitment from Flinders University and Hamburg University. The patient cohort included n=15 patients (age 62±8, 13/15 male(87%), BMI 28.7±3.8 kg/m^2, non-paroxysmal 12/15(80%); **S1, Supplemental Table 2**). Basket catheter and HD-grid recordings were performed as previously.[23] 64-electrode basket catheters (Constellation, Boston Scientific, 48mm (Δx = 4mm), 60mm (Δx = 5mm) and HD-grid a 3-3-3 mm catheter with 16 electrodes to record human AF (Δx = 3mm) were utilized to record unipolar electrograms (1-500 Hz, 2000Hz sampling) and surface electrocardiograms during spontaneous or induced AF lasting >1 minutes. Spontaneously terminating AF episodes were

defined as those that terminated before the full 1-minute without requiring defibrillation. Signal

filtering, cleaning, and processing are detailed further in Supplement S3.

Spatial autocorrelation dimension estimation

To calculate the spatial autocorrelation dimension, D_i, atrial and ventricular volumes were first

calculated using the convex hull of electrode co-ordinates from one of the following i) mapping catheter

(for atrial fibrillation), ii) epicardial sock (ventricular fibrillation) or iii) grid nodes (computer

simulations). This volume was then used to estimate the radius, R, which was related to the two-point

correlation, ξ_2, to give the spatial cross-correlation as given by equation (2).[18]

Specifically, ξ_2 gives the association between the maximum cross correlation and distance between two

pairwise signals recorded at different electrodes, which are spatially distanced Δx apart. This can often

be modelled by an exponential relationship[24]:

$$C(r) = e^{-r/\xi} \ (3)$$

The cross-correlation function between two sequences of normalized activations were calculated over

windows shifted throughout the duration of the recording, with a range of lag values encompassing

slightly more than one cycle length. The maximum cross-correlation was given by the correlation

coefficient for the sequence of activations between the two corresponding locations. This process was

repeated in a pairwise fashion to calculate the correlation between all possible combinations of

locations, with locations corresponding to electrodes for AF and VF data, or mesh vertices for computer

simulations.

Assessing the effect of pharmacological intervention

To understand the effect by which anti-arrhythmic medicines facilitate termination of

fibrillation, we studied the action of prototype sodium-channel blockade as archetypal

pharmacologic rhythm modulation strategy on spatial autocorrelation dimension. This was

assessed in: i) n=9 computer simulated epochs of AF with varying levels of I_{Na} inhibition (20%-

60%) leading to AF termination; and ii) n=21 epochs from 7 AF patients administered

flecainide during an electrophysiology study, mapped using the HD-grid catheter.

To simulate the effect of sodium-channel block in computer simulated AF, the two-dimensional

Courtemanche-Ramirez-Nattel cell model was implemented as previously described (run on

7x6cm square grids).[25] Differential equations were solved with a 25ms time-step, with

simulations run for 5 seconds. AF epochs were initiated by a standard S1-S2 cross-shock

protocol. To achieve spontaneous AF termination, ACh gradients were altered to a range of

values to achieve self-termination before the full 5 seconds (**Supplement S2**).

Comparison with other metrics characterising arrythmia dynamics

Electrogram-based metrics

To evaluate whether D_i provides new information, we evaluated whether established

electrogram-based metrics used to characterise arrythmia dynamics would associate with

spontaneously terminating AF and VF. This was assessed by calculating four metrics using the

HD-grid human AF data: i) Dominant Frequency (DF), Shannon Entropy (ShEn), mean bipolar

voltage, and peak-peak bipolar voltage. Details on the calculation of these metrics are provided

in **Supplement S4**.

Renewal rate constants

A common motif of extensive spatiotemporally disordered systems is the continuous creation

and annihilation of topological defects known as phase singularities (PS) PS.[23, 26] The average

rate of PS formation (given by λ_f) and PS destruction (λ_d) can be measured, which has shown

to decrease in cases of spontaneously terminating AF and VF.[21, 23] This decrease can be

measured using the '*mixing rate*', governed by the interaction λ_f and λ_d as reflected by the

second largest eigenvalue modulus (SLEM) of the Markov transition matrix (Supplement S4).

Here, we build on this observation and hypothesized that changes in λ_f, λ_d, the mixing rate

should associate with changes in D_i. The average rate of phase singularity (PS) formation

(given by λ_f) and destruction (λ_d) can be measured by calculating the exponential decay

constant from the inter-event time distribution as previously described[22] (further detailed in

Supplement S4). The interaction of λ_f and λ_d is given by the '*mixing rate*', which represents

the time taken to reach the stationary state distribution (**Supplement S4**). In this study, λ_f and

λ_d were calculated as previously described from pre-processed and Hilbert transformed

unipolar electrograms, with the PS detection algorithm further detailed **Supplement S5**. [21, 23]

Statistical Analysis

Differences in spatial autocorrelation dimension, correlation length, λ_f, λ_d and mixing rates

between cases of sustained and spontaneously terminating AF, VF, and computer simulated

fibrillation was assessed using generalized linear mixed models (significance at P<0.05)

(**Supplement S6**). A similar approach was used to study spatial autocorrelation dimension and

correlation length across different stages of VF and post flecainide administration. To study the

relationship between λ_f and λ_d with spatial autocorrelation dimension and correlation length,

a k-means clustering approach was implemented using 10 replicates and a city-block algorithm

as the distance metric. Additional details are provided in **Supplement S6**.

RESULTS

Spatial autocorrelation dimension reflects the likelihood AF/VF self-termination and observed epoch length

Sustained episodes of basket-catheter mapped human AF were associated with lower spatial autocorrelation dimension (D_i) (mean terminating = 7.10,95%CI:4.77,9.44; mean sustained = 34.38,95%CI:16.87,51.88; P<0.001). This was also seen in HD-grid mapped human AF (mean sustained = 78.78,95%CI:63.35,94.21; mean terminating = 9.41,95%CI:4.49,14,33; P=0.001) and human VF (mean terminating = 10.30,95%CI:-1.90,22.50; mean sustained = 52.80,95%CI:27.13,78.48; P<0.001) (**Figure 1A-C**). Higher D_i also associated with increasing epoch lengths that was best fit to a power-law relationship in basket-mapped human AF (R^2 = 0.940; P<0.001) and human VF (R^2 = 0.817; P=0.035) when compared to a first-degree polynomial (linear) and exponential relationship. However, in HD-grid human AF, the relationship between epoch length and the spatial autocorrelation dimension fit well to both a power (R^2 = 0.909; P<0.001) and exponential (R^2 = 0.906; P<0.001) relationship (**Fig.1D-F**).

Relationship to the evolution of human atrial and ventricular fibrillation

D_i were also further assessed in 380 epochs arising from n=24 paroxysmal AF patients, and 282 epochs arising from n=18 persistent human AF patients. Epochs from paroxysmal AF patients were associated with consistently lower D_i in paroxysmal AF (25.47,95%CI:25.47,36.70) when compared to persistent AF (57.73,95%CI:43.93,71.52) (P=0.002; **Figure 2A**).

Human ventricular fibrillation episodes from n=12 patients undergoing cardiac surgery were also studied. Three distinct stages of VF were studied: i) the perfusion stage (initial 30 seconds); ii) ischemia (following 150 seconds); and iii) reflow (last 30 seconds). **Extended Data Fig.1** depicts the propagation of electrical activity as VF evolves throughout the three stages, which can be seen to depict more turbulent behaviour as VF progresses. This evolution

was associated with larger D_i (mean perfusion = 14.78,95%CI:5.33,24.67); mean ischemia = 45.48,95%CI:33.36,57.12); mean reflow = 82.06,95%CI:41.78,122.22). Specifically, D_i between the perfusion and ischemia (P=0.012), and perfusion and reflow (P<0.001), and ischemia and reflow stages were significantly different (P=0.011) (**Figure 2B**).

Relationship to other fibrillatory metrics

Rates of PS formation and destruction

In all model systems, the rate of PS formation, λ_f, was seen to decrease in spontaneously terminating basket-mapped human AF (terminating = 0.017,95%CI:0.015,0.019); sustained = 0.027,95%CI:0.025,0.030), HD-grid human AF (terminating = 0.0022,95%CI:0.0017,0.0028; sustained = 0.007,95%CI:0.0066,0.0075), and human VF (terminating = 0.011,95%CI:0.010,0.012; sustained = 0.023,95%CI:0.024,0.025). Similar trends were seen for the rate of PS destruction, λ_d in basket-mapped human AF (terminating = 0.016,95%CI:0.014,0.017; sustained = 0.022,95%CI:0.021,0.23), HD-grid human AF (terminating = 0.026,95%CI:0.021,0.031); sustained = 0.060,95%CI:0.055,0.065), and human VF (terminating = 0.0103,95%CI:0.009,0.0106; sustained = 0.46,95%CI:0.42,0.50).

When related to the spatial autocorrelation dimension and assessed using k-means clustering, two distinct clusters and their centroids were identified: i) a cluster of self-terminating epochs situated towards the upper right-hand side of graph (orange), and ii) a cluster of sustained epochs on the lower left-hand side of the graph (blue) (**Fig.3A-C i and ii**). In all model systems, the two clusters gave large positive silhouette values (**Fig.3A-C iii and iv**), indicating that the clusters are well separated. The changes seen in λ_f and λ_d also led to larger SLEMs and slower mixing rates characterizing episodes of spontaneously terminating fibrillation in all model systems (**Fig.4A-D**).

Electrogram-based measures

When applied to human AF data, it was found that DF showed a trend towards lower values in

in spontaneously terminating epochs, however, no statistically significant differences were

found when compared to sustained AF epochs (mean sustained = 7.11Hz,95%CI:6.46,7.76;

mean terminating = 6.522Hz,95%CI:5.74,7.31; P = 0.26) (**Fig.3D**). Similar observations were

made for other electrogram-based metrics, with a downward trend in values associated with

terminating episodes, but no statistically significant differences observed for ShEn (mean

sustained = 4.59 bits,95%CI:4.02,5.16; mean terminating = 4.31 bits,95%CI:3.85,4.77;

P=0.49), mean bipolar voltage (mean sustained = 4.5×10^{-4} mV,95%CI:0.26×10^{-3},0.63×10^{-3};

mean terminating = 4.20×10^{-4} mV,95%CI:0.14×10^{-3},0.70×10^{-3}; P=0.85), and peak-peak bipolar

voltage (mean sustained = 0.57 mV,95%CI:0.41,0.73; mean terminating = 0.28

mV,95%CI:0.22,0.35; P=0.08) (**Fig.3D**).

Changes in response to anti-arrhythmic drug modulation

In cases of computer simulated AF, generalized linear mixed modelling identified a decreasing

trend in spatial autocorrelation dimension (D_i) (**Figure 6A ii**), rate of PS formation (λ_f), Figure

6A iii) and PS number (Figure **6C iv**) with increasing Na+ block, but no significant change was

seen for the rate of PS destruction (λ_d). Statistically significant differences were also observed

in control versus Na$^+$-channel block simulations for D_i (P<0.001) and PS number (P<0.001),

but not the PS formation rate λ_f (P=0.19).

In cases of human AF, the generalized linear mixed model identified a decreasing trend in D_i

(**Figure 6B ii**), rate of PS formation (λ_f, **Figure 6B iii**) and PS number (**Figure 6B iv**) in post-

flecainide recordings over time, but no significant change in the rate of PS destruction (λ_d).

Statistically significant differences were also observed pre- versus post-flecainide for D_i

(P<0.001), λ_f (P<0.001) and PS number (P=0.003).

DISCUSSION

Key insights from this study

In this study we aimed to evaluate the hypothesis that the spatial autocorrelation dimension (D_i) could represent a potentially useful approach to evaluating the potential for persistence or termination in atrial fibrillation. In multiple human systems studied, we observed that D_i was consistently higher in episodes that were more sustained (lasting the full period of observation) relative to episodes that self-terminated. This trend was also observed in patients classified using the clinical persistent (lasting >7 days) definition of AF, and further supported in episodes of AF where no anti-arrhythmic drugs were administered. These findings demonstrate that differences in D_i may be a useful tool to characterise the potential for persistence or termination of individual AF and VF episodes.

Electrophysiologic characterisation of human AF & VF is an important clinical need

Despite extensive research, the optimal approach to electrophysiologically characterise human fibrillatory dynamics remains incompletely defined. At present, in AF in particular, fundamental challenges remain such as why AF progresses from paroxysmal to persistent in some but not all patients,[10] why a proportion of patients respond to anti-AF medicines but others do not,[27] why AF persists despite durable pulmonary vein isolation in some patients,[8, 9] and why substrate and linear ablation has not led to dramatically improved clinical outcomes.[28] These problems have led many in the clinical field to suggest that understanding of the biophysical processes sustaining fibrillation is not complete, providing an important rationale to support the exploration of new strategies for electrophysiologic characterization of human fibrillatory dynamics.

Spatial autocorrelation dimension may add to available tools to characterise AF and VF dynamics

D_i is an estimator of local electrophysical spatial synchronization in AF and VF, in comparison to overall system size, with the key finding that this metric appears to be a potentially important predictor of the potential for temporal persistence of fibrillatory episodes. Here, we use the term estimator in the statistical sense, of acting as a rule or criterion for calculating an estimate of a parameter from observable data.

D_i offer a number of potential advantages for the purpose of characterisation of human fibrillatory dynamics, that may make this metric provide useful information in addition to established quantitative tools to explore fibrillatory dynamics. Firstly, the information captured by D_i leverages the inherent spatial properties of the AF and VF. While traditional metrics of fibrillatory dynamics, such as dominant frequency, entropy, voltage and fractionation are primarily reliant on local timing and electrogram properties, D_i is an integrated measure over spatially distributed data points, taking into account the overall system size. This may potentially provide important nuanced insights into the system's underlying dynamics. The spatially informed perspective provided by D_i potentially allow for more accurate consideration of the multi-scale and nonlinear behavior, providing insight into the structural and functional organisation of AF and VF as biophysical processes.

Secondly, D_i is attractive because it is conceptually plausible as potential metric to explain the persistence of fibrillation. In its provenance, D_i has clear theoretical foundations in nonlinear systems theory, although fascinatingly, in the Egolf and Greenside study that motivated this paper D_i was not formally named or defined, despite the hypothesis that this estimator could potentially be used to characterise spatial synchronization in fibrillation being being presented as a potential use case.[17] Mechanistically, D_i is dependent on three factors: (i) the system size; (ii) correlation length; (iii) and spatial dimensionality, analogous to the aspect ratio in extensive chaotic systems.[29] Furthermore, the foundations for D_i are supported by long-standing empirically derived principles in cardiac fibrillation. The critical mass hypothesis has long been a principle[30][31,32] underlying the persistence of fibrillation, and the clinical association of increasing chamber size with the probability of AF maintenance is a generally accepted clinical principle.[31,32] Separately, the association of spatial correlation as tool to differentiate paroxysmal from persistent forms of AF has historically been independently clinically validated.[33,34] D_i, as an integrated estimator of spatial synchronisation relative to system size, has the potential to usefully add to available tools for the electrophysiologic characterisation of fibrillation.

Thirdly, a key strength of strength of D_i is that it is potentially able to be measured using clinically available data and has the potentially to leverage recent advances in catheter mapping technologies. Quantification of other similar metrics such as wavelength, has been a challenge for in vivo clinical

implementation due to the need for the precise calculation of conduction velocity (CV), effective refractory period (ERP) or action potential duration (APD), difficult to obtain in human patients .[35] In contrast, D_i can be rapidly calculated from commonly recorded electrical signals such as the intracardiac unipolar & bipolar electrograms, which are routinely collected in electrophysiology studies.

Recent advances in electroanatomic have expanded the potential applications of spatial synchronisation metrics. Novel array catheters[36-38] allow simultaneous measurement of electrical synchronisation spatial dispersed electrogram information in a variety of potential configurations, which was historically limited in the era of linear mapping catheters.[33] Additionally, advances in 3D electroanatomic mapping allow rapid and accurate volumetric estimation of cardiac chamber dimensions, allowing system size to be determined.[39] With new systems integrating real-time export of electrogram information, measurement of D_i has the potential to be clinically available on existing commercial mapping platforms.

It is important to note that because D_i is dependent on system size, correlation length and dimension, the numerical value will vary among different systems studied. However, an interesting and important point is that D_i still accurately separates terminating from sustained fibrillation, and more persistent forms of AF and VF from the less severe. It also associated with the termination of AF by anti-arrhythmic drugs. Another important point to note is that the relationship between D_i and the observed time until AF and VF termination (epoch length) best fit to a power-law, potentially indicating that D_i more accurately predicts shorter lasting episodes of fibrillation with a higher likelihood of impeding termination, as opposed to more temporally persistent episodes where termination will occur much later. However, D_i may still be clinically useful in applications such as stratifying episodes of AF/VF likely to resolve on their own in the short-term, versus those likely to persist for some time.

Clinically, the spatial autocorrelation dimension could provide an important additional quantitative metric to those currently available to measure fibrillatory dynamics.[34, 35] Firstly, a potential benefit of this approach is that it could contribute to an overarching explanation of why cardiac fibrillation self-terminates. Existing contemporary clinical theories of fibrillation, such as the multiple wavelet re-entry,

361 rotor, or focal source, or critical mass theories of AF and VF[40], do not specifically quantitatively and

362 directly address why any individual episode of should have a propensity for self-termination. Similarly,

363 existing quantitative metrics of fibrillatory dynamics such as dominant frequency, entropy,

364 fractionation, or voltage-based substrate characterisation, also leave a potential gap as to why individual

365 epochs of AF and VF self-terminate.[41]

Links to spatiotemporal turbulence in other systems

An important point to mention is that D_i as presented here to quantify the propensity of AF and VF

terminate is linked to concepts that have been extensively explored in theoretical models that have been

used to study transient spatiotemporal chaos.[42-45] The termination of such transient spatiotemporal chaos

has been proposed to occur due to underlying differences in the system phase space, which describes

how a system tends to evolve under a variety of initial conditions.[46] For example, transient

spatiotemporal turbulence occurs when a *'chaotic saddle'* or *'non-attracting chaotic set'* is present,

which allows turbulent behaviour to eventually terminate when the system escapes into the asymptotic

attractor.[19, 47] D_i represents an index of complexity that approximates the degree of turbulence and

number of independent regions in the system[12], which potentially relates to the likelihood of the system

escaping into such an asymptotic attractor.

Translational perspective

The results presented here may be conceptually useful for understanding and predicting the propensity

for the persistence of fibrillation in a variety of clinical contexts. Clinical studies are underway to

validate the role of cardiac fibrillation in objectively determining the association of clinically measured

D_i with atrial fibrillation persistence, and clinical response to ablation.[48] D_i has the potential to

contribute to the development of new metrics to predict anti-arrhythmic drug response. Clinical studies

have recently been commenced to determine the association of D_i and the long-term efficacy of anti-

AF treatment. D_i may also have a potentially important role in determining response to AF ablation. An

important incompletely resolved clinical question is the role of what if further substrate or linear

compartmentalisation should occur in patients beyond pulmonary vein isolation.[21] It is possible that

cardiac D_i could play a role in determining which patients could require further ablation beyond PVI,

and used as an objective metric to guide clinical responses. Future investigations are underway to answer

these questions.

CONCLUSION

Lower spatial autocorrelation dimension, D_i, characterizes the termination of fibrillation. This

observation could provide insights into the factors governing termination and persistence of fibrillation

and add to improved therapeutic approaches based on spatial autocorrelation dimension modulation

Figure 1 Spatial correlation dimension reflects the likelihood of human AF/VF termination and epoch length

397   ***Figure 1: Spatial autocorrelation dimension reflects the likelihood of AF/VF termination and epoch length***

398   (**1A-C**): Terminating epochs of AF and VF are associated with lower spatial autocorrelation dimensions (1A-Cii). Observed epoch lengths

399   (which gives time until termination) increased with larger spatial autocorrelation dimensions, with a relationship that best fit to a power-

400   law (**1D-F**).

Figure 2: Relationship to the evolution of AF and VF

(**2A**): Paroxysmal AF was associated with lower spatial autocorrelation dimensions compared to persistent AF. (**2B**): As ischemic VF evolves and becomes more turbulent and disordered, spatial autocorrelation dimension increased.

Figure 3 Spatial correlation dimension vs. renewal rates and comparison to other fibrillatory metrics

Figure 3: Spatial autocorrelation dimension vs. renewal rates and comparison to other metrics

(**3A-C**): Rates of PS formation (λ_f) and destruction (λ_d) reduced with lower spatial correlation (D_i) in terminating epochs of AF and VF. When compared to sustained AF and VF epochs, two clusters were identified by k-means clustering. This separated and spontaneously terminating (blue) and sustained (orange) simulations ((i) and (ii)). The large positive silhouette values indicate good separate of these clusters ((iii) and (iv)). (**3D-G**): Electrogram-based metrics such as dominant frequency (DF), Shannon entropy, mean bipolar voltage, and peak-peak bipolar voltage showed a trend towards lower values in terminating AF/VF, but no statistically significant differences when compared in sustained versus spontaneously terminating AF and VF epochs.

Figure 4 · Changes in SLEM and mixing rate

Figure 4: Changes in SLEM and mixing rate

(**4A**): Decreases in λ_f and λ_d are reflected by increases in the second largest eigenvalue modulus (SLEM) of the Markov transition matrix. In all model systems, cases of sustained fibrillation demonstrated lower SLEM compared to self-terminating episodes, indicated by the greater spread of eigenvalues on the eigenvalue spectrum plot. (**4B-D**): The higher SLEM seen in self-terminating episodes also lead to a lower mixing rate when compared to sustained episodes in all model systems.

Figure 5 | Changes in response to anti-arrythmic drug modulation

441   *Figure 5: Changes in response to anti-arrhythmic drug modulation*

442   (**5A**): A decrease in spatial autocorrelation dimension D_i, λ_f and PS number was observed over

443   time in Na$^+$ block simulations, but not λ_d. (**5B**): In episodes of flecainide-infused human AF, a

444   decrease in spatial autocorrelation dimension D_i, λ_f and PS number was also seen, but not λ_d.

REFERENCES

1.   Gray R and Jalife JJCAIJoNS. Ventricular fibrillation and atrial fibrillation are two different beasts. *Chaos: An Interdisciplinary Journal of Nonlinear Science*. 1998;8:65-78.

2.   Weisfeldt ML and Becker LBJJ. Resuscitation after cardiac arrest: a 3-phase time-sensitive model. 2002;288:3035-3038.

3.   Ikeda T, Uchida T, Hough D, Lee JJ, Fishbein MC, Mandel WJ, Chen P-S and Karagueuzian HSJC. Mechanism of spontaneous termination of functional reentry in isolated canine right atrium: evidence for the presence of an excitable but nonexcited core. 1996;94:1962-1973.

4.   Clayton R, Murray A, Higham P and Campbell RJTL. Self-terminating ventricular tachyarrhythmias—a diagnostic dilemma? 1993;341:93-95.

5.   Rostock T, Salukhe TV, Steven D, Drewitz I, Hoffmann BA, Bock K, Servatius H, Müllerleile K, Sultan A and Gosau NJHr. Long-term single-and multiple-procedure outcome and predictors of success after catheter ablation for persistent atrial fibrillation. 2011;8:1391-1397.

6.   Buch E, Share M, Tung R, Benharash P, Sharma P, Koneru J, Mandapati R, Ellenbogen KA and Shivkumar KJHR. Long-term clinical outcomes of focal impulse and rotor modulation for treatment of atrial fibrillation: A multicenter experience. 2016;13:636-641.

7.   Takahashi Y, O'Neill MD, Hocini M, Dubois R, Matsuo S, Knecht S, Mahapatra S, Lim K-T, Jaïs P and Jonsson AJJotACoC. Characterization of electrograms associated with termination of chronic atrial fibrillation by catheter ablation. 2008;51:1003-1010.

8.   Benali K, Barré V, Hermida A, Galand V, Milhem A, Philibert S, Boveda S, Bars C, Anselme F, Maille BJCA and Electrophysiology. Recurrences of Atrial Fibrillation Despite Durable Pulmonary Vein Isolation: The PARTY-PVI Study. 2023:e011354.

9.   Ganesan AN, Shipp NJ, Brooks AG, Kuklik P, Lau DH, Lim HS, Sullivan T, Roberts‐Thomson KC and Sanders PJJotAHA. Long‐term outcomes of catheter ablation of atrial fibrillation: a systematic review and meta‐analysis. *Journal of the American Heart Association*. 2013;2:e004549.

10.   Lubitz SA, Benjamin EJ, Ruskin JN, Fuster V and Ellinor PTJNRC. Challenges in the classification of atrial fibrillation. 2010;7:451-460.

11.   Cheskes S, Verbeek PR, Drennan IR, McLeod SL, Turner L, Pinto R, Feldman M, Davis M, Vaillancourt C and Morrison LJJNEJoM. Defibrillation strategies for refractory ventricular fibrillation. 2022;387:1947-1956.

12.   Egolf DA, Melnikov IV, Pesch W and Ecke REJN. Mechanisms of extensive spatiotemporal chaos in Rayleigh–Bénard convection. 2000;404:733-736.

13.   Ecke RE, Hu YJPASM and Applications i. Spiral defect chaos in Rayleigh-Be´nard convection: defect population statistics. 1997;239:174-188.

14.   Winfree ATJS. Spiral waves of chemical activity. 1972;175:634-636.

15.   Tan TH, Liu J, Miller PW, Tekant M, Dunkel J and Fakhri NJNP. Topological turbulence in the membrane of a living cell. 2020;16:657-662.

16.   Huang X, Xu W, Liang J, Takagaki K, Gao X and Wu J-yJN. Spiral wave dynamics in neocortex. 2010;68:978-990.

17.   Egolf DA and Greenside HS. Relation between fractal dimension and spatial correlation length for extensive chaos. *Nature*. 1994;369:129-131.

18.   Egolf DA and Greenside HSJN. Relation between fractal dimension and spatial correlation length for extensive chaos. 1994;369:129-131.

19.   Lai Y-CJPLA. Persistence of supertransients of spatiotemporal chaotic dynamical systems in noisy environment. 1995;200:418-422.

20.   Nash MP, Mourad A, Clayton RH, Sutton PM, Bradley CP, Hayward M, Paterson DJ and Taggart PJC.
Evidence for multiple mechanisms in human ventricular fibrillation. *Circulation*. 2006;114:536-542.

21.   Dharmaprani D, Jenkins E, Aguilar M, Quah JX, Lahiri A, Tiver K, Mitchell L, Kuklik P, Meyer C, Willems
S, Clayton R, Nash M, Nattel S, McGavigan AD and Ganesan AN. M/M/Infinity Birth-Death Processes – A
Quantitative Representational Framework to Summarize and Explain Phase Singularity and Wavelet Dynamics
in Atrial Fibrillation. *Frontiers in Physiology*. 2021;11.

22.   Dharmaprani D, Schopp M, Kuklik P, Chapman D, Lahiri A, Dykes L, Xiong F, Aguilar M, Strauss B,
Mitchell LJCA and Electrophysiology. Renewal theory as a universal quantitative framework to characterize
phase singularity regeneration in mammalian cardiac fibrillation. *Circulaiton: Arrhythmia and
Electrophysiology*. 2019;12:e007569.

23.   Dharmaprani D, Jenkins EV, Quah JX, Lahiri A, Tiver K, Mitchell L, Bradley CP, Hayward M, Paterson DJ
and Taggart PJHR. A governing equation for rotor and wavelet number in human clinical ventricular fibrillation:
Implications for sudden cardiac death. 2021.

24.   Botteron GW and Smith JMJC. Quantitative assessment of the spatial organization of atrial fibrillation
in the intact human heart. 1996;93:513-518.

25.   Courtemanche M, Ramirez RJ and Nattel S. Ionic mechanisms underlying human atrial action potential
properties: insights from a mathematical model. *American Journal of Physiology - Heart and Circulatory
Physiology*. 1998;275:H301-H321.

26.   Winfree AT. *When time breaks down: the three-dimensional dynamics of electrochemical waves and
cardiac arrhythmias*: Princeton University Press; 1987.

27.   Hohnloser SH, Kuck K-H and Lilienthal JJTL. Rhythm or rate control in atrial fibrillation—
Pharmacological Intervention in Atrial Fibrillation (PIAF): a randomised trial. 2000;356:1789-1794.

28.   Brooks AG, Stiles MK, Laborderie J, Lau DH, Kuklik P, Shipp NJ, Hsu L-F and Sanders PJHr. Outcomes of
long-standing persistent atrial fibrillation ablation: a systematic review. *Heart Rhythm*. 2010;7:835-846.

29.   Cross M and Greenside H. *Pattern formation and dynamics in nonequilibrium systems*: Cambridge
University Press; 2009.

30.   Kim YH, Garfinkel A, Ikeda T, Wu TJ, Athill CA, Weiss JN, Karagueuzian HS and Chen PS. Spatiotemporal
complexity of ventricular fibrillation revealed by tissue mass reduction in isolated swine right ventricle. Further
evidence for the quasiperiodic route to chaos hypothesis. *J Clin Invest*. 1997;100:2486-500.

31.   Zipes DP, Fischer J, King RM, Nicoll Ad and Jolly WWJTAjoc. Termination of ventricular fibrillation in
dogs by depolarizing a critical amount of myocardium. 1975;36:37-44.

32.   Garrey WEJAJoP-LC. The nature of fibrillary contraction of the heart.—Its relation to tissue mass and
form. 1914;33:397-414.

33.   Botteron GW and Smith JM. Quantitative assessment of the spatial organization of atrial fibrillation in
the intact human heart. *Circulation*. 1996;93:513-8.

34.   Hoekstra B, Diks C, Allessie M, DeGoede JJAop and biochemistry. Spatial correlation analysis of atrial
activation patterns during sustained atrial fibrillation in conscious goats. 2000;108:313-331.

35.   Jacquemet V, Virag N and Kappenberger LJEE. Wavelength and vulnerability to atrial fibrillation:
Insights from a computer model of human atria. 2005;7:S83-S92.

36.   Hong KL, Redfearn D, Chacko S, Baley J, Baranchuk A and Glover BM. High-resolution mapping of the
atria using the HD Grid catheter. *HeartRhythm Case Reports*. 2019;5:351-353.

37.   Sroubek J, Rottmann M, Barkagan M, Leshem E, Shapira‐Daniels A, Brem E, Fuentes‐Ortega C,
Malinaric J, Basu S, Bar‐Tal M and Anter E. A novel octaray multielectrode catheter for high‐resolution atrial
mapping: Electrogram characterization and utility for mapping ablation gaps. *J Cardiovasc Electrophysiol*.
2019;30:749-757.

38.   Yavin HD, Bubar ZP, Higuchi K, Sroubek J, Yarnitsky J and Anter E. Propagation Vectors Facilitate
Differentiation Between Conduction Block, Slow Conduction, and Wavefront Collision. *Circulation: Arrhythmia
and Electrophysiology*. 2021;14.

39.   Borlich M and Sommer P. Cardiac Mapping Systems: Rhythmia, Topera, EnSite Precision, and CARTO.
Card Electrophysiol Clin. 2019;11:449-458.

40.   Nattel S and Dobrev D. Controversies about atrial fibrillation mechanisms: aiming for order in chaos
and whether it matters. *Circulation Research*. 2017;120:1396-1398.

41.   Baumert M, Sanders P and Ganesan A. Quantitative-electrogram-based methods for guiding catheter
ablation in atrial fibrillation. *Proceedings of the IEEE*. 2016;104:416-431.

42.   Braun R and Feudel FJPRE. Supertransient chaos in the two-dimensional complex Ginzburg-Landau
equation. 1996;53:6562.

43.   Kaneko KJPLA. Supertransients, spatiotemporal intermittency and stability of fully developed
spatiotemporal chaos. 1990;149:105-112.

44.   Bohr T, Pedersen AW and Jensen MHJPRA. Transition to turbulence in a discrete Ginzburg-Landau
model. 1990;42:3626.

45.   Huber G, Alstrøm P and Bohr TJPrl. Nucleation and transients at the onset of vortex turbulence.
1992;69:2380.

46.   Lilienkamp T and Parlitz UJCAIJoNS. Terminating transient chaos in spatially extended systems<? A3B2
show [editpick]?>. 2020;30:051108.

47.   Crutchfield JP and Kaneko KJPrl. Are attractors relevant to turbulence? 1988;60:2715.

48.   Quah J, Dharmaprani D, Lahiri A, Schopp M, Mitchell L, Selvanayagam JB, Perry R, Chahadi F, Tung M
and Ahmad W. Prospective cross‐sectional study using Poisson renewal theory to study phase singularity
formation and destruction rates in atrial fibrillation (RENEWAL‐AF): Study design. *Journal of Arrhythmia*.
2020;36:660-667.

560