Predictable genetic recruitment for luciferin sulfation in the

convergent evolution of bioluminescence

Emily S Lau[1*], Jessica A Goodheart[1,2], Nolan T Anderson[3], Vannie L Liu[3], Arnab

Mukherjee[3,4,5,6*], Todd H Oakley[1*]

[1] Department of Ecology, Evolution, and Marine Biology, University of California Santa

Barbara, Santa Barbara, CA 93106

[2] Division of Invertebrate Zoology, American Museum of Natural History, New York, NY

10025

[3] Department of Chemical Engineering, University of California Santa Barbara, Santa

Barbara, CA 93106

[4] Department of Biological Engineering, University of California Santa Barbara, Santa

Barbara, CA 93106

[5] Department of Chemistry, University of California Santa Barbara, Santa Barbara, CA

93106

[6] Neuroscience Research Institute, University of California Santa Barbara, Santa

Barbara, CA 93106

[*]**Corresponding authors:** Emily Lau (emily.lau@lifesci.ucsb.edu); Arnab Mukherjee

(arnabm@ucsb.edu); Todd Oakley (oakley@lifesci.ucsb.edu)

Abstract

Determinism and contingency shape evolutionary trajectories and the extent to which evolution is predictable depends on the relative contribution of each. However, the causes of patterns of predictable evolution are poorly understood. Here we propose evolution may be more deterministic — and therefore predictable — if functional evolution is constrained to fewer genetic solutions. We examine the bioluminescence system of ostracods and synthesize our findings with those reported previously for fireflies and sea pansies to show one aspect of bioluminescence, substrate metabolism, evolved convergently by recruiting homologous enzymes. We use gene expression, recombinant protein expression, and *in vitro* functional analyses to identify genes capable of substrate metabolism in ostracods. Our results, taken together with knowledge of firefly and sea pansy bioluminescence systems, provide empirical evidence supporting that these convergent mechanisms for substrate metabolism repeatedly evolved by co-opting sulfotransferases, a ubiquitous family of enzymes with broad substrate specificity. From this synthesis, we propose the lack of diverse genetic solutions for sulfation constrains evolution, resulting in more evolutionary determinism and increasing our ability to predict evolution.

Keywords: convergent evolution, complex traits, gene expression, bioluminescence, sulfotransferase

Introduction

Introduction

Convergent evolution — the repeated evolution of similar phenotypes — provides

replicates useful for investigating long-standing biological questions, such as to what

extent evolution is predictable (Orgogozo 2015; Stayton 2015). Evolution is more

predictable and deterministic when similar features are shared across convergently

evolved phenotypes, but less predictable and more contingent when similar features are

not shared (Blount et al. 2018). Importantly, since many convergent phenotypes are

complex, the extent of convergence may differ across levels of biological organization,

or "multi-level convergent evolution" (Lau and Oakley 2020). In some cases, phenotypic

convergence can be similar at one level of biological organization but radically different

at other levels (Rosenblum et al. 2014). As one example, many vertebrates

convergently evolved viviparity — the ability to produce live young — using structurally

and functionally similar placentas, yet also using non-homologous genes (Foster et al.

2022). For viviparity, structural and functional evolution are more predictable than

genetic evolution. In some other cases, convergence extends across multiple levels of

biological organization. For example, two squid species convergently evolved

bioluminescence using structurally similar light-producing organs that express many

genes homologous between the species. Therefore, squid light-producing organs show

strong predictability of evolution across multiple levels of organization (Pankey et al.

2014). Fully understanding evolutionary predictability requires comparing convergent

phenotypes across multiple biological levels of organization. Furthermore,

understanding general rules of evolutionary predictability will require going beyond

individual case studies to determine causes of these patterns — when and why is

convergent evolution predictable?

For several reasons, bioluminescence — the biological production of light by a living

organism — provides an outstanding system for studying evolutionary predictability.

First, bioluminescence convergently evolved at least 94 times across the tree of life

(Lau and Oakley 2020). Second, genes used to produce bioluminescence can be

reliably tested via *in vitro* functional experiments (Shimomura 2019). Third, across

origins of bioluminescence the extent of evolutionary convergence at different levels of

biological organization is variable, leading to great diversity in form and function

(Haddock et al. 2010; Widder 2010). Bioluminescence is used for many biological

functions: deterring predation, attracting prey, camouflaging, intraspecific

communication, or courtship (Haddock et al. 2010). Bioluminescence is also produced

under diverse conditions. In bioluminescent systems light may be produced

intracellularly or extracellularly using a variety of complex structures (Harvey 1952),

including glandular organs that may secrete (Dilly and Herring 1978; Takenaka et al.

2012) or house bioluminescence (Morin 1983), modified fin rays that house luminous

bacteria (Hulet and Musil 1968), or luminous tubules and ducts (Thacker and Roje

2009). At a molecular level, bioluminescence is produced when enzymes called

luciferases oxidize small molecule substrates called luciferins (Shimomura 2019). To

harness bioluminescence, luminous organisms may modulate luciferase-luciferin

reactions by housing luciferases and luciferins in different cellular or subcellular

locations (Clarke et al. 1962; Huvard 1993), or inactivating luciferins by using enzymes

to bind to or biochemically modify them (Lee et al. 1993; Inouye 2007; Viviani et al.

2020). However, compared to knowledge of the function of bioluminescence and

luciferase enzymes, little is known about how luciferins are metabolized for storage,

transport, or modulation of light.

Fireflies, sea pansies, and cypridinid ostracods convergently evolved the same

biochemical mechanism, sulfation — or the transfer of sulfate (SO_3^-) from the donor 3′-

phosphoadenosine-5′-phosphosulfate (PAPS) to a substrate (Falany 1997) — for

modulating the chemical state of their luciferins (Hori et al. 1972; Nakamura et al. 2014;

Fallon et al. 2016; Tzertzinis et al. 2022) (Figure 1, Supplemental Figure S1). Once

sulfated, these luciferins are unable to react with their respective luciferases (Fallon et

al. 2016), and in the cases of sea pansies and ostracods, sulfation prevents non-

specific oxidation of their luciferins (Cormier and Kazuo 1964; Nakamura et al. 2014).

Sulfation is catalyzed by a family of enzymes called sulfotransferases that contain a

sulfotransferase domain with two highly conserved motifs necessary for binding to

PAPS and transferring sulfate to a substrate (Driscoll et al. 1995; Bidwell et al. 1999).

Sulfotransferases may exhibit "substrate promiscuity", which we define as an enzyme's

ability to catalyze the same biochemical reaction in more than one substrate.

Sulfotransferase genes underlying luciferin metabolism have been identified and

functionally tested in fireflies (Fallon et al. 2016) and recently in sea pansies (Tzertzinis

et al. 2022), but not in ostracods. Previous work identified the presence of sulfated

luciferin in the ostracod *Vargula hilgendorfii* (Nakamura et al. 2014; Nakamura et al.

2019). This work also found that crude protein extracts from the whole organism could

use PAPS to enzymatically sulfate luciferin — suggesting the presence of a luciferin

sulfotransferase — to produce an inactive form of luciferin that could not react with

luciferase and was less vulnerable to non-specific oxidation. Additionally, crude protein

extracts of *V. hilgendorfii* can remove sulfate from sulfated luciferin to produce active

luciferin (Nakamura et al. 2014; Nakamura et al. 2019) Since crude protein extracts

from *V. hilgendorfii* can reversibly sulfate luciferin, this suggests sulfation may serve as

a mechanism for luciferin metabolism in the organism (Nakamura et al. 2014; Nakamura

et al. 2019). However, the genetic sequences of functional luciferin sulfotransferases in

ostracods have yet to be identified.

Although fireflies, sea pansies, and cypridinid ostracods convergently evolved the same

mechanism for luciferin sulfation, the extent of evolutionary convergence varies across

levels of biological organization. These taxa convergently evolved bioluminescence as a

defense mechanism — which was later co-opted for courtship in fireflies (Powell et al.

2022) and ostracods (Ellis et al. 2022) — but produce light by using structurally diverse

organs or tissue (Peterson and Buck 1968; Spurlock and Cormier 1975; Huvard 1993),

different luciferins (Tsarkova 2021), and non-homologous luciferases (Cormier and

Kazuo 1964; Oba et al. 2003; Oakley 2005) (Figure 1). Understanding how these taxa

genetically encode luciferin sulfation and synthesizing these findings with those from

other levels of biological organization — namely, organismal, organ, and tissue — will

reveal the extent to which convergently evolved phenotypes may vary across levels of

biological organization.

Here, we propose how limited genetic solutions for certain biochemical functions may constrain convergent evolution, which may result in deterministic and predictable genetic evolution across complex traits that evolved convergently. Because bioluminescent ostracods sulfate their luciferin (Nakamura et al. 2014; Nakamura et al. 2019) and because sulfotransferases are the only known genetic mechanism for sulfation, we hypothesize luminous ostracods produce sulfotransferases capable of sulfating ostracod luciferin. To test our hypothesis, we use gene expression analyses, recombinant protein expression, and *in vitro* biochemical assays to identify and functionally test five candidate sulfotransferase genes highly expressed in the light-producing organ of the cypridinid ostracod *Vargula tsujii*. We perform two types of functional tests to show that all five candidates exhibit luciferin sulfation activity as evidenced by at least one assay. Our results, taken together with functional evidence of luciferin sulfotransferases previously published on sea pansies and fireflies (Fallon et al. 2016; Tzertzinis et al. 2022), provide empirical evidence supporting independent recruitment of paralogous genes from a single gene family — sulfotransferases — in the convergent evolution of luciferin metabolism from three phylogenetically distant organisms with distinct luciferin substrates. Because there are no other mechanisms known for sulfation, we assume there are limited genetic solutions, which leads us to propose predictable genetic evolution of luciferin sulfation.

Results

Four candidates highly expressed in the upper lip exhibit significant luciferin

sulfotransferase activity in vitro

We identified 40 putative sulfotransferases expressed in the upper lip (the light-

producing organ) of *Vargula tsujii*. To identify and functionally verify luciferin

sulfotransferase genes in the ostracod *V. tsujii*, we hypothesized that, similar to fireflies

(Fallon et al. 2016), sulfotransferases highly expressed in the light-producing organ are

capable of sulfating luciferin in a PAPS-dependent manner. We selected the five most

highly expressed sulfotransferase genes in the light organ (Figure 2, Supplemental

Figure 3), hereafter referred to as ST1-5, to recombinantly express and test for luciferin

sulfation function.

A universal sulfotransferase assay kit showed four candidate sulfotransferase genes —

ST1, ST3, ST4, ST5 — exhibit significant sulfation activity on luciferin compared to the

negative control, which lacked sulfotransferase (ST1: $p = 3.75e\text{-}3$, ST2: $p = 9.381e\text{-}2$,

ST3: $p = 4.9e\text{-}7$, ST4: $p = 1.4e\text{-}5$, ST5: $p = 1.2e\text{-}4$, N = 6, Dunnett's test) (Figure 3A).

This assay quantifies formation of phosphate from 3'-phosphoadenosine-5'-phosphate

(PAP) — a metabolic by-product of sulfotransferases — as a proxy for sulfation activity.

By avoiding a buildup of PAP in the reaction volume, this assay also mitigates potential

product inhibition, which could limit sulfation activity *in vitro*.

Three candidates exhibit sulfation activity sufficient to modulate bioluminescence in vitro

Three of five candidate luciferin sulfotransferases — ST2, ST3, and ST5 — showed

significant decreases in bioluminescence compared to PAPS-free controls (ST1: $p =$

2.91e-1, ST2: p = 3.79e-2, ST3: p = 2.05e-5, ST4: p = 9.6e-1, ST5: p = 2.16e-5, N = 3,

unpaired 2-sample Student's t-test) (Figure 3B), suggesting the activities of ST2, ST3,

and ST5 are sufficient to modulate bioluminescence because once sulfated, luciferins

are unable to react with their corresponding luciferases to produce light.

Four candidates exhibit significant sulfation activity with p-nitrophenol

Four candidates — ST2, ST3, ST4, and ST5 — catalyzed sulfation of p-nitrophenol

relative to PAPS-free controls (ST1: p = 1.45e-1, ST2: p = 1.17e-2, ST3: p = 4.22e-3,

ST4: p = 6.91e-5, ST5: p = 2.92e-2, N = 3, unpaired 2-sample Student's t-test) (Figure

4). Since many sulfotransferases sulfate a range of small molecules, we hypothesized

ST1-5 may sulfate small molecules besides luciferin. To test this for ST1-5, we

examined their ability to catalyze sulfation of p-nitrophenol, a xenobiotic substrate

commonly sulfated by cytosolic sulfotransferases. These results support substrate

promiscuity in the sulfotransferase activities of ST2, ST3, ST4, and ST5.

Ostracods, fireflies, and sea pansies sulfate luciferin by using homologous genes

Ostracods, fireflies, and sea pansies convergently evolved sulfation as a mechanism for

metabolizing luciferin by independently recruiting sulfotransferase genes (Figure 5).

Phylogenetic analysis of functionally characterized sulfotransferases place the luciferin

sulfotransferases of ostracods and fireflies in a clade of arthropod and mammalian

sulfotransferases. This clade contains a mixture of sulfotransferases that may exhibit

substrate promiscuity or specificity towards endogenous and xenobiotic small

molecules, such as p-nitrophenol. Similarly, the luciferin sulfotransferase of sea pansies

falls in a separate clade, which also contains sulfotransferases with substrate

promiscuity or specialized activity towards substrates such as hormones and xenobiotic

substrates.

Discussion

Convergently evolved phenotypes often result from a mixture of similar and different

features across and within levels of biological organization, providing excellent

opportunities to explore general rules of evolutionary predictability. As one general rule,

molecular evolution may be more predictable when biochemical functions have fewer

possible genetic solutions, and less predictable when genetic redundancy permits many

solutions. Here, we examine multi-level convergent evolution of bioluminescence in

ostracods, fireflies, and sea pansies. These taxa similarly use bioluminescence to deter

predation (Kastendiek 1976; Branham and Wenzel 2003; Ellis et al. 2022), but use

structurally different organs and tissues (Peterson and Buck 1968; Satterlie and Case

1979; Huvard 1993) and non-homologous luciferase enzymes to oxidize luciferins for

light production (Lorenz et al. 1991; Oba et al. 2003; Oakley 2005). At the same time,

these taxa evolved a similar mechanism, sulfation, for luciferin metabolism (Hori et al.

1972; Nakamura et al. 2014; Fallon et al. 2016). We provide functional evidence for

sulfotransferases capable of sulfating luciferin in the ostracod *Vargula tsujii*. We

synthesize our results with those previously described in fireflies and sea pansies,

leading us to propose luciferin sulfation convergently evolved by recruiting paralogous

genes from the same gene family. These differences between sulfation recruiting

homologous genes and light production recruiting different genes raises the question:

How might biochemical function shape the genetic evolution of complex biological

systems like bioluminescence?

The evolutionary convergence of bioluminescence systems demonstrates how function

may inform determinism and contingency in molecular evolution. Evolution may be

deterministic and predictable when convergently evolved systems share functions

constrained to homologous genetic solutions, and may be contingent and less

predictable when functions can arise from many non-homologous genetic solutions. For

example, since many marine luciferins are intrinsically sensitive to oxygen (Shimomura

2019), their luciferin-oxidizing functions may have convergently evolved using proteins,

or "proto-luciferases", that at first provided hydrophobic environments to facilitate the

oxidation of luciferin to produce some light (Rees et al. 1998; Adams and Miller 2020). If

increasingly higher amounts of light production continued to enhance survival and

reproduction, selection could have led to the evolution of extant luciferases. This

hypothesis proposes luciferin oxidation functions may be accomplished by many

different genetic solutions, which may help to explain why many luciferases evolved

from non-homologous ancestral proteins (Delroisse et al. 2021). In contrast, sulfation in

biological systems — as far as we know — is catalyzed by sulfotransferases, which

contain two highly conserved motifs used to bind to PAPS and to transfer sulfate to a

substrate (Driscoll et al. 1995; Bidwell et al. 1999). Moreover, many sulfotransferases

exhibit substrate promiscuity and can sulfate a range of substrates (Allali-Hassani et al.

2007), making these enzymes amenable to evolutionary recruitment. Selection may act

on this promiscuity, enabling the divergence and evolution of new functions (Tokuriki

and Tawfik 2009), such as sulfation of luciferin. As such, the evolution of luciferin

sulfation may have been constrained to a homologous genetic solution with a

predisposition to functional evolution. Taken together, we propose molecular evolution

of light production is less predictable due to genetic redundancy, and the evolution of

sulfation-mediated luciferin metabolism is more deterministic and predictable because

of genetic constraint.

We show the luminous ostracod *V. tsujii* expresses multiple sulfotransferases capable

of sulfating luciferin *in vitro*, supporting the hypothesis that sulfotransferases sulfate

luciferin in bioluminescent cypridinid ostracods. To test whether there are luciferin

sulfotransferases in *V. tsujii*, we performed two types of functional tests for assessing

luciferin sulfation activity, tested for significant activity as compared with control

reactions, and then corrected for multiple comparisons. The sulfotransferase assay

demonstrated significant luciferin sulfation activity for four of the five candidates — ST1,

ST3, ST4, and ST5 — as compared with the negative control by using Dunnett's test.

The luciferase assay identified three of the five candidates — ST2, ST3, and ST5 — to

have significant luciferin sulfation activities capable of modulating bioluminescence *in*

vitro. However, after accounting for multiple comparisons by using the Benjamini-

Hochberg procedure, only ST3 and ST5 were statistically significant. This discrepancy

between the results of the sulfotransferase and luciferase assays could be explained by

experimental differences between the two assays. The sulfotransferase assay required

a high concentration of luciferin substrate (100 µM) to generate enough product to

produce detectable absorbance measurements. In the luciferase assay, the

concentration of luciferin (10 µM) was ten times less than that of the sulfotransferase

assay. We found ostracod luciferase exhibits substrate inhibition (Supplemental Figure

S6), which limited the concentration of luciferin that could be used in this assay. Since

the rate of sulfation reactions depends on substrate concentrations, (Atkinson 1966;

James 2021), the differences in substrate concentrations between the two assays may

have led to the observed differences in product formation of sulfated luciferin. Inhibition

of sulfotransferases by PAP (Chapman et al. 2004), a by-product of sulfation produced

in both assays but removed only in the sulfotransferase assay, may have also

contributed to the observed differences in sulfation. Nevertheless, results from both

assays support the presence of multiple sulfotransferases capable of sulfating luciferin

and modulating bioluminescence in the light-producing organ of *V. tsujii*. Future

research on the *in vivo* conditions of bioluminescence — such as concentrations of

luciferin, luciferase, PAPS, and PAP — may further refine these *in vitro* assays by

confirming physiologically relevant testing conditions. These results — taken together

with those from firefly and sea pansy bioluminescence systems — support the repeated

co-option of sulfotransferases in the convergent evolution of luciferin sulfation, leading

us to propose how genetic constraints on functional evolution may result in the

predictable and deterministic evolution of this mechanism.

Functional evolution may be predictable if constrained by genetic solutions, but

functional specialization of the recruited genes may be unpredictable. Functional

evolution does not necessarily require functional specialization of recruited genes

(O'Brien and Herschlag 1999; Khersonsky et al. 2006). New functions may be catalyzed

by multi-functional and promiscuous enzymes, or by functionally specialized enzymes.

For example, the luciferases of fireflies and click beetles convergently evolved by

recruiting homologous genes via gene duplication and neofunctionalization. In fireflies,

both copies of the gene retained the ancestral function and one copy gained luciferase

function (Oba et al. 2003; Oba et al. 2006). In contrast, click beetles only have one copy

of the gene that retained the ancestral function, while the other copy gained and

specialized in luciferase function while losing the ancestral function (Oba et al. 2010). In

line with this, our findings suggest that, while homologous sulfotransferase genes are

recruited in the evolution of luciferin sulfation, functional specialization in these genes

may differ across convergent systems. Out of the luciferin sulfotransferases identified by

at least one assay, four — ST2, ST3, ST4, and ST5 — can also sulfate p-nitrophenol.

Conversely, in fireflies, Fallon et al. (2016) demonstrated only one of the three highly

expressed sulfotransferase genes from the light-producing organ was capable of

sulfating firefly luciferin and could not sulfate p-nitrophenol. Two other sulfotransferases

readily sulfated p-nitrophenol but not firefly luciferin (Fallon et al. 2016). In sea pansies,

Tzertzinis et al. showed a luciferin sulfotransferase gene encodes multiple isoforms, two

of which can function as luciferin sulfotransferases (Tzertzinis et al. 2022). Therefore,

although fireflies might use a functionally specialized luciferin sulfotransferase to sulfate

luciferin, ostracods may use many promiscuous luciferin sulfotransferases, and sea

pansies may use one or both luciferin sulfotransferase isoforms. To test these

hypotheses, further work, such as gene knockout experiments and gene co-expression

analyses, will be necessary for confirming which sulfotransferases are associated with

bioluminescence function *in vivo*.

Understanding how complex systems convergently evolved across levels of biological

organization may inform general rules of predictability in evolution. The convergent

bioluminescence systems of fireflies, sea pansies, and ostracods illustrate how a shared

organismal function — using bioluminescence to deter predation — uses structurally

different organs and tissues, and a mixture of non-homologous and homologous genes

to produce and modulate bioluminescence. Therefore, molecular evolution of

bioluminescence may sometimes be more predictable than structural evolution,

depending on the range of genetic solutions to a function. These ideas may be broadly

extended to the general rule that evolution may be predictable if functions are limited to

one or a few genetic solutions and may be far less predictable if redundancy permits

many non-homologous genetic solutions.

Materials and Methods

We purchased TRIzol^{TM} reagent, Terrific Broth, and the pCMV-Cypridina Luc vector

(encoding ostracod luciferase) from Thermo Fisher Scientific (Waltham, MA). We

purchased the pQE80L *E. coli* expression vector and Ni-NTA agarose from Qiagen

(Hilden, Germany). The pET-6xHis-SUMO-MarathonRT plasmid (source of SUMO tag)

was a gift from Dr. Anna Pyle (Addgene plasmid # 109029). We purchased *Cypridina*

(ostracod) luciferin from Prolume LTD (Pinetop, AZ) and prepared single-use aliquots by

dissolving 0.5 mg luciferin in 100 % methanol to a stock concentration of 6.15 mM, then

stored the aliquots at -80 °C. We obtained 3′-phosphoadenosine 5′-phosphosulfate

(PAPS) lithium salt from Santa Cruz Biotechnology (Dallas, TX), dissolved it in

phosphate buffered saline (pH 7.4) to prepare 2 mM aliquots, and stored aliquots at -80

°C. We purchased the sulfotransferase assay kit and lithium-free PAPS substrate from

R&D Systems (Minneapolis, MN). We obtained p-nitrophenol from MP Biomedicals

(Santa Ana, CA), dissolved it in 20 mM tris hydrochloride (pH 7.4) to prepare a stock

solution of 10 mM, and stored the stock at 4 °C. We purchased Chinese hamster ovary

(CHO) cell lines from Takara Bio (San Jose, CA), oligonucleotide primers and

gBlocks™ from Integrated DNA Technologies (Coralville, IA), and Q5 DNA polymerase,

agarose gel extraction kit, reagents for Gibson assembly, competent *E. coli* 10-beta

cells and NEB Stable cells (for plasmid propagation), and competent *E. coli* BL21 cells

(for protein expression) from New England Biolabs (Ipswich, MA). We purchased

reagents for Quick Start™ Bradford protein assay and denaturing polyacrylamide gels

(Mini-PROTEAN TGX stain-free protein gels) from BioRad (Hercules, CA), Amicon®

ultracentrifugal filter units (30 kDa and 10 kDa molecular weight cutoff) from Sigma

Aldrich (St. Louis, MO), and reagents for purifying plasmid DNA from Promega

(Madison, WI). We purchased linear polyethyleneimine (25 kDa) transfection reagent

from Polysciences (Warrington, PA), dissolved it in deionized water to prepare stock

solutions at a concentration of 0.258 mg/mL, and stored stock solutions at -20 °C.

Construction of a high-quality reference transcriptome for Vargula tsujii

We used PacBio full-length isoform sequencing (Iso-Seq) technology to generate a

high-quality reference transcriptome for the California luminous ostracod, *Vargula tsujii*.

We collected total RNA from 88 *V. tsujii* specimens across various instar stages

(carapace lengths, measured from the rostrum to the keel, ranging from 0.68 to 2.02

mm) (Supplemental Table S1), including adult males and brooding females

(Supplemental Figure S2). We extracted RNA using TRIzol™ reagent, following the

manufacturer's protocol with two minor modifications. First, we used a higher

centrifugation speed (16,100 x g) to pellet cell debris and separate the organic and

aqueous layers. Second, we increased precipitation time to ~ 1 hour to increase RNA

yield. We shipped purified RNA to the DNA Sequencing Center at Brigham Young

University for library preparation and Single Molecule Real-Time (SMRT) sequencing

(PacBio Sequel II). To maximize coverage of different sized transcripts, we used two

fractions for sequencing: one that was resolved by pulsed field electrophoresis

(BluePippin, Sage Science, Beverly, MA) to isolate 4-10 kb transcripts; and another that

was not size-selected. We clustered and polished circular consensus sequencing (CCS)

reads using the IsoSeq v3 pipeline (https://github.com/PacificBiosciences/IsoSeq). We

combined the ensuing long reads with Illumina short reads from NCBI (SRR21201581)

(Ellis et al. 2022) using the transcriptome assembler, rnaSPAdes, with the following

flags: --rna, --only-assembler, and --trusted-contigs (Bushmanova et al. 2019). To

reduce duplicated transcripts, we used cd-hit-est (Li and Godzik 2006; Fu et al. 2012) to

cluster sequences with ≥ 95 % identity. We assessed the completeness of the final

assembly by using Benchmarking Universal Single-Copy Orthologs (BUSCO) v.3.0.2

(Simão et al. 2015; Waterhouse et al. 2018). BUSCO analysis identified 91.1 % of the

total orthologs from the arthropod gene set, 80% of which was classified as complete

and 11.1 % fragmented.

Gene expression profiling and identification of sulfotransferase candidates in Vargula

tsujii

To identify the most highly expressed sulfotransferase sequences in *Vargula tsujii*, we

collected RNA from the light producing organ (known as the upper lip) of ostracod

specimens maintained in the Oakley lab culture (Goodheart et al. 2020). We extracted

upper lips from nine ostracods (five males and four females) by dissection under a

stereomicroscope, immediately mixed them with TRIzol^{TM}, and stored the samples at -

80 °C until further use. We extracted RNA — using the same extraction protocol as

described above, for constructing the Iso-Seq transcriptome — and shipped purified

RNA samples to the Genomic Sequencing and Analysis Facility at The University of

Texas at Austin for library preparation and gene expression profiling using Tag-Seq

(Illumina HiSeq 2500, SR100), resulting in an average of 1.1 million reads (range: 0.61-

2.1 million) per sample after filtering (Supplemental Table 2) (Meyer et al. 2011). We

processed Tag-Seq reads using a modified version of the Tag-Seq pipeline freely

available on GitHub (https://github.com/z0on/tag-based_RNAseq) (Meyer et al. 2011),

which mapped the reads to a reference transcriptome — the Iso-Seq transcriptome of

V. tsujii — using Bowtie2 v.2.3.4.3 (Langmead and Salzberg 2012). On average, 0.87

million reads per sample (range: 0.45-1.5 million) produced unambiguous alignments to

the reference transcriptome (Supplemental Table 2). We translated the reference

transcriptome using TransDecoder v.5.5.0 (Haas, BJ.

https://github.com/TransDecoder/TransDecoder) and used HMMER v.3.2.1 (hmmer.org)

to identify protein sequences containing a sulfotransferase domain (Pfam ID: PF00685)

with an e-value score less than or equal to 1e-3. We used MAFFT (v.7.453) (Katoh and

Standley 2013) to align protein sequences and used IQ-TREE v.1.6.12 (Nguyen et al.

2015) to infer a maximum likelihood phylogeny, assuming the best-fit substitution model

(LG+I+G4) — according to Bayesian Information Criterion — as determined with

ModelFinder (Kalyaanamoorthy et al. 2017). To midpoint root the phylogeny and

visualize normalized gene expression counts, we used the R package phytools v.1.0.3

(Revell 2012).

Cloning, expression, and purification of ostracod (Cypridina noctiluca) luciferase

To facilitate protein purification, we used Gibson assembly (Supplemental Table S3) to

incorporate 3 x FLAG and hexahistidine tags at the C-terminus of an ostracod

(*Cypridina noctiluca*) luciferase encoding gene in the pCMV-Cypridina Luc vector

(Supplemental Figure S4). We propagated this modified vector by transforming

chemically competent NEB Stable *E. coli* cells. After DNA sequencing (Genewiz, South

Plainfield, NJ) to confirm successful cloning, we co-transfected CHO cells with this

vector and a marker plasmid encoding green fluorescent protein (GFP) by transient

transfection of ~ 70% confluent cultures using polyethylenimine. We used fluorescence

microscopy (Olympus CKX-41) to confirm transfection success by monitoring the

production of GFP, then we harvested the cell culture media — containing secreted

ostracod luciferase — 48 hours post transfection, and incubated the media with Ni-NTA

agarose beads in 10 % (v/v) sample buffer (500 mM NaH_2PO_4, 1.5 M NaCl, 100 mM

imidazole, pH 8) on a rotary shaker overnight at 4 °C. Subsequently, we loaded the Ni-

NTA agarose slurry in a gravity-flow column, collected the flowthrough, and washed the

column twice with 3 mL wash buffer (20 mM NaH_2PO_4, 500 mM NaCl, 10 mM imidazole,

pH 7.4) to remove nonspecifically bound proteins. Then, we eluted the column twice

with 2 mL elution buffer (20 mM NaH_2PO_4, 500 mM NaCl, 500 mM imidazole, pH 7.4).

To confirm the presence of ostracod luciferase throughout purification, we added 20 µL

of each purification step — namely, the flowthrough, wash, and each elution step — to

50 µL of 20 mM tris hydrochloride pH 8.0. We then tested for luciferase activity by

adding 5 µL of 50 µM luciferin to each sample and recorded luminescence using a

microplate reader (Tecan Spark) with an integration time of 2 seconds. After confirming

the presence of luciferase in our eluate samples (Supplemental Figure S5), we

concentrated the eluate and exchanged it into a storage buffer (20 mM tris

hydrochloride, 20 % glycerol, pH 8.0) by spin filtration using a 30 kDa molecular weight

cutoff filter. We assessed the purity of recombinant luciferase by performing denaturing

polyacrylamide gel electrophoresis, visualized the bands by activating the stain-free

gels with UV (BioRad Gel Doc), (Supplemental Table S4, Supplemental Figure S5) and

quantified protein yield using the Bradford assay. We flash-froze aliquots of recombinant

luciferase in dry ice and stored them at -80 °C until use.

Characterizing ostracod (Cypridina noctiluca) luciferase kinetics

To characterize the enzyme kinetics of ostracod luciferase, we added 65 µL of purified

ostracod luciferase to 10 µL of luciferin, resulting in a total volume of 75 µL and final

concentrations of 250 nM luciferase and varying final concentrations — 0 µM, 0.39 µM,

0.78125 µM, 1.5625 µM, 3.125 µM, 6.25 µM, 18.75 µM, 37.5 µM, 75 µM, and 150 µM —

of luciferin. We incubated the reactions at room temperature for 12 minutes to stabilize

the reading and quantified luminescence using a microplate reader (Tecan Spark) with

an integration time of 2 seconds (N = 3). Since we observed decreasing light production

at final substrate concentrations equal to or greater than 3.125 µM, we plotted and fitted

these data using a Michaelis-Menten equation modified for modeling substrate inhibition

(Copeland 2000), where v = velocity, V_{max} = maximum velocity, S = substrate

concentration, K_m = Michaelis-Menten constant, and K_i = the dissociation constant for

the inactive ternary complex formed when another substrate molecule binds to an

enzyme-substrate complex (Supplemental Figure S6):

12
$$v \ = \ \frac{V_{max}\,[S]}{K_m \ + \ [S]\left(1 + \frac{[S]}{K_i}\right)}$$

Cloning, expression, and purification of Vargula tsujii sulfotransferase candidates

To express and purify candidate luciferin sulfotransferase proteins, we codon-optimized

and synthesized DNA sequences corresponding to ST1-5 as gBlocks™ (Integrated

DNA Technologies). We used Gibson assembly (Supplemental Table S3) to clone these

fragments into a bacterial expression vector, pQE80L, allowing expression of the

resulting proteins with an N-terminal hexahistidine tag for purification (Supplemental

Figure S4). All plasmid constructs were propagated by electroporation of competent *E.*

coli NEB® 10-beta cells, verified by DNA sequencing (Genewiz, South Plainfield, NJ),

and subsequently electroporated in competent *E. coli* BL21 cells for protein expression.

Initial efforts to express these proteins in *E. coli* BL21 resulted in soluble expression

only for ST3 and ST4 (Supplemental Table S4, Supplemental Figure S7). For the

remaining proteins — namely ST1, ST2, and ST5 — we attempted to improve

expression by cloning the genes into a different expression vector, pET SUMO, which

appended a small ubiquitin-like modifier (SUMO) tag at the N-terminus to improve

protein solubility (Butt et al. 2005) (Supplemental Figure S4). Upon re-expression in *E.*

coli BL21, the SUMO-tagged constructs showed clearly detectable bands at the

expected size in a denaturing gel (Supplemental Table S4, Supplemental Figure S8).

We transformed *E. coli* BL21 cells with the sulfotransferase expressing plasmids and

grew cells in 50 mL Terrific Broth at 37 ºC to mid-log phage (optical density at 600 nm ~

0.4). Then, we induced protein expression by adding 50 µL isopropyl-β-D-

thiogalactopyranoside (IPTG) to a final concentration of 1 mM, moved induced cultures

to an orbital shaker at room temperature, and continued protein expression for 18

hours. To harvest bacterial cells, we centrifuged the cell cultures (5000 x g, 10 minutes)

and froze cell pellets at -20 °C for 3 hours. We resuspended frozen cell pellets in lysis

buffer (20 mM tris hydrochloride, 500 mM NaCl, 30 mM imidazole, 0.5 % tween 20, 5

mM β-mercaptoethanol, pH 7.4), lysed cells by sonication (5 cycles of 30 s on, 10 s off

at 30 % amplitude), pelleted cellular debris by centrifuging the crude lysate (12,000 x g,

30 minutes) at 4 °C, and incubated the clarified supernatant with Ni-NTA agarose beads

at 4 °C overnight on a rotary shaker. Subsequently, we loaded the Ni-NTA slurry in a

gravity-flow chromatography column and purified recombinant protein by washing twice

with 2 mL wash buffer (20 mM tris hydrochloride, 500 mM NaCl, 30 mM imidazole pH

7.4), and eluting with 4 mL elution buffer (20 mM tris hydrochloride, 500 mM NaCl, 500

mM imidazole, pH 7.4). The eluates were concentrated and exchanged into a storage

buffer (20 mM tris hydrochloride, 300 mM NaCl, 5 mM DTT, 20 % glycerol, pH 8) by

spin filtration using 10 kDa molecular weight cutoff filters. We assessed protein purity by

running a denaturing polyacrylamide gel, visualizing the bands by activating the stain-

free gels with UV (BioRad Gel Doc), and performing densitometry analysis of protein

bands on a denaturing polyacrylamide gel (Supplemental Figure S8), then quantified the

yield via Bradford assay. Using this approach, we successfully purified all five candidate

proteins, achieving > 90 % homogeneity for the purified fractions, as determined by

densitometry analysis of protein bands resolved by denaturing gel electrophoresis

(Supplemental Table S5).

Assaying for sulfotransferase activity based on PAP formation

We freshly purified and tested sulfotransferase candidates by using a sulfotransferase

assay kit (R&D Systems) that examines the formation of 3′-phosphoadenosine-5′-

phosphate (PAP), which is released as a byproduct during biochemical sulfation. In this

assay, a coupling phosphatase is used to hydrolyze PAP, releasing phosphate, which is

quantified by adding malachite green and monitoring the increase in absorbance at 620

nm due to the formation of a complex between malachite green and phosphate. As

recommended by the kit, we prepared all reagents in 50 mM Tris, 15 mM magnesium

chloride, pH 7.5, hereafter referred to as "assay buffer". We set up the assay by

incubating 5 μL coupling phosphatase with 10 μL ostracod luciferin, 10 μL PAPS, and

25 μL sulfotransferase, resulting in a final volume of 50 μL and working concentrations

of 10 ng/µL coupling phosphatase, 100 µM ostracod luciferin, 200 µM PAPS, and 10 µM

sulfotransferase. We set up control reactions — as described above — but omitted

PAPS to prevent sulfation activity (Supplemental Figure S9). We included a positive

control to confirm the enzyme activity of the coupling phosphatase by incubating 20 µL

assay buffer with 5 µL coupling phosphatase (100 ng/µL) and 25 µL PAP (100 µM) in a

total volume of 50 µL with working concentrations of 10 ng/µL coupling phosphatase

and 50 µM PAP. Importantly, since coupling phosphatase exhibits activity with PAPS

and PAP at about a ratio of 1:350, we ran a negative control to measure this

background activity by incubating 5 µL coupling phosphatase with 10 µL ostracod

luciferin, 10 µL PAPS, and 25 µL of assay buffer, resulting in a final volume of 50 µL

and working concentrations of 10 ng/µL coupling phosphatase, 100 µM ostracod

luciferin, and 200 µM PAPS. Lastly, we included an assay blank, which contained 50 µL

of the assay buffer. We incubated the reactions at room temperature for 3 hours, then

measured the formation of PAP — catalyzed by the coupling phosphatase — by adding

malachite green and measuring the increase in absorbance at 620 nm with a microplate

reader (Tecan Spark). To test whether the sulfotransferase candidates exhibited

significant sulfation activity using the luciferin substrate, we used Dunnett's test to

compare the absorbance measurements from each of the five experimental groups with

the absorbance measurements from the negative control.

In vitro modulation of bioluminescence by luciferin sulfation

We incubated freshly purified sulfotransferase candidates with ostracod luciferin and

PAPS, then measured their ability to enzymatically modulate bioluminescence by

incubating the reaction product with ostracod luciferase and recording light production.

To set up this experiment, we mixed 10 µL of 10 µM ostracod luciferin and 100 uM

PAPS prepared in 20 mM tris hydrochloride pH 7.4, with 1 µL of each purified protein

(final concentrations ST1: 15.16 µM, ST2: 21.79 µM, ST3: 51.22 µM, ST4: 31.59 µM,

ST5: 24.63 µM). We set up control reactions — as described above — but omitted

PAPS to prevent sulfation activity. We incubated the reactions at 25 °C for 3 hours and

terminated the reaction by heating at 65 °C for 5 minutes. To measure the decrease in

bioluminescence due to the conversion of luciferin to a sulfated form, we cooled and

added 90 µL of 20 mM tris hydrochloride pH 7.4 to dilute the reaction products. We then

added 10 µL of the diluted reaction product to 55 µL purified ostracod luciferase (final

concentration ~ 136 nM), then measured luminescence using a microplate reader

(Tecan Spark) with an integration time of 2 seconds. To test whether each candidate

sulfotransferase exhibited significant sulfation activity with the luciferin substrate and

PAPS, as compared with the control reaction, we performed Student's t-tests. To

conclude whether *V. tsujii* contains a sulfotransferase gene with significant sulfation

activity towards luciferin, we corrected for multiple comparisons by performing the

Benjamini-Hochberg procedure to control the false discovery rate.

In vitro p-nitrophenol sulfation assay

We tested sulfotransferase candidates for their ability to sulfate p-nitrophenol, a

common substrate for cytosolic sulfotransferases. To set up this assay, we mixed 25 µL

purified luciferin sulfotransferase, 50 µL p-nitrophenol, and 25 µL PAPS resulting in a

total reaction volume of 100 µL and working concentrations of 4 µM purified

sulfotransferase, 100 µM p-nitrophenol, and 500 µM PAPS. We set up control reactions

— as described above — but omitted PAPS to prevent sulfation. We incubated the

reactions at room temperature for 15 hours and monitored p-nitrophenol sulfation by

measuring the decrease in absorbance at 405 nm with a microplate reader (Tecan

Spark). We used Student's t-test to test whether each candidate sulfotransferase

exhibited significant sulfation activity with the p-nitrophenol substrate as compared with

the control reaction.

Phylogenetic analysis of functionally tested sulfotransferases

Based on literature searches, we compiled sequence data from functionally tested

sulfotransferases (Supplemental Table S6, Supplemental References), and used

MAFFT (v.7.453) (Katoh and Standley 2013) to align protein sequences. We inferred a

maximum likelihood phylogeny using IQ-TREE (v. 1.6.12) (Nguyen et al. 2015),

assuming the best-fit substitution model (LG+R5) — according to Bayesian Information

Criterion — as determined with ModelFinder (Kalyaanamoorthy et al. 2017). The

phylogeny was midpoint rooted and visualized via the Interactive Tree of Life v6

(Letunic and Bork 2021).

Acknowledgements

We thank members of the Oakley Lab, especially RM Varney and NM Hensley, and CM

Pomory for discussion and feedback. This work was supported by National Science

Foundation (NSF) awards (IOS-1754770 and DEB-1457754 to THO), by 2019 (THO)

and 2021 (AM) UCSB Academic Senate Faculty Research Grants, a R35 Maximizing

Investigators' Research Award from the National Institutes of Health (5R35GM133530

to AM), a Discovery Award from the Department of Defense Congressionally Directed

Medical Research Program (PR191154 to AM), and by the US Army Research Office

under Cooperative Agreement Numbers W911NF-19-D-0001-0009 and ICB-2019-SSB-

12 for the Institute for Collaborative Biotechnologies. ESL was supported by the NSF

Graduate Research Fellowship Program (NSF GRFP 1650114) and JAG by the NSF

Postdoctoral Fellowship in Biology (BIO PRFB-1711201). The authors acknowledge the

use of the Biological Nanostructures Laboratory within the California NanoSystems

Institute, supported by the University of California, Santa Barbara and the University of

California, Office of the President. Use was made of computational facilities purchased

with funds from the National Science Foundation (CNS-1725797) and administered by

the Center for Scientific Computing (CSC). The CSC is supported by the California

NanoSystems Institute and the Materials Research Science and Engineering Center

(MRSEC; NSF DMR 1720256) at UC Santa Barbara.

Author Contributions

ESL: Conceptualization, Methodology, Software, Formal analysis, Investigation, Data

Curation, Writing - Original Draft and Review & Editing, Visualization. THO:

Conceptualization, Resources, Writing - Review & Editing, Supervision, Project

administration, Funding acquisition. AM: Methodology, Resources, Writing - Review &

Editing, Supervision, Project administration, Funding acquisition. JAG: Software, Formal

analysis, Investigation, Writing - Review & Editing, Resources, Data Curation. NTA:

Methodology, Investigation. VLL: Investigation.

Data Availability

All raw sequencing data are available on NCBI (BioProject: PRJNA935772).

Sulfotransferase sequences are available on GenBank (Accessions: OQ626349,

OQ626350, OQ626351, OQ626352, OQ626353).

Figure 1. Convergently evolved bioluminescence systems of fireflies, sea

pansies, and ostracods may vary in the extent of convergence across and within

levels of biological organization. These taxa use bioluminescence as a defense

mechanism but produce light by using structurally different organs and tissues, non-

homologous luciferases (PF00501 = AMP-dependent synthetase/ligase, PF13193 =

AMP-binding enzyme C-terminal domain, PF00561 = alpha/beta hydrolase fold, SIG =

signal peptide, PF00094 = von Willebrand factor type D domain), and different luciferins.

However, they use the same mechanism — sulfation — for modulating their luciferins.

Luciferin sulfation is catalyzed by a luciferin sulfotransferase (LST), which transfers a

sulfate from the donor 3′-phosphoadenosine-5′-phosphosulfate (PAPS) to luciferin and

produces 3'-phosphoadenosine-5'-phosphate (PAP) and sulfated luciferin, which is

unable to react with their respective luciferase. The sulfate may be removed, either by a

LST or sulfatase, to produce the active form of luciferin, which reacts with luciferase to

produce light.

Figure 2. Maximum-likelihood inferred phylogeny and normalized gene expression of sulfotransferase-domain containing proteins in *Vargula tsujii*. We inferred a maximum-likelihood phylogeny using candidate sulfotransferase sequences identified from the Iso-Seq reference transcriptome. The five most highly expressed candidates (labeled ST1-5) were selected for further biochemical testing. The color bar represents the average (N = 9 ostracods, Supplemental Figure S3) of raw transcript counts normalized to the total number of mapped reads in each ostracod sample.

Figure 3. Candidate sulfotransferases from *Vargula tsujii* exhibit luciferin

sulfation activity. Sulfotransferase activity was assayed by incubating ostracod

luciferin and PAPS with each of the 5 candidate enzymes for 3 hours (circles =

control reactions, triangles = experimental reactions). (A) The sulfotransferase

assay assesses sulfotransferase activity by using a coupling phosphatase to hydrolyze

PAP — a metabolic by-product of sulfation — to form phosphate. At the end of the

reaction, malachite green is added to bind free phosphate, forming a complex that can

be quantified via absorbance. To quantify background absorbance due to the assay

buffer, we included a "blank" measurement. We included a positive (+) control — which

consisted of coupling phosphatase and PAP — to confirm the enzyme activity of the

coupling phosphatase. Compared to the negative (–) control reactions lacking

sulfotransferase, four sulfotransferases — ST1, ST3, ST4, ST5 — showed significant

luciferin sulfation activity (N = 6). (B) The luciferase assay uses luminescence to assess

luciferin sulfation activity. If a candidate's luciferin sulfation activity is sufficient to

modulate bioluminescence, the product of their sulfation reaction should produce less

bioluminescence upon addition of ostracod luciferase. Compared to the control

reactions lacking PAPS, three candidates — ST2, ST3, and ST5 — exhibit decreased

light production indicative of sulfation activity sufficient to modulate bioluminescence *in*

vitro (N = 3). Each datum is an average of three technical replicates.

Figure 4. Four candidates sulfate p-nitrophenol. Sulfation of p-nitrophenol, a

common substrate for many mammalian sulfotransferases, was assayed by measuring

change in absorbance at 405 nm following a 15 hour incubation with purified

sulfotransferase proteins in the absence (circles) and presence (triangles) of PAPS.

Four candidates — ST2, ST3, ST4, and ST5 — catalyze significant p-nitrophenol

sulfation under the conditions tested (N = 3).

Figure 5. Maximum-likelihood inferred phylogeny of functionally tested

sulfotransferases. Luciferin sulfotransferases from luminous organisms are denoted by

colored branches (firefly = red, ostracods = blue, sea pansies = purple). Invertebrate

sulfotransferases are highlighted in gray boxes. Colored squares (red = firefly luciferin,

blue = ostracod luciferin, purple = sea pansy luciferin, orange = p-nitrophenol, green =

other) indicate functionally demonstrated substrates for each sulfotransferase; gray

squares indicate molecules that cannot be sulfated by the respective sulfotransferase;

white squares indicate molecules that have not yet been tested. Accession numbers for

each sulfotransferase are listed on the right.

References

Adams ST Jr, Miller SC. 2020. Enzymatic promiscuity and the evolution of

bioluminescence. *FEBS J.* 287:1369–1380.

Allali-Hassani A, Pan PW, Dombrovski L, Najmanovich R, Tempel W, Dong A, Loppnau

P, Martin F, Thornton J, Edwards AM, et al. 2007. Structural and chemical profiling

of the human cytosolic sulfotransferases. *PLoS Biol.* 5:e97.

Atkinson DE. 1966. Regulation of Enzyme Activity. *Annu. Rev. Biochem.* 35:85–124.

Bidwell LM, McManus ME, Gaedigk A, Kakuta Y, Negishi M, Pedersen L, Martin JL.

1999. Crystal structure of human catecholamine sulfotransferase. *J. Mol. Biol.*

293:521–530.

Blount ZD, Lenski RE, Losos JB. 2018. Contingency and determinism in evolution:

Replaying life's tape. *Science* [Internet] 362. Available from:

http://dx.doi.org/10.1126/science.aam5979

Branham MA, Wenzel JW. 2003. The origin of photic behavior and the evolution of

sexual communication in fireflies (Coleoptera: Lampyridae). *Cladistics* 19:1–22.

Bushmanova E, Antipov D, Lapidus A, Prjibelski AD. 2019. rnaSPAdes: a de novo

transcriptome assembler and its application to RNA-Seq data. *GigaScience*

[Internet] 8. Available from: http://dx.doi.org/10.1093/gigascience/giz100

Butt TR, Edavettal SC, Hall JP, Mattern MR. 2005. SUMO fusion technology for difficult-

to-express proteins. *Protein Expr. Purif.* 43:1–9.

Chapman E, Best MD, Hanson SR, Wong C-H. 2004. Sulfotransferases: structure,

mechanism, biological activity, inhibition, and synthetic utility. *Angew. Chem. Int. Ed*

Engl. 43:3526–3548.

Clarke GL, Conover RJ, David CN, Nicol JAC. 1962. Comparative studies of

luminescence in copepods and other pelagic marine animals*. *J. Mar. Biol. Assoc.*

U. K. 42:541–564.

Copeland RA. 2000. Enzymes: A Practical Introduction to Structure, Mechanism, and

Data Analysis. John Wiley & Sons

Cormier MJ, Kazuo H. 1964. Studies on the bioluminescence of Renilla reniformis.

Biochimica et Biophysica Acta (BBA) - Specialized Section on Biophysical Subjects

88:99–104.

Delroisse J, Duchatelet L, Flammang P, Mallefet J. 2021. Leaving the Dark Side?

Insights Into the Evolution of Luciferases. *Frontiers in Marine Science* [Internet] 8.

Available from: https://www.frontiersin.org/articles/10.3389/fmars.2021.673620

Dilly PN, Herring PJ. 1978. The light organ and ink sac of Heteroteuthis dispar

(Mollusca: Cephalopoda). *J. Zool.* 186:47–59.

Driscoll WJ, Komatsu K, Strott CA. 1995. Proposed active site domain in estrogen sulfotransferase as determined by mutational analysis. *Proc. Natl. Acad. Sci. U. S. A.* 92:12328–12332.

Ellis EA, Goodheart JA, Hensley NM, González VL, Reda NJ, Rivers TJ, Morin JG, Torres E, Gerrish GA, Oakley TH. 2022. Sexual signals persist over deep time: ancient co-option of bioluminescence for courtship displays in cypridinid ostracods. *Syst. Biol.* [Internet]. Available from: http://dx.doi.org/10.1093/sysbio/syac057

Falany CN. 1997. Enzymology of human cytosolic sulfotransferases. *FASEB J.* 11:206–216.

Fallon TR, Li F-S, Vicent MA, Weng J-K. 2016. Sulfoluciferin is Biosynthesized by a Specialized Luciferin Sulfotransferase in Fireflies. *Biochemistry* 55:3341–3344.

Foster CSP, Van Dyke JU, Thompson MB, Smith NMA, Simpfendorfer CA, Murphy CR, Whittington CM. 2022. Different Genes are Recruited During Convergent Evolution of Pregnancy and the Placenta. *Mol. Biol. Evol.* 39:msac077.

Fu L, Niu B, Zhu Z, Wu S, Li W. 2012. CD-HIT: accelerated for clustering the next-generation sequencing data. *Bioinformatics* 28:3150–3152.

Goodheart JA, Minsky G, Brynjegard-Bialik MN, Drummond MS, David Munoz J, Fallon TR, Schultz DT, Weng J-K, Torres E, Oakley TH. 2020. Laboratory culture of the California Sea Firefly Vargula tsujii (Ostracoda: Cypridinidae): Developing a model system for the evolution of marine bioluminescence. *Sci. Rep.* 10:1–15.

Haddock SHD, Moline MA, Case JF. 2010. Bioluminescence in the Sea. *Ann. Rev. Mar. Sci.* 2:443–493.

Harvey EN. 1952. Bioluminescence Academic Press. *New York, NY USA* 649.

Hori K, Nakano Y, Cormier MJ. 1972. Studies on the bioluminescence of Renilla

reniformis. XI. Location of the sulfate group in luciferyl sulfate. *Biochim. Biophys.*

Acta 256:638–644.

Hulet WH, Musil G. 1968. Intracellular Bacteria in the Light Organ of the Deep Sea

Angler Fish, Melanocetus murrayi. *Copeia* 1968:506–512.

Huvard AL. 1993. Ultrastructure of the light organ and immunocytochemical localization

of luciferase in luminescent marine ostracods (Crustacea: Ostracoda:

Cypridinidae). *J. Morphol.* 218:181–193.

Inouye S. 2007. Expression, purification and characterization of calcium-triggered

luciferin-binding protein of Renilla reniformis. *Protein Expr. Purif.* 52:66–73.

James MO. 2021. Enzyme Kinetics of PAPS-Sulfotransferase. *Methods Mol. Biol.*

2342:285–300.

Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. 2017.

ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat.*

Methods 14:587–589.

Kastendiek J. 1976. Behavior of the sea pansy Renilla kollikeri Pfeffer (Coelenterata:

Pennatulacea) and its influence on the distribution and biological interactions of the

species. *Biol. Bull.* 151:518–537.

Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7:

improvements in performance and usability. *Mol. Biol. Evol.* 30:772–780.

Khersonsky O, Roodveldt C, Tawfik DS. 2006. Enzyme promiscuity: evolutionary and

mechanistic aspects. *Curr. Opin. Chem. Biol.* 10:498–508.

Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nat.*

Methods 9:357–359.

Lau ES, Oakley TH. 2020. Multi-level convergence of complex traits and the evolution of

bioluminescence. *Biol. Rev. Camb. Philos. Soc.* [Internet]. Available from:

https://onlinelibrary.wiley.com/doi/10.1111/brv.12672

Lee DH, Mittag M, Sczekan S, Morse D, Hastings JW. 1993. Molecular cloning and

genomic organization of a gene for luciferin-binding protein from the dinoflagellate

Gonyaulax polyedra. *J. Biol. Chem.* 268:8842–8850.

Letunic I, Bork P. 2021. Interactive Tree Of Life (iTOL) v5: an online tool for

phylogenetic tree display and annotation. *Nucleic Acids Res.* 49:W293–W296.

Li W, Godzik A. 2006. Cd-hit: a fast program for clustering and comparing large sets of

protein or nucleotide sequences. *Bioinformatics* [Internet] 22:1658–1659. Available

from: http://dx.doi.org/10.1093/bioinformatics/btl158

Lorenz WW, McCann RO, Longiaru M, Cormier MJ. 1991. Isolation and expression of a

cDNA encoding Renilla reniformis luciferase. *Proc. Natl. Acad. Sci. U. S. A.*

88:4438–4442.

Meyer E, Aglyamova GV, Matz MV. 2011. Profiling gene expression responses of coral

larvae (Acropora millepora) to elevated temperature and settlement inducers using

a novel RNA-Seq procedure. *Mol. Ecol.* 20:3599–3616.

Morin JG. 1983. Coastal Bioluminescence: Patterns and Functions. *Bull. Mar. Sci.*

33:787–817.

Nakamura M, Matsuda K, Nakamura M, Yamashita K, Suzuki T, Inouye S. 2019.

Enzymatic Conversion of Cypridina Luciferyl Sulfate to Cypridina Luciferin with

Coenzyme A as a Sulfate Acceptor in Cypridina (Vargula) hilgendorfii. *Photochem.*

Photobiol. 95:1376–1386.

Nakamura M, Suzuki T, Ishizaka N, Sato J-I. 2014. Identification of 3-enol sulfate of

Cypridina luciferin, Cypridina luciferyl sulfate, in the sea-firefly Cypridina (Vargula)

hilgendorfii. *Tetrahedron* 70:2161–2168.

Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. 2015. IQ-TREE: a fast and

effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol.*

Biol. Evol. 32:268–274.

Oakley TH. 2005. Myodocopa (Crustacea: Ostracoda) as models for evolutionary

studies of light and vision: multiple origins of bioluminescence and extreme sexual

dimorphism. *Hydrobiologia* 538:179–192.

Oba Y, Kumazaki M, Inouye S. 2010. Characterization of luciferases and its paralogue

in the Panamanian luminous click beetle Pyrophorus angustus: A click beetle

luciferase lacks the fatty acyl-CoA synthetic activity. *Gene* 452:1–6.

Oba Y, Ojika M, Inouye S. 2003. Firefly luciferase is a bifunctional enzyme: ATP-

dependent monooxygenase and a long chain fatty acyl-CoA synthetase. *FEBS Lett.*

540:251–254.

Oba Y, Sato M, Ohta Y, Inouye S. 2006. Identification of paralogous genes of firefly

luciferase in the Japanese firefly, Luciola cruciata. *Gene* 368:53–60.

O'Brien PJ, Herschlag D. 1999. Catalytic promiscuity and the evolution of new

enzymatic activities. *Chem. Biol.* 6:R91–R105.

Orgogozo V. 2015. Replaying the tape of life in the twenty-first century. *Interface Focus*

5:20150057.

Pankey MS, Minin VN, Imholte GC, Suchard MA, Oakley TH. 2014. Predictable

transcriptome evolution in the convergent and complex bioluminescent organs of squid. *Proc. Natl. Acad. Sci. U. S. A.* 111:E4736–E4742.

Peterson MK, Buck J. 1968. Light organ fine structure in certain Asiatic fireflies. *Biol. Bull.* 135:335–348.

Powell GS, Saxton NA, Pacheco YM, Stanger-Hall KF, Martin GJ, Kusy D, Felipe Lima Da Silveira L, Bocak L, Branham MA, Bybee SM. 2022. Beetle bioluminescence outshines extant aerial predators. *Proceedings of the Royal Society B: Biological Sciences* 289:20220821.

Rees JF, De Wergifosse B, Noiset O, Dubuisson M, Janssens B, Thompson EM. 1998. The origins of marine bioluminescence: Turning oxygen defence mechanisms into deep-sea communication tools. *J. Exp. Biol.* 201:1211–1221.

Revell LJ. 2012. phytools: an R package for phylogenetic comparative biology (and other things). *Methods in Ecology and Evolution* 3:217–223.

Rosenblum EB, Parent CE, Brandt EE. 2014. The Molecular Basis of Phenotypic Convergence. *Annu. Rev. Ecol. Evol. Syst.* 45:203–226.

Satterlie RA, Case JF. 1979. Development of bioluminescence and other effector responses in the pennatulid coelenterate Renilla köllikeri. *Biol. Bull.* 157:506–523.

Shimomura O. 2019. Bioluminescence: Chemical Principles and Methods. World Scientific

Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. 2015. BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics* 31:3210–3212.

Spurlock BO, Cormier MJ. 1975. A fine structure study of the anthocodium in Renilla

mülleri: Evidence for the existence of a bioluminescent organelle, the luminelle. *J. Cell Biol.* 64:15–28.

Stayton CT. 2015. What does convergent evolution mean? The interpretation of convergence and its implications in the search for limits to evolution. *Interface Focus* 5:20150039.

Takenaka Y, Yamaguchi A, Tsuruoka N, Torimura M, Gojobori T, Shigeri Y. 2012. Evolution of bioluminescence in marine planktonic copepods. *Mol. Biol. Evol.* 29:1669–1681.

Thacker CE, Roje DM. 2009. Phylogeny of cardinalfishes (Teleostei: Gobiiformes: Apogonidae) and the evolution of visceral bioluminescence. *Mol. Phylogenet. Evol.* 52:735–745.

Tokuriki N, Tawfik DS. 2009. Protein dynamism and evolvability. *Science* 324:203–207.

Tsarkova AS. 2021. Luciferins Under Construction: A Review of Known Biosynthetic Pathways. *Frontiers in Ecology and Evolution* [Internet] 9. Available from: https://www.frontiersin.org/articles/10.3389/fevo.2021.667829

Tzertzinis G, Baker B, Benner J, Brown E, Corrêa IR Jr, Ettwiller L, McClung C, Schildkraut I. 2022. Coelenterazine sulfotransferase from Renilla muelleri. *PLoS One* 17:e0276315.

Viviani VR, Silva JR, Amaral DT, Bevilaqua VR, Abdalla FC, Branchini BR, Johnson CH. 2020. A new brilliantly blue-emitting luciferin-luciferase system from Orfelia fultoni and Keroplatinae (Diptera). *Sci. Rep.* 10:9608.

Waterhouse RM, Seppey M, Simão FA, Manni M, Ioannidis P, Klioutchnikov G, Kriventseva EV, Zdobnov EM. 2018. BUSCO Applications from Quality

1   Assessments to Gene Prediction and Phylogenomics. *Mol. Biol. Evol.* 35:543–548.

2   Widder EA. 2010. Bioluminescence in the Ocean: Origins of Biological, Chemical, and

3   Ecological Diversity. *Science* [Internet] 328:704–708. Available from:

4   http://dx.doi.org/10.1126/science.1174269