Parallel shifts in flight-height associated with altitude across incipient

Heliconius species

David F Rivas-Sánchez[1,2], Carlos H. Gantiva-Q[3], Carolina Pardo-Diaz[3], Camilo Salazar[3], Stephen H

Montgomery[1]*, Richard M Merrill[2]*

Affiliations:

[1] School of Biological Sciences, University of Bristol, 24 Tyndall Avenue, Bristol, UK

[2] Division of Evolutionary Biology, Ludwig-Maximilians-Universität München, Grosshaderner Strasse 2, 82152

Planegg-Martinsried, Germany.

[3] Department of Biology, Faculty of Natural Sciences, Universidad del Rosario, Carrera 24 # 63C-69, Bogotá,

111221, Colombia

* Contributed equally

Corresponding authors:

DFRS: gb20421@bristol.ac.uk

SHM: s.montgomery@bristol.ac.uk

RMM: merrill@bio.lmu.de

18
19
Abstract

Vertical gradients in microclimate, resource availability and interspecific interactions

are thought to underly stratification patterns in tropical insect communities. However,

only a few studies have explored the adaptive significance of vertical space use

during early population divergence. We analysed flight-height variation across

speciation events in *Heliconius* butterflies representing parallel instances of

divergence between low and high-altitude populations. We measured flight-height in

wild *H. erato venus* and *H. chestertonii*, lowland and mountain specialists

respectively, and found that *H. chestertonii* is consistently flies at a lower height. We

compared these data with previously published results for *H. e. cyrbia* and *H.

himera,* the latter of which flies lower and like *H. chestertonii*, recently colonised

high-altitude, dry forests. We provide evidence that these repeated trends are largely

driven by shared sources of selection across equivalent environments, producing

similar trait-shifts in *H. himera* and *H. chestertonii*. Although our results imply a

signature of local adaptation, we did not find an association between resource

distribution and flight-height in *H. e. venus* and *H. chestertonii*. We discuss how

these results may be explained by variation in forest structure and the microclimate.

Overall, our findings underscore the importance of behavioural adjustments during

early divergence mediated by altitude-shifts.

39
40
41 ## Key words

42 *Heliconius*, local adaptation, flight-height, parallel evolution.
43
Introduction

The steep environmental gradients that characterise the upper and lower vegetation
layers of tropical forests are a key factor determining species abundance and
diversity [1-5]. In arthropods, particular species are consistently found in some
vertical strata while being less common in others due to variation in abiotic
conditions, behaviour, and the distribution of resources [6, 7]. For example, fruit-
feeding and nectar-feeding butterflies, respectively decrease and increase in
abundance towards the canopy, ostensibly tracking their food sources [8-10]. This
has broad implications for the structure of communities, with consequences for wider
trophic interactions [11, 12].
54
Comparatively, less is known about vertical segregation occurring between closely
related populations. Only a handful of studies have examined divergence in the use
of vertical space at this level, but these provide a range of evidence implicating flight
differences in early divergence. For instance, closely related *Heliconius* species that
that fly in separate vertical strata have different wing morphologies [13]. Similarly, the
Nymphalid *Archaeopreponae demophon* includes coexisting canopy and understory
populations with independent demographic dynamics and low levels of gene-flow,
despite their similarity and proximity [14]. Divergence in resource use, combined with
changes in larval behaviour, have also been linked to species divergence in
Colobura dirce [15], which can be split into two separate species based on
differences between their larvae: *C. annulata* are gregarious and feed on mature,
canopy *Cecropia* leaves while *C. dirce* are solitary and consume understory saplings
[15]. These examples suggest a role of vertical space in local adaptation, but the
underlying mechanisms remain unclear.
69
Recently, heritable flight-height differences were demonstrated between the
Heliconius species *H. e. cyrbia* and *H. himera* [16], which are distributed along a
gradient from wet, low-altitude to dry, high-altitude forests, with a narrow zone of
overlap where the two hybridise [17]. This correlates with the distribution of
resources in their habitats such that *H. himera* flies lower than *H. e. cyrbia*, tracking
the height of hostplants and flowers that are positioned lower at high-altitude [16].
This pattern may result from divergent ecological selection maximising hostplant
encounter rates and foraging efficiency across forests with different structure,
potentially contributing, alongside other factors [18-21], to the evolution of
reproductive isolation.
80
In the western slope of the Colombian Cordillera Occidental, *H. e. venus* and *H.
chestertonii* are distributed along a similar ecological gradient (Rivas-Sánchez et al.,
in press) (Figure 1 d-e). *H. e. venus* occupies wet forests at sea level, whereas its
close relative *H. chestertonii* lives in drier, scrub forests in the mountains [22, 23].
Because *H. chestertonii* is spatially and genetically independent from, but
ecologically comparable to *H. himera*, it can be studied as a convenient replica to
explore the relationship between vertical segregation and divergence during the
early stages of speciation. Here, we tested whether *H. e. venus* and *H. chestertonii*
fly at different heights and if this correlates with the location of larval and adult-stage
resources in their respective habitats. Under the assumption that flight-height
patterns constitute adaptive traits, we predicted that flight-height variation between
H. e. venus and *H. chestertonii* would mirror that between their ecological
equivalents *H. e. cyrbia* and *H. himera*.
94

Methods

i) Habitat characterisation

We investigated *Heliconius* butterflies in four sites representing the habitats of *H. e. venus* (Buenaventura, N 03º50'0.04" W77º15'45.1" and La Barra, N 03°57.558420 W 77°22.705080) and *H. chestertonii* (Montañitas N 03º40'36.3" W 76º31'19.4" and El Saladito N 03º29'11.5" W 76º36'39.8") in January 2021. At each of these sites, we characterised the local forest structure by measuring the trunk diameter at breast height (m) and the average distance (m) between neighbouring trees of common tree species. Additionally, we used clinometers and measuring tapes to estimate the height (m) of the canopy, the understory, and the shrubs in each site.

ii) Vertical distribution of resources

We explored whether the distribution of adult and larval resources differed between habitats. To do this, we measured the height (m) of all the *Passiflora* (Passifloraceae) hostplants that were visually identified along the trails used to collect butterflies. These plants were subsequently identified to the species level [24, 25] and filtered so that our dataset consisted exclusively of species known to be used by butterflies in the *erato* complex [26, 27], to which our focal species belong. We also measured the height (m) of the flowering plants that adult butterflies were observed feeding from at least once. Note that these methods differ slightly from Dell'Aglio, Mena [16], where plants were identified by following butterflies until they landed on a flower or hostplant. Due to the trail and forest structure, this approach was not possible here.

iii) Patterns of flight-height divergence

We measured the flight-height (m) of *H. e. venus* (n=33) and *H. chestertonii* (n=39) at the point where individuals were first encountered using measuring tapes. The butterflies were collected, sexed and marked, ensuring no repeated observations. To quantify levels of parallelism in flight-height with butterflies from an independent, ecologically equivalent pair, we included flight-height measures of *H. e. cyrbia* (n=78) and *H. himera* (n=86) in our dataset, available in Dell'Aglio, Mena [16].

iv) Statistical analyses

To test for significant differences in the diameter at the breast height, the average distance between neighbouring trees, and the height of different forest layers between habitats, we used Wilcoxon rank sum tests implemented in R [28]. We also modelled variation in the height of hostplants and flowers separately using linear mixed-effect models (implemented in *lme4* [29]) with "habitat" as a fixed factor and "site" as a random effect. This was followed by Wald χ^2 tests (implemented with the *Anova* function in the package *CAR* [30]). To analyse flight in *H. e. venus* and *H. chestertonii*, we used linear-mixed models where variation in flight-height was explained by the fixed factors "subspecies" and "sex", as well as their interaction and the random factor "site". We subsequently used the *drop1* function in base R to

142 remove model terms unless their deletion was detectable via likelihood tests. This
143 resulted in a reduced model with subspecies as the only main effect.
144
Finally, to test for parallel shifts in flight-height across independent divergence
events, we used a variance partitioning approach [31] combining data on *H. e.*
venus, *H. chestertonii*, *H. e. cyrbia* and *H. himera*. In detail, in our model we included
the main factors "habitat" (similarity of selection regimes across independent
divergence events) and "locality" (variation in flight-height that responds to the
unique properties of each divergence), as well as a "habitat*locality" interaction term
(inconsistency in magnitude and/or direction of divergence patterns of the two pairs
of species). We assessed the significance of each of these terms via likelihood (*drop*
1 function) and Wald χ^2 tests between alternative models with and without specific
factors. We then estimated semi-partial R^2 coefficients (*r2beta* function, Kenward
Roger method in the R package r2glmm [32]) in our final model, providing a rough
approximation of the proportion of variance explained by each term.
157
Results
159
The habitats of *H. e. venus* and *H. chestertonii* broadly differed (Figure 1a-c). In
particular, the trunk diameter at the breast height (W=32, p<0.05) and the distance
between neighbouring trees (W=15, p<0.01) were both significantly higher in the
habitat of *H. e. venus*. In addition, the height of the canopy and the shrubs forest
layers were not significantly different between the two forest types, but the
understory layer was significantly shorter in the habitat of *H. chestertonii* (W=30,
p<0.01). in contrast, we were unable to confirm significant differences in the vertical
distribution of hostplants (n= 37, χ_1^2=0.3949; p=0.5297) or flower resources (n=107,
χ_1^2=0.0096; p=0.9219; Figure 2a-b) between different habitats.
169
On average, *H. chestertonii* flew ~0.5 m lower than *H. e. venus* (n=74, χ_1^2=8.8601;
p< 0.01; Figure 2c) suggesting a parallel shift in flight behaviour occurred in *H.*
chestertonii and *H. himera*. Indeed, in our full model combining data from *H. e.*
venus, *H. chestertonii*, *H. e. cyrbia* and *H. himera*, the proportion of variance
attributable to "habitat" was higher than the attributable to differences in localities or
habitat*locality interactions (Table 1). Moreover, likelihood ratio comparisons
between alternative models showed that the effect of "habitat" was significant
(n=240, χ_1^2=21.725; p<0.001) but not the effect of "locality" nor the interaction
between them, suggesting a close correspondence between the patterns of
divergence in flight-height in *H. e. venus-H. chestertonii* and *H. e. cyrbia-H. himera*
(Figure 2d). In our reduced model with "habitat" as the only main factor, "habitat"
explained more than half of the variation flight-height between species (Table 1). .
182
183
Discussion
185
We leveraged two separate events of evolutionary divergence across ecologically
equivalent *Heliconius* taxa pairs in Colombia and Ecuador to explore local adaptation
in the context of altitude shifts. We demonstrate that the mountain specialist *H.*
chestertonii flies at a lower height compared to its ancestral-proxy, low-altitude
population, *H. e. venus*, closely mirroring a pattern formerly reported for *H. himera*
(high-altitude) and *H. e. cyrbia* (low-altitude) [16]. This strongly implies a role for

shared sources of divergent selection across habitat types driving local adaptation in
flight-height. However, we were not able to formally link shifts in flight-height in *H.*
chestertoniini to the distribution of plant resources.
195
Adaptation can be inferred when phenotypic differentiation takes place in parallel
across replicate populations from similar environments [31, 33, 34]. The derived
populations of *H. chestertonii* and *H. himera* independently colonised similar
mountainous habitats in the Eastern Andes, as evidenced from genomic data which
isolates these species in separate genetic clusters [22, 35]. Although we measured
flight height in wild individuals, the differences we found may be heritable, as
demonstrated in common garden experiments with *H. e. cyrbia* and *H. e. himera*
[16], and therefore subject to selection. Indeed, our tests for parallelism suggest
these independent shifts in flying behaviour are largely driven by shared sources of
divergent selection rather than stochastic evolutionary processes (e.g., genetic drift
or gene-flow) [31, 34, 36] and might thus constitute a signature of repeated local
adaptation (e. g. [37-41]).
208
In contrast to *H. e. cyrbia* and *H. himera* [16], we found no evidence for an
association between flight-height and the distribution of larval or adult resources in
H. e. venus or *H. chestertonii*. In other species, such correlations have been
interpreted as evidence that butterflies fly at a height that maximises encounter rates
with resources [8, 9, 42, 43]. Passion vines are strongly influenced by the forest
architecture because they rely on multiple plant species for structural support [25]. *P.*
auriculata was the only suitable hostplant in the *H. e. venus* habitat, whereas we
found both *P. biflora* and *P. suberosa* in the *H. chestertonii* habitat. These three
species belong to the same subgenus (*Decaloba*) and grow to modest dimensions
[24], likely explaining their similar heights in the wild. This lack of differentiation
conflicts with the hypothesis that flight height differences are a result of selection to
maximise foraging and oviposition opportunities. Interestingly, in the lower altitude *H.*
e. venus La Barra site, we found a few plants of a fourth species, *P. micropetala,*
which grow up to 6 m tall. We did not include these plants in our analyses (although
their inclusion does not qualitatively effect our results due to small sample size), as it
is unknown if *H. erato* uses *P. micropetala*. Nevertheless, *P. micropetala* is a very
near relative to *P. biflora* [24] and *Heliconius erato* are known to switch hosts
seasonally when their preferred hostplant is depleted [44-47]. Regardless, the
presence of these taller *Passiflora* in the *H. e. venus* habitat resonates with the idea
that the lowland, forest understory offers support for taller *Passiflora* [16, 25]. Further
studies could confirm whether *P. micropetala* is indeed used by *H. e. venus*, which
may relate to the flight-height differences we observed in respect to *H. chestertonii*.
231
232
In Colombia, the structure of high and low-altitude forests differs such that the
mountainous habitat of *H. chestertonii* has generally shorter trees at higher densities.
This high-altitude forest is also colder and drier (~ 7 °C and ~3% RH differences)
(Rivas et al., in press). These differences mirror those between the habitats of *H. e.*
cyrbia and *H. himera* [16] and may potentially result in alternative sources of
divergent selection across both pairs of taxa. For instance, oviposition behaviour can
be affected by microclimatic conditions [48, 49]. Using dummy models, Merckx, Van
Dongen [50] also showed that the woodland butterfly, *Pararge aegeria*, can warm up
faster when ambient temperatures are higher, and experiences less convective

cooling at lower flight-heights and in forests with a more sheltered structure.
Pupation height is also partly driven by desiccation avoidance, which is lower closer
to the ground [51]. It is therefore plausible that the butterflies in our study are flight-
height adjusted to compensate for habitat-differences that affect temperature-
dependent traits, rather than to match resource availability.
247
In conclusion, we showed that two derived mountain specialist *Heliconius* butterflies
have independently evolved flight behaviours via trait-shifts that are similar in
magnitude and direction, suggesting underlying repeated local adaptations in forests
at different elevations. In Colombia, this pattern does not appear to be driven by
differences in the vertical distribution of hostplants or nectar and pollen rich flowers.
We suggest that these parallel differences may instead relate to effects of variable
microclimatic conditions between forest types. These results add to a limited list of
studies exploring potential mechanisms explaining flight-height differences between
closely related insect taxa in the tropics.
257
258
259
260 **Figures and table**
261
262
263

264
265
Figure 1. Broad characteristics of low-altitude (blue) and high-altitude (red) habitats. Comparison of a)
diameter at the breast height (DBH), b) average distance between neighbouring trees (ADNT), c)
canopy, understory, and shrubs heights, d) average temperature and e) average relative humidity
throughout the day. In a-c), box plots show the median and upper and lower quartiles, each point is
an individual measure. Asterisks convey significance levels of comparisons (Wilcoxon test). NS, not

significant. * p < 0.05. ** p < 0.01. *** P < 0.001. In d-e) (modified from Rivas *et al* 2022 in press),

each line represents microclimate data point recorded by a data logger deployed in low-altitude (blue)

or high-altitude (red) forest.

277
278
279
Figure 2. Comparisons in resource distribution and flying-heights in *Heliconius* species. a)

Comparison of hostplant heights, b) adult-food source heights, c) flight-height and d) levels of

parallelism in flight-height between species from different clines. For a) circles are *Passiflora*

auriculata, squares are *P. micropetala*, pluses are *P. suberosa* and triangles *P. biflora*. For a-c) blue

H. e. venus is blue and *H. chestertonii* is red. For d). *H. e. venus* is blue and *H. chestertonii* is red, *H.*

e. cyrbia is dodger blue and *H. himera* is firebrick. Box plots show the median and upper and lower

quartiles, each point is an individual measure. Asterisks convey significance levels of comparisons

(Wilcoxon test). NS, not significant. * p < 0.05. ** p < 0.01. *** P < 0.001.

Table 1. Semi-partial R2 measures of model terms in models explaining variation in flight-height

across *H. himera*, *H. e. cyrbia*, *H. chestertonii* and *H. e. venus*.

Semi-partial R² measures (95% CI)				
Model	**Factor**	**Semi-partial R²**	**Lower CI**	**Upper CI**
	Habitat	0.451	0.087	0.784
Flight-height~Habitat:Locality	Locality	0.234	0.003	0.661
	Habitat x Locality	0.162	0.001	0.605
Flight-height~Habitat	Habitat	0.546	0.233	0.8

292
293

References

1. Scheffers, B.R., et al., *Increasing arboreality with altitude: a novel biogeographic dimension.* Proceedings of the Royal Society B: Biological Sciences, 2013. **280**(1770): p. 20131581.
2. Basset, Y., et al., *Stratification and diel activity of arthropods in a lowland rainforest in Gabon.* Biological Journal of the Linnean Society, 2001. **72**(4): p. 585-607.
3. Basset, Y., et al., *Arthropod Distribution in a Tropical Rainforest: Tackling a Four Dimensional Puzzle.* PLOS ONE, 2015. **10**(12): p. e0144110.
4. Basham, E.W., et al., *Vertical stratification patterns of tropical forest vertebrates: a meta-analysis.* Biological Reviews, 2022.
5. Jucker, T., et al., *A research agenda for microclimate ecology in human-modified tropical forests.* Frontiers in Forests and Global Change, 2020. **2**: p. 92.
6. Basset, Y., et al., *Vertical stratification of arthropod assemblages.* Arthropods of tropical forests, 2003: p. 17-27.
7. Wardhaugh, C.W., *The spatial and temporal distributions of arthropods in forest canopies: uniting disparate patterns with hypotheses for specialisation.* Biological Reviews, 2014. **89**(4): p. 1021-1041.
8. Schulze, C.H., K.E. Linsenmair, and K. Fiedler, *Understorey versus Canopy: Patterns of Vertical Stratification and Diversity among Lepidoptera in a Bornean Rain Forest.* Plant Ecology, 2001. **153**(1/2): p. 133-152.
9. De Vries, P., *Stratification offruit-feeding nymphalid butterflies in.* Journal of Research on the Lepidoptera, 1988. **26**(1-4): p. 98-108.
10. Schulze, C.H., *Habitat preference and flight activity of Morphinae butterflies in a Bornean rainforest, with a note on sound production by adult Zeuxidia (Lepidoptera: Nymphalidae).* Malayan Nature Journal, 1998. **52**: p. 163-176.
11. Elias, M., et al., *Mutualistic interactions drive ecological niche convergence in a diverse butterfly community.* PLoS biology, 2008. **6**(12): p. e300.
12. Willmott, K.R., et al., *Maintaining mimicry diversity: Optimal warning colour patterns differ among microhabitats in Amazonian clearwing butterflies.* Proceedings of the Royal Society B: Biological Sciences, 2017. **284**(1855).
13. Mena, S., et al., *Forest stratification shapes allometry and flight morphology of tropical butterflies.* Proceedings of the Royal Society B, 2020. **287**(1937): p. 20201071.
14. Nice, C.C., et al., *Vertical differentiation in tropical forest butterflies: a novel mechanism generating insect diversity?* Biology letters, 2019. **15**(1): p. 20180723.
15. Willmott, K., L. Constantino, and J. Hall, *A review of Colobura (Lepidoptera: Nymphalidae) with comments on larval and adult ecology and description of a sibling species.* Annals of the Entomological Society of America, 2001. **94**(2): p. 185-196.
16. Dell'Aglio, D.D., et al., *Divergence in Heliconius flight behaviour is associated with local adaptation to different forest structures.* Journal of Animal Ecology, 2022. **91**(4): p. 727-737.

17. Jiggins, C.D., et al., *What can hybrid zones tell us about speciation? The case of Heliconius erato and H. himera (Lepidoptera: Nymphalidae).* Biological Journal of the Linnean Society, 1996. **59**(3): p. 221-242.

18. McMillan, W.O., C.D. Jiggins, and J. Mallet, *What initiates speciation in passion-vine butterflies?* Proceedings of the National Academy of Sciences, 1997. **94**(16): p. 8628-8633.

19. Davison, A., et al., *Behavioral and Physiological Differences between Two Parapatric Heliconius Species 1.* Biotropica, 1999. **31**(4): p. 661-668.

20. Merrill, R.M., A. Chia, and N.J. Nadeau, *Divergent warning patterns contribute to assortative mating between incipient Heliconius species.* Ecology and Evolution, 2014. **4**(7): p. 911-917.

21. Montgomery, S. and R. Merrill, *Divergence in brain composition during the early stages of ecological specialization in Heliconius butterflies.* Journal of Evolutionary Biology, 2017. **30**(3): p. 571-582.

22. Arias, C.F., et al., *A hybrid zone provides evidence for incipient ecological speciation in Heliconius butterflies.* Molecular ecology, 2008. **17**(21): p. 4699-4712.

23. Brower, A.V., *Parallel race formation and the evolution of mimicry in Heliconius butterflies: a phylogenetic hypothesis from mitochondrial DNA sequences.* Evolution, 1996. **50**(1): p. 195-221.

24. Ulmer, T. and J.M. MacDougal, *Passiflora: passionflowers of the world*. 2004: Timber Press (OR).

25. Pérez, J.O., *Diversity of Colombian Passifloraceae: biogeography and an updated list for conservation.* Biota Colombiana, 2007. **8**(1).

26. Jiggins, C.D., *The ecology and evolution of Heliconius butterflies*. 2017: Oxford University Press.

27. Robinson, G.S., et al., *HOSTS-a Database of the World's Lepidopteran Hostplants.* Natural History Museum, London, 2010.

28. Team, R.C., *R: A language and environment for statistical computing. R Foundation for Statistical Computing.* 2020.

29. Bates, D., et al., *Fitting linear mixed-effects models using lme4.* arXiv preprint arXiv:1406.5823, 2014.

30. Fox, J. and S. Weisberg, *An R companion to applied regression*. 2018: Sage publications.

31. Langerhans, R.B. and T.J. DeWitt, *Shared and unique features of evolutionary diversification.* The American Naturalist, 2004. **164**(3): p. 335-349.

32. Jaeger, B., *Package 'r2glmm'.* R Found Stat Comput Vienna available CRAN R-project org/package= R2glmm, 2017.

33. Schluter, D., *The ecology of adaptive radiation: Oxford University Press.* Oxfors, UK, 2000.

34. Losos, J.B., et al., *Contingency and determinism in replicated adaptive radiations of island lizards.* Science, 1998. **279**(5359): p. 2115-2118.

35. Van Belleghem, S.M., et al., *Selection and isolation define a heterogeneous divergence landscape between hybridizing Heliconius butterflies.* Evolution, 2021. **75**(9): p. 2251-2268.

36. Schluter, D., *Evidence for ecological speciation and its alternative.* Science, 2009. **323**(5915): p. 737-741.

37. Colosimo, P.F., et al., *Widespread parallel evolution in sticklebacks by repeated*
fixation of ectodysplasin alleles. Science, 2005. **307**(5717): p. 1928-1933.
38. Johnson, M.S., J. Murray, and B. Clarke, *Parallel evolution in Marquesan partulid land*
snails. Biological Journal of the Linnean Society, 2000. **69**(4): p. 577-598.
39. Projecto-Garcia, J., et al., *Repeated elevational transitions in hemoglobin function*
during the evolution of Andean hummingbirds. Proceedings of the National Academy
of Sciences of the United States of America, 2013. **110**(51): p. 20669-20674.
40. Cresko, W.A., et al., *Parallel genetic basis for repeated evolution of armor loss in*
Alaskan threespine stickleback populations. Proceedings of the National Academy of
Sciences of the United States of America, 2004. **101**(16): p. 6050-6055.
41. Gillespie, R.G., et al., *Repeated Diversification of Ecomorphs in Hawaiian Stick*
Spiders. Current Biology, 2018. **28**(6): p. 941-947.e3.
42. Beccaloni, G.W., *Vertical stratification of ithomiine butterfly (Nymphalidae:*
Ithomiinae) mimicry complexes: the relationship between adult flight height and
larval host-plant height. Biological journal of the Linnean Society, 1997. **62**(3): p. 313-
341.
43. DeVries, P.J., C.M. Penz, and R.I. Hill, *Vertical distribution, flight behaviour and*
evolution of wing morphology in Morpho butterflies. Journal of Animal Ecology,
2010. **79**(5): p. 1077-1085.
44. Benson, W.W., *Resource Partitioning in Passion Vine Butterflies.* Evolution, 1978.
32(3): p. 493-518.
45. Ramos, R.R., D. Rodrigues, and A.V. Freitas, *Oviposition preference and larval*
performance in a Heliconius erato phyllis (Lepidoptera: Nymphalidae) population
from southeastern Brazil: is there a positive relationship? Journal of Natural History,
2012. **46**(11-12): p. 669-681.
46. Rodrigues, D. and G.R. Moreira, *Seasonal variation in larval host plants and*
consequences for Heliconius erato (Lepidoptera: Nymphalidae) adult body size.
Austral Ecology, 2004. **29**(4): p. 437-445.
47. Kerpel, S.M. and G.R.P. Moreira, *Absence of Learning and Local Specialization on*
Host Plant Selection by Heliconius erato. Journal of Insect Behavior, 2005. **18**(3): p.
433-452.
48. Eilers, S., L.B. Pettersson, and E. Öckinger, *Micro-climate determines oviposition site*
selection and abundance in the butterfly Pyrgus armoricanus at its northern range
margin. Ecological Entomology, 2013. **38**(2): p. 183-192.
49. Turlure, C. and H. Van Dyck, *On the consequences of aggressive male mate-locating*
behaviour and micro-climate for female host plant use in the butterfly Lycaena
hippothoe. Behavioral Ecology and Sociobiology, 2009. **63**(11): p. 1581-1591.
50. Merckx, T., et al., *Thermal flight budget of a woodland butterfly in woodland versus*
agricultural landscapes: An experimental assessment. Basic and Applied Ecology,
2008. **9**(4): p. 433-442.
51. Schnebel, E.M. and J. Grossfield, *The influence of light on pupation height in*
Drosophila. Behavior genetics, 1986. **16**: p. 407-413.
427
428