Lassa virus NP DEDDh 3'-5' exoribonuclease activity is required for optimal viral RNA

replication.

Cheng Huang[1]*, Emily Mantlo[1,2,] Slobodan Paessler[1],

[1]Department of Pathology, Galveston National Laboratory and Institute for Human

Infections and Immunity, University of Texas Medical Branch, Galveston, TX, USA

[2] Current address: Department of Diagnostic Medicine/Pathobiology, College of

Veterinary Medicine, Kansas State University, Manhattan, Kansas, USA.

*Corresponding Author: Cheng Huang, chhuang@utmb.edu Tel: 409-266-6934

Abstract

Mammarenavirus include several pathogens that can cause severe and fatal zoonotic

diseases in humans. Lassa virus (LASV) is the causative agent for Lassa fever (LF)

currently endemic in West Africa. There is no approved vaccines and antivirals against

LASV infection. Despite the substantial threat of LASV to public health, important

questions in the basic biology of LASV still exist. LASV nucleoprotein has a DEDDH 3'-

to-5' exoribonuclease motif (ExoN), which is known to degrade virus-derived double-

stranded RNA and thereby is key to LASV evasion of interferon response. Intriguingly,

the NP ExoN motif is highly conserved in mammarenaviruses, regardless of

pathogenicity, suggesting that NP ExoN has important function in virus life cycle in

addition to immune evasion. By using reverse genetics system, we rescued an ExoN-

deficient LASV mutant (ExoN- rLASV). Compared to wild-type rLASV, the ExoN- rLASV

demonstrated impaired virus growth and decreased viral RNA production in interferon-

deficient cells. For the first time, we found that abrogation of LASV ExoN activity led to

increased aberrant viral RNA production in infection. Furthermore, NP ExoN deficiency

increased LASV sensitivity to mutagenic nucleoside analogue. Our study provided

evidence for an additional role of LASV NP ExoN in viral RNA replication besides its

recognized function in immune evasion. LASV ExoN activity may be required for optimal

viral RNA synthesis and control of aberrant viral RNA and viral RNA mutation.

Significance

LASV NP contains a DEDDh ExoN motif, through which LASV is capable of

degrading virus-derived, immunostimulatory dsRNA to eliminate the danger signal and

inhibit interferon response. Intriguingly, the NP ExoN motif is highly conserved among

arenavirus family, regardless of pathogenicity and viral capability to suppress host

innate immune response. Therefore, arenavirus NP ExoN may have yet-to-be-identified

function in virus infection. In this study, we discovered that abrogation of LASV NP

ExoN activity impaired viral RNA replication in interferon-deficient cells and enhanced

virus sensitivity to mutagenic nucleoside analogue. Our new findings indicate that LASV

NP ExoN participates in optimal viral RNA replication and open new directions for future

studies. Elucidating the function of LASV NP ExoN may promote our understanding of

the basics of LASV replication and inform antiviral and vaccine development.

Introduction

Arenaviridae consists of four genera, *Mammarenavirus*, *Reptarenavirus*,

Hartmanivirus, and *Antennavirus (1, 2)*. Mammarenviruses (referred to as arenavirus

hereafter) include several pathogens of major clinical importance. These arenaviruses

mainly infect their rodent hosts, often persistently without causing overt disease signs.

Spillover to humans could occur through aerosol inhalation leading to severe or fatal

zoonotic diseases (3-6). The Old World Lassa virus (LASV) causes Lassa fever (LF)

with sever morbidity and high case fatality and is endemic in West Africa (7). The recent

LF outbreaks in Nigeria featured high fatality rates of up to 25.4% (8, 9). LF is also

associated with a high incidence of sequelae: approximately one-third of LF survivors

suffer from sudden-onset sensorineural hearing loss, which constitutes a neglected

public health and social burden (10). The New World arenavirus Junín virus (JUNV) and

Machupo virus (MACV) sporadically cause Argentine hemorrhagic fever (AHF) and

Bolivian hemorrhagic fever (BHF) in South America (4, 11, 12). Vaccines and

treatments are limited for these highly pathogenic arenaviruses. Accordingly, LASV,

JUNV and MACV are classified as Category A Priority Pathogens in the USA. The

World Health Organization has listed LF in the Blueprint list of priority diseases for

which there is an urgent need for accelerated research and development (13).

Arenaviruses are negative-sense RNA virus with a single-stranded, bi-segmented

RNA genome. The large (L) segment genomic RNA (L gRNA) and the small (S)

segment genomic RNA (S gRNA) is around 7.2 kb and 3.4 kb, respectively, in length

(Fig 1A). Each genomic RNA contains two coding regions separated by a highly

structured, GC-rich intergenic region (IGR) (Fig 1A). The L segment encodes the RNA-

dependent RNA polymerase (RdRp) L protein and a small, zinc finger protein (Z), which

drives the assembly and budding of virus particles. The S segment encodes the viral

glycoprotein (GP), which mediates virus entry, and the nucleoprotein (NP), which is the

major structural component of the nucleocapsid (14). L protein and NP are both

required for viral RNA replication (15) in virus-induced, discrete cytosolic structures (16).

After virus entry, NP mRNA and L mRNA are first synthesized from S gRNA and L

gRNA, respectively. mRNA transcription is terminated at IGR, which is the bona fade

termination signal (Fig 1B). Arenavirus mRNA lacks a poly-A tail that is essential for the

stability of conventional mRNA in host cells. Histone and HCV mRNAs, which also lack

the poly-A tail, are stabilized by stem-loop structures at 3'-ends (17, 18). It is believed

that arenavirus mRNA could be stabilized by IGR element at the 3'-end in a similar

manner. Anti-genomic S RNA (S agRNA) and anti-genomic L RNA (L agRNA) are also

synthesized from S gRNA and L gRNA. Later in infection, GPC mRNA and Z mRNA are

transcribed from S agRNA and L agRNA, respectively (Fig 1B). Of note, IGR element is

also required for efficient incorporation of arenavirus genomic RNA into progeny virus

particles (19).

Arenavirus NP contains two structurally and functionally separated domains. The N-

terminal half (aa 1-340 in LASV NP) binds to viral genomic RNA to form vRNP (20). In

the C-terminal half of LASV NP, residues D389, E391, D466, D533 and H528 constitute

a DEDDh 3'-5' exoribonuclease (ExoN)-like motif. Structural and biochemical studies

have experimentally proved that LASV NP DEDDh motif harbors exoribonuclease

activity specificity for double-stranded RNA (dsRNA). Mutation of any of these critical

residues abolishes the ExoN activity (21-23). These studies have also revealed ExoN

activity is essential for LASV NP inhibition of Sendai virus-induced interferon response

(21, 23), which is in line with LASV suppression of innate immune response *in vitro* and

in vivo. RNA virus replication often generates dsRNA products, which can be sensed by

host Pattern Recognition Receptors (RIG-I, MDA5 and PKR etc.) and trigger signaling

cascades that lead to an anti-viral innate immune response mediated by IFN/cytokines.

Thus, the LASV NP ExoN activity to degrade virus-derived dsRNA is believed key to

LASV immune evasion and pathogenicity (24, 25).

Interestingly, the NP DEDDh motif and its 3'-5' ExoN activity are highly conserved in

arenaviruses regardless of pathogenicity. For instance, NPs of non-pathogenic

arenavirus Tacaribe virus (TCRV), Pichinde virus (PICV) and Mopeia virus (MOPV) also

have the ExoN activity (22, 26-28). In addition, arenaviral NP ExoN activity is not

correlated with viral capability to suppress IFN response. MOPV and TCRV infection

stimulate IFN and cytokine responses and therefore are used as non-pathogenic

arenavirus models in comparison with their closely related pathogenic counterparts, i.e.

LASV and JUNV, respectively (29, 30). This raises a question whether the highly

conserved NP ExoN activity has additional function besides immune evasion.

Furthermore, LASV ExoN- NP mutant protein demonstrated decreased capability to

support LASV minigenome replication (31, 32), implying NP ExoN participates in LASV

replication. Interestingly, positive-sense RNA virus coronavirus also possesses the

same DEDDh ExoN motif in nsp14, which is essential for the fidelity of coronavirus RNA

replication (33-35). Therefore, further studies are needed to investigate whether LASV

NP ExoN motif has additional role in virus life cycle.

In present study, we rescued an ExoN-deficient LASV mutant (ExoN- rLASV) using

reverse genetics system and found that loss of ExoN activity resulted in impaired virus

growth and viral RNA replication in interferon-deficient cells. We also provide the

evidence that abrogation of LASV NP ExoN increased aberrant viral RNA production in

infection. Furthermore, ExoN-deficient LASV was more sensitive to mutagenic

nucleoside analogue as compared with wild type LASV. Overall, our study indicates that

LASV NP ExoN has previously unrecognized function in viral RNA replication besides

its role in immune evasion. Our new finding may enhance our understanding of the

basic virology of this important human pathogen.

Results

Abrogation of NP ExoN activity impaired LASV multiplication in interferon-

deficient cells.

Biochemical studies have demonstrated that mutation of any of the NP D389, E391,

D466, D533 and H528 residues diminished LASV NP ExoN activity. In LASV

minigenome replication system, NP D389A and D389AG392A mutants exhibited an

impaired capability to support reporter gene expression (31, 32), suggesting NP ExoN

activity is required for optimal LASV replication. To investigate the role of LASV ExoN in

virus infection, we attempted to rescue recombinant LASV (rLASV, Josiah strain)

harboring NP D389A and NP D389AG392A mutations by using murine Pol I promoter

based LASV reverse genetics system. To minimize the likelihood of reverse mutation,

two nucleotide substitution were introduced to mutate NP D380 and NP G392 residue.

Specifically, for NP D380A mutant, nt $GA_{1266}C_{1267}$ in S agRNA was mutated to

$GC_{1266}T_{1267}$. For NP D389AG392A mutant, $GA_{1266}C_{1267}$ (D389) and $GG_{1275}A_{1276}$ (G392)

were mutated to $GC_{1266}T_{1267}$ (A389) and $GC_{1275}T_{1276}$ (A392), respectively. The rLASV

NPD389A mutant (ExoN- rLASV hereafter) was successfully rescued. Sequencing

analysis confirmed the introduced mutation is stable after 3 passages in Vero cells (Fig

2A). On the other hand, the rLASV NP D389AG392A mutant was not rescued despite

repeated attempts, indicating rLASV with the NP D389AG392A mutation is not viable.

Next, we assessed the multiplication of ExoN- rLASV in interferon-deficient Vero cells

(Fig 2B). Compared with wild-type (wt) rLASV, ExoN- rLASV demonstrated attenuated

growth in Vero cells. At 48 hours post-infection (hpi), the virus titer of ExoN- rLASV was

51-fold and 835-fold lower at MOI 0.01 and MOI 1, respectively, than wt rLASV.

Consistently, ExoN- rLASV also formed smaller plaques in Vero cells in comparison

with wt rLASV (Fig 2C). Thus, abrogation of ExoN activity impaired rLASV propagation

in Vero cells.

NP ExoN deficiency impaired LASV RNA replication in interferon-deficient cells.

As Vero cells are IFN-deficient (36, 37), the impaired multiplication of ExoN- rLASV in

Vero cells attests to a role of NP ExoN in LASV replication other than IFN evasion.

Given the role of coronavirus nsp14 DEDDh ExoN activity in coronavirus RNA

replication, we reasoned that LASV NP ExoN is also involved in viral RNA replication.

Vero cells were infected with wt rLASV and ExoN- rLASV at MOI 0.01 for 72 hr or at

MOI 1 for 24 hr. The virus titer of ExoN- rLASV was 51.4-fold and 6671-fold lower at

MOI 0.01 and MOI 1, respectively, than wt rLASV, (Fig. 3A). To assess viral RNA level,

total RNA was purified from infected cells and reverse transcribed to cDNA using

random hexamer primers. Then, a qPCR assay was performed with NP specific primers

to quantitate the viral RNA level at the NP locus, which measures the total RNA level of

SgRNA, S agRNA and S mRNA (Fig 3B). Similarly, a qPCR assay with primers specific

for the L locus was performed to measure the total RNA level of LgRNA, L agRNA and

L mRNA (Fig 3C). At MOI 0.01, the ExoN- rLASV RNA level was 45.5-fold lower

(P<0.01, student t test) at NP locus and 25.2-fold lower (P=0.04, student t test) at L

locus than wt rLASV. At MOI 1, the ExoN- rLASV RNA level was 1637-fold lower

(P<0.01, student t test) at NP locus and 122-fold lower (P<0.01) at L locus than wt

rLASV.

We further examined the level of S gRNA and L gRNA. Reverse transcription (RT)

was performed with primers targeting the 3'-end of S genomic RNA and L genomic RNA

using high fidelity reverse transcriptase SuperScript IV (Invitrogen), followed by qPCR

assays targeting NP locus on S gRNA and L locus on L gRNA as described above. At

MOI 0.01, the level of ExoN- rLASV S gRNA and L gRNA were 62.3-fold lower (P<0.01,

t test) and 16.7-fold lower (P<0.02), respectively, than wt rLASV. At MOI 1, the level of

ExoN- rLASV SgRNA and Lg RNA were 1181-fold (P<0.01, t test) lower and 387-fold

(P<0.02) lower as compared with wt rLASV (Fig 3D and 3E). Overall, our data

demonstrated that abrogation of ExoN led to substantially decreased LASV RNA

production, to an extent largely comparable with virus titer reduction in IFN-deficient

Vero cells.

Formation of aberrant viral RNA in ExoN- rLASV infection

Next, we performed RT-PCR to examine S gRNA and L gRNA in ExoN- rLASV

infection. Vero cells were infected with wt rLASV and ExoN- rLASV at MOI 0.01 for 72

hrs or at MOI 1 for 24 hrs. Intracellular RNA samples were purified and reverse

transcribed to cDNA using primers targeting the 3'-ends of S gRNA and L gRNA. Close-

to-full-length S gRNA (3391 nt) and L gRNA (7265 nt) was amplified with high fidelity

Platinum SuperFi DNA Polymerase (Invitrogen) to minimize PCR error. In agarose

electrophoresis, we observed an aberrant S RNA product (indicated as S' in Fig 4A)

migrating slightly faster than the standard S gRNA product (3391nt) in ExoN- rLASV

samples both at MOI 0.01 and MOI 1. The aberrant S' product was detected specifically

in ExoN- rLASV infection in repeated experiments. In L gRNA PCR, we also observed

several aberrant L gRNA products (approximately 3.5 kb to 6 kb) in ExoN- rLASV

samples both at MOI 0.01 and MOI 1 (Fig 4B). The result indicated that abrogation of

NP ExoN activity increased aberrant Sg RNA and Lg RNA formation in LASV infection.

Abrogation of LASV NP ExoN activity affected the integrity of Sg RNA

To determine the sequence of the aberrant S' RNA in ExoN- rLASV infection, we

performed agarose gel electrophoresis with ExoN- rLASV and wt rLASV PCR samples

(MOI 1) and purified the PCR amplicons around the full-size of S segment. Purified

amplicons were cloned into cloning vector (pSMART, Lucigen). Sanger sequencing of

10 colonies identified two types of aberrant S RNAs formed in ExoN- rLASV infection.

Three out of 10 colonies had a 37-nt deletion (S agRNA nt1824-1860), which is shown

as the region from site I to site I' in Fig 5A and 5B. The 37-nt deletion is located in the

67-nt-long IGR and therefore could disrupt the stem-loop structure of IGR. In addition,

five colonies contained a 231-nt deletion spanning the IGR and NP coding region (S

agRNA nt1616-1846, from site II to II' in Fig 5C). The 231-nt deletion consists of 42 nts

in IGR and 189 nts at the 3' end of NP gene (Fig 5C), which could disrupt IGR structure

and cause reading frame shift from NP residue 507. In comparison, only 2 out of 10

colonies from wt rLASV S samples had the 37-nt deletion in IGR. These data indicated

that LASV lacking NP ExoN activity is more prone to form structural deletion around S

RNA IGR.

PacBio Single Molecule, Real-Time (SMRT) long-read sequencing to characterize

aberrant S gRNA formed in ExoN- rLASV infection.

To systematically assess the spectrum of aberrant S gRNA formed in ExoN- rLASV

infection, we utilize PacBio Single Molecule, Real-Time (SMRT) long-read sequencing

technology (38) to characterize LASV SgRNA at single molecule level. In PacBio SMRT

long-read sequencing, a DNA polymerase continues to read single circularized-cDNA-

template for multiple rounds and provides deep sequencing data for each molecule. It

can determine the circular consensus sequence (CCS) of full-length cDNA up to 10kb

long at single molecule level with up to 99.99% accuracy. PacBio SMRT long-read

sequencing is ideal for analyzing RNA quasi-species, long amplicons, and RNA

structural variations. It has been successfully used in research works on RNA virus, for

example, studies on defective interference RNA in influenza virus infection and hepatitis

C virus variants post drug treatment (39, 40).

We gel purified close-to-full-length S RNA PCR amplicon from wt- and ExoN- rLASV-

infected cells (MOI 1) and performed PacBio SMRT long-read sequencing (Sequel II). A

total of 402,410 CCS reads and 381,134 CCS reads was obtained for wt rLASV and

ExoN- rLASV samples, respectively (Fig 6A). The mean number of read passes for

each cDNA is 26 times for wt rLASV with an average length of 2319 nt and 28 times for

ExoN- rLASV with an average length of 2319 nt (Fig 6A). The mean read score of the

assay is >0.999. CCS reads with size of 3100-3391 bp were selected using Filter

FASTQ and aligned with LASV reference sequence using *minimap2*. RNA variants with

>3% frequency was called using *ivar*. In ExoN- rLASV sample, 88.9% and 4.9% of S

RNA had the 231-nt and the 37-nt deletion (Fig 6B and 6C). In comparison, 7.1% and

18.2% of the S RNA had the 231-nt and the 37-nt deletion, respectively, in wt rLASV

sample. The data also confirmed that greater than 99.93% of ExoN- rLASV harbored

the intended NP D389A mutation. In addition, an A-to-G substitution at nt 393 (in

antigenomic sense) was identified in S RNA in ExoN- rLASV sample at a frequency of

39.14%. A C-to-U substitution at nt 1518 in S RNA was found in wt rLASV sample with

a frequency of 3.07%. Overall, the PacBio SMRT sequencing data confirmed that

structural variations occurred at a substantially higher frequency in ExoN- rLASV S

gRNA than wt rLASV S gRNA.

ExoN- rLASV was more sensitive to nucleoside analogue treatment.

Coronavirus nsp14 DEDDh 3'-5' ExoN activity proofreads newly synthesized viral

RNA by removing mis-incorporated nucleotide. Abrogation of nsp14 ExoN activity

rendered coronavirus susceptible to lethal mutagenesis when treated with nucleoside

analogues such as 5-Fluorouracil (5-FU) (41). For prototype arenavirus LCMV, 5-FU

also could be incorporated into nascent RNA in viral RNA synthesis and introduces

mutations affecting LCMV fitness (42). It is possible that loss of ExoN activity could

increase arenavirus sensitivity to mutagenic nucleoside analogues. Thus, we evaluated

the sensitivity of wt rLASV and ExoN- rLASV to 5-FU in Vero cells (MOI 0.1). We first

verified that 5-FU treatment alone did not affect cell viability at the concentrations

administrated in the experiment (Fig. 7A, CellTiter-Glo Viability Assay, Promega). The

result indicated that abrogation of NP ExoN activity substantially increased LASV

sensitivity to 5-FU treatment starting from 100 µM (Fig 7B). At 400 µM, the ExoN-

rLASV titer deceased by 1053-fold compared with the titer in mock-treated cells, which

was 13.5-time greater than the 78-fold decrease for wt rLASV (Fig 7B).

EIDD-1931 (*N4*-hydroxycytidine or NHC) is a cytidine analogue that increases the

mutation frequency of a broad range RNA viruses (43). We further assessed the

sensitivity of ExoN- rLASV and wt rLASV to EIDD-1931. Treatment of Vero cells with

EIDD-1931 (5 µM -100 µM) alone did not substantially affect cell viability (Fig 7A,

CellTiter-Glo Viability Assay, Promega). At 100 µM EIDD-1931, the titer of ExoN- rLASV

was decreased by 47.6-fold, greater than the 6-fold reduction for wt rLASV (Fig 7C).

Collectively, our data indicated that abrogation of NP ExoN activity increased LASV

sensitivity to mutagenic nucleoside analogues.

Discussion

LASV NP DEDDh motif has been known essential for LASV evasion of host innate

immune and viral pathogenicity attributable to its ExoN activity to degrade dsRNA

generated in virus RNA replication. Interestingly, the DEDDh ExoN motif is highly

conserved in arenavirus family, regardless of viral pathogenicity, implying the ExoN

activity has yet-to-be-identified function(s) in arenavirus life cycle. Curiously, positive-

sense RNA virus coronavirus contains the same DEDDh ExoN motif in nsp14, which

has proofreading activity in viral RNA replication. In this study, we found that abrogation

of NP ExoN attenuated LASV growth in IFN-deficient Vero cells. Furthermore, ExoN-

rLASV demonstrated impaired viral RNA replication, including decreased S gRNA and L

gRNA production, in IFN-deficient cells. For the first time, we found that ExoN- rLASV

was prone to produce aberrant viral RNAs. Sanger sequencing and SMRT long-read

sequencing of S RNA revealed that ExoN- rLASV is more likely to form structural

deletions in S RNA, suggesting an impact of NP ExoN deficiency on viral genomic RNA

integrity. In addition, we also found an increased sensitivity of ExoN- rLASV to

mutagenic nucleoside analogue, suggesting NP ExoN has a role in reducing LASV RNA

mutation. Taken together, our study provide evidence for previously unrecognized

function(s) of LASV NP ExoN in optimal LASV RNA replication and control of aberrant

viral RNA production and viral RNA mutation.

The data that NP ExoN deficiency affected the propagation and the plaque size of

LASV in IFN-deficient Vero cells (Fig 2) clearly indicates an important role of NP ExoN

in LASV fitness. This is also highlighted by our data that the NP D389AG392A rLASV

mutant (Josiah strain) was not rescuable. Other studies have also documented that NP

ExoN deficiency leads to attenuated growth of rPICV, rMOPV and rLASV (AV strain) in

Vero cells (26, 27, 31). In addition, reversion to wt virus have been reported for ExoN-

rPICV and ExoN- rLASV (AV strain) in IFN-deficient cells. PICV possessing NP D380A

mutation (equivalent to the NP D389A mutation for LASV) forms smaller plaques. After

two passages of the PICV NP ExoN- mutant in Vero cell, wt revertant PICV, which

forms large plaques, could be readily identified (27). For LASV AV strain, an NP D389A

mutant was found partially reverted to wt sequence after two passages in Vero cells

(31). Overall, ours and others' data indicated NP ExoN activity is required for optimal

arenavirus replication.

There seems to be some differences in terms of the genetic stability of LASV NP

ExoN- mutants in our study with the LASV Josiah strain and the study with LASV AV

strain (31). For LASV AV strain, an NP D389AG392A mutant was rescued and

genetically stable, while an NP D389A mutant partially reverted to wt sequence. In

comparison, our data with the Josiah strain indicated the LASV NP D389AG392A

mutant could not be rescued and the NP D389A mutation is stable at least after 3

passages in Vero cells. The reason for these discrepancies is unclear, probably due to

virus strain difference. It is also possible that the two-nucleotide substitution in NP

D389A mutant in present study reduced the likelihood of reversion to wt sequence.

Nevertheless, both studies generally support for the involvement of NP ExoN in optimal

LASV replication and fitness.

Our data further showed that abrogation of NP ExoN activity impaired LASV RNA

replication, including S gRNA and L gRNA accumulation, in IFN-deficient cells (Fig 3).

These findings are in line with previous studies, in which NP D389A mutant protein and

NP D389AG392A mutant protein exhibited an impaired capability to support reporter

gene expression in LASV minigenome replication systems (31, 32). Future studies are

necessary to determine whether NP ExoN deficiency affects LASV genomic RNA

replication, mRNA transcription, or both. The RNA replication of RNA virus is error

prone. Nucleotide mis-incorporation could cause viral RdRp stop or pause RNA

elongation. As MOPV NP and LCMV NP are able to excise mismatched nucleotide at

the 3'-end of dsRNA substrate (28), it is possible that NP ExoN is capable of removing

mis-incorporated nucleotide in viral RNA synthesis and facilitating viral RNA synthesis.

In this scenario, NP ExoN deficiency may affect the yield or the rate of viral RNA

synthesis, which may partly explain the impaired viral RNA replication in ExoN- rLASV

infection. Other than LASV, NP ExoN activity is also required for optimal reporter gene

expression in PICV minigenome replication (27). Hence, it is interesting to study

whether NP ExoN is required for optimal viral RNA replication for other arenaviruses.

An important finding of the present study is that abrogation of NP ExoN increased

aberrant LASV RNA formation in LASV, which links NP ExoN activity to the integrity of

viral genomic RNA. For the first time, we detected increased aberrant S gRNA and Lg

RNA production in ExoN- rLASV-infected cells in repeated experiments (Fig 4). Sanger

sequencing of close-to-full-length S gRNA amplicon revealed two type of deletion

mutants in ExoN- rLASV samples. Three of 10 colonies had a 37-nt deletion (nt1824-

1860, S agRNA) located in the 67-nt-long IGR (site I to site I' shown in Fig 5A and 5B).

Five colonies contained a 231-nt deletion (nt1616-1846 in S agRNA, from II to II' shown

in Fig 5C) in the IGR and NP coding region. The 37-nt deletion could disrupt the stem-

loop structure in IGR. The 231-nt deletion could also disrupt IGR structure and cause

reading frame shift from NP residue 507. In comparison, only 2 out of 10 colonies from

wt rLASV S samples had the 37-nt deletion in IGR. The biological implication of these

deletions remains to be investigated in further studies. Of note, arenavirus IGR is the

termination signal of viral mRNA transcription and also an viral RNA element for efficient

viral genomic RNA packaging into progeny virus particles. Hence, abrogation of NP

ExoN activity may indirectly affect proper viral mRNA synthesis as well as the efficacy of

viral RNA packaging into virions.

To better understand the spectrum of LASV SgRNA variants, we further utilized

PacBio SMRT long-read sequencing to characterize LASV SgRNA amplicon at single

molecule level. Short-read Illumina NGS has low sensitivity or high false positive rates

in solving complex Structural Variations (SVs) with deletions and insertions at least 50

nt in size (44). PacBio SMRT long-read sequencing can determine the circular

consensus sequence of full-length cDNA up to 10kb long at single molecule level, which

is ideal for analyzing RNA quasi-species, long amplicons, and SVs. In addition, SMRT

long-read technology is ideal for sequencing through highly repetitive, GC-rich

sequences that are presented in the IGR of arenavirus RNA. To ensure the accuracy of

the SMRT assay, raw reads with a minimum number of passes greater than 3 and

higher than 99% accuracy were used to generate the CCS reads. The mean number of

read passes for each cDNA was 26 times for wt rLASV with an average length of 2319

nt and 28 times for ExoN- rLASV with an average length of 2319 nt (Fig 6A). The mean

reading score of the SMRT assay was >99.9%. Variants of >3% frequency was selected

in data analysis to ensure the reliability of the reads. The result indicated that 88.9%

and 4.9% of the S genomic RNA contained the 231-nt deletion and the 37-nt deletion in

ExoN- rLASV sample. In comparison, only 7.1% and 18.2% of the S RNA in wt rLASV

sample has the 231-nt deletion and the 37-nt deletion. The data suggests NP ExoN

helps to reduce structural deletion in viral genomic RNA. In addition, we found that

higher than 99.93% of ExoN- rLASV S gRNA harbored the intended NP D389A

mutation, which confirmed that the NP D389A mutation was maintained in ExoN-

rLASV.

The molecular basis for these structural deletions formed in ExoN- rLASV S RNA is

unclear. Interestingly, sequence analysis revealed the presence of homologous

sequences flanking the 37-nt deletion and the 231-nt deletion in S RNA (Fig 8, boxed

sequences, sites I/I' for the 37-nt deletion, sites II/II' for the 231-nt deletion).

Accordingly, we propose a stop-and-realign model for the mechanism of increased S

RNA structural deletions in ExoN- rLASV infection (Fig 8). MOPV NP ExoN and LCMV

NP ExoN excise mismatched nucleotide at the 3'-end of dsRNA substrate (28). When

viral RNA synthesis error occurs, LASV NP ExoN may also excise mismatched

nucleotide like MOPV and LCMV counterparts and facilitate RdRp L protein resume

RNA elongation. Without NP ExoN activity, viral RNA synthesis may stop as shown in

Fig 8A for the 37nt-deletion and Fig 8B for the 231nt-deletion (Step 1, mis-incorporation

and stop). Depending on the nucleotide mis-incorporated, LASV RdRp L protein may

realign the 3'-end of nascent RNA (at site I in Fig 8A and site II in Fig 8B) with

downstream homologous sequences on the template RNA (sites I' and II' in Fig 8A and

Fig 8B, respectively) and resume RNA elongation (Step 2, realign and resume)).

Consequently, the regions between these homologous sequences are skipped by RdRp

L protein, leading to the 37-nt deletion (Fig 8A) and the 231-nt deletion (Fig 8B) in S

RNA. Thus, LASV NP ExoN may act to reduce these structural deletions around IGR by

enhancing the fidelity of LASV RNA synthesis. In this regard, it is worth noting that RNA

viruses with low-fidelity RdRp, such as Sindbis virus or tombusvirus, exhibited an

increased production of defective viral genomic RNA correlated with an enhanced rate

of viral RNA recombination (45-47). Further studies are warranted to investigate the

molecular base of increased aberrant RNA formation associated with LASV NP ExoN

deficiency.

Our nucleoside analogue data support for a role of LASV NP ExoN in reducing RNA

mutation. The RNA replication of RNA virus is generally error prone, which increases

the diversity of virus population that could be advantageous in evolution. However, the

mutation rate should also be controlled as even a moderate 1.1-2.8-fold increase in

mutation rate may drastically affect virus fitness (48). How arenavirus controls mutation

is largely unknown. In LCMV infection, 5-FU could be incorporated into nascent viral

RNA and cause mutations that severely affect LCMV fitness. We found that abrogation

of NP ExoN activity increased LASV sensitivity to 5-FU as well as another mutagenic

nucleoside analogue EIDD-1931 (Fig 7). These data suggest LASV NP ExoN activity

contributes to reduce viral RNA mutation and enhance the fidelity of viral RNA

replication.

In summary, the present study provided evidence that LASV NP ExoN is

multifunctional, which is required for optimal viral RNA replication in addition to its

recognized role in immune evasion. Abrogation of LASV NP ExoN activity leads to

decreased viral RNA replication, increased aberrant viral RNA formation, and enhanced

sensitivity to mutagenic nucleotide analogue. These findings provide novel and

important insights into the role of NP ExoN motif in LASV life cycle. Future studies are

warranted to investigate whether the NP ExoN of other arenaviruses also play roles in

viral RNA replication. Hemorrhagic fever-causing arenaviruses continue to pose a threat

to public health with pandemic potential. A better understanding of the function of NP

ExoN might inform vaccine and antiviral development against these important human

pathogens.

Methods

Cells and Viruses.

Vero cells (CCL-81, ATCC) were maintained in Dulbecco's modified eagle medium

(Hyclone) supplemented with 10% FBS (Invitrogen) and 1% penicillin and streptomycin

solution (Hyclone). The LASV (Josiah strain) used in studies was recombinant virus that

were rescued using reverse genetic systems in BHK21 cells and passaged only once

(P1) in Vero cells as previously described (49). All plasmids used for LASV rescue were

sequencing confirmed. Site-directed mutagenesis on p-mPol I-LASV Sag plasmid was

performed using QuikChange II kit (Agilent) according to the instruction of manufacturer.

To construct the rLASV NPD380A mutant, forward primer 5'-CCAAATGCTAAGACCTG

GATGGCTATTGAAGGAAGACCTGAAGATC-OH, and reverse primer 5'-GATCTTCAG

GTCTTCCTTCAATAGCCATCCAGGT CTTAGCATTTGG-OH were used. To construct

the rLASV NP D389AG392A mutant, forward primer 5'-AAATGCTAAGACCTGGATGG

CTATTGAAGCTAGACCTGAAGATCCAGTGG-OH, and reverse primer 5'-CCACTGGA

TCTTCAGGTCTAGCTTCAATAGCCATCCAGGTCTTAGCATTT-OH were used. The

sequence of mutant rLASV was confirmed by Sanger Sequencing after RT-PCR

amplification of viral RNA extracted from infected cells. All infection work with

pathogenic arenaviruses was performed at the BSL4 facilities in Galveston National

Laboratory in the University of Texas Medical Branch in accordance with institutional

health and safety guidelines and federal regulations.

RNA Extraction, RT-PCR and real-time RT-qPCR

RNA lysates were prepared using the TRIzol reagent (Life Technology). RNA was purified using the RNeasy Minikit (Qiagen) and treated with DNase I (Qiagen) as previously reported (32, 50). Equal amount of total RNA (0.5 to 1μg) was reverse transcribed to cDNA at 55 $^{\circ}$C for 60 minutes using high fidelity reverse transcriptase SuperScript IV (Invitrogen) per manufacturer's protocol. The high-fidelity reverse transcriptase SuperScript IV has low RNase H activity and is suitable for long cDNA synthesis. Random hexamer primers and sequence specific primers were used in RT as specified in each experiment. cDNA samples were treated with RNase H. In RT of S gRNA, an S gRNA 3'UTR specific primer (5'-CGCACAGTGGATCCTAGGCTA-OH) was used. In RT of L gRNA, an L gRNA 3'UTR specific primer (5'- CACCGAGGATCCTAGG CATTAAGGCTATC-OH) was used in cDNA synthesis. PCR amplification of close to full-length Sg cDNA (3391bp) was conducted with Platinum SuperFi DNA Polymerase (Invitrogen) using forward primer 5'-ATCCTAGGCATTTTT GGTTGC-OH and reverse primer 5'-CGCACAGTGGATCCTAGGCTA-OH. PCR amplification of close to full-length Lg cDNA (7265 bp) was conducted with Platinum SuperFi DNA Polymerase using forward primer 5'-CACCGAGGATCCTAGG CATTAAGGCTATC-OH and reverse primer 5'-ATCCTAGGCAATTTGGTTGTTCTTTTTTGAG-OH. Real-time quantitative PCR (qPCR) was performed with SsoAdvanced Universal SYBR green Supermix (BioRad) on a CFX96 real-time PCR detection system (BioRad) as described previously (32, 50). To measure the viral RNA level at NP locus and L locus, cDNA synthesized with random primers was used in qPCR assay. LASV NP forward primer (5′-GAAGGGCCTG GGAAAACACT-OH) and LASV NP reverse primer (5′-AGGTAAGCCCAGCGGTAAAC-

OH) were used for NP locus. LASV L forward primer (5′-CAGCAGGTCAGACGAAGTG

T-OH) and LASV L reverse primer (5′-GTTGTGCATAGGGGAGGCTT-OH) were used

for L locus. To measure LASV Sg RNA and Lg RNA level with qPCR, cDNA

synthesized with SgRNA- and LgRNA-specific primers as indicated above was used in

qPCR assay. The qPCR data was analyzed with CFX Manager software (BioRad). The

RNA level of each target gene was normalized to that of housekeeping gene β-actin

with validated primers from BioRad. All experiments were performed separately in

triplicates.

PacBio SMRT long-read sequencing of LASV SgRNA

Vero cells was infected by ExoN- rLASV and wt rLASV at MOI 1. At 48 hpi, total RNA

was purified from infected cells. RT-PCR amplification of the close-to-full-length SgRNA

was performed as described above. PCR amplicons (approximately 3.4 kb) were gel

purified with Monarch DNA gel extraction kit (NEB). Library construction, PacBio SMRT

long-read sequencing and CCS determination was performed by Azenta Life Sciences

(USA). At least 1μg amplicon was ligated to barcoded adapters (Pacific Biosciences) for

SMRTbell library construction and then sequenced on the PacBio Sequel IIe platform

with v3.0 chemistry. Raw reads with a minimum number of passes greater than 3 were

used to generate the CCS reads using PacBio SMRTLINK v.10.1. HiFi reads (>=99%

accuracy) were extracted. Data analysis was performed utilizing *Galaxy*, an open

source, web-based platform supported by NIH, NSF and the Texas Advanced

Computing Center. CCS reads were filtered for those with size 3100-3394bp and high-

quality reads (Phred>20) using Filter FASTQ and mapped to the LASV reference seq

(MH358389) with *minimap2* (PacBio HiFi reads vs reference mapping (-k19 -w19 -

U50,500 -g10k -A1 -B4 -O6,26 -E2,1 -s200)). Variants with cutoff >3% frequency were

called with *ivar* and visualized using the Integrated Genomics Viewer (IGV).

Mutagenic nucleotide analogue treatment

5'-fluorouracil (Cat# 03738, Sigma) and N4-hydroxycytidine (EIDD-1931) (Cat#9002958,

Cayman) were dissolved in DMSO (cell culture grade, Sigma) and further diluted in

DMEM media contain 2% FBS and 0.1% DMSO. Vero cells were mock-treated or

treated with 5-FU or EIDD-1931 at different concentrations as indicated in each

experiment for 4 hrs and then infected with wt rLASV and ExoN- rLASV (MOI 0.1) for 1

hr. Virus inoculums were removed and replaced with fresh media (DMEM + 2% FBS

and 0.1% DMSO) containing 5-FU or EIDD-1931 of different concentrations as

indicated. At 48 hpi, the supernatants of infected cells were harvested and subjected to

plaque assay to determine virus titers. The viability of Vero cells treated with 5-FU and

EIDD-1931 alone were assessed using CellTiter-Glo Viability Assay (Promega).

Acknowledgments

This work was supported in part by Public Health Service grants 1R21AI166985 (to CH)

RO1AI093445 (to SP) and RO1AI129198 (to SP). CH was also supported by Institute

for Human Infections & Immunity, UTMB Pilot Study Fund (P86035) and would like to

acknowledge Galveston National Laboratory (supported by the Public Health Service

award 5UC7AI094660) for support of research activity. Research work in Dr. Paessler's

lab was also supported by the John. S. Dunn Distinguished Chair in Biodefense

endowment. E.K.M was supported by National Institutes of Health T32 training grant

AI060549.

Figure legends

Figure 1. (A). Schematic diagram of arenavirus S and L genomic RNAs. On each

genomic RNA, an Intergenic Region (IGR) separates two ORFs. The highly structured

IGRs are the termination signals of viral mRNA transcription and required for efficient

packaging of viral genomic RNA into progeny viruses. **(B).** Genomic RNA replication

and mRNA transcription of arenavirus S RNA. After virus entry, L protein and NP

transcribe NP mRNA from S gRNA. S antigenomic RNA (S agRNA) is also synthesized

from S gRNA, which serves as the template for GPC mRNA and S gRNA synthesis later

in virus infection. Arenavirus mRNA lacks a poly-A tail and contains the structured IGR

at 3'-end.

Figure 2. Abrogation of NP ExoN activity impaired LASV multiplication in

interferon-deficient cells.

A recombinant LASV (rLASV) with NP D389A mutation was rescued (ExoN- rLASV).

(A). The D389A mutation was maintained after 3 passages in Vero cells (Sanger

sequencing). (B). The growth curve of wild-type rLASV (wt) and the ExoN- rLASV (NP

ExoN-) in Vero cells at multiplicity of infection (MOI) 0.01 and MOI 1. (C) The plaque

morphology of wt rLASV (wt) and ExoN- rLASV.

Figure 3. Abrogation of NP ExoN activity affected LASV RNA replication.

Vero cells were infected with wt rLASV and ExoN- rLASV (MOI=0.01 and 1.0). At 72 hpi

(MOI 0.01) or 24 hpi (MOI 1.0), virus titer was determined by plaque assay **(A)**. RT-

qPCR assay for viral RNA level at the NP locus **(B)** and L locus **(C)**. The viral RNA level

for S gRNA **(D)** and L gRNA **(E)** were also determined by qRT-PCR. The viral RNA level

was normalized to the level of host β-actin mRNA and are presented as the fold

changes relative to the level of ExoN- rLASV samples (set as 1.0). The data presents

the mean and SEM of three independent experiments. (*: $P<0.01$; **: $P<0.02$; ***:

$P<0.05$ with Student *t*-test).

Figure 4. Aberrant viral RNA formation in ExoN- rLASV infection

Vero cells were infected with wt rLASV (wt) and ExoN- rLASV (ExoN-) at MOI 0.01 for

72 hr and at MOI 1 for 24 hr. RNA samples were purified from infected cells and

transcribed to cDNA with primers targeting the 3'-end of S gRNA and L gRNA,

respectively. High-fidelity PCR was performed to amplify close-to-full-length S gRNA

(3391 nt) and L gRNA (7265 nt) followed by agarose gel electrophoresis. (**A**). RT-PCR

result of S gRNA (S). S' is a slightly fast-migrating aberrant S RNA product in ExoN-

rLASV samples. (**B**). RT- PCR result of L gRNA. Several aberrant L gRNA products in

ExoN- rLASV samples (3.5 kb to 6 kb) are observed. The aberrant RNA products are

indicated by asterisk (*). Plasmid: PCR product using a plasmid containing LASV L

segment as template.

Figure 5. Sanger sequencing of aberrant S RNA formed in ExoN- rLASV infection.

(A). Predicted RNA structure of IGR of S RNA by *mFold.* **(B).** In ExoN- rLASV, a 37-nt

deletion (nt1824-1860 in S agRNA) from site I to site I' disrupted IGR structure. **(C)** a

231-nt deletion (nt1616-1846 in S agRNA) from C-terminus of NP at site II' to site II in

IGR (refer to A) was identified in S gRNA sample of ExoN- rLASV.

Figure 6. PacBio SMRT long-read sequencing of S gRNA in rLASV samples.

Vero cells were infected with wt rLASV and ExoN- rLASV (MOI 1.0). RT-PCR product of

the close-to-full-length SgRNA (3391 nt) were purified after agarose gel electrophoresis

and subjected to PacBio SMRT long-read sequencing (Sequel IIe) with v3.0 chemistry.

(A). HiFi CCS reads were generated using PacBio SMRTLINK v.10.1. **(B).** CCS reads

with size 3100-3391bp were selected with Filter FASTQ and mapped to LASV reference

seq (GenBank MH358389) with *minimap2.* Variants were called with *ivar* and visualized

using the Integrated Genomics Viewer. **(C)**. The percentage of the 231-nt deletion and

the 37-nt deletion in S RNA of wt rLASV and ExoN- rLASV sample.

Figure 7. NP ExoN deficiency enhanced LASV sensitivity to mutagenic nucleotide

analogues.

(A). Vero cells were treated with 5'-FU and EIDD-1931 at indicated concentrations. At

48 hr post treatment, cell viability was measured using CellTiter-Glo Viability Assay

(Promega). Data shown are the average (n=4) and SEM. **(B) and (C).** Vero cells were

treated with 5-FU and EIDD-1931 at different concentrations as indicated. Cells were

infected with wt rLASV (wt) and ExoN- rLASV (ExoN-) at MOI 0.1. At 48 hpi, virus titers

were determined by plaque assay. Log10 virus titer changes relative to the virus titer of

non-treated cells are shown. Data presented are the mean and the SEM of three

independent experiments.

Figure 8. A stop-and-realign model for the 37-nt deletion **(A)** and the 231-nt deletion **(B)**

in S RNA of ExoN- rLASV. Homologous sequences flanking the deletion sites are

indicated as boxed sequences (sites I/I' for the 37-nt deletion, sites II/II' for the 231-nt

deletion). Sites I and II are located in IGR stem-loop structure. In viral RNA synthesis,

NP ExoN may remove the mis-incorporated nucleotide and helps LASV RdRp L protein

to resume viral RNA elongation. In the absence of NP ExoN activity, viral RNA

elongation may terminate (Step 1). Depending on the nucleotide mis-incorporated (i.e.,

an A instead of C as shown in A and B), viral RdRp L protein may realign the 3'-end of

nascent RNA (site I or II) with downstream homologous sequences on the template

RNA (Sites I' and II') and resume RNA elongation (Step 2). Thus, the 37-nt deletion and

the 237-nt deletion in S RNA may be derived from viral RdRp skipping the regions

between the homologous sequences, an error could be controlled by functional NP

ExoN.

<u>References</u>

584
1.   Maes P, Adkins S, Alkhovsky SV, Avsic-Zupanc T, Ballinger MJ, Bente DA, Beer

M, Bergeron E, Blair CD, Briese T, Buchmeier MJ, Burt FJ, Calisher CH, Charrel

RN, Choi IR, Clegg JCS, de la Torre JC, de Lamballerie X, DeRisi JL, Digiaro M,

Drebot M, Ebihara H, Elbeaino T, Ergunay K, Fulhorst CF, Garrison AR, Gao GF,

Gonzalez JJ, Groschup MH, Gunther S, Haenni AL, Hall RA, Hewson R, Hughes

HR, Jain RK, Jonson MG, Junglen S, Klempa B, Klingstrom J, Kormelink R,

Lambert AJ, Langevin SA, Lukashevich IS, Marklewitz M, Martelli GP, Mielke-

Ehret N, Mirazimi A, Muhlbach HP, Naidu R, Nunes MRT, et al. 2019. Taxonomy

of the order Bunyavirales: second update 2018. Arch Virol 164:927-941.

2.   Shi M, Lin XD, Chen X, Tian JH, Chen LJ, Li K, Wang W, Eden JS, Shen JJ, Liu

L, Holmes EC, Zhang YZ. 2018. The evolutionary history of vertebrate RNA

viruses. Nature 556:197-202.

3.   Geisbert TW, Jahrling PB. 2004. Exotic emerging viral diseases: progress and

challenges. Nat Med 10:S110-121.

4.   Grant A, Seregin A, Huang C, Kolokoltsova O, Brasier A, Peters C, Paessler S.

2012. Junin virus pathogenesis and virus replication. Viruses 4:2317-2339.

5.   Gunther S, Lenz O. 2004. Lassa Virus, vol 41, p 339-390, Critical Reviews in

Clinical Laboratory Science.

6.   Lukashevich IS. 2012. Advanced vaccine candidates for Lassa fever. Viruses

4:2514-2557.

7.   McCormick JB, Webb PA, Krebs JW, Johnson KM, Smith ES. 1987. A

prospective study of the epidemiology and ecology of Lassa fever. J Infect Dis

155:437-444.

8.   Dan-Nwafor CC, Furuse Y, Ilori EA, Ipadeola O, Akabike KO, Ahumibe A,

Ukponu W, Bakare L, Okwor TJ, Joseph G, Mba NG, Akano A, Olayinka AT,

Okoli I, Okea RA, Makava F, Ugbogulu N, Oladele S, Namara G, Muwanguzi EN,

Naidoo D, Mutbam SK, Okudo I, Woldetsadik SF, Lasuba CL, Ihekweazu C.

2019. Measures to control protracted large Lassa fever outbreak in Nigeria, 1

January to 28 April 2019. Euro Surveill 24.

9.   Sattler RA, Paessler S, Ly H, Huang C. 2020. Animal Models of Lassa Fever.

Pathogens 9.

10.   Mateer EJ, Huang C, Shehu NY, Paessler S. 2018. Lassa fever-induced

sensorineural hearing loss: A neglected public health and social burden. PLoS

Negl Trop Dis 12:e0006187.

11.   Gomez RM, Jaquenod de Giusti C, Sanchez Vallduvi MM, Frik J, Ferrer MF,

Schattner M. 2011. Junin virus. A XXI century update. Microbes Infect 13:303-

311.

12.   Patterson M, Grant A, Paessler S. 2014. Epidemiology and pathogenesis of

623   Bolivian hemorrhagic fever. Curr Opin Virol 5:82-90.

624   13.   World Health Organization WH. 2018. 2018 Annual review of diseases

625   prioritized under the Research and Development Blueprint.

626   http://www.who.int/blueprint/priority-diseases/en/

627   14.   Buchmeier MJ, de la Torre JC, Peters CJ. 2007. Arenaviridae: the viruses and

628   their replication, 5th ed, vol 2. Wolter Kluwer Lippincott Williams & Wilkins,

629   Philadelphia, PA, USA.

630   15.   Pinschewer DD, Perez M, de la Torre JC. 2003. Role of the virus nucleoprotein in

631   the regulation of lymphocytic choriomeningitis virus transcription and RNA

632   replication. J Virol 77:3882-3887.

633   16.   Baird NL, York J, Nunberg JH. 2012. Arenavirus infection induces discrete

634   cytosolic structures for RNA replication. J Virol 86:11301-11310.

635   17.   Dickson AM, Wilusz J. 2011. Strategies for viral RNA stability: live long and

636   prosper. Trends Genet 27:286-293.

637   18.   Ross J. 1995. mRNA stability in mammalian cells. Microbiol Rev 59:423-450.

638   19.   Pinschewer DD, Perez M, de la Torre JC. 2005. Dual role of the lymphocytic

639   choriomeningitis virus intergenic region in transcription termination and virus

640   propagation. J Virol 79:4519-4526.

641   20.   Hastie KM, Liu T, Li S, King LB, Ngo N, Zandonatti MA, Woods VL, Jr., de la

642   Torre JC, Saphire EO. 2011. Crystal structure of the Lassa virus nucleoprotein-

643   RNA complex reveals a gating mechanism for RNA binding. Proc Natl Acad Sci

644   U S A 108:19365-19370.

645 21. Hastie KM, Kimberlin CR, Zandonatti MA, MacRae IJ, Saphire EO. 2011.

646   Structure of the Lassa virus nucleoprotein reveals a dsRNA-specific 3' to 5'

647   exonuclease activity essential for immune suppression. Proc Natl Acad Sci U S A

648   108:2396-2401.

649 22. Jiang X, Huang Q, Wang W, Dong H, Ly H, Liang Y, Dong C. 2013. Structures of

650   arenaviral nucleoproteins with triphosphate dsRNA reveal a unique mechanism

651   of immune suppression. J Biol Chem 288:16949-16959.

652 23. Qi X, Lan S, Wang W, Schelde LM, Dong H, Wallat GD, Ly H, Liang Y, Dong C.

653   2010. Cap binding and immune evasion revealed by Lassa nucleoprotein

654   structure. Nature 468:779-783.

655 24. Mantlo E, Paessler S, Huang C. 2019. Differential Immune Responses to

656   Hemorrhagic Fever-Causing Arenaviruses. Vaccines (Basel) 7.

657 25. Meyer B, Ly H. 2016. Inhibition of Innate Immune Responses Is Key to

658   Pathogenesis by Arenaviruses. J Virol 90:3810-3818.

659 26. Carnec X, Mateo M, Page A, Reynard S, Hortion J, Picard C, Yekwa E, Barrot L,

660   Barron S, Vallve A, Raoul H, Carbonnelle C, Ferron F, Baize S. 2018. A Vaccine

661   Platform against Arenaviruses Based on a Recombinant Hyperattenuated

662   Mopeia Virus Expressing Heterologous Glycoproteins. J Virol 92:J Virol

663   92:e0223017. https://doi.org/10.1128/JVI.0202230-0223017.

664 27. Huang Q, Shao J, Lan S, Zhou Y, Xing J, Dong C, Liang Y, Ly H. 2015. In vitro

665   and in vivo characterizations of pichinde viral nucleoprotein exoribonuclease

666   functions. J Virol 89:6595-6607.

28.   Yekwa E, Aphibanthammakit C, Carnec X, Coutard B, Picard C, Canard B, Baize

S, Ferron F. 2019. Arenaviridae exoribonuclease presents genomic RNA edition

capacity. bioRxiv doi:10.1101/541698:541698.

29.   Groseth A, Hoenen T, Weber M, Wolff S, Herwig A, Kaufmann A, Becker S.

2011. Tacaribe virus but not junin virus infection induces cytokine release from

primary human monocytes and macrophages. PLoS Negl Trop Dis 5:e1137.

30.   Pannetier D, Faure C, Georges-Courbot MC, Deubel V, Baize S. 2004. Human

macrophages, but not dendritic cells, are activated and produce alpha/beta

interferons in response to Mopeia virus infection. J Virol 78:10516-10524.

31.   Carnec X, Baize S, Reynard S, Diancourt L, Caro V, Tordo N, Bouloy M. 2011.

Lassa virus nucleoprotein mutants generated by reverse genetics induce a

robust type I interferon response in human dendritic cells and macrophages. J

Virol 85:12093-12097.

32.   Mateer EJ, Maruyama J, Card GE, Paessler S, Huang C. 2020. Lassa Virus, but

Not Highly Pathogenic New World Arenaviruses, Restricts Immunostimulatory

Double-Stranded RNA Accumulation during Infection. J Virol 94:e02006-02019.

https://doi.org/10.1128/JVI.02006-02019.

33.   Denison MR, Graham RL, Donaldson EF, Eckerle LD, Baric RS. 2011.

Coronaviruses: an RNA proofreading machine regulates replication fidelity and

diversity. RNA Biol 8:270-279.

34.   Minskaia E, Hertzig T, Gorbalenya AE, Campanacci V, Cambillau C, Canard B,

Ziebuhr J. 2006. Discovery of an RNA virus 3'->5' exoribonuclease that is

critically involved in coronavirus RNA synthesis. Proc Natl Acad Sci U S A

103:5108-5113.

35.   Ogando NS, Ferron F, Decroly E, Canard B, Posthuma CC, Snijder EJ. 2019.

The Curious Case of the Nidovirus Exoribonuclease: Its Role in RNA Synthesis

and Replication Fidelity. Front Microbiol 10:1813.

36.   Osada N, Kohara A, Yamaji T, Hirayama N, Kasai F, Sekizuka T, Kuroda M,

Hanada K. 2014. The genome landscape of the african green monkey kidney-

derived vero cell line. DNA Res 21:673-683.

37.   Desmyter J, Melnick JL, Rawls WE. 1968. Defectiveness of interferon production

and of rubella virus interference in a line of African green monkey kidney cells

(Vero). J Virol 2:955-961.

38.   Rhoads A, Au KF. 2015. PacBio Sequencing and Its Applications. Genomics

Proteomics Bioinformatics 13:278-289.

39.   Lui WY, Yuen CK, Li C, Wong WM, Lui PY, Lin CH, Chan KH, Zhao H, Chen H,

To KKW, Zhang AJX, Yuen KY, Kok KH. 2019. SMRT sequencing revealed the

diversity and characteristics of defective interfering RNAs in influenza A (H7N9)

virus infection. Emerg Microbes Infect 8:662-674.

40.   Yamashita T, Takeda H, Takai A, Arasawa S, Nakamura F, Mashimo Y, Hozan

M, Ohtsuru S, Seno H, Ueda Y, Sekine A. 2020. Single-molecular real-time deep

sequencing reveals the dynamics of multi-drug resistant haplotypes and

structural variations in the hepatitis C virus genome. Sci Rep 10:2651.

41. Smith EC, Blanc H, Surdel MC, Vignuzzi M, Denison MR. 2013. Coronaviruses

lacking exoribonuclease activity are susceptible to lethal mutagenesis: evidence

for proofreading and potential therapeutics. PLoS Pathog 9:e1003565.

42. Ruiz-Jarabo CM, Ly C, Domingo E, de la Torre JC. 2003. Lethal mutagenesis of

the prototypic arenavirus lymphocytic choriomeningitis virus (LCMV). Virology

308:37-47.

43. Agostini ML, Pruijssers AJ, Chappell JD, Gribble J, Lu X, Andres EL, Bluemling

GR, Lockwood MA, Sheahan TP, Sims AC, Natchus MG, Saindane M,

Kolykhalov AA, Painter GR, Baric RS, Denison MR. 2019. Small-Molecule

Antiviral beta-d-N(4)-Hydroxycytidine Inhibits a Proofreading-Intact Coronavirus

with a High Genetic Barrier to Resistance. J Virol 93:e01348-01319.

https://doi.org/10.1128/JVI.01348-01319.

44. Sedlazeck FJ, Rescheneder P, Smolka M, Fang H, Nattestad M, von Haeseler A,

Schatz MC. 2018. Accurate detection of complex structural variations using

single-molecule sequencing. Nat Methods 15:461-468.

45. Panaviene Z, Nagy PD. 2003. Mutations in the RNA-binding domains of

tombusvirus replicase proteins affect RNA recombination in vivo. Virology

317:359-372.

46. Poirier EZ, Mounce BC, Rozen-Gagnon K, Hooikaas PJ, Stapleford KA,

Moratorio G, Vignuzzi M. 2015. Low-Fidelity Polymerases of Alphaviruses

Recombine at Higher Rates To Overproduce Defective Interfering Particles. J

Virol 90:2446-2454.

47.   Vignuzzi M, Lopez CB. 2019. Defective viral genomes are key drivers of the

virus-host interaction. Nat Microbiol 4:1075-1087.

48.   Holland JJ, Domingo E, de la Torre JC, Steinhauer DA. 1990. Mutation

frequencies at defined single codon sites in vesicular stomatitis virus and

poliovirus can be increased only slightly by chemical mutagenesis. J Virol

64:3960-3962.

49.   Yun NE, Seregin AV, Walker DH, Popov VL, Walker AG, Smith JN, Miller M, de

la Torre JC, Smith JK, Borisevich V, Fair JN, Wauquier N, Grant DS, Bockarie B,

Bente D, Paessler S. 2013. Mice lacking functional STAT1 are highly susceptible

to lethal infection with Lassa virus. J Virol 87:10908-10911.

50.   Huang C, Kolokoltsova OA, Mateer EJ, Koma T, Paessler S. 2017. Highly

Pathogenic New World Arenavirus Infection Activates the Pattern Recognition

Receptor Protein Kinase R without Attenuating Virus Replication in Human Cells.

J Virol 91:e01090-01017. https://doi.org/10.1128/JVI.01090-01017.

A.

```
CCCAAATGCTAAGACCTGGATGGACATTGAAGGAAGACCTGAAGATC
|||||||||||||||||||||||||||||||||||||||||||||||
GGGTTTACGATTCTGGACCTACCTGTAACTTCCTTCTGGACTTCTAG
            385         390         395
    P   N   A   K   T   W   M   D   I   E   G   R   P   E   D
                                    N →
```

```
CCCAAATGCTAAGACCTGGATGGACATTGAAGGAAGACCTGAAGATC
|||||||||||||||||||||||||||||||||||||||||||||||
CCCAAATGCTAAGACCTGGATGGCTAATTGAAGGAAGACCTGAAGATC
```

B.

C.

A

I

II

I'

IGR structure (S ag)

B

S agRNA

NP

I

I'

GPC

5'   3'

37-nt deletion

C

S agRNA

NP

II'

II

GPC

5'   3'

231-nt deletion

A.

Sample ID	CCS Reads	Number of CCS Bases	CCS Read Score (mean)	Number of Passes (mean)
Wt rLASV	402,410	933,554,242	0.999	28
ExoN- rLASV	381,134	994,863,239	0.999	26

B.

C.

Sample	231-nt del	37-nt del
Wt rLASV	7.1%	18.2%
ExoN- rLASV	88.9%	4.9%

A. The 37-nt deletion in LASV NP ExoN-

(**1**) Mis-incorporation/stop

```
5′-UAAAUGGACGCCC CCGUGAC A*
     ||||||||||||| |||||||
3′-AUUUACCUGCGGG GGCACUG g.........gggacug UCCCAGAGUA
                  Site I                  Site I′
```

(**2**) Re-align/resume

```
                    5′-UAAAUGGACGCCC CCGUGAC A------→
                                     |||||||  |
3′-AUUUACCUGCGGG GGCACUG g.........ccccc gggacug UUCCCAGAGUA
```

B. The 231-nt deletion in LASV NP ExoN-

(**1**) Mis-incorporation/stop

```
              NP        Site II′              Site II          IGR
SagRNA 5′-AAAGACUUAU GCCAC aug.........g GUCAC GGGGGCCCUGACAAGGGUC
                                         ||||| |||||||||||||||||||||
                                   *A CAGUG CCCCCGGGACUGUUCCCAG-5′
```

(**2**) Re-align/resume

```
         5′-AAAGACUUAU GCCAC aug.........g GUCAC GGGGGCCCUGACAAGGGUC
                       |||||
             ←----- A CAGUG CCCCCGGGACUGUUCCCAG-5′
```