Endocardial HDAC3 is required for myocardial trabeculation

Jihyun Jang[1,2], Mette Bentsen[3], Ye Jun Kim[4], Erick Kim[4], Vidu Garg[1,2], Chen-Leng Cai[5],

Mario Looso[3], Deqiang Li[1, 2]

[1]Center for Cardiovascular Research, Abigail Wexner Research Institute, Nationwide

Children's Hospital, Columbus, OH 43215, USA

[2]Department of Pediatrics, The Ohio State University College of Medicine, Columbus, OH

43215, USA

[3]Bioinformatics Core Unit (BCU), Max Planck Institute for Heart and Lung Research,

61231, Bad Nauheim, Germany

[4]Department of Surgery, University of Maryland School of Medicine, Baltimore, MD 21201,

USA

[5]Department of Pediatrics, Herman Wells Center for Pediatric Research, Indiana

University School of Medicine, Indianapolis, IN 46201, USA

Address correspondence to:

Deqiang Li, MD&PhD

575 Children's Drive, Room WB4329,

Columbus, OH 43215, USA

Phone: 614-355-5826

Email: Deqiang.Li@nationwidechildrens.org

Abstract

Failure of proper ventricular trabeculation is often associated with congenital heart disease (CHD). Support from endocardial cells, including the secretion of extracellular matrix (ECM) and growth factors is critical for trabeculation. However, it is poorly understood how the secretion of ECM and growth factors is initiated and regulated by endocardial cells.

We found that genetic knockout (KO) of histone deacetylase 3 (*Hdac3*) in the endocardium in mice resulted in early embryo lethality and ventricular hypotrabeculation. Single cell RNA sequencing identified significant downregulation of ECM components in *Hdac3* KO endocardial cells. Secretome from cultured *Hdac3* KO mouse cardiac endothelial cells lacked transforming growth factor ß3 (TGFß3) and showed significantly reduced capacity in stimulating cultured cardiomyocyte proliferation, which was remarkably rescued by TGFß3 supplementation. Mechanistically, we identified that HDAC3 induced *Tgfß3* expression through repressing microRNA (miR)-129-5p.

Our findings provide novel insights into the pathogenesis of CHD and conceptual strategies to promote myocardial regeneration.

Introduction

Congenital heart disease (CHD) is the most common congenital disorder, and the leading cause of mortality from birth defects[1]. Defective myocardial growth is either the primary manifestation or associated with many forms of CHD[2, 3]. In the early primitive heart (mouse embryonic day 8-10), inner cardiac endothelial cells (CECs), also called endocardial cells, and outer cardiomyocytes (CMs), which are separated by extracellular matrix (ECM), co-contribute to early myocardial trabeculation (i.e., formation of myocardial meshwork extending to cardiac chamber)[4, 5, 6]. CECs juxtapose myocardial trabeculae and lead myocardial trabeculation by secretion of growth factors such as Neuregulin 1 (NRG1) and transforming growth factor ß (TGFß)[2, 3, 5]. Meanwhile, CECs support trabeculation by secreting ECM (e.g., various collagens, versican) that facilitate the travelling of CEC secreted growth factors to trabecular myocardium and support trabecular CM proliferation[4, 7]. Surprisingly, disruption of ECM components alone (e.g., hyaluronan synthase-2, versican) prevents the proper formation of trabeculae in mice[8, 9]. Although CECs play such important roles during ventricular trabeculation, it remains poorly understood how these ECM proteins and growth factors are regulated in the developing CECs.

TGFß signaling pathway is involved in many fundamental cellular processes in both adults and developing embryos including cell proliferation, differentiation, migration, and apoptosis[10, 11]. The TGFß superfamily includes many different ligands (e.g., Bone morphogenetic proteins, Growth and differentiation factors, Activin, Nodal, TGFß). These ligands bind to their specific type II receptors, which triggers phosphorylation of type I receptor, eventually leads to phosphor-Smad2/3 localized to the nucleus and drives a

variety of downstream target gene expression, which is the canonical TGFß signaling

pathway[12]. In addition, the binding of ligands to receptors can also directly activate

multiple non-Smad mediated intracellular signaling pathways such as MAP kinase

pathways and phosphatidylinositol-3-kinase/AKT pathways[13, 14]. These non-Smad

(noncanonical) TGFß signaling pathways are equally critical for driving some specific

cellular processes (e.g., cell proliferation, cell survival, endothelial/epithelial

mesenchymal transition)[13, 14]. During cardiovascular development, unsurprisingly, TGFß

signaling is implicated in most morphological events including cardiac lineage

specification, cardiac looping, valvogenesis, and cardiomyogenesis[15]. The TGFß family

includes TGFß1, TGFß2, and TGFß3. Although TGFß1, TGFß2 and TGFß3 are

abundantly expressed in the developing endocardium and epicardium, it appears that

they do not perform redundant functions: individual knockout results in different

phenotypes[15]. For instance, global knockout of either *Tgfß2* or *Tgfß3* result in early

perinatal lethality, presenting hypoplastic hearts with thin myocardium[16, 17], suggesting

that they play critical roles for myocardial growth. In contrast, global knockout of *Tgfß1*

lead to partial embryonic lethality with survivors appearing phenotypically normal in first

the two weeks of life[18]. Nonetheless, it is poorly understood how the expression of TGFß

ligands are regulated in the early developing heart.

Epigenetic modulators such as histone deacetylases (HDACs) are critical for early heart

development as key regulators of regulating gene expresison[19, 20]. There are 18 known

human HDACs, which can be subdivided into four classes based on their function and

DNA sequence similarity. Class I HDACs include HDAC1, HDAC2, HDAC3, and HDAC8,

and are predominantly nuclear and ubiquitously expressed[21]. Although class I HDACs

have high degree of homology, global knockout of each causes early embryonic lethality[19],

suggesting distinct roles during early heart development. Global deletion of *Hdac3* leads

to early embryonic lethality at gastrulation[22]. In the developing heart, lack of *Hdac3* in

cardiac progenitors disrupts secondary heart field development[23], lymphovenous valve

morphogenesis[24], and CM lineage specification[25]. Recently, we reported that HDAC3 in

the developing epicardium plays a critical role in supporting compact myocardial wall

expansion[26]. It is unclear whether or how HDAC3 in the developing endocardium impacts

myocardial trabeculation.

In the current study, we investigated the role of HDAC3 in the developing endocardium

during early heart development.

Results

Endothelial/Endocardial cell specific deletion of *Hdac3* results in embryonic lethality and hypotrabeculation.

To study the role of HDAC3 in the cardiac endothelium during early heart development, *Hdac3* was specifically deleted in the endothelial cells using *Tie2-Cre*[27], while its expression in other cells including CMs remains intact (Supplemental Figure 1). Note: since *Tie2-Cre* is transiently expressed in oocytes[28], the mating scheme was always *Hdac3*$^{f/+}$; *Tie2-Cre/+* (♂) and *Hdac3*$^{f/f}$ (♀). By recovering the progenies at a series of embryonic and postnatal stages, we identified that *Hdac3* KO in the developing endothelium resulted in early embryonic lethality starting at E12.5 (Supplemental Table 1). To investigate the potential cause of early embryonic lethality, we performed histological analyses. Interestingly, in Tie2 endothelial *Hdac3* KO embryos (*Hdac3*^{tko}), there was no noticeable phenotype in the various endothelial cell derived tissue where the *Tie2*-lineage contributes, such as endocardium (Figure 1, Supplemental Figure 2) and blood vessels (Supplemental Figure 3), given that *Tie2-Cre* is active in all endothelial cells in the body[29]. In contrast, there were myocardial defects in *Hdac3*^{tko} embryos starting at E11.5: *Hdac3*^{tko} trabeculae appeared to be significantly sparser and shorter as compared to the thick and long webbing like trabecular network in littermate control hearts (Supplemental Figure 2). This cardiac phenotype in *Hdac3*^{tko} mice was more prominent at E12.5 (Figure 1A). The compact layer was also significantly thinner in *Hdac3*^{tko} hearts. To further validate the myocardial trabecular phenotype in endothelial *Hdac3* deleted hearts, we specifically deleted *Hdac3* in the endocardium using *Nfatc1*$^{Cre/+}$, endocardial specific Cre[30], which also resulted in embryonic lethality (Supplemental Table 2).

Hdac3^{nko} (*Hdac3$^{f/f}$; Nfatc1$^{Cre/+}$*) hearts presented a similar cardiac phenotype, albeit at a

later stage (E14.5): smaller and sparser trabeculae, thinner compact wall (Supplemental

Figure 4). This is consistent with the achievement of late deletion of *Hdac3* in the

endocardium by *Nfatc1$^{Cre/+}$* (Supplemental Figure 4). Nonetheless, these results confirm

that the cardiac phenotypes resulting from *Tie2* endothelial *Hdac3* deletion are primarily

from the endocardium, and not secondary to another endothelium (e.g., vascular).

Hdac3^{tko} embryos were used for all subsequent mechanistic studies.

Endothelial *Hdac3* knockout hearts display decreased trabecular CM proliferation.

Next, we investigated whether altered cell proliferation and/or apoptosis contribute to the

hypotrabeculation phenotypes in *Hdac3^{tko}* hearts. At E11.5, the percentage of either p-

H3+ or BrdU+ CMs is significantly lower in *Hdac3^{tko}* trabecular and compact myocardium

as compared to littermate control hearts (Figure 1B). This finding is consistent with the

hypoplastic cardiac phenotypes seen in *Hdac3^{tko}* hearts (Figure 1A). In contrast, there

was no significant difference in cell apoptosis between *Hdac3^{tko}* hearts and littermate

control hearts (Supplemental Figure 5). Decreased CM proliferation of *Hdac3^{tko}* hearts

may result from disrupted communication from the endocardium to the trabecular

myocardium. We performed single cell RNA-sequencing (scRNA-seq) on E11.5 *Hdac3^{tko}*

hearts and littermate control hearts (Figure 2A). We obtained high-quality scRNA-seq

profiles for 23,468 and 25,547 cells from 4 *Hdac3^{tko}* (*Hdac3$^{f/f}$; Tie2-Cre/+; R26$^{eYFP/+}$*) and

4 control (3 of *Hdac3$^{f/+}$; Tie2-Cre/+; R26$^{eYFP/+}$* and 1 of *Hdac3$^{f/+}$; Tie2-Cre/+*) heart single

cell suspensions, respectively. Unsupervised clustering successfully separated 6 distinct

clusters including endocardial cells and CMs in these samples (Figure 2B and

Supplemental Figure 6). Referring to a recent transcriptome profiling methodology for

clustering embryonic CMs[31], we re-clustered CMs into "trabecular" and "compact" clusters

based on the overall scores after applying a few known trabecular and compact markers

(Trabecular: *Bmp10*, *Nppa* and *Thbs4*; Compact: *Hey2*, *Mycn* and *Tnnt1*) (Supplemental

Figure 7A and Figure 2C). Among them, *Bmp10+* CMs nicely represent the trabecular

cluster (Figure 2C), and thus we focused on this population for subsequent analyses since

Hdac3^{tko} hearts mainly display hypotrabeculation phenotypes (Figure 1). Overall, the total

number of *Bmp10+* trabecular CMs were not significantly different between *Hdac3^{tko}* and

littermate control hearts (Supplemental Figure 7B). Note: 10,000 cells/sample was

sequenced. Next, we performed differential gene expression (DGE) analyses on *Hdac3^{tko}*

and littermate control trabecular clusters. We found that the expression of eukaryotic

translation elongation factor 1 alpha 2 (*Eef1a2*) and early growth response 1 (*Egr1*) are

significantly down regulated in *Hdac3^{tko}* trabeculae as compared to control trabeculae

(Figure 2D). *Eef1a2* and *Egr1* are known to be critical for cell growth[32, 33, 34]. These

findings are consistent with reduced proliferation index seen in *Hdac3^{tko}* hearts (Figure

1B), suggesting myocardial growth retardation account for the hypotrabeculation

phenotypes in *Hdac3^{tko}* hearts.

Endothelial HDAC3 induces ECM gene expression.

Among all scRNA-seq samples, 7 of them carried *R26$^{eYFP/+}$* reporters (3 CTL and 4

Hdac3^{tko}). This enabled us to perform a lineage specific analysis. In the single cell

suspension, 8.6% cells in CTL hearts and 11.2% cells in *Hdac3^{tko}* hearts were YFP+

(Supplemental Figure 8A). As expected, the expression of *Hdac3* in YFP+ populations in

Hdac3^{tko} hearts was significantly decreased compared to YFP+ populations in CTL hearts

(Supplemental Figure 8B). During early heart development, endothelial progenitors can

also give rise to mesenchymal cells through endothelial to mesenchymal transition[6, 35].

As expected, YFP+ cells (*Tie2* lineage) were primarily CECs with a minority of

mesenchymal cells (Figure 2E). Also, as expected, *Hdac3* expression was absent in

Hdac3^{tko} YFP+ cells. Subsequent DGE analysis did not reveal significant gene expression

differences between *Hdac3^{tko}* and control mesenchymal cells (Figure 2E). Instead, in

CEC clusters, we found that the expression of genes regulating cellular component

organization was highly enriched in the control YFP+ CEC cluster (Figure 2F). In

particular, the expression of ECM associated genes, including *Col13a1, Col1a2, Tgfbi,*

Timp2, Col26a1, Col3a1, Col4a5, *Timp3*, and *P4ha3*, was significantly reduced in

Hdac3^{tko} YFP+ CEC cluster as compared to control YFP+ CEC cluster (Figure 2G).

Col1a1, Col1a2, Col2a1, and *Col3a1* and *P4ha3* DGE was further validated in *Hdac3^{tko}*

and control heart lysates by qRT-PCR (Figure 2H). These results suggest that cardiac

endothelial HDAC3 induces the expression of ECM associated genes, which are critical

for CM proliferation and trabeculation growth (Figure 2I).

183 Cardiac endothelial HDAC3 induces *Tgfβ3* in driving CM proliferation.

Trabeculation and ECM remodeling are coordinated during early heart development, and

both processes are regulated by growth signaling[36]. To better identify potential growth

signaling pathways regulated by HDAC3 in the endocardium, we used a mouse cardiac

endothelial cell (MCEC) line[37]. Using CRISPR-Cas9 gene editing technology and the

same set of gRNAs we have recently used[26], we successfully knocked out *Hdac3* in the

MCECs (Figure 3A). *Hdac3* KO MCECs grew slower than *Hdac3* empty vector (EV)

control MCECs. To determine whether the secretome from *Hdac3* KO endothelial cells is

competent to induce CM proliferation, we treated cultured primary embryonic CMs

isolated from E13.5 *Tnnt2$^{nGFP/+38}$* hearts with supernatant from either *Hdac3* KO MCECs

or *Hdac3* EV MCECs, respectively. *Hdac3* KO supernatant treatment resulted in

significant decrease of the percentage of Ki67+ CMs and the total number of CMs, as

compared to *Hdac3* EV supernatant treatment (Figure 3B). These results suggest that

growth signals derived from CECs required to support CM proliferation may be

inadequate/deficient in *Hdac3* KO MCECs, and the incompetent secretome from *Hdac3*

KO endothelial cells may contribute to the hypotrabeculation phenotype. Indeed,

treatment of wildtype MCEC supernatants but not control media (DMEM/F-12 without any

serum or growth factor supplementation) successfully rescued the hypotrabeculation

phenotype in *ex vivo* cultured E11.5 *Hdac3^{tko}* hearts (Supplemental Figure 11). To

unbiasedly search for such growth signals, we performed bulk RNA-Seq analyses on

Hdac3 KO and EV MCECs; we identified 484 downregulated and 239 upregulated genes.

Gene ontology pathway analyses revealed that a number of endocardium-initiated

signaling pathways were significantly downregulated in *Hdac3* KO MCECs (Supplemental

Figure 9A). Through DGE analysis, we found that growth factors including *Igf1, Tgfß3,*

Tgfß2, Tgfß1, Igf1, Cxcl12, Nrg1 and Vegfß were significantly downregulated in *Hdac3*

KO MCECs (Supplemental Figure 9B). We attempted to validate these results at the

protein level by western blot. Surprisingly, we found that only TGFβ3 protein expression

was significantly decreased in *Hdac3* KO MCECs as compared to *Hdac3* EV MCECs

(Supplemental Figure 9C), and thus we chose TGFβ3 to pursue further. Interestingly,

TGFβ3 was also significantly decreased when the deacetylase activity of HDAC3 was

inhibited by RGFP966 (a HDAC3 selective inhibitor) treatment (Figure 3C and

Supplemental Figure 9D). To assess *Tgfβ3* expression in E11.5 CECs, we sorted Tie2-

derived population from $Hdac3^{tko}$ ($Hdac3^{f/f}$; *Tie2-Cre/+*; $R26^{eYFP/+}$) and CTL ($Hdac3^{f/+}$;

Tie2-Cre/+; $R26^{eYFP/+}$) hearts by performing fluorescence activated cell sorting on YFP

signal (Supplemental Figure 10). We found that *Tgfβ3* mRNA level was also significantly

downregulated in E11.5 *Hdac3* KO CECs compared to littermate CTL CECs (Figure 3D).

TGFβ3 downregulation in E11.5 $Hdac3^{tko}$ hearts was further validated at the protein level

by western blot (Figure 3E). Next, we determined whether TGFβ3 deficiency in *Hdac3*

KO MCECs contributed to the CM proliferation deficit following *Hdac3* KO supernatant

treatment. First, we quantified TGFβ3 and found that it was significantly decreased in the

supernatants from *Hdac3* KO MCECs compared to that from *Hdac3* EV MCECs (Figure

3F). To determine whether TGFβ3 is an essential secretory component of MCEC

supernatants in stimulating CM proliferation, we generated *Tgfβ3* KO MCECs using

CRISPR-Cas9 gene editing technology (Supplemental Figure 12A). We found that *Tgfβ3*

KO significantly decreased the capacity of MCEC supernatants to induce CM proliferation

(percentage of Ki67+GFP+/GFP+ CMs) compared to CTL MCEC supernatant

(Supplemental Figure 12B). Further, we found that *Tgfβ3* KO significantly compromised

the capacity of MCEC supernatants in supporting myocardial trabeculation and

compaction when treating *ex vivo* cultured E11.5 CTL hearts with *Tgfβ3* KO MCEC

supernatants as compared to control MCEC supernatants (Supplemental Figure 12C).

These results altogether suggest that endocardial TGFβ3 is critical for myocardial growth.

Next, we evaluated CM proliferation after we supplemented *Hdac3* KO supernatants with

recombinant mouse TGFβ3 protein. Strikingly, TGFβ3 supplementation successfully

rescued CM proliferation defects by *Hdac3* KO supernatant treatment (Figure 3G).

Further, we found that TGFβ3 supplementation also rescued myocardial growth deficit in

ex vivo cultured E11.5 *Hdac3*[tko] hearts (Supplemental Figure 13). These results suggest

that cardiac endothelial HDAC3 supports myocardial growth by inducing *Tgfβ3*

expression.

Cardiac endothelial HDAC3 induces *Tgfβ3* expression through repressing

microRNA (miR)-129-5p.

Recently, we reported that HDAC3 regulates growth factors by modulating the miR

pathway in the developing epicardium[26]. To identify the potential HDAC3 downstream

miR targets in CECs, we performed miR sequencing in *Hdac3* KO and EV MCECs.

Through DGE analyses, we found that 5 miRs were significantly upregulated in *Hdac3*

KO MCECs (Figure 4A) (cutoff threshold: Log2 fold change greater than 1.5 and adjusted

P<0.01). To identify which miR(s) may regulate *Tgfβ3* expression, we treated MCECs

with mimics of each of the 5 miRs and found that treatment with miR-129-5p mimics

significantly inhibited the expression of *Tgfβ3* (Figure 4B). This finding was validated by

western blot (Figure 4C). We further validated the upregulation of miR-129-5p in both

Hdac3 KO MCECs and E11.5 *Hdac3*[tko] hearts (Figure 4D and 4E). Interestingly, miR-

129-5p was similarly significantly upregulated when the deacetylase activity of HDAC3 is

inhibited by RGFP966 treatment (Figure 4D), suggesting that the inhibition of miR-129-

5p expression by HDAC3 is deacetylase dependent. To determine whether miR-129-5p

upregulation in MCECs has potential impact on myocardial growth, we treated cultured

Tnnt2[nGFP/+38] CMs or cultured whole E11.5 wildtype hearts with supernatants from

MCECs transfected with either miR-129-5p mimics or scramble RNAs. We found that

miR-129-5p supernatant treatment significantly decreased its capacity to induce CM

proliferation and myocardial growth (Supplemental Figure 14). Lastly, to determine

whether HDAC3 induces *Tgfβ3* expression through repressing miR-129-5p, we knocked

down miR-129-5p by miRZip lentivirus in *Hdac3* KO MCECs. Remarkably, miR-129-5p

knockdown significantly restored the TGFβ3 expression in *Hdac3* KO MCECs (Figure 4G).

Further, the supernatants from miR-129-5p knockdown *Hdac3* KO MCECs significantly

restored the CM proliferation promoting capacity from *Hdac3* KO MCEC supernatants

(Figure 4H). These results suggest that HDAC3 promotes the expression of TGFβ3

through repressing miR-129 in supporting trabecular myocardial growth (Figure 4I).

Discussion

Myocardial growth/expansion, one of the most critical cardiac developmental events, is accompanied by two separate but closely associated processes: compaction and trabeculation. Many forms of congenital heart disease are associated with abnormal cardiac trabeculation and compaction[2, 3]. For proper myocardial development, besides CMs, the surrounding non-CMs including endocardial cells can influence myocardial growth by sending growth signals to the neighboring myocardia[39], as well as secreting ECM proteins that serve as signal transmission media[2, 40]

Epigenetic regulation can define a tissue-specific and developmental stage-specific transcription network during early heart development[41, 42], mainly achieved by posttranslational modifications of histones, around which DNA winds. There are several different types of histone modifications (e.g., methylation/demethylation, acetylation/deacetylation, phosphorylation/dephosphorylation, ubiquitination/ deubiquitination). By alternating between any of these states, the associated genes can be dynamically poised to be transcriptionally active/repressed. Histone deacetylases (HDACs) are family of enzymes that remove acetyl groups from a lysine residue on a histone, allowing the histones to wrap the DNA more tightly to repress gene expression. Class I HDACs, includes HDAC3, are widely expressed, but play specific roles during early embryogenesis and organogenesis[19]. Mesodermal or global KO of *Hdac3* results in myogenic defects and early embryonic lethality[22, 43]. Interestingly, specific deletion of *Hdac3* in the myocardium does not generate cardiac morphogenic phenotypes during early heart development, but rather compromises cardiac function at later postnatal stages[43, 44]. These early findings suggest that the function of HDAC3 in nonmyocyte

compartments may be critical for early cardiac morphogenesis and myocardial

development. Our recent work demonstrated that epicardial HDAC3 is critical for driving

myocardial compact wall expansion[26]. Our current study suggests that endocardial

HDAC3 is important for myocardial trabeculation: endothelial/endocardial deletion of

Hdac3 resulted in hypotrabeculation (Figure 1 and Supplemental Figure 4). We note that

Tie2-Cre turns on around E8.0-8.5 in all endothelial cells[29], whereas $Nfatc1^{Cre/+}$ is active

around E9.5 only in endocardial cells[30] and only achieves efficient *Hdac3* deletion by

E14.5 likely due to inefficient IRES-mediated Cre transcription[45] (Supplemental Figure 4).

Nonetheless, these two deletion models generated similar cardiac phenotypes of

hypotrabeculation, although we did observe differences in the severity of the myocardial

phenotypes and time of lethality, which we attribute to the different temporal and spatial

activities and/or efficiencies of these two Cres. We also observed that both E12.5

$Hdac3^{tko}$ and E14.5 $Hdac3^{nko}$ hearts exhibited thin compact wall phenotype. This is likely

attributable to *Hdac3* deficiency in the subepicardial endothelial population which form

coronary arteries invading and supporting compact wall[46]. This late thin compact wall

phenotype is also consistent with the similar phenotype resulted from epicardial deletion

of $Hdac3^{26}$. An earlier report of conditional deletion of *Hdac3* in endothelial cells using a

different *Tie2-Cre* driver primarily targeting hemogenic endothelial cells[27] resulted in mid-

gestational lethality (E14.5) presenting predominantly a lymphovenous and lymphatic

valve malformation phenotype[47]. However, it is unknown what phenotypes were present

in the myocardium in that study as none were reported. Nonetheless, we deemed it

inapplicable to evaluate potential lymphatic valve phenotypes in our pan-endothelial

Hdac3 KO embryos since these mice died around E12.5 (Supplemental Table 1), while

lymphovenous and lymphatic valve formation starts at E13.5[48]. However, it is interesting

to note that the role of HDAC1 and HDAC2 in the developing endocardium appears to be

dispensable, as deletion of either *Hdac1* or *Hdac2* by *Tie2-Cre* do not elicit apparent

cardiac phenotypes[24]. These results suggest that HDAC3 plays distinctive roles from

HDAC1 and HDAC2, although they share high homology and are grouped in the same

HDAC subfamily[19].

TGFß signaling is involved in many processes during early heart development such as

ECM production, endothelial to mesenchymal transition, and myogenesis[11, 49, 50, 51].

Specifically, TGFß signaling plays a critical role during early cardiac myocardial

development[52, 53, 54, 55]. For instance, knockout of either TGFß2 or TGFß3 results in

hypoplastic ventricular myocardium[15], while TGFß1 is reported to inhibit CM

proliferation[55, 56], suggesting the complexity and specificity of TGFß isoforms during early

heart development. In the current study, we found that TGFß3 expression was

significantly decreased in *Hdac3* deleted endocardium, and TGFß3 treatment vigorously

stimulated CM proliferation (Figure 3). Interestingly, inhibition of deacetylase activity of

HDAC3 alone significantly decreased TGFß3 expression. Further, supplementation of

TGFß3 to cardiac endothelial *Hdac3* KO secretome significantly rescued CM proliferation

and myocardial growth defect (Figure 3), suggesting that cardiac endothelial HDAC3

supports trabeculation through inducing TGFß3 in a deacetylase dependent manner.

Interestingly, HDAC3 is also reported to inhibit TGFß1 expression in the second heart

field development in a deacetylase independent manner[23], whereas HDAC3 deteriorates

colorectal cancer progression by inducing TGFß1 through inhibiting miR-296[57]. These

divergent findings illustrate that HDAC3 can either promote or inhibit TGFß signaling

pathway through different molecular mechanisms in a tissue/cell dependent manner.

Besides functioning as growth signals to CMs, TGFß signaling is predominantly known

for its induction on ECMs[50, 58]. During early heart development, ECM is mainly secreted

from endocardium and mediates signal transduction from endocardium to myocardium

supporting trabeculation[57]. Our scRNA-Seq data revealed that ECM is significantly down

regulated in *Hdac3* endothelial KO hearts (Figure 2). P4ha3, a major enzyme for collagen

synthesis[59] and a TGFß downstream target[60], was also significantly decreased in *Hdac3*

endothelial knockout hearts (Figure 2). These results suggest that ECM downregulation

may be secondary to decreased TGFß signaling to contribute to the hypotrabeculation

phenotypes in *Hdac3* endothelial/endocardial knockout hearts.

We identified that HDAC3 suppresses miR-129-5p in inducing TGFß3 (Figure 4). miR-

129-5p plays important roles in cancer by either promoting or inhibiting tumorigenesis

through regulating a variety of downstream targets and pathways[61, 62]. Several studies

have shown that miR-129-5p protects adult hearts from ischemia/reperfusion injuries[63,

[64], while another study demonstrated that treating cultured H9c2 with HDAC pan-inhibitor

Trichostatin A results in decreased CM proliferation with enhanced miR-129-5p

expression[65]. This is consistent with our current findings in the developing endocardium

(Figure 4). In our recent study, we found that HDAC3 induces the expression of IGF2 and

FGF9 by suppressing miR-322/503 in the developing epicardium[26]. Again, these findings

altogether suggest that HDAC3 can induce different sets of growth signals by switching

their downstream miR targets in a tissue/cell context dependent manner. On the other

hand, HDAC3, as a histone modifier, may have many downstream targets that participate

in other signaling pathways and exert various other biological effects. For instance,

several Wnt ligands (e.g., *Wnt4, Wnt5a,* and *Wnt11*) and Hedgehog signaling

components (*Smo, EVC,* and *Gli2*) were also significantly downregulated in *Hdac3* KO

MCECs. The disruption of these signaling pathways may affect cell adhesion, migration

and ciliogenesis[66, 67], thus also contribute to the overall *Hdac3^tko* cardiac phenotypes. We

will be actively investigating these possibilities in our future studies.

In summary, our current study demonstrates that endocardial HDAC3 promotes

myocardial trabeculation by inducing TGFß signaling through repressing miR-129-5p

during early heart development. This finding, along with others, exemplifies the

importance of endocardial signaling for driving myocardial morphogenesis, and thus

provides new insights on the pathogenesis of congenital heart disease. The dual

epigenetic inhibition mechanism may be leveraged in some other contexts such as adult

heart repair/regeneration.

Methods

Mice

Hdac3$^{flox/+}$, *Tie2-Cre*, *Nfatc1$^{Cre/+}$*, *R26^{eYFP}* and *Tnnt2$^{nGFP/+}$* mice were previously described[27, 30, 38, 68, 69]. B6.Cg-Tg(*Tek-cre*)12Flv/J (abbreviated as *Tie2-Cre* in this manuscript), *Hdac3$^{flox/+}$*, and *R26^{eYFP}* mice are available at the Jackson Laboratory (Stock numbers: #004128, #024119 and #006148, respectively). All animal protocols were approved by the Nationwide Children's Institutional Animal Care and Use Committee (IACUC #AR22-00194).

Histology and immunofluorescence staining

All embryo specimens were fixed in 4% paraformaldehyde overnight, dehydrated through an ethanol series, paraffin embedded, and sectioned (6-7μm). Primary antibodies (Major Resources Table) were incubated at 4°C overnight and secondary antibodies (Alexa 488 or Alexa 555, Thermo Fisher Scientific, Waltham, MA) were incubated at room temperature for one hour. Stained slides were imaged on a Zeiss LSM 710 confocal Microscope or a Leica DM6 fluorescence microscope. Epifluorescence and gross images of embryos were imaged on a Leica M205 FCA stereo fluorescence microscope. The number of phosphor-Histone 3 (p-H3)+ cells, Bromodeoxyuridine (BrdU)+ cells, and TUNEL+ cells was quantified using ImageJ software.

Embryonic cardiomyocyte (CM) culture and proliferation assessment

E13.5 ventricular CMs were isolated from dissected embryonic hearts and cultured as previously described[70]. To assess cell proliferation, CMs were fixed with 4%

paraformaldehyde and stained with Ki67 or p-H3 antibody followed by Alexa 594-

conjugated secondary antibody (Thermo Fisher Scientific, Waltham, MA). The

immunostaining was visualized and imaged on a Keyence BZ-X800 microscope. The

percentages of Ki67+GFP+/GFP+, p-H3+GFP+/GFP+ and total number of GFP+ CMs

were quantified using ImageJ software.

Endothelial cell culture, transient transfection, lentiviral infection, and luciferase

assay

Mouse cardiac endothelial cells (MCECs) were purchased from Cedarlane Labs

(Catalogue #CLU510, Burlington, NC). MCECs were cultured in 10% fetal bovine serum

(FBS) supplemented DMEM at 37°C in a humidified incubator with 5% CO_2. To generate

stable *Hdac3* KO MCECs, as previously described[26], we cloned *Hdac3* sgRNA into

lentiCRISPR v2 vector (Addgene #52961; Watertown, MA). Seventy-percent confluent

Lenti-X™ 293T cells (Takara Bio USA, Inc., Mountain View, CA) were transfected with

Hdac3 lentiviral plasmids (*Hdac3* KO or empty vector control [EV])[26], *Tgfβ3* lentiviral

plasmids (*Tgfβ3* KO or empty vector control [EV]) were purchased from VectorBuilder,

PsPAX2 (Addgene #12260) and PMD2G (Addgene #12259). Seventy-two hours after

transfection, supernatants containing lentivirus were collected and filtered through a

40µm cell strainer. Mir-129-5p, miR-129b-3p, miR-129b-5p, mir-129-2-3p, miR-708-3p

mimics and scramble control mimics were custom-ordered from ThermoFischer Scientific

(Waltham, MA). MCECs were transfected with miRNA mimics or control mimics (30 pmol)

using Lipofectamine RNAiMAX (Thermo Fisher Scientific, Waltham, MA). 72 hours after

miR mimics treatment, supernatants were collected. miRZip anti-miR precursor

constructs for miRZip-129-5p and pGreenPuro Scramble Hairpin control constructs were

purchased from System Biosciences (Palo Alto, CA), and lentiviruses were generated

according to the manufacturer's protocol.

Ex Vivo embryonic heart culture

E11.5 embryonic hearts were dissected in cold PBS, and then cultured in DMEM/F-12

media (Thermofisher Scientific, Waltham, MA, #11039021). Embryonic cardiac explants

were treated with either TGFß3 recombinant protein (Sigmal Aldrich, Saint Louis, MO,

#SRP6552, final concentration: 250 ng/ml), *Tgfß3* KO MCEC supernatants, miR-129-5p

mimics-treated MCEC supernatants and their corresponding control media or control

MCEC supernatants for 24 hours at 37 °C. Hearts were then fixed in 4%

paraformaldehyde overnight at 4 °C and processed for paraffin embedding and

sectioning.

Fluorescence activated cell sorting of YFP+ Tie2 derivative cardiac cells

E11.5 hearts were dissected and dissociated with 0.5% trypsin–EDTA into a single cell

suspension. Cell suspension was washed with PBS and YFP+ cells were gated using

their autofluoresence signal and sorted and using a BD Influx cell sorter (BD Biosciences).

The harvested YFP+ cells were used for subsequent gene expression analysis.

RNA isolation, quantitative real-time PCR (qRT-PCR), and bulk RNA-Seq

MCECs and embryonic day (E) 11.5 hearts were used for RNA isolation. E11.5 hearts

were microdissected in cold PBS and snap frozen in liquid nitrogen. The RNeasy Plus

442 Mini Kit (QIAGEN, Germantown, MD) was used to extract total RNA, and cDNAs were

443 generated using the Superscript III kit (Thermo Fisher Scientific, Waltham, MA). To detect

444 mRNA, SYBR Green qRT-PCR was performed. PCR primers for genes of interest are

445 listed in Major Resources Table. To perform mature microRNA expression analyses,

446 purified total RNAs were converted to cDNAs using TaqMan™ MicroRNA Reverse

447 Transcription Kit (Catalogue #4366596, Thermo Fisher Scientific). qRT-PCR reactions

448 were run on StepOne Plus Real-Time PCR System (Applied Biosystems, Waltham, MA).

449 The probe sequences for mmu-miR-129-5p and U6 snRNA were purchased from Thermo

450 Fisher Scientific. For bulk RNA and miRNA-Sequencing, samples were prepared

451 following the provider's guidelines (Novogene Corporation Inc, Sacramento, CA) and

452 sequenced on an Illumina NextSeq500 (San Diego, CA). Sequencing reads were aligned

453 to the UCSC mm10 reference genome using tophat2 and bowtie2 in R. Differential

454 expression of transcripts was calculated using the cufflinks suite in R analyses.

456 ***Single cell RNA-Sequencing (scRNA-seq)***

457 E11.5 heart tissues were dissected in cold PBS and digested with 0.25% trypsin at 37°C

458 for 15 minutes. Digested cardiac cells were filtered with 70 μm cell strainers to remove

459 remaining clumps. Subsequently, 10,000 live cells were captured with Chromium Next

460 GEM Single Cell 3' Reagent Kits v3.1 (10X Genomics, Pleasanton, CA) following

461 manufacturer's instructions. Briefly, single cells were partitioned into nanoliter-scale Gel

462 Beads-in-emulsion (GEMs) using Chromium Next GEM Chip G in the chromium controller.

463 Immediately following GEM generation, the Gel Bead was dissolved, primers were

464 released, and co-partitioned cell was lysed, and its mRNA were reverse transcribed into

barcoded cDNA. After further cleanup and amplification, the cDNA was enzymatically

fragmented and amplified via PCR to generate sufficient mass for library construction.

After end repair, A-tailing, adaptor ligation and sample index PCR, the library comprised

constructs containing the sample index, UMI sequences, barcode sequences, and

Illumina standard sequencing primers P5 and P7 at both ends. The library was sequenced

with Illumina NextSeq S4 platform to a sequencing depth of 140-270 million reads per

sample. Reads were aligned to the Ensembl 104 mouse reference genome including the

sequence for eYFP as a separate contig using STARsolo (version 2.7.10b)[71]. The

resulting count tables (cells x genes) were converted into Annotated Data (anndata)

format for analysis in Python. For quality control, all cells with less than 1500 or more than

9000 genes expressed were excluded. Additionally, cell doublets were filtered out using

the Scrublet method[72]. Gene counts were normalized per cell to a total target sum and

transformed using log1p. Highly variable genes were identified using the Seurat method[73]

and these genes were utilized for dimension reduction with PCA. Finally, cell embedding

was performed with the UMAP algorithm (umap-learn version 0.5.3) and cells were

clustered using leiden (leidenalg version 0.9.1) with resolution 0.2. Scoring of trabecular

and compact CM subtypes was performed using 'score_genes' and differentially

expressed genes between groups were obtained using 'rank_genes_groups', both

methods from the SCANPY package[74].

Western blotting

Cell or tissue lysates were prepared in lysis buffer (20 mM Tris-HCl [pH 7.5], 15 mM NaCl,

1 mM Na_2EDTA, 1 mM EGTA, 1% Triton X-100, 1 µg/ml leupeptin, 2.5 mM sodium

pyrophosphate, 1 mM Na$_3$VO$_4$, and 1 mM β-glycerophosphate) with protease inhibitor

cocktail (Roche, Indianapolis, IN) and 1 mM phenylmethylsulfonyl fluoride. Protein

samples were resolved on 4–12% SDS-PAGE acrylamide gels before transferring to

polyvinylidene fluoride membranes. Primary antibodies were visualized by

chemiluminescence using HRP-conjugated secondary antibodies. Antibodies are listed

in Major Resources Table. Western blots were normalized to loading controls and

densitometric analysis was performed using Image J software.

Statistical analysis

All experiments were independently repeated at least three times, and the number of

samples (n) is stated in figure legends. Results are reported as the mean ± SD. For

statistical analyses, normal distribution of data was assessed by Shapiro-Wilk's test

(when n≥6). For data with small sample size (n<6), non-parametric tests were applied.

For parametric data with normal distribution, the statistical significance of the difference

between means was assessed using the unpaired two tailed Student's t-test between two

groups and the Tukey or Dunnett post hoc test for one-way ANOVA. For non-parametric

data or data with small sample size (n<6), the Mann-Whitney U test and Kruskal-Wallis

test for one-way ANOVA followed by the Dunn post hoc test were performed for

comparisons between two groups. Specific statistical tests are stated in figure legends.

RNA-seq data were background-corrected, variance-stabilized, normalized, and count

matrix extracted through quality control using lumi package for the R programming

language   (http://bioconductor.org/packages/release/bioc/html/lumi.html).   Differential

expression   analysis   was   performed   using   DESeq2   R   package

(https://bioconductor.org/packages/release/bioc/html/DESeq2. html). The resulting *p*-

values were adjusted using the Benjamini and Hochberg's approach for controlling the

False Discovery Rate (FDR)[75]. Genes or miRs with an adjusted *p*-value<0.05 found by

DESeq2 were assigned as differentially expressed. Gene ontology (GO) enrichment

analyses were performed using the Database for Annotation, Visualization and Integrated

Discovery (DAVID) bioinformatics resources (https://david.ncifcrf.gov/). For relative gene

expression analysis of qRT-PCR data, the expression of a gene of interest relative to an

internal loading control gene was calculated using the comparative CT method (-$\Delta\Delta$CT)[76].

The relative protein expression on western blots was quantified by densitometry, Fold

changes for relative gene or protein expression were derived through division of values

in the experimental group by the mean values from the control group. *P*-value<0.05 was

considered statistically significant. All statistical analyses were performed using

GraphPad Prism 8 (GraphPad Prism software, Inc). The images with the value closest to

the mean value were selected as representative images.

Data Availability

All data associated with this study are present in the main text or the supplementary materials. The scRNA-seq, RNA-seq and miRNA-seq data generated in this study have been deposited in the GEO database under accession code GSE229661. All raw data are available from the authors upon reasonable request.

References:

1. Zaidi S, Brueckner M. Genetics and Genomics of Congenital Heart Disease. *Circulation*
research **120**, 923-940 (2017).

2. Qu X, Harmelink C, Baldwin HS. Endocardial-Myocardial Interactions During Early
Cardiac Differentiation and Trabeculation. *Front Cardiovasc Med* **9**, 857581 (2022).

3. Wu M. Mechanisms of Trabecular Formation and Specification During Cardiogenesis.
Pediatric cardiology **39**, 1082-1089 (2018).

4. Lin W, Li D. Zinc and Zinc Transporters: Novel Regulators of Ventricular Myocardial
Development. *Pediatric cardiology* **39**, 1042-1051 (2018).

5. Zhang W, Chen H, Qu X, Chang CP, Shou W. Molecular mechanism of ventricular
trabeculation/compaction and the pathogenesis of the left ventricular noncompaction
cardiomyopathy (LVNC). *American journal of medical genetics Part C, Seminars in*
medical genetics **163C**, 144-156 (2013).

6. Zhang H, Lui KO, Zhou B. Endocardial Cell Plasticity in Cardiac Development, Diseases
and Regeneration. *Circulation research* **122**, 774-789 (2018).

7. Andries LJ, Sys SU, Brutsaert DL. Morphoregulatory interactions of endocardial
endothelium and extracellular material in the heart. *Herz* **20**, 135-145 (1995).

8.   Camenisch TD*, et al.* Disruption of hyaluronan synthase-2 abrogates normal cardiac

morphogenesis and hyaluronan-mediated transformation of epithelium to mesenchyme. *J*

Clin Invest **106**, 349-360 (2000).

9.   Hatano S*, et al.* Versican/PG-M is essential for ventricular septal formation subsequent to

cardiac atrioventricular cushion development. *Glycobiology* **22**, 1268-1277 (2012).

10.   Chen JL, Colgan TD, Walton KL, Gregorevic P, Harrison CA. The TGF-beta Signalling

Network in Muscle Development, Adaptation and Disease. *Adv Exp Med Biol* **900**, 97-131

(2016).

11.   Sorensen DW, van Berlo JH. The Role of TGF-beta Signaling in Cardiomyocyte

Proliferation. *Curr Heart Fail Rep* **17**, 225-233 (2020).

12.   Zhang Y, Yang X. The Roles of TGF-beta Signaling in Cerebrovascular Diseases. *Front*

Cell Dev Biol **8**, 567682 (2020).

13.   Moustakas A, Heldin CH. Non-Smad TGF-beta signals. *J Cell Sci* **118**, 3573-3584 (2005).

14.   Zhang YE. Non-Smad pathways in TGF-beta signaling. *Cell Res* **19**, 128-139 (2009).

15.   Azhar M*, et al.* Transforming growth factor beta in cardiovascular development and

function. *Cytokine Growth Factor Rev* **14**, 391-407 (2003).

16.   Koo SH, Cunningham MC, Arabshahi B, Gruss JS, Grant JH, 3rd. The transforming growth

factor-beta 3 knock-out mouse: an animal model for cleft palate. *Plast Reconstr Surg* **108**,

938-948; discussion 949-951 (2001).

17.   Sanford LP*, et al.* TGFbeta2 knockout mice have multiple developmental defects that are

non-overlapping with other TGFbeta knockout phenotypes. *Development* **124**, 2659-2670

(1997).

18.   Kulkarni AB, Karlsson S. Transforming growth factor-beta 1 knockout mice. A mutation in

one cytokine gene causes a dramatic inflammatory disease. *Am J Pathol* **143**, 3-9 (1993).

19.   Haberland M, Montgomery RL, Olson EN. The many roles of histone deacetylases in

development and physiology: implications for disease and therapy. *Nature reviews*

Genetics **10**, 32-42 (2009).

20.   Dsilva P, Pai P, Shetty MG, Babitha KS. The role of histone deacetylases in embryonic

development. *Mol Reprod Dev* **90**, 14-26 (2023).

21.   Wang Z, Zhao YT, Zhao TC. Histone deacetylases in modulating cardiac disease and their

clinical translational and therapeutic implications. *Exp Biol Med (Maywood)* **246**, 213-225

(2021).

22.   Bhaskara S*, et al.* Deletion of histone deacetylase 3 reveals critical roles in S phase

progression and DNA damage control. *Mol Cell* **30**, 61-72 (2008).

23.   Lewandowski SL, Janardhan HP, Trivedi CM. Histone Deacetylase 3 Coordinates

Deacetylase-independent Epigenetic Silencing of Transforming Growth Factor-beta1

(TGF-beta1) to Orchestrate Second Heart Field Development. *The Journal of biological*

chemistry **290**, 27067-27089 (2015).

24.   Janardhan HP, Milstone ZJ, Shin M, Lawson ND, Keaney JF, Jr., Trivedi CM. Hdac3

regulates lymphovenous and lymphatic valve formation. *J Clin Invest* **127**, 4193-4206

(2017).

25.   Poleshko A*, et al.* Genome-Nuclear Lamina Interactions Regulate Cardiac Stem Cell

Lineage Restriction. *Cell* **171**, 573-587 e514 (2017).

26.   Jang J*, et al.* Epicardial HDAC3 Promotes Myocardial Growth Through a Novel MicroRNA

Pathway. *Circulation research* **131**, 151-164 (2022).

27.   Koni PA, Joshi SK, Temann UA, Olson D, Burkly L, Flavell RA. Conditional vascular cell

adhesion molecule 1 deletion in mice: impaired lymphocyte migration to bone marrow.

The Journal of experimental medicine **193**, 741-754 (2001).

28.   de Lange WJ, Halabi CM, Beyer AM, Sigmund CD. Germ line activation of the Tie2 and

SMMHC promoters causes noncell-specific deletion of floxed alleles. *Physiol Genomics*

35, 1-4 (2008).

29.   Kisanuki YY, Hammer RE, Miyazaki J, Williams SC, Richardson JA, Yanagisawa M. Tie2-

Cre transgenic mice: a new model for endothelial cell-lineage analysis in vivo. *Dev Biol*

230, 230-242 (2001).

30.   Wu B, *et al.* Endocardial cells form the coronary arteries by angiogenesis through

633   myocardial-endocardial VEGF signaling. *Cell* **151**, 1083-1096 (2012).

635   31.   Wu T, *et al.* PRDM16 Is a Compact Myocardium-Enriched Transcription Factor Required

636   to Maintain Compact Myocardial Cardiomyocyte Identity in Left Ventricle. *Circulation* **145**,

637   586-602 (2022).

639   32.   Hassan MK, Kumar D, Patel SA, Dixit M. EEF1A2 triggers stronger ERK mediated

640   metastatic program in ER negative breast cancer cells than in ER positive cells. *Life Sci*

641   **262**, 118553 (2020).

643   33.   Fish JE, *et al.* Dynamic regulation of VEGF-inducible genes by an ERK/ERG/p300

644   transcriptional network. *Development* **144**, 2428-2444 (2017).

646   34.   Thornton S, Anand N, Purcell D, Lee J. Not just for housekeeping: protein initiation and

647   elongation factors in cell growth and tumorigenesis. *J Mol Med (Berl)* **81**, 536-548 (2003).

649   35.   Anbara T, Sharifi M, Aboutaleb N. Endothelial to Mesenchymal Transition in the

650   Cardiogenesis and Cardiovascular Diseases. *Curr Cardiol Rev* **16**, 306-314 (2020).

652   36.   Del Monte-Nieto G, *et al.* Control of cardiac jelly dynamics by NOTCH1 and NRG1 defines

653   the building plan for trabeculation. *Nature* **557**, 439-445 (2018).

655   37.   Barbieri SS, Weksler BB. Tobacco smoke cooperates with interleukin-1beta to alter beta-

656   catenin trafficking in vascular endothelium resulting in increased permeability and

induction of cyclooxygenase-2 expression in vitro and in vivo. *FASEB J* **21**, 1831-1843

(2007).

38.　Yan J*, et al.* A series of robust genetic indicators for definitive identification of

cardiomyocytes. *J Mol Cell Cardiol* **97**, 278-285 (2016).

39.　Tian Y, Morrisey EE. Importance of myocyte-nonmyocyte interactions in cardiac

development and disease. *Circulation research* **110**, 1023-1034 (2012).

40.　Silva AC, Pereira C, Fonseca A, Pinto-do OP, Nascimento DS. Bearing My Heart: The

Role of Extracellular Matrix on Cardiac Development, Homeostasis, and Injury Response.

Front Cell Dev Biol **8**, 621644 (2020).

41.　Paige SL*, et al.* A temporal chromatin signature in human embryonic stem cells identifies

regulators of cardiac development. *Cell* **151**, 221-232 (2012).

42.　Wamstad JA*, et al.* Dynamic and coordinated epigenetic regulation of developmental

transitions in the cardiac lineage. *Cell* **151**, 206-220 (2012).

43.　Lewandowski SL, Janardhan HP, Trivedi CM. Histone Deacetylase 3 Coordinates

Deacetylase-independent Epigenetic Silencing of Transforming Growth Factor-beta1

(TGF-beta1) to Orchestrate Second Heart Field Development. *The Journal of biological*

chemistry **290**, 27067-27089 (2015).

44.　Montgomery RL*, et al.* Maintenance of cardiac energy metabolism by histone deacetylase

3 in mice. *J Clin Invest* **118**, 3588-3597 (2008).

45.   Artap S*, et al.* Endocardial Hippo signaling regulates myocardial growth and
685   cardiogenesis. *Dev Biol* **440**, 22-30 (2018).

687   46.   Tian X*, et al.* Subepicardial endothelial cells invade the embryonic ventricle wall to form
688   coronary arteries. *Cell Res* **23**, 1075-1090 (2013).

690   47.   Celikyurt U, Karauzum K, Sahin T, Agacdiken A, Vural A, Ural D. Association between
691   resolution of fragmented QRS and response to cardiac resynchronization therapy. *Ann*
692   *Noninvasive Electrocardiol* **20**, 126-131 (2015).

694   48.   Yang Y, Oliver G. Development of the mammalian lymphatic vasculature. *J Clin Invest*
695   **124**, 888-897 (2014).

697   49.   Conway SJ, Doetschman T, Azhar M. The inter-relationship of periostin, TGF beta, and
698   BMP in heart valve development and valvular heart diseases. *ScientificWorldJournal* **11**,
699   1509-1524 (2011).

701   50.   Frangogiannis NG. Transforming growth factor-beta in myocardial disease. *Nat Rev*
702   *Cardiol* **19**, 435-455 (2022).

704   51.   Lee JH, Massague J. TGF-beta in developmental and fibrogenic EMTs. *Semin Cancer*
705   *Biol* **86**, 136-145 (2022).

707   52.   Chakrabarti M*, et al.* Transforming Growth Factor Beta3 is Required for Cardiovascular
708   Development. *J Cardiovasc Dev Dis* **7**, (2020).

53.   Dunker N, Krieglstein K. Tgfbeta2 -/- Tgfbeta3 -/- double knockout mice display severe

midline fusion defects and early embryonic lethality. *Anat Embryol (Berl)* **206**, 73-83

(2002).

54.   Kruithof BP, Kruithof-De-Julio M, Poelmann RE, Gittenberger-De-Groot AC, Gaussin V,

Goumans MJ. Remodeling of the myocardium in early trabeculation and cardiac valve

formation; a role for TGFbeta2. *Int J Dev Biol* **57**, 853-863 (2013).

55.   Kodo K*, et al.* iPSC-derived cardiomyocytes reveal abnormal TGF-beta signalling in left

ventricular non-compaction cardiomyopathy. *Nat Cell Biol* **18**, 1031-1042 (2016).

56.   Engelmann GL, Boehm KD, Birchenall-Roberts MC, Ruscetti FW. Transforming growth

factor-beta 1 in heart development. *Mech Dev* **38**, 85-97 (1992).

57.   Li J*, et al.* HDAC3 deteriorates colorectal cancer progression via microRNA-296-

3p/TGIF1/TGFbeta axis. *J Exp Clin Cancer Res* **39**, 248 (2020).

58.   Algeciras L, Palanca A, Maestro D, RuizdelRio J, Villar AV. Epigenetic alterations of

TGFbeta and its main canonical signaling mediators in the context of cardiac fibrosis. *J*

Mol Cell Cardiol **159**, 38-47 (2021).

59.   Van Den Diepstraten C, Papay K, Bolender Z, Brown A, Pickering JG. Cloning of a novel

prolyl 4-hydroxylase subunit expressed in the fibrous cap of human atherosclerotic plaque.

Circulation **108**, 508-511 (2003).

60.   Luo Y, *et al.* A novel profibrotic mechanism mediated by TGFbeta-stimulated collagen

736   prolyl hydroxylase expression in fibrotic lung mesenchymal cells. *J Pathol* **236**, 384-394

737   (2015).

739   61.   Deng B, Tang X, Wang Y. Role of microRNA-129 in cancer and non-cancerous diseases

740   (Review). *Exp Ther Med* **22**, 918 (2021).

742   62.   Xu S, *et al.* The Role of miR-129-5p in Cancer: A Novel Therapeutic Target. *Curr Mol*

743   *Pharmacol* **15**, 647-657 (2022).

745   63.   Ma R, Chen X, Ma Y, Bai G, Li DS. MiR-129-5p alleviates myocardial injury by targeting

746   suppressor of cytokine signaling 2 after ischemia/reperfusion. *Kaohsiung J Med Sci* **36**,

747   599-606 (2020).

749   64.   Xing J, Liu J, Liu J, Xu Z. miR-129-5p ameliorates ischemia-reperfusion injury by targeting

750   HMGB1 in myocardium. *Gen Physiol Biophys* **39**, 461-470 (2020).

752   65.   Majumdar G, Raghow R. Trichostatin A induces a unique set of microRNAs including miR-

753   129-5p that blocks cyclin-dependent kinase 6 expression and proliferation in H9c2 cardiac

754   myocytes. *Mol Cell Biochem* **415**, 39-49 (2016).

756   66.   Quadri N, Upadhyai P. Primary cilia in skeletal development and disease. *Exp Cell Res*

757   **431**, 113751 (2023).

759   67.   Flaherty MP, Kamerzell TJ, Dawn B. Wnt signaling and cardiac differentiation. *Prog Mol*

760   *Biol Transl Sci* **111**, 153-174 (2012).

68. Mullican SE*, et al.* Histone deacetylase 3 is an epigenomic brake in macrophage alternative activation. *Genes & development* **25**, 2480-2488 (2011).

69. Srinivas S*, et al.* Cre reporter strains produced by targeted insertion of EYFP and ECFP into the ROSA26 locus. *BMC Dev Biol* **1**, 4 (2001).

70. Beall MC, Li D, Jang J. Isolation of Embryonic Cardiomyocytes and Cell Proliferation Assay Using Genetically Engineered Reporter Mouse Model. *Bio Protoc* **13**, e4802 (2023).

71. Dobin A*, et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21 (2013).

72. Wolock SL, Lopez R, Klein AM. Scrublet: Computational Identification of Cell Doublets in Single-Cell Transcriptomic Data. *Cell Syst* **8**, 281-291 e289 (2019).

73. Satija R, Farrell JA, Gennert D, Schier AF, Regev A. Spatial reconstruction of single-cell gene expression data. *Nat Biotechnol* **33**, 495-502 (2015).

74. Wolf FA, Angerer P, Theis FJ. SCANPY: large-scale single-cell gene expression data analysis. *Genome Biol* **19**, 15 (2018).

75. Wang L, Feng Z, Wang X, Wang X, Zhang X. DEGseq: an R package for identifying differentially expressed genes from RNA-seq data. *Bioinformatics* **26**, 136-138 (2010).

76. Schmittgen TD, Livak KJ. Analyzing real-time PCR data by the comparative C(T) method. *Nature protocols* **3**, 1101-1108 (2008).

Acknowledgments

We thank Dr. Dennis Lewandowski in the Heart Center at the Nationwide Children's Hospital for his critical reading and editing of the manuscript.

Author contributions

J.J. and D.L. conceived and designed the study, analyzed the experiments, wrote the paper. M.B and M.L performed scRNA-Seq data analysis. Y.J.K and E.K. performed the experiments. C.C. provided $Tnnt2^{nGFP/+}$ mice. V.G., J.J. and D.L. edited manuscript. All authors reviewed the results and approved the final version of the manuscript.

Competing Interests

None.

Sources of Funding

This work was partially supported by the National Heart, Lung, and Blood Institute R01 grant HL153406 (D. L.), American Heart Association (AHA)-Career Development Award 23CDA1046244 and AHA-Innovative Project Award 23IPA1048560 (J. J.).

Figures and Figure legends

Figure 1. Endothelial specific deletion of *Hdac3* results in hypotrabeculation.

(A) Hematoxylin and eosin staining on cross sections of an E12.5 *Hdac3^tko^* embryo (*Hdac3^f/f^; Tie2-Cre/+*) and a littermate control (CTL) embryo (*Hdac3^f/f^*). Arrows mark endocardium and arrowheads point to trabeculae. Quantifications are shown on the bottom (CTL: n=8, *Hdac3^tko^*: n=7). Scale bars, 200 μm (main panels) and 25 μm (insets).

(B) Decreased cardiomyocyte (CM) proliferation in E11.5 *Hdac3^tko^* hearts. Immunofluorescence staining of Bromodeoxyuridine/5-bromo-2'-deoxyuridine (BrdU) and phospho-Histone H3 (p-H3) staining. High-power Images in the trabecular myocardium (Tra) and in the compact myocardium (Com) are highlighted in white and yellow boxes respectively. Scale bars, 200 μm. Quantification of proliferation index in the trabecular

820   and compact myocardium respectively are shown on the bottom. BrdU (CTL: n=8,

821   *Hdac3^{tko}*: n=7); p-H3 (n=6 in each group). *P*-values were determined by unpaired two

822   tailed Student's *t*-test.

Figure 2. Single-cell RNA sequencing reveals decreased expression of ECM genes in *Hdac3^tko^* hearts.

(A) Schematic workflow of scRNA-Seq experimental design. Representative micrographs

of GFP immunofluorescence staining of E11.5 *Hdac3^{tko}* embryo (*Hdac3^{f/f}; Tie2-Cre/+;*

R26R^{eYFP/+}) and a littermate control (CTL) embryo (*Hdac3^{f/+}; Tie2-Cre/+; R26R^{eYFP/+}*).

Scale bars, 250 μm. **(B)** UMAP visualization of cell populations in an E11.5 control heart.

(C) Trabecular and compact cardiomyocyte (CM) score were calculated within *Tnni*+ CM

population. Left UMAP showed CMs split into a lower cluster that has a high trabecular

score, and an upper cluster that has a high compact score. Right UMAP shows the

expression level of *Bmp10*+ per cell. **(D)** Dot plots of differentially expressed genes in

Hdac3^{tko} and CTL trabecular clusters. Violin plots of expression of *Egr1* and *Eef1a2* in

Hdac3^{tko} and CTL trabecular clusters. **(E)** The UMAP on the left shows that YFP+ cells

are distributed in cardiac endothelial cell (CEC, *Cdh*+), mesenchymal cell 1 (MC1,

Pdgfra+), and mesenchymal cell 2 (MC2, *Dcn*+) clusters. The overlapping UMAPs of

Hdac3^{tko} and CTL YFP+ cells are shown on the right. **(F)** Gene Ontology pathway analysis

of top marker genes in CTL-specific YFP+ CEC cluster (Cut off value: -log (FDR)>5). **(G)**

Heatmap of extracellular matrix (ECM) associated genes in E11.5 *Hdac3^{tko}* and CTL

hearts. Data were extracted from YFP+ CEC clusters in the scRNA-Seq dataset. **(H)** qRT-

PCR analysis of *Col1a1*, *Col1a2*, *Col2a1*, *Col3a1* and *P4ha3* in E11.5 *Hdac3^{tko}* heart as

compared to CTL hearts (n=8 in each group for *Col1a1*, *Col1a2*, *Col2a1* and *Col3a1,* n=5

in each group for *P4ha3*). *P*-values were determined by unpaired two tailed Student's *t*-

test. **(I)** Illustration of decreased expression of proliferation related genes in trabecular

CMs and ECM-associated genes in CECs of *Hdac3^{tko}* hearts.

Figure 3. Cardiac endothelial *Hdac3* knockout results in downregulation of TGFβ3

and decreased cardiomyocyte proliferation.

(A) Generation of *Hdac3* knockout (KO) and empty vector control (EV) mouse cardiac

endothelial cells (MCECs) by CRISPR/Cas9. Deletion of *Hdac3* was verified by western

blot. Quantification is shown below (n=6 in each group). **(B)** The effects of *Hdac3* KO

MCEC supernatants on E13.5 *Tnnt2^{nGFP/+}* cardiomyocytes (CMs) proliferation.

Representative immunofluorescence micrographs are shown. Scale bar, 275 μm.

Percentage of Ki67+ CMs and total number of GFP+ CMs were quantitated (n=4 in each

group). **(C)** Quantification of *Tgfβ3* expression in *Hdac3* KO MCECs, or RGFP966-treated

MCECs (10 uM) by qRT-PCR. *Gapdh* was used as cDNA loading control (n=4 in each

group). **(D)** Downregulation of *Tgfβ3* gene expression in *Hdac3* deficient cardiac

endothelial cells. YFP+ E11.5 cardiac endothelial cells (CECs) from *Hdac3^{tko}* (*Hdac3^{f/f};*

Tie2-Cre/+; R26^{eYFP/+}) and CTL (*Hdac3^{f/+}; Tie2-Cre/+; R26^{eYFP/+}*) hearts were sorted by

fluorescence-activated cell sorting. Quantification of *Hdac3* and *Tgfβ3* expression by

qPCR was shown on the right. n=4 for each group. *Gapdh* was used as cDNA loading

control (n=4 in each group). **(E)** Representative Western blot and quantification of TGFβ3

expression in E11.5 *Hdac3^{tko}* hearts (n=12) and littermate CTL hearts (n=8). **(F)** Secretion

of TGFβ3 from *Hdac3* KO and EV MCECs. Coomassie brilliant blue staining of total

extracted proteins from supernatants served as protein loading controls. TGFβ3 in the

MCEC supernatants were detected by western blot. Quantifications are shown on the

right (n=4 in each group). **(G)** Rescue of CM proliferation defect by *Hdac3* KO MCECs

supernatant supplemented with recombinant TGFβ3 (working concentration: 250 ng/ml).

Representative immunofluorescence micrographs are shown. Scale bar, 200 μm.

Percentage of Ki67+ CMs and total number of GFP+ CMs were quantitated (n=7 in each

group). *P*-values were determined by unpaired two tailed Student's *t*-test for **(A)** and **(E)**,

Mann-Whitney U test for **(C)**, **(D)** and **(F)** and one-way ANOVA followed by the Tukey

post hoc test for **(B) and (G)**.

Figure 4. HDAC3 induces Tgfβ3 by repressing miR-129-5p.

(A) Heatmap of microRNA (miR) sequencing of *Hdac3* knockout (KO) and empty vector control (EV) mouse cardiac endothelial cells (MCECs). MiRs with reads less than 100 were discarded and miRNA expression levels were normalized by TPM (transcript per million) values (TPM = [miRNA total reads/total clean reads] × 10^6). Log_2 fold changes were calculated by TPM per miR in the KO group divided by the mean TPM per miR in the EV group then followed by calculation of Log (fold change, 2). Significantly downregulated miRs are shown in blue, and significantly up-regulated miRs are shown in red. Cut-off criteria: Log2>1.5 and adjusted *P*-value<0.01. **(B)** Quantification of the expression of *Tgfβ3* in MCECs after miR mimics treatment (final concentration 10 nM) by qRT-PCR. *Gapdh* was used as a cDNA loading control (n=6 in each group). Relative gene expression was compared to the miR scrambles (scCTL) group. **(C)** Representative Western blot and quantification of the expression of TGFβ3 after miR-129-5p mimics treatment by western blot (n=8 in each group). **(D)** (Left) Quantification of *Tgfβ3* expression in *Hdac3* KO and EV MCECs (n=4 in each group). (Right) Quantification of the expression of *Tgfβ3* in MCECs after Hdac3 inhibitor treatment (RGFP966, final concentration 5 uM) (n=4 in each group). **(E)** Quantification of the expression of *Tgfβ3* in *Hdac3^tko^* heart lysates (n=9 in each group). **(F)** Quantification of the expression of miR-129-5p by qRT-PCR**.** *Hdac3* KO or EV MECs were transfected with LentimiRa-GFP-miRZip129 (miRZIP129) or pGreenPuro Scramble Hairpin control lentivirus (scCTL) and transfected cells were selected using puromycin (n=4 in each group)**. (G)** Representative Western blot and quantification of the expression of TGFβ3 after treatment with miRZIP129 or scCTL in *Hdac3* KO and EV MCECs (n=6 in each group). **(H)** Knockdown

of miR-129-5p rescues CM proliferation deficit induced by *Hdac3* KO MCEC supernatants.

Representative immunofluorescence micrographs are shown. Scale bar, 200 μm.

Quantification of percentage of phospho-histone H3 (p-H3)+ CMs was shown on the right.

Independent samples: n=9 in each group. The protein expression of TGFβ3 *P*-values

were determined by unpaired two tailed Student's *t*-test for **(C)** and **(E)**, the Mann-Whitney

U test for **(D)** and the one-way ANOVA followed by Tukey post hoc test for **(B)**, **(F)**, **(G)**

and **(H)**. **(I)** Working model: in the developing endocardium, HDAC3 represses the

expression of miR-129-5p to release their suppression on the expression of TGFβ3.

When *Hdac3* is deleted, the expression of miR-129-5p is increased, which subsequently

suppresses the expression of TGFβ3 to a stronger extent, and the decrease of TGFβ3,

together with decreased ECM, leads to hypotrabeculation.