Endocardial HDAC3 is required for myocardial trabeculation

Jihyun Jang[1,2], Mette Bentsen[3], Ye Jun Kim[4], Erick Kim[4], Vidu Garg[1,2], Chen-Leng Cai[5],

Mario Looso[3], Deqiang Li[1, 2]

[1]Center for Cardiovascular Research, Abigail Wexner Research Institute, Nationwide

Children's Hospital, Columbus, OH 43215, USA

[2]Department of Pediatrics, The Ohio State University College of Medicine, Columbus, OH

43215, USA

[3]Bioinformatics Core Unit (BCU), Max Planck Institute for Heart and Lung Research,

61231, Bad Nauheim, Germany

[4]Department of Surgery, University of Maryland School of Medicine, Baltimore, MD 21201,

USA

[5]Department of Pediatrics, Herman Wells Center for Pediatric Research, Indiana

University School of Medicine, Indianapolis, IN 46201, USA

Address correspondence to:

Deqiang Li, MD&PhD

575 Children's Drive, Room WB4329,

Columbus, OH 43215, USA

Phone: 614-355-5826

Email: Deqiang.Li@nationwidechildrens.org

Abstract

Failure of proper ventricular trabeculation is often associated with congenital heart disease (CHD). Support from endocardial cells, including the secretion of extracellular matrix (ECM) and growth factors is critical for trabeculation. However, it is poorly understood how the secretion of ECM and growth factors is initiated and regulated by endocardial cells.

We found that genetic knockout (KO) of histone deacetylase 3 (*Hdac3*) in the endocardium in mice resulted in early embryo lethality and ventricular hypotrabeculation. Single cell RNA sequencing identified significant downregulation of ECM components in *Hdac3* KO endocardial cells. Secretome from cultured *Hdac3* KO mouse cardiac endothelial cells lacked transforming growth factor ß3 (TGFß3) and showed significantly reduced capacity in stimulating cultured cardiomyocyte proliferation, which was remarkably rescued by TGFß3 supplementation. Mechanistically, we identified that HDAC3 induced *Tgfß3* expression through repressing microRNA(miR)-129-5p.

Our findings provide novel insights into the pathogenesis of CHD and conceptual strategies to promote myocardial regeneration.

Introduction

Congenital heart disease (CHD) is the most common congenital disorder, and the leading

cause of mortality from birth defects[1]. Defective myocardial growth is either the primary

manifestation or associated with many forms of CHD[2, 3]. In the early primitive heart

(mouse embryonic day 8-10), inner cardiac endothelial cells (CECs), also called

endocardial cells, and outer cardiomyocytes (CMs), which are separated by extracellular

matrix (ECM), co-contribute to early myocardial trabeculation (i.e., formation of

myocardial meshwork extending to cardiac chamber)[4-6]. CECs juxtapose myocardial

trabeculae and lead myocardial trabeculation by secretion of growth factors such as

Neuregulin 1 (NRG1) and transforming growth factor ß (TGFß)[2, 3, 5]. Meanwhile, CECs

support trabeculation by secreting ECM (e.g., various collagens, versican) that facilitate

the travelling of CEC secreted growth factors to trabecular myocardium and support

trabecular CM proliferation[4, 7]. Surprisingly, disruption of ECM components alone (e.g.,

hyaluronan synthase-2, versican) prevents the proper formation of trabeculae in mice[8, 9].

Although CECs play such important roles during ventricular trabeculation, it remains

poorly understood how these ECM proteins and growth factors are regulated in the

developing CECs.

TGFß signaling pathway is involved in many fundamental cellular processes in both

adults and developing embryos including cell proliferation, differentiation, migration, and

apoptosis[10, 11]. The TGFß superfamily includes many different ligands (e.g., Bone

morphogenetic proteins, Growth and differentiation factors, Activin, Nodal, TGFß). These

ligands bind to their specific type II receptors, which triggers phosphorylation of type I

receptor, eventually leads to phosphor-Smad2/3 localized to the nucleus and drives a

variety of downstream target gene expression, which is the canonical TGFß signaling

pathway[12]. In addition, the binding of ligands to receptors can also directly activate

multiple non-Smad mediated intracellular signaling pathways such as MAP kinase

pathways and phosphatidylinositol-3-kinase/AKT pathways[13, 14]. These non-Smad

(noncanonical) TGFß signaling pathways are equally critical for driving some specific

cellular processes (e.g., cell proliferation, cell survival, endothelial/epithelial

mesenchymal transition)[13, 14]. During cardiovascular development, unsurprisingly, TGFß

signaling is implicated in most morphological events including cardiac lineage

specification, cardiac looping, valvogenesis, and cardiomyogenesis[15]. The TGFß family

includes TGFß1, TGFß2, and TGFß3. Although TGFß1, TGFß2 and TGFß3 are

abundantly expressed in the developing endocardium and epicardium, it appears that

they do not perform redundant functions: individual knockout results in different

phenotypes[15]. For instance, global knockout of either *Tgfß2* or *Tgfß3* result in early

perinatal lethality, presenting hypoplastic hearts with thin myocardium[16, 17], suggesting

that they play critical roles for myocardial growth. In contrast, global knockout of *Tgfß1*

lead to partial embryonic lethality with survivors appearing phenotypically normal in first

the two weeks of life[18]. Nonetheless, it is poorly understood how the expression of TGFß

ligands are regulated in the early developing heart.

Epigenetic modulators such as histone deacetylases (HDACs) are critical for early heart

development as key regulators of regulating gene expresison[19, 20]. There are 18 known

human HDACs, which can be subdivided into four classes based on their function and

DNA sequence similarity. Class I HDACs include HDAC1, HDAC2, HDAC3, and HDAC8,

and are predominantly nuclear and ubiquitously expressed[21]. Although class I HDACs

have high degree of homology, global knockout of each causes early embryonic lethality[19],

suggesting distinct roles during early heart development. Global deletion of *Hdac3* leads

to early embryonic lethality at gastrulation[22]. In the developing heart, lack of *Hdac3* in

cardiac progenitors disrupts secondary heart field development[23], lymphovenous valve

morphogenesis[24], and CM lineage specification[25]. Recently, we reported that HDAC3 in

the developing epicardium plays a critical role in supporting compact myocardial wall

expansion[26]. It is unclear whether or how HDAC3 in the developing endocardium impacts

myocardial trabeculation.

In the current study, we investigated the role of HDAC3 in the developing endocardium

during early heart development.

Results

Endothelial/Endocardial cell specific deletion of *Hdac3* results in embryonic lethality and hypotrabeculation.

To study the role of HDAC3 in the cardiac endothelium during early heart development, *Hdac3* was specifically deleted in the endothelial cells using *Tie2-Cre*[27], while its expression in other cells including CMs remains intact (Supplemental Figure 1). Note: since *Tie2-Cre* is transiently expressed in oocytes[28], the mating scheme was always *Hdac3^f/+^; Tie2-Cre/+* (♂) and *Hdac3^f/f^* (♀). By recovering the progenies at a series of embryonic and postnatal stages, we identified that *Hdac3* KO in the developing endothelium resulted in early embryonic lethality starting at E12.5 (Supplemental Table 1). To investigate the potential cause of early embryonic lethality, we performed histological analyses. Interestingly, in Tie2 endothelial *Hdac3* KO embryos (*Hdac3^tko^*), there was no noticeable phenotype in the various endothelial cell derived tissue where the Tie2-lineage contributes, such as endocardium (Figure 1, Supplemental Figure 2) and blood vessels (Supplemental Figure 3), given that *Tie2-Cre* is active in all endothelial cells in the body[29]. In contrast, there were myocardial defects in *Hdac3^tko^* embryos starting at E11.5: *Hdac3^tko^* trabeculae appeared to be significantly sparser and shorter as compared to the thick and long webbing like trabecular network in littermate control hearts (Supplemental Figure 2). This cardiac phenotype in *Hdac3^tko^* mice was more prominent at E12.5 (Figure 1A). The compact layer was also significantly thinner in *Hdac3^tko^* hearts. To further validate the myocardial trabecular phenotype in endothelial *Hdac3* deleted hearts, we specifically deleted *Hdac3* in the endocardium using *Nfatc1^Cre/+^*, endocardial specific Cre[30], which also resulted in embryonic lethality (Supplemental Table 2).

Hdac3^{nko} (*Hdac3^{f/f}; Nfatc1^{Cre/+}*) hearts presented a similar cardiac phenotype, albeit at a

later stage (E14.5): smaller and sparser trabeculae, thinner compact wall (Supplemental

Figure 4). This is consistent with the achievement of late deletion of *Hdac3* in the

endocardium by *Nfatc1^{Cre/+}* (Supplemental Figure 4). Nonetheless, these results confirm

that the cardiac phenotypes resulting from *Tie2* endothelial *Hdac3* deletion are primarily

from the endocardium, and not secondary to another endothelium (e.g., vascular).

Hdac3^{tko} embryos were used for all subsequent mechanistic studies.

Endothelial *Hdac3* knockout hearts display decreased trabecular CM proliferation.

Next, we investigated whether altered cell proliferation and/or apoptosis contribute to the

hypotrabeculation phenotypes in *Hdac3^{tko}* hearts. At E11.5, the percentage of either p-

H3+ or BrdU+ CMs is significantly lower in *Hdac3^{tko}* trabecular and compact myocardium

as compared to littermate control hearts (Figure 1B). This finding is consistent with the

hypoplastic cardiac phenotypes seen in *Hdac3^{tko}* hearts (Figure 1A). In contrast, there

was no significant difference in cell apoptosis between *Hdac3^{tko}* hearts and littermate

control hearts (Supplemental Figure 5). Decreased CM proliferation of *Hdac3^{tko}* hearts

may result from disrupted communication from the endocardium to the trabecular

myocardium. We performed single cell RNA-sequencing (scRNA-seq) on E11.5 *Hdac3^{tko}*

hearts and littermate control hearts (Figure 2A). We obtained high-quality scRNA-seq

profiles for 23,468 and 25,547 cells from 4 *Hdac3^{tko}* (*Hdac3^{f/f}; Tie2-Cre/+; R26^{eYFP/+}*) and

4 control (3 of *Hdac3^{f/+}; Tie2-Cre/+; R26^{eYFP/+}* and 1 of *Hdac3^{f/+}; Tie2-Cre/+*) heart single

cell suspensions, respectively. Unsupervised clustering successfully separated 6 distinct

clusters including endocardial cells and CMs in these samples (Figure 2B and

Supplemental Figure 6). Referring to a recent transcriptome profiling methodology for

clustering embryonic CMs[31], we re-clustered CMs into "trabecular" and "compact" clusters

based on the overall scores after applying a few known trabecular and compact markers

(trabecular: *Bmp10*, *Nppa* and *Thbs4*; compact: *Hey2*, *Mycn* and *Tnnt1*) (Supplemental

Figure 7A and Figure 2C). Among them, *Bmp10+* CMs nicely represent the trabecular

cluster (Figure 2C), and thus we focused on this population for subsequent analyses since

Hdac3^{tko} hearts mainly display hypotrabeculation phenotypes (Figure 1). Overall, the total

number of *Bmp10+* trabecular CMs were not significantly different between *Hdac3^{tko}* and

littermate control hearts (Supplemental Figure 7B). Note: 10,000 cells/sample was

sequenced. Next, we performed differential gene expression (DGE) analyses on *Hdac3^{tko}*

and littermate control trabecular clusters. We found that the expression of eukaryotic

translation elongation factor 1 alpha 2 (*Eef1a2*) and early growth response-1 (*Egr-1*) are

significantly down regulated in *Hdac3^{tko}* trabeculae as compared to control trabeculae

(Figure 2D). *Eef1a2* and *Egr1* are known to be critical for cell growth[32-34]. These findings

are consistent with reduced proliferation index seen in *Hdac3^{tko}* hearts (Figure 1B),

suggesting myocardial growth retardation account for the hypotrabeculation phenotypes

in *Hdac3^{tko}* hearts.

Endothelial HDAC3 induces ECM gene expression.

Among all scRNA-seq samples, 7 of them carried *R26$^{eYFP/+}$* reporters (3 CTL and 4

Hdac3^{tko}). This enabled us to perform a lineage specific analysis. In the single cell

suspension, 8.6% cells in CTL hearts and 11.2% cells in *Hdac3^{tko}* hearts were YFP+

(Supplemental Figure 8A). As expected, the expression of *Hdac3* in YFP+ populations in

Hdac3^tko hearts was significantly decreased compared to YFP+ populations in CTL hearts

(Supplemental Figure 8B). During early heart development, endothelial progenitors can

also give rise to fibroblasts/mesenchymal cells through endothelial to mesenchymal

transition[6, 35]. As expected, YFP+ cells (Tie2 lineage) were primarily CECs with a minority

of fibroblasts (Figure 2E). Also, as expected, *Hdac3* expression was absent in *Hdac3^tko*

YFP+ cells. Subsequent DGE analysis did not reveal significant gene expression

differences between *Hdac3^tko* and control fibroblasts (Figure 2E). Instead, in CEC clusters,

we found that the expression of genes regulating cellular component organization was

highly enriched in the control YFP+ CEC cluster (Figure 2F). In particular, the expression

of ECM associated genes, including *Col13a1, Col1a2, Tgfbi, Timp2, Col26a1, Col3a1,*

Col4a5, Timp3, and *P4ha3*, was significantly reduced in *Hdac3^tko* YFP+ CEC cluster as

compared to control YFP+ CEC cluster (Figure 2G). *Col1a1, Col1a2, Col2a1*, and *Col3a1*

and *P4ha3* DGE was further validated in *Hdac3^tko* and control heart lysates by qRT-PCR

(Figure 2H). These results suggest that cardiac endothelial HDAC3 induces the

expression of ECM associated genes, which are critical for CM proliferation and

trabeculation growth (Figure 2I).

Cardiac endothelial HDAC3 induces *Tgfβ3* in driving CM proliferation.

Trabeculation and ECM remodeling are coordinated during early heart development, and

both processes are regulated by growth signaling[36]. To better identify potential growth

signaling pathways regulated by HDAC3 in the endocardium, we used a mouse cardiac

endothelial cell (MCEC) line[37]. Using CRISPR-Cas9 gene editing technology and the

same set of gRNAs we have recently used[26], we successfully knocked out *Hdac3* in the

MCECs (Figure 3A). *Hdac3* KO MCECs grew slower than *Hdac3* empty vector (EV)

control MCECs. To determine whether the secretome from *Hdac3* KO endothelial cells is

competent to induce CM proliferation, we treated cultured primary embryonic CMs

isolated from E13.5 *Tnnt2^{nGFP/+38}* hearts with supernatant from either *Hdac3* KO MCECs

or *Hdac3* EV MCECs, respectively. *Hdac3* KO supernatant treatment resulted in

significant decrease of the percentage of Ki67+ CMs and the total number of CMs, as

compared to *Hdac3* EV supernatant treatment (Figure 3B). These results suggest that

growth signals derived from CECs required to support CM proliferation may be

inadequate/deficient in *Hdac3* KO MCECs. To unbiasedly search for such growth signals,

we performed bulk RNA-Seq analyses on *Hdac3* KO and EV MCECs; we identified 484

downregulated and 239 upregulated genes. Gene ontology pathway analyses revealed

that a number of endocardium-initiated signaling pathways were significantly

downregulated in *Hdac3* KO MCECs (Supplemental Figure 9A). Through DGE analysis,

we found that growth factors including *Igf1, Tgfβ3, Tgfβ2, Tgfβ1, Igf1, Cxcl12, Nrg1 and*

Vegfβ were significantly downregulated in *Hdac3* KO MCECs (Supplemental Figure 9B).

We attempted to validate these results at the protein level by western blot. Surprisingly,

we found that only TGFβ3 protein expression was significantly decreased in *Hdac3* KO

MCECs as compared to *Hdac3* EV MCECs (Supplemental Figure 9C), and thus we chose

TGFβ3 to pursue further. Interestingly, TGFβ3 was also significantly decreased when the

deacetylase activity of HDAC3 was inhibited by RGFP966 (a HDAC3 selective inhibitor)

treatment (Figure 3C and Supplemental Figure 9D). To assess *Tgfβ3* expression in E11.5

CECs, we sorted Tie2-derived population from *Hdac3^{tko}* (*Hdac3^{f/f}; Tie2-Cre/+; R26^{eYFP/+}*)

and CTL (*Hdac3^{f/+}; Tie2-Cre/+; R26^{eYFP/+}*) hearts by performing fluorescence activated

cell sorting on YFP signal (Supplemental Figure 10). We found that *Tgfβ3* mRNA level

was also significantly downregulated in E11.5 *Hdac3* KO CECs compared to littermate

CTL CECs (Figure 3D). TGFβ3 downregulation in E11.5 *Hdac3^{tko}* hearts was further

validated at the protein level by western blot (Figure 3E). Next, we determined whether

TGFβ3 deficiency in *Hdac3* KO MCECs contributed to the CM proliferation deficit

following *Hdac3* KO supernatant treatment. First, we quantified TGFβ3 and found that it

was significantly decreased in the supernatants from *Hdac3* KO MCECs compared to that

from *Hdac3* EV MCECs (Figure 3F). To determine whether TGFβ3 is an essential

secretory component of MCEC supernatants in stimulating CM proliferation, we

generated *Tgfβ3* KO MCECs using CRISPR-Cas9 gene editing technology

(Supplemental Figure 11A). We found that *Tgfβ3* KO significantly decreased the capacity

of MCEC supernatants to induce CM proliferation (percentage of Ki67+GFP+/GFP+ CMs)

compared to CTL MCEC supernatant (Supplemental Figure 11B). Further, we found that

Tgfβ3 KO significantly compromised the capacity of MCEC supernatants in supporting

myocardial trabeculation and compaction when treating *ex vivo* cultured E11.5 wildtype

hearts with *Tgfβ3* KO MCEC supernatants as compared to control MCEC supernatants

(Supplemental Figure 11C). These results altogether suggest that endocardial TGFβ3 is

critical for myocardial growth. Next, we evaluated CM proliferation after we supplemented

Hdac3 KO supernatants with recombinant mouse TGFβ3 protein. Strikingly, TGFβ3

supplementation successfully rescued CM proliferation defects by *Hdac3* KO supernatant

treatment (Figure 3G). Further, we found that TGFβ3 supplementation also rescued

myocardial growth deficit in *ex vivo* cultured E11.5 *Hdac3^{tko}* hearts (Supplemental Figure

12). These results suggest that cardiac endothelial HDAC3 supports myocardial growth

by inducing *Tgfβ3* expression.

Cardiac endothelial HDAC3 induces *Tgfβ3* expression through repressing

microRNA (miR)-129-5p.

Recently, we reported that HDAC3 regulates growth factors by modulating the miR

pathway in the developing epicardium[26]. To identify the potential HDAC3 downstream

miR targets in CECs, we performed miR sequencing in *Hdac3* KO and EV MCECs.

Through DGE analyses, we found that 5 miRs were significantly upregulated in *Hdac3*

KO MCECs (Figure 4A) (cutoff threshold: Log2 fold change greater than 1.5 and adjusted

P<0.01). To identify which miR(s) may regulate *Tgfβ3* expression, we treated MCECs

with mimics of each of the 5 miRs and found that treatment with miR-129-5p mimics

significantly inhibited the expression of *Tgfβ3* (Figure 4B). This finding was validated by

western blot (Figure 4C). We further validated the upregulation of miR-129-5p in both

Hdac3 KO MCECs and E11.5 *Hdac3^{tko}* hearts (Figure 4D and 4E). Interestingly, miR-

129-5p was similarly significantly upregulated when the deacetylase activity of HDAC3 is

inhibited by RGFP966 treatment (Figure 4D), suggesting that the inhibition of miR-129-

5p expression by HDAC3 is deacetylase dependent. To determine whether miR-129-5p

upregulation in MCECs has potential impact on myocardial growth, we treated cultured

Tnnt2^{nGFP/+38} CMs or cultured whole E11.5 wildtype hearts with supernatants from

MCECs transfected with either miR-129-5p mimics or scramble RNAs. We found that

miR-129-5p supernatant treatment significantly decreased its capacity to induce CM

proliferation and myocardial growth (Supplemental Figure 13). Lastly, to determine

whether HDAC3 induces *Tgfβ3* expression through repressing miR-129-5p, we knocked

down miR-129-5p by miRZip lentivirus in *Hdac3* KO MCECs. Remarkably, miR-129-5p

knockdown significantly restored the TGFβ3 expression in *Hdac3* KO MCECs (Figure 4G).

Further, the supernatants from miR-129-5p knockdown *Hdac3* KO MCECs significantly

restored the CM proliferation promoting capacity from *Hdac3* KO MCEC supernatants

(Figure 4H). These results suggest that HDAC3 promotes the expression of TGFβ3

through repressing miR-129 in supporting trabecular myocardial growth (Figure 4I).

Discussion

Myocardial growth/expansion, one of the most critical cardiac developmental events, is accompanied by two separate but closely associated processes: compaction and trabeculation. Many forms of congenital heart disease are associated with abnormal cardiac trabeculation and compaction[2, 3]. For proper myocardial development, besides cardiomyocytes, the surrounding non-cardiomyocytes including endocardial cells can influence myocardial growth by sending growth signals to the neighboring myocardia[39], as well as secreting ECM proteins that serve as signal transmission media[2, 40]

Epigenetic regulation can define a tissue-specific and developmental stage-specific transcription network during early heart development[41, 42], mainly achieved by posttranslational modifications of histones, around which DNA winds. There are several different types of histone modifications (e.g., methylation/demethylation, acetylation/deacetylation, phosphorylation/dephosphorylation, ubiquitination/ deubiquitination). By alternating between any of these states, the associated genes can be dynamically poised to be transcriptionally active/repressed. Histone deacetylases (HDACs) are family of enzymes that remove acetyl groups from a lysine residue on a histone, allowing the histones to wrap the DNA more tightly to repress gene expression. Class I HDACs, includes HDAC3, are widely expressed, but play specific roles during early embryogenesis and organogenesis[19]. Mesodermal or global knockout of *Hdac3* results in myogenic defects and early embryonic lethality[22, 43]. Interestingly, specific deletion of *Hdac3* in the myocardium does not generate cardiac morphogenic phenotypes during early heart development, but rather compromises cardiac function at later postnatal stages[43, 44]. These early findings suggest that the function of HDAC3 in

nonmyocyte compartments may be critical for early cardiac morphogenesis and

myocardial development. Our recent work demonstrated that epicardial HDAC3 is critical

for driving myocardial compact wall expansion[26]. Our current study suggests that

endocardial HDAC3 is important for myocardial trabeculation: endothelial/endocardial

deletion of *Hdac3* resulted in hypotrabeculation (Figure 1 and Supplemental Figure 4).

We note that *Tie2-Cre* turns on around E8.0-8.5 in all endothelial cells[29], whereas

Nfatc1[Cre/+] is active around E9.5 only in endocardial cells[30] and only achieves efficient

Hdac3 deletion by E14.5 likely due to inefficient IRES-mediated Cre transcription[45]

(Supplemental Figure 4). Nonetheless, these two deletion models generated similar

cardiac phenotypes of hypotrabeculation, although we did observe differences in the

severity of the myocardial phenotypes and time of lethality, which we attribute to the

different temporal and spatial activities and/or efficiencies of these two Cres. We also

observed that both E12.5 *Hdac3[tko]* and E14.5 *Hdac3[nko]* hearts exhibited thin compact wall

phenotype. This is likely attributable to *Hdac3* deficiency in the subepicardial endothelial

population which form coronary arteries invading and supporting compact wall[46]. This late

thin compact wall phenotype is also consistent with the similar phenotype resulted from

epicardial deletion of *Hdac3*[26]. An earlier report of conditional deletion of *Hdac3* in

endothelial cells using a different Tie2-Cre driver primarily targeting hemogenic

endothelial cells[27] resulted in mid-gestational lethality (E14.5) presenting predominantly

a lymphovenous and lymphatic valve malformation phenotype[47]. However, it is unknown

what phenotypes were present in the myocardium in that study as none were reported.

Nonetheless, we deemed it inapplicable to evaluate potential lymphatic valve phenotypes

in our pan-endothelial *Hdac3* KO embryos since these mice died around E12.5

(Supplemental Table 1), while lymphovenous and lymphatic valve formation starts at

E13.5[48]. However, it is interesting to note that the role of HDAC1 and HDAC2 in the

developing endocardium appears to be dispensable, as deletion of either *Hdac1* or *Hdac2*

by Tie2-Cre do not elicit apparent cardiac phenotypes[24]. These results suggest that

HDAC3 plays distinctive roles from HDAC1 and HDAC2, although they share high

homology and are grouped in the same HDAC subfamily[19].

TGFß signaling is involved in many processes during early heart development such as

ECM production, endothelial to mesenchymal transition, and myogenesis[11, 49-51].

Specifically, TGFß signaling plays a critical role during early cardiac myocardial

development[52-55]. For instance, knockout of either TGFß2 or TGFß3 results in hypoplastic

ventricular myocardium[15], while TGFß1 is reported to inhibit CM proliferation[55, 56],

suggesting the complexity and specificity of TGFß isoforms during early heart

development. In the current study, we found that TGFß3 expression was significantly

decreased in *Hdac3* deleted endocardium, and TGFß3 treatment vigorously stimulated

CM proliferation (Figure 3). Interestingly, inhibition of deacetylase activity of HDAC3 alone

significantly decreased TGFß3 expression. Further, supplementation of TGFß3 to cardiac

endothelial Hdac3 KO secretome significantly rescued CM proliferation and myocardial

growth defect (Figure 3), suggesting that cardiac endothelial HDAC3 supports

trabeculation through inducing TGFß3 in a deacetylase dependent manner. Interestingly,

HDAC3 is also reported to inhibit TGFß1 expression in the second heart field

development in a deacetylase independent manner[23], whereas HDAC3 deteriorates

colorectal cancer progression by inducing TGFß1 through inhibiting miR-296[57]. These

divergent findings illustrate that HDAC3 can either promote or inhibit TGFß signaling

pathway through different molecular mechanisms in a tissue/cell dependent manner.

Besides functioning as growth signals to cardiomyocytes, TGFß signaling is

predominantly known for its induction on ECMs[50, 58]. During early heart development,

ECM is mainly secreted from endocardium and mediates signal transduction from

endocardium to myocardium supporting trabeculation[57]. Our scRNA-Seq data revealed

that ECM is significantly down regulated in *Hdac3* endothelial KO hearts (Figure 2).

P4ha3, a major enzyme for collagen synthesis[59] and a TGFß downstream target[60], was

also significantly decreased in *Hdac3* endothelial knockout hearts (Figure 2). These

results suggest that ECM downregulation may be secondary to decreased TGFß

signaling to contribute to the hypotrabeculation phenotypes in *Hdac3*

endothelial/endocardial knockout hearts.

We identified that HDAC3 suppresses miR-129-5p in inducing TGFß3 (Figure 4). miR-

129-5p plays important roles in cancer by either promoting or inhibiting tumorigenesis

through regulating a variety of downstream targets and pathways[61, 62]. Several studies

have shown that miR-129-5p protects adult hearts from ischemia/reperfusion injuries[63,

64], while another study demonstrated that treating cultured H9c2 cardiomyocytes with

HDAC pan-inhibitor Trichostatin A results in decreased cardiomyocyte proliferation with

enhanced miR-129-5p expression[65]. This is consistent with our current findings in the

developing endocardium (Figure 4). In our recent study, we found that HDAC3 induces

the expression of IGF2 and FGF9 by suppressing miR-322/503 in the developing

epicardium[26]. Again, these findings altogether suggest that HDAC3 can induce different

sets of growth signals by switching their downstream miR targets in a tissue/cell context

dependent manner.

In summary, our current study demonstrates that endocardial HDAC3 promotes

myocardial trabeculation by inducing TGFß signaling through repressing miR-129-5p

during early heart development. This finding, along with others, exemplifies the

importance of endocardial signaling for driving myocardial morphogenesis, and thus

provides new insights on the pathogenesis of congenital heart disease. The dual

epigenetic inhibition mechanism may be leveraged in some other contexts such as adult

heart repair/regeneration.

Methods

Mice

$Hdac3^{flox/+}$, *Tie2-Cre*, $Nfatc1^{Cre/+}$, $R26^{eYFP}$ and $Tnnt2^{nGFP/+}$ mice were previously described[27, 30, 38, 66, 67]. B6.Cg-Tg(*Tek-cre*)12Flv/J (abbreviated as *Tie2-Cre* in this manuscript), $Hdac3^{flox/+}$, and $R26^{eYFP}$ mice are available at the Jackson Laboratory (Stock numbers: #004128, #024119 and #006148, respectively). All animal protocols were approved by the Nationwide Children's Institutional Animal Care and Use Committee (IACUC #AR22-00194).

Histology and immunofluorescence staining

All embryo specimens were fixed in 4% paraformaldehyde overnight, dehydrated through an ethanol series, paraffin embedded, and sectioned (6-7µm). Primary antibodies (Major Resources Table) were incubated at 4°C overnight and secondary antibodies (Alexa 488 or Alexa 555, Thermo Fisher Scientific, Waltham, MA) were incubated at room temperature for one hour. Stained slides were imaged on a Zeiss LSM 710 confocal Microscope or a Leica DM6 fluorescence microscope. Epifluorescence and gross images of embryos were imaged on a Leica M205 FCA stereo fluorescence microscope. The number of phosphor-Histone 3 (p-H3)+ cells, Bromodeoxyuridine (BrdU)+ cells, and TUNEL+ cells was quantified using ImageJ software.

Embryonic cardiomyocyte culture and proliferation assessment

E13.5 ventricular cardiomyocytes were isolated from dissected embryonic hearts and cultured as previously described[68]. To assess cell proliferation, cardiomyocytes were

fixed with 4% paraformaldehyde and stained with Ki67 or p-H3 antibody followed by Alexa

594-conjugated secondary antibody (Thermo Fisher Scientific, Waltham, MA). The

immunostaining was visualized and imaged on a Keyence BZ-X800 microscope. The

percentages of Ki67+GFP+/GFP+, p-H3+GFP+/GFP+ and total number of GFP+

cardiomyocytes were quantified using ImageJ software.

Endothelial cell culture, transient transfection, lentiviral infection, and luciferase

assay

Mouse cardiac endothelial cells (MCECs) were purchased from Cedarlane Labs

(Catalogue #CLU510, Burlington, NC). MCECs were cultured in 10% fetal bovine serum

(FBS) supplemented DMEM at 37°C in a humidified incubator with 5% CO_2. To generate

stable *Hdac3* KO MCECs, as previously described[26], we cloned *Hdac3* sgRNA into

lentiCRISPR v2 vector (Addgene #52961; Watertown, MA). Seventy-percent confluent

Lenti-X™ 293T cells (Takara Bio USA, Inc., Mountain View, CA) were transfected with

Hdac3 lentiviral plasmids (*Hdac3* KO or empty vector control [EV])[26], *Tgfβ3* lentiviral

plasmids (*Tgfβ3* KO or empty vector control [EV]) were purchased from VectorBuilder,

PsPAX2 (Addgene #12260) and PMD2G (Addgene #12259). Seventy-two hours after

transfection, supernatants containing lentivirus were collected and filtered through a

40μm cell strainer. Mir-129-5p, miR-129b-3p, miR-129b-5p, mir-129-2-3p, miR-708-3p

mimics and scramble control mimics were custom-ordered from ThermoFischer Scientific

(Waltham, MA). MCECs were transfected with miRNA mimics or control mimics (30 pmol)

using Lipofectamine RNAiMAX (Thermo Fisher Scientific, Waltham, MA). 72 hours after

miR mimics treatment, supernatants were collected. miRZip anti-miR precursor

constructs for miRZip-129-5p and pGreenPuro Scramble Hairpin control constructs were

purchased from System Biosciences (Palo Alto, CA), and lentiviruses were generated

according to the manufacturer's protocol.

Ex Vivo embryonic heart culture

E11.5 embryonic hearts were dissected in cold PBS, and then cultured in DMEM/F-12

media (Thermofisher Scientific, Waltham, MA, #11039021). Embryonic cardiac explants

were treated with either TGFß3 recombinant protein (Sigmal Aldrich, Saint Louis, MO,

#SRP6552, final concentration: 250 ng/ml), TGFß3 KO MCEC supernatants, miR-129-5p

mimics-treated MCEC supernatants and their corresponding control media or control

MCEC supernatants for 24 h at 37 °C. Hearts were then fixed in 4% paraformaldehyde

overnight at 4 °C and processed for paraffin embedding and sectioning.

Fluorescence activated cell sorting of YFP+ Tie2 derivative cardiac cells

E11.5 hearts were dissected and dissociated with 0.5% trypsin–EDTA into a single cell

suspension. Cell suspension was washed with PBS and YFP+ cells were gated using

their autofluoresence signal and sorted and using a BD Influx cell sorter (BD Biosciences).

The harvested YFP+ cells were used for subsequent gene expression analysis.

RNA isolation, quantitative real-time PCR (qRT-PCR), and bulk RNA-Seq

MCECs and embryonic day (E) 11.5 hearts were used for RNA isolation. E11.5 hearts

were microdissected in cold PBS and snap frozen in liquid nitrogen. The RNeasy Plus

Mini Kit (QIAGEN, Germantown, MD) was used to extract total RNA, and cDNAs were

generated using the Superscript III kit (Thermo Fisher Scientific, Waltham, MA). To detect

mRNA, SYBR Green qRT-PCR was performed. PCR primers for genes of interest are

listed in Major Resources Table. To perform mature microRNA expression analyses,

purified total RNAs were converted to cDNAs using TaqMan™ MicroRNA Reverse

Transcription Kit (Catalogue #4366596, Thermo Fisher Scientific). qRT-PCR reactions

were run on StepOne Plus Real-Time PCR System (Applied Biosystems, Waltham, MA).

The probe sequences for mmu-miR-129-5p and U6 snRNA were purchased from Thermo

Fisher Scientific. For bulk RNA and miRNA-Sequencing, samples were prepared

following the provider's guidelines (Novogene Corporation Inc, Sacramento, CA) and

sequenced on an Illumina NextSeq500 (San Diego, CA). Sequencing reads were aligned

to the UCSC mm10 reference genome using tophat2 and bowtie2 in R. Differential

expression of transcripts was calculated using the cufflinks suite in R analyses.

Single cell RNA-Sequencing (scRNA-seq)

E11.5 heart tissues were dissected in cold PBS and digested with 0.25% trypsin at 37°C

for 15 minutes. Digested cardiac cells were filtered with 70 μm cell strainers to remove

remaining clumps. Subsequently, 10,000 live cells were captured with Chromium Next

GEM Single Cell 3' Reagent Kits v3.1 (10X Genomics, Pleasanton, CA) following

manufacturer's instructions. Briefly, single cells were partitioned into nanoliter-scale Gel

Beads-in-emulsion (GEMs) using Chromium Next GEM Chip G in the chromium controller.

Immediately following GEM generation, the Gel Bead was dissolved, primers were

released, and co-partitioned cell was lysed, and its mRNA were reverse transcribed into

barcoded cDNA. After further cleanup and amplification, the cDNA was enzymatically

fragmented and amplified via PCR to generate sufficient mass for library construction.

After end repair, A-tailing, adaptor ligation and sample index PCR, the library comprised

constructs containing the sample index, UMI sequences, barcode sequences, and

Illumina standard sequencing primers P5 and P7 at both ends. The library was sequenced

with Illumina NextSeq S4 platform to a sequencing depth of 140-270 million reads per

sample. Reads were aligned to the Ensembl 104 mouse reference genome including the

sequence for eYFP as a separate contig using STARsolo (version 2.7.10b)[69]. The

resulting count tables (cells x genes) were converted into Annotated Data (anndata)

format for analysis in Python. For quality control, all cells with less than 1500 or more than

9000 genes expressed were excluded. Additionally, cell doublets were filtered out using

the Scrublet method[70]. Gene counts were normalized per cell to a total target sum and

transformed using log1p. Highly variable genes were identified using the Seurat method[71]

and these genes were utilized for dimension reduction with PCA. Finally, cell embedding

was performed with the UMAP algorithm (umap-learn version 0.5.3) and cells were

clustered using leiden (leidenalg version 0.9.1) with resolution 0.2. Scoring of trabecular

and compact CM subtypes was performed using 'score_genes' and differentially

expressed genes between groups were obtained using 'rank_genes_groups', both

methods from the SCANPY package[72].

Western blotting

Cell or tissue lysates were prepared in lysis buffer (20 mM Tris-HCl [pH 7.5], 15 mM NaCl,

1 mM Na_2EDTA, 1 mM EGTA, 1% Triton X-100, 1 µg/ml leupeptin, 2.5 mM sodium

pyrophosphate, 1 mM Na_3VO_4, and 1 mM β-glycerophosphate) with protease inhibitor

cocktail (Roche, Indianapolis, IN) and 1 mM phenylmethylsulfonyl fluoride. Protein

samples were resolved on 4–12% SDS-PAGE acrylamide gels before transferring to

polyvinylidene fluoride membranes. Primary antibodies were visualized by

chemiluminescence using HRP-conjugated secondary antibodies. Antibodies are listed

in Major Resources Table. Western blots were normalized to loading controls and

densitometric analysis was performed using Image J software.

Statistical analysis

All experiments were independently repeated at least three times, and the number of

samples (n) is stated in figure legends. Results are reported as the mean ± SD. For

statistical analyses, normal distribution of data was assessed by Shapiro-Wilk's test

(when n≥6). For data with small sample size (n<6), non-parametric tests were applied.

For parametric data with normal distribution, the statistical significance of the difference

between means was assessed using the unpaired two tailed Student's t-test between two

groups and the Tukey or Dunnett post hoc test for one-way ANOVA. For non-parametric

data or data with small sample size (n<6), the Mann-Whitney U test and Kruskal-Wallis

test for one-way ANOVA followed by the Dunn post hoc test were performed for

comparisons between two groups. Specific statistical tests are stated in figure legends.

RNA-seq data were background-corrected, variance-stabilized, normalized, and count

matrix extracted through quality control using lumi package for the R programming

language (http://bioconductor.org/packages/release/bioc/html/lumi.html). Differential

expression analysis was performed using DESeq2 R package

(https://bioconductor.org/packages/release/bioc/html/DESeq2. html). The resulting *p*-

values were adjusted using the Benjamini and Hochberg's approach for controlling the

False Discovery Rate (FDR)[73]. Genes or miRs with an adjusted p-value<0.05 found by

DESeq2 were assigned as differentially expressed. Gene ontology (GO) enrichment

analyses were performed using the Database for Annotation, Visualization and Integrated

Discovery (DAVID) bioinformatics resources (https://david.ncifcrf.gov/). For relative gene

expression analysis of qRT-PCR data, the expression of a gene of interest relative to an

internal loading control gene was calculated using the comparative CT method ($-\Delta\Delta CT$)[74].

The relative protein expression on western blots was quantified by densitometry, Fold

changes for relative gene or protein expression were derived through division of values

in the experimental group by the mean values from the control group. P-value<0.05 was

considered statistically significant. All statistical analyses were performed using

GraphPad Prism 8 (GraphPad Prism software, Inc). The images with the value closest to

the mean value were selected as representative images.

Data Availability

All data associated with this study are present in the main text or the supplementary

materials. The scRNA-seq, RNA-seq and miRNA-seq data generated in this study have

been deposited in the GEO database under accession code GSE229661. All raw data

are available from the authors upon reasonable request.

References:

1. Zaidi S and Brueckner M. Genetics and Genomics of Congenital Heart Disease. *Circulation research*. 2017;120:923-940.

2. Qu X, Harmelink C and Baldwin HS. Endocardial-Myocardial Interactions During Early Cardiac Differentiation and Trabeculation. *Front Cardiovasc Med*. 2022;9:857581.

3. Wu M. Mechanisms of Trabecular Formation and Specification During Cardiogenesis. *Pediatric cardiology*. 2018;39:1082-1089.

4. Lin W and Li D. Zinc and Zinc Transporters: Novel Regulators of Ventricular Myocardial Development. *Pediatric cardiology*. 2018;39:1042-1051.

5. Zhang W, Chen H, Qu X, Chang CP and Shou W. Molecular mechanism of ventricular trabeculation/compaction and the pathogenesis of the left ventricular noncompaction cardiomyopathy (LVNC). *American journal of medical genetics Part C, Seminars in medical genetics*. 2013;163C:144-56.

6. Zhang H, Lui KO and Zhou B. Endocardial Cell Plasticity in Cardiac Development, Diseases and Regeneration. *Circulation research*. 2018;122:774-789.

7. Andries LJ, Sys SU and Brutsaert DL. Morphoregulatory interactions of endocardial endothelium and extracellular material in the heart. *Herz*. 1995;20:135-45.

8. Camenisch TD, Spicer AP, Brehm-Gibson T, Biesterfeldt J, Augustine ML, Calabro A, Jr., Kubalak S, Klewer SE and McDonald JA. Disruption of hyaluronan synthase-2 abrogates normal cardiac morphogenesis and hyaluronan-mediated transformation of epithelium to mesenchyme. *J Clin Invest*. 2000;106:349-60.

9. Hatano S, Kimata K, Hiraiwa N, Kusakabe M, Isogai Z, Adachi E, Shinomura T and Watanabe H. Versican/PG-M is essential for ventricular septal formation subsequent to cardiac atrioventricular cushion development. *Glycobiology*. 2012;22:1268-77.

10. Chen JL, Colgan TD, Walton KL, Gregorevic P and Harrison CA. The TGF-beta Signalling Network in Muscle Development, Adaptation and Disease. *Adv Exp Med Biol*. 2016;900:97-131.

11. Sorensen DW and van Berlo JH. The Role of TGF-beta Signaling in Cardiomyocyte

Proliferation. *Curr Heart Fail Rep*. 2020;17:225-233.

12. Zhang Y and Yang X. The Roles of TGF-beta Signaling in Cerebrovascular Diseases.

Front Cell Dev Biol. 2020;8:567682.

13. Moustakas A and Heldin CH. Non-Smad TGF-beta signals. *J Cell Sci*. 2005;118:3573-84.

14. Zhang YE. Non-Smad pathways in TGF-beta signaling. *Cell Res*. 2009;19:128-39.

15. Azhar M, Schultz Jel J, Grupp I, Dorn GW, 2nd, Meneton P, Molin DG, Gittenberger-de

Groot AC and Doetschman T. Transforming growth factor beta in cardiovascular development

and function. *Cytokine Growth Factor Rev*. 2003;14:391-407.

16. Koo SH, Cunningham MC, Arabshahi B, Gruss JS and Grant JH, 3rd. The transforming

growth factor-beta 3 knock-out mouse: an animal model for cleft palate. *Plast Reconstr Surg*.

2001;108:938-48; discussion 949-51.

17. Sanford LP, Ormsby I, Gittenberger-de Groot AC, Sariola H, Friedman R, Boivin GP,

Cardell EL and Doetschman T. TGFbeta2 knockout mice have multiple developmental defects

that are non-overlapping with other TGFbeta knockout phenotypes. *Development*.

1997;124:2659-70.

18. Kulkarni AB and Karlsson S. Transforming growth factor-beta 1 knockout mice. A mutation

in one cytokine gene causes a dramatic inflammatory disease. *Am J Pathol*. 1993;143:3-9.

19. Haberland M, Montgomery RL and Olson EN. The many roles of histone deacetylases in

development and physiology: implications for disease and therapy. *Nature reviews Genetics*.

2009;10:32-42.

20. Dsilva P, Pai P, Shetty MG and Babitha KS. The role of histone deacetylases in embryonic

development. *Mol Reprod Dev*. 2023;90:14-26.

21. Wang Z, Zhao YT and Zhao TC. Histone deacetylases in modulating cardiac disease and

their clinical translational and therapeutic implications. *Exp Biol Med (Maywood)*. 2021;246:213-

225.

22.   Bhaskara S, Chyla BJ, Amann JM, Knutson SK, Cortez D, Sun ZW and Hiebert SW.

Deletion of histone deacetylase 3 reveals critical roles in S phase progression and DNA damage

control. *Mol Cell*. 2008;30:61-72.

23.   Lewandowski SL, Janardhan HP and Trivedi CM. Histone Deacetylase 3 Coordinates

Deacetylase-independent Epigenetic Silencing of Transforming Growth Factor-beta1 (TGF-

beta1) to Orchestrate Second Heart Field Development. *The Journal of biological chemistry*.

2015;290:27067-27089.

24.   Janardhan HP, Milstone ZJ, Shin M, Lawson ND, Keaney JF, Jr. and Trivedi CM. Hdac3

regulates lymphovenous and lymphatic valve formation. *J Clin Invest*. 2017;127:4193-4206.

25.   Poleshko A, Shah PP, Gupta M, Babu A, Morley MP, Manderfield LJ, Ifkovits JL, Calderon

D, Aghajanian H, Sierra-Pagan JE, Sun Z, Wang Q, Li L, Dubois NC, Morrisey EE, Lazar MA,

Smith CL, Epstein JA and Jain R. Genome-Nuclear Lamina Interactions Regulate Cardiac Stem

Cell Lineage Restriction. *Cell*. 2017;171:573-587 e14.

26.   Jang J, Song G, Pettit SM, Li Q, Song X, Cai CL, Kaushal S and Li D. Epicardial HDAC3

Promotes Myocardial Growth Through a Novel MicroRNA Pathway. *Circulation research*.

2022;131:151-164.

27.   Koni PA, Joshi SK, Temann UA, Olson D, Burkly L and Flavell RA. Conditional vascular

cell adhesion molecule 1 deletion in mice: impaired lymphocyte migration to bone marrow. *The*

Journal of experimental medicine. 2001;193:741-54.

28.   de Lange WJ, Halabi CM, Beyer AM and Sigmund CD. Germ line activation of the Tie2

and SMMHC promoters causes noncell-specific deletion of floxed alleles. *Physiol Genomics*.

2008;35:1-4.

29.   Kisanuki YY, Hammer RE, Miyazaki J, Williams SC, Richardson JA and Yanagisawa M.

Tie2-Cre transgenic mice: a new model for endothelial cell-lineage analysis in vivo. *Dev Biol*.

2001;230:230-42.

30.   Wu B, Zhang Z, Lui W, Chen X, Wang Y, Chamberlain AA, Moreno-Rodriguez RA,

Markwald RR, O'Rourke BP, Sharp DJ, Zheng D, Lenz J, Baldwin HS, Chang CP and Zhou B.

Endocardial cells form the coronary arteries by angiogenesis through myocardial-endocardial

VEGF signaling. *Cell*. 2012;151:1083-96.

31.   Wu T, Liang Z, Zhang Z, Liu C, Zhang L, Gu Y, Peterson KL, Evans SM, Fu XD and Chen

J. PRDM16 Is a Compact Myocardium-Enriched Transcription Factor Required to Maintain

Compact Myocardial Cardiomyocyte Identity in Left Ventricle. *Circulation*. 2022;145:586-602.

32.   Hassan MK, Kumar D, Patel SA and Dixit M. EEF1A2 triggers stronger ERK mediated

metastatic program in ER negative breast cancer cells than in ER positive cells. *Life Sci*.

2020;262:118553.

33.   Fish JE, Cantu Gutierrez M, Dang LT, Khyzha N, Chen Z, Veitch S, Cheng HS, Khor M,

Antounians L, Njock MS, Boudreau E, Herman AM, Rhyner AM, Ruiz OE, Eisenhoffer GT,

Medina-Rivera A, Wilson MD and Wythe JD. Dynamic regulation of VEGF-inducible genes by an

ERK/ERG/p300 transcriptional network. *Development*. 2017;144:2428-2444.

34.   Thornton S, Anand N, Purcell D and Lee J. Not just for housekeeping: protein initiation

and elongation factors in cell growth and tumorigenesis. *J Mol Med (Berl)*. 2003;81:536-48.

35.   Anbara T, Sharifi M and Aboutaleb N. Endothelial to Mesenchymal Transition in the

Cardiogenesis and Cardiovascular Diseases. *Curr Cardiol Rev*. 2020;16:306-314.

36.   Del Monte-Nieto G, Ramialison M, Adam AAS, Wu B, Aharonov A, D'Uva G, Bourke LM,

Pitulescu ME, Chen H, de la Pompa JL, Shou W, Adams RH, Harten SK, Tzahor E, Zhou B and

Harvey RP. Control of cardiac jelly dynamics by NOTCH1 and NRG1 defines the building plan for

trabeculation. *Nature*. 2018;557:439-445.

37.   Barbieri SS and Weksler BB. Tobacco smoke cooperates with interleukin-1beta to alter

beta-catenin trafficking in vascular endothelium resulting in increased permeability and induction

of cyclooxygenase-2 expression in vitro and in vivo. *FASEB J*. 2007;21:1831-43.

38.   Yan J, Zhang L, Sultana N, Oh JG, Wu B, Hajjar RJ, Zhou B and Cai CL. A series of robust

genetic indicators for definitive identification of cardiomyocytes. *J Mol Cell Cardiol*. 2016;97:278-

85.

39.   Tian Y and Morrisey EE. Importance of myocyte-nonmyocyte interactions in cardiac

development and disease. *Circulation research*. 2012;110:1023-34.

40.   Silva AC, Pereira C, Fonseca A, Pinto-do OP and Nascimento DS. Bearing My Heart: The

Role of Extracellular Matrix on Cardiac Development, Homeostasis, and Injury Response. *Front*

Cell Dev Biol. 2020;8:621644.

41.   Paige SL, Thomas S, Stoick-Cooper CL, Wang H, Maves L, Sandstrom R, Pabon L,

Reinecke H, Pratt G, Keller G, Moon RT, Stamatoyannopoulos J and Murry CE. A temporal

chromatin signature in human embryonic stem cells identifies regulators of cardiac development.

Cell. 2012;151:221-32.

42.   Wamstad JA, Alexander JM, Truty RM, Shrikumar A, Li F, Eilertson KE, Ding H, Wylie JN,

Pico AR, Capra JA, Erwin G, Kattman SJ, Keller GM, Srivastava D, Levine SS, Pollard KS,

Holloway AK, Boyer LA and Bruneau BG. Dynamic and coordinated epigenetic regulation of

developmental transitions in the cardiac lineage. *Cell*. 2012;151:206-20.

43.   Lewandowski SL, Janardhan HP and Trivedi CM. Histone Deacetylase 3 Coordinates

Deacetylase-independent Epigenetic Silencing of Transforming Growth Factor-beta1 (TGF-

beta1) to Orchestrate Second Heart Field Development. *The Journal of biological chemistry*.

2015;290:27067-89.

44.   Montgomery RL, Potthoff MJ, Haberland M, Qi X, Matsuzaki S, Humphries KM,

Richardson JA, Bassel-Duby R and Olson EN. Maintenance of cardiac energy metabolism by

histone deacetylase 3 in mice. *J Clin Invest*. 2008;118:3588-97.

45.   Artap S, Manderfield LJ, Smith CL, Poleshko A, Aghajanian H, See K, Li L, Jain R and

Epstein JA. Endocardial Hippo signaling regulates myocardial growth and cardiogenesis. *Dev*

Biol. 2018;440:22-30.

46.   Tian X, Hu T, Zhang H, He L, Huang X, Liu Q, Yu W, He L, Yang Z, Zhang Z, Zhong TP,

Yang X, Yang Z, Yan Y, Baldini A, Sun Y, Lu J, Schwartz RJ, Evans SM, Gittenberger-de Groot

AC, Red-Horse K and Zhou B. Subepicardial endothelial cells invade the embryonic ventricle wall

to form coronary arteries. *Cell Res*. 2013;23:1075-90.

47.   Celikyurt U, Karauzum K, Sahin T, Agacdiken A, Vural A and Ural D. Association between

resolution of fragmented QRS and response to cardiac resynchronization therapy. *Ann*

Noninvasive Electrocardiol. 2015;20:126-31.

48.   Yang Y and Oliver G. Development of the mammalian lymphatic vasculature. *J Clin Invest*.

2014;124:888-97.

49.   Conway SJ, Doetschman T and Azhar M. The inter-relationship of periostin, TGF beta,

and BMP in heart valve development and valvular heart diseases. *ScientificWorldJournal*.

2011;11:1509-24.

50.   Frangogiannis NG. Transforming growth factor-beta in myocardial disease. *Nat Rev*

Cardiol. 2022;19:435-455.

51.   Lee JH and Massague J. TGF-beta in developmental and fibrogenic EMTs. *Semin Cancer*

Biol. 2022;86:136-145.

52.   Chakrabarti M, Al-Sammarraie N, Gebere MG, Bhattacharya A, Chopra S, Johnson J,

Pena EA, Eberth JF, Poelmann RE, Gittenberger-de Groot AC and Azhar M. Transforming

Growth Factor Beta3 is Required for Cardiovascular Development. *J Cardiovasc Dev Dis*. 2020;7.

53.   Dunker N and Krieglstein K. Tgfbeta2 -/- Tgfbeta3 -/- double knockout mice display severe

midline fusion defects and early embryonic lethality. *Anat Embryol (Berl)*. 2002;206:73-83.

54.   Kruithof BP, Kruithof-De-Julio M, Poelmann RE, Gittenberger-De-Groot AC, Gaussin V

and Goumans MJ. Remodeling of the myocardium in early trabeculation and cardiac valve

formation; a role for TGFbeta2. *Int J Dev Biol*. 2013;57:853-63.

55.   Kodo K, Ong SG, Jahanbani F, Termglinchan V, Hirono K, InanlooRahatloo K, Ebert AD,

Shukla P, Abilez OJ, Churko JM, Karakikes I, Jung G, Ichida F, Wu SM, Snyder MP, Bernstein D

and Wu JC. iPSC-derived cardiomyocytes reveal abnormal TGF-beta signalling in left ventricular

non-compaction cardiomyopathy. *Nat Cell Biol*. 2016;18:1031-42.

56.   Engelmann GL, Boehm KD, Birchenall-Roberts MC and Ruscetti FW. Transforming

growth factor-beta 1 in heart development. *Mech Dev*. 1992;38:85-97.

57.   Li J, Hu M, Liu N, Li H, Yu Z, Yan Q, Zhou M, Wang Y, Song Y, Pan G, Liang F and Chen

R. HDAC3 deteriorates colorectal cancer progression via microRNA-296-3p/TGIF1/TGFbeta axis.

J Exp Clin Cancer Res. 2020;39:248.

58.   Algeciras L, Palanca A, Maestro D, RuizdelRio J and Villar AV. Epigenetic alterations of

TGFbeta and its main canonical signaling mediators in the context of cardiac fibrosis. *J Mol Cell*

Cardiol. 2021;159:38-47.

59.   Van Den Diepstraten C, Papay K, Bolender Z, Brown A and Pickering JG. Cloning of a

novel prolyl 4-hydroxylase subunit expressed in the fibrous cap of human atherosclerotic plaque.

Circulation. 2003;108:508-11.

60.   Luo Y, Xu W, Chen H, Warburton D, Dong R, Qian B, Selman M, Gauldie J, Kolb M and

Shi W. A novel profibrotic mechanism mediated by TGFbeta-stimulated collagen prolyl

hydroxylase expression in fibrotic lung mesenchymal cells. *J Pathol*. 2015;236:384-94.

61.   Deng B, Tang X and Wang Y. Role of microRNA-129 in cancer and non-cancerous

diseases (Review). *Exp Ther Med*. 2021;22:918.

62.   Xu S, Li W, Wu J, Lu Y, Xie M, Li Y, Zou J, Zeng T and Ling H. The Role of miR-129-5p

in Cancer: A Novel Therapeutic Target. *Curr Mol Pharmacol*. 2022;15:647-657.

63.   Ma R, Chen X, Ma Y, Bai G and Li DS. MiR-129-5p alleviates myocardial injury by

targeting suppressor of cytokine signaling 2 after ischemia/reperfusion. *Kaohsiung J Med Sci*.

2020;36:599-606.

64.   Xing J, Liu J, Liu J and Xu Z. miR-129-5p ameliorates ischemia-reperfusion injury by

targeting HMGB1 in myocardium. *Gen Physiol Biophys*. 2020;39:461-470.

65.   Majumdar G and Raghow R. Trichostatin A induces a unique set of microRNAs including

miR-129-5p that blocks cyclin-dependent kinase 6 expression and proliferation in H9c2 cardiac

myocytes. *Mol Cell Biochem*. 2016;415:39-49.

66.   Mullican SE, Gaddis CA, Alenghat T, Nair MG, Giacomin PR, Everett LJ, Feng D, Steger

DJ, Schug J, Artis D and Lazar MA. Histone deacetylase 3 is an epigenomic brake in macrophage

alternative activation. *Genes & development*. 2011;25:2480-8.

67.   Srinivas S, Watanabe T, Lin CS, William CM, Tanabe Y, Jessell TM and Costantini F. Cre

reporter strains produced by targeted insertion of EYFP and ECFP into the ROSA26 locus. *BMC*

Dev Biol. 2001;1:4.

68.   Beall MC, Li D and Jang J. Isolation of Embryonic Cardiomyocytes and Cell Proliferation

Assay Using Genetically Engineered Reporter Mouse Model. *Bio Protoc*. 2023;13:e4802.

69.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M and

Gingeras TR. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*. 2013;29:15-21.

70.   Wolock SL, Lopez R and Klein AM. Scrublet: Computational Identification of Cell Doublets

in Single-Cell Transcriptomic Data. *Cell Syst*. 2019;8:281-291 e9.

71.   Satija R, Farrell JA, Gennert D, Schier AF and Regev A. Spatial reconstruction of single-

cell gene expression data. *Nat Biotechnol*. 2015;33:495-502.

72.   Wolf FA, Angerer P and Theis FJ. SCANPY: large-scale single-cell gene expression data

analysis. *Genome Biol*. 2018;19:15.

73.   Wang L, Feng Z, Wang X, Wang X and Zhang X. DEGseq: an R package for identifying

differentially expressed genes from RNA-seq data. *Bioinformatics*. 2010;26:136-8.

74.   Schmittgen TD and Livak KJ. Analyzing real-time PCR data by the comparative C(T)

method. *Nature protocols*. 2008;3:1101-8.

Acknowledgments

We thank Dr. Dennis Lewandowski in the Heart Center at the Nationwide Children's

Hospital for his critical reading and editing of the manuscript.

Author contributions

J.J. and D.L. conceived and designed the study, analyzed the experiments, wrote the

paper. M.B and M.L performed scRNA-Seq data analysis. Y.J.K and E.K. performed the

experiments. C.C. provided $Tnnt2^{nGFP/+}$ mice. V.G., J.J. and D.L. edited manuscript. All

authors reviewed the results and approved the final version of the manuscript.

Competing Interests

None.

Sources of Funding

This work was partially supported by the National Heart, Lung, and Blood Institute R01

grant HL153406 (D. L.), American Heart Association (AHA)-Career Development Award

23CDA1046244 and AHA-Innovative Project Award 23IPA1048560 (J. J.).

Figures and Figure legends

Figure 1. Endothelial specific deletion of *Hdac3* results in hypotrabeculation.

(A) Hematoxylin and eosin staining on cross sections of an E12.5 *Hdac3^{tko}* embryo (*Hdac3^{f/f}; Tie2-Cre/+*) and a littermate control (CTL) embryo (*Hdac3^{f/f}*). Arrows mark endocardium and arrowheads point to trabeculae. Quantifications are shown on the bottom (CTL: n=8, *Hdac3^{tko}*: n=7). Scale bars, 200 µm (main panels) and 25 µm (insets).

(B) Decreased cardiomyocyte (CM) proliferation in E11.5 *Hdac3^{tko}* hearts. Immunofluorescence staining of Bromodeoxyuridine/5-bromo-2'-deoxyuridine (BrdU) and phospho-Histone H3 (p-H3) staining. High-power Images in the trabecular myocardium (Tra) and in the compact myocardium (Com) are highlighted in white and yellow boxes respectively. Scale bars, 200 µm. Quantification of proliferation index in the trabecular

754   and compact myocardium respectively are shown on the bottom. BrdU (CTL: n=8,

755   *Hdac3^{tko}*: n=7); p-H3 (n=6 in each group). *P*-values were determined by unpaired two

756   tailed Student's *t*-test.

Figure 2. Single-cell RNA sequencing reveals decreased expression of ECM genes in *Hdac3^{tko}* hearts.

(A) Schematic workflow of scRNA-Seq experimental design. Representative micrographs of GFP immunofluorescence staining of E11.5 *Hdac3^{tko}* embryo (*Hdac3^{f/f}; Tie2-Cre/+; R26R^{eYFP/+}*) and a littermate control (CTL) embryo (*Hdac3^{f/+}; Tie2-Cre/+; R26R^{eYFP/+}*). Scale bars, 250 μm. **(B)** UMAP visualization of cell populations in an E11.5 control heart. **(C)** Trabecular and compact cardiomyocyte (CM) score were calculated within *Tnni*+ CM population. Left UMAP showed CMs split into a lower cluster that has a high trabecular score, and an upper cluster that has a high compact score. Right UMAP shows the expression level of *Bmp10*+ per cell. **(D)** Dot plots of differentially expressed genes in *Hdac3^{tko}* and CTL trabecular clusters. Violin plots of expression of *Egr1* and *Eef1a2* in *Hdac3^{tko}* and CTL trabecular clusters. **(E)** The UMAP on the left shows that YFP+ cells are distributed in cardiac endothelial cell (CEC, *Cdh*+), fibroblast cell 1 (FC1, *Pdgfra*+), and fibroblast cell 2 (FC2, *Dcn*+) clusters. The overlapping UMAPs of *Hdac3^{tko}* and CTL YFP+ cells are shown on the right. **(F)** Gene Ontology pathway analysis of top marker genes in CTL-specific YFP+ CEC cluster (Cut off value: -log (FDR)>5). **(G)** Heatmap of extracellular matrix (ECM) associated genes in E11.5 *Hdac3^{tko}* and CTL hearts. Data were extracted from YFP+ CEC clusters in the scRNA-Seq dataset. **(H)** qRT-PCR analysis of *Col1a1*, *Col1a2*, *Col2a1*, *Col3a1* and *P4ha3* in E11.5 *Hdac3^{tko}* heart as compared to CTL hearts (n=8 in each group for *Col1a1*, *Col1a2*, *Col2a1* and *Col3a1,* n=5 in each group for *P4ha3*). *P*-values were determined by unpaired two tailed Student's *t*-test. **(I)** Illustration of decreased expression of proliferation related genes in trabecular CMs and ECM-associated genes in CECs of *Hdac3^{tko}* hearts.

Figure 3. Cardiac endothelial *Hdac3* knockout results in downregulation of TGFβ3 and decreased cardiomyocyte proliferation.

(A) Generation of *Hdac3* knockout (KO) and empty vector control (EV) mouse cardiac endothelial cells (MCECs) by CRISPR/Cas9. Deletion of *Hdac3* was verified by western blot. Quantification is shown below (n=6 in each group). **(B)** The effects of *Hdac3* KO MCEC supernatants on E13.5 *Tnnt2$^{nGFP/+}$* cardiomyocytes (CM) proliferation. Representative immunofluorescence micrographs are shown. Scale bar, 275 μm. Percentage of Ki67+ CMs and total number of GFP+ CMs were quantitated (n=4 in each group). **(C)** Quantification of *Tgfβ3* expression in *Hdac3* KO MCECs, or RGFP966-treated MCECs (10 uM) by qRT-PCR. *Gapdh* was used as cDNA loading control (n=4 in each group). **(D)** Downregulation of *Tgfβ3* gene expression in *Hdac3* deficient cardiac endothelial cells. YFP+ E11.5 cardiac endothelial cells (CECs) from *Hdac3^{tko}* (*Hdac3$^{f/f}$; Tie2-Cre/+; R26$^{eYFP/+}$*) and CTL (*Hdac3$^{f/+}$; Tie2-Cre/+; R26$^{eYFP/+}$*) hearts were sorted by fluorescence-activated cell sorting. Quantification of *Hdac3* and *Tgfβ3* expression by qPCR was shown on the right. n=4 for each group. *Gapdh* was used as cDNA loading control (n=4 in each group). **(E)** Representative Western blot and quantification of TGFβ3 expression in E11.5 *Hdac3^{tko}* hearts (n=12) and littermate CTL hearts (n=8). **(F)** Secretion of TGFβ3 from *Hdac3* KO and EV MCECs. Coomassie brilliant blue staining of total extracted proteins from supernatants served as protein loading controls. TGFβ3 in the MCEC supernatants were detected by western blot. Quantifications are shown on the right (n=4 in each group). **(G)** Rescue of CM proliferation defect by *Hdac3* KO MCECs supernatant supplemented with recombinant TGFβ3 (working concentration: 250 ng/ml). Representative immunofluorescence micrographs are shown. Scale bar, 200 μm.

805   Percentage of Ki67+ CMs and total number of GFP+ CMs were quantitated (n=7 in each

806   group). *P*-values were determined by unpaired two tailed Student's *t*-test for **(A)** and **(E)**,

807   Mann-Whitney U test for **(C)**, **(D)** and **(F)** and one-way ANOVA followed by the Tukey

808   post hoc test for **(B) and (G)**.

Figure 4. HDAC3 induces Tgfβ3 by repressing miR-129-5p.

(A) Heatmap of microRNA (miR) sequencing of *Hdac3* knockout (KO) and empty vector control (EV) mouse cardiac endothelial cells (MCECs). MiRs with reads less than 100 were discarded and miRNA expression levels were normalized by TPM (transcript per million) values (TPM = [miRNA total reads/total clean reads] × 10^6). Log$_2$ fold changes were calculated by TPM per miR in the KO group divided by the mean TPM per miR in the EV group then followed by calculation of Log (fold change, 2). Significantly downregulated miRs are shown in blue, and significantly up-regulated miRs are shown in red. Cut-off criteria: Log2>1.5 and adjusted *P*-value<0.01. **(B)** Quantification of the expression of *Tgfβ3* in MCECs after miR mimics treatment (final concentration 10 nM) by qRT-PCR. *Gapdh* was used as a cDNA loading control (n=6 in each group). Relative gene expression was compared to the miR scrambles (scCTL) group. **(C)** Representative Western blot and quantification of the expression of TGFβ3 after miR-129-5p mimics treatment by western blot (n=8 in each group). **(D)** (Left) Quantification of *Tgfβ3* expression in *Hdac3* KO and EV MCECs (n=4 in each group). (Right) Quantification of the expression of *Tgfβ3* in MCECs after Hdac3 inhibitor treatment (RGFP966, final concentration 5 uM) (n=4 in each group). **(E)** Quantification of the expression of *Tgfβ3* in *Hdac3^{tko}* heart lysates (n=9 in each group). **(F)** Quantification of the expression of miR-129-5p by qRT-PCR**.** *Hdac3* KO or EV MECs were transfected with LentimiRa-GFP-miRZip129 (miRZIP129) or pGreenPuro Scramble Hairpin control lentivirus (scCTL) and transfected cells were selected using puromycin (n=4 in each group)**. (G)** Representative Western blot and quantification of the expression of TGFβ3 after treatment with miRZIP129 or scCTL in *Hdac3* KO and EV MCECs (n=6 in each group). **(H)** Knockdown

of miR-129-5p rescues CM proliferation deficit induced by *Hdac3* KO MCEC supernatants.

Representative immunofluorescence micrographs are shown. Scale bar, 200 µm.

Quantification of percentage of phospho-histone H3 (p-H3)+ CMs was shown on the right.

Independent samples: n=9 in each group. The protein expression of TGFβ3 *P*-values

were determined by unpaired two tailed Student's *t*-test for **(C)** and **(E)**, the Mann-Whitney

U test for **(D)** and the one-way ANOVA followed by Tukey post hoc test for **(B)**, **(F)**, **(G)**

and **(H)**. **(I)** Working model: in the developing endocardium, HDAC3 represses the

expression of miR-129-5p to release their suppression on the expression of TGFβ3.

When *Hdac3* is deleted, the expression of miR-129-5p is increased, which subsequently

suppresses the expression of TGFβ3 to a stronger extent, and the decrease of TGFβ3,

together with decreased ECM, leads to hypotrabeculation.