Genome editing in the mouse brain with minimally immunogenic Cas9 RNPs

Elizabeth C. Stahl[1-3], Jennifer K. Sabo[1*], Min Hyung Kang[1,5], Ryan Allen[1], Elizabeth Applegate[1],

Shin Eui Kim[1], Yoonjin Kwon[1,3], Anmol Seth[1,3], Nicholas Lemus[1,3], Viviana Salinas-Rios[1],

Katarzyna Soczek[1-3], Marena Trinidad[1], Linda T. Vo[1#], Chris Jeans[2], Anna Wozniak[4], Timothy

Morris[4], Athen Kimberlin[4], Thomas Foti[4], David F. Savage[1,3,5] and Jennifer A. Doudna[1-3,5-8†]

[1]Innovative Genomics Institute, University of California, Berkeley, California 94720, USA;

[2]California Institute for Quantitative Biosciences (QB3), University of California, Berkeley,

California 94720, USA; [3]Department of Molecular and Cell Biology, University of California,

Berkeley, California 94720, USA; [4]Aldevron LLC, Madison, Wisconsin 53719, USA; [5]Howard

Hughes Medical Institute, University of California, Berkeley, California 94720, USA; [6]Department

of Chemistry, University of California, Berkeley, CA, USA 94720; [7]MBIB Division, Lawrence

Berkeley National Laboratory, Berkeley, California 94720, USA; [8]Gladstone Institutes, University

of California, San Francisco, CA 94114

†Correspondence: doudna@berkeley.edu

*Current address: Scribe Therapeutics Inc. Alameda, CA, 94501

#Current address: Third Rock Ventures, Boston, MA, 02116

Abstract

Transient delivery of CRISPR-Cas9 ribonucleoproteins (RNPs) into the central nervous system (CNS) for therapeutic genome editing could avoid limitations of viral vector-based delivery including cargo capacity, immunogenicity, and cost. Here we tested the ability of cell penetrant Cas9 RNPs to edit the mouse striatum when introduced using a convection enhanced delivery system. These transient Cas9 RNPs showed comparable editing of neurons and reduced adaptive immune responses relative to one formulation of Cas9 delivered using AAV serotype 9. The production of ultra-low-endotoxin Cas9 protein manufactured at scale further improved innate immunity. We conclude that injection-based delivery of minimally immunogenic CRISPR genome editing RNPs into the CNS provides a valuable alternative to virus-mediated genome editing.

Introduction

Editing somatic cells directly *in vivo* is anticipated to be the next wave of therapeutics for many genetic diseases, especially those affecting the central nervous system (CNS)[1,2]. Clustered regularly interspaced short palindromic repeats (CRISPR) is a revolutionary tool adapted from bacterial immune systems for genome editing[3–5]. To achieve gene disruption, the functional endonuclease, Cas9, is directed by a guide RNA to a target site in DNA to generate a double strand break leading to insertions and deletions (indels). Unfortunately, despite many genetic disease indications, the brain remains a challenging target for genome editing.

To circumvent the blood-brain barrier (BBB), most genomic medicines rely on direct intracranial injection of viral vectors encoding the transgene of interest. Viral vectors, such as recombinant adeno-associated virus (AAV), have had great success in gene therapy and are less immunogenic than most viral vectors, however, they require re-manufacturing for each target and are hindered by costly production scale up. Additionally, AAV has a limited DNA packaging capacity, and is associated with immunogenicity in the brain from both the vector and expression of foreign transgenes[6–9]. Although the brain has been considered an immune-privileged site, green fluorescent protein can induce a strong inflammatory response and neuronal cell death three weeks after injection with AAV serotype 9 has been reported[6–9]. Additionally, Cas9-specific immune responses have been elicited following AAV delivery in mice[10–12] and pre-existing cellular and humoral immunity to Cas9 and AAVs are documented in humans[13–18]. Despite these drawbacks, AAVs are the most clinically relevant delivery systems currently in use for the CNS.

The development of transient, non-viral delivery systems that can effectively edit neurons throughout the brain with minimal immunogenicity would greatly facilitate future clinical applications. Previously, we developed cell penetrating Cas9 ribonucleoproteins (RNPs) capable of genome editing in mouse neurons both *in vitro* and *in vivo*[19]. To enable self-delivery of the Cas9 RNPs, four repeats of the positively charged Simian vacuolating virus 40 nuclear

localization sequences (SV40-NLS) were fused to the N-terminus along with two repeats to the

C-terminus of Cas9, a strategy that was also reported to enable delivery of zinc-finger

nucleases[20]. Using a single guide to turn on the tdTomato reporter from the lox-stop-lox (LSL-

Ai9[21]) mouse, we reported edited striatal volume of approximately 1.5mm^3 [19].

Here we report further optimization of cell penetrant Cas9 RNPs, demonstrating efficacy

in human primary cells and improved editing of the mouse striatum using a convection

enhanced delivery (CED) system. We compared the transient RNP complexes to AAV serotype

9 for Cas9 delivery to the CNS, to measure both editing efficiency and the host immune

response. We found that the Cas9-AAV was able to better diffuse throughout the brain, leading

to distally edited cells; while the cell penetrant Cas9-RNPs edited significantly more neurons

within the region near the injection site. Both groups elicited humoral responses, but vehicle-

specific antibodies in the Cas9-AAV group persisted at high levels out to 90-days. Cas9-AAV

treated brains were also associated with significantly elevated *Cd3e* gene expression at four

weeks, suggesting an ongoing adaptive immune response. Cas9-RNP treated brains showed

acute microglial activation that was mitigated by reducing endotoxin levels during protein

manufacturing scale up. Taken together, Cas9 RNPs are a promising strategy for future

therapeutic intervention in neurological disorders to address current limitations of viral delivery.

Results

Development of Cas9 cell penetrant RNP and AAV to measure genome editing with the

tdTomato reporter system

Creating a large deletion in the lox-stop-lox cassette in Ai9 mice with a single guide RNA (sgRNA) enables expression of tdTomato and efficient quantification of editing by fluorescent read out (Figure S1A). Cas9 from *Streptococcus pyogenes* (engineered with four copies of SV40 NLS on the N-terminus and two copies on the C-terminus (4x-SpyCas9-2x) to be cell penetrant) was first produced from recombinant *E. coli* in a laboratory setting, using a low-endotoxin method. Editing efficiency of the RNP was compared to Cas9 delivered by recombinant adeno-associated virus (AAV) (Figure 1A-B). Since SpyCas9 cannot be packaged within a single AAV with its guide RNA, we used clinically relevant AAV-SauCas9-sgRNA (derived from *Staphylococcus aureus*)[22–24]. AAV serotype 9 was produced using a baculovirus transfected into Sf9 insect cells[25,26]. To control for differences in the Cas9 orthologs, cell penetrant 4x-SauCas9-2x protein was also produced following the same expression and purification methods as 4x-SpyCas9-2x. Due to differences in PAM requirements between the two Cas9 orthologs (SpyCas9 NGG, SauCas9 NNGRRT), a new guide was designed for SauCas9 to target the tdTomato locus (Figure S1B).

We confirmed editing in neural precursor cells (NPCs) isolated from embryonic day 13.5 Ai9 mice with all constructs *in vitro* (Figure S1). Addition of 4x-SauCas9-2x RNP into the cell culture supernatant enabled editing of NPCs compared to 0x-SauCas9-2x RNPs, demonstrating that the four SV40-NLS peptides can mediate delivery of additional Cas9 orthologs. 4x-SauCas9-2x RNPs were slightly less efficacious than 4x-SpyCas9-2x RNPs, which could be explained in part by differences in the guide RNAs (Figure S1A-B). We observed editing of mouse NPCs with SauCas9 when delivered as both cell penetrant RNP (59 ± 6% tdTomato cells[+], Figure S1D) and AAV (42 ± 1% tdTomato cells[+], Figure S1E). The same titer of AAV9-

104   CMV-GFP resulted in 96 ± 2% GFP[+] cells, suggesting that editing lagged behind transduction;

105   and an increase in empty capsids in the AAV-CMV-SauCas9 group was noted (Figure S1F)[27,28].

106   To further examine the potential for cell penetrant Cas9 RNPs to edit difficult cells *in*

vitro, we tested delivery and editing with 4x-SpyCas9-2x in human neural precursor cells

derived from induced pluripotent stem cells (iPSCs)[29–31]. Human NPCs were treated with pre-

formed RNPs using an established guide RNA targeting EMX1[32]. In human cells, we detected

10-fold higher rates of editing with 4x-SpyCas9-2x compared to standard RNPs (0x-SpyCas9-

2x) delivered with commercial transfection reagents (Figure S2A-B).

Cas9 RNPs result in modest editing of brain parenchyma following delivery into cerebrospinal

fluid

115   To determine the optimal route of delivery for Cas9 RNPs into the mouse CNS, we

tested intraparenchymal injections into the striatum, as well as injection into the cerebrospinal

fluid (CSF), including intrathecal (IT) and intracerebroventricular (ICV) routes. Following IT

injection of cell penetrant RNPs, we observed edited glial cells and neurons in the cortex and

striatum of one hemisphere, but no editing within the spinal cord (Figure S3A). Following ICV

injection of Cas9 RNPs in neonatal p0 mice, we observed tdTomato[+] cells in the subventricular

zone and white matter, including glial cells and neural stem/progenitor cells expressing Ki67 and

DCX evaluated three weeks after delivery (Figure S3B-C). Editing post-ICV injection in adult

mice was restricted to the cells within the lateral ventricles, choroid plexus, subventricular zone,

and hippocampus in a subset of mice (Figure S3D-E). The total number of edited cells with RNP

delivery into the CSF was lower than with direct intraparenchymal injection (Figure S3F).

126   Therefore, we sought to further improve upon intraparenchymal injections using a

convection enhanced delivery system (CED), which generates a high-pressure gradient to aid in

biodistribution of macromolecules in the brain. CED has been used to increase the spread of

AAV in the brains of large animal models and humans by infusing relatively high injection

volumes at high rates[33,34]. Additionally, we tested two needle designs to enable CED with Cas9-

RNPs. We found that the CED enabled robust editing in the mouse striatum (Figure S4B) and

the step-cannula reduced reflux of RNP from the needle-injection track, as reported previously[35]

(Figure 1D). Furthermore, tdTomato⁺ neurons edited by Cas9-RNPs within the striatum were

observed to extend along the basal ganglia circuit into the globus pallidus and substantia nigra

along the anterior-posterior axis (Figure 1C-E).

Convection enhanced delivery of Cas9 RNPs and AAVs mediates robust editing in the mouse

striatum

Using bilateral CED injections into the striatum, we compared edited tissue volume with

the 4x-SpyCas9-2x RNP, 4x-SauCas9-2x RNP, and AAV9-SauCas9-sgRNA in adult Ai9 mice at

three weeks post-injection. Despite performing well *in vitro*, 4x-SauCas9-2x RNPs

underperformed *in vivo* when tested at two different doses and additional NLS configurations

(Figure S5A-G, S6D). Therefore, we performed our primary comparison with two orthologous

systems: 4x-SpyCas9-2x RNP (hereafter referred to as Cas9-RNP) and AAV9-SauCas9-sgRNA

(hereafter referred to as Cas9-AAV, which serves as a positive control, Figure 1F).

First, we tested Cas9-AAV injection with CED at two doses, 3×10^8 vg/μL and 3×10^9

vg/μL (1.5×10^9-1.5×10^{10} vg/hemisphere)[36], and proceeded with the higher dose for subsequent

studies *in vivo* (Figure S6A-B). We also tested several doses of Cas9-RNP ranging from 10μM

to 100μM (50-500 pmol/hemisphere). Interestingly, there was no significant difference in editing

when delivering RNPs in this concentration range (Figure 1H, Figure S6C). We chose the 25μM

RNP concentration (4.15 mg/mL or approximately 1.75 mg/kg Cas9) group for further study as it

had the highest maximal editing rate. Above 25μM in the RNP group, we observed a decrease

in NeuN staining and an increase in GFAP staining out to 90-days in the Cas9-RNP group,

suggesting dose-limiting effects (Figure 1G).

At both 21 and 90-days post-injection, the Cas9-AAV group outperformed the Cas9-RNP

group when quantifying total edited striatal volume (n=8 at 21-days, n=4 at 90-days, p<0.05,

Figure 1I). The volume of edited cells was relatively stable in the Cas9-AAV group between 21

and 90-days at approximately 47 ± 3% (covering approximately 13.4 mm^3 of striatum), while the

Cas9-RNP group had editing levels of 22 ± 3% (approximately 6.2 mm^3 of striatum) between 21

and 90-days (increased from previous report of editing 1.5mm^3 striatal volume[19]). Edited cells

were observed further along the rostral-caudal axis in the Cas9-AAV group (-2.12 mm to 2.5

mm relative to Bregma), demonstrating better diffusion of the editor away from the injection site

(Figure S6E-F).

Since large deletions in the tdTomato locus make on-target editing difficult to assess

using short-read next-generation sequencing (NGS), we developed an NHEJ droplet digital

PCR assay (ddPCR) to measure drop-off of HEX-labeled probes over the cut sites, in relation to

distal reference FAM-labeled probes. Genomic DNA was isolated from 2-mm thick sections of

each injected hemisphere, covering multiple brain sub-structures. Loss of the HEX probe

reached 2 ± 1% in the Cas9-RNP group and 15 ± 10% in the Cas9-AAV group, indicating edited

alleles, when measured at 28-days (Figure S7).

We also quantified the percentage of edited NeuN$^+$ neurons within the tdTomato$^+$ region

of interest (ROI) per hemisphere between Cas9-AAV and Cas9-RNP at the 21-day timepoint.

We found that Cas9-RNP edited significantly more NeuN$^+$ neurons per ROI (36 ± 10%)

compared to Cas9-AAV (20 ± 2%) (Figure 1J, n=6-8 injections, p<0.05). Within the ROI,

neurons were the most frequently edited cell type in both groups, including DARPP-32$^+$ medium

spiny neurons (Figure S8). Additionally editing of ALDH1L1$^+$ and OLIG2$^+$ glial cells was noted in

both groups (approximately 2% of edited cells within the ROI in the Cas9-RNP group and 8% of

cells in the Cas9-AAV group). Therefore, Cas9-RNPs were able to edit comparable numbers of

neurons and glia as Cas9-AAVs in a given area of striatal tissue.

Comparison of local and peripheral immune response between Cas9 RNPs and AAVs in the Ai9

reporter mouse

We next examined the local and peripheral immune response following delivery of Cas9 RNPs and AAVs into the brain. Using immunofluorescent staining for Iba1 (Figure 2A), we observed a significant increase in percent Iba1$^+$ area in the 25µM Cas9-RNP group from sham-treated animals (Figure 2B, n=6 replicates, p<0.05). Staining for CD45 showed dim expression on Iba1$^+$ microglia and high expression on CD3$^+$ T-cells, which were slightly increased in the 25µM Cas9-RNP group compared to the sham and Cas9-AAV, but not significantly different, at three weeks post-injection (Figure 2C-D, n=6-12 replicates).

In addition to the immune response at the local site of injection, circulating IgG antibodies were measured at 28 and 90-days post-injection. We found that sham-treated animals had no pre-existing antibodies to either SpyCas9, SauCas9, nor AAV9 capsids. At 28-days following striatal injection, there was a 1.6e4-fold increase in anti-SpyCas9 IgG in the 25µM Cas9-RNP group, a 1.3e4-fold increase in anti-AAV9 capsid IgG in the Cas9-AAV group, and an 8.9e1-fold increase in anti-SauCas9 IgG in the Cas9-AAV group (i.e., humoral response against transgene) (Figure 2E, n=3-5 biological replicates). No cross-reactivity was observed between ortholog RNPs, as described previously[11], nor were any anti-AAV capsid antibodies detected in the RNP group. At 90-days, the levels of IgG fell to a 5.4e2-fold increase in the 25µM Cas9 RNP group and 1.2e4-fold increase in the Cas9 AAV group from the sham controls, demonstrating greater maintenance of systemic antibodies against the capsid in the AAV group.

The cellular and humoral immune response to Cas9 RNPs was dose-dependent and a significant increase in CD45$^+$ cells was observed at the 100µM RNP dose compared to sham, Cas9-AAV, and Cas9-RNP at 25µM (Figure S9). Cas9-reactive cells were also identified in the spleen by interferon-gamma (IFN-γ) ELISpot assay (Figure S9E) at both 25µM and 100µM doses of Cas9-RNPs, but not in sham treated animals.

Since the mice had no pre-existing antibodies to SpyCas9, we tested how the immune

response would differ in the RNP group by first exposing the mice to a single subcutaneous

injection of 4x-SpyCas9-2x protein and adjuvant (AddaVax^{TM}) four-weeks prior to stereotaxic

surgery with Cas9-RNPs. We found that pre-exposing the mice to Cas9 with adjuvant had a

synergistic effect on both serum IgG and activation of IFN-γ^+ cells in the spleen (Figure S9F-I).

Mice that received surgery maintained tdTomato$^+$ cells in the brain to the measured time point.

Additional studies using this immunization strategy may help to further characterize the immune

response to Cas9-RNPs by modeling pre-existing immunity relevant to humans.

Finally, we measured gene expression changes near the injection site in mice that

received Cas9-RNP and AAV at 3 and 28-days post-injection using RT-qPCR. At three days,

the Cas9-AAV group had a modest but significant increase in *Fas* (1.19-fold) and *Fasl* (1.85-

fold) compared to the sham group (Figure 2F). At 28-days post-injection, both Cas9-RNP and -

AAV had a significant increase in *Fas* (1.61 and 1.89-fold respectively). In addition, the Cas9-

AAV group had a significant increase in *CD3e* gene expression (5.45-fold, n=4 replicates,

$p<0.05$), closely followed by *CD8a* (2.06-fold, ns, p=0.06), while Cas9-RNP had a slight but non-

significant increase in *CD3e* (2.15-fold, ns, p=0.06), compared to the sham group.

There were no detectable off-target editing events at 1 and 4-months post-injection in

any of the experimental groups at the evaluated sites (Figure S10A-C). In the Cas9-AAV group,

the Cas9 transgene was expressed in the brain out to 4-months, the last tested time point, as

expected (Figure S10D). Additionally, few genes were differentially expressed between the

groups at 4-months, except for *Fas* (1.54-fold, p<0.05), which was significantly elevated in the

Cas9-AAV group compared to the sham (Figure S10E-F). We used long read sequencing to

examine whether any fragments of the viral genome had been integrated near the cut site in the

tdTomato locus, as previously reported[37–40]. We also observed partial integrations of viral

fragments in our amplicon, although our *in vivo* editing rates and sequencing depth were

relatively low (Figure S11).

Overall, delivery of Cas9 by either AAV or RNP resulted in significant activation of

immune responses in the brain and periphery, although with generally small effect sizes

compared to sham injected mice. The increased Iba1$^+$ cells near tdTomato$^+$ cells in the striatum

of the 25µM Cas9-RNP group raised the question of whether the response was due to Cas9

itself or impurities within the protein product. We hypothesized that the local immune response

may be due to endotoxins from *E. coli* in the RNP complexes.

Production and testing of ultra-low endotoxin 4x-SpyCas9-2x protein

To examine the impact of endotoxin on the immune response to RNPs, we partnered

with a commercial producer of Cas9 protein and were able to significantly scale up

manufacturing to produce a large quantity of ultra-low endotoxin 4x-SpyCas9-2x protein using

an industrial tag-free expression and purification system (Figure 3A).

Using the limulus amebocyte lysate (LAL) assay, we measured an endotoxin

concentration of 0.035 EU/mg in the industrial-produced protein compared to 0.2 EU/mg in the

lab-produced protein (Figure S12A). Interestingly, using the same assay, we found that guide

RNA could be an unexpected source of endotoxin contamination. Endotoxin was present in at

least three unopened vials of lyophilized RNA that had been stored at -80°C from a 2018 lot, but

not in a more recently purchased lot from the same vendor when resuspended simultaneously

(Figure S12A-C). To rule out false positives due to reaction of LAL with beta-glucans, we

performed the Recombinant Factor C (rFC) assay. We measured a similar level of endotoxins in

the guides between the LAL and rFC assays, demonstrating the positive signal was from

contamination with endotoxin and not beta-glucans (Figure S12D-E).

To measure the physiological impact of endotoxin in our samples, we used HEK293

cells that were engineered to produce secreted embryonic alkaline phosphatase (SEAP)

downstream of NF-κB activation resulting from human toll-like receptor 4 stimulation (hTLR4)

with endotoxin/lipopolysaccharide (LPS, Figure S13A-E). The lab-produced protein stimulated

NF-κB in HEK293 cells significantly greater than the industrial produced protein (p<0.01,

unpaired t-test). Treatment with the industrially produced protein led to similar levels of SEAP

between hTLR4 cells and the parental cell line (Null2), demonstrating that most of the NF-κB

stimulation was downstream of other pattern-recognition receptors (such as TLR3, TLR5, or

nucleotide-binding oligomerization domain-containing protein 1 (NOD-1) activation) and not due

to LPS signaling through hTLR4 (Figure S13A-E). When combined with sg298 from the 2018 or

2022 lot, absorbance levels of SEAP further increased in RNP complexes made with lab

produced protein, while the industrial protein with either guide did not induce a response (Figure

S13F). Furthermore, guide RNA alone did not stimulate NF-κB in HEK293 cells (Figure S13D).

Finally, we measured endotoxins in the "optimized" formulation of RNPs, comprised of

the industrially produced 4x-SpyCas9-2x protein and 2022 sgRNA, using the LAL assay.

Estimating delivery of 10μL per mouse, the endotoxin burden was 0.44 EU/kg when RNPs were

formulated at 25μM. These data suggest that RNPs could be delivered below the 0.2 EU/kg

FDA threshold for intrathecal delivery[41] when formulated at 10μM without significant loss of

editing (Figure 3B and Figure 1H).

Optimized RNP formulation reduces immune response

We performed CED bilateral intrastriatal injections to test if reducing endotoxins would

improve the host immune response to RNPs *in vivo*. In this experiment, we compared the

optimized RNP formulation (industrially produced 4x-SpyCas9-2x NLS protein with sg298 2022)

to the standard formulation used in Figures 1 and 2 (laboratory produced 4x-SpyCas9-2x NLS

protein with sg298 2018) at 25μM. The standard RNP induced a significant increase in Iba1$^+$

area, consistent with our previous measurements (Figure 3C and Figure 1A); however, the

optimized Cas9-RNP formulation did not induce microglial activation. Additionally, there was no

increase in CD45$^+$ and CD3$^+$ cells from the sham in the optimized RNP group (Figure 3D). Of

note, the standard RNP edited an average of 31 ± 3% striatal volume (greater than values

reported in Figure 1, possibly due to differences between protein lots or injections), while the

optimized RNP edited an average of 23 ± 8% striatal volume (Figure 3F). When tested *in vitro,*

the optimized RNP performed slightly better at direct delivery than nucleofection compared to

the standard formulation, which could explain in part the differences *in vivo* (Figure S14A).

Interestingly, the standard RNP also lead to significantly greater anti-Cas9 IgG responses at

three-weeks post-injection, possibly due to endotoxin boosting the adaptive immune response

(Figure 3G). Taken together, we found that reducing endotoxins in both the guide RNA and

protein components of the RNP leads to a reduced innate immune response, comparable to the

sham, while maintaining high on-target editing. Furthermore, cell penetrant Cas9 proteins are

amenable to expedited manufacturing of large quantities suitable for *in vivo* experiments.

In conclusion, our results establish complementary genome editing and immunogenicity

outcomes between the two tested Cas9 delivery strategies (Figure S14B). To enable high-levels

of editing in neurons within a localized brain region, minimizing adaptive immune responses,

and timely and affordable manufacturing scale up, the RNP offers an effective alternative

delivery system to viral vectors.

Discussion

In the present study, we demonstrate that cell penetrant Cas9 RNPs edit a significant

volume of the mouse striatum using convection enhanced delivery. Furthermore, the 4x-NLS

modification enables self-delivery of Cas9 orthologous proteins *in vitro* to both mouse and

human cells, demonstrating cross-species compatibility of the system for the first time. We also

show that Cas9 RNPs have dose-dependent effects on the immune response, which can be

mitigated by using ultra-low endotoxin protein produced in an industrial non-GMP setting. These

experiments are informative for the design of future therapeutic applications of Cas9 RNP

editors in mice and larger animal models.

Several studies have reported non-viral delivery of Cas9 into the mouse brain. The

"CRISPR-Gold" Cas9 nanoparticle delivery system induced 14% edited glial cells near the

injection site, sufficient to reduce repetitive behaviors in a mouse model of fragile X syndrome[42].

Additionally, incubating RNPs with R7L10, an arginine and leucine rich cationic peptide, induced

45% indels in the CA3 region of the hippocampus, leading to behavioral improvements in an

Alzheimer's disease mouse model[43]. Efficient editing of DARPP-32 medium spiny neurons in

the striatum was achieved here and in recent work by others using RNPs packaged in

biodegradable PEGylated nanocapsules[44]. Interestingly the PEGylated nanocapsules have a

neutral charge, while 4x-Cas9-2x NLS RNPs have a net-positive charge, suggesting the

mechanism of entry may differ between the two strategies. Systemic delivery of genome editors

with glucose-conjugated silica nanoparticles and AAV9 can also lead to modest levels of editing

in the brain, sufficient for therapeutic benefit [45,46]. Despite the need for direct injection, the

simplicity of the 4x-Cas9-2x RNP makes it ideal from a manufacturing perspective compared to

other viral and nanoparticle formulations. The RNP formulation buffer could be further

supplemented with polymers, such as polyethylene glycol (PEG)[47,48], to possibly improve

biodistribution with CED in the future.

Several studies show correlation between editing at the tdTomato locus and subsequent

editing at endogenous sites[42,49,45]. It is important to note that expression of the tdTomato protein

in the Ai9 mouse model underreports the actual genome editing efficiency, as double strand

breaks that result in indels and small deletions are not sufficient to turn on the reporter[19]. We

sought to further resolve editing outcomes with a ddPCR assay, since the tdTomato locus is too

large for Illumina-based NGS. In the ddPCR probe drop-off assay we detected approximately a

6-fold increase in editing with AAV compared to RNP (approximately 15% versus 2.5% indels),

whereas our immunofluorescent measurement showed a 2-fold increase. We speculate that the

difference in reported editing efficiency between the two assays is based on the sampling

methodology. The image analysis workflow quantified the volume of striatal tissue containing

edited cells (Figure 1I), where 20-36% of neurons were edited within 25-50% of the striatum

(Figure 1J). Therefore, the image quantification reflects the maximal biodistribution of edited

cells in the striatum. The ddPCR assay utilized genomic DNA from 2-mm thick tissues dissected

from the expected injection site including the striatum, as well as the cortex and corpus

callosum, which likely favors editing events in the Cas9-AAV group due to its enhanced diffusion

through the brain. While dose escalating the RNP did not improve editing levels, suggesting a

saturating dose was already achieved as measured by image quantification, editing at different

doses was not assessed using ddPCR. Furthermore, increasing the dose of Cas9-AAV may

have further increased editing levels. Studies suggest that correcting pathological mutations in

20-30% of striatal neurons expressing mutant huntingtin protein is sufficient to significantly

improve the disease pathology, therefore even modest editing levels in the striatum could

enable therapeutic benefit[50]. Ultimately, behavioral assays are needed to further determine

therapeutic benefit of the RNP approach in a disease-relevant model.

We hypothesized that cell penetrant Cas9-RNPs would be less immunogenic than Cas9-

AAVs due to their transient expression. As the dose of 4x-SpyCas9-2x RNPs increased from

25µM to 100µM, there was an increase in $CD45^+$ and $GFAP^+$ cells, and a decrease in $NeuN^+$

cells. As such, subsequent experiments were performed at 25µM, which was well-tolerated and

resulted in similar levels of editing as the higher dose. The 25µM Cas9-RNP led to lower levels

of vehicle-specific antibodies by 90-days post-injection compared to AAVs and did not

upregulate gene signatures of T-cells at 28-days as measured by RT-qPCR, supporting our

hypothesis. By 90 days, the levels of anti-Cas9 antibodies in the Cas9-AAV group were only

elevated in three out of five mice, demonstrating that the kinetics of Cas9 antibody persistence

were similar between RNPs and AAVs, despite stable, intracellular Cas9 expression. Reducing

endotoxin in both the Cas9 protein and guide RNA prevented microglial reactions and reduced

humoral responses at 21-days.

In the Cas9-AAV group, few immune cells (CD45, Iba1, or CD3) were observed in the

striatum by immunostaining, however *CD3e* gene expression was significantly upregulated in

explanted tissue, closely followed by an increase in *CD8a*. This finding could indicate

accumulation of cytotoxic T-cells trafficking into the parenchyma from the blood vessels or

ventricles. Additionally, no changes in NeuN, GFAP, and CD45 expression were observed in the

Cas9-AAV group out to 4 months, demonstrating that the AAV delivery strategy was well-

tolerated overall in naïve mice. A study by Li et al. found that mice immunized against SauCas9

with Freund's adjuvant one week prior to intravenous delivery of AAV8-SauCas9-sgRNA

resulted in accumulation of cytotoxic T-cells in the liver and subsequent removal of edited

hepatocytes[12]. Therefore, the host immune response to Cas9-AAV in mice with pre-existing

immunity would likely be different than what we observed in naïve mice. In the Cas9-RNP

group, we found that pre-exposing mice to SpyCas9 protein with AddaVax[TM] adjuvant 4 weeks

prior to stereotaxic surgery synergistically increased systemic adaptive immune responses.

Further studies are needed to assess the immune response to Cas9-AAV and RNP in models

with pre-existing immunity, but how well these immunized mouse models recapitulate pre-

existing immunity in humans is not clear. Furthermore, breakdown of the BBB in the context of

neurodegenerative disease or strong expression of the tdTomato fluorescent reporter could also

impact the host immune response[51,52].

In this study, we used a strong CMV-promoter to drive expression of SauCas9 from the

AAV, which allowed us to assess all subsets of edited cells in the striatum, in comparison to the

RNP, which is not inherently designed to be neuron-specific. Although the SauCas9 transgene

was still expressed 4-months post-delivery, editing at predicted off-target sites was not detected.

Further work to experimentally determine guide-specific off-target sites, such as Guide-Seq[53] or

Circle-Seq[54], was not performed. To prevent potential genotoxic side-effects due to long-term

Cas9 expression, we recommend applying additional safeguards, such as AAV self-inactivation

strategies and neuron-specific promoters, such as human synapsin 1 (hSyn), to increase cell

specificity[55–57]. While self-inactivating AAVs may improve safety, they may not be sufficient to

reduce partial integration of the viral genome at the Cas9 cut site, which has been reported[37–40].

Strategies to mitigate the host response to genome editors include providing

immunosuppressants with CRISPR-Cas9 infusion and screening for pre-existing immunity prior

to dosing when translating *in vivo* editing to humans[58].

While SpyCas9 is generally the most efficacious and widely used Type II CRISPR

protein to date, it is too large to be packaged in a single AAV with sgRNA. Therefore, in this

study, we used AAV-CMV-SauCas9-U6-sgRNA as a positive control to benchmark delivery of

RNPs, as similar AAV constructs have been used extensively in the literature and in clinical

trials[24,23,12,59]. *In vivo* delivery of 4x-SauCas9-2x RNP did not lead to significant editing, thus,

most comparisons were performed between two orthologous genome editors and delivery

systems with different NLS configurations, which may impact the interpretation of this work. *In*

vivo editing with 4x-SauCas9-2x RNP was possibly hindered by the reduced thermostability of

the protein, as reported previously[60]. Employing Cas9 RNPs from a more thermostable

organism could be advantageous in the future. Since RNPs are not restricted by cargo

packaging, one advantage of the cell penetrant RNP technology is the ability to use SpyCas9

for future therapeutic applications in the brain.

In conclusion, the cell penetrant 4x-Cas9-2x NLS fusion protein enables simple and

effective delivery of Cas9 RNPs into neurons *in vitro* and *in vivo* Our study is the first to

comprehensively profile the host immune response to Cas9 in the brain, benchmark an RNP

delivery strategy against the gold-standard for gene delivery in the CNS, and demonstrate

feasibility of large-scale manufacturing. Given that Cas9-RNPs excel at editing high levels of

neurons within a localized region of the brain, this is a promising modality to characterize

therapeutic benefit in disease models in the future.

Materials & Methods

Plasmid construction

Cloning of several spacers into a plasmid encoding SauCas9 was performed as

previously described. Oligonucleotides encoding sgRNAs were custom synthesized (Integrated

DNA Technologies; IDT, Coralville, IA) and phosphorylated by T4 polynucleotide kinase (New

England Biolabs; NEB, Ipswich, MA) for 30 min at 37°C. Oligonucleotides were annealed for 5

min at 95°C, cooled to room temperature and ligated into the BsmBI restriction sites of

pSTX8,pKLT7.1_SaCas9prot_SaCas9guide plasmid. The following 23nt spacer sequences

were cloned into the plasmid (spo 1: TGGTATGGCTGATTATGATCCTC; spo2:

TCCCCCTGAACCTGAAACATAAA; spo3: GATGAGTTTGGACAAACCACAAC; spo4:

TCCAGACATGATAAGATACATTG; spo5: CTCATCAATGTATCTTATCATGT), and plasmids

were used for editing in mouse neural precursor cells *in vitro*. The best performing SauCas9

spacer (spo4: TCCAGACATGATAAGATACATTG) was then cloned into an AAV2 backbone

plasmid. pX601-AAV-CMV::NLS-SaCas9-NLS-3xHA-bGHpA;U6::BsaI-sgRNA was a gift from

Feng Zhang (Addgene plasmid # 61591 ; http://n2t.net/addgene:61591 ;

RRID:Addgene_61591). Briefly, the plasmid was digested using BbsI and a pair of annealed

oligos were cloned into the guide RNA destination site by Golden Gate assembly. Correct

construction of all plasmids was verified by Sanger sequencing (UC Berkeley DNA Sequencing

Facility).

Recombinant adeno-associated virus (AAV) Production

The custom AAV9-CMV-61591-HA-Bgh vectors were produced at Virovek (Hayward,

CA) in insect Sf9 cells by dual infection with rBV-inCap9-inRep-hr2 and rBV-CMV-61591-HA-

Bgh. The AAV9-CMV-GFP vectors were produced by dual infection with rBVinCap9-inRep-hr2

and rBV-CMV-GFP. The vectors were purified through two rounds of cesium chloride (CsCl)

ultracentrifugation. The CsCl was removed through buffer exchange with two PD-10 desalting

columns. Viral titer (approximately 2e13 vg/mL) and purity were confirmed by nanodrop

spectrophotometer, real-time PCR, and SDS-PAGE protein gel analysis. The vectors were

passed through 0.2um sterilized filters, tested for endotoxins (< 0.6 EU/mL), as well as

baculovirus and Sf9 DNA contamination (not detected).

Purification of low-endotoxin proteins in a laboratory setting

Protein expression and purification was performed in the QB3 Macrolab at UC Berkeley

using a custom low-endotoxin workflow. Briefly, the plasmid, 4xNLS-pMJ915v2 (Addgene

plasmid # 88917; http://n2t.net/addgene:88917; RRID:Addgene_88917), was transformed into

E. coli Rosetta2(DE3)pLysS cells (Novagen) and an overnight culture was used to inoculate 1 L

flasks (12-24 L total per batch). Cells were grown for approximately 3 hours at 37°C then

cooled to 16°C. At OD 0.8-0.9, cells were induced and harvested after 16-18 hours growth.

Cells were lysed by homogenization in a buffer containing 1mM $MgCl_2$ and benzonase (1:1000)

to help reduce viscosity and centrifuged to remove insoluble material. Purification by Ni affinity

(10 mL Ni resin for every 6 L cell lysate) was performed, and the bound protein was washed

with 10 column volumes of buffer containing 0.1% Triton-X114 at 4°C to help reduce

endotoxins. Tag removal with TEV protease (1:100) was performed overnight at 4°C, then

heparin affinity was used to concentrate each batch of protein which was then flash frozen and

stored at -80°C. A Sephacryl S300 size-exclusion column (SEC) and flow path were sanitized

with 0.5 M NaOH overnight, then washed with up to 3 column volumes of buffer to rinse and

equilibrate the system. Frozen samples were thawed, combined, and adjusted to 4.5 mL, and

the S300 standard protocol was performed for size-exclusion. Samples were refrigerated

overnight, and sanitation and size-exclusion were repeated the next day to further reduce

endotoxin contamination. Peak fractions were pooled, concentrated to 40µM, aliquoted at 50µL,

flash frozen in liquid nitrogen, and stored at -80°C in sterile, endotoxin-free Buffer 1 (25 mM

NaP (pH 7.25), 300 mM NaCl, 200 mM trehalose (Sigma Aldrich #T5251, St. Louis, MO)). Final

average protein yield was 1 mg per 1 L cells. Plasmids for 2xNLS-SauCas9-2xNLS, 3xNLS-

SauCas9-2xNLS, and 4xNLS-SauCas9-2xNLS were created by deletion mutagenesis using the

existing 4xNLS construct as a template. The genes were fully sequenced to confirm no

additional mutations were introduced during the mutagenesis procedure.

Purification of ultra-low endotoxin proteins in an industrial setting

4x-SpyCas9-2x NLS protein was manufactured according to Aldevron proprietary

workflows for expression and purification of gene editing nucleases. Briefly, the gene for 4x-

SpyCas9-2x NLS was synthesized (ATUM Bio, Sunnyvale, CA) and cloned into a pD881

expression vector (ATUM). Expression-ready plasmid DNA was transformed into E. coli

BL21(DE3) (New England Biolabs) culture in animal-free TB media. At the appropriate OD600,

expression was induced with 2.0% (w/v) Rhamnose and growth culture was harvested by

centrifugation. Cells were lysed via dual-pass high-pressure homogenization and clarified via

centrifugation. The clarified lysate was purified via multi-step chromatography using

standard/commercially available resins. In the final chromatography step, the product is eluted

via step elution and pooled to maximize final protein purity and minimize endotoxin. Product was

dialyzed into the final formulation buffer, underwent three (3) exchanges of buffer, and was

pooled into a sterile vessel for final filtration and dispensing. Product was evaluated for key

quality attributes including endotoxin via PTS Endosafe assay (Charles River Labs, Cambridge,

MA).

Quantification of endotoxins in Cas9 RNPs

Proteins, guide RNAs, and pre-formed RNP complexes were subjected to several

assays to quantify endotoxin burden according to the manufacturer's instructions. All assays

were performed with autoclaved or certified pyrogen-free plasticware and endotoxin (ET)-free

water. The plate-reader based LAL assay was performed with the Endosafe Endochrome-K kit

(Charles River, #R1708K), where a control standard endotoxin (CSE) was diluted from 5 EU/mL

to 0.005 EU/mL. Samples were diluted 1:100 and plated in triplicate. An equal volume of LAL

was added to each well. A Tecan Spark plate reader (Tecan, #30086376, M□nnendorf,

Switzerland) with SparkControl magellan V 2.2 software was used at 37°C to read absorbance

at 405nm every 30 seconds for 100 cycles. Time at which absorbance crossed optical density

(OD) of 0.1 was recorded and used to determine endotoxin levels.

The cartridge-based LAL assay was performed using an Endosafe nexgen-PTS machine

with R&D cartridges as recommended (Charles River, cat # PTS2005, 0.05 EU/mL sensitivity).

Briefly, samples were diluted 1:50 in a large volume of ET-free water, vortexed, and 25µL was

loaded into each of the four lanes of the cartridge, where two lanes contain CSE spike-in to

calculate efficiency of the assay, which is valid from 50%-200% recovery. The final valid ET

value was recorded from the duplicate measurement from a single cartridge.

The PyroGene Recombinant Factor C Endpoint Fluorescent Assay (Lonza, Walkersville,

MD, cat # 50-658U) was performed as recommended. Kit-supplied CSE was diluted from 5

EU/mL to 0.005 EU/mL and samples were diluted 1:100 in ET-free water and added to a plate in

triplicate. The plate was heated at 37°C for 10 minutes, then an equal volume of working

reagent was added to each well. Fluorescence was read immediately at time 0 and again after

incubating for 60 minutes. Relative fluorescence units (RFUs) of the ET-free water only blank

wells were subtracted from all measurements, then delta RFUs between the two time points

was calculated, and a linear regression was applied to the standard curve to calculate EUs in

the samples. Fluorescence measurements were performed on a Cytation5 with Gen 5 3.04

software (BioTek, Winooski, VT).

HEK-Blue cells (hTLR4 and Null2) were purchased from Invivogen (San Diego, CA) and

were grown under BSL2 conditions (37°C with 5% CO_2) to measure SEAP production

downstream of NFkappaB activation following treatment with Cas9 proteins, guide RNAs, and

RNPs *in vitro* as recommended. Cells were grown in T-75 flasks with supplied antibiotic

selection markers and passaged at 70% confluency. Cells were detached with gentle scraping

in 1x PBS, centrifuged, counted, and plated for experiments in freshly prepared HEK-Blue

Detection Media at approximately 32,000 cells per well in a 96-well plate. 180µL of cell

suspension was plated directly into 20µL of diluted CSE (5 to 0.078 ng) or samples (diluted to

10µM) and incubated overnight at 37°C. Absorbance was read at 620nm in a Tecan Spark plate

reader (Tecan, #30086376, M□nnendorf, Switzerland).

Neural progenitor cell (NPC) line creation and culture

Neural progenitor cells were isolated from Ai9-tdTomato homozygous mouse embryos

(day 13.5) by microdissection of cortical tissues into Hibernate E (#HECA, Brain Bits, LLC,

Springfield, IL) and processing with the Neural Dissociation Kit with papain (#130-092-628,

Miltenyi, Bergisch Gladbach, Germany) according to the manufacturer's instruction. Single cells

grew into non-adherent neurospheres, which were maintained in culture media (DMEM/F12

(ThermoFisher #10565-018, Waltham, MA), B-27 supplement (#12587-010), N-2 supplement

(#17502-048), MEM non-essential amino acids (#11140-050), 10 mM HEPES (#15630-080),

1000X 2-mercaptoethanol (#21985-023), 100X Pen/Strep (#15140-122)) supplemented with

growth factors (FGF-basic (Biolegend #579606) and EGF (ThermoFisher #PHG0311) to a final

concentration of 20 ng/mL in media. Neurospheres were passaged every six days using the

Neural Dissociation Kit to approximately 1.5 million cells per 10-cm dish and growth factors

were refreshed every 3 days. Cells were authenticated by immunofluorescent staining for Nestin

and GFAP, routinely tested for mycoplasma, and were used for experiments between passages

2 and 20. Dissociated cells were grown in monolayers in 96-well plates pre-coated with poly-DL-

ornithine (SigmaAldrich, #P8638), laminin (SigmaAldrich #11243217001) and fibronectin

(SigmaAldrich #F4759) at 10,000-30,000 cells per well for direct delivery and nucleofection

experiments.

Human induced pluripotent stem cell differentiation into NPCs and culture

MSC-iPSC1 cells were a generous gift from Boston Children's Hospital. iPSCs were differentiated into NPCs based on dual SMAD inhibition as previously described. Briefly, iPSCs were plated onto Matrigel in the presence of 10µM Y-27632 (Sigma #Y0503) at a density of 200,000 cells/cm^2. The next day (day 0) media was changed to KSR media (Knockout DMEM (ThermoFisher #10829018), 15% Knockout serum replacement (ThermoFisher #10828010), L-glutamine (1mM), 1% MEM Non-essential amino acids, and 0.1mM B-mercaptoethanol). Media was changed daily during differentiation and gradually changed from KSR media to N2/B27 media (Neurobasal medium (ThermoFisher #21103049), GlutaMAX Supplement (ThermoFisher #35050061), N-2 supplement (ThermoFisher #17502048) and B-27 supplement (ThermoFisher #17504044)) by increasing N2/B27 media to 1/3 on day 4, 2/3 on day 6 and full N2/B27 media on day 8. For the first 12 days of differentiation media was supplemented with 100nM LDN193189 (Sigma #SML0559) and 10µM SB431542 (Tocris Bioscience #1614, Bristol, England). For the first 4 days media was also supplemented with 2µM XAV939 (Tocris Bioscience #3748). On day 19, NPCs were dissociated with StemPro Accutase (ThermoFisher #A1110501) and replated onto Matrigel for expansion. NPCs were passaged every 6 days and maintained in NPC media (DMEM/F12, N2 supplement, B27 supplement and 20ng/ml bFGF (Corning, #354060, Corning, NY)) with media changes every other day. For direct delivery experiments, 12,000 cells were seeded in Matrigel in a 96-well plate and treated in triplicate with 100pmol of 4xNLS-SpyCas9-2xNLS RNPs with the EMX1 guide RNA (spacer: 5' GAGTCCGAGCAGAAGAAGAA) or non-targeting guide RNA (spacer: 5' AACGACTAGTTAGGCGTGTA). In the Lipofectamine^{TM} CRISPRmax group (ThermoFisher, #CMAX00003), 3µg of 0xNLS-SpyCas9-2xNLS protein (18 pmol) was mixed with sgRNA (1:1 molar ratio) in 8µL OptiMEM with 6µL of Cas9 Plus Reagent (1 µg protein: 2µL reagent) and was mixed with a second tube containing 3.6µL CRISPRmax reagent in 8µL OptiMEM,

incubated for at least 5 minutes and was immediately distributed to cells in triplicate (1µg RNP

per well), according to the manufacturer's recommendations.

Cas9 ribonucleoprotein (RNP) assembly and delivery to cells

For cell culture experiments, RNPs were prepared immediately before use at a 1.2:1

molar ratio of single guide RNA (Synthego, Redwood City, CA) to protein (QB3 Macrolab or

Aldevron). The solution was incubated for 5-10 minutes at room temperature. For nucleofection,

RNPs were formed at 10µM in 10µL of pre-supplemented buffer (Lonza P3 Primary Cell 96-well

Kit, #V4SP-3096). A 15µL suspension of 250,000 mouse NPCs was mixed with 10µL RNP

solution and added to the nucleofection cuvette. Nucleofection was performed using the 4D

Nucleofector X Unit (Lonza, #AAF-1003X) with pulse code EH-100 and cells were recovered

with 75 µL media per well approximately 2 minutes post-nucleofection. Nucleofected cells were

then transferred to 100 µL fresh media in 96-well plates in triplicate and allowed to grow for 5

days at 37°C before analysis by flow cytometry for tdTomato expression. For direct delivery,

RNPs were formed at 10µM in 10µL of sterile Buffer 1 (25 mM sodium phosphate pH (7.25),

100 mM NaCl, 200 mM trehalose). After NPCs were grown for two days in an adherent

monolayer, 10µL was added to cell monolayer ("direct delivery") for a final concentration of 1µM

(100pmol RNP/100µL media). Media was changed 48-hours post-treatment and cells were

collected 5 days post-treatment for analysis by flow cytometry for tdTomato expression or 4

days post-treatment for DNA sequencing.

For *in vivo* experiments, RNPs were prepared similarly at 10µM concentration in Buffer 1

and were incubated at 37°C for 10 minutes. RNPs were sterile filtered by centrifuging through

0.22µm Spin-X cellulose acetate membranes (Corning CoStar, #32119210) at 15,000xg for 1

minute at 4°C. RNPs were then concentrated using 100kDa Ultra-0.5 ml Centrifugal Filter Unit

(Amicon, #, Burlington, MA) at 14,000xg at 4°C until the final desired concentration was reached

(25-100µM, minimum 50µL volume) and collected by centrifuging at 1,000xg for 2 minutes.

RNPs were then divided into single-use 20µL aliquots, flash frozen in liquid nitrogen, and stored

at -80°C until the experiment. Prior to intracranial injection, RNPs were thawed, pipetted to mix,

loaded into a 25µL syringe (Hamilton, #7654-01, Reno, NV) and injected with custom 29-gauge

CED cannulas.

AAV9 transduction

A single 50µL aliquot of AAV9-CMV-SauCas9-U6-sgRNA or AAV9-CMV-GFP (Virovek)

was thawed from -80°C and stored at 4°C. AAV9 was diluted in 1x PBS without calcium or

magnesium to the desired concentration. For *in vivo* experiments, AAV was diluted on the day

of surgeries to 3e8-3e9 vg/µL and stored on ice until loaded into the syringe. 5µL was injected in

each hemisphere to a final dose of 1.5e9-1.5e10 vg per hemisphere using the CED. For cell

culture experiments, serial dilutions were performed from 1.6e9 vg/µL to 2e8 vg/µL (lowest MOI

= 200,000) and 10µL of each were added into 96-well plates in triplicate and maintained for 3

days or 9 days prior to flow cytometry for GFP expression (transduction) and tdTomato

expression (genome editing). AAV9 was added at the same time as NPC seeding for optimal

transduction.

Empty capsid quantification by cryo-electron microscopy

AAV samples were frozen using FEI Vitrobot Mark IV cooled down to 8°C at 100%

humidity. Briefly, 4 µl of AAV9 capsids containing GFP or Cas9 cargo was deposited on 2/2 400

mesh C-flat grids (Electron Microscopy Sciences, Hatfield, PA, #CF224C-50), which were

previously glow discharged at 15 mA for 15 s on PELCO easyGLOW instrument. Grids were

blotted for 3 s with blot force 8 and wait time 2.5 s. Micrographs were collected manually on

Talos Arctica operated at 200 kV and magnification 36,000x (pixel size 1.14 Å/pix) using a

super-resolution camera setting (0.57 Å/pix) on K3 Direct Electron Detector. Micrographs were

collected using SerialEM v. 3.8.7 software. Capids were counted manually by three blinded

reviewers for each image and the three counts were averaged and reported as percentage

empty capsids between the two groups.

RNP size and aggregation measurement by dynamic light scattering (DLS)

RNPs were prepared at 25µM, according to the methods described above for *in vivo*

experiments. Approximately 40µL of RNP was added to a disposable cuvette (ZEN0040) and

inserted into the Zetasizer Nano (ZSP, Malvern, United Kingdom) then 173⁰ scattering angle

was measured at 25⁰C and again at 37⁰C after incubation for 30 minutes. Size distribution by

mass histograms are shown with estimated peak size in nanometers (nm). Each sample was

analyzed 3 times, resulting in a histogram generated from 10 individual measurements, for a

total protocol run time of approximately 10 minutes.

Analysis of editing in vitro

tdTomato positivity was assessed by flow cytometry using IGI facilities on the Attune

NxT (Thermo Fisher, AFC2). Briefly, mouse NPCs were washed once with 1x PBS, harvested

with 0.25% trypsin, neutralized with DMEM containing 10% FBS, and resuspended in 150uL of

1x PBS per well of a round-bottom 96-well plate for analysis. The percentage of tdTomato⁺ cells

from each well was recorded. For analysis of genomic DNA (gDNA), media was removed, cells

were rinsed once with 1X PBS, then incubated with 100 µL QuickExtract solution (Lucigen

CorporationSupplier Diversity Partner QuickExtract DNA Extraction Solution 1.0, Fisher,

#QE09050) at 37°C for 5 minutes. The cell lysate was then moved to a thermal cycler and

incubated at 65°C for 20 minutes and 95°C for 20 minutes. gDNA was used in PCR reactions to

generate amplicons of approximately 150-300bp for Illumina sequencing. A list of primers used

for NGS is provided in Supplementary Table 1. Sequencing was performed with Illumina MiSeq

in the IGI Center for Translational Genomics and reads were analyzed in CRISPResso

(website).

Stereotaxic infusion of Cas9 RNPs and AAVs

Ai9 mice (Jackson Laboratory, #007909, Bar Harbor, ME) were group housed at the

University of California, Berkeley with a 12-hour light-dark cycle and allowed to feed and drink

ad libitum. Housing, maintenance, and experimentation of the mice were carried out with strict

adherence to ethical regulations set forth by the Animal Care and Use Committee (ACUC) at the

University of California, Berkeley. Cas9-RNP and AAVs were prepared on-site at the University

of California, Berkeley for injection into male and female tdTomato Ai9 mice between 2 to 5

months of age. All tools were autoclaved and injected materials were sterile. Mice anesthetized

with 2% isoflurane, given pre-emptive analgesics, and were arranged on Angle-two stereotactic

frame (Leica, Nussloch, Germany). The incision area was swabbed with three alternating wipes

of 70% ethanol and betadine scrub with sterile applicators prior to performing minimally

damaging craniotomies. The stereotaxic surgery coordinates used for targeting the striatum,

relative to bregma, were +0.74 mm anteroposterior, ±1.90 mm mediolateral, –3.37 mm

dorsoventral. Bilateral CED infusion of Cas9 RNPs (10- 100µM) or Cas9 AAVs (3e8-3e9 vg/µL)

was performed with a syringe pump to deliver 5µL at 0.5 µL per minute (Model 310 Plus, KD

Scientific, Holliston, MA) with a step or non-step cannula. For intracerebroventricular (ICV)

infusion of Cas9 RNPs, cannulas were placed at –0.7 mm anteroposterior, +/–1.2 mm

mediolateral, and –2.5 mm dorsoventral according to Paxinos atlas of the adult mouse. Post-

infusion, the syringes were left in position for 2 minutes before slow removal from the injection

site, which was then cleaned, sutured, and surgically glued. Throughout the procedure, mice

were kept at 37°C for warmth and Puralube Vet Ointment (Dechra, NDC #17033-211-38,

Northwich, England) was applied to the outside of the eyes. For ICV injection of p0 neonatal

mice, anesthesia was induced by hypothermia then 4µL of 100µM Cas9-RNP was injected with

a hand-held 33-gauge needle unilaterally with 10% Fast Green dye to visualize distribution from

one ventricle throughout the CNS. The needle was inserted 2 mm deep at a location

approximately 0.25 mm lateral to the sagittal suture and 0.50-0.75 mm rostral to the neonatal

coronal suture. RNP was slowly injected, then the needle was held in place for 15 seconds, and

mice were monitored until recovery. For intrathecal injection, anesthetized mice received a 5,

25, or 50 µL bolus injection of Cas9 RNP at 300µM. The 29-gauge needle was inserted at the

L6-S1 vertebral junction and angled slightly rostrally for the injection. Mice were allowed to fully

recover before being transferred back to their housing. Recovery weight following all procedures

was monitored daily for one week and mice were housed without further disruption for various

time periods until tissue collection.

Tissue collection and immunostaining

At the defined study endpoints (3, 21, and 90-days post-injection), mice were placed

under anesthesia and tissues were perfused with 10mL of cold PBS and 5mL of 4%

paraformaldehyde (PFA, Electron Microscopy Sciences, #15710, Hatfield, PA). Brains were

post-fixed overnight in 4% PFA at 4°C, rinsed 3x with PBS, then cryoprotected in a 10% sucrose

in PBS solution for approximately 3 days. Brains were embedded in optimal cutting temperature

(OCT, Thermo Fisher, #23-730-571) media, and stored at -80°C. Brains were cut at 20-35 µm-

thick sections using a cryostat (Leica CM3050S) and transferred to positively charged

microscope slides. For immunohistochemical analysis, tissues were blocked with solution (0.3%

TritonX-100, 1% bovine serum albumin (SigmaAldrich #A9418), 5% normal goat serum

(SigmaAldrich, #G9023)) before 4°C incubation overnight with primary antibody in blocking

solution. The next day, tissues were washed three times with PBS and incubated with

secondary antibodies for one hour at room temperature. After three PBS washes, samples were

incubated with DAPI solution (0.5 ug/mL, Roche LifeScience, Penzberg, Germany) as a DNA

fluorescence probe for 10 minutes, washed three times with PBS, submerged once in deionized

water, and mounted with glass coverslips in Fluoromount-G slide mounting medium

(SouthernBiotech, Birmingham, AL). Primary antibodies included rabbit polyclonal anti-S100β

(1:500, Abcam, #ab41548, Cambridge, England), rabbit polyclonal anti-Olig-2 (1:250, Millipore

Sigma, #AB9610, Burlington, MA), rabbit polyclonal anti-doublecortin (1:800, Cell Signaling

Technology, #4604, Danvers, MA), rabbit polyclonal anti-Ki67 (1:100, Abcam, #ab15580),

mouse monoclonal anti-NeuN (1:500, Millipore Sigma, #MAB377), rabbit polyclonal anti-

DARPP-32 (1:100, Cell Signaling Technology, #2302), rabbit polyclonal anti-Iba1 (1:100, Wako

Chemicals, #019-19741, Richmond, VA), mouse monoclonal anti-glial fibrillary acidic protein

(1:1000, Millipore Sigma, #MAB3402), rat monoclonal anti-CD45 (1:200, Thermo Fisher, #RA3-

6B2), and rabbit polyclonal anti-CD3 (1:150, Abcam, #ab5690). Secondary antibodies included

donkey anti-rat 488 (1:500, Thermo Fisher, #A-21208), goat anti-rabbit 488 (1:500, Thermo

Fisher, #A32731), goat anti-rabbit 647 (1:500, Thermo Fisher, #A21245), and goat anti-mouse

IgG1 647 (1:500, Thermo Fisher, #A-21240).

Fluorescent imaging and image quantification

Whole brain sections were imaged and stitched using the automated AxioScanZ1 (Zeiss,

Oberkochen, Germany) with a 20x objective in the DAPI and tdTomato channels. Images

generated from slide scanning were viewed in ZenLite software (v 3.6 blue edition) as CZI files.

Images were then exported to FIJI (v1.53q), Imaris (v 9.9.1), or QuPath (v 0.3.2) for

quantification by authors blinded to the sample identity. The area of reflux from CED and blunt

needles was calculated directly in ZenLite using the shape and area analysis tools.

Immunostained cells and tissues were imaged on the Evos Revolve widefield microscope using

a 20x objective or Stellaris 5 confocal microscope (Leica) with a 10x or 25x water-immersion

objective to capture data in DAPI, FITC, tdTomato and CY5 channels. Approximately four

images were taken at 20-25x per hemisphere across multiple sections for image quantification

of CD45, Iba1, and CD3 (8-12 images quantified and averaged per injection). Approximately

four to six z-stack images were captured and stitched per hemisphere for qualitative images of

Iba1 and for quantification of NeuN, DARPP-32, ALDH1L1, and OLIG2 with tdTomato at

1024x1024 pixel resolution with a scanning speed of 100-200.

Measurements of striatal editing by volume were conducted using QuPath software

(version 0.3.2) from images obtained from the Zeiss AxioscanZ1. Briefly, regions of interest

(ROIs) were drawn to outline the border of each striatum and the inner area of tdTomato editing

using the polygon tool to create annotations. All coronal plane areas were automatically

calculated. Dorsoventral coordinates (relative to bregma) were then estimated in millimeters by

consulting the Mouse Brain Atlas (C57BL/6J Coronal). Approximate tissue volume was

calculated by averaging outlined areas between consecutive sections to represent the mean

area across a dorsoventral segment and multiplying by the difference in dorsoventral

coordinates. Edited striatal volumes were then divided by total striatal volumes to obtain percent

editing. Additional tdTomato$^+$ cell count measurements were conducted in Imaris software

version 10.0 (Oxford Instruments, Abingdon, UK). Briefly, ROIs were drawn over each

hemisphere (including cells in all brain sub-structures) using the "Segment only a Region of

Interest" tool, and positive cells detected using the automated "Spots" tool to provide cell counts.

Positivity thresholds were adjusted for each image to accurately capture edited cells manually.

Counts of tdTomato cells on each image were then related back to approximate coordinates

relative to bregma using the Mouse Brain Atlas (C57BL/6J Coronal) to quantify the distribution

of edited cells.

Cell type specific measurements were conducted using QuPath software (version 0.3.2)

on images obtained from Stellaris 5 z-stack maximal projections. ROIs were again drawn

around areas of observed tdTomato editing, using the polygon tool to create a single annotation

per image. Cell count calculations were performed using the "Cell Detection" and "Positive Cell

Detection" tools, adjusting "Cell Mean" thresholds accordingly for each channel and image.

Percent area and intensity measurements were performed in Fiji/ImageJ software (version

2.1.0/1.53c). Images were converted to 32-bit, and thresholds were adjusted to detect the

corresponding stained area. Measurements were set to include area, minimum, maximum, and

mean gray value, and area fraction, as well as to limit to threshold. All image quantification was

performed on 2-5 serial sections with 3-10 independent injections per group for each analysis.

Cell counts, area, intensity, and volume measurements were in general averaged from serial

sections and were then grouped with other biological replicates, including independent

injections, to report the treatment group average with standard deviation displayed by bar graph

or box and whisker plot.

Serum collection and ELISA

Blood was collected from mice at the time of euthanasia, allowed to clot at room

temperature for 15-30 minutes, then centrifuged for 5 minutes at 2,000xg. Sera was collected

and placed immediately on dry ice then stored at -80°C. Enzyme-linked immunosorbent assays

were performed using the SeraCare Protein Detector™ HRP Microwell Anti-Mouse ELISA Kit,

#5110-0011 (54-62-18) according to the manufacturer's recommendations. First, 96-well plates

were coated with antigens of interest (0.5 µg protein per well for SauCas9 and SpyCas9 (4xNLS

protein variants) and approximately 1e9 empty AAV capsids per well) overnight at 4°C. Wells

were washed three times and blocked at room temperature for one hour. Serum samples were

then incubated in wells at varying concentrations (1:50 to 1:10,000 dilution) in 1X blocking buffer

for four hours at room temperature, along with monoclonal antibody controls to generate a

standard curve. Standards included CRISPR/Cas9 Monoclonal Antibody 7A9 (Epigentek, #A-

9000-050, Farmingdale, NY), GenCRISPR□□ SaCas9 Antibody 26H10 (GenScript, #A01952,

Piscataway, NJ), and Anti-Adeno-associated Virus 9 Antibody clone HL2374 (Millipore Sigma,

#MABF2326-25UG). Following three additional washes, the HRP secondary antibody was

added at 1:500 in 1x blocking buffer and incubated for one hour. Wells were then washed three

more times, and peroxidase substrate solutions were added. Absorbance was recorded at a

wavelength of 405nm with Cytation5 plate reader with Gen 5 3.04 software (BioTek). Serum

antibody concentrations were calculated using five-parameter logistic curve (5PL) data analysis

at MyAssays.com and normalized to sham controls.

Splenocyte collection and enzyme linked immunospot (ELISpot)

Spleens were collected at the time of euthanasia and stored in media composed of

RPMI 1640 (ThermoFisher, #11875-119) with 10% FBS (VWR, #89510-186, Radnor, PA) and

1% P/S (ThermoFisher, #15140-122). Briefly, spleens were physically dissociated by forcing

through a 100µm cell strainer in 10mL of media then single cells were passed through a 70µm

strainer and centrifuged at 200xg for 5 minutes. Cells were resuspended in 5mL of 1x RBC

Lysis Buffer (Miltenyi # 130-094-183) for approximately 3 minutes, then centrifuged again and

resuspended in media for counting. The mouse interferon-gamma (IFN-γ) ELISpot kit (R&D

Systems, #EL485, Minneapolis, MN) was used according to the manufacturer's instructions to

assess activation of splenocytes, containing T-cells, in response to treatment with Cas9

proteins. Briefly, the plate was pre-washed and 200µL of media for at least 20 minutes in the

incubator, prior to adding cells at 300,000 per well in 100µL media. Treatments at 2x dose were

prepared in media and 100µL was added to wells in triplicate (final 5 µg/mL concentration).

Plates were wrapped in foil and incubated for 48 hours without disturbing. Concanavalin A

(SigmaAldrich, #C5275) was used as a positive control for cell mediated IFN-γ production (final

4 µg/mL concentration). After 48-hours, cells were removed and the secreted analyte was

detected with immunostaining using the kit-provided biotinylated monoclonal antibody specific

for mouse IFN-γ, streptavidin-conjugated alkaline phosphatase, and stabilized detection mixture

of 5-Bromo-4-Chloro-3'Indolylphosphate-p-Toluidine Salt (BCIP) and Nitro Blue Tetrazolium

Chloride (NBT). After staining, plates were dried overnight and spot forming units were imaged

and counted on the ImmunoSpot S6 Macro Analyzer (Cellular Technology Limited, Shaker

Heights, OH).

Immunization of mice to Cas9 with adjuvant

AddaVax™ (Invivogen, vac-adx-10), a squalene-based oil-in-water nano-emulsion, was

mixed with an equal volume containing 25µg of 4x-SpyCas9-2x protein diluted in sterile buffer at

room temperature for a final injection volume of 50µL. Mice received two 25µL subcutaneous

injections of the AddaVax:Cas9 mixture (immunized) or AddaVax:Buffer alone (sham) with a 30-

gauge insulin syringe into each flank. After four weeks, stereotaxic surgery with bilateral

injections of 5µL of 25µM Cas9-RNPs was performed in a subset of mice. Mice showed no

signs of pain or distress following treatment with AddaVax and no acute events were noted after

surgery. Mice that received AddaVax, with or without surgery, were euthanized 6-weeks post-

subcutaneous injections. Brains, serum, and spleens were collected for analysis of adaptive

immune responses against repeated exposure to Cas9.

DNA/RNA extraction from brain tissue slices and quantitative RT-PCR, droplet digital PCR, and

long-read sequencing

Brains were collected at 3, 14, and 28-days or 4-months for DNA and RNA analysis.

Briefly, mice were placed under anesthesia and perfused with cold PBS. Brains were harvested

and cut into 2-mm sections using a matrix around the injection site (Zivic Instruments,

Pittsburgh, PA). The slices were transferred onto chilled glass slides and further trimmed to

approximate 30mg tissue weight (1−1.25 mm wide × 2 mm long). Tissues were flash frozen in

liquid nitrogen then stored at -80°C until processing. DNA and RNA were collected from tissues

using the AllPrep DNA/RNA Mini Kit (Qiagen, #80204, Venlo, Netherlands) according to the

manufacturer's instructions. Briefly, brains were homogenized in 1.5 mL tubes with a disposable

pestle directly in RLT lysis buffer supplemented with 2-mercaptoethanol, then passed through

Qiashredder columns to further homogenize prior to adding directly to the DNA and RNA

binding columns. DNA was eluted in 100µL of EB, and RNA was eluted in 40 µL RNAse-free

water. Concentrations of nucleic acids were measured by nanodrop spectrophotometer and

samples were stored at -20°C.

Gene expression was quantified across multiple samples using a Custom RT^2 PCR

Array (Qiagen, #330171, CLAM45824) and analyzed using the RT² Profiler PCR Data Analysis

Tool on GeneGlobe (Qiagen). For reverse transcription, the RT^2 First Strand Kit (Qiagen,

#330404) was used according to the manufacturer's instructions. cDNA was diluted in water and

added to the RT² SYBR Green qPCR Mastermix (Qiagen, #330502) then distributed across the

24-wells containing verified assay primers and controls (PCR array reproducibility control,

reverse transcription efficiency control, genomic DNA contamination control, two house-keeping

genes, and 19 experimental genes). Quantitative real-time PCR was performed on the CFX96

Touch Real-Time PCR System (BioRad). cDNA was also used in a droplet digital PCR reaction

to measure SauCas9 expression at the 4-month time-point. qPCR assay IDs are included in

Tables S1 and S2.

DNA was also used for PCR amplicon sequencing of predicted off-target sites and for

droplet digital PCR (ddPCR). Off-target sites were predicted using Cas-OFFinder

(http://www.rgenome.net/cas-offinder/)[61]. Predicted off-targets are described further in Tables

S3 and S4. Primers were designed using NCBI Primer Blast with an amplicon size of 250-

300bp, listed in Table S1. Sequencing was performed with Illumina MiSeq in the IGI Center for

Translational   Genomics   and   reads   were   analyzed   in   CRISPResso2

(http://crispresso.pinellolab.org)[62].

For droplet digital PCR (ddPCR), a custom NHEJ ddPCR assays were generated using

the online Bio-Rad design tool (Table S2). Assays for SauCas9 and SpyCas9 contain both the

primers and probes (HEX-probe spanning the cut-site and a distal reference FAM-probe). To

prepare the reactions, 110 ng of gDNA was combined with the 20x assay, 2x ddPCR Supermix

for Probes (No dUTP), 1µL of of SmaI restriction enzyme (2 units per reaction), and water up to

22µL. Then 20 µl of each reaction mix was added to DG8TM Cartridges (Bio-Rad #1864008,

Hercules, CA) followed by 70 µl of Droplet Generation Oil for Probes (Bio-Rad #1863005) and

droplets were formed in the QX200 Droplet Generator. Droplets were then transferred to a 96-

well plate and thermal cycled according to the manufacturer's recommendation with a 3-minute

annealing/extension step. After thermal cycling, the sealed plate was placed in the QX200

Droplet Reader and data was acquired and analyzed in the QuantaSoft Analysis Pro Software

using the "Drop-Off" analysis, manually setting the thresholds for cluster calling (FAM+ only,

FAM+ HEX+ cluster, FAM-HEX- cluster), and exporting fractional abundance calculations.

Long-read sequencing of the tdTomato locus was performed on DNA isolated from the

treated mouse brains. Briefly, PCR amplicons were generated on 16 samples from the 4-month

treatment groups using primers with unique barcodes for sample de-multiplexing. The KAPA

HiFi Hotstart PCR Kit (Roche, KK2502) was used to amplify the 1100 bp product and reactions

were cleaned with AMPure XP magnetic beads (Beckman Coulter Inc., Brea, CA) prior to

analysis by Qubit and Bioanalyzer with the DNA 7500 Kit (Agilent, #5067-1506, Santa Clara,

CA). Samples were combined and 1µg of pooled amplicons and submitted (> 20ng/µL) for

sequencing with one PacBio Sequel 8M SMRT Cell at the QB3 Vincent Coates Genomic

Sequencing Lab, yielding approximately 110,000 reads per sample. Data were analyzed using a

custom pipeline to identify viral fragment trapping during DNA repair. Briefly, PacBio circular

consensus reads were trimmed with Cutadapt (Version 4.1)[63], then aligned to the AAV vector

using NGMLR (Version 0.2.7)[64] to generate BAM files. Soft-clipped regions of aligned reads

were extracted using PySam (Version 0.18.0, https://github.com/pysam-developers/pysam) to

parse CIGAR strings, then realigned to the tdTomato locus with NGMLR to verify integration

within 200 bp of the cut site. Confirmed integrations were visualized along the AAV genome

using pyGenomeTracks (Version 3.3) and coverage statistics were summarized using

PySam[65,66].

Statistical analyses

The data presented in bar graphs and box and whisker plots are averages across

multiple technical and biological replicates and error bars represent the standard deviation.

Sample sizes are indicated in the text and figure legends and generally refer to technical

injection replicates (two technical replicates, i.e., bilateral injections, per one biological

replicate). When comparing two groups with normal distribution, an unpaired student's t-test

was performed in Prism 9 (GraphPad Software version 9.4.1). When comparing multiple groups,

a one-way ANOVA with Tukey's multiple comparison test was performed in Prism 9 (GraphPad

Software version 9.4.1). The RT-qPCR experiments used Student's t-test of the experimental

group compared to the sham control (Qiagen GeneGlobe RT^2 Profiler PCR Data Analysis). $p \leq$

0.05 was considered significant

Data Availability Statement

Long-read sequencing (BAM files from PacBio circular consensus sequence, CCS) are

available in Sequence Read Archive (SRA). Accession number: [#]. All additional data is

available upon request.

Acknowledgements

Thank you to Netravathi Krishnappa and Christopher Hann-Soden for library preparation,

sequencing, and analysis at the IGI Center for Translational Genomics and the QB3 Vincent J.

Coates Genomics Sequencing Lab. Thank you to members of the Doudna lab at IGI, particularly

Drs. Matthew Kan, Jennifer Hamilton, Cole Urnes, and Talia Wenger, as well as Brett Staahl,

Ross Wilson, and Fyodor Urnov for intellectual contributions and support. Thank you to

Aldevron, LLC for collaboration on this effort, especially Samantha Foti, Allison Pappas, David

Yoder, and Max Sellman. Thank you to Denise Schichnes and Steven Ruzin at the CNR

Berkeley Imaging Facility at the University of California, Berkeley as well as Feather Ives, Anita

Flynn, and Holly Aaron at the CRL Microscopy Imaging Core (RRID:SCR_017852) at the

University of California, Berkeley. Thank you to Eva Harris and lab members for the use of the

ELISpot plate reader. Thank you to Kathy Snow and Ethan Saville at Jackson Laboratories for

discussion about QuPath image quantification software. Thank you to the Bankiewicz lab, in

particular Drs. Victor Van Laar, Lluis Samaranch, for discussion about CED. Thank you to Drs.

Greg Barton and David Raulet for intellectual discussions regarding endotoxins and immune

response. Thank you to Biogen Inc., especially Robin Kleinmen and Anirvan Ghosh, for funding

and intellectual contributions. Thank you to Ruth L. Kirschstein F32 NIGMS for funding (E.C.S).

Author Contributions

E.C.S performed the experiments, analysis, and manuscript preparation and oversaw

contributions from researchers (R.A., E. A., S.E.K., A.S., N.L., Y.K.) who assisted with tissue

processing, cloning, ELISAs, immunohistochemistry, and image analysis. E.C.S., J.K.S., and

M.H.K. designed *in vivo* experiments, performed stereotaxic surgeries, confocal microscopy,

and quantitative PCR. M.T. performed long-read NGS sequencing analysis. E.C.S. performed

isolation and culture of mouse primary cells, flow cytometry, and ELISpot assays. K.S.

performed electron microscopy. V.S.R. and L.T.V. performed human stem cell culture and

differentiation. C.J., A.W., T.M., A.K., and T.F. performed custom low-endotoxin protein

expression and purification. D.F.S. and J.A.D. approved the experiments, provided intellectual

contributions, and co-wrote the manuscript.

Declaration of Interests Statement

J.A.D. is a cofounder of Caribou Biosciences, Editas Medicine, Scribe Therapeutics, Intellia

Therapeutics and Mammoth Biosciences. J.A.D. is a scientific advisory board member of

Vertex, Caribou Biosciences, Intellia Therapeutics, eFFECTOR Therapeutics, Scribe

Therapeutics, Mammoth Biosciences, Synthego, Algen Biotechnologies, Felix Biosciences, The

Column Group and Inari. J.A.D. is a Director at Johnson & Johnson and Tempus and has

research projects sponsored by Biogen, Pfizer, Apple Tree Partners and Roche. Patent

applications have been filed relating to the technologies described herein. The indicated authors

are employees of Aldevron, LLC, which offers proteins, pDNA, mRNA and reagents for sale

similar to some of the compounds described in this manuscript.

Keywords (5-10)

CRISPR-Cas9, Genome Editing, Viral Vectors, Non-viral Delivery, Mouse, Brain, Host Immune

Response, Neurons, Microglia, Endotoxin/LPS

936
937
938

References

939
940
1. Heidenreich, M., and Zhang, F. (2016). Applications of CRISPR–Cas systems in neuroscience.
Nat Rev Neurosci *17*, 36–44. 10.1038/nrn.2015.2.

2. The SCGE Consortium, Saha, K., Sontheimer, E.J., Brooks, P.J., Dwinell, M.R., Gersbach, C.A.,
Liu, D.R., Murray, S.A., Tsai, S.Q., Wilson, R.C., et al. (2021). The NIH Somatic Cell Genome
Editing program. Nature *592*, 195–204. 10.1038/s41586-021-03191-1.

3. Jinek, M., Chylinski, K., Fonfara, I., Hauer, M., Doudna, J.A., and Charpentier, E. (2012). A
Programmable Dual-RNA–Guided DNA Endonuclease in Adaptive Bacterial Immunity.
Science *337*, 816–821. 10.1126/science.1225829.

4. Jinek, M., East, A., Cheng, A., Lin, S., Ma, E., and Doudna, J. (2013). RNA-programmed
genome editing in human cells. eLife *2*, e00471. 10.7554/eLife.00471.

5. Doudna, J.A. (2020). The promise and challenge of therapeutic genome editing. Nature *578*,
229–236. 10.1038/s41586-020-1978-5.

6. Hadaczek, P., Forsayeth, J., Mirek, H., Munson, K., Bringas, J., Pivirotto, P., L McBride, J.,
Davidson, B.L., and Bankiewicz, K.S. (2009). Transduction of Nonhuman Primate Brain with
Adeno-Associated Virus Serotype 1: Vector Trafficking and Immune Response. Human Gene
Therapy *20*, 225–237. 10.1089/hum.2008.151.

7. Ciesielska, A., Hadaczek, P., Mittermeyer, G., Zhou, S., Wright, J.F., Bankiewicz, K.S., and
Forsayeth, J. (2013). Cerebral Infusion of AAV9 Vector-encoding Non-self Proteins Can Elicit
Cell-mediated Immune Responses. Molecular Therapy *21*, 158–166. 10.1038/mt.2012.167.

8. Samaranch, L., Sebastian, W.S., Kells, A.P., Salegio, E.A., Heller, G., Bringas, J.R., Pivirotto, P.,
DeArmond, S., Forsayeth, J., and Bankiewicz, K.S. (2014). AAV9-mediated Expression of a
Non-self Protein in Nonhuman Primate Central Nervous System Triggers Widespread
Neuroinflammation Driven by Antigen-presenting Cell Transduction. Molecular Therapy *22*,
329–337. 10.1038/mt.2013.266.

9. Agustín-Pavón, C., Mielcarek, M., Garriga-Canut, M., and Isalan, M. (2016). Deimmunization
for gene therapy: host matching of synthetic zinc finger constructs enables long-term
mutant Huntingtin repression in mice. Mol Neurodegeneration *11*, 64. 10.1186/s13024-
016-0128-x.

10. Chew, W.L., Tabebordbar, M., Cheng, J.K.W., Mali, P., Wu, E.Y., Ng, A.H.M., Zhu, K., Wagers,
A.J., and Church, G.M. (2016). A multifunctional AAV–CRISPR–Cas9 and its host response.
Nat Methods *13*, 868–874. 10.1038/nmeth.3993.

11. Moreno, A.M., Palmer, N., Alemán, F., Chen, G., Pla, A., Jiang, N., Chew, W.L., Law, M., and
Mali, P. (2019). immune-orthogonal orthologues of AAV capsids and of Cas9 circumvent the

immune response to the administration of gene therapy. Nat Biomed Eng *3*, 806–816.
10.1038/s41551-019-0431-2.

12. Li, A., Tanner, M.R., Lee, C.M., Hurley, A.E., De Giorgi, M., Jarrett, K.E., Davis, T.H., Doerfler,
A.M., Bao, G., Beeton, C., et al. (2020). AAV-CRISPR Gene Editing Is Negated by Pre-existing
Immunity to Cas9. Molecular Therapy *28*, 1432–1441. 10.1016/j.ymthe.2020.04.017.

13. Simhadri, V.L., McGill, J., McMahon, S., Wang, J., Jiang, H., and Sauna, Z.E. (2018).
Prevalence of Pre-existing Antibodies to CRISPR-Associated Nuclease Cas9 in the USA
Population. Molecular Therapy - Methods & Clinical Development *10*, 105–112.
10.1016/j.omtm.2018.06.006.

14. Charlesworth, C.T., Deshpande, P.S., Dever, D.P., Camarena, J., Lemgart, V.T., Cromer, M.K.,
Vakulskas, C.A., Collingwood, M.A., Zhang, L., Bode, N.M., et al. (2019). Identification of
preexisting adaptive immunity to Cas9 proteins in humans. Nat Med *25*, 249–254.
10.1038/s41591-018-0326-x.

15. Wagner, D.L., Amini, L., Wendering, D.J., Burkhardt, L.-M., Akyüz, L., Reinke, P., Volk, H.-D.,
and Schmueck-Henneresse, M. (2019). High prevalence of Streptococcus pyogenes Cas9-
reactive T cells within the adult human population. Nat Med *25*, 242–248. 10.1038/s41591-
018-0204-6.

16. Ferdosi, S.R., Ewaisha, R., Moghadam, F., Krishna, S., Park, J.G., Ebrahimkhani, M.R., Kiani,
S., and Anderson, K.S. (2019). Multifunctional CRISPR-Cas9 with engineered
immunosilenced human T cell epitopes. Nat Commun *10*, 1842. 10.1038/s41467-019-
09693-x.

17. Tang, X.-Z.E., Tan, S.X., Hoon, S., and Yeo, G.W. (2022). Pre-existing adaptive immunity to
the RNA-editing enzyme Cas13d in humans. Nat Med *28*, 1372–1376. 10.1038/s41591-022-
01848-6.

18. Mingozzi, F., and High, K.A. (2013). Immune responses to AAV vectors: overcoming barriers
to successful gene therapy. Blood *122*, 23–36. 10.1182/blood-2013-01-306647.

19. Staahl, B.T., Benekareddy, M., Coulon-Bainier, C., Banfal, A.A., Floor, S.N., Sabo, J.K., Urnes,
C., Munares, G.A., Ghosh, A., and Doudna, J.A. (2017). Efficient genome editing in the
mouse brain by local delivery of engineered Cas9 ribonucleoprotein complexes. Nat
Biotechnol *35*, 431–434. 10.1038/nbt.3806.

20. Liu, J., Gaj, T., Wallen, M.C., and Barbas, C.F. (2015). Improved Cell-Penetrating Zinc-Finger
Nuclease Proteins for Precision Genome Engineering. Molecular Therapy - Nucleic Acids *4*,
e232. 10.1038/mtna.2015.6.

21. Madisen, L., Zwingman, T.A., Sunkin, S.M., Oh, S.W., Zariwala, H.A., Gu, H., Ng, L.L.,
Palmiter, R.D., Hawrylycz, M.J., Jones, A.R., et al. (2010). A robust and high-throughput Cre

reporting and characterization system for the whole mouse brain. Nat Neurosci *13*, 133–
140. 10.1038/nn.2467.

22. Ran, F.A., Cong, L., Yan, W.X., Scott, D.A., Gootenberg, J.S., Kriz, A.J., Zetsche, B., Shalem, O.,
Wu, X., Makarova, K.S., et al. (2015). In vivo genome editing using Staphylococcus aureus
Cas9. Nature *520*, 186–191. 10.1038/nature14299.

23. Maeder, M.L., Stefanidakis, M., Wilson, C.J., Baral, R., Barrera, L.A., Bounoutas, G.S.,
Bumcrot, D., Chao, H., Ciulla, D.M., DaSilva, J.A., et al. (2019). Development of a gene-
editing approach to restore vision loss in Leber congenital amaurosis type 10. Nat Med *25*,
229–233. 10.1038/s41591-018-0327-9.

24. Yin, C., Zhang, T., Qu, X., Zhang, Y., Putatunda, R., Xiao, X., Li, F., Xiao, W., Zhao, H., Dai, S.,
et al. (2017). In Vivo Excision of HIV-1 Provirus by saCas9 and Multiplex Single-Guide RNAs
in Animal Models. Molecular Therapy *25*, 1168–1186. 10.1016/j.ymthe.2017.03.012.

25. Gao, G., Vandenberghe, L.H., Alvira, M.R., Lu, Y., Calcedo, R., Zhou, X., and Wilson, J.M.
(2004). Clades of Adeno-Associated Viruses Are Widely Disseminated in Human Tissues. J
Virol *78*, 6381–6388. 10.1128/JVI.78.12.6381-6388.2004.

26. Smith, R.H., Levy, J.R., and Kotin, R.M. (2009). A Simplified Baculovirus-AAV Expression
Vector System Coupled With One-step Affinity Purification Yields High-titer rAAV Stocks
From Insect Cells. Molecular Therapy *17*, 1888–1896. 10.1038/mt.2009.128.

27. Chen, C., Akerstrom, V., Baus, J., Lan, M.S., and Breslin, M.B. (2013). Comparative analysis
of the transduction efficiency of five adeno associated virus serotypes and VSV-G
pseudotype lentiviral vector in lung cancer cells. Virol J *10*, 86. 10.1186/1743-422X-10-86.

28. Gaj, T., Staahl, B.T., Rodrigues, G.M.C., Limsirichai, P., Ekman, F.K., Doudna, J.A., and
Schaffer, D.V. (2017). Targeted gene knock-in by homology-directed genome editing using
Cas9 ribonucleoprotein and AAV donor delivery. Nucleic Acids Research *45*, e98–e98.
10.1093/nar/gkx154.

29. Park, I.-H., Zhao, R., West, J.A., Yabuuchi, A., Huo, H., Ince, T.A., Lerou, P.H., Lensch, M.W.,
and Daley, G.Q. (2008). Reprogramming of human somatic cells to pluripotency with
defined factors. Nature *451*, 141–146. 10.1038/nature06534.

30. Qi, Y., Zhang, X.-J., Renier, N., Wu, Z., Atkin, T., Sun, Z., Ozair, M.Z., Tchieu, J., Zimmer, B.,
Fattahi, F., et al. (2017). Combined small-molecule inhibition accelerates the derivation of
functional cortical neurons from human pluripotent stem cells. Nat Biotechnol *35*, 154–163.
10.1038/nbt.3777.

31. Chambers, S.M., Fasano, C.A., Papapetrou, E.P., Tomishima, M., Sadelain, M., and Studer, L.
(2009). Highly efficient neural conversion of human ES and iPS cells by dual inhibition of
SMAD signaling. Nat Biotechnol *27*, 275–280. 10.1038/nbt.1529.

32. Fu, Y., Foden, J.A., Khayter, C., Maeder, M.L., Reyon, D., Joung, J.K., and Sander, J.D. (2013).
High-frequency off-target mutagenesis induced by CRISPR-Cas nucleases in human cells.
Nat Biotechnol *31*, 822–826. 10.1038/nbt.2623.

33. Bobo, R.H., Laske, D.W., Akbasak, A., Morrison, P.F., Dedrick, R.L., and Oldfield, E.H. (1994).
Convection-enhanced delivery of macromolecules in the brain. Proc. Natl. Acad. Sci. U.S.A.
91, 2076–2080. 10.1073/pnas.91.6.2076.

34. Hadaczek, P., Kohutnicka, M., Krauze, M.T., Bringas, J., Pivirotto, P., Cunningham, J., and
Bankiewicz, K. (2006). Convection-Enhanced Delivery of Adeno-Associated Virus Type 2
(AAV2) into the Striatum and Transport of AAV2 Within Monkey Brain. Human Gene
Therapy *17*, 291–302. 10.1089/hum.2006.17.291.

35. Krauze, M.T., Saito, R., Noble, C., Tamas, M., Bringas, J., Park, J.W., Berger, M.S., and
Bankiewicz, K. (2005). Reflux-free cannula for convection-enhanced high-speed delivery of
therapeutic agents: Technical note. Journal of Neurosurgery *103*, 923–929.
10.3171/jns.2005.103.5.0923.

36. Aschauer, D.F., Kreuz, S., and Rumpel, S. (2013). Analysis of Transduction Efficiency,
Tropism and Axonal Transport of AAV Serotypes 1, 2, 5, 6, 8 and 9 in the Mouse Brain. PLoS
ONE *8*, e76310. 10.1371/journal.pone.0076310.

37. Hanlon, K.S., Kleinstiver, B.P., Garcia, S.P., Zaborowski, M.P., Volak, A., Spirig, S.E., Muller,
A., Sousa, A.A., Tsai, S.Q., Bengtsson, N.E., et al. (2019). High levels of AAV vector
integration into CRISPR-induced DNA breaks. Nat Commun *10*, 4439. 10.1038/s41467-019-
12449-2.

38. Nelson, C.E., Wu, Y., Gemberling, M.P., Oliver, M.L., Waller, M.A., Bohning, J.D., Robinson-
Hamm, J.N., Bulaklak, K., Castellanos Rivera, R.M., Collier, J.H., et al. (2019). Long-term
evaluation of AAV-CRISPR genome editing for Duchenne muscular dystrophy. Nat Med *25*,
427–432. 10.1038/s41591-019-0344-3.

39. Ferrari, S., Jacob, A., Cesana, D., Laugel, M., Beretta, S., Varesi, A., Unali, G., Conti, A.,
Canarutto, D., Albano, L., et al. (2022). Choice of template delivery mitigates the genotoxic
risk and adverse impact of editing in human hematopoietic stem cells. Cell Stem Cell *29*,
1428-1444.e9. 10.1016/j.stem.2022.09.001.

40. Simpson, B.P., Yrigollen, C.M., Izda, A., and Davidson, B.L. (2023). Targeted long-read
sequencing captures CRISPR editing and AAV integration outcomes in brain. Molecular
Therapy, S1525001623000047. 10.1016/j.ymthe.2023.01.004.

41. Zink McCullough, K. Calculating Endotoxin Limits for Drug Products (American
Pharmaceutical Review).

42. Lee, B., Lee, K., Panda, S., Gonzales-Rojas, R., Chong, A., Bugay, V., Park, H.M., Brenner, R.,
Murthy, N., and Lee, H.Y. (2018). Nanoparticle delivery of CRISPR into the brain rescues a

mouse model of fragile X syndrome from exaggerated repetitive behaviours. Nat Biomed
Eng *2*, 497–507. 10.1038/s41551-018-0252-8.

43. Park, H., Oh, J., Shim, G., Cho, B., Chang, Y., Kim, S., Baek, S., Kim, H., Shin, J., Choi, H., et al.
(2019). In vivo neuronal gene editing via CRISPR–Cas9 amphiphilic nanocomplexes alleviates
deficits in mouse models of Alzheimer's disease. Nat Neurosci *22*, 524–528.
10.1038/s41593-019-0352-0.

44. Metzger, J.M., Wang, Y., Neuman, S.S., Snow, K.J., Murray, S.A., Lutz, C.M., Bondarenko, V.,
Felton, J., Gimse, K., Xie, R., et al. (2023). Efficient in vivo neuronal genome editing in the
mouse brain using nanocapsules containing CRISPR-Cas9 ribonucleoproteins. Biomaterials
293, 121959. 10.1016/j.biomaterials.2022.121959.

45. Wang, Y., Wang, X., Xie, R., Burger, J.C., Tong, Y., and Gong, S. (2022). Overcoming the
Blood–Brain Barrier for Gene Therapy via Systemic Administration of GSH-Responsive Silica
Nanocapsules. Advanced Materials, 2208018. 10.1002/adma.202208018.

46. Yan, S., Zheng, X., Lin, Y., Li, C., Liu, Z., Li, J., Tu, Z., Zhao, Y., Huang, C., Chen, Y., et al. (2023).
Cas9-mediated replacement of expanded CAG repeats in a pig model of Huntington's
disease. Nat. Biomed. Eng. 10.1038/s41551-023-01007-3.

47. He, M., Li, J., Han, H., Borges, C.A., Neiman, G., Røise, J.J., Hadaczek, P., Mendonsa, R.,
Holm, V.R., Wilson, R.C., et al. (2020). A traceless linker for aliphatic amines that rapidly and
quantitatively fragments after reduction. Chem. Sci. *11*, 8973–8980. 10.1039/D0SC00929F.

48. Saito, R., Krauze, M.T., Noble, C.O., Tamas, M., Drummond, D.C., Kirpotin, D.B., Berger,
M.S., Park, J.W., and Bankiewicz, K.S. (2006). Tissue affinity of the infusate affects the
distribution volume during convection-enhanced delivery into rodent brains: Implications
for local drug delivery. Journal of Neuroscience Methods *154*, 225–232.
10.1016/j.jneumeth.2005.12.027.

49. Wei, T., Cheng, Q., Min, Y.-L., Olson, E.N., and Siegwart, D.J. (2020). Systemic nanoparticle
delivery of CRISPR-Cas9 ribonucleoproteins for effective tissue specific genome editing. Nat
Commun *11*, 3232. 10.1038/s41467-020-17029-3.

50. Oura, S., Noda, T., Morimura, N., Hitoshi, S., Nishimasu, H., Nagai, Y., Nureki, O., and Ikawa,
M. (2021). Precise CAG repeat contraction in a Huntington's Disease mouse model is
enabled by gene editing with SpCas9-NG. Commun Biol *4*, 771. 10.1038/s42003-021-02304-
w.

51. Sweeney, M.D., Sagare, A.P., and Zlokovic, B.V. (2018). Blood–brain barrier breakdown in
Alzheimer disease and other neurodegenerative disorders. Nat Rev Neurol *14*, 133–150.
10.1038/nrneurol.2017.188.

52. Huang, L., Bommireddy, R., Munoz, L.E., Guin, R.N., Wei, C., Ruggieri, A., Menon, A.P., Li, X.,
Shanmugam, M., Owonikoko, T.K., et al. (2021). Expression of tdTomato and luciferase in a

murine lung cancer alters the growth and immune microenvironment of the tumor. PLoS
ONE *16*, e0254125. 10.1371/journal.pone.0254125.

53. Tsai, S.Q., Zheng, Z., Nguyen, N.T., Liebers, M., Topkar, V.V., Thapar, V., Wyvekens, N.,
Khayter, C., Iafrate, A.J., Le, L.P., et al. (2015). GUIDE-seq enables genome-wide profiling of
off-target cleavage by CRISPR-Cas nucleases. Nat Biotechnol *33*, 187–197.
10.1038/nbt.3117.

54. Tsai, S.Q., Nguyen, N.T., Malagon-Lopez, J., Topkar, V.V., Aryee, M.J., and Joung, J.K. (2017).
CIRCLE-seq: a highly sensitive in vitro screen for genome-wide CRISPR–Cas9 nuclease off-
targets. Nat Methods *14*, 607–614. 10.1038/nmeth.4278.

55. Ibraheim, R., Tai, P.W.L., Mir, A., Javeed, N., Wang, J., Rodríguez, T.C., Namkung, S., Nelson,
S., Khokhar, E.S., Mintzer, E., et al. (2021). Self-inactivating, all-in-one AAV vectors for
precision Cas9 genome editing via homology-directed repair in vivo. Nat Commun *12*, 6267.
10.1038/s41467-021-26518-y.

56. Li, A., Lee, C.M., Hurley, A.E., Jarrett, K.E., De Giorgi, M., Lu, W., Balderrama, K.S., Doerfler,
A.M., Deshmukh, H., Ray, A., et al. (2019). A Self-Deleting AAV-CRISPR System for In Vivo
Genome Editing. Mol Ther Methods Clin Dev *12*, 111–122. 10.1016/j.omtm.2018.11.009.

57. Merienne, N., Vachey, G., de Longprez, L., Meunier, C., Zimmer, V., Perriard, G., Canales, M.,
Mathias, A., Herrgott, L., Beltraminelli, T., et al. (2017). The Self-Inactivating KamiCas9
System for the Editing of CNS Disease Genes. Cell Rep *20*, 2980–2991.
10.1016/j.celrep.2017.08.075.

58. Ewaisha, R., and Anderson, K.S. (2023). Immunogenicity of CRISPR therapeutics—Critical
considerations for clinical translation. Front. Bioeng. Biotechnol. *11*, 1138596.
10.3389/fbioe.2023.1138596.

59. Mancuso, P., Chen, C., Kaminski, R., Gordon, J., Liao, S., Robinson, J.A., Smith, M.D., Liu, H.,
Sariyer, I.K., Sariyer, R., et al. (2020). CRISPR based editing of SIV proviral DNA in ART
treated non-human primates. Nat Commun *11*, 6065. 10.1038/s41467-020-19821-7.

60. Harrington, L.B., Paez-Espino, D., Staahl, B.T., Chen, J.S., Ma, E., Kyrpides, N.C., and Doudna,
J.A. (2017). A thermostable Cas9 with increased lifetime in human plasma. Nat Commun *8*,
1424. 10.1038/s41467-017-01408-4.

61. Bae, S., Park, J., and Kim, J.-S. (2014). Cas-OFFinder: a fast and versatile algorithm that
searches for potential off-target sites of Cas9 RNA-guided endonucleases. Bioinformatics
30, 1473–1475. 10.1093/bioinformatics/btu048.

62. Clement, K., Rees, H., Canver, M.C., Gehrke, J.M., Farouni, R., Hsu, J.Y., Cole, M.A., Liu, D.R.,
Joung, J.K., Bauer, D.E., et al. (2019). CRISPResso2 provides accurate and rapid genome
editing sequence analysis. Nat Biotechnol *37*, 224–226. 10.1038/s41587-019-0032-3.

63. Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing
reads. EMBnet j. *17*, 10. 10.14806/ej.17.1.200.

64. Sedlazeck, F.J., Rescheneder, P., Smolka, M., Fang, H., Nattestad, M., von Haeseler, A., and
Schatz, M.C. (2018). Accurate detection of complex structural variations using single-
molecule sequencing. Nat Methods *15*, 461–468. 10.1038/s41592-018-0001-7.

65. Ramírez, F., Bhardwaj, V., Arrigoni, L., Lam, K.C., Grüning, B.A., Villaveces, J., Habermann, B.,
Akhtar, A., and Manke, T. (2018). High-resolution TADs reveal DNA sequences underlying
genome organization in flies. Nat Commun *9*, 189. 10.1038/s41467-017-02525-w.

66. Lopez-Delisle, L., Rabbani, L., Wolff, J., Bhardwaj, V., Backofen, R., Grüning, B., Ramírez, F.,
and Manke, T. (2021). pyGenomeTracks: reproducible plots for multivariate genomic
datasets. Bioinformatics *37*, 422–423. 10.1093/bioinformatics/btaa692.

1162

1163

1164   **Figure Legends**

1165

Figure 1. In vivo editing at tdTomato locus with viral and non-viral Cas9 delivery

strategies. (A) Schematic of 4x-SpyCas9-2x cell-penetrating protein expression and purification

systems, (B) AAV9-SauCas9-sgRNA expression and purification systems, (C) and expected

edited brain regions in the partial basal ganglia direct circuit shown in sagittal view (top) and

coronal view (bottom). Neurons extend from the striatum into the globus pallidus (GP) and

substantia nigra (SNr). Created with BioRender.com. (D) Comparison of convection enhanced

delivery (CED) of cell-penetrant 4x-SpyCas9-2x RNP with step and non-step cannulas. The

step-cannula significantly reduced reflux of material in the needle injection track (n=3-6

injections per group, unpaired t-test, ** $p<0.01$.) Scale bar: 1 mm. (E) Serial sections of single

hemisphere sagittal view of edited tdTomato+ cells in the basal ganglia circuit after injection of

Cas9 RNP with CED into the striatum, with signal detected near the GP and SNr. Scale bar: 1

mm. (F) Representative coronal section of the striatum of mice that received Cas9 RNPs and

AAVs at 21 days post-injection, showing the distribution of tdTomato+ edited cells. Scale bar: 1

mm. (G) Co-staining of tdTomato with NeuN and GFAP in the striatum at 90-days post-injection.

Scale bar: 50 μm. (H) Volume of edited striatal tissue as the concentration of injected Cas9

RNPs was increased from 10 to 100 μM (n=4-6 injections, one-way ANOVA, ns). (I)

Quantification of editing following treatment with Cas9 AAV (3e9 vg/ μL, 1.5e10 vg/hemisphere)

and Cas9 RNPs (25 μM, 125pmol/hemisphere) at 21 and 90-days (n=4-6 injections, one-way

ANOVA, * $p< 0.05$). (J) Co-localization of tdTomato and NeuN quantified per regions of interest

(ROI), e.g., edited area per hemisphere (n=4-6 injections, one-way ANOVA, ** $p< 0.01$). Scale

bar: 250 μm.

1187

Figure 2. Immune response following in vivo editing with viral and non-viral Cas9

delivery strategies. (A) Representative immunostaining of Iba1 (microglia, green) with

tdTomato and DAPI using confocal microscopy. Scale bar: 50 μm. (B) Quantification of Iba1$^+$

staining intensity and percent area (n=4-6 technical replicates, one-way ANOVA, *p<0.05). (C)

Quantification of CD45$^+$ and CD3$^+$ cells per image (n=3-6 replicates, one-way ANOVA, ns). (D)

Representative images of CD45, CD3, and Iba1 showing co-expression of CD45 (green) with

both Iba1 (microglia, red) and CD3 (T-cells, red) cells and differential cell morphology. Merged

images include DAPI (gray) and tdTomato (magenta). Scale bar: 50 μm. (E) Quantification of

IgG antibodies against Cas9 or AAV capsid proteins measured 28 and 90-days after bilateral

intrastriatal injections by ELISA (n=3-5 biological replicates). (F) Heat map summarizing RT-

qPCR results of gene expression from homogenized brain tissue near the injection site (striatum

and cortex) at two time-points. *Ppih* was used as a housekeeping control for delta-delta Ct

analysis and compared to the sham group using Qiagen analysis portal (n=4, *p < 0.05).

1201

Figure 3. Optimized, low-endotoxin RNP formulation reduces local immune response. (A)

Schematic of manufacturing scale up to produce industrial ultra-low endotoxin 4x-SpyCas9-2x

protein using a tag-free expression and purification system. (B) Endotoxin levels calculated on a

per mouse basis between the standard (laboratory 4x-SpyCas9-2x with sg298 2018) and

optimized (industrial 4x-SpyCas9-2x protein with sg298 2022) RNP formulations at 25μM

measured by LAL assay. The optimized RNP had a final endotoxin level of 0.44 EU/kg. Dotted

line: FDA recommendation of 0.2 EU/kg/hr for drug products administered intrathecally in

humans. (C) Quantification of Iba1$^+$ staining intensity and percent area (n=6-10, one-way

ANOVA, *p<0.05). (D) Quantification of CD45$^+$ and (E) CD3$^+$ cells per image (n=6-10, one-way

ANOVA, ns). (F) Percent volume of edited striatal tissue for Cas9 RNPs injected at 25μM (n=6-

10 injections). (G) Quantification of IgG antibodies against Cas9 or AAV capsid proteins

measured 21-days after bilateral intrastriatal injections by ELISA (n=3-5 biological replicates).

1214

Figure 1. In vivo editing at tdTomato locus with viral and non-viral Cas9 delivery strategies

Figure 2. Immune response to in vivo editing with viral and non-viral Cas9 delivery strategies

Figure 3. Optimized, low-endotoxin RNP formulation reduces local immune response