Longitudinal Changes in Value-based Learning in Middle Childhood: Distinct Contributions of Hippocampus and Striatum

Johannes Falck[1], Lei Zhang[2,3,4], Laurel Raffington[5], Johannes J. Mohn[6,7], Jochen Triesch[8], Christine Heim[6,9] & Yee Lee Shing[1]

[1]*Department of Psychology, Goethe University Frankfurt, 60629 Frankfurt am Main, Germany*

[2]*Social, Cognitive and Affective Neuroscience Unit, Department of Cognition, Emotion, and Methods in Psychology, Faculty of Psychology, University of Vienna, 1010 Vienna, Austria*

[3]*Centre for Human Brain Health, School of Psychology, University of Birmingham, Birmingham B15 2TT, UK*

[4]*Institute for Mental Health, School of Psychology, University of Birmingham, Birmingham B15 2TT, UK*

[5]*Center for Lifespan Psychology, Max Planck Institute for Human Development, 14195 Berlin, Germany*

[6]*Charité – Universitätsmedizin Berlin, Institute of Medical Psychology, 10117 Berlin, Germany*

[7]*Max Planck School of Cognition, Max Planck Institute for Human Cognitive and Brain Sciences, 04103 Leipzig, Germany*

[8]*Frankfurt Institute for Advanced Studies (FIAS), 60439 Frankfurt am Main, Germany*

[9]*Center for Safe & Healthy Children, The Pennsylvania State University, State College, PA 16802, USA*

Abstract

The hippocampal-dependent memory system and striatal-dependent memory system modulate reinforcement learning depending on feedback timing in adults, but their contributions during development remain unclear. In a 2-year longitudinal study, 6-to-7-year-old children performed a reinforcement learning task in which they received feedback immediately or with a short delay following their response. Children's learning was found to be sensitive to feedback timing modulations in their reaction time and inverse temperature parameter, which quantifies value-guided decision-making. They showed longitudinal improvements towards more optimal value-based learning, and their hippocampal volume showed protracted maturation. Better delayed model-derived learning covaried with larger hippocampal volume longitudinally, in line with the adult literature. In contrast, a larger striatal volume in children was associated with both better immediate and delayed model-derived learning longitudinally. These findings show, for the first time, an early hippocampal contribution to the dynamic development of reinforcement learning in middle childhood, with neurally less differentiated and more cooperative memory systems than in adults.

Introduction

As children enter school during middle childhood, they must learn to act appropriately in new situations through feedback. For example, children must learn to raise their hand before speaking during class. The teacher may reinforce this behavior immediately or with a delay, which raises the question whether feedback timing modulates their learning. Here, reinforcement learning (RL)[1] provides a useful mechanistic framework to describe such feedback-driven value-based learning and decision-making. RL models allow to explicitly test for the influence of separate components during value-based learning, such as model-free and model-based learning[2], social and non-social learning[3,4], or the contribution of different memory systems[5–7].

The role of feedback timing has previously been studied in relation to memory sytems. The memory systems account is a theoretical framework that proposes that different types of memory are supported by distinct neural systems in the brain. Specifically, this account suggests that there are two memory systems: a hippocampal-dependent system and a striatal-dependent system. These systems modulate memory and value-based learning, and their interactive development has been of particular interest to developmental research[8,9]. In adults, the hippocampal-dependent memory system has been shown to contribute to episodic memory during reinforcement learning and is more engaged during feedback that is presented with a delay[6,10,11], as opposed to the striatal-dependent memory system, which is more engaged after immediate feedback and supports habitual memory[5,12–14]. Specifically, hippocampal activation was greater during delayed feedback than during immediate feedback, whereas striatal activation was greater during immediate feedback than during delayed feedback[5]. The engagement of the hippocampus during delayed feedback was further supported by enhanced episodic memory for incidentally presented objects compared to objects presented with immediate feedback. Taken together, findings from adult studies suggest that feedback timing modulates the engagement of the hippocampal and striatal memory systems during value-based learning. Given the differential developmental trajectories of these systems and the impact the systems have on reinforcement learning and memory, it is important to understand whether children would show similar feedback timing modulations as previously shown in adults. In addition, whether such feedback timing modulation changes over time remains largely unexplored. To this end, in this study, we examined the contributions of hippocampal and striatal structural volumes during the longitudinal development of reinforcement learning across two years in 6-to-7-year-old children. We will introduce the key parameters in reinforcement learning and then we review the existing literature on developmental trajectories in reinforcement learning as well as on hippocampus and striatum, our two brain regions of interest.

Reinforcement learning behavior modulated by feedback timing can be modeled computationally using at least three parameters that reflect feedback-based learning and decision-making. For feedback-based learning, a learning rate parameter determines the extent to which the reward prediction error, defined as the difference between the received reward and the expected reward, influences the update of the future choice values. A higher learning rate emphasizes recent outcomes,

whereas a lower learning rate reflects learning integrated over a longer outcome history[15]. Value updates

may further depend on an outcome sensitivity parameter that scales the individual magnitude of received

rewards. Finally, in decision-making, the inverse temperature parameter plays a key role in determining

the tendency to select the more valuable choice and quantifies choice stochasticity. A higher inverse

temperature reflects more value-guided, deterministic choice behavior compared to a lower inverse

temperature reflecting more random choices. Learning rates and inverse temperature have been studied

extensively across development, mainly with cross-sectional studies showing mixed findings regarding

their age gradients[16]. One study reported lower learning rates in children compared to adolescents[17],

while other studies found no differences[18,19] or even higher learning rates in children[8,20]. Developmental

differences regarding the inverse temperature parameter are slightly more consistent, with studies

reporting no differences[8,21–23] or higher inverse temperature with age that suggests that behavior is

increasingly value-guided and less explorative[17–19,24]. To the best of our knowledge, outcome sensitivity

has not been modeled computationally across development. However, studies that linked striatal reward

activation to self-reported reward sensitivity showed increasing sensitivity from childhood to

adolescence[25,26].

In general, the inconsistencies regarding developmental differences in parameters may be due

to their dependency on model and task properties[27], which could be reconciled by comparing

developmental changes to simulation-based optimal learning[15]. Such comparisons acknowledge that

optimal parameter values vary depending on the context, and it has been suggested that humans develop

towards more optimal parameter values from childhood into adulthood[16]. Importantly, to our knowledge

previous reinforcement learning studies with children were cross-sectional, and only two studies

investigated children under 8 years of age[17,28]. Cross-sectional studies, in which developmental change

is inferred as a between-subject factor, do not capture the dynamics in middle childhood if individual

differences are large, whereas longitudinal studies test development as a within-subject factor, which is

crucial for uncovering change across time. Thus, longitudinal changes in reinforcement learning in

middle childhood as well as their putative striatal and hippocampal associations remain unknown. To

this end, learning rates, outcome sensitivity and inverse temperature are relevant computational

parameters to study longitudinal changes in striatal and hippocampal systems during value-based

learning.

Striatal and hippocampal contributions to reinforcement learning during middle childhood may

differ as these brain regions undergo major developmental changes. Whereas earlier structural studies

with relatively small sample sizes showed large developmental variability and a tendency for an earlier

volume peak in the striatum than in the hippocampus[29–35], a recent cross-sectional large-scale study was

able to contrast striatal and hippocampal trajectories with greater granularity[36]. These data showed

striatal volume peaks in the first decade which then declined throughout later developmental periods,

whereas hippocampal volume showed a more protracted inverted-U-shaped trajectory that peaked in

adolescence. Based on these structural findings, striatal and hippocampal systems are expected to

develop functionally at different rates[37], with habit memory depending on the earlier developing striatum

and episodic memory depending on the later developing hippocampus[38]. A direct investigation of the

longitudinal development of both memory systems in childhood would shed light on whether the

memory systems show a differential engagement similar to that of adults[5]. Such knowledge could be

useful to structure learning processes according to the developmental status. For example, children's

ability to learn from delayed feedback may depend on how well their hippocampus has developed. In

the same study sample, we previously reported that children's hippocampal volume was related to their

family's income level[39]. Additionally, previous research has shown that stress can reduce the

effectiveness of the hippocampal-dependent memory system[11]. This suggests that environmental factors

such as income and stress may play a role in shaping how well children learn from delayed feedback,

particularly through their impact on hippocampal development. By identifying the specific

environmental factors that impact children's learning and brain development, we can identify risk groups

and tailor interventions to ameliorate adverse effects.

This study aimed to explore the development of value-based learning in children and its

relationship with structural brain development over time. We hypothesized that the timing of feedback

would modulate children's learning from reinforcement, and that such modulation can be captured by

reinforcement learning (RL) model parameters. Additionally, we predicted that children's value-based

longitudinal development would shift towards more optimal learning behavior. Regarding structural

brain development, we expected the striatum to be relatively mature by middle childhood compared to

the protracted hippocampal maturation. Our second objective was to investigate the relationship between

value-based learning and structural brain development using longitudinal structural equation modeling.

We anticipated that there would be differentiated brain-cognition links between brain volume and value-

based learning. Specifically, we predicted that immediate feedback learning would be more strongly

associated with striatal volume, whereas hippocampal volume would be more closely linked to delayed

feedback and the facilitation of episodic memory encoding. Finally, we examined how these brain-

cognition dynamics would change over time by analyzing their longitudinal changes.

Method

Participants

Children and their parents took part in 2 waves of data collection with an interval of about 2 years (*mean*

= 2.07, *SD* = 0.17, *range* = 1.69 – 2.68). The inclusion criteria for wave 1 were children attending first

or second grade, no psychiatric or physical health disorders, at least one parent speaking fluent German,

and born full-term (≥ 37 weeks of gestation). At wave 1, 142 children (46% female, age *mean* = 7.19,

SD = 0.46, *Range* = 6.07 - 7.98) and their parents or caregivers participated in the study. 140 children

were included in the analysis (one child did not complete the probabilistic learning task, and another

child was later excluded due to technical problems during the task). A subgroup of 90 children (49%

female, 100% right-handed), who was randomly selected, completed magnetic resonance imaging

(MRI) scanning at wave 1, and 82 of them contributed to structural data after removing scans with

excessive movement. At wave 2, 127 children (46% female, age *mean* = 9.25, *SD* = 0.45, *Range* = 8.30

-10.2) continued taking part in the study, while families of the remaining children were unable to be

contacted or decided not to return to the study. 126 children at wave 2 completed the reinforcement

learning task and were included in the analysis. All children at wave 2 were invited for MRI scanning,

and 104 of them completed scanning (45% female, 92% right-handed). 99 children contributed to

structural data, after removing scans with excessive movement. In total, 73 children contributed to the

longitudinal MRI data and 126 children contributed to the longitudinal learning data. As previously

reported for this study sample, we found no systematic bias due to wave 2 dropout[39].

Procedure

The study consisted of a series of cognitive tasks tested during two behavioral sessions, including a

reinforcement learning task, and one MRI session at wave 1[39,40]. Two years later, the children underwent

one behavioral and one MRI session. MRI scanning was performed within three weeks of the behavioral

task session. Each session lasted between 150 and 180 minutes and was scheduled either on weekdays

between 2 p.m. and 6 p.m. or during weekends. Before participation, the parents provided written

informed consent and children's verbal assent at both waves. All children were compensated with an

honorarium of 8 euro per hour.

Measures

Reinforcement learning task. Children completed an adapted reinforcement learning task[5] in which they

learned the preferred associations between four cues (cartoon characters) and two choices (round-shaped

or square-shaped lolli) through probabilistic feedback (87.5 % contingent and 12.5 % non-contingent

reward probability). In each trial, after an initial inter-trial interval of 0.5 s, a cue and its choice options

were presented for up to 7 s until the child made a choice (Figure 1, choice phase). In the delay phase,

we manipulated feedback timing. For two cues, the selected choice remained visible for 1 s (immediate

feedback condition), whereas for the other two cue characters, it remained visible for 5 s before feedback

was given (delayed feedback condition). A final feedback phase of 2 s indicated a reward by a green

frame, and a punishment by a red frame. Inside each frame, a unique object picture was shown, which

was incidentally encoded and irrelevant to the task. The child was instructed to pay attention to the

feedback indicated by the frame color. In an initial practice phase of 32 trials, the child practiced the

task with a fifth cartoon character not included in the actual task to avoid practice effects. The

experimenter instructed the child to select the choice that was most likely to result in a reward. The

Experimenter checked whether the child learned the more rewarded choice during practice and let it

repeat the practice task otherwise to ensure understanding of the task. In the actual task, 128 trials were

presented in four blocks and with small breaks in between. Cues were presented in a mixed, pseudo-

randomized order. A total of 64 unique objects were shown in the feedback phase, each one twice within

the same feedback condition. In both delay phases, contingent choice and choice location remained the

same for each cue within the task, but were balanced across participants by using four different task

versions. At wave 2, four new cues replaced the previous ones to rule out memory effects.

Object recognition test. At wave 1, children were additionally tested for recognition memory on the

object pictures that were incidentally encoded during reinforcement learning. A total of 80 objects (48

old objects and 32 new objects) were presented in randomized order. The 48 old objects (24 for each

feedback condition) were selected from the 64 old objects shown during learning based on two lists to

balance the shown and omitted old objects across task versions. Each old object was shown twice during

learning, but if the child failed to respond during learning, no feedback or object was shown in the trial,

so some objects only appeared once. These objects were excluded at the individual level (individually

missing object *mean* = 2.71). At recognition, children had 4 response options ('old sure', 'old unsure',

'new unsure', 'new sure') with up to 7 s to respond. The children answered verbally, and the

experimenter entered their response. At wave 2, this test was excluded due to time constraints.

Figure 1. (A) Depiction of two example trials of immediate and delayed feedback conditions presented

at wave 1. For immediate feedback (top panel), between choice response and feedback, cue and choice

were presented for 1 s. At feedback, a green frame around the incidentally encoded object indicated a

positive outcome, which appeared in 87.5% of the trials when selecting the squard-shaped lolli for this

example cue. For delayed feedback (bottom panel), the delay phase between choice response and

feedback lasted for 5 s. The red frame around the object indicated a negative outcome and appeared in

87.5% of the trials when selecting the squard-shaped lolli for this example cue. (B) For each feedback

condition, two action-outcome contingencies were learned to balance a potential choice bias. With the

four task versions, the cues and outcome contingencies were counterbalanced across participants.

Brain volume. We extracted the bilateral brain volumes for our regions of interest, which were striatum

and hippocampus. The striatum regions included nucleus accumbens, caudate and putamen. For our

imaging data, structural MRI images were acquired on a Siemens Magnetom TrioTim syngo 3 Tesla
scanner with a 12-channel head coil (Siemens Medical AG, Erlangen, Germany) using a 3D T1–
weighted Magnetization Prepared Rapid Gradient Echo (MPRAGE) sequence, with the following
parameters: 192 slices; field of view $= 256$ mm, voxel size $= 1$ mm^3, TR $= 2500$ ms; TE $= 3.69$ ms,
flip angle $= 7°$, TI $= 1100$ ms. Volumetric segmentation was performed using the Freesurfer 6.0.0
image analysis suite[41]. Previous studies suggested that software tools based on adult brain templates
provide inaccurate segmentation for pediatric samples, which can be improved through the use of study–
specific template brains[42,43]. Thus, we created two study-specific template brains (one for each wave)
using Freesurfer's "make_average_subject" command. This pipeline utilized the default adult template
brain registrations of the "recon–all–all" command to average surfaces, curvatures, and volumes from
all subjects into a study–specific template brain. All subjects were then re–registered to this study-
specific template brain to improve segmentation accuracy. Segmented images were manually inspected
for accuracy and 8 cases at wave 1 and 5 cases at wave 2 were excluded for inaccurate or failed
registration due to excessive motion.

Data analysis

Behavioral learning performance. As a first step, we calculated learning outcomes diretly from the raw
data, which where learning accuracy, win-stay and lose-shift behavior as well as reaction time. Learning
accuracy was defined as the proportion to choose the more rewarding option, while win-stay and lose-
shift refer to the proportion of staying with the previously chosen option after a reward and switching
to the alternative choice after receiving a punishment, respectively. We used these outcomes as our
dependent variables to examine the effect of the predictors feedback timing (immediate, delayed), wave
(1, 2), wave 1 age, and sex (girls, boys), utilizing generalized linear mixed models (GLMM) with the R
package lme4[44]. All reported models included random slopes for within-subject factors feedback timing
and wave (see Supplementary Material 2 for the model structure). We systematically tested main effects
and interactions between the predictors and their interaction had to statistically improve the predictive
ability of the model to be included in the final reported model. All predictor variables were grand-mean-
centered to interpret the interaction effects independent from other predictors.

Reinforcement learning models. As a next step, we used computational modeling to compare the
learning models of basic heuristic strategies and value-based learning and to determine the model that
could best capture children's trial-by-trial learning behavior. For heuristic strategies, we considered
models that reflected a Win-stay-lose-shift (wsls) or a Win-stay (ws) strategy. Win-stay is a heuristic
strategy in which the same action is repeated if it leads to a positive outcome in the previous trial, and
Win-stay-lose-shift additionally switches to a different action if the previous outcome is negative. Note
that these model-based outcomes are not identical to the win-stay and lose-shift behavior that were

calculated from the raw data. The use of such model-based measure offers the advantage in discerning

the underlying hidden cognitive process with greather nuance, in contrast to classical approaches that

directly use raw behavioral data. The models quantified the learning behavior for each individual I for

each cue c and trial t. The heuristic models consisted of a weight w that reflected its degree in strategy

use. In the case of reward $r = 1$, w was equal to 1 for the chosen option (eg. choice A), and 0 for the

unchosen option (e.g. choice B), thus maximizing win-stay, i.e., choosing A at the subsquent trial $t + 1$:

$$w_{i,c,t+1,A|r = 1} = 1 \text{ and } w_{i,c,t+1,B|r = 1} = 0 \tag{1}$$

For trials $r = 0$ (applicable only to the wsls model), model weights were the opposite, maximizing lose-

shift:

$$w_{i,c,t+1,A|r = 0} = 0; \; w_{i,c,t+1,B|r = 0} = 1 \tag{2}$$

The initial weights for both choices were set to $w_{i,c,t=1} = 0.5$. The weight w then scaled the parameter

τ_wsls or τ_ws to estimate the individual strategy use during decision-making. The choice probabilities

were calculated using the softmax function, eg., for the chosen option A:

$$p(A) = \frac{exp^{w_{i,c,t,A}*\tau\_wsls_i}}{exp^{w_{i,c,t,A}*\tau\_wsls_i} + exp^{w_{i,c,t,B}*\tau\_wsls_i}} \tag{3}$$

Thus, a higher probability of strategy use was reflected by a larger value of τ_wsls or τ_ws.

For value-based learning, we considered a Rescorla-Wagner model and several variants based on our

theoretical conceptions. The baseline value-based model vbm_1 updated the value v of the selected choice

(A or B) for the next trial t. This value update was determined by calculating the difference between the

received reward r and the expected value v of the selected choice, which was the reward prediction error.

The value update was further scaled by a learning rate α ($0 < \alpha < 1$):

$$v_{i,c,t+1,A} = v_{i,c,t,A} + \alpha_i(r_{i,c,t} - v_{i,c,t,A}) \tag{4}$$

When the outcome sensitivity parameter ρ ($0 < \rho < 20$) was included, the reward was additionally

scaled at the value update:

$$v_{i,c,t+1,A} = v_{i,c,t,A} + \alpha_i(\rho_i * r_{i,c,t} - v_{i,c,t,A}) \tag{5}$$

The inverse temperature parameter τ ($0 < \tau < 20$) was included in the softmax function to compute

choice probabilities:

$$p(A) = \frac{exp^{v_{i,c,t,A}*\tau_i}}{exp^{v_{i,c,t,A}*\tau_i} + exp^{v_{i,c,t,B}*\tau_i}} \tag{6}$$

Note, however, that outcome sensitivity and inverse temperature are difficult to fit simultaneously due

to non-identifiability issues[45]. Therefore, models including the inverse temperature fixed outcome

sensitivity at 1 (inverse temperature model family), assuming no individual differences in outcome

sensitivity. For the outcome sensitivity model family, outcome sensitivity was freely estimated, and the

inverse temperature was fixed at 1, asssuming the same degree of value-based decision behavior across

individuals. Even though outcome sensitivity is usually restricted to an upper bound of 2 to not inflate

outcomes at value update, this configuration led to ceiling effects in outcome sensitivity and non-

converging model results. Further, this issue was not resolved when we fixed the inverse temperature at

the group mean of 15.47 of the winning inverse temperature family model. It may be that in children,

individual differences in outcome sensitivity are more pronounced, leading to more extreme values.

Therefore, we decided to extend the upper bound to 20, parallel to the inverse temperature, and all our

models converged with Rhat < 1.1. Each model family consisted of 4 model variants vbm_{1-4} ($1\alpha1\tau$,

$2\alpha1\tau, 1\alpha2\tau, 2\alpha2\tau$) and vbm_{5-8} ($1\alpha1\rho$, $2\alpha1\rho, 1\alpha2\rho, 2\alpha2\rho$), in which each parameter was either

separated by feedback timing or kept as a single parameter across feedback conditions. Our baseline

value-based model vbm_1 included a single learning rate and a single inverse temperature ($1\alpha1\tau$).

Parameter estimation. All choice data were fitted in a hierarchical Bayesian analysis using the Stan

language in R[46,47] adopted from the hBayesDM package[48]. Posterior parameter distributions were

estimated using Markov chain Monte Carlo (MCMC) sampling running 4 chains each with 3,000

iterations, using the first half of the chain as warmup, and group-level parameters and individual-level

parameters were estimated simoultaneously. The hierarchical Bayesian approach provides more stable

and reliable parameter estimates as opposed to point-estimation approaches like maximum likelihood

estimation[49]. Each model fit both wave 1 and wave 2 data at once, considering the correlation structure

of the same parameter across waves, to account for within-subject dependency using the Cholesky

decomposition. The Cholesky decomposition used a Lewandowski-Kurowicka-Joe prior of 2, and all

other group-level parameters had a prior normal distribution, Normal (0, 0.5). Non-response trials (wave

1 = 2.41%, wave 2 = 0.97% on average) were excluded in advance.

Model simulation and model-derived learning score. To appropriately interpret the parameter results

with respect to the optimal parameter combination of the winning model, we simulated 5,000,000

individual datasets using 10,000 different parameter value combinations (covering the whole range of

each parameter) to identify the optimal parameter combination of the winning model that was selected

by model comparison. In addition, we computed the model-derived mean choice probability of the

contingent, i.e., the more rewarded option, and we referred to it as the model-derived learning score.

This model-derived choice probability differs from the observed empirical choice probability (i.e., the

accuracy of selecting the more rewarded option), because the model-derived learning score combines

the model with the data by incorporating latent information carried out by key learning parameters. Thus,

the learning score captures observed behavior based on trial-by-trial latent processes predicted by value-

based models. We used this as metric to interpret the fitted posterior parameters in relation to the optimal

parameter combination of our probabilistic learning task.

Model selection and validation. We conducted a 2-step sequential procedure for the model development

and model selection. As a first step, we compared model evidence for the baseline value-based model

that does not separate learning rate and inverse temperature by feedback timing (vbm_1:$1\alpha, 1\tau$) to the

non-value-based, heuristic strategy models that reflect Win-stay or Win-stay-lose-shift strategy behavior

($ws, wsls$). As a second step, we compared model evidence for 8 value-based model variants, 4 of the

model family with learning rate and inverse temperature ($1\alpha1\tau$, $2\alpha1\tau$, $1\alpha2\tau$, $2\alpha2\tau$) and 4 of the model

family with learning rate and outcome sensitivity ($1\alpha1\rho$, $2\alpha1\rho$, $1\alpha2\rho$, $2\alpha2\rho$). This allowed us to

compare whether children showed separable effects of feedback timing on one of the model parameters.

We compared the model fit using Bayesian leave-one-out cross-validation and obtained the expected

log pointwise predictive density ($elpd_{loo}$) using the R package loo[50]. We further computed the model

weights (*Pseudo-BMA+*) using Pseudo Bayesian model averaging stabilized by Bayesian bootstrap with

100,000 iterations[51]. To validate our models, we estimated predictive accuracy by comparing one-step-

ahead model predictions with the choice data[15,52]. We performed parameter recovery for the winning

model and model recovery by comparing it to a set of models used during model comparison

(Supplementary Material 1)[53].

Episodic memory at wave 1

We predicted the individual corrected recognition memory (hits-false alarms) by feedback condition in

a linear mixed effects model using the R package lme4[44]. Only confident ("sure") ratings were included

in the analysis, which were 98.1 % of all given responses. A total of 140 children completed the

recognition memory test and 138 were included in the analysis, with two being excluded due to negative

corrected recognition memory value (i.e., poor recognition memory). Age and sex were controlled for

as covariates.

Longitudinal brain-cognition links

We used latent change score (LCS) models to examine the longitudinal relationships between brain and

learning score measures. LCS models are longitudinal structural equation models that have been widely

applied to estimate developmental changes and coupling effects across domains such as the brain and

cognition[54,55]. LCS models allow the definition of specific paths between multiple variables to test

explicit hypotheses and estimate latent change from the observed variables that account for measurement

error and increase testing power[56]. We compiled univariate LCS models for each variable separately

(learning scores and brain volumes) to examine whether there was significant individual variance and

change, which could be related within a multivariate LCS model as a next step. Model fit had to be at

least acceptable, with a comparative fit index (*CFI*) > 0.95, standardized root mean square residual

(*SRMR*) < .08 and root mean square error of approximation (*RMSEA*) < .08[57]. Age and sex were included

as covariates at wave 1, as well as the estimated total intracranial volume (eTIV) when brain volume

was included in the model. Multivariate LCS models allow to estimate meaningful brain-cognition

relationships: a wave 1 covariance between brain and cognition, brain predicting change onto cognition,

or vice versa, and a covariance in both brain and cognition change scores (wave 1 to wave 2). Before

compiling the variables into an LCS model, they were checked for outliers ± 4 *SD* around the mean. We

identified one outlier for the learning rate at wave 2, which was removed for the explorative LCS model

that included model parameters. There were no further outliers in other cognitive variables or brain

volumes. Continuous variables were standardized to the wave 1 measure so that wave 2 values represent

the change from wave 1, sex was contrast-coded (girls = 1, boys = -1).

364   Results

Behavioral results

First, we were interested in whether children showed behavioral differences between waves and

feedback timing. A descriptive overview is provided in Table 1 and Figure 2. The details of the reported

GLMM models, including the random effects structure and the effects of age and sex, are described in

the Supplementary Material 2. Since some children were poor learners who failed to reach 50 % average

accuracy in their last 20 trials (13 children at wave 1 and 6 children at wave 2), we also performed

behavioral analyses with a reduced dataset in which results remained unchanged (Supplementary

Materials 6).

Table 1. Descriptive behavioral results of dependent variables Accuracy (ACC, probability correct),

win-stay probability (WS), lose-shift probability (LS), and reaction time (RT, in seconds), as well as

mixed model fixed effects that predicted these dependent variables.

	Descriptive Results				Mixed Model Effects	
	Wave 1		Wave 2		Wave	Feedback
	Ime	Del	Ime	Del		
ACC	0.69 (0.46)	0.70 (0.46)	0.79 (0.41)	0.80 (0.40)	↑ W2	–
WS	0.81 (0.39)	0.80 (0.40)	0.88 (0.32)	0.88 (0.32)	↑ W2	–
LS	0.47 (0.50)	0.50 (0.50)	0.42 (0.49)	0.42 (0.49)	↓ W2	–
RT	2.10 (1.31)	2.07 (1.29)	1.70 (1.02)	1.67 (1.00)	↓ W2	↓ Del

Note. Mean (standard deviation) of the variables, split by wave and feedback timing, is reported in the

table. Mixed model effects and their directionality (increasing ↑ or decreasing ↓) predicting the

dependent variables. W2 = Wave 2, Ime = Immediate feedback, Del = Delayed feedback.

Children's learning improved between waves. With the complete dataset, we found that increased

learning accuracy (i.e., the probability of choosing the more rewarding option) was predicted at wave 2

compared to wave 1, but there were no differences in accuracy by feedback timing ($\beta_{wave=2}$ = .550, *SE*

= .061, z = 8.97, p < .001, $\beta_{feedback=delayed}$ = .013, *SE* = .024, z = 0.54, p = .590). Furthermore, win-

stay probability increased and lose-shift probability decreased longitudinally, again without differences

by feedback timing (WS: $\beta_{wave=2}$ = .586, *SE* = .071, z = 8.22, p < .001, $\beta_{feedback=delayed}$ = .023, *SE*

= .033, z = 0.69, p = .489; LS: $\beta_{wave=2}$ = -.252, SE = .037, z = -6.87, p < .001, $\beta_{feedback=delayed}$ = .030,

SE = .022, z = 1.37, p = .169). Reaction times were faster at wave 2 compared to wave 1, and they were

faster for delayed compared to immediate feedback trials ($\beta_{wave=2}$ = -221, SE = 22.8, $t(df_{\text{Satterthwaite}}$ =

135) = -9.70, p < .001, $\beta_{feedback=delayed}$ = -13.8, SE = 6.59, $t(df_{\text{Satterthwaite}}$ = 136) = -2.10, p = .038). To

summarize, children's average accuracy improved over 2 years, while their win-stay probability

increased and their lose-shift probability decreased between waves. Children were able to respond faster

to cues paired with delayed feedback compared to cues paired with immediate feedback, and they

became faster in their decision-making across waves (see mixed model effects overview in Table 1). Of

note, reaction times were largely uncorrelated with accuracy and switching behavior (win-stay, lose-

shift), while accuracy and switching behavior showed significant correlations at both waves (Figure 2D).

Figure 2. Individual differences in the behavioral reinforcement learning outcomes and their longitudinal

change. (A) Accuracy did not differ by feedback timing and increased between waves. (B) Win-stay and

lose-shift proportion did not differ by feedback timing, and win-stay increased and lose-shift proportion

decreased between waves. (C) Reaction time differed by feedback timing, in which decisions for cues

learned with delayed feedback were faster, and reaction times were faster at wave 2 compared to wave

1. (D) Correlations between behavioral outcomes reveal that learning accuracy was primarily correlated

with the win-stay and lose-shift probabilities both within and between waves, but was uncorrelated to

reaction time. Significant correlations are circled, p-values were adjusted for multiple comparisons using

bonferroni correction.

Modeling results

Children's behavior was best described by value-based learning. We conducted a 2-step sequential

procedure for model development and model selection. Model comparison using leave-one-out cross

validation showed evidence in favor of the value-based learning model, reflected in the highest expected

log pointwise predictive density and highest model weights, confirming that children's learning

behavior in the longitudinal data can generally be better described by a value-based rather than by a

heuristic strategy model ($elpd_{loo}$ = -15154.9, *pseudo-BMA+ = 1*, Table 2). Children whose individual

fit was better for a heuristic model ($wsls$) than for the value-based model (vbm_1), were at both waves

more likely to be poor learners (defined as an accuracy below 50% in the last 20 trials). Taken together,

children's learning behavior was best described by a value-based model, and a heuristic strategy model

captured more poor learners compared to a value-based model.

Table 2. Model comparison results.

Model	Parameters	$\Delta elpd_{loo}$ [SE]	$\Sigma elpd_{loo}$ [mean]	pseudo-BMA+
Step 1: heuristic strategy models and value-based learning model				
vbm_1	1α, 1τ	0 [0]	-15154.9 [-0.45]	1
ws	$1\tau_{ws}$	-1327.7 [159.5]	-16482.7 [-0.49]	<0.01
$wsls$	$1\tau_{wsls}$	-4247.3 [284.8]	-19402.3 [-0.58]	0
Step 2: value-based learning models				
vbm_3	**1α, 2τ**	**0 [0]**	**-15045.3 [-0.45]**	**0.73**
vbm_7	1α, 2ρ	-2.93 [2.92]	-15048.2 [-0.45]	0.24
vbm_6	2α, 1ρ	-24.34 [8.85]	-15069.6 [-0.45]	<0.01
vbm_8	2α, 2ρ	-29.71 [15.95]	-15075.0 [-0.45]	0.02
vbm_4	2α, 2τ	-43.34[14.89]	-15088.6 [-0.45]	<0.01
vbm_2	2α, 1τ	-46.45 [13.97]	-15091.7 [-0.45]	<0.01
vbm_5	1α, 1ρ	-59.01 [7.59]	-15104.3 [-0.45]	<0.01
vbm_1	1α, 1τ	-109.63 [11.98]	-15154.9 [-0.45]	<0.01

Note. Model = heuristic (ws, $wsls$) and value-based models (vbm_{1-8}) that were compared against each

other. Parameters = corresponding model parameters learning rate α, inverse temperature τ and

outcome sensitivity ρ. $\Delta elpd_{loo}$[SE] = difference in the Bayesian leave-one-out cross-validation

estimate of the expected log pointwise predictive density relative to the winning model and its standard

errors. $\Sigma elpd_{loo}$[mean] = sum of expected log pointwise predictive density of all 33,460 trials,

including all participants and waves, and trial mean. *Pseudo-BMA+* = model weight for relative model

evidence using Bayesian model averaging stabilized by Bayesian bootstrap using 100,000 iterations.

434   *Feedback timing modulated choice stochasticity.* Model vbm_3 ($1\alpha 2\tau$) showed the largest model

435   evidence, reflected in the highest expected log pointwise predictive density and highest model weights

436   and suggests that feedback timing affected the inverse temperature, but not the learning rate or outcome

437   sensitivity ($elpd_{loo}$ = -15045.3, *pseudo-BMA+* = 0.73, Table 2). Table 3 and Figure 3A provide a

438   descriptive overview of the winning model parameters. Of note, there were only small differences in

439   model fit ($elpd_{loo}$) to the second-best model (vbm_7, $1\alpha 2\rho$, $\Delta elpd_{loo}$ = -2.93, $elpd\_SE_{loo}$ = 2.92,

440   *pseudo-BMA+* = 0.24), which suggests a potential separable feedback timing effect on outcome

441   sensitivity. We also performed the model comparison with a reduced dataset in which the winning model

442   remained the same (Supplementary Materials 6). The average inverse temperature did not differ by

443   feedback condition, but showed large within-person condition differences at both waves, indicating

444   individual differences in feedback timing modulation (wave 1: $\Delta\tau_{del-ime}$ *Mean* = 0.22, *SD* = 3.80,

445   *Range* = 21.74, wave 2: $\Delta\tau_{del-ime}$ *Mean* = 0.35, *SD* = 3.70, *Range* = 24.03). The correlations between

446   the parameters are shown in Supplementary Material 3.

447   Since reaction times were predicted by feedback timing behaviorally, and inverse temperature is

448   assumed to reflect decision-making, we were interested in whether differences in reaction time were

449   related to inverse temperature differences. Indeed, at both waves, children who responded faster during

450   delayed compared to immediate feedback had a higher inverse temperature at delayed compared to

451   immediate feedback (wave 1: r = -.261, t(*df* = 138 = -3.18, p = .002, wave 2: r = -.345, t(*df* = 124) = -

452   4.10, p < .001, Figure 3B). Taken together, children's learning behavior was best described by a value-

453   based model, where feedback timing modulated individual differences in the choice rule during value-

454   based learning. Interestingly, the differences in the choice rule and reaction time were correlated.

455   Specifically, more value-guided choice behavior (i.e., higher inverse temperature) was related to faster

456   responses during delayed feedback relative to immediate feedback, suggesting a link between model

457   parameter and behavior in relation to feedback timing.

458

Figure 3. (A) Individual differences in the learning rate and inverse temperature of the winning model and their longitudinal change. The inverse temperature τ but not learning rate α was separated by feedback timing, and both increased between waves in their values (top panel). The condition difference in the inverse temperature did not differ on average, but showed individual differences (bottom left panel). (B) The condition differences in the inverse temperature correlated with reaction time, i.e., higher delayed compared to immediate inverse temperature was related to faster delayed compared to immediate reaction time.

Table 3. Description of model parameters from the winning value-based model vbm_3.

	Wave 1					Wave 2				
	α	τ_{Ime}	τ_{Del}	ls_{Ime}	ls_{Del}	α	τ_{Ime}	τ_{Del}	ls_{Ime}	ls_{Del}
Mean	0.02	14.6	14.8	0.73	0.73	0.05	16.2	16.5	0.82	0.82
SD	0.02	2.04	2.37	0.12	0.13	0.04	2.37	2.21	0.13	0.13
Min	<0.01	6.73	5.25	0.53	0.53	<0.01	4.37	6.85	0.53	0.53
Max	0.09	17.5	17.9	0.94	0.94	0.22	18.6	18.7	0.96	0.96

Note. α = learning rate across feedback timing, τ_{Ime}/ ls_{Ime} = inverse temperature and learning score for immediate feedback, τ_{Del}/ ls_{Del} = inverse temperature and learning score for delayed feedback.

Children's value-based learning became more optimal. Next, we compared the parameter space according to model simulation (Figure 4A) with the empirical posterior parameters fitted by the winning model (Table 3, Figure 4B) to determine whether children increased their value-based learning towards more optimal parameter combinations. Both fitted and simulated parameter combinations allowed us to derive a learning score that captured learning performance according to the winning value-based model.

Note that the learning score was defined as the average choice probability for the more rewarded choice

option. We refer to these model-derived choice probabilities as learning score, since they reflect value-

based learning and combine information of learned values, that depend on the learning rate, and values

translated into choice probabilities, that depend on the inverse temperature. Thus, a higher learning score

reflects more optimal value-based learning. We simulated 10,000 parameter combinations and created

a learning score map according to each parameter combination (Figure 4A). The optimal parameter

combination was at a learning rate $\alpha = 0.29$, and an inverse temperature $\tau = 19.8$, and with an average

learning score of 96.5 % (Figure 4A). Children's fitted learning rates ranged 0.01 – 0.22 and inverse

temperature 6.73 – 18.70 and were outside the parameter space of a learning score above 96 % (Table

3 and Figure 4A). The average longitudinal increases in learning rate and inverse temperature were

mirrored by average increases in the learning scores, confirming our prediction that their parameters

developed towards optimal value-based learning (arrow in Figure 4B). We further found that the average

longitudinal change in win-stay and lose-shift proportion also developed towards more optimal value-

based learning (Supplementary Material 4).

Model validation. To validate our winning model vbm_3, we estimated its predictive accuracy by

comparing one-step-ahead model predictions with the choice data. The one-step ahead predictions of

the winning model captured children's choices overall well, with predictive accuracies of 65.3 % at

wave 1 and 75.7 % at wave 2 (Figure 4C). Further, our winning model showed a good parameter

recovery for learning rate ($r = 0.85$) and inverse temperature ($r = 0.75 – 0.77$). Our winning model

showed excellent on the group level (100%) when comparing it to a set of models used during model

comparison ($vbm_1, vbm_7, wsls$). The individual model recovery was lower (58%), with 35% of the

simulated winning model fitting best on our baseline model vbm_1 with a single inverse temperature,

which likely reflects the noisy property of the inverse temperature (Supplementary Material 1).

Figure 4. (A) The model simulation depicts parameter combinations and simulation-based average learning scores. The cyan "X" in the middle top depicts the optimal parameter combination where average learning scores were at 96.5 %, and the cyan rectangle depicts the space of the fitted parameter combinations, (B) Enlarged view of the space of fitted parameter combinations. The colored arrows depict mean change (bold arrow) and individual change (transparent arrows) of the fitted parameters. The greyscale gradient-filled dots, that are connected by the arrows, depict the individual learning score, while the the greyscale gradient in the background depicts the simulated average learning score. The mean change reveals an overall change towards the higher, i.e., more optimal, learning scores. (C) One-step-ahead posterior predictions of the winning model for each wave. The colored lines depict averaged trial-by-trial task behavior for each feedback condition, and a cyan ribbon indicates the 95% highest density interval of the one-step-ahead prediction using the entire posterior distribution.

Longitudinal brain-cognition links

Significant longitudinal change in brain and cognition. We first performed univariate LCS model analyses to estimate a latent change score of immediate and delayed learning scores as well as striatal and hippocampal volumes (see descriptive changes in Figure 5B-C). All four variables of interest showed significant positive mean changes and variances, and all univariate models provided a good fit to the data (Supplementary Material 5). This allowed us to further relate the differences in structural brain changes to changes in learning.

Hippocampal volume exhibited more protracted development during middle childhood. We next fitted

a bivariate LCS model to compare striatal and hippocampal change scores. We theorized that by middle

childhood, the striatum would be relatively mature, whereas the hippocampus continues to develop. We

progressively constructed multiple LCS models to test this idea. First, the bivariate LCS model provided

a good data fit (χ^2 (14) = 10.09, *CFI* = 1.00, *RMSEA (CI)* = 0 (0-.06), *SRMR* = .04). We then further

fitted two constrained models, to see whether setting the mean striatal change or the mean hippocampal

change to 0 would lead to a drop in the model fit. Compared to the unrestricted model, the constrained

model that assumed no striatal change did not lead to a drop in model fit ($\Delta\chi^2$ (1) = 2.74, *p* = .098),

whereas the model that assumed hippocampal change dropped in model fit ($\Delta\chi^2$ (1) = 12.69, *p* < .001).

Finally, we tested a more stringent assumption of equal change for striatal and hippocampal volumes,

in which the model dropped in model fit compared to the unrestricted model ($\Delta\chi^2$ (1) = 18.04, *p* < .001)

and suggests that striatal and hippocampal change differed. Together, these results support our

postulation of separable maturational brain trajectories in our study sample, suggesting that the

hippocampus continued to grow in middle childhood, whereas striatal volume increased less.

Hippocampal and striatal volume showed distinct associations to learning. We fitted a four-variate LCS

model to test our prediction of selective brain-cognition links. Specifically, we assumed a larger

contribution of striatal volume at immediate learning, and a larger contribution of hippocampal volume

at delayed learning. The LCS model provided good data fit (χ^2 (27) = 15.4, *CFI* = 1.00, *RMSEA (CI)* =

0 (0 − .010, *SRMR* = .045), and all relevant paths are shown in Figure 5D (see Table 4 for a detailed

model overview). For the striatal associations to cognition, we found that wave 1 striatal volume

covaried with both immediate learning score and delayed learning score ($\phi_{STR_{w1},LS_{i,w1}}$= 0.19, *z* = 2.52,

SE = 0.07, *p* = .012, $\phi_{STR_{w1},LS_{d,w1}}$= 0.18, *z* = 2.37, *SE* = 0.07, *p* = .018). Constraining the striatal

association to immediate learning to 0 worsened the model fit relative to the unrestricted model ($\Delta\chi^2$ (1)

= 5.66, *p* = .017), which was the same when constraining the striatal association to delayed learning to

0 ($\Delta\chi^2$ (1) = 5.14, *p* = .023). In summary, larger striatal volume was associated with better learning

scores for both immediate and better delayed feedback. This pattern remained the same in the results of

the reduced dataset (Supplementary Material 6).

Hippocampal volume, on the other hand, only covaried with delayed learning at wave 1 ($\phi_{HPC_{w1},LS_{d,w1}}$=

0.14, *z* = 2.05, *SE* = 0.07, *p* = .041), not with immediate learning score ($\phi_{HPC_{w1},LS_{i,w1}}$= 0.12, *z* = 1.68,

SE = 0.07, *p* = .092). Fixing the path between hippocampal volume and delayed learning to 0 worsened

the model fit relative to the unrestricted model ($\Delta\chi^2$ (1) = 4.19, *p* = .041), but not when its path to

immediate learning was constrained to 0 ($\Delta\chi^2$ (1) = 2.94, *p* = .086). This suggests that larger hippocampal

volume was specifically associated with better delayed learning. In the results of the reduced dataset,

the hippocampal association to the delayed learning score was no longer significant, suggesting a

weakened pattern when excluding poor learners (Supplementary Material 6). It is likely that the

exclusion reduced the group variance for hippocampal volume and delayed learning score in the model.

As a next step, the associations between striatum and hippocampus to immediate or delayed learning

was directly compared against each other. A model equal-constraining striatal and hippocampal paths

to immediate learning ($\Delta\chi^2$ (1) = 0.41, p = .521) and another model equal-constraining these paths to

delayed learning ($\Delta\chi^2$ (1) = 0.14, p = .707) did not lead to a worse model fit compared to the unrestricted

model, which suggests that the brain-cognition links have considerable overlap. This is in line with the

high wave 1 covariance and change-change covariance within the brain and cognition domain (see Table

4). We found no longitudinal links between the brain and cognition domains, which suggests that the

found brain-cognition links at wave 1 remained longitudinally stable (see Supplementary Material 5 for

an exploratory LCS model that related the model parameters to striatal and hippocampal volume).

Taken together, the confirmatory LCS model results were in line with our predictions of a relatively

larger involvement of the hippocampus during delayed feedback learning, but the findings on striatal

volume disconfirmed a selective association with immediate feedback learning and suggest a more

general role of the striatum in both learning conditions.

No evidence for enhanced episodic memory during delayed feedback. Finally, we investigated whether

a hippocampal contribution at delayed feedback would selectively enhance episodic memory. Episodic

memory, as measured by individual corrected object recognition memory (hits - false alarms) of

confident ("sure") ratings, showed at trend better memory for items shown in the delayed feedback

condition ($\beta_{feedback=delayed}$ = .009, *SE* =.005, *t(df* = 137) = 1.80, *p* = .074, see Figure 5A). Note that

in the reduced dataset, delayed feedback predicted enhanced item memory significantly (Supplementary

Material 6). The inclusion of poor learners in the complete dataset may have weakend this effect because

their hippocampal function was worse and was not involved in learning (nor encoding), regardless of

feedback timing. To summarize, there was inconclusive support for enhanced episodic memory during

delayed compared to immediate feedback, calling for future study to test the postulation of a selective

association between hippocampal volume and delayed feedback learning.

Figure 5. (A) Recognition memory (corrected recognition = hits - false alarms) for objects presented during delayed feedback was only enhanced at trend. (B) Learning scores depicted here were used in the LCS analyses. Learning scores were the model-derived choice probability of the contingent choice using fitted posterior parameters. (C) Hippocampal and striatal volumes increased between waves, while hippocampal volume increased most. (D) A four-variate latent change score (LCS) model that included striatal and hippocampal volumes as well as immediate and delayed learning scores. Depicted are significant paths cross-domain (brain-cognition, dashed lines) and within-domain (brain or cognition, solid lines), other paths are omitted for visual clarity and are summarized in Table 4. Depicted brain-cognition links included $\phi_{STR_{w1},LS_{ime,w1}}$ (covariance between striatal volume and immediate learning score at wave 1), as well as $\phi_{HPC_{w1},LS_{del,w1}}$ and $\phi_{STR_{w1},LS_{del,w1}}$ (covariances between hippocampal and striatal volumes and delayed learning score at wave 1). Brain links included $\phi_{STR_{w1},HPC_{w1}}$ and $\rho_{\Delta STR,\Delta HPC}$ (wave 1 covariance and change-change covariance), and similarly, cognition links included $\phi_{LS_{ime,w1},LS_{del,w1}}$ and $\rho_{\Delta LS_{ime},\Delta LS_{del}}$. Covariates included age, sex and estimated total intracranial volume. ** denotes significance at $\alpha < .001$, * at $\alpha < .05$.

603 Table 4. Parameter estimates of a four-variate latent change score model that includes brain (striatal and

604 hippocampal volume) and cognition domains (immediate and delayed learning score)

	STR	LS_{ime}	HPC	LS_{del}
Model fit: $\chi^2 = 15.4$, $df = 27$, $CFI = 1$, $RMSEA$ $(CI) = 0$ (0-0.01), $SRMR = 0.045$				
Mean change Δ	0.06* (0.03)	0.76** (0.08)	0.38** (0.04)	0.75** (0.08)
wave 1 variance σ	fixed to 1	fixed to 1	fixed to 1	fixed to 1
change variance σ_Δ	0.07** (0.01)	0.88** (0.10)	0.18* (0.07)	0.83** (0.10)
Intercept-change regression β	-0.04 (0.04)	-0.83* (0.29)	-0.16* (0.06)	-0.73* (0.27)
Wave 1 covariates				
age onto Intercept ϕ	0.19 (0.10)	-0.05 (0.08)	0.29* (0.08)	0.08 (0.08)
sex onto Intercept ϕ	-0.42** (0.07)	-0.14 (0.07)	-0.47** (0.07)	-0.11 (0.07)
eTIV onto Intercept ϕ	0.68** (0.05)	–	0.70** (0.05)	–
Brain-cognition links (cross-domain)	$STR-LS_{ime}$	$STR-LS_{del}$	$HPC-LS_{ime}$	$HPC-LS_{del}$
wave 1 covariation ϕ	**0.19* (0.07)**	**0.18* (0.07)**	0.12 (0.07)	**0.14* (0.07)**
change-change covariance ρ	<0.01 (0.03)	<0.01 (0.03)	-0.06 (0.05)	-0.07 (0.05)
wave 1 brain onto cognition change γ	0.25 (0.13)	0.22 (0.12)	0.05 (0.11)	0.06 (0.10)
wave 1 cognition onto brain change γ	-0.19 (0.13)	0.21 (0.13)	0.05 (0.10)	<0.01 (0.10)
Brain links (within-domain)	$STR-HPC$			
wave 1 covariation ϕ	**0.53** (0.07)**			
change-change covariance ρ	**0.03* (0.01)**			
wave 1 striatum onto hippocampal change γ	0.06 (0.05)			
wave 1 hippocampus onto striatal change γ	0.02 (0.03)			
Cognition links (within-domain)	$LS_{ime}-LS_{del}$			
wave 1 covariation ϕ	**0.95** (0.10)**			
change-change covariance ρ	**0.82** (0.10)**			
wave 1 LS_{ime} into LS_{del} change γ	-0.07 (0.27)			
wave 1 LS_{del} into LS_{ime} change γ	0.06 (0.28)			

Parameter estimates in bold are the paths of interest depicted in Figure 5D. Standard errors are shown in

parentheses. eTIV = estimated total intracranial volume. ** denotes significance at $\alpha < .001$, * at $\alpha < .05$. sex

coded as 1 = girls, -1 = boys.

Discussion

In this study, we examined the longitudinal development of value-based learning in middle childhood

and its associations with striatal and hippocampal volumes that were predicted to differ by feedback

timing. Children improved their learning in the 2-year study period. Behaviorally, learning was

improved by an increase in accuracy and a reduction in reaction time (i.e., faster responses). Further,

children's switching behavior improved by an increase in win-stay and a decrease in lose-shift behavior.

Computationally, learning was enhanced by an increase in learning rate and inverse temperature, which together constituted more optimal value-based learning. Further, feedback timing modulated specifically the inverse temperature. In terms of brain structures, we found that longitudinal changes in hippocampal volume were larger compared to striatal volume, which suggests more protracted hippocampal maturation. The brain-cognition links were longitudinally stable and partially confirmed our hypotheses. In line with previous adult literature and our assumption, hippocampal volume was more strongly associated with delayed feedback learning. Contrary to our expectations, episodic memory performance was not enhanced under delayed feedback compared to immediate feedback. Furthermore, striatal volume unexpectedly was associated with both immediate and delayed feedback learning, suggesting a common involvement of the striatum during value-based learning in middle childhood across timescales.

Children's learning improvement between waves was described behaviorally by increased win-stay and decreased lose-shift behavior. Our finding is in line with cross-sectional studies in the developmental literature that reported increased learning accuracy and win-stay behavior[58,59]. Our longitudinal dataset with younger children further suggests that learning change is not only accompanied by increased win-stay, but also decreased lose-shift behavior. We found lower learning performance and less optimal switching behavior in girls compared to boys, which could point to sex differences for reinforcement learning during middle childhood (Supplementary Material 2). Previous studies have found both male and female advantages depending on their age and the type of learning task[38,60,61]. Alternatively, sex differences may have been driven by confounding variables not included in the analysis.

Computationally, we found longitudinally increased and more optimal learning rate and inverse temperature, as shown by simulation data, that add to the growing literature of developmental reinforcement learning[16]. Adult studies that examined feedback timing during reinforcement learning reported average learning rates range from 0.12 to 0.34[5,13,14], which are much closer to the simulated optimal learning rates of 0.29 than children's average learning rates of 0.02 and 0.05 at wave 1 and 2 in our study. Therefore, it is likely that individuals approach adult-like optimal learning rates later during adolescence. However, the differences in learning rate across studies have to be interpreted with caution. The differences in the task and the analysis approach may limit their comparability[15,27]. Task proporties such as the trial number per condition differed across studies. Our study included 32 trials per cue in each condition, while in adult studies, the trials per condition ranged from 28 to 100[5,13,14]. Optimal learning rates in a stable learning environment were at around 0.25 for 10 to 30 trials[15], another study reported a lower optimal learning rate of around 0.08 for 120 trials[62]. This may partly explain why in our case of 32 trials per condition and cue, optimal learning rates called for a relatively high optimal learning rate of 0.29, while in other studies, optimal learning rates may be lower. Regarding differences in the analysis approach, the hierarchical bayesian estimation approach used in our study produces more reliable results in comparison to maximum likelihood estimation[49], which had been used in some of the

previous adult studies and may have led to biased results towards extreme values. Taken together, our

study underscores the importance of using longitudinal data to examine developmental change as well

as the importance of simulation-based optimal parameters to interpret the direction of developmental

change.

Despite a relatively immature hippocampal structure in middle childhood, our results confirmed a

longitudinally stable association between hippocampal volume and delayed feedback learning. However,

episodic memory in this learning condition was not enhanced. This suggests a developmentally early

hippocampal contribution to value-based learning during delayed feedback, which does not modulate

episodic memory as much as compared to adults. Therefore, our study partially extends the findings

from the adult literature to middle childhood[5,12–14]. The reduced effect of delayed feedback on episodic

memory may be due to the protracted development of hippocampal maturation. In an aging study with

a similar task, older adults failed to exhibit enhanced episodic memory for objects presented during

delayed feedback trials, and they showed no enhanced hippocampal activation during delayed feedback

and[14]. Therefore, the findings converge nicely at both childhood and older adulthood, during which the

structural and functional integrity of hippocampus are known to be less optimal than at younger

adulthood[63–65].

Our brain-cognition links were only partially confirmed, as striatal volumes exhibited associations with

not just immediate learning scores, as we predicted, but also with delayed learning scores. This result

suggests that the striatum may be important for value-based learning in general rather than exhibiting a

selective association with immediate feedback learning. This is also what we found in an explorative

analysis that related the striatum to learning rate in general and further predicted longitudinal change in

learning rate (Supplemental Material 5). This overall reduced brain-behavior specificity could reflect

less differentiated memory systems during development, similar to findings from aging research. Here,

older adults exhibited stronger striatal and hippocampal co-activation during both implicit and explicit

learning, compared to more dissociable brain-behavior relationships in younger adults[66]. Interestingly,

even in young adults, clear dissociations between memory systems such as in non-human lesion studies

are uncommon, and factors like stress modulate their cooperative interaction[6,10,11,67,68]. Further, there are

methodological differences to previous studies that could explain why striatal volumes were not

uniquely associated with immediate learning in our study. For example, previous studies related reward

prediction errors to striatal and hippocampal activation[5,13,14], whereas we examined individual

differences in brain structure and the model-derived learning scores. Future functional neuroimaging

studies with children could further clarify whether children's memory systems are indeed less

differentiated and explain the attenuated modulation by feedback timing. Taken together, compared to

the adult literature, our results with children showed that the hippocampal structure was associated with

delayed feedback learning, but did not enhance episodic memory encoding, while the striatum generally

supported value-based learning. These findings point towards a developmental effect of less
differentiated and more cooperative memory systems in middle childhood.

Our computational modeling results revealed a separable effect of feedback timing on inverse
temperature, which suggests that the memory systems modulated learning during decision-making. The
reported behavioral differences in reaction time and their correlation to the inverse temperature further
support the idea of a decision-related mechanism, as we found children to respond faster during delayed
feedback trials and faster responding children also exhibited more value-guided choice behavior (i.e.
higher inverse temperature) during delayed compared to immediate feedback. The hippocampus may
contribute to a decision-related effect in the delayed feedback condition by facilitating the encoding and
retrieval of learned values[69]. This is in contrast to previous event-related fMRI and EEG studies
reporting feedback timing modulations at value update[5,13,14], which may be due to at least two reasons.
First, we did not include a functional brain measure to examine its differential engagement during the
choice and feedback phases. Second, in such a reinforcement learning task, disentangling model
parameters from the choice and feedback phases can be challenging, such as for the inverse temperature
and outcome sensitivity[70]. Taken together, hippocampal engagement at delayed feedback may enhance
outcome sensitivity as well as facilitate choice behavior through improved retrieval of action-outcome
associations. A mechanism facilitating retrieval seems especially relevant in our paradigm, where
multiple cues were learned and presented in a mixed order, thus creating a high memory load. To
summarize, our study results suggest that feedback timing could modulate decision-making in addition
to or as alternative to a mechanism at value update. However, disentangling the effects of inverse
temperature and outcome sensitivity is challenging and warrants careful interpretation. Future studies
might shed new light by examining neural activations at both task phases, by additionally modeling
reaction times using a drift-diffusion approach, or by choosing a task design that allows independent
manipulations of these phases and associated model parameters, e.g., by using different reward
magnitudes during reinforcement learning, or by studying outcome sensitivity without decision-making.

One aim of developmental investigations is to identify the emergence of brain and cognition dynamics,
such as the hippocampal-dependent and striatal-dependent memory systems, which have been shown to
engage during reinforcement learning depending on the delay in feedback delivery. Our longitudinal
study partially confirmed these brain-cognition links in middle childhood but with less specificity as
previously found in adults.
An early existing memory system dynamic, similar to that of adults, is relevant for applying
reinforcement learning principles at different timescales. In scenarios such as in the classroom, a teacher
may comment on a child's behavior immediately after the action or some moments later, in par with our
experimental manipulation of 1 second versus 5 seconds. Within such short range of delay in teachers'
feedback, children's learning ability during the first years of schooling may function equally well and

depend on the striatal-dependent memory system. However, we anticipate that the reliance on the

hippocampus will become even more pronounced when feedback is further delayed for longer time.

Children's capacity for learning over longer timescales relies on the hippocampal-dependent memory

system, which is still under development. This knowledge could help to better structure learning

according to their development. Furthermore, probabilistic learning from delayed feedback may be a

potential diagnostic tool to examine the hippocampal-dependent memory system during learning in

children at risk. Environmental factors such as stress[11] and socioeconomic status[39,71] have been shown

to affect hippocampal structure and function and may contribute to a heightened risk for

psychopathology in the long term[72–74]. Deficits in hippocampal-dependent learning may be particularly

relevant to psychopathology since dysfunctional behavior may arise from a tendency to prioritize short-

term consequences over long-term ones[75,76] and from the maladaptive application of previously learned

behavior in inappropriate contexts[77]. Interestingly, poor learners showed relatively less value-based

learning in favor of stronger simple heuristic strategies, and excluding them modulated the hippocampal-

dependent associations to learning and memory in our results. More studies are needed to further clarify

the relationship between hippocampus and psychopathology during cognitive and brain development.

Another key question is whether developmental trajectories observed cross-sectionally are also

confirmed by longitudinal results, such as for the learning rate and inverse temperature. Our results show

developmental improvements in these learning parameters in only two years. This suggests that the

initial two years of schooling constitute a dynamic period for feedback-based learning, in which

contingent feedback is important in shaping behavior and development.

Additional Information

Funding. This study was supported by the Jacobs Foundation [grant 2014–1151] to YLS and CH. The work of YLS was also supported by the European Union (ERC-2018-StG-PIVOTAL-758898), the Deutsche Forschungsgemeinschaft (German Research Foundation, Project ID 327654276, SFB 1315,'Mechanisms and Disturbances in Memory Consolidation: From Synapses to Systems'), and the Hessisches Ministerium für Wissenschaft und Kunst (HMWK; project 'The Adaptive Mind').

Acknowledgments. We thank the Max Planck Institute for Human Development and all members of the Jacobs study team for their vital contribution, and all participants and family members for taking part in the study.

Conflicts of interest. The authors declare no competing financial interests.

Ethics approval. This study was approved by the "Deutsche Gesellschaft für Psychologie" ethics committee (YLS_012015).

Availability of data and code. https://osf.io/pju65/

Author ORCIDs.
JF: https://orcid.org/0000-0003-0505-0798
LZ: https://orcid.org/0000-0002-9586-595X
LR: https://orcid.org/0000-0002-0144-5605
JJM: https://orcid.org/0000-0002-3893-8008
JT: https://orcid.org/0000-0001-8166-2441
CH: https://orcid.org/0000-0002-6580-6326
YLS: https://orcid.org/0000-0001-8922-7292

References

1. Sutton, R. S. & Barto, A. G. *Reinforcement learning: An introduction*. (MIT press, 2018).

2. Gläscher, J., Daw, N., Dayan, P. & O'Doherty, J. P. States versus Rewards: Dissociable neural

prediction error signals underlying model-based and model-free reinforcement learning.

Neuron **66**, 585 (2010).

3. Bolenz, F., Reiter, A. M. F. & Eppinger, B. Developmental Changes in Learning:

Computational Mechanisms and Social Influences. *Front. Psychol.* **0**, 2048 (2017).

4. Zhang, L. & Gläscher, J. A brain network supporting social influences in human decision-

making. *Sci. Adv.* **6**, 1–20 (2020).

5. Foerde, K. & Shohamy, D. Feedback Timing Modulates Brain Systems for Learning in

Humans. *J. Neurosci.* **31**, 13157–13167 (2011).

6. Packard, M. G. & Goodman, J. Factors that influence the relative use of multiple memory

systems. *Hippocampus* **23**, 1044–1052 (2013).

7. Goodman, J. & Packard, M. G. Memory Systems of the Basal Ganglia. *Handb. Behav.*

Neurosci. **24**, 725–740 (2016).

8. Davidow, J. Y., Foerde, K., Galván, A. & Shohamy, D. An Upside to Reward Sensitivity: The

Hippocampus Supports Enhanced Reinforcement Learning in Adolescence. *Neuron* **92**, 93–99

(2016).

9. Hartley, C. A., Nussenbaum, K. & Cohen, A. O. Interactive Development of Adaptive

Learning and Memory. 1–27 (2021).

10. Packard, M. G., Goodman, J. & Ressler, R. L. Emotional modulation of habit memory: neural

mechanisms and implications for psychopathology. *Curr. Opin. Behav. Sci.* **20**, 25–32 (2018).

11. Schwabe, L. & Wolf, O. T. Stress and multiple memory systems: from 'thinking' to 'doing'.

Trends Cogn. Sci. **17**, 60–68 (2013).

12. Foerde, K., Race, E., Verfaellie, M. & Shohamy, D. A role for the medial temporal lobe in

feedback-driven learning: Evidence from amnesia. *J. Neurosci.* **33**, 5698–5704 (2013).

13. Höltje, G. & Mecklinger, A. Feedback timing modulates interactions between feedback

processing and memory encoding: Evidence from event-related potentials. *Cogn. Affect. Behav.*

Neurosci. 2020 202 **20**, 250–264 (2020).

14. Lighthall, N. R., Pearson, J. M., Huettel, S. A. & Cabeza, R. Feedback-Based Learning in

Aging: Contributions and Trajectories of Change in Striatal and Hippocampal Systems. *J.*

Neurosci. **38**, 8453–8462 (2018).

15. Zhang, L., Lengersdorff, L., Mikus, N., Gläscher, J. & Lamm, C. Using reinforcement learning

models in social neuroscience: Frameworks, pitfalls and suggestions of best practices. *Soc.*

Cogn. Affect. Neurosci. **15**, 695–707 (2020).

16. Nussenbaum, K. & Hartley, C. A. Reinforcement learning across development: What insights

can we draw from a decade of research? *Developmental Cognitive Neuroscience* **40**, (2019).

17.   Decker, J. H., Lourenco, F. S., Doll, B. B. & Hartley, C. A. Experiential reward learning
819   outweighs instruction prior to adulthood. *Cogn. Affect. Behav. Neurosci.* **15**, 310–320 (2015).

820   18.   Javadi, A. H., Schmidt, D. H. K. & Smolka, M. N. Differential representation of feedback and
821   decision in adolescents and adults. *Neuropsychologia* **56**, 280–288 (2014).

822   19.   Palminteri, S., Kilford, E. J., Coricelli, G. & Blakemore, S. J. The Computational Development
823   of Reinforcement Learning during Adolescence. *PLoS Comput. Biol.* **12**, 1–25 (2016).

824   20.   Master, S. L. *et al.* Distentangling the systems contributing to changes in learning during
825   adolescence. *Dev. Cogn. Neurosci.* **41**, 100732 (2020).

826   21.   Hauser, T. U., Iannaccone, R., Walitza, S., Brandeis, D. & Brem, S. Cognitive flexibility in
827   adolescence: Neural and behavioral mechanisms of reward prediction error processing in
828   adaptive decision making during development. *Neuroimage* **104**, 347–354 (2015).

829   22.   Moutoussis, M. *et al.* Change, stability, and instability in the Pavlovian guidance of behaviour
830   from adolescence to young adulthood. *PLoS Comput. Biol.* **14**, (2018).

831   23.   Van Den Bos, W., Cohen, M. X., Kahnt, T. & Crone, E. A. Striatum-medial prefrontal cortex
832   connectivity predicts developmental changes in reinforcement learning. *Cereb. Cortex* **22**,
833   1247–1255 (2012).

834   24.   Rodriguez Buritica, J. M., Heekeren, H. R. & van den Bos, W. The computational basis of
835   following advice in adolescents. *J. Exp. Child Psychol.* **180**, 39–54 (2019).

836   25.   Galván, A. The Teenage Brain: Sensitivity to Rewards. *Curr. Dir. Psychol. Sci.* **22**, 88–93
837   (2013).

838   26.   van Duijvenvoorde, A. C. K. *et al.* A cross-sectional and longitudinal analysis of reward-
839   related brain activation: Effects of age, pubertal stage, and reward sensitivity. *Brain Cogn.* **89**,
840   3–14 (2014).

841   27.   Eckstein, M. K., Wilbrecht, L. & Collins, A. G. E. What do RL Models Measure ? Interpreting
842   Model Parameters in Cognition and Neuroscience. *Curr. Opin. Behav. Sci.* **41**, 128–137 (2021).

843   28.   Cohen, A. O., Nussenbaum, K., Dorfman, H. M., Gershman, S. J. & Hartley, C. A. The rational
844   use of causal inference to guide reinforcement learning strengthens with age. *npj Sci. Learn.* **5**,
845   1–9 (2020).

846   29.   Raznahan, A. *et al.* Longitudinal four-dimensional mapping of subcortical anatomy in human
847   development. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 1592 (2014).

848   30.   Wierenga, L. *et al.* Typical development of basal ganglia, hippocampus, amygdala and
849   cerebellum from age 7 to 24. *Neuroimage* **96**, 67–72 (2014).

850   31.   Giedd, J. N. Structural Magnetic Resonance Imaging of the Adolescent Brain. *Ann. N. Y. Acad.*
851   *Sci.* **1021**, 77–85 (2004).

852   32.   Uematsu, A. *et al.* Developmental Trajectories of Amygdala and Hippocampus from Infancy to
853   Early Adulthood in Healthy Individuals. *PLoS One* **7**, e46970 (2012).

854   33.   Giedd, J. N. *et al.* Child Psychiatry Branch of the National Institute of Mental Health

855   Longitudinal Structural Magnetic Resonance Imaging Study of Human Brain Development.

856   *Neuropsychopharmacology* **40**, 43 (2015).

857   34.   Goodman, J., Marsh, R., Peterson, B. S. & Packard, M. G. Annual research review: The

858   neurobehavioral development of multiple memory systems--implications for childhood and

859   adolescent psychiatric disorders. *J. Child Psychol. Psychiatry.* **55**, 582–610 (2014).

860   35.   Goddings, A. L. *et al.* The influence of puberty on subcortical brain development. *Neuroimage*

861   **88**, 242–251 (2014).

862   36.   Dima, D. *et al.* Subcortical volumes across the lifespan: Data from 18,605 healthy individuals

863   aged 3–90 years. *Hum. Brain Mapp.* 1–18 (2021). doi:10.1002/hbm.25320

864   37.   Lavenex, P. & Banta Lavenex, P. Building hippocampal circuits to learn and remember:

865   Insights into the development of human memory. *Behavioural Brain Research* **254**, 8–21

866   (2013).

867   38.   Mandolesi, L., Petrosini, L., Menghini, D., Addona, F. & Vicari, S. Children' s radial arm

868   maze performance as a function of age and sex. *Int. J. Dev. Neurosci.* **27**, 789–797 (2009).

869   39.   Raffington, L. *et al.* Stable longitudinal associations of family income with children's

870   hippocampal volume and memory persist after controlling for polygenic scores of educational

871   attainment. *Dev. Cogn. Neurosci.* **40**, 100720 (2019).

872   40.   Raffington, L. *et al.* Effects of stress on 6- and 7-year-old children's emotional memory differs

873   by gender. *J. Exp. Child Psychol.* **199**, 104924 (2020).

874   41.   Fischl, B. FreeSurfer. *Neuroimage* **62**, 774–781 (2012).

875   42.   Phan, T. V., Smeets, D., Talcott, J. B. & Vandermosten, M. Processing of structural

876   neuroimaging data in young children: Bridging the gap between current practice and state-of-

877   the-art methods. *Dev. Cogn. Neurosci.* **33**, 206–223 (2018).

878   43.   Schoemaker, D. *et al.* Hippocampus and amygdala volumes from magnetic resonance images

879   in children: Assessing accuracy of FreeSurfer and FSL against manual segmentation.

880   *Neuroimage* **129**, 1–14 (2016).

881   44.   Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting Linear Mixed-Effects Models Using

882   lme4. *J. Stat. Softw.* **67**, 1–48 (2015).

883   45.   Brown, V. M. *et al.* Reinforcement Learning Disruptions in Individuals with Depression and

884   Sensitivity to Symptom Change following Cognitive Behavioral Therapy. *JAMA Psychiatry*

885   (2021). doi:10.1001/jamapsychiatry.2021.1844

886   46.   Stan Development Team. RStan: the R interface to Stan. R package version 2.21.2. *http//mc-*

887   *stan. org* (2021).

888   47.   R Core Team. R: A Language and Environment for Statistical Computing. (2021).

889   48.   Ahn, W.-Y., Haines, N. & Zhang, L. Revealing Neurocomputational Mechanisms of

890   Reinforcement Learning and Decision-Making With the hBayesDM Package. *Comput.*

891   *Psychiatry* **1**, 24 (2017).

892   49.   Brown, V. M., Chen, J., Gillan, C. M. & Price, R. B. Improving the Reliability of
893   Computational Analyses: Model-Based Planning and Its Relationship With Compulsivity. *Biol.*
894   *Psychiatry Cogn. Neurosci. Neuroimaging* **5**, 601–609 (2020).

895   50.   Vehtari, A., Gelman, A. & Gabry, J. Practical Bayesian model evaluation using leave-one-out
896   cross-validation and WAIC. *Stat. Comput.* **27**, 1413–1432 (2017).

897   51.   Yao, Y., Vehtari, A., Simpson, D. & Gelman, A. Using Stacking to Average Bayesian
898   Predictive Distributions (with Discussion). *Bayesian Anal.* **13**, 917–1007 (2018).

899   52.   Crawley, D. *et al.* Modeling flexible behavior in childhood to adulthood shows age-dependent
900   learning mechanisms and less optimal learning in autism in each age group. *PLoS Biol.* **18**, 1–
901   25 (2020).

902   53.   Wilson, R. C. & Collins, A. G. E. Ten simple rules for the computational modeling of
903   behavioral data. *Elife* **8**, 1–33 (2019).

904   54.   Kievit, R. A. *et al.* Developmental cognitive neuroscience using latent change score models: A
905   tutorial and applications. *Dev. Cogn. Neurosci.* **33**, 99–117 (2018).

906   55.   Ferrer, E. & McArdle, J. J. Longitudinal modeling of developmental changes in psychological
907   research. *Curr. Dir. Psychol. Sci.* **19**, 149–154 (2010).

908   56.   Sluis, S. van der, Verhage, M., Posthuma, D. & Dolan, C. V. Phenotypic Complexity,
909   Measurement Bias, and Poor Phenotypic Resolution Contribute to the Missing Heritability
910   Problem in Genetic Association Studies. *PLoS One* **5**, e13929 (2010).

911   57.   Little, T. *Longitudinal structural equation modeling*. (Guilford Press, 2013).
912   doi:10.2/JQUERY.MIN.JS

913   58.   Chierchia, G. *et al.* Confirmatory reinforcement learning changes with age during adolescence.
914   *Dev. Sci.* e13330 (2021). doi:10.1111/desc.13330

915   59.   Habicht, J., Bowler, A., Moses-Payne, M. E. & Hauser, T. U. Children are full of optimism, but
916   those rose-tinted glasses are fading – reduced learning from negative outcomes drives
917   hyperoptimism in children. (2021).

918   60.   Overman, W. H. Sex differences in early childhood, adolescence, and adulthood on cognitive
919   tasks that rely on orbital prefrontal cortex. *Brain Cogn.* **55**, 134–147 (2004).

920   61.   Evans, K. L. & Hampson, E. Sex-dependent effects on tasks assessing reinforcement learning
921   and interference inhibition. *Front. Psychol.* **6**, 1–10 (2015).

922   62.   Behrens, T. E. J., Woolrich, M. W., Walton, M. E. & Rushworth, M. F. S. Learning the value
923   of information in an uncertain world. *Nat. Neurosci.* **10**, 1214–1221 (2007).

924   63.   Shing, Y. L. *et al.* Episodic memory across the lifespan: The contributions of associative and
925   strategic components. *Neurosci. Biobehav. Rev.* **34**, 1080–1091 (2010).

926   64.   Keresztes, A. *et al.* Hippocampal maturity promotes memory distinctiveness in childhood and
927   adolescence. *Proc. Natl. Acad. Sci. U. S. A.* **114**, 9212–9217 (2017).

928   65.   Ghetti, S. & Bunge, S. A. Neural changes underlying the development of episodic memory

during middle childhood. *Dev. Cogn. Neurosci.* **2**, 381–395 (2012).

66. Dennis, N. A. & Cabeza, R. Age-related dedifferentiation of learning systems: An fMRI study of implicit and explicit learning. *Neurobiol. Aging* **32**, 2318.e17-2318.e30 (2011).

67. Ferbinteanu, J. Contributions of Hippocampus and Striatum to Memory-Guided Behavior Depend on Past Experience. *J. Neurosci.* **36**, 6459–6470 (2016).

68. White, N. M. & McDonald, R. J. Multiple Parallel Memory Systems in the Brain of the Rat. *Neurobiol. Learn. Mem.* **77**, 125–184 (2002).

69. Shadlen, M. N. N. & Shohamy, D. Decision Making and Sequential Sampling from Memory. *Neuron* **90**, 927–939 (2016).

70. Browning, M., Paulus, M. & Huys, Q. J. M. What is computational psychiatry good for? *Biol. Psychiatry* **0**, (2022).

71. Hackman, D. A., Farah, M. J. & Meaney, M. J. Socioeconomic status and the brain: mechanistic insights from human and animal research. *Nat. Rev. Neurosci. 2010 119* **11**, 651–659 (2010).

72. Frodl, T. *et al.* Childhood stress, serotonin transporter Gene and Brain structures in major depression. *Neuropsychopharmacology* **35**, 1383–1390 (2010).

73. Lucassen, P. J., Korosi, A., Krugers, H. J. & Oomen, C. A. *Early Life Stress- and Sex-Dependent Effects on Hippocampal Neurogenesis. Stress: Neuroendocrinology and Neurobiology* **2**, (Elsevier Inc., 2017).

74. Rahman, M. M., Callaghan, C. K., Kerskens, C. M., Chattarji, S. & O'Mara, S. M. Early hippocampal volume loss as a marker of eventual memory deficits caused by repeated stress. *Sci. Rep.* **6**, 1–15 (2016).

75. Levin, M. E., Haeger, J., Ong, C. W. & Twohig, M. P. An Examination of the Transdiagnostic Role of Delay Discounting in Psychological Inflexibility and Mental Health Problems. *Psychol. Rec.* **68**, 201–210 (2018).

76. Von Siebenthal, Z. *et al.* Decision-making impairments following insular and medial temporal lobe resection for drug-resistant epilepsy. *Soc. Cogn. Affect. Neurosci.* **12**, 128–137 (2017).

77. Maren, S., Phan, K. L. & Liberzon, I. The contextual brain: Implications for fear conditioning, extinction and psychopathology. *Nat. Rev. Neurosci.* **14**, 417–428 (2013).