RBP-J regulates homeostasis and function of circulating Ly6C[lo] monocytes

Tiantian Kou[1,2,3,6], Lan Kang[1,3,6], Bin Zhang[1,3], Jiaqi Li[1], Baohong Zhao[4,5], Wenwen Zeng[1,2,3] and Xiaoyu Hu[1,2,3,7, *]

Affiliations:

[1]Institute for Immunology and School of Medicine, Tsinghua University, Beijing, China.

[2]Tsinghua-Peking Center for Life Sciences, Tsinghua University, Beijing, China.

[3]Beijing Key Laboratory for Immunological Research on Chronic Diseases, Beijing, China.

[4]Arthritis and Tissue Degeneration Program and the David Z. Rosensweig Genomics Research Center, Hospital for Special Surgery, New York, NY.

[5]Department of Medicine, Weill Cornell Medicine, New York, NY.

[6]These authors contributed equally.

[7]Lead Contact

[*]Correspondence: xiaoyuhu@tsinghua.edu.cn.

Summary

Notch-RBP-J signaling plays an essential role in maintenance of myeloid homeostasis. However, its role in monocyte cell fate decisions is not fully understood. Here we showed that conditional deletion of transcription factor RBP-J in myeloid cells resulted in marked accumulation of blood Ly6C^{lo} monocytes that highly expressed chemokine receptor CCR2. Bone marrow transplantation and parabiosis experiments revealed a cell intrinsic requirement of RBP-J for controlling blood Ly6C^{lo}CCR2^{hi} monocytes. RBP-J-deficient Ly6C^{lo} monocytes exhibited enhanced capacity competing with wildtype counterparts in blood circulation. In accordance with alterations of circulating monocytes, RBP-J deficiency led to markedly increased population of lung tissues with Ly6C^{lo} monocytes and CD16.2^{+} interstitial macrophages. Furthermore, RBP-J deficiency-associated phenotypes could be genetically corrected by further deleting *Ccr2* in myeloid cells. These results demonstrate that RBP-J functions as a crucial regulator of blood Ly6C^{lo} monocytes and thus derived lung-resident myeloid populations, at least in part through regulation of CCR2.

Introduction

Monocytes are integral components of the mononuclear phagocyte system (MPS) that develop from monocyte precursors in the bone marrow[1]. Several functionally and phenotypically distinct subsets of blood monocytes have been defined on the basis of expression of surface markers[2-5]. In mice, Ly6C[hi] monocytes (also called inflammatory or classical monocytes) characterized by Ly6C[hi]CCR2[hi]CX3CR1[lo] exhibit a short half-life, and are recruited to tissue and differentiate into macrophages and dendritic cells[6]. Ly6C[hi] monocytes are analogous to CD14[+] human monocytes based on gene expression profiling[7]. A second subtype of monocytes is called Ly6C[lo] monocytes (also termed patrolling monocytes or non-classical monocytes), which express high level of CX3CR1 and low level of CCR2, equivalent to CD14[lo]CD16[+] human monocytes. Ly6C[lo] monocytes are regarded as blood-resident macrophages that patrol blood vessels and scavenge microparticles attached to the endothelium under physiological conditions, and exhibit a long half-life in the steady-state[8-10]. Ly6C[lo] monocytes may exert a protective effect by suppressing atherosclerosis in mice, while high levels of nonclassical monocytes have been associated with more advanced vascular dysfunction and oxidative stress in patients with coronary artery disease[11-13]. In the inflammatory settings, Ly6C[lo] monocytes often show anti-inflammatory properties, yet also exhibit a proinflammatory role under certain circumstances[5,9,14-18]. In addition, Ly6C[lo] monocytes have demonstrated protective functions in tumorigenesis, such as engulfing tumor materials to prevent cancer metastasis, activating and recruiting NK cells to the lungs[19,20]. While the functions of Ly6C[lo] monocytes are complex and sometimes contradictory depending on the animal models and experimental approaches, it is clear that they play a crucial role in both health and disease.

The generation and maintenance of Ly6C[lo] monocytes are regulated by several transcription factors, including Nr4a1 and CEBPβ[21,22]. The colony-stimulating factor 1 receptor (CSF1R) signaling pathway is important for the generation and survival of Ly6C[lo] monocytes, and neutralizing antibodies against CSF1R can reduce their numbers[23]. Interestingly, recent studies have suggested that different subsets of

75   monocytes may be supported by distinct cellular sources of CSF-1 within bone marrow

niches, in which targeted deletion of *Csf1* from sinusoidal endothelial cells selectively

reduces Ly6C[lo] monocytes but not Ly6C[hi] monocytes[24]. LAIR1 has been shown to be

activated by stromal protein Colec12, and LAIR1 deficiency leads to aberrant

proliferation and apoptosis in bone marrow non-classical monocytes[25]. These findings

80   further underscore the complexity of the mechanisms that regulate Ly6C[lo] monocyte,

and suggest that multiple factors are involved in this process.

    Recombination signal-binding protein for immunoglobulin kappa J region (RBP-J;

also named CSL in humans) is commonly known as the master nuclear mediator of

85   canonical Notch signaling[26]. In the absence of Notch intracellular domain, RBP-J

associates with co-repressor proteins to repress transcription of downstream target

genes. In the immune system, the best studied functions of Notch signaling are its roles

in regulating the development of lymphocytes, such as T cells and marginal zone B

cells[26]. Notch signaling also has been reported to regulate the differentiation and

90   function of myeloid cells including granulocyte/monocyte progenitors, osteoclasts and

dendritic cells [27-30]. In osteoclasts, RBP-J represses osteoclastogenesis, particularly

under inflammatory conditions[30]. In dendritic cells, Notch-RBP-J signaling controls the

maintenance of CD8[-] DC[28]. In addition, Notch2-RBP-J pathway is essential for

development of CD8[-]ESAM[+] DC in the spleen and CD103[+]CD11b[+] DC in the lamina

95   propria of the intestine[29]. However, the role of Notch-RBP-J signaling pathway in

regulating monocyte subsets remains elusive. Here, we demonstrated that RBP-J

controlled homeostasis of blood Ly6C[lo] monocytes in a cell intrinsic manner. RBP-J

deficiency led to a drastic increase in blood Ly6C[lo] monocytes, lung Ly6C[lo] monocytes

and CD16.2[+] IM. Our results suggest that RBP-J plays a critical role in regulating the

100   population and characteristics of Ly6C[lo] monocytes in the blood and lung.

Results

RBP-J is essential for maintenance of blood Ly6C[lo] monocytes

To investigate the role of RBP-J in monocyte subsets, we utilized mice with RBP-J

specific deletion in the myeloid cells (*Rbpj*^{fl/fl}*Lyz2*^{cre/cre} mice). Efficient deletion of *Rbpj*
in blood monocytes was confirmed by quantitative real-time PCR (qPCR) (Figure 1 -
figure supplement 1A). Flow cytometric analysis revealed that RBP-J deficient mice
had a significant increase in the proportion of Ly6C^{lo} monocytes, but not Ly6C^{hi}
monocytes, in blood, compared to age-matched control mice with the genotype
Rbpj^{+/+}*Lyz2*^{cre/cre} (Figure 1A). In contrast to circulating monocytes, RBP-J deficient
mice exhibited minimal alterations in the percentages of monocyte subsets in bone
marrow (BM) and spleen (Figure 1B; Figure 1 - figure supplement 1C). Additionally,
RBP-J deficiency in myeloid cells did not appear to have an effect on neutrophils
(Figure 1- figure supplement 1B). Therefore, among circulating myeloid populations,
RBP-J selectively controlled the subset of Ly6C^{lo} monocytes.

BM progenitors that give rise to circulating monocytes are monocyte-dendritic cell
progenitors (MDPs) and common monocyte progenitors (cMoPs)[21,31-33]. Next, we
analyzed BM progenitors, and found that the percentages of MDPs and cMoPs were
equivalent in control and RBP-J deficient mice (Figure 1C). These results in
conjunction with the observations of normal BM monocyte populations suggest that
alterations of peripheral blood Ly6C^{lo} monocytes may not originate from BM. Next, we
examined whether RBP-J may influence the egress of Ly6C^{lo} monocytes from BM, and
measured BM exit rate of Ly6C^{lo} monocytes using *in vivo* labeling of sinusoidal cells
as previously described (Figure 1D)[34]. The results showed that RBP-J deficiency did
not affect egress of Ly6C^{lo} monocytes from BM (Figure 1E). Taken together, these data
revealed RBP-J as a critical regulator controlling homeostasis of peripheral blood
Ly6C^{lo} monocytes.

RBP-J deficiency does not affect Ly6C^{hi} monocyte conversion or Ly6C^{lo} monocyte
survival and proliferation

Next, we aimed to determine whether the increase of blood Ly6C^{lo} monocytes in RBP-
J deficient mice was due to decreased cell death or enhanced proliferation. We first
stained monocytes with Annexin V and 7-AAD to identify apoptotic cells and observed

comparable percentages of apoptotic blood Ly6C[lo] monocytes in control and RBP-J deficient mice (Figure 2A). Given the crucial role of Nr4a1 in the survival of Ly6C[lo] monocytes[21], we detected the expression of Nr4a1, which was similar in Ly6C[lo] monocytes from control and RBP-J deficient mice (Figure 2- figure supplement 1A). We then assessed the proliferative capacity by analyzing Ki-67 expression as well as *in vivo* EdU incorporation. Ki-67 levels in blood monocytes displayed no differences between control and RBP-J deficient mice (Figure 2B). The percentage of EdU[+] monocytes did not significantly differ between control and RBP-J deficient mice at the indicated time points, implying that the turnover of Ly6C[lo] monocytes was normal in RBP-J deficient mice (Figure 2C). Fluorescent latex beads as particulate tracers could be phagocytosed by monocytes after intravenous injection, and stably label Ly6C[lo] monocytes[35]. Thus, we intravenously injected fluorescent latex beads into control and RBP-J deficient mice to track circulating monocytes (Figure 2D). By day 7 post injection, only Ly6C[lo] monocytes were latex[+] as previously reported[35], whereas control and RBP-J deficient mice groups presented a similar frequency of latex[+] monocytes (Figure 2E). Together, these data indicated that RBP-J did not influence monocyte survival and proliferation.

Previous studies have shown that Ly6C[lo] monocytes are observed in recipient mice following the adoptive transfer of Ly6C[hi] monocytes, indicating that Ly6C[hi] monocytes can convert into Ly6C[lo] monocytes[10,31]. We next wished to evaluate whether conversion of Ly6C[hi] monocytes was regulated by RBP-J by isolating bone marrow GFP[+]Ly6C[hi] monocytes from *Lyz2*[cre/cre]*Cx3cr1*[gfp/+] control or *Rbpj*[fl/fl]*Lyz2*[cre/cre]*Cx3cr1*[gfp/+] mice and adoptively transferring them into *Ccr2*[RFP/RFP] recipients (Figure 2F). Sixty hours after transfer, a subset of cells from donors were converted into Ly6C[lo] monocytes (Figure 2 - figure supplement 1B), and equal percentages of Ly6C[lo] monocytes were derived from control and *Rbpj*[fl/fl]*Lyz2*[cre/cre]*Cx3cr1*[gfp/+] donors (Figure 2G). Thus, the conversion of Ly6C[hi] monocytes into Ly6C[lo] monocytes was not affected by RBP-J deficiency.

RBP-J regulates blood Ly6C[lo] monocytes in a cell intrinsic manner

165   We next wondered whether the increase of Ly6C^{lo} monocytes in RBP-J deficient mice was bone marrow-derived and cell intrinsic. We performed bone marrow transplantation by engrafting lethally irradiated mice with a 1:4 mixture of *Rbpj$^{+/+}$Lyz2$^{cre/cre}$* control and *Rbpj$^{fl/fl}$Lyz2$^{cre/cre}$* BM cells (CD45.2) and *Cx3cr1$^{gfp/+}$* BM cells (CD45.1) (Figure 3A). Eight weeks after transplantation, we analyzed the

170   frequencies of CD45.2^{+} donor cells in the bone marrow and blood of recipient mice and found that the frequencies of CD45.2^{+} cells within total cells were similar to the mixture ratio of bone marrow cells for Ly6C^{hi} monocytes and neutrophils in both BM and blood (Figure 3B; Figure 3 - figure supplement 1A-B). Specifically, more blood Ly6C^{lo} monocytes were derived from RBP-J deficient donors than control donors (Figure 3B),

175   reflecting that the contribution of RBP-J deficient cells in blood Ly6C^{lo} cells was significantly higher than that of control cells. These results implied that RBP-J deficient BM cells were highly efficient in generating blood Ly6C^{lo} monocytes.

      Given that RBP-J deficient mice had more Ly6C^{lo} monocytes in blood than did

180   control mice, we performed parabiosis experiments, a surgical union of two organisms, which allowed parabiotic mice to share their blood circulation. The contribution of circulating cells from one animal in another can be estimated by measuring the percentage of blood cells that originated from each animal[36]. We joined a CD45.1, *Cx3cr1$^{gfp/+}$* mouse with an age and sex-matched *Rbpj$^{+/+}$Lyz2$^{cre/cre}$* control or

185   *Rbpj$^{fl/fl}$Lyz2$^{cre/cre}$* mouse (CD45.2) (Figure 3C). Four weeks after the procedure, about 50% of B cells and T cells in parabiotic mice displayed efficient exchange of their circulation (Figure 3 - figure supplement 1C). As expected, RBP-J deficient mice exhibited higher percentages of Ly6C^{lo} monocytes than control animals (Figure 3D). Intriguingly, in the parabiotic *Cx3cr1$^{gfp/+}$* mice, RBP-J deficient cells still constituted

190   significantly higher proportion of circulating Ly6C^{lo}, but not Ly6C^{hi} monocytes, than RBP-J sufficient counterparts as indicated by the percentages of the GFP-negative population (Figure 3E). These results implicated that the enhanced ability of Ly6C^{lo} monocytes to circulate in the peripheral blood as a result of RBP-J deficiency was cell intrinsic.

RBP-J regulates phenotypical marker genes in blood Ly6C^{lo} monocytes

To further study the consequences of RBP-J loss of function in blood Ly6C^{lo} monocytes, we performed gene expression profiling by RNA-seq, which revealed that Ly6C^{lo} monocytes in RBP-J deficient mice exhibited low expression of *Itgax* but high expression of *Ccr2*, in comparison to those in control mice (Figure 4A). These differential expression patterns of phenotypic marker genes were also confirmed by qPCR and flow cytometry analysis (Figure 4B). However, Ly6C^{hi} blood monocytes and BM monocytes displayed normal levels of CD11c and CCR2 (Figure 4B; Figure 4 - figure supplement 1A). Moreover, principal component analysis showed distinct expression pattern of monocytes between control and RBP-J deficient mice (Figure 4C). To determine whether the regulation of phenotypic markers by RBP-J in blood Ly6C^{lo} monocytes was cell-intrinsic, we performed BM transplantation as described above (Figure 3A). The findings indicated that Ly6C^{lo} monocytes derived from RBP-J deficient BM cells expressed high levels of CCR2 but low levels of CD11c, in comparison to those derived from control BM cells, whereas no significant changes were observed in blood Ly6C^{hi} and BM monocytes, as expected (Figure 4D; Figure 4 - figure supplement 1B). These results collectively suggested that RBP-J regulated the expression of CCR2/CD11c in a cell-intrinsic manner.

RBP-J-mediated control of blood Ly6C^{lo} monocytes is CCR2 dependent

Given that RBP-J deficiency led to enhanced CCR2 expression in Ly6C^{lo} monocytes and that CCR2 is essential for monocyte functionality under various inflammatory and non-inflammatory conditions[37], we wished to genetically investigate the role of CCR2 in the RBP-J deficient background. We generated RBP-J/CCR2 double deficient (DKO) mice with the genotype *Rbpj^{fl/fl}Lyz2^{cre/cre}Ccr2^{RFP/RFP}* with *Lyz2^{cre/cre}Ccr2^{RFP/+}* and *Rbpj^{fl/fl}Lyz2^{cre/cre}Ccr2^{RFP/+}* mice serving as controls. As expected, the expression of CCR2 in Ly6C^{lo} monocytes was reduced in DKO mice compared to RBP-J deficient mice (Figure 5A), confirming the successful deletion of the *Ccr2* gene. DKO mice showed a lower percentage of both Ly6C^{hi} and Ly6C^{lo} monocytes than RBP-J deficient

mice (Figure 5B-C). Notably, the percentage of Ly6C^{lo} monocytes in DKO mice was comparable to that observed in the control mice (Figure 5B-C), implicating that deletion of CCR2 corrected RBP-J deficiency-associated phenotype of increased Ly6C^{lo} monocytes. Whereas, both RBP-J deficient and DKO mice exhibited lower expression levels of CD11c in their Ly6C^{lo} monocyte than control mice, suggesting that the regulation of CD11c expression was independent of CCR2 (Figure 5D). These results suggested that RBP-J regulated Ly6C^{lo} monocytes, at least in part, through CCR2.

Lung Ly6C^{lo} monocytes are accumulated in RBP-J deficient mice

In mice, alveolar macrophages (AM) are maintained by local self-renewal, and arise from fetal liver derived precursors, whereas interstitial macrophages (IM) are probably originated from monocytes [38-40]. Joey et al identified 3 subpopulations of IM according to expression of CD206 and CD16.2 and Ly6C^{lo} monocytes are proposed to give rise to CD64$^+$CD16.2$^+$ IM[41]. We next compared the populations of lung monocytes and IM between control and RBP-J deficient mice. The conditional deletion of RBP-J resulted in a significant increase in the absolute and relative numbers of Ly6C^{lo} monocytes and CD16.2$^+$ IM, while the numbers of Ly6C^{hi} monocytes, CD206$^-$ IM, and CD206$^+$ IM remained unchanged (Figure 6A-B). To confirm these findings, lung sections from *Lyz2*$^{cre/cre}$*Cx3cr1*$^{gfp/+}$ control and *Rbpj*$^{fl/fl}$*Lyz2*$^{cre/cre}$*Cx3cr1*$^{gfp/+}$ mice were stained with an anti-GFP antibody, which revealed a significant increase in the number of GFP$^+$ cells in RBP-J deficient mice (Figure 6C). In addition, we evaluated the proliferative capacity of Ly6C^{lo} monocytes and CD16.2$^+$ IM, but did not observe enhanced EdU incorporation in Ly6C^{lo} monocytes and CD16.2$^+$ IM from RBP-J deficient mice (Figure 6 - figure supplement 1A-B). These findings suggested that the augmented population of these cells might not be resulted from increased *in situ* proliferation. Previous reports have indicated that LPS can increase the population of IM in a CCR2-dependent manner[40]. We thus challenged mice with LPS, and analyzed monocytes and IM at day 0 and day 4. LPS exposure induced an increase in numbers of Ly6C^{lo} monocytes, CD16.2$^+$ IM and CD206$^-$ IM compared to baseline both in control and RBP-J deficient mice (Figure 6D-E). There was a trend toward higher Ly6C^{hi} monocyte after LPS treatment, although

255 this observation was not statistically significant. Of note, RBP-J deficient mice exhibited robustly elevated numbers of Ly6C^{lo} monocytes and CD16.2$^+$ IM compared with control mice after LPS treatment (Figure 6D-E), suggesting that RBP-J deficient Ly6C^{lo} monocytes were recruited to inflamed tissue in large numbers and differentiated into CD16.2$^+$ IM.

To investigate whether the elevated levels of lung monocytes and CD16.2$^+$ IM in RBP-J deficient mice were derived from blood Ly6C^{lo} monocytes, we examined monocytes and IM in *Lyz2*$^{cre/cre}$*Ccr2*$^{RFP/+}$ control, *Lyz2*$^{cre/cre}$*Ccr2*$^{RFP/RFP}$, *Rbpj*$^{fl/fl}$*Lyz2*$^{cre/cre}$*Ccr2*$^{RFP/+}$, and *Rbpj*$^{fl/fl}$*Lyz2*$^{cre/cre}$*Ccr2*$^{RFP/RFP}$ (DKO) mice. The deletion

265 of CCR2 led to a decrease in Ly6C^{hi} monocytes, and the Ly6C^{lo} monocytes in DKO mice were reduced to a level similar to that of control mice (Figure 7A, B). While the DKO mice had more CD16.2$^+$ IM compared to control mice, these cells were severely reduced compared to RBP-J deficient mice (Figure 7A, C). The above data supported the notion that increased lung Ly6C^{lo} monocytes and CD16.2$^+$ IM in RBP-J deficient

270 mice were derived from Ly6C^{lo} blood monocytes. In summary, these results suggested that in RBP-J deficient mice, recruitment of blood Ly6C^{lo} monocytes to the lung was markedly facilitated by increased cell number as well as heightened expression of CCR2, leading to the increase of lung Ly6C^{lo} monocytes and CD16.2$^+$ IM.

275 **Discussion**

In this study, we identified RBP-J as a pivotal factor in regulating blood Ly6C^{lo} monocyte cell fate. Our results showed that mice with conditional deletion of RBP-J in myeloid cells exhibited a robust increase in blood Ly6C^{lo} monocytes, and subsequently accumulated lung Ly6C^{lo} monocytes and CD16.2$^+$ IM under steady-state conditions.

280 Bone marrow transplantation experiments in which RBP-J deficient cells were transplanted into recipient mice as well as the parabiosis experiment showed similar increase in blood Ly6C^{lo} monocytes, demonstrating that RBP-J was an intrinsic factor required for the regulation of Ly6C^{lo} monocytes. Further analysis revealed that RBP-J regulated the expression of CCR2 and CD11c on the surface of Ly6C^{lo} monocytes.

285   Moreover, the phenotype of elevated blood Ly6C[lo] monocytes and their progeny was further ameliorated by deleting *Ccr2* in an RBP-J deficient background.

Notch-RBP-J signaling have been shown to play a role in regulating functional polarization and activation of macrophages, and regulated formation of Kupffer cells
290   and macrophage differentiation from Ly6C[hi] monocytes in ischemia[42-48]. At steady-state, interaction of DLL1 with Notch 2 regulates conversion of Ly6C[hi] monocytes into Ly6C[lo] monocytes in special niches of the BM and spleen[49]. Under inflammatory conditions, Notch2 and TLR7 pathways independently and synergistically promote conversion of Ly6C[hi] monocytes into Ly6C[lo] monocytes[50]. However, in our study, the
295   percentage of blood Ly6C[lo] monocytes increased in RBP-J deficient mice. These results somewhat differ from what was observed in mice with conditional deletion of Notch2[49]. Actually, 4 members of the Notch family have been identified in mammals[26]. Thus, Notch receptors seem to be non-redundant in regulating monocyte cell fate, and distinct Notch receptors are likely to act in a concerted manner to coordinate the monocyte
300   differentiation program.

Colony-stimulating factor 1 receptor (CSF1R) signal, CX3CR1, C/EBPβ and Nr4a1 have been suggested to be involved in the generation and survival of Ly6C[lo] monocytes. The lifespan of Ly6C[lo] monocytes is acutely shortened after blockade of CSF1R, and
305   the numbers of these cells are reduced, and the percentage of dead cells increased in mice with endothelial cell-specific depletion of Csf1[23,24]. CX3CR1 and C/EBPβ-knockout mice show accelerated death of Ly6C[lo] monocyte, display the decreased level of Ly6C[lo] monocytes[22,51]. Nr4a1 knockout mice had significantly fewer monocytes in blood, bone marrow, and spleen, due to differentiation deficiency in MDP and
310   accelerated apoptosis of Ly6C[lo] monocytes in bone marrow[21]. RBP-J deficient mice had normal MDP and Ly6C[lo] monocytes, expressed normal level of Nr4a1 and Ki-67, exhibited normal half-life and turnover rate, which suggested that RBP-J may not regulate Ly6C[lo] monocytes through these factors. Further studies are required to elucidate the precise molecular mechanisms of RBP-J in Ly6C[lo] monocytes under

315   steady state.

At the steady state, Ly6C[lo] monocytes patrol endothelium of blood vessel. After R848 induced endothelial injury, Ly6C[lo] monocytes recruit neutrophils to mediate focal endothelial necrosis and fuel vascular inflammation, and subsequently, the Ly6C[lo]

320   monocytes remove cellular debris[9,52]. In response to *Listeria monocytogenes* infection, Ly6C[lo] monocytes can extravasate but do not give rise to macrophages or DCs[8]. Additional evidence shows that Ly6C[lo] monocytes do not merely act as luminal blood macrophages. In lung, exposure to unmethylated CpG DNA expands CD16.2[+] IM, which originate from Ly6C[lo] monocytes and spontaneously produce IL-10, thereby

325   preventing allergic inflammation[41]. A recent study shows that Ly6C[lo] monocytes give rise to CD9[+] macrophages, which provide an intracellular replication niche for *Salmonella* Typhimurium in spleen, and Ly6C[lo]-depleted mice are more resistant to *Salmonella* Typhimurium infection, suggesting that Ly6C[lo] monocytes exert certain functions in systemic infection[53]. Our data provide evidence for RBP-J in the function

330   of blood Ly6C[lo] monocytes, as RBP-J deficient Ly6C[lo] monocytes exhibited enhanced competition in blood circulation, as well as gave rise to increased numbers of lung CD16.2[+] IM. In summary, we showed that RBP-J acted as a fundamental regulator of the maintenance of blood Ly6C[lo] monocytes and their descendent, at least in part through regulation of CCR2. These results provide insights into understanding the

335   mechanisms that regulate monocyte homeostasis and function.

References

1. Guilliams, M., Mildner, A., and Yona, S. (2018). Developmental and Functional Heterogeneity of Monocytes. Immunity *49*, 595-613. 10.1016/j.immuni.2018.10.005.

2. Geissmann, F., Jung, S., and Littman, D.R. (2003). Blood Monocytes Consist of Two Principal Subsets with Distinct Migratory Properties. Immunity *19*, 71-82. 10.1016/s1074-7613(03)00174-2.

3. Passlick, B., Flieger, D., and Ziegler-Heitbrock, H.W. (1989). Identification and characterization of a novel monocyte subpopulation in human peripheral blood. Blood *74*, 2527-2534.

4. Weber, C., Belge, K.U., von Hundelshausen, P., Draude, G., Steppich, B., Mack, M., Frankenberger, M., Weber, K.S., and Ziegler-Heitbrock, H.W. (2000). Differential chemokine receptor expression and function in human monocyte subpopulations. J Leukoc Biol *67*, 699-704. 10.1002/jlb.67.5.699.

5. Cros, J., Cagnard, N., Woollard, K., Patey, N., Zhang, S.Y., Senechal, B., Puel, A., Biswas, S.K., Moshous, D., Picard, C., et al. (2010). Human CD14dim monocytes patrol and sense nucleic acids and viruses via TLR7 and TLR8 receptors. Immunity *33*, 375-386. 10.1016/j.immuni.2010.08.012.

6. Ginhoux, F., and Jung, S. (2014). Monocytes and macrophages: developmental pathways and tissue homeostasis. Nat Rev Immunol *14*, 392-404. 10.1038/nri3671.

7. Ingersoll, M.A., Spanbroek, R., Lottaz, C., Gautier, E.L., Frankenberger, M., Hoffmann, R., Lang, R., Haniffa, M., Collin, M., Tacke, F., et al. (2010). Comparison of gene expression profiles between human and mouse monocyte subsets. Blood *115*, e10-19. 10.1182/blood-2009-07-235028.

8. Auffray, C., Fogg, D., Garfa, M., Elain, G., Join-Lambert, O., Kayal, S., Sarnacki, S., Cumano, A., Lauvau, G., and Geissmann, F. (2007). Monitoring of blood vessels and tissues by a population of monocytes with patrolling behavior. Science *317*, 666-670. 10.1126/science.1142883.

9. Carlin, L.M., Stamatiades, E.G., Auffray, C., Hanna, R.N., Glover, L., Vizcay-Barrena, G., Hedrick, C.C., Cook, H.T., Diebold, S., and Geissmann, F. (2013). Nr4a1-dependent Ly6C(low) monocytes monitor endothelial cells and orchestrate their disposal. Cell *153*, 362-375. 10.1016/j.cell.2013.03.010.

10. Yona, S., Kim, K.W., Wolf, Y., Mildner, A., Varol, D., Breker, M., Strauss-Ayali, D., Viukov, S., Guilliams, M., Misharin, A., et al. (2013). Fate mapping reveals origins and dynamics of monocytes and tissue macrophages under homeostasis. Immunity *38*, 79-91. 10.1016/j.immuni.2012.12.001.

11. Hanna, R.N., Shaked, I., Hubbeling, H.G., Punt, J.A., Wu, R., Herrley, E., Zaugg, C., Pei, H., Geissmann, F., Ley, K., and Hedrick, C.C. (2012). NR4A1 (Nur77) deletion polarizes macrophages toward an inflammatory phenotype and increases atherosclerosis. Circ Res *110*, 416-427. 10.1161/CIRCRESAHA.111.253377.

12. Hamers, A.A., Vos, M., Rassam, F., Marinkovic, G., Kurakula, K., van Gorp, P.J., de Winther, M.P., Gijbels, M.J., de Waard, V., and de Vries, C.J. (2012).

390   Bone marrow-specific deficiency of nuclear receptor Nur77 enhances atherosclerosis. Circ Res *110*, 428-438. 10.1161/CIRCRESAHA.111.260760.

13.   Urbanski, K., Ludew, D., Filip, G., Filip, M., Sagan, A., Szczepaniak, P., Grudzien, G., Sadowski, J., Jasiewicz-Honkisz, B., Sliwa, T., et al. (2017). CD14(+)CD16(++) "nonclassical" monocytes are associated with endothelial dysfunction in patients with coronary artery disease. Thromb Haemost *117*, 971-
395   980. 10.1160/TH16-08-0614.

14.   Amano, H., Amano, E., Santiago-Raber, M.L., Moll, T., Martinez-Soria, E., Fossati-Jimack, L., Iwamoto, M., Rozzo, S.J., Kotzin, B.L., and Izui, S. (2005). Selective expansion of a monocyte subset expressing the CD11c dendritic cell marker in the Yaa model of systemic lupus erythematosus. Arthritis Rheum *52*,
400   2790-2798. 10.1002/art.21365.

15.   Nahrendorf, M., Swirski, F.K., Aikawa, E., Stangenberg, L., Wurdinger, T., Figueiredo, J.L., Libby, P., Weissleder, R., and Pittet, M.J. (2007). The healing myocardium sequentially mobilizes two monocyte subsets with divergent and complementary functions. J Exp Med *204*, 3037-3047. 10.1084/jem.20070885.

405 16.   Misharin, A.V., Cuda, C.M., Saber, R., Turner, J.D., Gierut, A.K., Haines, G.K., 3rd, Berdnikovs, S., Filer, A., Clark, A.R., Buckley, C.D., et al. (2014). Nonclassical Ly6C(-) monocytes drive the development of inflammatory arthritis in mice. Cell Rep *9*, 591-604. 10.1016/j.celrep.2014.09.032.

17.   Puchner, A., Saferding, V., Bonelli, M., Mikami, Y., Hofmann, M., Brunner, J.S.,
410   Caldera, M., Goncalves-Alves, E., Binder, N.B., Fischer, A., et al. (2018). Non-classical monocytes as mediators of tissue destruction in arthritis. Ann Rheum Dis *77*, 1490-1497. 10.1136/annrheumdis-2018-213250.

18.   Brunet, A., LeBel, M., Egarnes, B., Paquet-Bouchard, C., Lessard, A.J., Brown, J.P., and Gosselin, J. (2016). NR4A1-dependent Ly6C(low) monocytes
415   contribute to reducing joint inflammation in arthritic mice through Treg cells. Eur J Immunol *46*, 2789-2800. 10.1002/eji.201646406.

19.   Hanna, R.N., Cekic, C., Sag, D., Tacke, R., Thomas, G.D., Nowyhed, H., Herrley, E., Rasquinha, N., McArdle, S., Wu, R., et al. (2015). Patrolling monocytes control tumor metastasis to the lung. Science *350*, 985-990.
420   10.1126/science.aac9407.

20.   Thomas, G.D., Hanna, R.N., Vasudevan, N.T., Hamers, A.A., Romanoski, C.E., McArdle, S., Ross, K.D., Blatchley, A., Yoakum, D., Hamilton, B.A., et al. (2016). Deleting an Nr4a1 Super-Enhancer Subdomain Ablates Ly6C(low) Monocytes while Preserving Macrophage Gene Function. Immunity *45*, 975-
425   987. 10.1016/j.immuni.2016.10.011.

21.   Hanna, R.N., Carlin, L.M., Hubbeling, H.G., Nackiewicz, D., Green, A.M., Punt, J.A., Geissmann, F., and Hedrick, C.C. (2011). The transcription factor NR4A1 (Nur77) controls bone marrow differentiation and the survival of Ly6C-monocytes. Nat Immunol *12*, 778-785. 10.1038/ni.2063.

430 22.   Tamura, A., Hirai, H., Yokota, A., Kamio, N., Sato, A., Shoji, T., Kashiwagi, T., Torikoshi, Y., Miura, Y., Tenen, D.G., and Maekawa, T. (2017). C/EBPbeta is required for survival of Ly6C(-) monocytes. Blood *130*, 1809-1818.

10.1182/blood-2017-03-772962.

23. MacDonald, K.P., Palmer, J.S., Cronau, S., Seppanen, E., Olver, S., Raffelt, N.C., Kuns, R., Pettit, A.R., Clouston, A., Wainwright, B., et al. (2010). An antibody against the colony-stimulating factor 1 receptor depletes the resident subset of monocytes and tissue- and tumor-associated macrophages but does not inhibit inflammation. Blood *116*, 3955-3963. 10.1182/blood-2010-02-266296.

24. Emoto, T., Lu, J., Sivasubramaniyam, T., Maan, H., Khan, A.B., Abow, A.A., Schroer, S.A., Hyduk, S.J., Althagafi, M.G., McKee, T.D., et al. (2022). Colony stimulating factor-1 producing endothelial cells and mesenchymal stromal cells maintain monocytes within a perivascular bone marrow niche. Immunity *55*, 862-878 e868. 10.1016/j.immuni.2022.04.005.

25. Keerthivasan, S., Senbabaoglu, Y., Martinez-Martin, N., Husain, B., Verschueren, E., Wong, A., Yang, Y.A., Sun, Y., Pham, V., Hinkle, T., et al. (2021). Homeostatic functions of monocytes and interstitial lung macrophages are regulated via collagen domain-binding receptor LAIR1. Immunity *54*, 1511-1526 e1518. 10.1016/j.immuni.2021.06.012.

26. Radtke, F., MacDonald, H.R., and Tacchini-Cottier, F. (2013). Regulation of innate and adaptive immunity by Notch. Nat Rev Immunol *13*, 427-437. 10.1038/nri3445.

27. Klinakis, A., Lobry, C., Abdel-Wahab, O., Oh, P., Haeno, H., Buonamici, S., van De Walle, I., Cathelin, S., Trimarchi, T., Araldi, E., et al. (2011). A novel tumour-suppressor function for the Notch pathway in myeloid leukaemia. Nature *473*, 230-233. 10.1038/nature09999.

28. Caton, M.L., Smith-Raska, M.R., and Reizis, B. (2007). Notch-RBP-J signaling controls the homeostasis of CD8- dendritic cells in the spleen. J Exp Med *204*, 1653-1664. 10.1084/jem.20062648.

29. Lewis, K.L., Caton, M.L., Bogunovic, M., Greter, M., Grajkowska, L.T., Ng, D., Klinakis, A., Charo, I.F., Jung, S., Gommerman, J.L., et al. (2011). Notch2 receptor signaling controls functional differentiation of dendritic cells in the spleen and intestine. Immunity *35*, 780-791. 10.1016/j.immuni.2011.08.013.

30. Zhao, B., Grimes, S.N., Li, S., Hu, X., and Ivashkiv, L.B. (2012). TNF-induced osteoclastogenesis and inflammatory bone resorption are inhibited by transcription factor RBP-J. J Exp Med *209*, 319-334. 10.1084/jem.20111566.

31. Varol, C., Landsman, L., Fogg, D.K., Greenshtein, L., Gildor, B., Margalit, R., Kalchenko, V., Geissmann, F., and Jung, S. (2007). Monocytes give rise to mucosal, but not splenic, conventional dendritic cells. J Exp Med *204*, 171-180. 10.1084/jem.20061011.

32. Hettinger, J., Richards, D.M., Hansson, J., Barra, M.M., Joschko, A.C., Krijgsveld, J., and Feuerer, M. (2013). Origin of monocytes and macrophages in a committed progenitor. Nat Immunol *14*, 821-830. 10.1038/ni.2638.

33. Liu, Z., Gu, Y., Chakarov, S., Bleriot, C., Kwok, I., Chen, X., Shin, A., Huang, W., Dress, R.J., Dutertre, C.A., et al. (2019). Fate Mapping via Ms4a3-Expression History Traces Monocyte-Derived Cells. Cell *178*, 1509-1525 e1519. 10.1016/j.cell.2019.08.009.

34. Debien, E., Mayol, K., Biajoux, V., Daussy, C., De Aguero, M.G., Taillardet, M., Dagany, N., Brinza, L., Henry, T., Dubois, B., et al. (2013). S1PR5 is pivotal for the homeostasis of patrolling monocytes. Eur J Immunol *43*, 1667-1675. 10.1002/eji.201343312.

35. Tacke, F., Ginhoux, F., Jakubzick, C., van Rooijen, N., Merad, M., and Randolph, G.J. (2006). Immature monocytes acquire antigens from other cells in the bone marrow and present them to T cells after maturing in the periphery. J Exp Med *203*, 583-597. 10.1084/jem.20052119.

36. Liu, K., Waskow, C., Liu, X., Yao, K., Hoh, J., and Nussenzweig, M. (2007). Origin of dendritic cells in peripheral lymphoid organs of mice. Nat Immunol *8*, 578-583. 10.1038/ni1462.

37. Shi, C., and Pamer, E.G. (2011). Monocyte recruitment during infection and inflammation. Nat Rev Immunol *11*, 762-774. 10.1038/nri3070.

38. Guilliams, M., De Kleer, I., Henri, S., Post, S., Vanhoutte, L., De Prijck, S., Deswarte, K., Malissen, B., Hammad, H., and Lambrecht, B.N. (2013). Alveolar macrophages develop from fetal monocytes that differentiate into long-lived cells in the first week of life via GM-CSF. J Exp Med *210*, 1977-1992. 10.1084/jem.20131199.

39. Hashimoto, D., Chow, A., Noizat, C., Teo, P., Beasley, M.B., Leboeuf, M., Becker, C.D., See, P., Price, J., Lucas, D., et al. (2013). Tissue-resident macrophages self-maintain locally throughout adult life with minimal contribution from circulating monocytes. Immunity *38*, 792-804. 10.1016/j.immuni.2013.04.004.

40. Sabatel, C., Radermecker, C., Fievez, L., Paulissen, G., Chakarov, S., Fernandes, C., Olivier, S., Toussaint, M., Pirottin, D., Xiao, X., et al. (2017). Exposure to Bacterial CpG DNA Protects from Airway Allergic Inflammation by Expanding Regulatory Lung Interstitial Macrophages. Immunity *46*, 457-473. 10.1016/j.immuni.2017.02.016.

41. Schyns, J., Bai, Q., Ruscitti, C., Radermecker, C., De Schepper, S., Chakarov, S., Farnir, F., Pirottin, D., Ginhoux, F., Boeckxstaens, G., et al. (2019). Non-classical tissue monocytes and two functionally distinct populations of interstitial macrophages populate the mouse lung. Nat Commun *10*, 3964. 10.1038/s41467-019-11843-0.

42. Wang, Y.C., He, F., Feng, F., Liu, X.W., Dong, G.Y., Qin, H.Y., Hu, X.B., Zheng, M.H., Liang, L., Feng, L., et al. (2010). Notch signaling determines the M1 versus M2 polarization of macrophages in antitumor immune responses. Cancer Res *70*, 4840-4849. 10.1158/0008-5472.CAN-10-0269.

43. Kang, L., Zhang, X., Ji, L., Kou, T., Smith, S.M., Zhao, B., Guo, X., Pineda-Torra, I., Wu, L., and Hu, X. (2020). The colonic macrophage transcription factor RBP-J orchestrates intestinal immunity against bacterial pathogens. J Exp Med *217*. 10.1084/jem.20190762.

44. Hu, X., Chung, A.Y., Wu, I., Foldi, J., Chen, J., Ji, J.D., Tateya, T., Kang, Y.J., Han, J., Gessler, M., et al. (2008). Integrated regulation of Toll-like receptor responses by Notch and interferon-gamma pathways. Immunity *29*, 691-703.

10.1016/j.immuni.2008.08.016.

45. Foldi, J., Shang, Y., Zhao, B., Ivashkiv, L.B., and Hu, X. (2016). RBP-J is required for M2 macrophage polarization in response to chitin and mediates expression of a subset of M2 genes. Protein Cell 7, 201-209. 10.1007/s13238-016-0248-7.

46. Xu, H., Zhu, J., Smith, S., Foldi, J., Zhao, B., Chung, A.Y., Outtz, H., Kitajewski, J., Shi, C., Weber, S., et al. (2012). Notch-RBP-J signaling regulates the transcription factor IRF8 to promote inflammatory macrophage polarization. Nat Immunol 13, 642-650. 10.1038/ni.2304.

47. Sakai, M., Troutman, T.D., Seidman, J.S., Ouyang, Z., Spann, N.J., Abe, Y., Ego, K.M., Bruni, C.M., Deng, Z., Schlachetzki, J.C.M., et al. (2019). Liver-Derived Signals Sequentially Reprogram Myeloid Enhancers to Initiate and Maintain Kupffer Cell Identity. Immunity 51, 655-670 e658. 10.1016/j.immuni.2019.09.002.

48. Krishnasamy, K., Limbourg, A., Kapanadze, T., Gamrekelashvili, J., Beger, C., Hager, C., Lozanovski, V.J., Falk, C.S., Napp, L.C., Bauersachs, J., et al. (2017). Blood vessel control of macrophage maturation promotes arteriogenesis in ischemia. Nat Commun 8, 952. 10.1038/s41467-017-00953-2.

49. Gamrekelashvili, J., Giagnorio, R., Jussofie, J., Soehnlein, O., Duchene, J., Briseno, C.G., Ramasamy, S.K., Krishnasamy, K., Limbourg, A., Kapanadze, T., et al. (2016). Regulation of monocyte cell fate by blood vessels mediated by Notch signalling. Nat Commun 7, 12597. 10.1038/ncomms12597.

50. Gamrekelashvili, J., Kapanadze, T., Sablotny, S., Ratiu, C., Dastagir, K., Lochner, M., Karbach, S., Wenzel, P., Sitnow, A., Fleig, S., et al. (2020). Notch and TLR signaling coordinate monocyte cell fate and inflammation. Elife 9. 10.7554/eLife.57007.

51. Landsman, L., Bar-On, L., Zernecke, A., Kim, K.W., Krauthgamer, R., Shagdarsuren, E., Lira, S.A., Weissman, I.L., Weber, C., and Jung, S. (2009). CX3CR1 is required for monocyte homeostasis and atherogenesis by promoting cell survival. Blood 113, 963-972. 10.1182/blood-2008-07-170787.

52. Imhof, B.A., Jemelin, S., Ballet, R., Vesin, C., Schapira, M., Karaca, M., and Emre, Y. (2016). CCN1/CYR61-mediated meticulous patrolling by Ly6Clow monocytes fuels vascular inflammation. Proc Natl Acad Sci U S A 113, E4847-4856. 10.1073/pnas.1607710113.

53. Hoffman, D., Tevet, Y., Trzebanski, S., Rosenberg, G., Vainman, L., Solomon, A., Hen-Avivi, S., Ben-Moshe, N.B., and Avraham, R. (2021). A non-classical monocyte-derived macrophage subset provides a splenic replication niche for intracellular Salmonella. Immunity 54, 2712-2723 e2716. 10.1016/j.immuni.2021.10.015.

Acknowledgements:

We thank the core facility at Institute for Immunology, Tsinghua University for valuable technical assistance.

Funding:

This research was supported by National Natural Science Foundation of China grants (31821003, 31991174, 32030037 and 82150105 to X. Hu), a Ministry of Science and Technology of China grant (2020YFA0509100 to X. Hu), and funds from Tsinghua-Peking Center for Life Sciences and Institute for Immunology at Tsinghua University (to X. Hu).

Author contributions:

T. Kou designed and performed experiments, analyzed and interpreted data, and wrote the manuscript. L. Kang designed and performed experiments, analyzed and interpreted data. B. Zhang provided advice on RNA-seq analyses. J. Li performed experiments. B. Zhao provided advice on experiments. W. Zeng provided advice on experiments. X. Hu conceptualized the project, supervised experiments, interpreted data, and wrote the manuscript.

Declaration of interests:

The authors declare no competing interests.

Data availability:

Sequencing data sets are deposited in the Genome Expression Omnibus with assigned accession numbers as follows: RNA-seq (GEO accession no. GSE208772).

Materials and methods
Mice

Cx3cr1gfp/gfp mice (JAX stock 005582; Jung et al., 2000) and *Ccr2RFP/RFP* mice (JAX stock 017586) were purchased from the Jackson Laboratory. Mice with a myeloid-

specific deletion of the *Rbpj* were generated by crossing *Rbpj^{fl/fl}* mice to *Lyz2*-Cre mice as described previously (Hu et al., 2008). *Rbpj^{fl/fl}Lyz2^{cre/cre}* were crossed to *Cx3cr1^{gfp/gfp}* mice to obtain *Lyz2^{cre/cre}Cx3cr1^{gfp/+}* and *Rbpj^{fl/fl}Lyz2^{cre/cre}Cx3cr1^{gfp/+}* mice. *Cx3cr1^{gfp/+}* mice were obtained by crossing *Cx3cr1^{gfp/gfp}* with C57/BL6 CD45.1^+ mice. *Rbpj^{fl/fl}Lyz2^{cre/cre}* mice were crossed with *Ccr2^{RFP/RFP}* mice to obtain *Lyz2^{cre/cre}Ccr2^{RFP/+}*, *Lyz2^{cre/cre}Ccr2^{RFP/RFP}*, *Rbpj^{fl/fl}Lyz2^{cre/cre}Ccr2^{RFP/+}* and *Rbpj^{fl/fl}Lyz2^{cre/cre}Ccr2^{RFP/RFP}* mice. All mice were maintained under specific pathogen-free conditions. All animal experimental protocols were approved by the Institutional Animal Care and Use Committees of Tsinghua University. Gender- and age-matched mice were used at 7–12 weeks old for experiments.

Quantitative RT-PCR

Blood monocytes were sorted by flow cytometry. The total RNA was extracted using total RNA purification kit (GeneMarkbio) and reversely transcribed to cDNA by M-MLV Reverse Transcriptase (Takara). qPCR was performed on a real-time PCR system (StepOnePlus; Applied Biosystems) using FastSYBR mixture (CWBIO). *Gapdh* messenger RNA was used as internal control to normalize the expression of target genes. Primer sequences are provided in supplemental Table 1.

Table 1: Primers sequences for regular qPCR used in this study

Gene	Forward primer	Reverse primer
Gapdh	ATCAAGAAGGTGGTGAAGCA	AGACAACCTGGTCCTCAGTGT
Rbpj	ACCCCTGTGCCTGTCGTAGAA	TCCCGGAATGCAGAAATGTC
Nr4a1	TTGAGCTTGAATACAGGGCA	AGTTGGGGGAGTGTGCTAGA
Ccr2	CCTTGGGAATGAGTAACTGTGTGAT	ATGGAGAGATACCTTCGGAACTTCT

RNA-seq Analysis

Ly6C^{hi} and Ly6C^{lo} blood monocytes were isolated from *Rbpj^{+/+}Lyz2^{cre/cre}* and *Rbpj^{fl/fl}Lyz2^{cre/cre}* mice, and total RNA was extracted using total RNA purification kit (GeneMarkbio). RNA was converted into RNA-seq libraries, which were sequenced

with the pair-end option using an Illumina-HiSeq2500 platform at Beijing Genomics Institute (BGI), China. The significantly down-regulated genes were identified with P value <0.05 and (FPKM + 1) fold changes ≤0.2, and significantly up-regulated genes were identified with P value <0.05 and (FPKM + 1) fold changes ≥5.7. The RNA-seq data are deposited in Gene Expression Omnibus under accession number GSE208772

Annexin V staining

Annexin V and 7-amino-actinomycin D (7-AAD) were used for identification of apoptotic monocytes by flow cytometry. Blood cells were stained with annexin V (eBioscience) and 7-AAD (Biolegend) according to the manufacturers' protocols.

EdU pulsing and latex beads labeling

Mice were injected intravenously with a single 1mg EdU (Thermo Scientific). BM and blood cells were collected and stained with Fluorescence-conjugated mAb against CD45, CD11b, Ly6G, CD115 and Ly6C. Cells were then fixed, permeabilized and stained with reaction cocktail using EdU Assay Kit. Labeled cells were analyzed by flow cytometry.

Mice were injected intravenously with a single 10 µL Latex beads (Polysciences) in 250 µl PBS. Blood cells were harvested at indicated time. Cells were then stained for CD45, CD11b, Ly6G, CD115, Ly6C, and analyzed by flow cytometry.

In vivo labeling of sinusoidal leukocytes

Mice were injected intravenously with 1 µg of PE-conjugated anti-CD45 antibodies. Two minutes after antibody injection, mice were sacrificed, and bone marrow were harvested. BM cells were then stained and analyzed by flow cytometry.

Generation of BM chimera

C57/BL6 CD45.2$^+$ mice were lethally irradiated in two doses of 5.5Gy 2 h apart. 0.8 × 10^5 BM cells from $Rbpj^{+/+}Lyz2^{cre/cre}$ mice (CD45.2$^+$) or $Rbpj^{fl/fl}Lyz2^{cre/cre}$ mice (CD45.2$^+$) were mixed with 3.2× 10^5 BM cells from $Cx3cr1^{gfp/+}$ mice (CD45.1$^+$), and injected

intravenously into recipient mice. Mice were used for experiments 8 weeks after irradiation.

Adoptive transfers

BM GFP$^+$Ly6C^{hi} monocytes were sorted from *Lyz2*$^{cre/cre}$*Cx3cr1*$^{gfp/+}$ or *Rbpj*$^{fl/fl}$*Lyz2*$^{cre/cre}$*Cx3cr1*$^{gfp/+}$ mice, and transferred to *Ccr2*$^{RFP/RFP}$ mice. Blood cells were collected 60 hours later for flow cytometry analyses.

Parabiosis

Cx3cr1$^{gfp/+}$ CD45.1$^+$ mice were surgically joined with age-matched female *Rbpj*$^{fl/fl}$*Lyz2*$^{cre/cre}$ or *Rbpj*$^{+/+}$*Lyz2*$^{cre/cre}$ CD45.2$^+$ mice at the age of 6 weeks. Bloods were obtained via cardiac puncture, and cell populations were analyzed by flow cytometry at

660 4 weeks after the surgery.

Intranasal instillations of LPS

Rbpj$^{+/+}$*Lyz2*$^{cre/cre}$ or *Rbpj*$^{fl/fl}$*Lyz2*$^{cre/cre}$ mice were anesthetized with isoflurane and intranasally instilled with 10 μg LPS in 25 μl of PBS. Lungs were harvested 4 days later

665 for flow cytometry analyses.

Immunofluorescence histology

Lungs from *Rbpj*$^{fl/fl}$*Lyz2*$^{cre/cre}$*Cx3cr1*$^{gfp/+}$ and *Lyz2*$^{cre/cre}$*Cx3cr1*$^{gfp/+}$ mice were fixed in 1% paraformaldehyde and incubated in 30% sucrose separately overnight at 4°C. The

670 samples were then incubated in the mixture of 30% sucrose and OTC compound (Sakura Finetek) overnight at 4°C. The tissues were embedded and frozen in OCT compound and then cut at 10-μm thickness. Tissue sections were dried for 10 min at 50°C and then fixed in 1% paraformaldehyde at room temperature for 10 min, permeabilized in PBS/0.5% Triton X-100/0.3 M glycine for 10 min and blocked in

675 PBS/5% goat serum for 1 h at room temperature. Sections were stained with rabbit anti-mouse GFP antibodies (1:1000; EASYBIO) overnight at 4°C, and washed with PBS/0.1% Tween-20 for 30 min three times at room temperature. Sections were then incubated

with AF488-conjugated goat anti-rabbit antibodies (1:200; proteintech) for 2 h at room temperature, and washed with PBS/0.1% Tween-20 for 30 min three times at room temperature. Sections were stained with DAPI (Solarbio) for 7 min, washed in PBS for 8 min two times at room temperature and mounted with SlowFade Diamond Antifade Mountant (Life Technologies).

Cell isolation and Flow cytometry

Peripheral blood was sampled by eyeball extirpating, spleens were mashed through a 70-μm strainer, and bone marrow cells were collected from femurs. Lungs were perfused with 5ml of HBSS (MACGENE) through the right ventricle, excised, and digested in HBSS containing 5% FBS, 1 mg/ml collagenase type I (Sigma-Aldrich) and 0.05 mg/ml DNase I (Sigma-Aldrich) for 1 h at 37 °C. The digested tissues were homogenized by shaking, passed through a 70-μm cell strainer to create a single cell suspension. The suspension was enriched in mononuclear cells and harvested from the 1.080:1.038 g ml-1 interface using a density gradient (Percoll from GE Healthcare). BM, Spleen, PB and lung cells were stained with fluorescently conjugated antibodies. The absolute number of cells was counted by using CountBright™ Absolute Counting Beads (Invitrogen).

Antibodies against CD45 (30-F11), Ly6C (HK1.4), CD64 (X54-5/7.1), CD206 (C068C2), CD16.2 (9E9), CD45.1 (A20), F4/80 (BM8), Biotin, CCR2 (SA203G11) and CD3ε (145-2C11) were purchased from BioLegend. Antibodies against Ly6G (1A8-Ly6g), CD11b (M1/70), CD115(AFS98), CD117(2B8), CD135 (A2F10), CD11c (N418), CD206 (MR6F3), CD45.2 (104), Nur77 (12.14), Ki-67 (SolA15) and lineage maker were purchased from eBioscience. Fluorescence-conjugated mAb against Ly6C (AL-21) were purchased from BD Biosciences. Isotype-matched antibodies (eBioscience) were used for control staining. All antibodies were used in 1:400 dilutions, and surface antigens were stained on ice for 30 min.

For intracellular staining, cells were stained with antibodies to surface antigens, fixed

and permeabilized with the Cytofix/Cytoperm Fixation/Permeabilization Solution Kit (BD Biosciences), and stained with antibodies or isotype control diluted in permeabilization buffer separately for 30 min at room temperature.

Cells were analyzed on FACSFortessa or FACSAria III flow cytometer (BD Biosciences) using FlowJo software.

Figure Legends:

Figure 1. RBP-J deficient mice display more Ly6C^{lo} blood monocytes.

(A) Blood Ly6C^{hi} and Ly6C^{lo} monocytes in *Rbpj$^{+/+}$Lyz2*$^{cre/cre}$ control and *Rbpj$^{fl/fl}$Lyz2*$^{cre/cre}$ mice were determined by flow cytometry analyses (FACS). Representative FACS plots (left) and cumulative data of cell ratio (right) are shown.

(B and C) Representative FACS plots and cumulative data quantitating percentages of BM monocyte subsets (B) and myeloid progenitor cells (C) (CD45$^+$CD11b$^+$Ly6G$^-$CD115$^+$ Ly6C^{hi} monocyte; CD45$^+$CD11b$^+$Ly6G$^-$CD115$^+$Ly6C^{lo} monocyte; MDP, CD45$^+$Lin$^-$CD117$^+$CD115$^+$CD135$^+$ Ly6C$^-$; cMoP, CD45$^+$Lin$^-$CD11b$^-$CD117$^+$CD115$^+$CD135$^-$Ly6C$^+$). Lin: CD3, B220, Ter119, and Gr-1, CD11b.

(D) Experimental outline for panel (E).

(E) Cumulative data quantitating percentages of sinusoidal monocytes (CD45$^+$) within total BM Ly6C^{lo} monocytes.

Data are pooled from at least 2 independent experiments; n ≥ 4 in each group. Data are shown as mean ± SEM; n.s., not significant; **, P < 0.01 (two-tailed Student's unpaired t test). Each symbol represents an individual mouse.

Figure 2. Monocyte subsets in RBP-J deficient mice display normal cell death.

(A) Representative FACS plots of monocyte subsets in blood stained with 7-AAD and Annexin V.

(B) FACS analysis of Ki-67 expression in *Rbpj$^{+/+}$Lyz2*$^{cre/cre}$ control and *Rbpj$^{fl/fl}$Lyz2*$^{cre/cre}$ blood monocyte subsets. Black lines represent control mice and red lines represent RBP-J deficient mice.

(C) Analysis of time course of EdU incorporation of monocyte subsets in BM and blood after a single 1mg EdU pulsing. The percentages of EdU$^+$ cells among the indicated monocyte subsets are shown.

(D) Experimental outline for panel (E).

(E) Analysis of time course of latex beads incorporation of monocyte subsets in blood after Latex beads injection. The percentages of latex$^+$ cells among the indicated monocyte subsets are shown.

(F) Cartoon depicting the adoptive transfer. BM GFP$^+$Ly6C^{hi} monocytes were sorted from *Lyz2*$^{cre/cre}$*Cx3cr1*$^{gfp/+}$ or *Rbpj*$^{fl/fl}$*Lyz2*$^{cre/cre}$*Cx3cr1*$^{gfp/+}$ mice and transferred into *Ccr2*$^{RFP/RFP}$ recipient mice. Sixty hours after transfer, cell fate was analyzed.

(G) Representative FACS plots are shown in the left panel, and the frequencies of GFP$^+$ Ly6C^{lo} monocytes within total Ly6C^{lo} monocytes are shown in the right panel.

Data are pooled from 2 independent experiments (G); n ≥ 2 in each group (C, E and G). Data are shown as mean ± SEM; n.s., not significant; (two-tailed Student's unpaired t test). Each symbol represents an individual mouse (E and G).

Figure 3. The role of RBP-J in blood Ly6C^{lo} monocytes is cell intrinsic.

(A) Cartoon depicting the bone marrow transplantation. BM recipient 6-week-old male C57BL/6 mice (CD45.2) were lethally irradiated. BM cells from donor mice (*Rbpj*$^{+/+}$*Lyz2*$^{cre/cre}$ or *Rbpj*$^{fl/fl}$*Lyz2*$^{cre/cre}$ and CD45.1, *Cx3cr1*$^{gfp/+}$) were collected and transferred into recipient mice. Mice were used after 8 weeks of bone marrow reconstitution.

(B) Representative FACS plots (left) and cumulative data (right) quantitating the frequency of *Rbpj*$^{+/+}$*Lyz2*$^{cre/cre}$ and *Rbpj*$^{fl/fl}$*Lyz2*$^{cre/cre}$ donor cells among Ly6C^{hi} and Ly6C^{lo} monocytes in the blood of recipient mice.

(C) Cartoon depicting the generation of *Rbpj*$^{+/+}$*Lyz2*$^{cre/cre}$ control or *Rbpj*$^{fl/fl}$*Lyz2*$^{cre/cre}$ and *Cx3cr1*$^{gfp/+}$ parabiotic pairs.

(D) Representative FACS plots (left) and cumulative data (right) quantitating percentages of monocyte subsets derived from control or RBP-J deficient mice in control or RBP-J deficient mice.

(E) Representative FACS plots (left) and cumulative data (right) quantitating percentages of monocyte subsets derived from control or RBP-J deficient mice in *Cx3cr1*$^{gfp/+}$ mice.

Data are pooled from at least 2 independent experiments; n ≥ 6 in each group. Data are shown as mean ± SEM; n.s., not significant; ***, P < 0.001; ****, P < 0.0001 (two-tailed Student's unpaired t test). Each symbol represents an individual mouse. SSC-A, side scatter area.

Figure 4. Phenotypic markers in Ly6C^{lo} monocytes are changed in RBP-J deficient mice.

(A) Heatmap of RNA-seq dataset showing the top 20 downregulated and upregulated genes in blood Ly6C^{lo} monocytes from *Rbpj$^{fl/fl}$Lyz2*$^{cre/cre}$ versus *Rbpj$^{+/+}$Lyz2*$^{cre/cre}$ control mice. Blue and red font indicates downregulated and upregulated genes in RBP-J deficient Ly6C^{lo} monocytes respectively.

(B) Representative FACS plots, cumulative MFI, and qPCR analysis of CCR2/CD11c expression in control and RBP-J deficient blood monocyte subsets. Shaded curves represent isotype control, black lines represent control mice and red lines represent RBP-J deficient mice.

(C) PCA of indicated cell types.

(D) Representative FACS plots and cumulative MFI of CCR2/CD11c expression in blood monocyte subsets derived from control or RBP-J deficient mice. Black lines represent control mice, and red lines represent RBP-J deficient mice.

Data are pooled from 2 independent experiments (B and D); n ≥ 6 in each group. Data are shown as mean ± SEM; n.s., not significant; **, P < 0.01; ****, P < 0.0001 (two-tailed Student's unpaired t test). Each symbol represents an individual mouse.

Figure 5. Blood Ly6C^{lo} monocytes are decreased in DKO mice.

(A) Representative FACS plots and cumulative MFI of CCR2 expression in *Lyz2$^{cre/cre}$Ccr2$^{RFP/+}$* control, *Rbpj$^{fl/fl}$Lyz2$^{cre/cre}$Ccr2$^{RFP/+}$* and *Rbpj$^{fl/fl}$Lyz2$^{cre/cre}$Ccr2$^{RFP/RFP}$* (DKO) blood Ly6C^{hi} and Ly6C^{lo} monocytes are shown.

(B and C) Blood monocyte subsets in control, RBP-J deficient and DKO mice were determined by FACS. Representative FACS plots (B) and cumulative data of cell ratio (C) are shown.

(D) Representative FACS plots and cumulative MFI of CD11c expression in control, RBP-J deficient and DKO blood Ly6C^{hi} and Ly6C^{lo} monocytes are shown.

Data are pooled from at least 2 independent experiments; n ≥ 5 in each group. Data are shown as mean ± SEM; n.s., not significant; **, P < 0.01; ***, P < 0.001; ****, P < 0.0001 (two-tailed Student's unpaired t test). Each symbol represents an individual

mouse.

Figure 6. RBP-J deficient mice exhibit more lung Ly6C^{lo} monocytes and CD16.2^{+} IM.

(A and B) Indicated populations in the lungs of *Rbpj$^{+/+}$Lyz2$^{cre/cre}$* and *Rbpj$^{fl/fl}$Lyz2$^{cre/cre}$* mice were determined by FACS. Representative FACS plots (A) and cumulative data of cell ratio and absolute numbers (B) are shown.

(C) Immunofluorescence staining for GFP^{+} cells in the lung from *Lyz2$^{cre/cre}$Cx3cr1$^{gfp/+}$* and *Rbpj$^{fl/fl}$Lyz2$^{cre/cre}$Cx3cr1$^{gfp/+}$* mice (CX3CR1 [green]; DAPI [blue]). Scale bars represent 50 µm.

(D and E) *Rbpj$^{+/+}$Lyz2$^{cre/cre}$* and *Rbpj$^{fl/fl}$Lyz2$^{cre/cre}$* mice were instilled intranasally with PBS or PBS containing LPS, and lungs were harvested at the indicated time points. Representative FACS plots (D) and cumulative data of cell ratio and absolute numbers (E) are shown.

Data are pooled from at least 2 independent experiments; n ≥ 4 in each group. Data are shown as mean ± SEM; n.s., not significant; *, P < 0.05; **, P < 0.01; ****, P < 0.0001 (two-tailed Student's unpaired t test). Each symbol represents an individual mouse.

Figure 7. DKO mice lack lung Ly6C^{lo} monocytes and CD16.2^{+} IM.

(A) Representative FACS plots of lung monocyte and IM subsets in *Lyz2$^{cre/cre}$Ccr2$^{RFP/+}$*, *Rbpj$^{fl/fl}$Lyz2$^{cre/cre}$Ccr2$^{RFP/+}$*, *Lyz2$^{cre/cre}$Ccr2$^{RFP/RFP}$* and *Rbpj$^{fl/fl}$Lyz2$^{cre/cre}$Ccr2$^{RFP/RFP}$* mice.

(B and C) Cumulative data of cell ratio and absolute numbers of monocyte (B) and IM (C) subsets.

Data are pooled from 2 independent experiments; n ≥ 4 in each group. Data are shown as mean ± SEM; n.s., not significant; *, P < 0.05; **, P < 0.01; ***, P < 0.001; ****, P < 0.0001 (two-tailed Student's unpaired t test). Each symbol represents an individual mouse.

Figure 1 - Figure supplement 1. Control and RBP-J deficient mice show similar neutrophils.

(A) qPCR analysis of *Rbpj* expression in sorted monocyte subsets from control and RBP-J deficient mice.

(B) Blood and BM CD45$^+$CD11b$^+$Ly6G$^+$ neutrophils were determined by FACS. Cumulative data of cell ratio are shown.

(C) Spleen monocyte subsets were determined by FACS. Representative FACS plots (left) and cumulative data of cell ratio and absolute numbers (right) are shown.

Data are pooled from at least 2 independent experiments; n ≥ 2 in each group. Data are shown as mean ± SEM; n.s., not significant; *, P < 0.05 (two-tailed Student's unpaired t test). Each symbol represents an individual mouse.

Figure 2 - Figure supplement 1. The conversion of Ly6C^{hi} monocytes is identical in control and RBP-J deficient mice.

(A) Representative FACS plots, cumulative MFI and qPCR analysis of Nr4a1 expression in control and RBP-J deficient blood monocyte subsets. Shaded curves represent isotype control, black lines represent control mice, and red lines represent RBP-J deficient mice.

(B) Representative FACS plots of GFP$^+$ Ly6C^{hi} monocytes in recipient.

Data are pooled from 2 independent experiments (A); n ≥ 4 in each group. Data are shown as mean ± SEM; n.s., not significant (two-tailed Student's unpaired t test). Each symbol represents an individual mouse.

Figure 3 - Figure supplement 1. Cell-intrinsic requirement of RBP-J for Ly6C^{lo} maintenance.

(A) Representative FACS plots (left) and cumulative data (right) quantitating percentages of blood neutrophils in recipient mice.

(B) BM neutrophils and Ly6C^{hi} monocytes in recipient mice were determined by FACS.

(C) CD4$^+$ T cells and CD19$^+$ B cells were analyzed by FACS in parabiotic mice.

Data are pooled from at least 2 independent experiments (A and C); n ≥ 4 in each group. Data are shown as mean ± SEM; n.s., not significant (two-tailed Student's unpaired t test). Each symbol represents an individual mouse.

Figure 4 - Figure supplement 1. Normal expression of CCR2 and CD11c in BM monocytes.

890   (A) Cumulative MFI of CCR2 expression in BM monocyte subsets.

(B) FACS plots of CCR2 and CD11c in BM Ly6C^{hi} and Ly6C^{lo} from control and RBP-J deficient mice.

Data are pooled from 3 independent experiments (A); n ≥ 7 in each group. Data are shown as mean ± SEM; n.s., not significant (two-tailed Student's unpaired t test). Each

895   symbol represents an individual mouse.

Figure 6 - Figure supplement 1. RBP-J is not required for turnover of lung Ly6C^{lo} monocytes and CD16.2$^+$ IM.

(A and B) Incorporation of EdU was assessed 24 hours after injection. BM monocyte

900   subsets were used as controls. FACS plots (A) and representative data (B) are shown. Each symbol represents an individual mouse.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 1- figure supplement 1

Figure 3- figure supplement 1

A

BM

Ly6C^{hi} Ly6C^{lo}

CCR2 MFI

○ *Rbpj*$^{+/+}$*Lyz2*$^{cre/cre}$
● *Rbpj*$^{fl/fl}$*Lyz2*$^{cre/cre}$

n.s.   n.s.

B

BMT BM

— *Rbpj*$^{+/+}$*Lyz2*$^{cre/cre}$ (CD45.2^{+})
— *Rbpj*$^{fl/fl}$*Lyz2*$^{cre/cre}$ (CD45.2^{+})

Ly6C^{hi} Ly6C^{lo}

% of Max

CCR2   CD11c

Figure 4- figure supplement 1

A

BM

| | Ly6C^hi | Ly6C^lo | Lung Ly6C^hi | Ly6C^lo | CD16.2^+ |

Rbpj^{+/+}*Lyz2*^{cre/cre}

55.4 | 6.29 | 3.16 | 3.88 | 2.02

Rbpj^{fl/fl}*Lyz2*^{cre/cre}

54.5 | 5.08 | 8.16 | 3.25 | 1.30

% of Max

EdU

B

BM | Lung

%EdU^+ — Ly6C^hi | Ly6C^lo | Ly6C^hi | Ly6C^lo | CD16.2^+

○ *Rbpj*^{+/+}*Lyz2*^{cre/cre}
● *Rbpj*^{fl/fl}*Lyz2*^{cre/cre}

Figure 6- figure supplement 1

Blood

Wildtype

RBP-J deficient

Lung

Ly6C^{lo} momocyte

CD16.2^{+} IM

CCR2