Elevated temperature alters microbial communities, but not decomposition rates, during

three years of in-situ peat decomposition

Spencer Roth[1], Natalie A. Griffiths[2,3], Randall K. Kolka[4], Keith C. Oleheiser[2,3], Alyssa A.

Carrell[1], Dawn M. Klingeman[1], Angela Seibert[5], Jeffrey P. Chanton[6], Paul J. Hanson[2,3], and

Christopher W. Schadt[1,3,7,*]

[1]Biosciences Division, Oak Ridge National Laboratory, Oak Ridge, TN, USA 37830

[2]Environmental Sciences Division, Oak Ridge National Laboratory, Oak Ridge, TN, USA 37830

[3]Climate Change Science Institute, Oak Ridge National Laboratory, Oak Ridge, TN, USA 37830

[4]Northern Research Station, USDA Forest Service, Grand Rapids, MN, USA 55744

[5]Department of Geosciences, Boise State University, Boise, ID, USA 83725

[6]Department of Earth, Ocean, and Atmospheric Science, Florida State University, Tallahassee,

FL, USA 32306

[7]Department of Microbiology, University of Tennessee, Knoxville, TN, USA 37996

*Author for correspondence: Christopher Schadt, schadtcw@ornl.gov

Key words: Peatlands, climate change, microbiome, organic matter decomposition,

This manuscript has been authored by UT-Battelle, LLC under Contract No. DE-AC05-00OR22725 with the U.S.
Department of Energy. The United States Government retains and the publisher, by accepting the article for
publication, acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-
wide license to publish or reproduce the published form of this manuscript, or allow others to do so, for United
States Government purposes. The Department of Energy will provide public access to these results of federally
sponsored research in accordance with the DOE Public Access Plan (http://energy.gov/downloads/doe-public-
access-plan).

Abstract:

Peatlands store approximately one-third of the global terrestrial carbon and are historically considered carbon sinks due to primary production outpacing microbial decomposition of organic matter. Climate change has the potential to alter the rate at which peatlands store or release carbon, and results from the Spruce and Peatland Responses Under Changing Environments (SPRUCE) experiment have shown net losses of organic matter and increased greenhouse gas production from a boreal peatland in response to whole-ecosystem warming. In this study, we utilized the SPRUCE sites to investigate how warming and elevated CO_2 impact peat microbial communities and peat soil decomposition. We deployed peat soil decomposition ladders across warming and CO_2 treatment enclosures for three years, after which we characterized bacterial, archaeal, and fungal communities through amplicon sequencing and measured peat mass and compositional changes across four depth increments. Microbial diversity and community composition were significantly impacted by soil depth, temperature, and CO_2 treatment. Bacterial/archaeal α-diversity increased significantly with increasing temperature, and fungal α-diversity was significantly lower under elevated CO_2 treatment. Trans-domain microbial networks showed higher complexity (nodes, edges, degree, betweenness centrality) of microbial communities in decomposition ladders from warmed enclosures, and the number of highly connected, hub taxa within the networks was positively correlated with temperature. Methanogenic hubs were identified in the networks constructed from the warmest enclosures, indicating increased importance of methanogenesis in response to warming. Microbial community responses were not however reflected in measures of peat soil decomposition, as warming and elevated CO_2 had no significant short-term effects on soil mass loss or composition. Regardless of treatment, on average only 4.5% of the original soil mass was

lost after three years and variation between replicates was high, potentially masking treatment

effects. Many previous studies from the SPRUCE experiment have shown that warming is

accelerating organic-matter decomposition and CO_2 and CH_4 production, and our results suggest

that these changes may be driven by warming-induced shifts in microbial communities.

1. Introduction:

Despite covering less than 10% of the Earth's surface, peatlands contain approximately one-

third of all global terrestrial organic matter (OM) [1-3]. Peatland organic soil deposits can be

several meters deep, the result of thousands of years of net primary production outpacing OM

mineralization. Microorganisms are primarily responsible for OM degradation in peatlands [4,

5]; however, anoxic, acidic, oligotrophic, and cold conditions that are common across northern

peatlands greatly constrain microbial activity [4].

Climate change has the potential to alter peatland biogeochemistry, especially at northern

latitudes where warming is occurring at an accelerated pace compared to equatorial regions [6].

The large carbon stocks in northern peatlands that have built up over millennia may therefore be

vulnerable to climate change [7]; however, the effects of warming and increased atmospheric

CO_2 on peatland ecosystems remain to be fully described [8, 9]. The Spruce and Peatland

Responses Under Changing Environments (SPRUCE) experiment is a long-term warming and

elevated CO_2 experiment investigating peatland responses to climate change on an ecosystem

level (https://mnspruce.ornl.gov/). Since 2016, whole-ecosystem warming up to +9 ºC above

ambient has been applied to a boreal peatland in a regression-based design. In addition, elevated

air partial pressure of CO_2 has been applied to half of the 10 SPRUCE experimental enclosures.

Results from the SPRUCE experiment have shown significant, rapid loss of carbon from the

peatland with increasing temperature [10] concomitant with a large decline and death of

Sphagnum sp. at the highest temperatures, that contribute the largest share of Gross Primary

Production (GPP) in these ecosystems [11]. In addition, porewater concentrations of CO_2 and

CH_4 have been shown to correlate with temperature treatment at SPRUCE [12], and radiocarbon

analysis of soil suggests that warming is promoting microbial respiration of solid-phase peat

[13]. Similar results have also been obtained in studies of other peatlands [14-16], and in

incubation studies that show increasing CH_4 and CO_2 production with warming [17, 18].

Net losses of carbon from the SPRUCE sites have been attributed to increased degradation of

OM, rather than a reduction in primary production [10]. Under anoxic conditions, microbial

degradation of OM involves multiple steps including hydrolysis, fermentation, and anaerobic

respiration. Mineralization of carbon to CO_2 and CH_4 in peatlands therefore relies on microbial

metabolic interactions which may be altered by climate change. Incubation experiments have

demonstrated that microbial activities [19, 20] and metabolic interactions are significantly altered

by warming [15], although impacts on community structure and diversity vary [17, 18, 21, 22].

In situ results showing increasing $CH_4:CO_2$ with warming suggest that microbial interactions in

the SPRUCE sites may be altered to favor increased methanogenesis [12, 23].

Previous studies have investigated the effects of warming on peat soil decomposition through

incubations or whole-ecosystem assessments [10, 12, 13, 17, 18, 24]. Valuable insights have

been gained from these experiments; however, incubation studies do not fully reflect

environmental conditions, and "bottle effects" may influence microbial community structure

[25]. Conversely, assessing decomposition from *in situ* environmental measurements is complex

and may be influenced by other ecosystem processes such as changes in primary productivity. To

overcome these limitations, we utilized new peat soil decomposition ladders, which are peat litter

bags attached to a rigid frame, to assess the impacts of temperature, and CO_2 treatments on peat

soil decomposition at four depths. This approach allows for *in situ* investigation of

decomposition while controlling for the effects of primary productivity and excluding fresh litter

inputs. While studies of fresh plant litter decomposition using similar methods are quite common

across many forest ecosystem types[26], studies of soil and particularly peat decomposition

using decomposition bag methods appear to be absent from the literature. Peat decomposition

ladders were deployed in the top 40 cm of peat in the 10 SPRUCE experimental enclosures, as

changes in OM mineralization have been most pronounced in the surface and intermediate layers

of peat [12, 24, 27]. Following a three-years of *in situ* incubation, we measured changes in peat

soil mass and chemical composition and characterized microbial communities in the

decomposition bags through amplicon sequencing and network analyses. We hypothesized that

decomposition of peat soil would increase with increasing temperature, driven by changes in

their microbial communities.

2. Methods:

2.1 Site description

The SPRUCE experiment is located on the S1 bog (low pH, acid organic soil environment) at

the USDA Forest Service Marcell Experimental Forest, MN, USA. The SPRUCE site

description, experimental design, warming, and CO_2 treatments have been previously described

in detail [28]. Briefly, above- and belowground, whole-ecosystem warming has been applied in a

regression-based design to 10 open-air enclosures on the S1 bog since August 2015. Enclosures

are duplicated for each level of warming (+0 ºC, +2.25 ºC, +4.5 ºC, +6.75 ºC and +9 ºC above

ambient), and half of the enclosures receive an elevated CO_2 atmosphere (+500 ppm).

2.2 Peat ladder construction, deployment, and retrieval

Organic soil used in the decomposition ladders was collected from within the S1 bog but

outside of the footprint of the SPRUCE enclosures. Soil was collected from four depths using a

Macaulay peat corer (0-10, 10-20, 20-30, and 30-40 cm), brought back to the laboratory, and air

dried. After air drying, soil from each depth was homogenized by breaking up the dried soil

using sieves, and large fragments of vegetation (i.e., roots) were removed. Soil was then weighed

(2.4-4.2 g air-dry weight, corresponding to an approximate wet weight of 20 g) and placed into

fine-mesh bags (7-µm mesh size) which were then heat sealed [29, 30].

The fine-mesh bags were placed into decomposition ladders which were made of

acrylonitrile butadiene styrene plastic (design from J. Patrick Megonigal, personal

communication, 2016). The ladders were placed vertically in the peat profile to allow for depth-

specific measurements of soil decomposition. Each ladder had four openings (6.5 cm wide by 9

cm tall) corresponding to the four soil depths (Figure S1). Each ladder was constructed by

placing four, soil-filled, fine-mesh bags, one from each depth, between two plastic ladder

holders. The ladders were lined with mesh (0.5 x 1.0 mm mesh size) to reduce abrasion of the

fine-mesh bags during deployment and retrieval. The two ladder holders were closed with plastic

fasteners.

The ladders were deployed on October 2, 2017 in the ten SPRUCE enclosures. Ladders

were placed vertically into the peat so that the peat collected from 0-10 cm was incubated at that

same depth. Three replicate ladders were deployed per enclosure and retrieval date. The replicate

ladders were placed in three different hollow locations that are associated with companion litter

and wood decomposition studies within an enclosure. Three replicate ladders per enclosure were

deployed and immediately retrieved from the peatland for measurement of initial mass and

chemistry (t=0 y; T_0). On October 15, 2020 (t=3 y; T_f), three replicate ladders per enclosure were

retrieved to measure soil mass loss, changes in chemistry, and microbial community

composition.

2.3 Peat soil mass loss and carbon, nitrogen, and phosphorus analyses

After retrieval, ladders were placed into individual plastic bags and were stored at 4 °C

until processing (within 24 hours). Next, each fine-mesh bag was carefully removed from the

ladder and the soil was weighed to determine wet mass. For the T_f bags, soil from each mesh bag

was sub-sampled so that ~1/2 was retained for dry mass and chemistry measurements, ~1/4 was

frozen at -20 °C for microbial community analyses, and the remainder was frozen (-20 °C) for

archival purposes. Soil collected at T_0 was not sub-subsampled as only dry mass and chemistry

measurements were conducted.

Soil samples for mass loss and chemistry were air dried in a drying room (humidity

<30%) for 1-2 months (until the change in mass was <5%) and then weighed. A subsample from

each T_f sample was oven-dried to calculate an air-dry to oven-dry conversion factor and to

calculate percent peat mass loss on an oven-dry basis. Peat used in construction of

decomposition bags were not oven-dried prior to installation (T_0) into the enclosures to prevent

changes in organic matter quality, however, initial masses were corrected based on the air-dry to

oven-dry conversion calculated from each sample at T_f. Each air-dried sample from T_0 and T_f

was ground (IKA Tube Mill grinder, Wilmington, NC, USA) and a subsample was analyzed for

carbon and nitrogen content using a combustion elemental analyzer (LECO-CHN628 analyzer,

St. Joseph, MI, USA).

2.4 FTIR analysis

Fourier Transform Infrared Spectroscopy (FTIR) analysis was performed on the peat

ladders to analyze the response of peat soil organic carbon fractions to enclosure treatments. The

peat soil was ground into a homogenous powder using a Spex Sampleprep 5100 mixer-mill.

FTIR spectra were collected using a JASCO 6800 FT-IR Spectrometer. Approximately 0.003 g

of sample powder was secured onto the quartz crystal (Si/CaF2), and infrared light from

wavenumbers 4000 cm^{-1} – 650 cm^{-1} was transmitted onto the sample at a resolution of 4 cm^{-1}.

Each spectrum was attenuated total reflection corrected and baseline corrected to account for

variability in the beam penetration depth. To produce the functional data for molecular

composition analysis, eight spectra per sample were averaged.

The spectra data were analyzed using Hodgkins' normalization method [31]. Instrument

and matrix variation impact on sample spectra absorbance were accounted for by dividing the

baseline-corrected peak heights by the total integrated area of the spectrum. Using the maximum

baseline-corrected absorbance between peak endpoints, the aromatics and carbohydrates

functional group locations were identified. The normalized aromatics spectral peak heights were

located at 1510 cm^{-1} and 1615 cm^{-1}. The normalized carbohydrate spectral peak height was at

1040 cm^{-1}. Each of these peak heights was used to calculate the percent of aromatics and

carbohydrates in each sample.

2.5 DNA extraction and sequencing

DNA was extracted from ~0.2 g of field-wet peat soil from all samples using the Qiagen

DNeasy 96 PowerSoil Pro QIAcube HT Kit (Qiagen, Venlo, The Netherlands) following the

manufacturer's protocol. Amplicon metagenomic sequencing libraries were prepared as

described in the Illumina 16S metagenomic sequencing library preparation guide (Part 15044223

Rev B) with a mixture of 515F and 806R primers for archaea/bacteria and custom primers

designed to the ITS2 gene region for fungi [32, 33]. Pooled libraries for each sample type were

validated on an Agilent Bioanalyzer (Agilent, Santa Clara, CA) using a DNA7500 chip, and the

final library pool concentration was determined on an Invitrogen Qubit (Waltham, MA) with the

broad range double stranded DNA assay. Paired end sequencing (2x 251x 8x 8) was competed

on an Illumina MiSeq instrument (Illumina, San Diego, CA) using v2 chemistry. Due to low base

diversity of the amplicons, PhiX control DNA was included in the sequencing run.

2.6 Sequence analysis

Demultiplexed 16S rRNA (V4 region) and ITS2 paired-end sequences were imported into

QIIME2 (v. 2021.4) [34]. ITS2 primer sequences were removed using the *cutadapt* plugin [35],

and all sequencing runs were individually denoised using the dada2 plugin [36]. Resulting

feature tables and representative sequences were merged for downstream analyses. Taxonomy

was assigned using the silva database (16S) [37] and UNITE database (ITS) [38]. Sequences and

feature tables were filtered based on taxonomic assignments to include only Bacteria and

Archaea, removing Chloroplast and Mitochondria sequences (16S) and Fungi with taxonomy

assigned to the phylum level at minimum (ITS). A rooted phylogenetic tree was built for each

dataset using the *align-mafft-to-fasttree* pipeline [39] in the *phylogeny* plugin.

2.7 Statistical analyses

All statistical analyses and figures were produced in R v. 4.1.0 with the *vegan* [40], *car*,

phyloseq [41], *ggplot2* [42], *microbiome*, *SpeicEasi* [43], *igraph* [44], and *hilldiv* [45] packages.

ITS and 16S feature tables, taxonomy, rooted phylogeny, and associated metadata were imported

and analyzed as *phyloseq* objects. Based on the regression design of SPRUCE, temperature

treatment (+0 ºC, +2.25 ºC, +4.5 ºC, +6.75 ºC, +9 ºC) was used as a continuous variable for all

statistical analyses. Soil depth and CO_2 treatment were treated as categorical variables, and

depth-specific subsets of the data were generated to investigate the effects of temperature and

CO_2 at a given depth. Samples were rarefied for α-diversity analyses (16S rarefaction depth =

9500; ITS rarefaction depth = 1000). The *hilldiv* package was used to calculate α-diversity

metrics as Hill numbers (effective number of species) [46], and two-way ANOVA analysis was

used to investigate the effects of temperature, soil depth, CO_2 treatment, and the interaction

between temperature and depth on α-diversity. Bray-Curtis dissimilarities [47] of total sum

scaled data were calculated and used to assess the effects of temperature, depth, CO_2 treatment,

and the interaction between temperature, CO_2, and soil depth on community composition by

PERMANOVA. Principal coordinates analysis plots of Bray-Curtis dissimilarities were

generated to visualize community composition. The *betadisper* function was used to assess

differences in β-dispersion across depths, temperature treatments, and CO_2 treatments.

Trans-domain networks were generated using the *SpiecEasi* and *iGraph* packages in R.

Networks were constructed for each temperature treatment and included all depths within a given

temperature treatment (n=24 network^{-1}). Phyloseq objects were filtered by temperature treatment,

and only amplicon sequence variants (ASVs) with a total sum of > 5 (16S) or > 3 (ITS) and an

occurrence in > 20% of the samples were included in the network analyses. If a sample was

lacking either a 16S or ITS library after filtering, the sample was excluded from analysis.

Networks parameters were set as method=mb, nlambda=50, lambda.min.ratio=1e-3, and

thresh=0.01. Empty nodes were removed, and networks were visualized using *phyloseq*. The

number of nodes, edges, node degree and betweenness centrality were calculated for each

network with the *igraph* package. Network hubs were identified by selecting nodes that had

degree and betweenness centrality measures in the 90[th] percentile, indicating high connectedness

and centrality in the network. Pearson correlation was used to investigate the relationship

between network topology and SPRUCE temperature treatments.

Kruskal-Wallis test was used to compare peat soil mass and composition at T_0 to T_f. Effect

sizes for T_0-T_f comparisons were calculated as $(\chi^2 - 1)/(n - 2)$, where χ^2 is the Kruskal-Wallis test

result and n is the number of samples. Linear models were used to assess the effect of depth,

temperature treatment, and CO_2 treatment on soil mass loss and compositional changes.

2.8 Data availability

Amplicon sequences generated in this study have been deposited in the NCBI SRA under

BioProject ID PRJNA941900. All code used in statistical analysis of the data and generation of

figures is available at https://github.com/swroth/Peat_Decomp_SPRUCE.

3.   Results:

3.1 Community summary

Bacterial/archaeal communities were dominated by the phylum *Acidobacteriota* across soil

depths and temperature treatments (28-46% mean relative abundance) (Figure S2). Near the

surface (0-10 cm and 10-20 cm), *Actinobacteriota*, *Proteobacteria*, and *Planctomycetota*

comprised approximately 35-45% of the bacterial/archaeal communities, while accounting for <

20% of the total community at 20-30 cm and 30-40 cm (Figure S2). Dominant phyla of the

fungal communities also varied across depth, with *Basidiomycota* in highest relative abundance

at 0-10 cm (~62% average relative abundance), and *Ascomycota* dominating in the deeper three

depths (~54-68% average relative abundance) (Figure S3).

3.2 Microbial community composition and α-diversity are significantly influenced by

temperature and CO_2 treatments

Bacterial/Archaeal and fungal community compositions were significantly impacted by

depth, temperature treatment, CO_2 treatment, and the interactive effects of these variables

(Figure 1, Table 1). Of the variables investigated, depth was the most influential factor driving

compositional differences in the bacterial/archaeal and fungal communities, explaining 25% and

11% of the variation, respectively (Table 1). Temperature treatment explained 6.1% of the

variation in the bacterial/archaeal community and 4.9% in the fungal community, and CO_2

treatment accounted for 1.9% and 2.7% of the variation for both bacterial/archaeal and fungal

communities respectively (Table 1). Significant interactive effects between soil depth and

temperature treatment, and between temperature treatment and CO_2 treatment were observed for

the bacterial/archaeal communities (Table 1). Significant interactions between all tested variables

were observed for fungal communities, such that temperature and CO_2 treatment only influenced

fungal community composition at specific depths (Table 1). No significant differences in β-

dispersion were observed across soil depth, temperature treatment, or CO_2 treatment.

Bacterial/archaeal α-diversity was significantly highest near the surface and declined with

depth across all levels of q (species richness at q=0, Shannon entropy at q=1, Inverse Simpson at

q=2) (Figure 2A, Figure S4, Table 2). Temperature treatment also significantly impacted α-

diversity, with highest diversity measured in decomposition ladders from the +9 ºC enclosures

across all α-diversity metrics (Figure 2A, Figure S4, Table 2). Bacterial/archaeal α-diversity was

significantly higher in elevated CO_2 enclosures when compared to ambient CO_2 at q=0 and q=1

(Figure 2A, Figure S4A, Table 2). At q=2, no effect of CO_2 on α-diversity was observed,

suggesting that CO_2 treatment has a limited effect on abundant bacterial/archaeal taxa (Figure

S4B, Table 2). In contrast, fungal α-diversity was significantly lower in enclosures with elevated

CO_2 when compared to ambient CO_2 enclosures across all α-diversity metrics measured (Figure

2B, Figure S5, Table 2). Soil depth only significantly impacted fungal species richness (q=0),

with highest richness observed at 0-10 cm, and we observed a significant effect of temperature

treatment at q=2 (Figure 2B, Figure S5, Table 2) and significant interactive effects of

temperature and depth at q=1 and q=2, such that temperature effects on fungal α-diversity were

depth specific.

3.3 Trans-Domain networks

We constructed temperature treatment-specific, trans-domain networks to investigate the

impact of temperature treatment on the connectedness of microbial communities through

comparisons of network topology (Figure S6). The microbial network from +0 ºC enclosures

across all depths had the lowest number of nodes (taxa), edges (connections between taxa), and

average degree (mean number of edges per node), collectively indicating lower complexity of the

microbial communities in +0 ºC enclosures when compared to warmed enclosures (Figure 3).

The number of nodes and edges and the average degree peaked in the network built from +4.5 ºC

enclosures (Figure 3A,B,C). Networks built from all the temperature treatments included

bacterial, archaeal, and fungal nodes, with the highest number of archaeal nodes present in the +9

ºC network (35 nodes), and the most fungal nodes occurring in the +4.5 ºC network (126 nodes)

(Figure 3B).

Within each temperature treatment network, we identified hub taxa as those with degree and

betweenness centrality measures in the 90[th] percentile. The number of hub taxa was significantly

correlated with temperature (R^2 = 0.975, p = 0.005), with a high of 71 hubs in the +9 ºC network

(Figure 3D). Archaeal hubs were present in the +4.5 ºC, +6.75 ºC and +9 ºC networks, and

fungal hubs were observed in all networks except the +2.25 ºC network (Figure 3D). Class-level

taxonomic distribution of the hub taxa revealed that *Acidobacteriae* were the most prominent

across all treatments and depths, followed by the *Verrucomicrobiae* (Figure 4). In the +9 ºC

network, six unique classes were represented in the hubs, including the methanogenic class

Methanosarcinia (Figure 4). Two unique classes were present in the +0 ºC network hubs,

Myxococcia and *Dehalococcoidia* (Figure 4). Only three of the fungal hubs could be assigned to

a guild by FUNguild analysis, including possible ectomycorrhizal and saprotrophic

Agaricomycetes hubs and a *Sordariomycetes* hub that is a probable plant pathogen.

3.4 Peat soil mass loss and compositional changes are highly variable

On average, the mass of peat soil in decomposition ladders was significantly lower after three

years of treatment ($\bar{x} \pm$ SD mass loss = 4.50% $\pm$ 11.07%; Kruskal-Wallis χ^2 = 6.7158, p =

0.0095) (Figure 5A); however, no significant effects of depth, temperature, CO_2 treatment, or

their interactions on peat soil mass loss were observed (Figure 6A; Table S1). Peat C and N

content decreased significantly over the course of the experiment (Figure S7), and C:N of the

peat was significantly higher at T_f compared to T_0 (Kruskal-Wallis χ^2 = 7.0841, p = 0.0078)

(Figure 5B), indicating a more rapid loss of N compared to C. Similar to soil mass loss, neither

depth nor the treatment variables had a significant effect on the change in C:N (Figure 6B; Table

S1).

Results of Fourier-transform infrared spectroscopy (FTIR) analyses further showed limited

effects of the treatments on peat soil composition after three years of incubation. Aromatics (%)

decreased significantly over the course of the experiment (Figure 5C), but there was no

significant effect of soil depth or treatment on the change in aromatics (Figure 6C; Table S1).

Depth significantly impacted the change in % carbohydrates, with largest change in the %

carbohydrates at 0-10 cm. However, there was no significant difference in % carbohydrates

between T_0 and T_f (Figure 5D), and temperature and CO_2 treatment did not significantly impact

carbohydrate content (Figure 6D; Table S1).

4. Discussion:

The effects that climate change will have on peatland microbial communities and the

resulting impacts on soil decomposition have yet to be fully resolved but could have outsized

effects due to the massive stores of carbon in peatlands. Previous results from SPRUCE have

shown rapid loss of carbon presumably driven by increased decomposition in response to

elevated temperature [10]. In this study, we utilized decomposition ladders to investigate the

effects of warming and elevated CO_2 on peat microbial communities and decomposition while

limiting inputs from primary productivity. Our results show that bacterial/archaeal and fungal

communities are significantly impacted by the SPRUCE treatments; however, in contrast to

previous research, we did not observe a significant effect of temperature, elevated CO_2, or soil

depth on mass loss or compositional changes. An average of less than 4.5% of the initial mass

was lost over the three-year experiment regardless of treatment, demonstrating the high

recalcitrance of organic soils in peatlands, and suggesting that these prior reported results may

driven by turnover of more recently fixed C rather than the historic C stocks of peat studied here.

These small differences also illustrate just how inherently difficult measure such decomposition

is to measure given unavoidable experimental variability between replicates and other sources of

error, combined with small mass losses of these recalcitrant substrates.

Bacterial/archaeal and fungal community compositions were significantly influenced by the

SPRUCE treatments (Figure 1, Table 1), indicating that increased temperature and atmospheric

CO_2 as a result of climate change may alter microbial ecology in peatlands. Consistent with

previous research, bacterial/archaeal community responses to temperature treatment were more

pronounced than fungal responses [22, 48, 49]. Bacterial/archaeal α-diversity was significantly

highest in the warmest SPRUCE enclosures across all soil depths, whereas the influence of

temperature treatment on fungal α-diversity was depth specific and only observed at q=2 (Figure

2, Table 2). Ecosystem disturbance and environmental stress have been negatively correlated

with microbial diversity [50], thus higher bacterial/archaeal diversity in warmed enclosures

suggests that warming may alleviate environmental stress on prokaryotic communities in

peatlands. Higher bacterial/archaeal diversity may be a direct cause of warming; however,

contrasting results have also been observed in anaerobic peat soil microcosms that investigated

direct warming effects [17, 21]. Indirect effects such as warming-induced increases in substrate

and nutrient availability [51] are therefore more likely driving changes in bacterial/archaeal

diversity by partially relieving nutrient stress. Our experimental design did not allow us to

delineate between direct and indirect effects of warming, as porewater total organic carbon and

nutrient concentrations were largely correlated with temperature treatment on average over the

course of the experiment (Table S2). However, previous research has demonstrated that labile

substrate and nutrient availability and microbial diversity are positively correlated which

supports our results.

Network analyses further revealed the impact of increased temperature on peat microbial

community structure. The number of nodes and edges were highest in the networks from warmed

enclosures (Figure 3), and a positive correlation between the number of input ASVs and edge

and node counts indicates that this is likely a reflection of species richness. Species richness and

changes in network topology such as a decreased ratio of positive to negative edges and

increased modularity in microbial networks have been shown to correspond to higher community

stability [50, 52, 53], potentially through increased microbial functional redundancy [54]. Paired

with diversity measures, our network results indicate that warming may positively influence

microbial community stability.

The abundance of microbial hub taxa (those taxa which are highly connected within the

network) was positively correlated with temperature treatment (Figure 3D). The number of hub

taxa within microbial networks has been associated with functional potential of the microbial

community [55, 56], and hub taxa may exert strong influence over microbiome structure and

ecosystem function regardless of their relative abundance within the community [57, 58].

Therefore, our results suggest that warming may promote increased microbial functional

potential in these peatland ecosystems.

We observed *Methanomicrobia*, *Methanobacteria*, and *Methanosarcinia* hubs in the +6.75

ºC and +9 ºC networks (Figure 4), supporting results that show methanogenesis is an

increasingly important function in peatland carbon-cycle response to warming. The relative

abundance of methanogenic taxa has been shown to increase in response to increasing

temperature in incubations [17], and rates of methane production have largely been shown to

increase with warming [59], including in incubation studies of soil from the S1 bog [17].

Detection of an acetoclastic methanogen in the +9 ºC network corresponds to isotopic analysis of

CH_4 in the SPRUCE enclosures indicating that acetoclastic methanogenesis is increasing with

warming [59]. A significant linear relationship between *in situ* porewater concentrations of CH_4

and temperature treatment has been observed in the top 25 cm of soil at SPRUCE [12], further

supporting our results. Methane has a global warming potential of 28 times that of CO_2 on a 100-

year time span [6], and higher potential for methanogenesis in response to climate change may

fuel a positive feedback loop.

Methanogenesis can be supported by syntrophic interactions, especially under nutrient-

limiting conditions that are observed in bogs [60, 61]. Two known syntrophic taxa, *Syntrophia*

[62] and *Syntrophorhabdia* [63], were identified as hubs within the networks (Figure 4),

suggesting the potential importance of syntrophy within the sites. Other hubs identified in the

networks from warmed enclosures including *Bathyarchaeia* and *Holophagae* may further

support increased acetoclastic methanogenesis, as these taxa have been previously shown to have

the potential for acetogenesis in anoxic environments [64, 65].

Studies of peatland responses to disturbance often investigate fungal and bacterial/archaeal

communities independent of one-another despite knowledge of the complex interplay across

microbial domains [66, 67]. Here, we observed the highest number of fungal nodes and hub taxa

in networks from decomposition ladders in warmed enclosures, suggesting that peatland

warming may promote trans-domain interactions, further arguing for such holistic approaches.

Possible saprotrophic and ectomycorrhizal fungal hub taxa were primarily observed in the

networks from warmed enclosures apart from an *Agaricomycetes* hub in the +0 ºC network

(Figure 4). Fungal communities play an important role in organic-matter (OM) decomposition in

peatlands [68], and warming may favor dominance of saprotrophic and mycorrhizal fungi from

Basidiomycota and *Ascomycota* [69], as partially observed in our study. Additionally, warming

treatments in these systems are inextricably linked to drying of the peat surface and increased

depths to water table. It is thus possible that peat drying and water table changes, more so than

warming, may be responsible for the shifts in fungal communities observed; however, the

average water table height was similar across all SPRUCE enclosures in the week prior to

termination of the experiment (0.17 m $\pm$ 2.47 cm) as well as the three prior months.

In contrast to bacterial/archaeal diversity, fungal α-diversity was significantly lower under

elevated CO_2 compared to ambient. Fungal responses to CO_2 treatment are likely mediated by

plant responses, as CO_2 treatment at SPRUCE is above-ground and is unlikely to directly alter

soil biogeochemistry. Interactions between plant roots and fungi are common, and fungi are most

prevalent (absolute abundance) near the surface of the peatland where active plant growth occurs

[70]. Our results are intriguing and suggest further investigation into plant-fungal interactions

under elevated CO_2 conditions.

Significant changes in the peat microbial communities in response to SPRUCE treatments

were not mirrored by the peat soil decomposition rates. We anticipated that increased

temperature would result in increased peat soil mass loss, as temperature treatment at SPRUCE

has resulted in rapid carbon loss that was presumed to be driven by enhanced decomposition

[10], increased CO_2 and CH_4 in porewaters [12], and increased microbial respiration of solid

phase peat [13]. However, our results showed that soil mass loss and C:N were not significantly

impacted by temperature or CO_2 treatment over the course of three years (Figure 6). The lack of

differences is likely driven by the short time scale and low initial mass of peat soil in the

decomposition bag study compared to the SPRUCE enclosures. Hanson et al. (2020) estimated

the rate of carbon loss from SPRUCE to be 31.3 g $C \cdot m^{-2} \cdot year^{-1} \cdot \,^oC^{-1}$ using approaches including

elevation changes and ecosystem CO_2 flux mass balances. Using this rate to estimate the

expected loss of carbon from the peat decomposition ladders suggests that differences in mass

loss across temperature treatments were on the order of milligrams over a three-year period, thus

likely requiring a level of precision that we were unable to be obtain in our litter bag-based

experiments.

On average, only 4.5% of the initial peat soil mass was lost over the course of the experiment

(Figure 5). The low mass loss demonstrated the recalcitrance of the organic soils in the

decomposition ladders is likely driven by a combination of factors including the anoxic, acidic,

and oligotrophic conditions of the sites, as well as the chemical composition of the peat. Peat

soils at SPRUCE are largely derived from *Sphagnum*, which is known to engineer acidic,

nutrient poor, waterlogged conditions [71, 72] and produces anti-microbial compounds and

metabolites [73, 74], thereby inhibiting microbial degradation processes. Even the mass loss of

fresh *Sphagnum* litter in decomposition bags has been previously shown to be similarly low with

only ~10% of initial mass lost after two years [75, 76], so these lower rates for peat soil should

not be unexpected.

Diffusion of exogenous dissolved organic matter (DOM) into the decomposition ladders may

have also help explain the differences in results between mass loss and community change.

Dissolved organic matter is preferentially mineralized by peatland microorganisms when

compared to solid-phase peat [77], and fresh plant inputs of DOM can even fuel microbial

respiration in deep peat soils [78]. Utilization of exogenous DOM may also partially explain

discrepancies between our results and previous results from SPRUCE that have shown increased

CO_2 and CH_4 production with warming. Porewater concentrations of total organic carbon were

highest in the + 9 ºC enclosures near the termination of our experiment, and higher inputs of

DOM may have led to increased OM mineralization and shifts in microbial community structure

without impacting peat soil mass in the decomposition ladders.

Similar to soil mass-loss observations, FTIR analysis indicated that temperature and CO_2

treatment had no effect on changes in peat soil composition. The relatively short duration of the

experiment may have masked temperature effects on the percent aromatics and carbohydrates of

the peat, although previous results have indicated that carbon at SPRUCE is compositionally

stable [79]. We are hopeful that our future planned ladder extractions at our site with their longer

field incubation periods may allow for better assessment of treatment effects on peat soil

decomposition.

Conclusions

While we did not observe changes in peat soil mass or composition across the SPRUCE

treatments in this study, much previous research from the SPRUCE experiment has shown loss

of OM near the surface and increased greenhouse gas production in response to elevated

temperatures. Our results suggest that these losses in OM may be driven by changes in microbial

community structure and dynamics, as we observed significant changes in microbial diversity

and network structure in response to warming. The apparent decoupling of changes in peat soil

mass and composition and microbial communities may be limited by the very slow peat

decomposition rates and precision of mass loss estimates in our study. Collectively, our results

and previous results from the SPRUCE experiment therefore suggest that climate change may

alter peatland microbial ecology however the ultimate effects of these changes in on rates of

degradation of OM and greenhouse gas production in boreal regions remains unclear at this time.

Acknowledgements

We would like to thank Randy Hedin and Evelyn Magner (US Forest Service Northern

Research Station) for their assistance in assembling and testing the peat decomposition ladders.

This research was sponsored by the Environmental Systems Science Program, U.S. Department

of Energy, Office of Science, Biological, and Environmental Research, as part of the Terrestrial

Ecosystem Science Scientific Focus Area at Oak Ridge National Laboratory. Oak Ridge

National Laboratory is managed by UT-Battelle, LLC, for the U. S. Department of Energy under

contract DE-AC05-00OR22725. This research used resources of the Compute and Data

Environment for Science (CADES) at the Oak Ridge National Laboratory, which is supported by

the Office of Science of the U.S. Department of Energy under Contract No. DE-AC05-

00OR22725.

487 Author contribution statement

488 Conceptualization – NAG, PJH, RKK, CWS; Investigation – SR, RKK, NAG, KCO, AAC, AS,

489 JPC, DMK; Formal analysis – SR; Original draft writing – SR; Draft revision – SR, RKK, NAG,

490 KCO, AAC, AS, JPC, DMK, CWS

492 ORCID:

SR: 0000-0001-9559-1154

NAG: 0000-0003-0068-7714

RKK: 0000-0002-6419-8218

PJH: 0000-0001-7293-3561

KCO: 0000-0002-0875-2720

AAC: 0000-0003-1142-4709

DMK: 0000-0002-4307-2560

AS: 0000-0001-8269-9270

JPC: 0000-0002-3303-9708

CWS: 0000-0001-8759-2448

Table 1: Results of permutational multivariate analysis of variance (PERMANOVA) of
bacterial/archaeal and fungal community compositions based on Bray-Curtis distances.
Significant factors and interactive effects ($p < 0.05$) are bolded.
507

Variable	Bacterial/Archaeal			Fungal		
	F-Statistic	R^2	p-value	F-Statistic	R^2	p-value
Depth	15.402	0.25323	**0.001**	4.0315	0.10959	**0.001**
Temperature Treatment	11.1683	0.06121	**0.001**	5.46	0.04948	**0.001**
CO_2 Treatment	3.3775	0.01851	**0.004**	3.0053	0.02723	**0.001**
Depth:Temperature Treatment	2.907	0.03766	**0.001**	1.9049	0.05178	**0.001**
Depth:CO_2 Treatment	1.2092	0.01988	0.176	1.6339	0.04442	**0.006**
Temperature treatment:CO_2 treatment	4.983	0.02731	**0.001**	2.6901	0.02438	**0.004**
Depth:Temperature Treatment:CO_2 Treatment	1.4101	0.02318	0.069	1.8305	0.04976	**0.001**

Table 2: Resulting p-values from ANOVA describing the effects of depth and SPRUCE treatments on bacterial/archaeal and fungal α-diversity across Hill numbers (q=0, q=1, and q=2). Significant values ($p < 0.05$) are bolded.

Variable	16S			ITS		
	q=0	q=1	q=2	q=0	q=1	q=2
Depth	**<0.001**	**<0.001**	**<0.001**	**<0.001**	0.802	0.296
Temperature Treatment	**0.002**	**<0.001**	**<0.001**	0.728	0.113	**0.034**
CO_2 Treatment	**<0.001**	**0.015**	0.133	**0.009**	**<0.001**	**0.002**
Depth:Temperature Treatment	0.978	0.907	0.856	0.094	**0.013**	**0.003**

Figure Captions:

Figure 1: Principal coordinates analysis of Bray-Curtis dissimilarities of bacterial/archaeal (A) and fungal (B) communities in decomposition ladders. Points are colored based on sample depth, filled based on enclosure CO_2 treatments and shaped based on enclosure temperature treatments.

Figure 2: Bacterial/Archaeal (A) and Fungal (B) α-diversity at q = 0 (species richness) across depths (vertical facets/colors), CO_2 treatments (horizontal facets), and temperature treatments (x-axis).

Figure 3: Summary of network topologies for temperature treatment microbial networks. The number of edges (A) represents the sum total connections between taxa (nodes; B) within the network. Mean degree (C) represents the average number of connections per taxon. Hubs (D) are defined as nodes within the network that are in the 90th percentile for degree and betweenness centrality and represent highly connected and centralized taxa.

Figure 4: Class-level distribution of network hubs across temperature treatments (color). Hubs were aggregated at class-level taxonomic annotations (y-axis) and faceted by kingdom-level taxonomy.

Figure 5: Comparisons of peat soil oven dry mass (A), carbon:nitrogen (B), percent aromatics (C), and percent carbohydrates (D) in peat decomposition ladders at the start of the experiment (T_0) and after three years of incubation in the SPRUCE enclosures (T_f). Box and whisker plots display the median (middle of box), quartiles (top and bottom of box), minimum and maximum values excluding outliers (end of whisker) and outliers (points).

Figure 6: Peat mass loss (% of starting mass lost) (A), percent C:N change ($T_f - T_0$) (B), percent aromatics change ($T_f - T_0$) (C), and percent carbohydrates change ($T_f - T_0$) (D) across depths (facets) and temperature treatments (x-axis).

References:

1. Turunen, J., et al., *Estimating carbon accumulation rates of undrained mires in Finland - application to boreal and subarctic regions.* Holocene, 2002. **12**(1): p. 69-80.
2. Nichols, J.E. and D.M. Peteet, *Rapid expansion of northern peatlands and doubled estimate of carbon storage.* Nature Geoscience, 2019. **12**(11): p. 917-+.
3. Vitt, D.H., et al., *Spatial and temporal trends in carbon storage of peatlands of continental western Canada through the Holocene.* Canadian Journal of Earth Sciences, 2000. **37**(5): p. 683-693.
4. Andersen, R., S.J. Chapman, and R.R.E. Artz, *Microbial communities in natural and disturbed peatlands: A review.* Soil Biology & Biochemistry, 2013. **57**: p. 979-994.
5. Bridgham, S.D., et al., *Methane emissions from wetlands: biogeochemical, microbial, and modeling perspectives from local to global scales.* Global Change Biology, 2013. **19**(5): p. 1325-1346.
6. IPCC, *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change.* 2013, Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press. 1535.
7. Crowther, T.W., et al., *Quantifying global soil carbon losses in response to warming.* Nature, 2016. **540**(7631): p. 104-+.
8. Davidson, E.A. and I.A. Janssens, *Temperature sensitivity of soil carbon decomposition and feedbacks to climate change.* Nature, 2006. **440**(7081): p. 165-173.
9. Bridgham, S.D., et al., *The carbon balance of North American wetlands.* Wetlands, 2006. **26**(4): p. 889-916.
10. Hanson, P.J., et al., *Rapid Net Carbon Loss From a Whole-Ecosystem Warmed Peatland.* Agu Advances, 2020. **1**(3).
11. Norby, R.J., et al., *Rapid loss of an ecosystem engineer: Sphagnum decline in an experimentally warmed bog.* Ecology and Evolution, 2019. **9**(22): p. 12571-12585.
12. Wilson, R.M., et al., *Soil metabolome response to whole-ecosystem warming at the Spruce and Peatland Responses under Changing Environments experiment.* Proceedings of the National Academy of Sciences of the United States of America, 2021. **118**(25).
13. Wilson, R.M., et al., *Radiocarbon Analyses Quantify Peat Carbon Losses With Increasing Temperature in a Whole Ecosystem Warming Experiment.* Journal of Geophysical Research-Biogeosciences, 2021. **126**(11).
14. Schadel, C., et al., *Potential carbon emissions dominated by carbon dioxide from thawed permafrost soils.* Nature Climate Change, 2016. **6**(10): p. 950-+.
15. Tveit, A.T., et al., *Metabolic and trophic interactions modulate methane production by Arctic peat microbiota in response to warming.* Proceedings of the National Academy of Sciences of the United States of America, 2015. **112**(19): p. E2507-E2516.
16. Yvon-Durocher, G., et al., *Methane fluxes show consistent temperature dependence across microbial to ecosystem scales.* Nature, 2014. **507**(7493): p. 488-491.
17. Kolton, M., et al., *Impact of Warming on Greenhouse Gas Production and Microbial Diversity in Anoxic Peat From a Sphagnum-Dominated Bog (Grand Rapids, Minnesota, United States).* Frontiers in Microbiology, 2019. **10**.
18. Kluber, L.A., et al., *Constraints on microbial communities, decomposition and methane production in deep peat deposits.* Plos One, 2020. **15**(2).

19.   Keiser, A.D., et al., *Peatland microbial community response to altered climate tempered*
by nutrient availability. Soil Biology & Biochemistry, 2019. **137**.
20.   AminiTabrizi, R., et al., *Elevated temperatures drive abiotic and biotic degradation of*
organic matter in a peat bog under oxic conditions. Science of the Total Environment,
2022. **804**.
21.   Yang, S.Z., et al., *Decoupling of microbial community dynamics and functions in Arctic*
peat soil exposed to short term warming. Molecular Ecology, 2021. **30**(20): p. 5094-
5104.
22.   Lulakova, P., et al., *High-Alpine Permafrost and Active-Layer Soil Microbiomes Differ in*
Their Response to Elevated Temperatures. Frontiers in Microbiology, 2019. **10**.
23.   Hopple, A.M., et al., *Massive peatland carbon banks vulnerable to rising temperatures.*
Nature Communications, 2020. **11**(1).
24.   Wilson, R.M., et al., *Stability of peatland carbon to rising temperatures.* Nature
Communications, 2016. **7**.
25.   Wilson, R.M., et al., *Functional capacities of microbial communities to carry out large*
scale geochemical processes are maintained during ex situ anaerobic incubation. Plos
One, 2021. **16**(2).
26.   Krishna, M.P. and M. Mohan, *Litter decomposition in forest ecosystems: a review.*
Energy, Ecology and Environment, 2017. **2**: p. 236-249.
27.   Tfaily, M.M., et al., *Vertical Stratification of Peat Pore Water Dissolved Organic Matter*
Composition in a Peat Bog in Northern Minnesota. Journal of Geophysical Research-
Biogeosciences, 2018. **123**(2): p. 479-494.
28.   Hanson, P.J., et al., *Attaining whole-ecosystem warming using air and deep-soil heating*
methods with an elevated CO2 atmosphere. Biogeosciences, 2017. **14**(4): p. 861-883.
29.   Weiss, J.V., D. Emerson, and J.P. Megonigal, *Rhizosphere iron(III) deposition and*
reduction in a Juncus effusus L.-dominated wetland. Soil Science Society of America
Journal, 2005. **69**(6): p. 1861-1870.
30.   Kirwan, M.L., et al., *The impact of sea-level rise on organic matter decay rates in*
Chesapeake Bay brackish tidal marshes. Biogeosciences, 2013. **10**(3): p. 1869-1876.
31.   Hodgkins, S.B., et al., *Tropical peatland carbon storage linked to global latitudinal*
trends in peat recalcitrance. Nature Communications, 2018. **9**.
32.   Dove, N.C., et al., *Microbiome Variation Across Two Hemlock Species With Hemlock*
Woolly Adelgid Infestation. Frontiers in Microbiology, 2020. **11**.
33.   Rogers, T.J., et al., *Exploring variation in phyllosphere microbial communities across*
four hemlock species. Ecosphere, 2018. **9**(12).
34.   Bolyen, E., et al., *Reproducible, interactive, scalable and extensible microbiome data*
science using QIIME 2. Nature Biotechnology, 2019. **37**(8): p. 852-857.
35.   Martin, M., *Cutadapt removes adapter sequences from high-throughput sequencing*
reads. 2011, 2011. **17**(1): p. 3.
36.   Callahan, B.J., et al., *DADA2: High-resolution sample inference from Illumina amplicon*
data. Nature Methods, 2016. **13**(7): p. 581-+.
37.   Quast, C., et al., *The SILVA ribosomal RNA gene database project: improved data*
processing and web-based tools. Nucleic Acids Research, 2013. **41**(D1): p. D590-D596.
38.   Nilsson, R.H., et al., *The UNITE database for molecular identification of fungi: handling*
dark taxa and parallel taxonomic classifications. Nucleic Acids Research, 2019. **47**(D1):
p. D259-D264.

39.   Price, M.N., P.S. Dehal, and A.P. Arkin, *FastTree 2-Approximately Maximum-Likelihood*
Trees for Large Alignments. Plos One, 2010. **5**(3).
40.   Oksanen, J., et al., *vegan: Community Ecology Package.* 2020.
41.   McMurdie, P.J. and S. Holmes, *phyloseq: An R Package for Reproducible Interactive*
Analysis and Graphics of Microbiome Census Data. Plos One, 2013. **8**(4).
42.   Villanueva, R.A.M. and Z.J. Chen, *ggplot2: Elegant Graphics for Data Analysis, 2nd*
edition. Measurement-Interdisciplinary Research and Perspectives, 2019. **17**(3): p. 160-
167.
43.   Kurtz, Z.D., et al., *Sparse and Compositionally Robust Inference of Microbial Ecological*
Networks. Plos Computational Biology, 2015. **11**(5).
44.   Csardi, G. and T. Nepusz, *The igraph software package for complex network research.*
InterJournal, 2006. **Complex Systems**: p. 1695.
45.   Alberdi, A. and M.T.P. Gilbert, *hilldiv: an R package for the integral analysis of diversity*
based on Hill numbers. bioRxiv, 2019: p. 545665.
46.   Chao, A.N., C.H. Chiu, and L. Jost, *Unifying Species Diversity, Phylogenetic Diversity,*
Functional Diversity, and Related Similarity and Differentiation Measures Through Hill
Numbers. Annual Review of Ecology, Evolution, and Systematics, Vol 45, 2014. **45**: p.
297-324.
47.   Bray, J.R. and J.T. Curtis, *An Ordination of the Upland Forest Communities of Southern*
Wisconsin. Ecological Monographs, 1957. **27**(4): p. 326-349.
48.   Xiong, J.B., et al., *Divergent Responses of Soil Fungi Functional Groups to Short-term*
Warming (vol 68, pg 708, 2014). Microbial Ecology, 2014. **68**(4): p. 890-890.
49.   Zhao, B.Y., P. Xing, and Q.L.L. Wu, *Interactions between bacteria and fungi in*
macrophyte leaf litter decomposition. Environmental Microbiology, 2021. **23**(2): p. 1130-
1144.
50.   Hernandez, D.J., et al., *Environmental stress destabilizes microbial networks.* Isme
Journal, 2021. **15**(6): p. 1722-1734.
51.   Iversen, C.M., et al., *Whole-Ecosystem Warming Increases Plant-Available Nitrogen and*
Phosphorus in an Ombrotrophic Bog. Ecosystems, 2022.
52.   Thebault, E. and C. Fontaine, *Stability of Ecological Communities and the Architecture of*
Mutualistic and Trophic Networks. Science, 2010. **329**(5993): p. 853-856.
53.   Tardy, V., et al., *Stability of soil microbial structure and activity depends on microbial*
diversity. Environmental Microbiology Reports, 2014. **6**(2): p. 173-183.
54.   Griffiths, B.S. and L. Philippot, *Insights into the resistance and resilience of the soil*
microbial community. Fems Microbiology Reviews, 2013. **37**(2): p. 112-129.
55.   Faust, K., et al., *Cross-biome comparison of microbial association networks.* Frontiers in
Microbiology, 2015. **6**.
56.   Shi, Y., et al., *Abundance of kinless hubs within soil microbial networks are associated*
with high functional potential in agricultural ecosystems. Environment International,
2020. **142**.
57.   Banerjee, S., K. Schlaeppi, and M.G.A. van der Heijden, *Keystone taxa as drivers of*
microbiome structure and functioning. Nature Reviews Microbiology, 2018. **16**(9): p.
567-576.
58.   Xun, W.B., et al., *Specialized metabolic functions of keystone taxa sustain soil*
microbiome stability. Microbiome, 2021. **9**(1).

| 680 | 59. | Gill, A.L., et al., *Deep peat warming increases surface methane and carbon dioxide emissions in a black spruce-dominated ombrotrophic bog.* Global Change Biology, 2017. **23**(12): p. 5398-5411. |

59. Gill, A.L., et al., *Deep peat warming increases surface methane and carbon dioxide*
emissions in a black spruce-dominated ombrotrophic bog. Global Change Biology, 2017.
23(12): p. 5398-5411.
60. McInerney, M.J., J.R. Sieber, and R.P. Gunsalus, *Syntrophy in anaerobic global carbon*
cycles. Current Opinion in Biotechnology, 2009. **20**(6): p. 623-632.
61. Morris, B.E., et al., *Microbial syntrophy: interaction for the common good.* Fems
Microbiology Reviews, 2013. **37**(3): p. 384-406.
62. McInerney, M.J., et al., *Physiology, ecology, phylogeny, and genomics of*
microorganisms capable of syntrophic metabolism. Incredible Anaerobes: From
Physiology to Genomics to Fuels, 2008. **1125**: p. 58-72.
63. Qiu, Y.L., et al., *Syntrophorhabdus aromaticivorans gen. nov., sp. nov., the first cultured*
anaerobe capable of degrading phenol to acetate in obligate syntrophic associations with
a hydrogenotrophic methanogen. Applied and Environmental Microbiology, 2008. **74**(7):
p. 2051-2058.
64. He, Y., et al., *Genomic and enzymatic evidence for acetogenesis among multiple lineages*
of the archaeal phylum Bathyarchaeota widespread in marine sediments. Nature
Microbiology, 2016. **1**(6).
65. Liesack, W., et al., *Holophaga Foetida Gen-Nov, Sp-Nov, a New, Homoacetogenic*
Bacterium Degrading Methoxylated Aromatic-Compounds. Archives of Microbiology,
1994. **162**(1-2): p. 85-90.
66. de Vries, F.T., et al., *Soil food web properties explain ecosystem services across*
European land use systems. Proceedings of the National Academy of Sciences of the
United States of America, 2013. **110**(35): p. 14296-14301.
67. Faust, K. and J. Raes, *Microbial interactions: from networks to models.* Nature Reviews
Microbiology, 2012. **10**(8): p. 538-550.
68. Thormann, M.N., *The Role of Fungi in Boreal Peatlands*, in *Boreal Peatland Ecosystems*,
R.K. Wieder and D.H. Vitt, Editors. 2006, Springer Berlin Heidelberg: Berlin,
Heidelberg. p. 101-123.
69. Asemaninejad, A., et al., *Climate change favours specific fungal communities in boreal*
peatlands. Soil Biology & Biochemistry, 2018. **120**: p. 28-36.
70. Lin, X.J., et al., *Microbial Community Stratification Linked to Utilization of*
Carbohydrates and Phosphorus Limitation in a Boreal Peatland at Marcell Experimental
Forest, Minnesota, USA. Applied and Environmental Microbiology, 2014. **80**(11): p.
3518-3530.
71. Vanbreemen, N., *How Sphagnum Bogs down Other Plants.* Trends in Ecology &
Evolution, 1995. **10**(7): p. 270-275.
72. Verhoeven, J.T.A. and W.M. Liefveld, *The ecological significance of organochemical*
compounds in Sphagnum. Acta Botanica Neerlandica, 1997. **46**(2): p. 117-130.
73. Fudyma, J.D., et al., *Untargeted metabolomic profiling of Sphagnum fallax reveals novel*
antimicrobial metabolites. Plant Direct, 2019. **3**(11).
74. Hajek, T., et al., *Cell-wall polysaccharides play an important role in decay resistance of*
Sphagnum and actively depressed decomposition in vitro. Biogeochemistry, 2011. **103**(1-
3): p. 45-57.
75. Scheffer, R.A., R.S.P. van Logtestijn, and J.T.A. Verhoeven, *Decomposition of Carex*
and Sphagnum litter in two mesotrophic fens differing in dominant plant species. Oikos,
2001. **92**(1): p. 44-54.

76.   Verhoeven, J.T.A. and E. Toth, *Decomposition of Carex and Sphagnum Litter in Fens -*
Effect of Litter Quality and Inhibition by Living Tissue-Homogenates. Soil Biology &
Biochemistry, 1995. **27**(3): p. 271-275.

77.   Hopple, A.M., et al., *Does dissolved organic matter or solid peat fuel anaerobic*
respiration in peatlands? Geoderma, 2019. **349**: p. 79-87.

78.   Tfaily, M.M., et al., *Organic matter transformation in the peat column at Marcell*
Experimental Forest: Humification and vertical stratification. Journal of Geophysical
Research-Biogeosciences, 2014. **119**(4): p. 661-675.

79.   Baysinger, M.R., et al., *Compositional stability of peat in ecosystem-scale warming*
mesocosms. Plos One, 2022. **17**(3).
736