Diversity and functions of soil microbial communities in the Trans-Himalayan grazing

ecosystem: influence of herbivores and abiotic factors

Shamik Roy[1,2], Pronoy Baidya[1], Jalmesh Karapurkar[1], M. Jose[1], Sumanta Bagchi[1,3] *

1 Centre for Ecological Sciences, Indian Institute of Science, Bangalore 560012, India.

2 School of Biological Sciences, University of East Anglia, Norwich NR4 7TJ, UK

3 Divecha Centre for Climate Change, Indian Institute of Science, Bangalore 560012, India

*Correspondence: sbagchi@iisc.ac.in

ORCID: SR, 0000-0002-9916-5651; SB, 0000-0002-4841-6748; MJ, 0000-0002-1012-0407;

JK, 0000-0003-1125-2468; PB, 0000-0003-1337-1372.

Author contributions: SR, JK, PB and SB conducted fieldwork. SR, MJ and JK conducted lab

analyses. SR and SB analyzed the data. SR and SB wrote the first draft. All authors

contributed to writing the manuscript.

Originality-Significance statement

Diversity-function relationships in soil microbial decomposers are an important aspect of

global biogeochemical processes. We study how microbial communities and functions

respond to large mammalian herbivores and seasonality, as well as the underlying roles of

abiotic conditions that link to functional redundancy.

Highlights

• Diversity-function relationships for microbial decomposers remain uncertain.

• Microbial diversity is positively related to functions

• This relationship is robust to grazer-exclusion.

• Microbial diversity-function relationship can be susceptible to altered precipitation

Summary

1. Diversity-function relationships are well established for producers and their productivity.

They are also evident among consumers. However, these are not well known for

microbial decomposers and decomposition processes in soil. Further, it also remains

unknown whether and how consumers, such as large mammalian herbivores who are a

major feature across more than one-third of the world's terrestrial realm, influence

microbial decomposer diversity-function relationships.

2. We used a14-year old long-term herbivore-exclusion experiment to answer two

questions: (a) whether microbial functions vary with microbial diversity (both species

richness and composition), and (b) whether herbivores alter the diversity-function

relationships among microbial decomposers. We measured functional richness, functional

dispersion, and multifunction from the utilization profiles of 30 metabolic substrates.

Alongside, we also measured species richness and species composition of soil microbes.

3. Data were from 60 16S rDNA gene amplicon sequences with 7.6 million reads covering

1937 potential species across 47 phyla, and 1800 catabolic profiles.

4. We found microbial functional diversity in soil was positively related (i.e., coupled) to

community composition, but not to species richness. This challenges the prevailing

paradigm of functional redundancy in hyper-diverse soil microbial communities where

changes in species richness is expected to not affect functional diversity. Instead, our

results indicate that certain combinations of species can outperform others. Therefore,

global change factors that alter microbial communities can also impact decomposition

processes and services. This coupling between diversity and functions was unaffected by

experimental herbivore-exclusion, indicating resilience and resistance among

decomposers.

5. Structural equation models suggested that the strength of this relationship is favoured by

availability of soil moisture. These also showed microbial functions varied more strongly

with temporal variables (e.g., seasonality) than with spatial variables (e.g., edaphic factors

such as soil texture and pH).

6. While ecosystem functions and services derived from microbial decomposers have

intrinsic resilience and resistance, they respond strongly to variability in water

availability. Decomposition in grazing ecosystems may be particularly susceptible to how

soil microbes respond to increased precipitation variability under ongoing and projected

climate change.

KEYWORDS:

Biodiversity; Ecosystem function; Functional redundancy; Microbial decomposers; Drylands;

Species richness; Distance-based structural equation model

Introduction

Species diversity begets ecological functions, and high taxonomic diversity promotes

functional diversity as well. Oppositely, depending on ecological resilience and resistance,

loss of diversity can deplete functions. The empirical evidence for such diversity-function

relationships mostly come from producers and their production (Schulze and Mooney, 1993;

Loreau et al., 2001; Hooper et al., 2005; Allison and Martiny, 2008; Reich et al., 2012;

Tilman et al., 2014). In comparison, diversity-function relationships for consumers are more

complex and less generalizable due to variability in nutrient sources across space and trophic

levels (Olff and Ritchie, 1998; Duffy, 2002; Duffy et al., 2007; Cardinale et al., 2011). It is

pertinent to ask whether the patterns and processes seen in producers and consumers can be

extrapolated to decomposers and decomposition processes. But the answer seems to be

contentious as both 'yes' and 'no' have been reported by studies examining diversity-

function relationships for decomposers (Wagg et al., 2014, 2019; Delgado-Baquerizo et al.,

2016a; Louca et al., 2016; Chen et al., 2022). Generally, experimental manipulations tend to

find positive diversity-function relationships in decomposers (Wagg et al., 2014, 2019;

Delgado-Baquerizo et al., 2016a). In contrast, when natural variation in species richness in

ecosystems is weighed against functions, studies tend to find no relationship due to functional

redundancy in decomposers (Louca et al., 2016; Chen et al., 2022). Such contradictory

evidence could be due to few species in experimental studies (c. 10^1 to 10^2 species) (Wagg et

al., 2014, 2019; Delgado-Baquerizo et al., 2016a) and many species in observational studies

(c. 10^3 species) (Louca et al., 2016; Chen et al., 2022). Increasingly, in natural communities

across marine and terrestrial ecosystems, microbial community composition—but not their

species richness—is increasingly seen to influence decomposition functions (Fierer et al.,

2013; Delgado-Baquerizo et al., 2017; Galand et al., 2018). Therefore, the question of

whether, or not, microbial decomposition process in soil is related (i.e., coupled) to

community composition and identity of the species, warrants further investigation. This is

critical for knowing whether diversity can buffer and ameliorate the effect of stressors that

act on decomposition functions and services.

Decomposition processes by microbes do not occur in isolation of other trophic

levels. As consumers of plant production, large mammalian herbivores simultaneously alter

both the quantity and quality of biomass-input to soil. Grazing can influence microbial

community composition and regulate taxonomic profiles of the decomposer community in

soil (Bardgett et al., 2001; Yang et al., 2019; Ylänne et al., 2021). Simultaneously, grazing

can also influence functional attributes of soil microbes (Bagchi et al., 2017; Roy and Bagchi,

2022; Roy et al., 2023). These effects are known to be highly variable across ecosystems, and

can be positive, negative, or neutral (Tanentzap and Coomes, 2012; McSherry and Ritchie,

2013; Wang and Tang, 2019; Eldridge et al., 2020; Roy and Bagchi, 2022; Roy et al., 2023).

Given such heterogeneity, it remains poorly understood whether grazer influence on

microbial functions in soil is mediated through their effects on microbial community

composition, or not. The role of large mammalian herbivores in mediating inter-dependency

between the taxonomic and functional aspects of soil microbial diversity has broad

implications for management and sustainability of grazing in dryland ecosystems.

Grazer effects on soil functions arise from their interactions with microbial

decomposers that can have far-reaching consequences for local, regional, and global

biogeochemical cycles (Bardgett and Wardle, 2003; Bagchi et al., 2017; Naidu et al., 2022;

Roy and Bagchi, 2022; Roy et al., 2023). In addition to grazers, underlying edaphic

conditions such as moisture availability (Bagchi et al., 2017; Zhang et al., 2022), soil pH

(Fierer et al., 2006; Delgado-Baquerizo et al., 2016b), and texture (Girvan et al., 2003) can

also affect microbial decomposers. Moreover, soil microbial communities and functions can

be influenced by seasonality as they undergo high degree of temporal change as well

(Frossard et al., 2012; Bagchi et al., 2017; Roy et al., 2023). Therefore, one can expect that

microbial diversity-function relationship can also vary temporally (Wang et al., 2023).

Here we assess microbial diversity-function relationship by measuring variation in

microbial diversity and functions at regular intervals through the growing season in the

Trans-Himalayan ecosystem of northern India. The specific questions addressed were: (1)

whether microbial diversity (both species richness and composition) and functions vary with

grazing and across the growing season (3 levels: early, mid, late); (2) whether microbial

community composition influences decomposition functions; (3) whether herbivores alter the

relationship between functions and species composition that is otherwise mediated by

edaphic conditions and seasonality.

Methods

Study site and sampling

We used a long-term replicated grazer-exclusion experiment started in 2005-06 in Spiti

region of Himachal Pradesh in northern India (32° N, 78° E) (Bagchi and Ritchie, 2010a).

We set up replicated herbivore exclosures (ungrazed plot), each 10×10 m^2, with adjacent

paired control grazed plots to understand the effect of large mammalian herbivores on the

ecosystem. Spiti region is a part of the larger Trans-Himalayan drylands across Central Asia

(Fig. A.1). This high-altitude grazing ecosystem (elevation 4400-4800 m) experiences cold

and semiarid climate with temperature dropping below -30 °C in winters. Vegetation

growing-season starts in May, peaks in July-August, and senescence begins in September.

Large mammalian herbivores comprise of native species (bharal *Pseudois nayaur*, ibex

Capra sibirica, and yak *Bos grunniens*) and livestock (goat, sheep, donkey, horse, cattle, and

yak-cattle hybrids) (Bagchi and Ritchie, 2010a, 2010b).

After 14 years of the grazer-exclusion experiment, in 2019, we sampled soil from

n=10 paired-adjacent plots, in May (early-season), July (peak-season), and September (end-

season). We used a 2.5 cm diameter and 20 cm depth corer for soils meant for microbial

functions and edaphic variables. A smaller sample was packaged in sterilized vials for DNA

analysis.

Soil microbial functions

We measured soil microbial functions and functional diversity using n = 30 metabolic

substrates (Degens and Harris, 1997; Degens, 1998). In this way, metabolic profiles were

obtained from 30×10×3×2=1800 measurements, plus corresponding controls (Fig. 2). We

used substrates from different chemical groups: amino acids (Alanine, Arginine, Serine,

Asparagine, Glutamine, Glutamate, Histidine), amines (Glucosamine, Putrescine), alcohols

(Xylitol, Mannitol, Sorbitol, Glycerol), carbohydrates (D-Glucose, Xylose, Mannose,

Sucrose, Inositol), carboxylic acids (α-Ketoglutaric acid, Fumaric acid, Succinic acid,

Tartaric acid, Oxalic acid, Malic acid, Ascorbic acid, Citric acid, Malonic acid), an aromatic

compound (Inosine), a phenolic compound (Gallic acid), and a polymer (Polysorbate). We

measured soil respiration after addition of substrate (substrate induced respiration, SIR)

compared to controls (basal respiration) (Degens and Harris, 1997). A substrate was judged

to be catabolized if the corresponding SIR was above the 95% confidence-interval of basal

respiration (i.e., treatment > control). We used the substrates concentrations such that equal

amounts of C was added across all treatments (Degens and Harris, 1997): 100 mM for

carboxylic acids and alcohols; 75 mM for carbohydrates; 40 mM for polymers; 15 mM for

aromatic, amino acids, amines, and phenols. We pre-incubated 5 g soil at 60% water holding

capacity at 28 °C for 24 h. Next, we added 1 ml of substrate solution and incubated soils in

air-tight containers along with of 1.5 ml Cresol Red indicator dye (prepared from 2.5 mM

NaHCO$_3$, 150 mM KCl, 12.5 μg ml^{-1} Cresol Red dye) at 28 °C for 15 h (Rowell, 1995). Next,

we estimated substrate utilization (mg C g^{-1} soil day^{-1} respired as CO_2) from colorimetric

measurement of Cresol Red indicator at 570 nm wavelength, alongside corresponding

controls and standards (Rowell, 1995).

Soil DNA extraction, amplification, and sequencing

We estimated microbial taxonomic diversity from Operational Taxonomic Units (OTUs) in

DNA extracted from soil. Taxonomic profiles were obtained from 10×3×2=60 samples, plus

corresponding controls (Fig. 2). We used MOBIO Power Soil DNA Extraction Kit

(Netherlands) following manufacturer instructions. DNA extracts were stored at -20 °C until

further molecular analysis. We performed amplification for 16S rRNA hyper-variable region

V3-V4 using 25 ng DNA extract, KAPA HiFi HotStart Ready Mix, and 100 nM of 341F (5'-

CCTACGGGNGGCWGCAG-3') and 785R (5'-GACTACHVGGGTATCTAATCC-3')

primers (Klindworth et al., 2013). Initial denaturation was at 95 °C for 5 min followed by 25

cycles of 95 °C for 30 s, 55 °C for 45 s and 72 °C for 30 s with a final extension at 72 °C for

7 min. Amplicons were sequenced using Illumina MiSeq platform and data were processed

using Mothur (Schloss et al., 2009) to obtain microbial taxonomy data. The raw sequences

were aligned to form contigs, and only contigs with length between 300 bp and 532 bp were

retained because they cover the targeted error-free sequence of the 16S rRNA region with

high confidence. Contigs were then checked for chimeras, and removed, using reference-

based chimera checking following UCHIME algorithm (Edgar et al., 2011). The contigs were

then categorized into OTUs with 97% sequence similarity using SILVA_v138 database

(Quast et al., 2013). As an additional level of quality filtering, OTUs with cumulative

abundance of less than 20 across all samples in the data set were removed. Controls showed

minimal absorbance at 260 nm (<1 ng/μl of DNA) compared to samples (24 to 70 ng/μl of

DNA), and we did not sequence them since they did not show any amplification.

Edaphic variables

We measured three key edaphic variables (Fig. A.2), namely soil moisture, pH, and texture

following standard procedures (Robertson et al., 1999) as these can influence variation in

microbial communities and in functions (Fierer and Jackson, 2006; Sinsabaugh et al., 2008).

Briefly, soil moisture was measured as volumetric water content with a time domain

reflectometry probe (Spectrum Technologies, USA), pH was measured from 1:2.5 (v/w)

aqueous suspension using a table-top pH meter, and soil texture was measured from 50 g soil

in 1000 ml of $(NaPO_3)_6$ solution using a Bouyoucos hydrometer.

Statistical analyses

We summarized microbial functions with three indices **a) Functional richness:** calculated as

the number of substrates metabolized. b) **Dispersion:** multivariate measure of functional

heterogeneity (dispersion) estimated as the distance of each sample relative to their respective

group centroid (Anderson et al., 2006). We used 'vegan' library in R 3.6.3 to estimate

richness and dispersion (Oksanen et al., 2019). **c) Multifunction:** measured as sum of SIR

across 30 catabolites (mg C g^{-1} soil hour^{-1}). This represents the decomposition potential of the

microbial communities. Multifunctionality is the ability of an ecosystem to perform multiple

functions, or provide several services, simultaneously (Hector and Bagchi, 2007).

We estimated species richness as the number of OTUs in a sample. We used Generalized

Linear Models (GLM) with grazing (grazed/ungrazed) and season (early, mid, and late) as

fixed-effects, alongside plot-identity as the random-effect to assess the effects of grazing and

season on species richness and functional diversity. This GLM accounts for the pair-wise

spatial structure in the data (Pinheiro and Bates, 2000), and we used 'nlme' library in R 3.6.3.

We verified whether the data met the assumptions of GLMs using theoretical and observed

quantiles of residuals, and the data did not require any transformations.

Microbial composition and function

We used ordination with principal coordinates analysis (PCoA) of pair-wise Bray-Curtis

dissimilarity to visualise species composition and decomposer functions across seasons and

grazing treatment in a multivariate space summarized as two axes (Legendre and Legendre,

2012).

We used kriging to visualize the relationship between species composition and microbial

functions. Kriging is an interpolation method that helps estimate the value of a variable (e.g.,

functional richness) over a continuous space of microbial species composition using an

already existing but a limited set of sampled data points (Dale and Fortin, 2014; Li et al.,

2023). We used PCoA axes 1 and 2 as a measure of species composition to interpolate the

values of functional indices for unmeasured values of species composition. Kriging is a two-

step process: first, a variogram is fitted to determine the spatial (species composition space)

covariance structure of the sampled data points; second, interpolate values for the unsampled

locations across a given species composition space from the weights derived from the spatial

covariance structure of the known data points (Dale and Fortin, 2014; Li et al., 2023). For

this, various variogram models were fitted by changing the trend removal parameter (no trend

removal, 1^{st} order, 2^{nd} order), model function (exponential, spherical, gaussian), and nugget

(fixed, unfixed). First step in fitting the variogram model is trend removal. Trend removal is

done by fitting a polynomial equation to the predictor values (PCoA axes 1 and 2) using

ordinary least squares, and therefore the variograms are computed using the residuals. 1^{st}

order trend removal will remove linear trends, whereas 2^{nd} order will remove quadratic

trends. Trend removal is followed by fitting a model to the empirically determined

semivariogram using any of the three model functions where nugget (y-intercept value) is

either fixed at zero or unfixed (variable). For each microbial functional index there were 18

different combinations (3 trend removal parameters $\times$ 3 model function $\times$ 2 nugget) to build a

variogram model. We selected the final combination for each functional index where the

resulting model had the least sum of squares (Table A.1). For all the functional indices we

selected 2nd order trend removal with exponential function and nugget fixed at zero. We

implemented kriging with "geoR" package in R (Justiniano {Ribeiro Jr} et al., 2022).

Distance-based structural equation models

We used distance-based structural equation models (dbSEM) to evaluate the relationship

between species composition and function, and whether grazers affect the relationship

directly or indirectly through edaphic factors. dbSEM uses pairwise-dissimilarity matrix for

both dependant and predictor variable to disentangle direct and indirect hypothesized causal

relationships (Fourtune et al., 2018; Jabot et al., 2020). We used permutation based dbSEM

where SEM is performed on the input matrices of variables where rows and columns are

randomly permuted to obtain an unbiased P-value for each hypothesized path (Fourtune et al.,

2018; Jabot et al., 2020). P-value is computed as the proportion where permuted path

coefficient values are less than or equal to the observed path coefficient value. The edaphic

variables were soil moisture, pH and texture. For soil texture we only included sand content

in the analysis because of autocorrelation with clay and silt content. The paths in the dbSEM

were motivated by important processes that shapes up microbial communities and known *a-*

priori. a) Large mammalian herbivore alter microbial communities (Grazing $\rightarrow$ Species

composition) (Sankaran and Augustine, 2004; Bagchi et al., 2017). b) Herbivore can regulate

decomposition process directly (Grazing $\rightarrow$ Function (functional richness, functional

composition, multifunction)), or indirectly through species composition (Grazing → Species

composition → Function) (Bagchi et al., 2017; Naidu et al., 2022; Roy and Bagchi, 2022). c)

Background edaphic variables can shape up the microbial communities (Edaphic (pH, sand,

moisture) → Species composition) (Girvan et al., 2003; Fierer and Jackson, 2006; Bagchi et

al., 2017). d) Variation in background edaphic variables can alter function directly (Edaphic

→ Function), or indirectly through changes in species composition (Edaphic → Species

composition → Function) (Girvan et al., 2003; Delgado-Baquerizo et al., 2016b; Bagchi et

al., 2017). e) both microbial community and function can vary temporally (Season → Species

composition, Season → Function) (Bardgett et al., 2005; Frossard et al., 2012; Delgado-

Baquerizo et al., 2013; Bagchi et al., 2017). f) Grazing can also influence soil moisture

(Grazing → Moisture) (Bagchi et al., 2017), that can in-turn alter pH (Moisture → pH). g)

Sand content can also alter moisture (Sand → Moisture) and pH (Sand → pH). We assessed

goodness of fit between the model and data with χ^2 test, comparative fit index (CFI), and

Standardized Root Mean Square Error of Approximation (RMSEA). We accepted the model

$P(\chi^2) > 0.05$, CFI ~ 1, and RMSEA < 0.05. We implemented paths for dbSEM using 'lavaan'

library in R (Rosseel, 2012).

Results

Microbial communities and functions –

We recorded 1937 unique OTUs across 47 known phyla and 7650894 reads.

Actinobacteriota, Acidobacteriota, Planctomycetota, Proteobacteria and Verrucomicrobiota

were most common (Fig. 1, A.3.). Species richness of OTUs ranged between 494 and 866 per

sample (mean 713±80 SD). Microbial activity was highest for carboxylic acids such as α-

Ketoglutaric acid, Malic acid, Citric acid, Succinic acid, and Malonic acid, Similarly,

functional richness (i.e., number of substrates metabolized) ranged between 6 and 30 (mean

21±7 SD, Fig. 1, A.3.).

Seasonality and grazer-effects on microbial species and functions –

Variation in species (OTUs) richness was not explained by season (Fig. 2a;

$F_{2,45}=0.02$, $P=0.98$), and by grazing (Fig. 2a; $F_{1,45}=1.61$, $P=0.21$). Microbial functional

richness varied with season (Fig. 2b; $F_{2,45}=38.96$, $P<0.001$) but did not vary with grazing

(Fig. 2b; $F_{1,45}=0.01$, $P=0.98$). Similarly, microbial functional dispersion varied with season

(Fig. 2c; $F_{2,45}=48.89$, $P<0.001$) but did not vary with grazing (Fig. 2c; $F_{1,45}=0.01$, $P=0.92$).

Multifunction also varied with season (Fig. 2d; $F_{2,45}=10.94$, $P<0.001$), but not with grazing

(Fig. 2d; $F_{1,45}=0.07$, $P=0.79$).

In multivariate space summarized with PCoA ordination on Bray-Curtis dissimilarity

matrix, microbial species composition varied with season (Fig. 3; $F_{2,54}=4.70$, $P=0.001$), but

not with grazing (Fig. 3; $F_{1,54}=1.33$, $P=0.206$). functional composition also varied with

season (Fig. 3; $F_{2,54}=4.83$, $P=0.001$), but not with grazing (Fig. 3; $F_{1,54}=0.87$, $P=0.555$).

Relationship between species composition and functions –

The relationship between species composition and microbial functions was visualized using

kriging (Fig. 4). Kriged maps of microbial functions indicates a dependence on microbial

species composition where different community composition support different magnitude of

functions.

Relationship between species composition and microbial functions were further

assessed using dbSEM. Overall, dbSEM for functional richness (Fig. 5), functional dispersion

(Fig. 5), and multifunction (Fig. 5) indicated good match between the data and hypothesized

paths (SRMR=0.005, RMSEA<0.001, P-value (RMSEA)=0.982, CFI>0.999, $\chi^2=1.34$, P-

value (χ^2)=0.512). Path coefficients suggested a significant positive relationship between

species composition and functional richness (Species composition → Functional richness;

path coefficient=0.08, P=0.06; Fig. 5), and with functional dispersion (Species composition

→ Dispersion; path coefficient=0.08, P=0.10; Fig. 5), but not with multifunction (Species

composition → Multifunction; path coefficient=-0.06, P=0.13; Fig. 5). Sand content had a

positive effect on species composition (Sand → Species composition; path coefficient=0.14,

P=0.001; Fig. 5), and a negative effect on multifunction (Sand → Multifunction; path

coefficient=-0.06, P=0.04; Fig. 5). Seasonality had positive relationship with species

composition (Season → Species composition; path coefficient=0.14, P=0.001; Fig. 5), and

soil moisture (Season → Moisture; path coefficient=0.29, P<0.001; Fig. 5), such that as the

season progresses from early to mid to late, species composition and soil moisture increases.

Moisture in turn had a positive effect on species composition (Moisture → Species

composition; path coefficient=0.12, P=0.02; Fig. 5). In this way seasonality had a direct

(Season → Species composition → Function), as well as indirect effect through moisture

(Season → Moisture → Species composition → Function), on the species composition-

function relationship. Grazing did not influence this relationship directly (Grazing → Species

composition → Function), or indirectly through edaphic factors (Grazing → Edaphic →

Species composition → Function).

Permutation based dbSEM for individual functions indicated good match between the

data and hypothesized paths (SRMR=0.005, RMSEA<0.001, P-value (RMSEA)=0.982,

CFI>0.999, χ^2=1.34, P-value (χ^2)=0.512). Significant species composition and function

relationship was found for 6 out of 30 functions (Species composition → Function, Fig. A.1).

The relationship was positive for 5 out of 6 functions, and negative for only Malic acid. For

all these 6 functions season had a significant effect on the relationship (Season → Species

composition → Function, $P<0.05$). Season also had an indirect effect on 5 out of 6 significant

species composition-function relationship through soil moisture (Season → Moisture →

Species composition → Function, $P<0.05$). There was significant temporal variation in 28 out

of 30 functions (Season → Function, $P<0.05$). Soil moisture and sand content also had

positive effect on the relationship for all 6 functions. Soil moisture had positive effect on 17

out of 30 functions and negative effect on 3 out of 30 functions (Moisture → Function,

$P<0.05$). Sand content had positive effect on 3 out of 30 functions and negative effect on only

1 function (Sand → Function, $P<0.05$). While grazing had a negative effect on 4 out of 30

functions (Grazing → Function, $P<0.05$), but grazing did not influence species composition-

function relationship directly (Grazing → Species composition → Function, $P>0.05$), or

indirectly through edaphic factors (Grazing → Edaphic → Species composition → Function,

$P>0.05$).

Discussion

Diversity-function relationship is known for producers and consumers but remains uncertain

for microbial decomposers and decomposition process. Here we investigate whether, or not,

microbial taxonomic diversity (structure) is related to the microbial decomposition process

(function) in soil. In the Trans-Himalayan drylands, we find three key results: (i) soil

microbial communities and functions are more variable across time than over space (Fig. 1, 2,

3), (ii) microbial species composition affects functional richness, but not functional

composition and multifunction (Fig. 4, 5), and (iii) large mammalian herbivores did not affect

the diversity-function relationship for microbes, whereas background edaphic variables and

season does affect the relationship (Fig. 5).

Taxonomy~Function relationship

A general paradigm in ecology is that taxonomy determines functions, such that greater

taxonomic diversity will favor functional diversity. This paradigm is supported for producers

and consumers, at both local (α-diversity) as well as landscape levels (β-diversity) (Loreau et

al., 2001; Hooper et al., 2005; Reich et al., 2012; Hautier et al., 2018), and can influence the

provisioning of ecosystem services. Unlike producers and consumers, soil microbial

decomposers are hyper-diverse. As a case in point, we recorded more than 700 OTUs on

average per sample, and 1937 OTUs overall across 7.6 million reads. Therefore, and

understandably, the data did not show any support for the relationship between species

richness and functions (Fig. A.5). For instance, where local richness is below 200 OTUs per

sample, there is evidence of coupling and therefore a lack of redundancy (Delgado-Baquerizo

et al., 2016a). But, further along at 700 OTUs per sample (Fig. 3-4) microbial diversity can

appear de-coupled from functions (Peter et al., 2011; Louca et al., 2016; Chen et al., 2022).

Thus, not surprisingly, different studies have encountered coupled as well as de-coupled

relationships between microbial species richness and functions (Peter et al., 2011; Delgado-

Baquerizo et al., 2016a; Louca et al., 2016; Chen et al., 2022), and our data emphasize how it

is important to accommodate species composition (Hooper et al., 2005).

Our results show that taxonomically dissimilar community can perform more

functions (richness) but has no effect on the magnitude (composition and multifunction) of

the functions. Although this result seems counterintuitive but is not trivial. The ability to

perform any function is tightly linked with the identity and relative abundance of each

species in the community, but which functions will operate and at what rate depends on the

environment. The interdependency between taxonomic composition and functions seems to

challenge "functional redundancy" or "functional equivalence" hypothesis. It proposes that

changes in microbial community composition exercise only proximate controls on functions

whereas the current environment ultimately determines functions. Alternatively, "functional

dissimilarity" hypothesis proposes that not 'only' environment, but a 'combination' of

environment and microbial community composition ultimately determine microbial functions

(Strickland et al., 2009). Indeed, we find that while seasonality and edaphic factors (moisture

and texture) together with community composition exerts strong control over microbial

functions.

Seasonality

We find temporal variation in microbial community structure and functions to be

more pronounced than spatial heterogeneity (Fig 2, 3, 5). This is consistent with other studies

where fluctuations in resource availability explained why microbes respond to time more

than space (Frossard et al., 2012; Bagchi et al., 2017; Roy and Bagchi, 2022; Roy et al.,

2023). Vegetation at our study site follows a unimodal distribution in the growing season and

the underlying temperature regime can cause a seasonal variation in resource availability for

microbes to metabolize (Millard and Singh, 2010; Roy et al., 2023). Such a stark difference

between temporal and spatial responsiveness of microbes lends further support to adage

"everything is everywhere but the environment selects" (Bass-Becking, 1934), and

encourages further discussion on the temporal aspect of soil microbes in soil and its

functional implications. Our study also highlights why biodiversity-ecosystem function

(BEF) studies, particularly for microbes, should incorporate temporal dynamics (days,

months, years) rather than restricting to a snapshot in time.

Edaphic factors

We find a consistent effect of soil moisture and texture on the decomposer diversity-

function relationship (Fig. 5). Moisture and texture can play an important role in habitat-

filtering (Bass-Becking, 1934) as they promote functions through taxonomic composition

(Fig. 5), resulting in the loss of functional redundancies among decomposers. On average, the

Trans-Himalayan drylands are getting wetter over decadal time-scales while experiencing

increased inter-annual variability (Murthy and Bagchi, 2018; Hu and Zhou, 2021). Under

such ongoing and projected precipitation regimes, the stability of ecosystem functions will

depend on how changing moisture availability affects coupling of taxonomy and function.

Further, strong association of texture with the functions can exacerbate the negative effects of

habitat filtering by altered precipitation regimes on decomposition. Variation in soil texture

had positive effect functional richness (Fig. 5), that invariably also reduces heterogeneity in

multifunction across the landscape (Fig. 5), leading to effective homogenization. Soil texture

can cause homogenization in decomposition process by altering soil water availability, pore

size distribution, surface area, and ultimately nutrient availability that in turn opens new

niches (increases access to newer substrates) for microbes to proliferate and reduces

heterogeneity in decomposition. While the effect of moisture and texture (Fig. 4) is consistent

with other studies across ecosystems (Girvan et al., 2003; Delgado-Baquerizo et al., 2016c),

unlike other studies (Fierer and Jackson, 2006; Sinsabaugh et al., 2008) we find no effect of

soil pH. This could be due to the invariability in soil pH across the landscape (space), and

season (time), but it can vary between different landscapes and ecosystems.

Large mammalian herbivores

We find that the relationship between microbial taxonomy and function is robust to

the presence or absence of large mammalian herbivores. This observation is in line with

patterns seen for producers and their production systems where herbivore removal did not

impact producer diversity and ecosystem stability relationship (Kohli et al., 2019). However,

contrary to the direct effect of consumers on producers, the mechanisms operating for

decomposers might be different (Bagchi et al., 2017; Naidu et al., 2022; Roy and Bagchi,

2022). There could be multiple interdependent paths that could have led to this observation:

1. Balance between direct and indirect herbivore effects: Net effect, or a lack thereof, on

440   the microbial decomposer diversity-function relationship can be a balance between

441   the direct and indirect effects of large-mammalian herbivores (Bagchi and Ritchie,

442   2010a; Cherif and Loreau, 2013; Naidu et al., 2022; Roy and Bagchi, 2022; Roy et al.,

443   2023). Direct effect of herbivores, through the quantity of nutrient input, on

444   decomposers is exerted by consumption of plant productions that diverts nutrient

445   away from the soil towards secondary production (Bagchi and Ritchie, 2010b). On the

446   other hand, indirect effect of herbivores, through the quality of nutrient input, on

447   decomposers is exerted by converting plant materials into dung and urine, and altering

448   plant community composition, that change the proportion of labile to recalcitrant

449   substrates (Bagchi et al., 2017; Naidu et al., 2022; Roy and Bagchi, 2022).

450   Equilibrium in the relative strengths of the direct and indirect influence of herbivores

451   can therefore manifest as no net effect on the decomposer diversity-function

452   relationship (Hooper et al., 2000). This can also explain the variability in microbial

453   responses to herbivores, with neutral, positive, negative effects in different

454   ecosystems (Tanentzap and Coomes, 2012; McSherry and Ritchie, 2013; Wang and

455   Tang, 2019; Eldridge et al., 2020).

456   2. Historical and evolutionary adaptations: The tight coupling between microbial

457   diversity and function, could have established before the start of the decadal-scale

458   grazer-exclusion experiment due to competition-cooperation interactions (Morrissey

459   et al., 2019; Lechón-Alonso et al., 2021). Once established, the interactions among

460   microbial decomposers may persist for long periods (Morrissey et al., 2019; López

Zieher et al., 2020), and can only respond to the prevailing edaphic constraints

(moisture, texture) (Fig. 5).

3. Effect on other groups of microbial decomposers: We recognize that we cannot

exclude the effect of herbivores on fungi or archaea that are also responsible for the

decomposition process. Indeed, previous studies have shown that grazers shift

fungal:bacterial ratio towards dominance by fungi that ultimately regulates soil-C

dynamics (Bagchi et al., 2017). Therefore, this would need further research where we

incorporate the fungal taxonomic information as well. Nonetheless, our study

suggests that spatiotemporal variability in the soil microclimate had a stronger effect

on the diversity-function relationship of microbial decomposers, than herbivory *per se*

(Hooper et al., 2000). This is in line with the studies that suggests high altitude

ecosystems to be more sensitive to variability in climate parameters than ecosystems

at lower elevations (Ohmura, 2012; Lamprecht et al., 2018; Li et al., 2018).

Overall, we find grazer-effect microbial functions are mediated through changes in

microbial community composition. Soil moisture controls most dryland ecosystem processes

(Millard and Singh, 2010; Delgado-Baquerizo et al., 2013; Maestre et al., 2016), and future

changes in soil moisture availability in drylands (Schlaepfer et al., 2017; Konapala et al.,

2020), particularly in the Trans-Himalaya and Central Asia's high-elevation drylands, might

challenge their conservation and sustainable management for ecosystem functions and

services.

Acknowledgements

SR was supported by a graduate fellowship from CSIR-India in IISc, and Discipline Hopping

for Discovery Science grant by NERC-UK in UEA; SB was supported by MHRD-India.

Funding support for fieldwork and lab-analyses were obtained from DST-SERB, DST-FIST,

DBT-IISc. Govt. of Himachal Pradesh and state departments have supported our research in

Spiti. We received generous support from Dorje Chhering, Dorje Chhewang, and Deepak

Shetti, in field and in lab.

References

Allison, S.D., Martiny, J.B.H., 2008. Resistance, resilience, and redundancy in microbial
communities. Proceedings of the National Academy of Sciences 105, 11512–11519.
doi:10.1073/pnas.0801925105

Anderson, M.J., Ellingsen, K.E., McArdle, B.H., 2006. Multivariate dispersion as a measure
of beta diversity. Ecology Letters 9, 683–693. doi:10.1111/j.1461-0248.2006.00926.x

Bagchi, S., Ritchie, M.E., 2010a. Introduced grazers can restrict potential soil carbon
sequestration through impacts on plant community composition. Ecology Letters 13,
959–968. doi:10.1111/j.1461-0248.2010.01486.x

Bagchi, S., Ritchie, M.E., 2010b. Herbivore effects on above- and belowground plant
production and soil nitrogen availability in the Trans-Himalayan shrub-steppes.
Oecologia 164, 1075–1082. doi:10.1007/s00442-010-1690-5

Bagchi, S., Roy, S., Maitra, A., Sran, R.S., 2017. Herbivores suppress soil microbes to
influence carbon sequestration in the grazing ecosystem of the Trans-Himalaya.
Agriculture, Ecosystems and Environment 239, 199–206.
doi:10.1016/j.agee.2017.01.033

Bardgett, R., Bowman, W., Kaufmann, R., Schmidt, S., 2005. A temporal approach to linking
aboveground and belowground ecology. Trends in Ecology & Evolution 20, 634–641.
doi:10.1016/j.tree.2005.08.005

Bardgett, R.D., Jones, A.C., Jones, D.L., Kemmitt, S.J., Cook, R., Hobbs, P.J., 2001. Soil
microbial community patterns related to the history and intensity of grazing in sub-
montane ecosystems. Soil Biology and Biochemistry 33, 1653–1664.
doi:10.1016/S0038-0717(01)00086-4

Bardgett, R.D., Wardle, D.A., 2003. Herbivore-mediated linkages between aboveground and
belowground communities. Ecology 84, 2258–2268. doi:10.1890/02-0274

Bass-Becking, L.G.M., 1934. Geobiologie of inleiding tot de milieukunde. Den Haag: W.P.
Van Stockum & Zoon.

Bell, C.W., Acosta-Martinez, V., McIntyre, N.E., Cox, S., Tissue, D.T., Zak, J.C., 2009.
Linking microbial community structure and function to seasonal differences in soil
moisture and temperature in a Chihuahuan Desert grassland. Microbial Ecology 58,
827–842. doi:10.1007/s00248-009-9529-5

Bengtsson, J., 1998. Which species? What kind of diversity? Which ecosystem function?
Some problems in studies of relations between biodiversity and ecosystem function.
Applied Soil Ecology 10, 191–199. doi:10.1016/S0929-1393(98)00120-6

Cardinale, B.J., Matulich, K.L., Hooper, D.U., Byrnes, J.E., Duffy, E., Gamfeldt, L.,
Balvanera, P., O'Connor, M.I., Gonzalez, A., 2011. The functional role of producer
diversity in ecosystems. American Journal of Botany 98, 572–592.
doi:10.3732/ajb.1000364

Cardinale, B.J., Srivastava, D.S., Emmett Duffy, J., Wright, J.P., Downing, A.L., Sankaran,
M., Jouseau, C., 2006. Effects of biodiversity on the functioning of trophic groups and
ecosystems. Nature 443, 989–992. doi:10.1038/nature05202

Chen, W., Wang, J., Chen, X., Meng, Z., Xu, R., Duoji, D., Zhang, J., He, J., Wang, Z.,
Chen, J., Liu, K., Hu, T., Zhang, Y., 2022. Soil microbial network complexity predicts
ecosystem function along elevation gradients on the Tibetan Plateau. Soil Biology and
Biochemistry 172, 108766. doi:10.1016/j.soilbio.2022.108766

Cherif, M., Loreau, M., 2013. Plant-herbivore-decomposer stoichiometric mismatches and
nutrient cycling in ecosystems. Proceedings of the Royal Society B: Biological Sciences
280, 20122453. doi:10.1098/rspb.2012.2453

Coleman, D.C., Whitman, W.B., 2005. Linking species richness, biodiversity and ecosystem function in soil systems, in: Pedobiologia. Elsevier GmbH, pp. 479–497. doi:10.1016/j.pedobi.2005.05.006

Dale, M.R.T., Fortin, M.-J., 2014. Spatial Analysis: A Guide For Ecologists. Cambridge University Press.

Degens, B.P., 1998. Microbial functional diversity can be influenced by the addition of simple organic substrates to soil. Soil Biology and Biochemistry 30, 1981–1988. doi:10.1016/S0038-0717(98)00070-4

Degens, B.P., Harris, J.A., 1997. Development of a physiological approach to measuring the catabolic diversity of soil microbial communities. Soil Biology and Biochemistry 29, 1309–1320. doi:10.1016/S0038-0717(97)00076-X

Delgado-Baquerizo, M., Eldridge, D.J., Ochoa, V., Gozalo, B., Singh, B.K., Maestre, F.T., 2017. Soil microbial communities drive the resistance of ecosystem multifunctionality to global change in drylands across the globe. Ecology Letters 20, 1295–1305. doi:10.1111/ele.12826

Delgado-Baquerizo, M., Giaramida, L., Reich, P.B., Khachane, A.N., Hamonts, K., Edwards, C., Lawton, L.A., Singh, B.K., 2016a. Lack of functional redundancy in the relationship between microbial diversity and ecosystem functioning. Journal of Ecology 104, 936–946. doi:10.1111/1365-2745.12585

Delgado-Baquerizo, M., Grinyer, J., Reich, P.B., Singh, B.K., 2016b. Relative importance of soil properties and microbial community for soil functionality: insights from a microbial swap experiment. Functional Ecology 30, 1862–1873. doi:10.1111/1365-2435.12674

Delgado-Baquerizo, M., Maestre, F.T., Gallardo, A., Bowker, M.A., Wallenstein, M.D., Quero, J.L., Ochoa, V., Gozalo, B., García-Gómez, M., Soliveres, S., García-Palacios, P., Berdugo, M., Valencia, E., Escolar, C., Arredondo, T., Barraza-Zepeda, C., Bran, D., Carreira, J.A., Chaieb, M., Conceicao, A.A., Derak, M., Eldridge, D.J., Escudero, A., Espinosa, C.I., Gaitán, J., Gatica, M.G., Gómez-González, S., Guzman, E., Gutiérrez, J.R., Florentino, A., Hepper, E., Hernández, R.M., Huber-Sannwald, E., Jankju, M., Liu, J., Mau, R.L., Miriti, M., Monerris, J., Naseri, K., Noumi, Z., Polo, V., Prina, A., Pucheta, E., Ramírez, E., Ramírez-Collantes, D.A., Romao, R., Tighe, M., Torres, D., Torres-Díaz, C., D. Ungar, E., Val, J., Wamiti, W., Wang, D., Zaady, E., 2013. Decoupling of soil nutrient cycles as a function of aridity in global drylands. Nature 502, 672–676. doi:10.1038/nature12670

Delgado-Baquerizo, M., Maestre, F.T., Reich, P.B., Jeffries, T.C., Gaitan, J.J., Encinar, D., Berdugo, M., Campbell, C.D., Singh, B.K., 2016c. Microbial diversity drives multifunctionality in terrestrial ecosystems. Nature Communications 7, 1–8. doi:10.1038/ncomms10541

Duffy, J.E., 2002. Biodiversity and ecosystem function: the consumer connection. Oikos 99, 201–219. doi:10.1034/j.1600-0706.2002.990201.x

Duffy, J.E., Cardinale, B.J., France, K.E., McIntyre, P.B., Thébault, E., Loreau, M., 2007. The functional role of biodiversity in ecosystems: incorporating trophic complexity. Ecology Letters 10, 522–538. doi:10.1111/j.1461-0248.2007.01037.x

Edgar, R.C., Haas, B.J., Clemente, J.C., Quince, C., Knight, R., 2011. UCHIME improves sensitivity and speed of chimera detection. Bioinformatics 27, 2194–2200. doi:10.1093/bioinformatics/btr381

Eldridge, D.J., Travers, S.K., Val, J., Wang, J.T., Liu, H., Singh, B.K., Delgado-Baquerizo, M., 2020. Grazing Regulates the Spatial Heterogeneity of Soil Microbial Communities Within Ecological Networks. Ecosystems 23, 932–942. doi:10.1007/s10021-019-00448-9

Fierer, N., Colman, B.P., Schimel, J.P., Jackson, R.B., 2006. Predicting the temperature
dependence of microbial respiration in soil: A continental-scale analysis. Global
Biogeochemical Cycles 20, GB3026. doi:10.1029/2005GB002644
Fierer, N., Jackson, R.B., 2006. The diversity and biogeography of soil bacterial
communities. Proceedings of the National Academy of Sciences 103, 626–631.
doi:10.1073/pnas.0507535103
Fierer, N., Ladau, J., Clemente, J.C., Leff, J.W., Owens, S.M., Pollard, K.S., Knight, R.,
Gilbert, J.A., McCulley, R.L., 2013. Reconstructing the microbial diversity and function
of pre-agricultural tallgrass prairie soils in the United States. Science 342, 621–624.
doi:10.1126/SCIENCE.1243768/SUPPL_FILE/FIERER.SM.PDF
Fourtune, L., Prunier, J.G., Paz-Vinas, I., Loot, G., Veyssière, C., Blanchet, S., 2018.
Inferring causalities in landscape genetics: An extension of wright's causal modeling to
distance matrices. American Naturalist 191, 491–508. doi:10.1086/696233
Frank, D. a., McNaughton, S.J., Tracy, B.F., 1998. The Ecology of the Earth's Grazing
Ecosystems. BioScience 48, 513–521. doi:10.2307/1313313
Frossard, A., Gerull, L., Mutz, M., Gessner, M.O., 2012. Disconnect of microbial structure
and function: enzyme activities and bacterial communities in nascent stream corridors.
The ISME Journal 6, 680–691. doi:10.1038/ismej.2011.134
Galand, P.E., Pereira, O., Hochart, C., Auguet, J.C., Debroas, D., 2018. A strong link
between marine microbial community composition and function challenges the idea of
functional redundancy. The ISME Journal 12, 2470–2478. doi:10.1038/s41396-018-
0158-1
Gessner, M.O., Swan, C.M., Dang, C.K., McKie, B.G., Bardgett, R.D., Wall, D.H.,
Hättenschwiler, S., 2010. Diversity meets decomposition. Trends in Ecology &
Evolution 25, 372–380. doi:10.1016/J.TREE.2010.01.010
Girvan, M.S., Bullimore, J., Pretty, J.N., Osborn, A.M., Ball, A.S., 2003. Soil type is the
primary determinant of the composition of the total and active bacterial communities in
arable soils. Applied and Environmental Microbiology 69, 1800–1809.
doi:10.1128/AEM.69.3.1800-1809.2003
Gossner, M.M., Lewinsohn, T.M., Kahl, T., Grassein, F., Boch, S., Prati, D., Birkhofer, K.,
Renner, S.C., Sikorski, J., Wubet, T., Arndt, H., Baumgartner, V., Blaser, S., Blüthgen,
N., Börschig, C., Buscot, F., Diekötter, T., Jorge, L.R., Jung, K., Keyel, A.C., Klein, A.-
M., Klemmer, S., Krauss, J., Lange, M., Müller, J., Overmann, J., Pašalić, E., Penone,
C., Perović, D.J., Purschke, O., Schall, P., Socher, S.A., Sonnemann, I., Tschapka, M.,
Tscharntke, T., Türke, M., Venter, P.C., Weiner, C.N., Werner, M., Wolters, V., Wurst,
S., Westphal, C., Fischer, M., Weisser, W.W., Allan, E., 2016. Land-use intensification
causes multitrophic homogenization of grassland communities. Nature 540, 266–269.
doi:10.1038/nature20575
Grman, E., Zirbel, C.R., Bassett, T., Brudvig, L.A., 2018. Ecosystem multifunctionality
increases with beta diversity in restored prairies. Oecologia 188, 837–848.
doi:10.1007/s00442-018-4248-6
Hautier, Y., Isbell, F., Borer, E.T., Seabloom, E.W., Harpole, W.S., Lind, E.M., MacDougall,
A.S., Stevens, C.J., Adler, P.B., Alberti, J., Bakker, J.D., Brudvig, L.A., Buckley, Y.M.,
Cadotte, M., Caldeira, M.C., Chaneton, E.J., Chu, C., Daleo, P., Dickman, C.R., Dwyer,
J.M., Eskelinen, A., Fay, P.A., Firn, J., Hagenah, N., Hillebrand, H., Iribarne, O.,
Kirkman, K.P., Knops, J.M.H., la Pierre, K.J., McCulley, R.L., Morgan, J.W., Pärtel,
M., Pascual, J., Price, J.N., Prober, S.M., Risch, A.C., Sankaran, M., Schuetz, M.,
Standish, R.J., Virtanen, R., Wardle, G.M., Yahdjian, L., Hector, A., 2018. Local loss
and spatial homogenization of plant diversity reduce ecosystem multifunctionality.
Nature Ecology and Evolution 2, 50–56. doi:10.1038/s41559-017-0395-0

Hector, A., Bagchi, R., 2007. Biodiversity and ecosystem multifunctionality. Nature 448, 188–190. doi:10.1038/nature05947

Hooper, D.U., Bignell, D.E., Brown, V.K., Brussard, L., Dangerfield, J.M., Wall, D.H., Wardle, D.A., Coleman, D.C., Giller, K.E., Lavelle, P., Van Der Putten, W.H., De Ruiter, P.C., Rusek, J., Silver, W.L., Tiedje, J.M., Wolters, V., 2000. Interactions between Aboveground and Belowground Biodiversity in Terrestrial Ecosystems: Patterns, Mechanisms, and Feedbacks. BioScience 50, 1049–1061. doi:10.1641/0006-3568(2000)050[1049:IBAABB]2.0.CO;2

Hooper, D.U., Chapin, F.S., Ewel, J.J., Hector, A., Inchausti, P., Lavorel, S., Lawton, J.H., Lodge, D.M., Loreau, M., Naeem, S., Schmid, B., Setälä, H., Symstad, A.J., Vandermeer, J., Wardle, D.A., 2005. Effects of biodiversity on ecosystem functioning: A consensus of current knowledge. Ecological Monographs 75, 3–35. doi:10.1890/04-0922

Hu, S., Zhou, T., 2021. Skillful prediction of summer rainfall in the Tibetan Plateau on multiyear time scales. Science Advances 7, eabf9395. doi:10.1126/sciadv.abf9395

Jabot, F., Laroche, F., Massol, F., Arthaud, F., Crabot, J., Dubart, M., Blanchet, S., Munoz, F., David, P., Datry, T., 2020. Assessing metacommunity processes through signatures in spatiotemporal turnover of community composition. Ecology Letters. doi:10.1111/ele.13523

Justiniano {Ribeiro Jr}, P., Diggle, P., Christensen, O., Schlather, M., Bivand, R., Ripley, B., 2022. geoR: Analysis of Geostatistical Data. R Package Version 1.9-2.

Klindworth, A., Pruesse, E., Schweer, T., Peplies, J., Quast, C., Horn, M., Glöckner, F.O., 2013. Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-generation sequencing-based diversity studies. Nucleic Acids Research 41, e1–e1. doi:10.1093/nar/gks808

Koerner, S.E., Smith, M.D., Burkepile, D.E., Hanan, N.P., Avolio, M.L., Collins, S.L., Knapp, A.K., Lemoine, N.P., Forrestel, E.J., Eby, S., Thompson, D.I., Aguado-Santacruz, G.A., Anderson, J.P., Anderson, T.M., Angassa, A., Bagchi, S., Bakker, E.S., Bastin, G., Baur, L.E., Beard, K.H., Beever, E.A., Bohlen, P.J., Boughton, E.H., Canestro, D., Cesa, A., Chaneton, E., Cheng, J., D'Antonio, C.M., Deleglise, C., Dembélé, F., Dorrough, J., Eldridge, D.J., Fernandez-Going, B., Fernández-Lugo, S., Fraser, L.H., Freedman, B., García-Salgado, G., Goheen, J.R., Guo, L., Husheer, S., Karembé, M., Knops, J.M.H., Kraaij, T., Kulmatiski, A., Kytöviita, M.M., Lezama, F., Loucougaray, G., Loydi, A., Milchunas, D.G., Milton, S.J., Morgan, J.W., Moxham, C., Nehring, K.C., Olff, H., Palmer, T.M., Rebollo, S., Riginos, C., Risch, A.C., Rueda, M., Sankaran, M., Sasaki, T., Schoenecker, K.A., Schultz, N.L., Schütz, M., Schwabe, A., Siebert, F., Smit, C., Stahlheber, K.A., Storm, C., Strong, D.J., Su, J., Tiruvaimozhi, Y. V., Tyler, C., Val, J., Vandegehuchte, M.L., Veblen, K.E., Vermeire, L.T., Ward, D., Wu, J., Young, T.P., Yu, Q., Zelikova, T.J., 2018. Change in dominance determines herbivore effects on plant biodiversity. Nature Ecology and Evolution 2, 1925–1932. doi:10.1038/s41559-018-0696-y

Kohli, M., Borer, E.T., Kinkel, L., Seabloom, E.W., 2019. Stability of grassland production is robust to changes in the consumer food web. Ecology Letters 22, 707–716. doi:10.1111/ele.13232

Konapala, G., Mishra, A.K., Wada, Y., Mann, M.E., 2020. Climate change will affect global water availability through compounding changes in seasonal precipitation and evaporation. Nature Communications 11, 1–10. doi:10.1038/s41467-020-16757-w

Lamprecht, A., Semenchuk, P.R., Steinbauer, K., Winkler, M., Pauli, H., 2018. Climate change leads to accelerated transformation of high-elevation vegetation in the central Alps. New Phytologist 220, 447–459. doi:10.1111/NPH.15290

Lechón-Alonso, P., Clegg, T., Cook, J., Smith, T.P., Pawar, S., 2021. The role of competition
versus cooperation in microbial community coalescence. PLOS Computational Biology
17, e1009584. doi:10.1371/journal.pcbi.1009584
Legendre, P., Legendre, L., 2012. Numerical Ecology, 3rd ed. Elsevier. doi:10.2307/2530779
Li, D., Wu, S., Liu, L., Zhang, Y., Li, S., 2018. Vulnerability of the global terrestrial
ecosystems to climate change. Global Change Biology 24, 4095–4106.
doi:10.1111/gcb.14327
Li, Y., Wang, X., Chen, Y., Gong, X., Yao, C., Cao, W., Lian, J., 2023. Application of
predictor variables to support regression kriging for the spatial distribution of soil
organic carbon stocks in native temperate grasslands. Journal of Soils and Sediments 23,
700–717. doi:10.1007/s11368-022-03370-1
López Zieher, X.M., Vivanco, L., Yahdjian, L., 2020. Soil bacterial communities remain
altered after 30 years of agriculture abandonment in Pampa grasslands. Oecologia 193,
959–968. doi:10.1007/s00442-020-04736-3
Loreau, M., Naeem, S., Inchausti, P., Bengtsson, J., Grime, J.P., Hector, A., Hooper, D.U.,
Huston, M.A., Raffaelli, D., Schmid, B., Tilman, D., Wardle, D.A., 2001. Biodiversity
and Ecosystem Functioning: Current Knowledge and Future Challenges. Science 294,
804–808. doi:10.1126/science.1064088
Louca, S., Parfrey, L.W., Doebeli, M., 2016. Decoupling function and taxonomy in the global
ocean microbiome. Science 353, 1272–1277. doi:10.1126/science.aaf4507
Maestre, F.T., Eldridge, D.J., Soliveres, S., Kéfi, S., Delgado-Baquerizo, M., Bowker, M.A.,
García-Palacios, P., Gaitán, J., Gallardo, A., Lázaro, R., Berdugo, M., 2016. Structure
and Functioning of Dryland Ecosystems in a Changing World. Annual Review of
Ecology, Evolution, and Systematics 47, 215–237. doi:10.1146/annurev-ecolsys-
121415-032311
McNaughton, S.J., 1976. Serengeti migratory wildebeest: Facilitation of energy flow by
grazing. Science 191, 92–94. doi:10.1126/science.191.4222.92
McSherry, M.E., Ritchie, M.E., 2013. Effects of grazing on grassland soil carbon: A global
review. Global Change Biology 19, 1347–1357. doi:10.1111/gcb.12144
Milchunas, D.G., Lauenroth, W.K., 1993. Quantitative effects of grazing on vegetation and
soils over a global range of environments. Ecological Monographs 63, 327–366.
doi:10.2307/2937150
Millard, P., Singh, B.K., 2010. Does grassland vegetation drive soil microbial diversity?
Nutrient Cycling in Agroecosystems 88, 147–158. doi:10.1007/s10705-009-9314-3
Morrissey, E.M., Mau, R.L., Hayer, M., Liu, X.J.A., Schwartz, E., Dijkstra, P., Koch, B.J.,
Allen, K., Blazewicz, S.J., Hofmockel, K., Pett-Ridge, J., Hungate, B.A., 2019.
Evolutionary history constrains microbial traits across environmental variation. Nature
Ecology and Evolution 3, 1064–1069. doi:10.1038/s41559-019-0918-y
Murthy, K., Bagchi, S., 2018. Spatial patterns of long-term vegetation greening and browning
are consistent across multiple scales: Implications for monitoring land degradation. Land
Degradation & Development 29, 2485–2495. doi:10.1002/ldr.3019
Naidu, D.G.T., Roy, S., Bagchi, S., 2022. Loss of grazing by large mammalian herbivores
can destabilize the soil carbon pool. Proceedings of the National Academy of Sciences
119, 1–7. doi:10.1073/pnas.2211317119
Ohmura, A., 2012. Enhanced temperature variability in high-altitude climate change.
Theoretical and Applied Climatology 110, 499–508. doi:10.1007/s00704-012-0687-x
Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., Mcglinn, D., Minchin,
P.R., O'Hara, R.B., Simpson, G.L., Solymos, P., Stevens, M.H.H., Szoecs, E., Wagner,
H., 2019. vegan: community ecology package. R package version 2.5-6.

Olff, H., Ritchie, M.E., 1998. Effects of herbivores on grassland plant diversity. Trends in
Ecology & Evolution 13, 261–265. doi:10.1016/S0169-5347(98)01364-0
Peter, H., Beier, S., Bertilsson, S., Lindström, E.S., Langenheder, S., Tranvik, L.J., 2011.
Function-specific response to depletion of microbial diversity. ISME Journal 5, 351–
361. doi:10.1038/ismej.2010.119
Pinheiro, J., Bates, D., 2000. Mixed-Effects Models in S and S-PLUS, 1st ed, Mixed-Effects
Models in S and S-PLUS. Springer, New York, NY. doi:10.1007/b98882
Purahong, W., Schloter, M., Pecyna, M.J., Kapturska, D., Däumlich, V., Mital, S., Buscot, F.,
Hofrichter, M., Gutknecht, J.L.M., Krüger, D., 2014. Uncoupling of microbial
community structure and function in decomposing litter across beech forest ecosystems
in Central Europe. Scientific Reports 4, 1–7. doi:10.1038/srep07014
Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., Glöckner,
F.O., 2013. The SILVA ribosomal RNA gene database project: Improved data
processing and web-based tools. Nucleic Acids Research 41, D590–D596.
doi:10.1093/nar/gks1219
Reich, P.B., Tilman, D., Isbell, F., Mueller, K., Hobbie, S.E., Flynn, D.F.B., Eisenhauer, N.,
2012. Impacts of biodiversity loss escalate through time as redundancy fades. Science
336, 589–592. doi:10.1126/science.1217909
Robertson, G.P., Coleman, D.C., Bledsoe, C.S., Sollins, P., 1999. Standard soil methods for
long-term ecological research. Oxford University Press, New York, NY.
doi:10.1016/s0016-7061(01)00063-5
Rosseel, Y., 2012. lavaan: An R Package for Structural Equation Modelin. Journal of
Statistical Software 48, 1–36.
Rowell, M.J., 1995. Colorimetric method for CO2 measurement in soils. Soil Biology and
Biochemistry 27, 373–375. doi:10.1016/0038-0717(94)00218-P
Roy, S., Bagchi, S., 2022. Large Mammalian Herbivores and the Paradox of Soil Carbon in
Grazing Ecosystems: Role of Microbial Decomposers and Their Enzymes. Ecosystems
25, 976–988. doi:10.1007/s10021-021-00696-8
Roy, S., Naidu, D.G.T., Bagchi, S., 2023. Functional substitutability of native herbivores by
livestock for soil carbon stock is mediated by microbial decomposers. Global Change
Biology 29, 2141–2155. doi:10.1111/gcb.16600
Sankaran, M., Augustine, D.J., 2004. Large herbivores suppress decomposer abundance in a
semiarid grazing ecosystem. Ecology 85, 1052–1061. doi:10.1890/03-0354
Schlaepfer, D.R., Bradford, J.B., Lauenroth, W.K., Munson, S.M., Tietjen, B., Hall, S.A.,
Wilson, S.D., Duniway, M.C., Jia, G., Pyke, D.A., Lkhagva, A., Jamiyansharav, K.,
2017. Climate change reduces extent of temperate drylands and intensifies drought in
deep soils. Nature Communications 8, 1–9. doi:10.1038/ncomms14196
Schloss, P.D., Westcott, S.L., Ryabin, T., Hall, J.R., Hartmann, M., Hollister, E.B.,
Lesniewski, R.A., Oakley, B.B., Parks, D.H., Robinson, C.J., Sahl, J.W., Stres, B.,
Thallinger, G.G., Van Horn, D.J., Weber, C.F., 2009. Introducing mothur: Open-source,
platform-independent, community-supported software for describing and comparing
microbial communities. Applied and Environmental Microbiology 75, 7537–7541.
doi:10.1128/aem.01541-09
Schulze, E.-D., Mooney, H.A., 1993. Biodiversity and Ecosystem Function. Springer Verlag,
New York.
Sinsabaugh, R.L., Lauber, C.L., Weintraub, M.N., Ahmed, B., Allison, S.D., Crenshaw, C.,
Contosta, A.R., Cusack, D., Frey, S., Gallo, M.E., Gartner, T.B., Hobbie, S.E., Holland,
K., Keeler, B.L., Powers, J.S., Stursova, M., Takacs-Vesbach, C., Waldrop, M.P.,
Wallenstein, M.D., Zak, D.R., Zeglin, L.H., 2008. Stoichiometry of soil enzyme activity
at global scale. Ecology Letters 11, 1252–1264. doi:10.1111/j.1461-0248.2008.01245.x

Strickland, M.S., Lauber, C., Fierer, N., Bradford, M.A., 2009. Testing the functional
significance of microbial community composition. Ecology 90, 441–451.
doi:10.1890/08-0296.1

Swan, C.M., Kominoski, J.S., 2012. Biodiversity and Ecosystem Function of Decomposition.
ELS. doi:10.1002/9780470015902.A0023601

Tanentzap, A.J., Coomes, D.A., 2012. Carbon storage in terrestrial ecosystems: Do browsing
and grazing herbivores matter? Biological Reviews 87, 72–94. doi:10.1111/j.1469-
185X.2011.00185.x

Tilman, D., Isbell, F., Cowles, J.M., 2014. Biodiversity and Ecosystem Functioning. Annual
Review of Ecology, Evolution, and Systematics 45, 471–493. doi:10.1146/annurev-
ecolsys-120213-091917

Wagg, C., Bender, S.F., Widmer, F., van der Heijden, M.G.A., 2014. Soil biodiversity and
soil community composition determine ecosystem multifunctionality. Proceedings of the
National Academy of Sciences 111, 5266–5270. doi:10.1073/pnas.1320054111

Wagg, C., Schlaeppi, K., Banerjee, S., Kuramae, E.E., van der Heijden, M.G.A., 2019.
Fungal-bacterial diversity and microbiome complexity predict ecosystem functioning.
Nature Communications 10, 1–10. doi:10.1038/s41467-019-12798-y

Wang, C., Henry, H.A.L., Miao, X., Shi, B., Song, Y., Liang, Q., Sun, W., 2023. Seasonal
variation modifies the spatial patterns of soil microbial community structure and enzyme
activity in a meadow steppe. Applied Soil Ecology 182, 104686.
doi:10.1016/j.apsoil.2022.104686

Wang, C., Tang, Y., 2019. A global meta-analyses of the response of multi-taxa diversity to
grazing intensity in grasslands. Environmental Research Letters 14, 114003.
doi:10.1088/1748-9326/ab4932

Wilson, C.H., Strickland, M.S., Hutchings, J.A., Bianchi, T.S., Flory, S.L., 2018. Grazing
enhances belowground carbon allocation, microbial biomass, and soil carbon in a
subtropical grassland. Global Change Biology 24, 2997–3009. doi:10.1111/gcb.14070

Yang, F., Niu, K., Collins, C.G., Yan, X., Ji, Y., Ling, N., Zhou, X., Du, G., Guo, H., Hu, S.,
2019. Grazing practices affect the soil microbial community composition in a Tibetan
alpine meadow. Land Degradation and Development 30, 49–59. doi:10.1002/ldr.3189

Ylänne, H., Madsen, R.L., Castaño, C., Metcalfe, D.B., Clemmensen, K.E., 2021. Reindeer
control over subarctic treeline alters soil fungal communities with potential
consequences for soil carbon storage. Global Change Biology 27, 4254–4268.
doi:10.1111/gcb.15722

Zhang, S., Chen, Y., Lu, Y., Guo, H., Guo, X., Liu, C., Zhou, X., Zhang, Y., 2022. Spatial
variability and driving factors of soil multifunctionality in drylands of China. Regional
Sustainability 3, 223–232. doi:10.1016/j.regsus.2022.10.001

Zhou, Z., Wang, C., Luo, Y., 2020. Meta-analysis of the impacts of global change factors on
soil microbial diversity and functionality. Nature Communications 11, 3072.
doi:10.1038/s41467-020-16881-7

 FIGURES

 Fig. 1

Figure 1. Treemaps summarizing the distribution of microbial taxonomic profile (phylum

level) and microbial functional profile (n=30 substrates) in the presence and absence of large

mammalian grazers (n=10 paired plots) averaged for all plots and across the growing-season

(Early, Mid, and Late). Area of each cell under taxonomy and function is proportional to the

total abundance of each phylum and microbial activity for each substrate, respectively.

Number denotes the percentage of the relative abundance (taxonomy) and relative microbial

activity (function).

Fig. 2

Figure 2. Species richness of microbial taxonomy (a), functional richness (b), functional

dispersion (c), and multifunction (mg C g^{-1} soil hour^{-1}; d) in the presence and absence of large

mammalian grazers across the growing-season. Species richness (mean ± 95% CI) did not

vary between grazed and ungrazed plots, and with the growing-season. Functional richness,

functional dispersion, and multifunction did not vary between grazed and ungrazed plots, but

varied across the growing-season. Data from n=10 plots.

Fig. 3

Figure 3. Principal coordinate analysis (PCoA) plot for microbial communities and functions

in the presence and absence of large mammalian grazers, and at different points in the

growing-season. The scatter plot shows principal coordinate 1 (Axis 1) versus principal

coordinate 2 (Axis 2). Percentage of variation explained by each axis for each plot is shown

in the axis labels.

Fig. 4

Figure 4. Kriged maps summarizing the relationship of microbial functional richness (a),

functional dispersion (b), and multifunction (mg C g-1 soil hour-1; c) with microbial species

composition. The colored space highlights the interpolation of microbial functions across the

extent of microbial species composition from the study site measured at different points in

the growing-season. Species composition is represented as axis 1 and axis 2 of principal

coordinate analysis (PCoA). Hollow circles denotes the sampled data points for species

composition. Value in the legend indicates the range of interpolated values of respective

functional index. Overall, microbial decomposer functions can vary with species

composition.

Fig. 5

Figure 5. Models representing the distance-based path analyses results that evaluates the

effects of grazing and season on species composition and function relationship via its

influence in soil abiotic properties. Thick arrows are statistically significant paths ($\alpha = 0.10$),

where blue arrows represent positive path coefficient and red arrow represent negative path

coefficient. Value above every significant path represents standardized path estimates.

Modelled paths from grazing and pH were statistically non-significant. Diagnostics for all the

models were SRMR=0.004, RMSEA<0.001, P-value (RMSEA)=0.982, CFI>0.999, χ^2=1.34,

df=2, P-value (χ^2)=0.512. Asterix represent statistical significance (***P$\leq$0.01, **P$\leq$0.05,

and *P$\leq$0.10). Overall, there was a positive relationship between microbial functions and

species composition.

APPENDIX

Diversity and functions of soil microbial communities in the Trans-Himalayan grazing

ecosystem: influence of herbivores and abiotic factors

Shamik Roy, Pronoy Baidya, Jalmesh Karapurkar, Jose Mathew, Sumanta Bagchi*

CONTENTS

 Additional Introduction

    Structure-function relationship documented from across a wide range of producers

  and consumers (Hooper et al., 2005; Reich et al., 2012; Grman et al., 2018; Hautier et al.,

  2018) have established a positive yet decelerating relationship (Fig. A.5.). This is also

  expected for microbial decomposers in soil (Coleman and Whitman, 2005; Delgado-

  Baquerizo et al., 2016a). From the curvilinear shape of the relationship, it follows that

  numbers, quantity and diversity of functions initially rise with taxonomic diversity and

  saturate thereafter (Fig. A.5.). Over the initial ascending part (Fig. A.5.), the structure-

  function relationship is coupled. Any increase in the taxonomic diversity will increase

  functional diversity, albeit at different rates in different microbial communities, and vice-

  versa. For the asymptotic region at high diversity (Fig. A.5.), one expects no relationship

  between the taxonomic and functional aspects. Here, these two aspects may appear decoupled

  (Bell et al., 2009; Frossard et al., 2012; Purahong et al., 2014) because microbial decomposer

  assemblages in soil have high levels of redundancy in their functions (Allison and Martiny,

  2008; Wagg et al., 2019). We know that order and magnitude of this curvilinear relationship

  will vary with ecosystems as scale for taxonomic and functional axes changes (Delgado-

  Baquerizo et al., 2016a). This is because different ecosystems provision for different

  ecological functions and services.

    Using experimental herbivore removal (see Methods), we study whether grazer

  effects on microbial functions are mediated through changes in microbial community

  structure. First, we determine whether grazer-exclusion alters soil microbial community

  structure and functions. Next, we evaluate the empirical support for alternative competing

  hypotheses on how grazer-exclusion may influence the degree of coupling between the

  taxonomic and functional aspects of soil microbial decomposers (Fig. A.5.). Grazer impacts

on functions can differ along a gradient from low to high taxonomic diversity, and this leads

to three competing hypotheses:

i)   Sensitivity Hypothesis (H_S): Over the initial ascending part, grazers can alter the slope

of the relationship (i.e., sensitivity, Fig. A.5.). When grazer-exclusion increases the

slope, it raises the sensitivity of functions to taxonomic diversity; loss of taxonomic

diversity will lead to strong declines in functions. This outcome amounts to large

decline in resilience under loss of herbivores. Otherwise, when grazers decrease the

slope of this relationship, then grazer-exclusion will lower the sensitivity of functions

to taxonomic diversity; loss of taxonomic diversity will lead to proportionately less

reduction in functions. This outcome amounts to greater resilience in absence of

grazing (Fig. A.5.).

ii)   Magnitude Hypothesis (H_M): Any response in functions does not result from changes

in slope of the relationship, but rather from change in the intercept (i.e., magnitude,

Fig. A.5.). If grazing favors functions, then grazer-exclusion will lower the intercept

such that lower levels of taxonomic diversity are needed to maintain functions under

grazing than under grazer-exclusion. However, if grazing lowers functions, then

grazer-exclusion will raise the intercept such that a higher taxonomic diversity is

needed to maintain functions under grazing than under grazer-exclusion. In the

asymptotic region, grazer-exclusion can either increase or decrease microbial

functions (Sankaran and Augustine, 2004; Bagchi et al., 2017), but this can be

independent of any response in taxonomic diversity.

iii) Sensitivity-Magnitude Hypothesis (H_{SM}): This is intermediate between H_S and H_M

whereby the grazer-exclusion will not only change the slope of the relationship

(sensitivity) but also the intercept (magnitude) (Fig. A.5.). This outcome has

consequences for both resilience and functional repertoire of microbial communities

in the absence of grazing.

The null hypothesis against H_S, H_M, and H_{SM} expects no difference in either slope or intercept

under grazing and under grazer-exclusion (Bengtsson, 1998).

Additional methods

Microbial richness and function: model comparison and SEM

We used a model selection and simplification approach with Akaike Information Criterion

(AIC, with ΔAIC $\leq$ -2 between competing models for parsimony) to compare GLMs and

assess support for the alternative hypotheses (Fig. A.5.; H_S, H_{M1}, H_{M2}, H_{SM}) in the data. The

omnibus model contained a 3-way interaction between species richness, grazing, and season

(m_0: Functions ~ Taxonomy × Grazing × Season) alongside the random-effect due to plot-

identity. We checked whether season affects the richness-function relation through 2-way

(Richness×Season) or 3-way (Richness×Grazing×Season) interaction. If season had an effect,

then we evaluated simplified GLMs for each season separately, otherwise we evaluated

GLMs on the full dataset. For these, we tested whether grazers altered the slope or intercept

following a three-step approach.

Step 1: We tested for sensitivity (H_S, H_{SM}) by comparing a model with interactive effects

between taxonomic diversity and grazing (m_1: Functions ~ Species richness × Grazing)

against a model with additive effects between them (m_2: Functions ~ Species richness +

Grazing). Support for m_1 in the data would suggest that H_S, H_{SM} prevails over H_{M1}, H_{M2}, and

over the null hypothesis (Fig. A.5.). To understand whether the data supported H_S or H_{SM} we

used analysis of variance (ANOVA) on the m_1 model. We checked whether grazing had a

statistically significant effect (α=0.05) on functions, either as main-effect or through two-way

interaction (Species richness × Grazing). If grazing had a statistically significant effect

($\alpha=0.05$) on functions through both main-effect and two-way interaction, then data indicates

H_{SM}. Else, if grazing affects function only through two-way interaction, then data indicates

H_S.

Step 2: If m_2 is supported over m_1 in step 1, then we tested for magnitude by comparing m_2

(Functions ~ Species richness + Grazing) and an intercept-only GLM (m_3: Functions ~

Grazing). Model m_2 indicates that both species richness and grazing affects functions,

whereas model m_3 indicates that grazer-effects on functions are independent of species

richness. Here, support for m_2 over m_3 would suggest that grazing-exclusion can alter

magnitude (H_{M1}) without alteration of positive function-species richness relationship.

Step 3: In m_3 is supported over m_2 in step 2, we compared m_3 (Functions ~ Grazing) and null

model (m_4: Functions ~ 1). Null model indicates that variation in microbial functions is

independent of grazing and microbial species richness. Here, support for m_3 would suggest

that grazing can alter magnitude (H_{M2}) without any function-species richness relationship.

GLMs were evaluated for all functional indices (Cardinale et al., 2006; Grman et al., 2018).

Additional results

Relationship between species richness and functions –

The omnibus models for all aspects of microbial functions showed there was a lack of

3-way interaction as well as no 2-way interactions involving season (Table A.2.). So, we

report results on effects of grazer-exclusion on the relationship between microbial functions

and species richness for all seasons together (Fig. A.6.). Model comparison for all functional

indices (m_1, m_2, m_3, m_4) showed no relationship between species richness and functions

across all seasons, and this was not altered by grazing (Fig. A.6; Table A.2., A.3.). Since

grazer-exclusion neither altered the slope nor the intercept of the taxonomy-function

relationship (Fig. A.6; Table A.2., A.3.), the null hypothesis was supported over H_S, H_{SM},

H_{M1}, and H_{M2}. Model comparison of individual functions suggests a significant relationship

between species richness and 8 out of 30 functions (Table A.4., A.5.). For alanine, inosine,

malic acid, and mannitol, the data supports H_S such that grazer-exclusion reduces the

sensitivity (slope) of the relationship. Data for glucosamine supports H_{M2} where there is no

relationship between species richness and the function, but grazer-exclusion increases the

magnitude of glucosamine. Data for mannose supports H_{M1} where grazer-exclusion increased

the magnitude with no effect on the positive relationship between species richness and the

function. For polysorbate and putrescine, the relationship varied with season due to

significant two-way interaction (Richness×Season, $P<0.05$; Table A.5.). Model comparison

for polysorbate find support for H_{M2} hypothesis in the mid-season and H_{M1} hypothesis in the

late-season (Table A.5.). Model comparison for putrescine find support for H_{M1} hypothesis in

the mid-season (Table A.5.).

1014

1015   Figure A.1. Map of sampling locations

1016

1017   Figure A.2. Edaphic conditions soil pH, soil moisture, and soil texture in in the presence and

1018   absence of large mammalian grazers (n=10 paired plots). Soil moisture is measured as

1019   volumetric water content (VWC, %) across the growing-season.

1020

1021

 Figure A.3. Soil microbial communities at the phylum level (covering >=90% of the total

 abundance) and relative microbial functional profile (n=30 substrates) in the presence and

 absence of large mammalian grazers (n=10 paired plots) across the growing-season.

Figure A.4. dbSEM for all functions

Figure A.5. Concept diagram to summarize hypothesized effects of microbial species

richness on microbial functions in the presence (solid line) or absence (dashed

line) of large mammalian grazers. Exclusion of grazers can alter the sensitivity of the

functions to species diversity by either increasing or decreasing the slope (Sensitivity

hypothesis; a). Alternatively, grazer-exclusion can alter the intercept (Magnitude hypothesis;

c, d). Grazer-exclusion can also alter slope and magnitude of the relationship simultaneously

(Sensitivity-Magnitude hypothesis; e).

1037

1038

1039

1040

1041   Figure A.6. Relationship between species richness and functional richness (a), functional

1042   dispersion (b), and multifunction (c) in the presence and absence of large mammalian grazers

1043   across the growing-season. Species richness was not related to microbial functions in both

1044   grazed and ungrazed plots across the growing-season. Units for 'Multifunction' was mg C g^{-1}

1045   soil hour^{-1}.

1046

1047

1048   Table A.1. Best model for kriging using sum of squares (SSQ)

	Trend	Nugget	Exponential (SSQ)	Spherical (SSQ)	Gaussian (SSQ)
Functional richness	No	Fixed	52551.11	59125.98	59125.98
		Estimated	52745.91	51583.67	59125.98
	1st	Fixed	33433.22	39481.94	39481.94
		Estimated	33433.24	39481.94	39481.94
	2nd	Fixed	**29508.12**	34531.09	34531.09
		Estimated	29508.12	34531.09	34531.09
Functional dispersion	No	Fixed	0.007484251	0.007484251	0.007484251
		Estimated	0.007484251	0.007484251	0.007484251
	1st	Fixed	0.005320591	0.005320591	0.005320591
		Estimated	0.005320591	0.005320591	0.005320591
	2nd	Fixed	**0.004776054**	0.004776054	0.004776054
		Estimated	0.004776054	0.004776054	0.004776054
Multifunction	No	Fixed	40569.47	40569.47	40569.47
		Estimated	40569.47	40569.47	40569.47
	1st	Fixed	29273.22	29273.22	29273.22
		Estimated	29273.22	29273.22	29273.22
	2nd	Fixed	**22815.05**	22815.05	22815.05
		Estimated	22815.05	22815.05	22815.05

1049

1050

1051   Table A.2. Summary of omnibus GLM model: Functions ~ Species richness * Grazing *

1052   Season

y-variable	x-variable	F-value	P-value
Functional richness	Species Richness	$F_{1,48}=0.20$	0.653
	Grazing	$F_{1,48}=0.01$	0.944
	Season	$F_{2,48}=36.82$	<0.001
	Taxonomy × Grazing	$F_{1,48}=1.94$	0.169
	Taxonomy × Season	$F_{2,48}=1.25$	0.296
	Grazing × Season	$F_{2,48}=0.28$	0.754
	Taxonomy × Grazing × Season	$F_{2,48}=0.34$	0.716
Functional dispersion	Species Richness	$F_{1,48}=0.07$	0.796
	Grazing	$F_{1,48}=0.01$	0.952
	Season	$F_{2,48}=42.18$	<0.001
	Taxonomy × Grazing	$F_{1,48}=0.11$	0.737
	Taxonomy × Season	$F_{2,48}=0.27$	0.767
	Grazing × Season	$F_{2,48}=0.18$	0.825
	Taxonomy × Grazing × Season	$F_{2,48}=0.51$	0.604
Multifunction	Species Richness	$F_{1,48}=0.15$	0.701
	Grazing	$F_{1,48}=0.11$	0.747
	Season	$F_{2,48}=10.80$	<0.001
	Taxonomy × Grazing	$F_{1,48}=1.41$	0.241
	Taxonomy × Season	$F_{2,48}=2.19$	0.122

	Grazing × Season	$F_{2,48}=0.34$	0.716
	Taxonomy × Grazing × Season	$F_{2,48}=0.31$	0.738

1053
1054
1055

1056   Table A.3. Summary of the GLM model comparison to test for alternative hypothesis (H_S,

1057 H_{M1}, H_{M2}, H_{SM}).

	Models	F-value	P-value	ΔAIC
Functional richness	1, 2	$F_{56,57}=3.17$	0.080	-1.31
	2, 3	$F_{57,58}=0.09$	0.760	1.90
	3, 4	$F_{58,59}=0.01$	>0.999	2
Functional dispersion	1, 2	$F_{56,57}=1.19$	0.28	0.74
	2, 3	$F_{57,58}=0.03$	0.873	1.97
	3, 4	$F_{58,59}=0.01$	0.948	1.99
Multifunction	1, 2	$F_{56,57}=2.63$	0.111	-0.75
	2, 3	$F_{57,58}=0.14$	0.711	1.85
	3, 4	$F_{58,59}=0.05$	0.822	1.95

1058

1059 Table A.4. Summary of omnibus GLM model for 30 functions: Functions ~ Species richness *

1060 Grazing * Season. R: Species richness, Grazing: Grazing, S: Season

Functions ↓	R		G		S		R:G		R:S		G:S		R:G:S	
	$F_{1,48}$	P	$F_{1,48}$	P	$F_{2,48}$	P	$F_{1,48}$	P	$F_{2,48}$	P	$F_{2,48}$	P	$F_{2,48}$	P
Alanine	0.01	0.924	1.72	0.195	0.07	0.931	**6.05**	**0.018**	2.29	0.112	2.1	0.133	0.23	0.792
Arginine	0.01	0.977	0.2	0.657	**15.31**	**0.001**	1.81	0.185	0.52	0.597	1.25	0.296	0.64	0.533
Ascorbic acid	0.34	0.562	0.08	0.779	**3.86**	**0.028**	2.71	0.106	0.05	0.954	1.37	0.263	2.39	0.102
Asparagine	0.01	0.951	0.07	0.793	0.69	0.505	1.01	0.32	0.86	0.431	0.1	0.906	0.77	0.471
Citric acid	0.05	0.825	0.16	0.688	**4.09**	**0.023**	0.47	0.497	0.96	0.39	0.51	0.607	1.91	0.159
Dextrose	1.5	0.226	1.81	0.185	**9.87**	**0.001**	0.97	0.33	1.98	0.149	0.01	0.997	2.43	0.099
Fumaric acid	1.57	0.217	0.17	0.685	1.35	0.269	0.79	0.379	0.45	0.638	0.71	0.495	0.91	0.408
Gallic acid	0.12	0.734	0.01	0.993	3.18	0.051	1.41	0.241	**3.34**	**0.044**	0.32	0.728	0.72	0.49
Glucosamine	0.77	0.384	**3.78**	**0.058**	2.68	0.079	0.46	0.499	0.2	0.816	1.35	0.27	0.52	0.596
Glutamic acid	0.01	0.912	0.32	0.575	0.61	0.547	1.04	0.314	1.1	0.341	0.53	0.593	0.07	0.933
Glutamine	0.27	0.603	0.62	0.435	2.44	0.098	0.01	0.946	**3.36**	**0.043**	0.08	0.92	0.85	0.435
Glycerol	0.92	0.343	**3.02**	**0.089**	1.23	0.3	0.66	0.421	0.21	0.815	0.93	0.403	0.08	0.924
Histidine	0.09	0.764	0.26	0.616	**3.86**	**0.028**	0.23	0.634	0.45	0.638	0.28	0.756	0.49	0.614
Inosine	0.01	0.982	0.35	0.555	**8.09**	**0.001**	3.2	0.08	0.74	0.48	1.13	0.332	0.57	0.569
Inositol	0.08	0.772	0.76	0.387	**5.37**	**0.008**	1.04	0.313	2.61	0.084	1.09	0.343	0.33	0.722
Ketoglutarate	1.21	0.277	0.12	0.725	0.79	0.462	0.79	0.38	0.49	0.615	0.83	0.441	0.65	0.526
Malic acid	0.45	0.505	2.35	0.132	**9.79**	**0.001**	3.56	0.065	1.22	0.304	1.42	0.251	0.29	0.746
Malonic acid	0.59	0.446	0.01	0.966	**11.14**	**0.001**	0.13	0.719	0.49	0.615	0.33	0.724	0.4	0.672
Mannitol	0.03	0.858	0.24	0.63	**6.65**	**0.003**	2.94	0.093	1.5	0.234	0.01	0.989	0.69	0.506
Mannose	**5.85**	**0.019**	1.67	0.202	**6.23**	**0.004**	0.06	0.815	0.64	0.531	2.19	0.123	0.23	0.792
Oxalic acid	0.08	0.778	0.4	0.53	0.91	0.409	0.77	0.384	0.12	0.89	0.8	0.456	0.47	0.63
Polysorbate	0.39	0.535	0.02	0.892	1.09	0.344	0.2	0.657	**4.35**	**0.018**	1.93	0.157	0.68	0.51
Putrescine	0.9	0.348	2.18	0.146	3.03	0.057	0.07	0.791	**4.5**	**0.016**	0.85	0.432	0.53	0.593
Serine	0.95	0.336	0.53	0.469	1.11	0.339	3.23	0.078	1.19	0.312	1.41	0.254	0.83	0.443
Sorbitol	0.02	0.898	0.56	0.459	**4.88**	**0.012**	0.82	0.37	0.12	0.891	0.18	0.835	0.38	0.688
Succinic acid	0.53	0.471	0.84	0.363	0.21	0.809	0.77	0.384	0.36	0.699	0.31	0.732	0.24	0.786
Sucrose	3.03	0.088	1.26	0.267	**6.72**	**0.003**	0.04	0.843	**3.23**	**0.048**	0.26	0.768	1.39	0.259
Tartaric acid	0.59	0.447	0.7	0.407	0.64	0.53	0.17	0.679	1.03	0.363	0.66	0.522	0.65	0.525
Xylitol	0.13	0.719	0.5	0.485	**9.14**	**0.001**	1.26	0.268	0.59	0.558	0.05	0.952	0.1	0.907
Xylose	0.11	0.742	1.43	0.238	0.25	0.78	0.42	0.521	1.04	0.362	0.63	0.538	0.12	0.89

1061

1062

1063 Table A.5. Summary of the GLM model comparison to test for alternative hypotheses (H_S,

1064 H_{M1}, H_{M2}, H_{SM}) in individual functions. For few functions the GLMs were evaluated for each

1065 season separately due to a significant effect of season as identified from the omnibus model

1066 summarized in Table A.4.

Function	Season	Models	F-value	P-value	ΔAIC
Alanine		**1, 2**	**$F_{56,57}=6.04$**	**0.017**	**-4.15**
		2, 3	$F_{57,58}=0.08$	0.778	1.92
		3, 4	$F_{58,59}=1.50$	0.226	0.47
Arginine		1, 2	$F_{56,57}=3.16$	0.081	-1.29
		2, 3	$F_{57,58}=0.01$	0.973	1.99
		3, 4	$F_{58,59}=0.13$	0.715	1.86
Ascorbic acid		1, 2	$F_{56,57}=1.34$	0.252	0.58
		2, 3	$F_{57,58}=0.33$	0.566	1.65
		3, 4	$F_{58,59}=0.03$	0.861	1.96
Asparagine		1, 2	$F_{56,57}=0.74$	0.395	1.21
		2, 3	$F_{57,58}=0.01$	0.917	1.99
		3, 4	$F_{58,59}=0.07$	0.796	1.93
Citric acid		1, 2	$F_{56,57}=0.11$	0.740	1.88
		2, 3	$F_{57,58}=0.07$	0.790	1.92
		3, 4	$F_{58,59}=0.12$	0.729	1.87
Dextrose		1, 2	$F_{56,57}=1.94$	0.169	-0.04
		2, 3	$F_{57,58}=0.74$	0.394	1.23
		3, 4	$F_{58,59}=1.70$	0.197	0.26
Fumaric acid		1, 2	$F_{56,57}=1.09$	0.301	0.84
		2, 3	$F_{57,58}=1.41$	0.240	0.54
		3, 4	$F_{58,59}=0.36$	0.549	1.62
Gallic acid	Early	1, 2	$F_{16,17}=0.09$	0.773	1.89
		2, 3	$F_{17,18}=3.67$	0.072	-1.91
		3, 4	$F_{18,19}=0.01$	0.992	1.99
	Mid	1, 2	$F_{16,17}=3.01$	0.102	-1.45
		2, 3	$F_{17,18}=1.69$	0.211	0.10
		3, 4	$F_{18,19}=0.15$	0.699	1.83
	Late	1, 2	$F_{16,17}=0.11$	0.744	1.86
		2, 3	$F_{17,18}=1.78$	0.199	0.01
		3, 4	$F_{18,19}=0.19$	0.669	1.79

Glucosamine		1, 2	$F_{56,57}=0.11$	0.741	1.88
		2, 3	$F_{57,58}=0.31$	0.578	1.67
		3, 4	**$F_{58,59}=4.22$**	**0.045**	**-2.21**
Glutamic acid		1, 2	$F_{56,57}=1.46$	0.232	0.45
		2, 3	$F_{57,58}=0.04$	0.839	1.96
		3, 4	$F_{58,59}=0.32$	0.576	1.67
Glutamine	Early	1, 2	$F_{16,17}=0.06$	0.806	1.92
		2, 3	$F_{17,18}=1.69$	0.211	0.11
		3, 4	$F_{18,19}=0.03$	0.874	1.97
	Mid	1, 2	$F_{16,17}=1.23$	0.283	0.51
		2, 3	$F_{17,18}=1.03$	0.324	0.82
		3, 4	$F_{18,19}=0.73$	0.404	1.20
	Late	1, 2	$F_{16,17}=1.55$	0.231	0.15
		2, 3	$F_{17,18}=3.88$	0.065	-2.11
		3, 4	$F_{18,19}=1.02$	0.325	0.89
Glycerol		1, 2	$F_{56,57}=0.35$	0.554	1.62
		2, 3	$F_{57,58}=1.58$	0.215	0.36
		3, 4	$F_{58,59}=2.59$	0.113	-0.62
Histidine		1, 2	$F_{56,57}=0.73$	0.398	1.23
		2, 3	$F_{57,58}=0.05$	0.829	1.95
		3, 4	$F_{58,59}=0.29$	0.589	1.69
Inosine		**1, 2**	**$F_{56,57}=4.66$**	**0.035**	**-2.79**
		2, 3	$F_{57,58}=0.01$	0.950	1.99
		3, 4	$F_{58,59}=0.28$	0.599	1.71
Inositol		1, 2	$F_{56,57}=1.71$	0.196	0.19
		2, 3	$F_{57,58}=0.02$	0.890	1.98
		3, 4	$F_{58,59}=0.70$	0.405	1.28
α-Ketoglutarate		1, 2	$F_{56,57}=1.06$	0.301	0.87
		2, 3	$F_{57,58}=1.37$	0.247	0.58
		3, 4	$F_{58,59}=0.03$	0.856	1.97
Malic acid		**1, 2**	**$F_{56,57}=4.35$**	**0.041**	**-2.49**
		2, 3	$F_{57,58}=0.13$	0.716	1.86
		3, 4	$F_{58,59}=1.97$	0.1666	-0.002
Malonic acid		1, 2	$F_{56,57}=0.08$	0.773	1.91
		2, 3	$F_{57,58}=0.459$	0.501	1.52
		3, 4	$F_{58,59}=0.01$	0.945	1.99
Mannitol		**1, 2**	**$F_{56,57}=4.37$**	**0.041**	**-2.51**

		2, 3	$F_{57,58}=0.01$	0.926	1.99
		3, 4	$F_{58,59}=0.217$	0.643	1.78
Mannose		1, 2	$F_{56,57}=0.44$	0.508	1.53
		2, 3	**$F_{57,58}=4.09$**	**0.047**	**-2.16**
		3, 4	$F_{58,59}=2.22$	0.141	-0.25
Oxalic acid		1, 2	$F_{56,57}=0.48$	0.491	1.49
		2, 3	$F_{57,58}=0.15$	0.697	1.84
		3, 4	$F_{58,59}=0.37$	0.547	1.62
Polysorbate	Early	1, 2	$F_{16,17}=0.07$	0.798	1.91
		2, 3	$F_{17,18}=2.41$	0.139	-0.65
		3, 4	$F_{18,19}=0.15$	0.699	1.83
	Mid	1, 2	$F_{16,17}=0.02$	0.886	1.97
		2, 3	$F_{17,18}=0.69$	0.417	1.20
		3, 4	**$F_{18,19}=5.18$**	**0.035**	**-3.06**
	Late	1, 2	$F_{16,17}=3.69$	0.073	-2.15
		2, 3	**$F_{17,18}=12.32$**	**0.003**	**-8.90**
		3, 4	$F_{18,19}=0.48$	0.495	1.47
Putrescine	Early	1, 2	$F_{16,17}=0.69$	0.416	1.15
		2, 3	$F_{17,18}=0.47$	0.504	1.46
		3, 4	$F_{18,19}=0.03$	0.875	1.97
	Mid	1, 2	$F_{16,17}=0.01$	0.947	1.99
		2, 3	$F_{17,18}=0.09$	0.767	1.89
		3, 4	$F_{18,19}=0.43$	0.520	1.53
	Late	1, 2	$F_{16,17}=0.79$	0.388	1.04
		2, 3	**$F_{17,18}=8.55$**	**0.009**	**-6.14**
		3, 4	$F_{18,19}=3.14$	0.093	-1.22
Serine		1, 2	$F_{56,57}=3.75$	0.058	-1.89
		2, 3	$F_{57,58}=1.09$	0.301	0.86
		3, 4	$F_{58,59}=31$	0.582	1.68
Sorbitol		1, 2	$F_{56,57}=1.76$	0.190	0.14
		2, 3	$F_{57,58}=0.06$	0.813	1.94
		3, 4	$F_{58,59}=0.49$	0.484	1.49
Succinic acid		1, 2	$F_{56,57}=1.04$	0.312	0.89
		2, 3	$F_{57,58}=0.83$	0.367	1.14
		3, 4	$F_{58,59}=0.71$	0.404	1.27
Sucrose	Early	1, 2	$F_{16,17}=0.79$	0.386	1.03
		2, 3	$F_{17,18}=3.25$	0.089	-1.50

		3, 4	$F_{18,19}=1.22$	0.284	0.69
	Mid	1, 2	$F_{16,17}=0.05$	0.832	1.94
		2, 3	$F_{17,18}=1.63$	0.218	0.16
		3, 4	$F_{18,19}=0.07$	0.796	1.92
	Late	1, 2	$F_{16,17}=2.41$	0.140	-0.81
		2, 3	$F_{17,18}=0.21$	0.652	1.75
		3, 4	$F_{18,19}=1.37$	0.26	0.53
Tartaric acid		1, 2	$F_{56,57}=0.33$	0.571	1.65
		2, 3	$F_{57,58}=0.83$	0.366	1.13
		3, 4	$F_{58,59}=0.53$	0.471	1.46
Xylitol		1, 2	$F_{56,57}=2.53$	0.117	-0.66
		2, 3	$F_{57,58}=0.05$	0.822	1.95
		3, 4	$F_{58,59}=0.47$	0.494	1.51
Xylose		1, 2	$F_{56,57}=0.39$	0.535	1.58
		2, 3	$F_{57,58}=0.02$	0.884	1.98
		3, 4	$F_{58,59}=1.68$	0.200	0.29

1067
1068

1069