Zn²⁺ is Essential for Ca²⁺ Oscillations in Mouse Eggs

Hiroki Akizawa[1], Emily Lopes[1,2], Rafael A. Fissore[1*]

[1]Department of Veterinary and Animal Sciences, University of Massachusetts Amherst,

661 North Pleasant Street, Amherst, Massachusetts, 01003, United States.

[2]Molecular and Cellular Biology Graduate Program, University of Massachusetts,

Amherst, Massachusetts, 01003, United States.

Running Tittle: Labile Zn²⁺ and Ca²⁺ release

Key Words: Fertilization, mammals, Ca²⁺, IP₃R1, oocytes, eggs, sperm, divalent cations, chelators

[*]Author for correspondence (rfissore@umass.edu)

*Rafael A. Fissore

661 North Pleasant Street

ISB-427A

Department of Veterinary and Animal Science

University of Massachusetts, Amherst, 01003

Phone:413-687-5773

Email: rfissore@umass.edu

Abstract

Changes in the intracellular concentration of free calcium (Ca^{2+}) underpin egg activation and initiation of development in animals and plants. In mammals, the Ca^{2+} release is periodical, known as Ca^{2+} oscillations, and mediated by the type 1 inositol 1,4,5-trisphosphate receptor (IP$_3$R1). Another divalent cation, zinc (Zn^{2+}), increases exponentially during oocyte maturation and is vital for meiotic transitions, arrests, and polyspermy prevention. It is unknown if these pivotal cations interplay during fertilization. Here, using mouse eggs, we showed that basal concentrations of labile Zn^{2+} are indispensable for sperm-initiated Ca^{2+} oscillations because Zn^{2+}-deficient conditions induced by cell-permeable chelators abrogated Ca^{2+} responses evoked by fertilization and other physiological and pharmacological agonists. We also found that chemically- or genetically generated Zn^{2+}-deficient eggs displayed reduced IP$_3$R1 sensitivity and diminished ER Ca^{2+} leak despite the stable content of the stores and IP$_3$R1 mass. Resupplying Zn^{2+} restarted Ca^{2+} oscillations, but excessive Zn^{2+} prevented and terminated them, hindering IP$_3$R1 responsiveness. The findings suggest that a permissive window of Zn^{2+} concentrations is required for Ca^{2+} responses and IP$_3$R1 function in eggs, ensuring optimal response to fertilization and egg activation.

Introduction

Vertebrate eggs are arrested at the metaphase stage of the second meiosis (MII) when ovulated because they have an active Cdk1/cyclin B complex and inactive APC/C^{Cdc20} (Heim et al., 2018). Release from MII initiates egg activation, the first hallmark of embryonic development (Ducibella et al., 2002; Schultz and Kopf, 1995). The universal signal of egg activation is an increase in the intracellular concentration of calcium (Ca^{2+}) (Ridgway et al., 1977; Stricker, 1999). Ca^{2+} release causes the inactivation of the APC/C inhibitor Emi2, which enhances cyclin B degradation and induces meiotic exit (Lorca et al., 1993; Shoji et al., 2006; Suzuki et al., 2010a). In mammals, the stereotypical fertilization Ca^{2+} signal, oscillations, consists of transient but periodical Ca^{2+} increases that promote progression into interphase (Deguchi et al., 2000; Miyazaki et al., 1986). The sperm-borne Phospholipase C zeta1 (PLCζ) persistently stimulates the production of inositol 1,4,5-trisphosphate (IP$_3$) (Matsu-ura et al., 2019; Saunders et al., 2002; Wu et al., 2001) that binds its cognate receptor in the endoplasmic reticulum (ER), IP$_3$R1 and causes Ca^{2+} release from the egg's main Ca^{2+} reservoir (Wakai et al., 2019). The intake of extracellular Ca^{2+} via plasma membrane channels and transporters ensures the persistence of the oscillations (Miao et al., 2012; Stein et al., 2020; Wakai et al., 2019, 2013).

Before fertilization, maturing oocytes undergo cellular and biochemical
modifications (see for review (Ajduk et al., 2008)). The nucleus of immature oocytes,
known as the germinal vesicle (GV), undergoes the breakdown of its envelope marking
the onset of maturation and setting in motion a series of cellular events that culminate
with the release of the first polar body, the correct ploidy for fertilization, and re-arrest at
MII (Eppig, 1996). Other organelles are also reorganized, such as cortical granules
migrate to the cortex for exocytosis and polyspermy block, mitochondria undergo
repositioning, and the cytoplasm's redox state becomes progressively reduced to promote
the exchange of the sperm's protamine load (Liu, 2011; Perreault et al., 1988; Wakai et
al., 2014). Wide-ranging adaptations also occur in the Ca^{2+} release machinery to produce
timely and protracted Ca^{2+} oscillations following sperm entry (Fujiwara et al., 1993;
Lawrence et al., 1998), including the increase in the content of the Ca^{2+} stores, ER
reorganization with cortical cluster formation, and increased IP$_3$R1 sensitivity (Lee et al.,
2006; Wakai et al., 2012). The total intracellular levels of zinc (Zn^{2+}) also remarkably
increase during maturation, amounting to a 50% rise, which is necessary for oocytes to
proceed to the telophase I of meiosis and beyond (Kim et al., 2010). Remarkably, after
fertilization, Zn^{2+} levels need to decrease, as Emi2 is a Zn^{2+}-associated molecule, and
high Zn^2 levels prevent MII exit (Bernhardt et al., 2012; Shoji et al., 2014; Suzuki et al.,
2010b). Following the initiation of Ca^{2+} oscillations, approximately 10 to 20% of the Zn^{2+}
accrued during maturation is ejected during the Zn^{2+} sparks, a conserved event in
vertebrates and invertebrate species (Kim et al., 2011; Que et al., 2019; Tokuhiro and
Dean, 2018; Wozniak et al., 2020; Zhang et al., 2016). The use of Zn^{2+} chelators such as
N,N,N,N-tetrakis (2-pyridinylmethyl)-1,2-ethylenediamine (TPEN) to create Zn^{2+}-
deficient conditions buttressed the importance of Zn^{2+} during meiotic transitions (Kim et
al., 2010; Suzuki et al., 2010b). However, whether the analogous dynamics of Ca^{2+} and
Zn^{2+} during maturation imply crosstalk and Zn^{2+} levels modulate Ca^{2+} release during
fertilization is unknown.
IP$_3$Rs are the most abundant intracellular Ca^{2+} release channel in non-muscle
cells (Berridge, 2016). They form a channel by assembling into tetramers with each
subunit of ~270kDa MW (Taylor and Tovey, 2010). Mammalian eggs express the type I
IP$_3$R, the most widespread isoform (Fissore et al., 1999; Parrington et al., 1998). IP$_3$R1 is
essential for egg activation because its inhibition precludes Ca^{2+} oscillations (Miyazaki
and Ito, 2006; Miyazaki et al., 1992; Xu et al., 2003). Myriad and occasionally cell-
specific factors influence Ca^{2+} release through the IP$_3$R1 (Taylor and Tovey, 2010). For
example, following fertilization, IP$_3$R1 undergoes ligand-induced degradation caused by
the sperm-initiated long-lasting production of IP$_3$ that effectively reduces the IP$_3$R1 mass

(Brind et al., 2000; Jellerette et al., 2000). Another regulatory mechanism is Ca^{2+}, a
universal cofactor, which biphasically regulates IP_3Rs' channel opening (Iino, 1990; Jean
and Klee, 1986), congruent with several Ca^{2+} and calmodulin binding sites on the
channel's sequence (Sienaert et al., 1997; Sipma et al., 1999). Notably, Zn^{2+} may also
participate in IP3R1 regulation. Recent studies using electron cryomicroscopy (cryoEM),
a technique that allows peering into the structure of IP3R1 with a near-atomic resolution,
have revealed that a helical linker (LNK) domain near the C-terminus mediates the
coupling between the N- and C-terminal ends necessary for channel opening (Fan et al.,
2015). The LNK domain contains a putative Zinc-finger motif proposed to be vital for
IP3R1 function (Fan et al., 2015; Paknejad and Hite, 2018). Therefore, the exponential
increase in Zn^{2+} levels in maturing oocytes, besides its essential role in meiosis
progression, may optimize the IP3R1 function, revealing hitherto unknown cooperation
between these cations during fertilization.
Here, we examined whether crosstalk between Ca^{2+} and Zn^{2+} is required to
initiate and sustain Ca^{2+} oscillations and maintain Ca^{2+} store content in MII eggs. We
found that Zn^{2+}-deficient conditions inhibited Ca^{2+} release and oscillations without
reducing Ca^{2+} stores, IP_3 production, IP3R1 expression, or altering the viability of eggs or
zygotes. We show instead that Zn^{2+} deficiency impaired IP3R1 function and lessened the
receptor's ability to gate Ca^{2+} release out of the ER. Remarkably, resupplying Zn^{2+} re-
established the oscillations interrupted by low Zn^{2+}, although persistent increases in
intracellular Zn^{2+} were harmful, disrupting the Ca^{2+} responses and preventing egg
activation. Together, the results show that besides contributing to oocyte maturation, Zn^{2+}
has a central function in Ca^{2+} homeostasis such that optimal Zn^{2+} concentrations ensure
IP3R1 function and the Ca^{2+} oscillations required for initiating embryo development.
119

Results
TPEN dose-dependently lowers intracellular Zn^{2+} and inhibits sperm-initiated Ca^{2+}
oscillations.
TPEN is a cell-permeable, non-specific chelator with a high affinity for transition
metals widely used to study their function in cell physiology (Arslan et al., 1985; Lo et
al., 2020). Mouse oocytes and eggs have exceedingly high intracellular concentrations of
Zn^{2+} (Kim et al., 2011, 2010), and the TPEN-induced defects in the progression of meiosis
have been ascribed to its chelation (Bernhardt et al., 2011; Kim et al., 2010). In support
of this view, the Zn^{2+} levels of cells showed acute reduction after TPEN addition, as
reported by indicators such as FluoZin-3 (Arslan et al., 1985; Suzuki et al., 2010b).
Studies in mouse eggs also showed that the addition of mM concentrations of TPEN

disrupted Ca^{2+} oscillations initiated by fertilization or $SrCl_2$ (Lawrence et al., 1998;
Suzuki et al., 2010b), but the mechanism(s) and target(s) of the inhibition remained
unknown. To gain insight into this phenomenon, we first performed dose-titration studies
to determine the effectiveness of TPEN in lowering Zn^{2+} in eggs. The addition of 2.5 mM
TPEN protractedly reduced Zn^{2+} levels, whereas 5 and 10 mM TPEN acutely and
persistently reduced FluoZin-3 fluorescence (**Fig. 1A)**. These concentrations of TPEN are
higher than the reported free Zn^{2+} concentrations in cells, but within range of those of
found in typical culture conditions (Lo et al., 2020; Qin et al., 2011). We next determined
the concentrations of TPEN required to abrogate fertilization-initiated oscillations.
Following intracytoplasmic sperm injection (ICSI), we monitored Ca^{2+} responses while
increasing TPEN concentrations. As shown in **Fig. 1B**, 5 and 10 µM TPEN effectively
blocked ICSI-induced Ca^{2+} oscillations in over half of the treated cells, and the remaining
eggs, after a prolonged interval, resumed lower-frequency rises (**Fig. 1B**-center panels).
Finally, 50 µM or greater concentrations of TPEN permanently blocked these oscillations
(**Fig. 1B**-right panel). It is noteworthy that at the time of addition, TPEN induces a sharp
drop in basal Fura-2 F340/ F380 ratios, consistent with Fura-2's high affinity for Zn^{2+}
(Snitsarev et al., 1996).

We next used membrane-permeable and -impermeable chelators to assess
whether TPEN inhibited Ca^{2+} oscillations by chelating Zn^{2+} from intracellular or
extracellular compartments. The addition of the high-affinity but cell-impermeable Zn^{2+}
chelators DTPA and EDTA neither terminated nor temporarily interrupted ICSI-induced
Ca^{2+} oscillations (**Fig. 1C**). Protractedly, Ca^{2+} oscillations slowed down, possibly because
of chelation and lowering of external Ca^{2+} (**Fig. 1C**). Conversely, TPA, a permeable Zn^{2+}
chelator, blocked the ICSI-initiated oscillations but required higher concentrations than
TPEN (**Fig. 1D**). Collectively, the data suggest that basal levels of labile Zn^{2+} are essential
to sustain the fertilization-initiated Ca^{2+} oscillations in eggs.

We next evaluated whether TPEN supplementation prevented the completion of
meiosis and pronuclear (PN) formation following fertilization (**Fig. 1E and Table 1**). All
fertilized eggs promptly extruded second polar bodies regardless of treatment (**Fig. 1E**).
TPEN, however, impaired PN formation, and by 4- or 7-hr. post-ICSI, most treated eggs
failed to show PNs, unlike controls (**Fig. 1E and Table 1**). Together, these results
demonstrate that depletion of Zn^{2+} terminates Ca^{2+} oscillations and delays or prevents
events of egg activation, including PN formation.

TPEN is a universal inhibitor of Ca^{2+} oscillations in eggs.
Mammalian eggs initiate Ca^{2+} oscillations in response to numerous stimuli and

conditions (Miyazaki and Ito, 2006; Wakai and Fissore, 2013). Fertilization and its release
of PLCz stimulate the phosphoinositide pathway leading to the production of IP_3 and Ca^{2+}
oscillations (Miyazaki, 1988; Saunders et al., 2002). Neurotransmitters such as
acetylcholine (Ach) and other G-protein coupled receptor agonists engage a similar
mechanism (Dupont et al., 1996; Kang et al., 2003), although in these cases, IP_3
production occurs at the plasma membrane and is short-lived (Kang et al., 2003; Swann
and Parrington, 1999). Agonists such as $SrCl_2$ and thimerosal generate oscillations by
sensitizing IP3R1 without producing IP_3. The mechanism(s) of $SrCl_2$ is unclear, although
its actions are reportedly directly on the IP3R1 (Hajnóczky and Thomas, 1997; Hamada
et al., 2003; Nomikos et al., 2015, 2011; Sanders et al., 2018). Thimerosal oxidizes dozens
of thiol groups in the receptor, which enhances the receptor's sensitivity and ability to
release Ca^{2+} (Bootman et al., 1992; Evellin et al., 2002; Joseph et al., 2018). We took
advantage of the varied points at which the mentioned agonists engage the
phosphoinositide pathway to examine TPEN's effectiveness in inhibiting their effects.
m*Plcz* mRNA injection, like fertilization, induces persistent Ca^{2+} oscillations, although
m*Plcz*'s tends to be more robust. Consistent with this, the addition of 10 and 25 µM TPEN
transiently interrupted or belatedly terminated oscillations, whereas 50 µM acutely
stopped all responses (**Fig. 2A**). By contrast, $SrCl_2$-initiated rises were the most sensitive
to Zn^{2+}-deficient conditions, with 2.5 µM TPEN nearly terminating all oscillations that 5
µM did (**Fig. 2B**). TPEN was equally effective in ending the Ach-induced Ca^{2+} responses
(**Fig. 2C**), but curbing thimerosal responses required higher concentrations (**Fig. 2D**).
Lastly, we ruled out that downregulation of IP3R1 was responsible for the slow-down or
termination of the oscillations by TPEN. To accomplish this, we examined the IP3R1 mass
in eggs (Jellerette et al., 2004) with and without TPEN supplementation and injection of
m*Plcz* mRNA. By 4-h post-injection, the mRNA induced the expected down-regulation
of IP3R1 reactivity vs. uninjected eggs, but in TPEN-treated and *Plcz* mRNA-injected
eggs, the decrease was insignificant (**Fig. 2F**). These findings together show that Zn^{2+}
deficiency inhibits the IP3R1-mediated Ca^{2+} oscillations independently of IP_3 production
or loss of receptor, suggesting a role of Zn^{2+} on IP3R1 function (**Fig. 2E**).
196

Zn^{2+} depletion reduces IP3R1-mediated Ca^{2+} release.
To directly assess the inhibitory effects of TPEN on IP3R1 function, we used caged
IP_3 (cIP_3) that, after short UV pulses, releases IP_3 into the ooplasm (Wakai et al., 2012;
Walker et al., 1987). To exclude the possible contribution of external Ca^{2+} to the responses,
we performed the experiments in Ca^{2+}-free media. In response to sequential cIP_3 release
5 min. apart, control eggs displayed corresponding Ca^{2+} rises that occasionally

transitioned into short-lived oscillations (**Fig. 3A**). The addition of TPEN after the third
cIP$_3$ release prevented the subsequent Ca^{2+} response and prematurely terminated the in-
progress Ca^{2+} rises (**Fig. 3B and inset)**. Pre-incubation of eggs with TPEN precluded
cIP$_3$-induced Ca^{2+} release, even after 5 sec. of UV exposure (**Fig. 3C**). The addition of
excess ZnSO$_4$ (100 µM) overcame TPEN's inhibitory effects, but only if added before
(Fig. 3E) and not after the addition of TPEN **(Fig. 3D)**. Similar concentrations of MgCl$_2$
or CaCl$_2$ failed to reverse TPEN effects **(Fig. 3F, G)**. Together, the results show that Zn^{2+}
is required for IP$_3$R1-mediated Ca^{2+} release downstream of IP$_3$ production, appearing to
interfere with receptor gating, as suggested by TPEN's rapid termination of in-progress
Ca^{2+} rises and ongoing oscillations.
ERp44 is an ER luminal protein of the thioredoxin family that interacts with the
IP$_3$R1, reportedly inhibiting its ability to mediate Ca^{2+} release (Higo et al., 2005). The
localization of ERp44 in the ER-Golgi intermediate compartment of somatic cells
correlates with Zn^{2+}'s availability and changes dramatically after TPEN treatment (Higo
et al., 2005; Watanabe et al., 2019). To rule out the possibility that TPEN suppresses the
function of IP$_3$R1 by modifying the subcellular distribution of ERp44, we overexpressed
ERp44 by injecting HA tagged-*Erp44* mRNA into MII eggs and monitored the effect on
Ca^{2+} release. TPEN did not alter the localization of ERp44 (**Supplementary Fig. 1A**),
and overexpression of ERp44 modified neither the Ca^{2+} oscillations induced by agonists
(**Supplementary Fig. 1B**) nor the effectiveness of TPEN to block them (data not shown).
Thus, TPEN and Zn^{2+} deficiency most likely inhibits Ca^{2+} release by directly interfering
with IP$_3$R1 function rather than modifying this particular regulator.
225
Zn^{2+} depletion diminishes the ER Ca^{2+} leak and increases Ca^{2+} store content.
Our above cIP$_3$ results that TPEN inhibited IP$_3$R1-mediated Ca^{2+} release and
interrupted in-progress Ca^{2+} rises despite the presence of high levels of environmental IP$_3$
suggest its actions are probably independent of IP$_3$ binding, agreeing with an earlier report
showing that TPEN did not modify IP$_3$'s affinity for the IP$_3$R (Richardson and Taylor,
1993). Additionally, the presence of a Zn^{2+}-binding motif near the C-term cytoplasmic
domain of the IP$_3$R1's channel, which is known to influence agonist-induced IP$_3$R1 gating
(Fan et al., 2015), led us to posit and examine that Zn^{2+} deficiency may be disturbing Ca^{2+}
release to the cytosol and out of the ER. To probe this possibility, we queried if pre-
treatment with TPEN inhibited Ca^{2+} release through IP$_3$R1. We first used Thapsigargin
(Tg), a Sarcoplasmic/ER Ca^{2+} ATPase pump inhibitor (Thastrup et al., 1990) that unmasks
a constitutive Ca^{2+} leak out of the ER (Lemos et al., 2021); in eggs, we have demonstrated
it is mediated at least in part by IP$_3$R1 (Wakai et al., 2019). Treatment with TPEN for 15

min. slowed the Tg-induced Ca^{2+} leak into the cytosol, resulting in delayed and lowered

amplitude Ca^{2+} responses (**Fig. 4A;** $P<0.05$). To test whether the reduced response to Tg

means that TPEN left a temporarily increased Ca^{2+} content in the ER after it, we added

the Ca^{2+} ionophore ionomycin (Io), which empties all stores independently of IP$_3$Rs. Io-

induced Ca^{2+} responses were 3.3-fold greater in TPEN-treated cells, supporting the view

that TPEN interferes with the ER Ca^{2+} leak **(Fig. 4A;** $P < 0.05$**)**. We further evaluated this

concept using *in vitro* aged eggs that often display reduced Ca^{2+} store content than freshly

collected counterparts (Abbott et al., 1998). After culturing eggs in the presence or

absence of TPEN for 2-hr., we added Io during Ca^{2+} monitoring, which in TPEN-treated

eggs induced bigger Ca^{2+} rises than in control eggs **(Fig. 4B;** $P <0.05$**)**. We confirmed that

this effect was independent of IP$_3$R1 degradation because TPEN did not change IP$_3$R1

reactivity in unfertilized eggs (**Fig. 4C**; $P <0.05$).

Next, we used the genetically encoded FRET sensor D1ER (Palmer et al., 2004) to

assess the TPEN's effect on the ER's relative Ca^{2+} levels changes following the additions

of Tg or Ach. TPEN was added 10 min. before 10 μM Tg or 50 μM Ach, and we

simultaneously monitored changes in cytosolic and intra-ER Ca^{2+} (**Fig. 4D, E**). For the

first three min., the Tg-induced decrease in Ca^{2+}-ER was similar between groups.

However, while the drop in Ca^{2+} content continued in control eggs, in TPEN-treated eggs,

it came to an abrupt halt, generating profound differences between the two groups (**Fig.**

4D; $P <0.05$). TPEN had even more pronounced effects following the addition of Ach,

leading to a reduced- and prematurely terminated- Ca^{2+} release from the ER in treated

eggs (**Fig. 4E;** $P <0.05$).

Lastly, we sought a cellular model where low labile Zn^{2+} occurred without

pharmacology. To this end, we examined a genetic model where the two non-selective

plasma membrane channels that could influx Zn^{2+} in maturing oocytes have been deleted

(Bernhardt et al., 2017; Carvacho et al., 2016, 2013), namely, the transient receptor

potential melastatin-7 (TRPM7) and TRP vanilloid 3 (TRPV3), both members of the TRP

superfamily of channels (Wu et al., 2010). We found that eggs from double knockout

females (dKOs) had lower labile Zn^{2+} levels (**Fig. 4F**), and the addition of Tg revealed an

expanded Ca^{2+} store content in these eggs vs. control WT eggs (**Fig. 4G**). Remarkably, in

dKO eggs, the Ca^{2+} rise induced by Tg showed a shoulder or inflection point before the

peak delaying the time to peak (**Fig. 4G**, **inset**; $P <0.001$). These results in dKO eggs

show a changed dynamic of the Tg-induced Ca^{2+} release, suggesting that Zn^{2+} deficient

levels modify ER Ca^{2+} release independently of chelators.

Ca^{2+} oscillations in eggs occur within a window of Zn^{2+} concentrations.

We next examined if resupplying Zn^{2+} could restart the Ca^{2+} oscillations terminated by Zn^{2+} depletion. Zn pyrithione (ZnPT) rapidly increases cellular Zn^{2+} upon extracellular addition (Barnett et al., 1977; Robinson, 1964). Dose titration studies and imaging fluorimetry revealed that 0.01 μM ZnPT caused subtle and protracted increases in Zn^{2+} levels, whereas 0.1 μM ZnPT caused rapid increases in eggs' Zn^{2+} baseline (**Fig. 5A**). We induced detectable Ca^{2+} oscillations by injection of m*Plcz* mRNA followed by 50 μM TPEN (**Fig. 5B**), which terminated them. After 30 min, we added 0.1 μM ZnPT, and within 15 min. the oscillations restarted in most TPEN-treated eggs (**Fig. 5C**). We repeated this approach using Thimerosal (**Fig. 5D, E**). Adding 0.1 μM ZnPT did not restore the Ca^{2+} oscillations retrained by TPEN, but 0.5 μM ZnPT did so (**Fig. 5E**). These results demonstrate that Zn^{2+} plays a pivotal, enabling role in the generation of Ca^{2+} oscillations in mouse eggs.

Excessive intracellular Zn²⁺ inhibits Ca²⁺ oscillations.

Zn^{2+} is necessary for diverse cellular functions, consistent with numerous amino acids and proteins capable of binding Zn^{2+} within specific and physiological ranges (Pace and Weerapana, 2014). Excessive Zn^{2+}, however, can cause detrimental effects on cells and organisms (Broun et al., 1990; Hara et al., 2022; Sikora and Ouagazzal, 2021). Consistent with the deleterious effects of Zn^{2+}, a previous study showed that high concentrations of ZnPT, ~50 μM, prevented $SrCl_2$-induced egg activation and initiation of development (Bernhardt et al., 2012; Kim et al., 2011). We examined how ZnPT and excessive Zn^{2+} levels influence Ca^{2+} oscillations. Our conditions revealed that pre-incubation or continuous exposure to 0.1 μM or 1.0 μM ZnPT delayed or prevented egg activation induced by m*Plcz* mRNA injection (**Supplementary Fig. 2**). We used these ZnPT concentrations to add it into ongoing oscillations induced by ICSI and monitored the succeeding Ca^{2+} responses. The addition of 0.05 to 10 μM ZnPT caused an immediate elevation of the basal levels of Fura-2 and termination of the Ca^{2+} oscillations (**Fig. 6A-D**). m*Plcz* mRNA-initiated Ca^{2+} responses were also interrupted by adding 0.1 μM ZnPT, whereas untreated eggs continued oscillating (**Fig. 6E, F**). ZnPT also inhibited IP₃R1-mediated Ca^{2+} release triggered by cIP₃, suggesting that excessive Zn^{2+} directly inhibits IP₃R1 function (**Fig. 6G**).

The increased basal ratios of Fura-2 caused by the addition of ZnPT are unlikely to represent changes in basal Ca^{2+} levels because they would have likely caused some cellular responses such as the release of the second polar body, egg fragmentation, or cell death, neither of which happened. It might reflect, instead, Fura-2's ability to report changes in Zn^{2+} levels, which was the case because the addition of TPEN lowered the

Fura-2 ratios without restarting the Ca^{2+} oscillations (**Fig. 6F**). To ensure the impact of

ZnPT abolishing Ca^{2+} oscillations was not an imaging artifact obscuring ongoing rises,

we simultaneously monitored cytoplasmic and ER Ca^{2+} levels with Rhod-2 and D1ER,

respectively. This approach allowed synchronously observing opposite Ca^{2+} changes in

the two compartments. In uninjected eggs, the fluorescent values remained unchanged,

and in m*Plcz* mRNA-injected eggs, the reporters' signals displayed the expected opposite

changes (**Fig. 6H, I**). The addition of ZnPT rapidly increased Rhod-2 signals in uninjected

and oscillating eggs, unlike D1ER's, suggesting it cannot detect changes in Zn^{2+} levels to

this extent despite faithfully reporting Ca^{2+} changes. In oscillating eggs following ZnPT

addition, D1ER progressively showed fewer and lower amplitude changes, consistent

with the diminishing and eventual termination of the Ca^{2+} oscillations. Noteworthy, the

basal fluorescent ratio of D1ER in these eggs remained unchanged after ZnPT, further

demonstrating its unresponsiveness to Zn^{2+} changes of this magnitude. We confirmed that

both reporters were still in working order, as the addition of Io triggered Ca^{2+} changes

detected by both reporters (**Fig. 6H, I**).

Discussion

The present study demonstrates that appropriate levels of labile Zn^{2+} are essential

for initiating and maintaining IP3R1-mediated Ca^{2+} oscillations in mouse eggs regardless

of the initiating stimuli. Both deficient and excessive Zn^{2+} compromise IP3R1 sensitivity,

diminishing and mostly terminating Ca^{2+} oscillations. The results demonstrate that IP3R1

and Zn^{2+} act in concert to modulate Ca^{2+} signals revealing previously unexplored

crosstalk between these ions at fertilization (**Fig. 7**).

Zn^{2+} is an essential micronutrient for living organisms (Kaur et al., 2014) and is

required for various cellular functions, such as proliferation, transcription, and

metabolism (Lo et al., 2020; Maret and Li, 2009; Yamasaki et al., 2007). Studies using

Zn^{2+} chelators have uncovered what appears to be a cell-specific, narrow window of Zn^{2+}

concentrations needed for cellular proliferation and survival (Carraway and Dobner,

2012; Lo et al., 2020). Further, TPEN appeared especially harmful, and in a few cell lines,

even low doses provoked oxidative stress, DNA fragmentation, and apoptosis (Mendivil-

Perez et al., 2012). We show here that none of the Zn^{2+} chelators, permeable or

impermeable, affected cell viability within our experimental observations, confirming

findings from previous studies that employed high concentrations of TPEN to interrupt

the Ca^{2+} oscillations (Lawrence et al., 1998) or inducing egg activation of mouse eggs

(Suzuki et al., 2010b). Our data demonstrating that ~2.5 mM is the threshold

concentration of TPEN in eggs that first causes noticeable changes in basal Zn^{2+}, as

revealed by FluoZin, is consistent with the ~2 to 5-mM Zn^{2+} concentrations in most

culture media without serum supplementation (Lo et al., 2020), and with the ~100 pM

basal Zn^{2+} in cells (Qin et al., 2011). Lastly, the effects on Ca^{2+} release observed here with

TPEN and other chelators were due to the chelation of Zn^{2+}, as pretreatment with $ZnSO_4$

but not with equal or greater concentrations of $MgCl_2$ or $CaCl_2$ rescued the inhibition of

the responses, which is consistent with results by others (Kim et al., 2010; Lawrence et

al., 1998).

To identify how Zn^{2+} deficiency inhibits Ca^{2+} release in eggs, we induced Ca^{2+}

oscillations using various stimuli and tested the effectiveness of membrane-permeable

and impermeable chelators to abrogate them. Chelation of extracellular Zn^{2+} failed to

terminate the Ca^{2+} responses, whereas membrane-permeable chelators did, pointing to

intracellular labile Zn^{2+} levels as essential for Ca^{2+} release. All agonists used here were

susceptible to inhibition by TPEN, whether their activities depended on IP_3 production or

allosterically induced receptor function, although the effective TPEN concentrations

varied across stimuli. Some agents, such as m*Plcz* mRNA or thimerosal, required higher

concentrations than $SrCl_2$, Ach, or cIP_3. The reason underlying the different agonists'

sensitivities to TPEN will require additional research, but the persistence of IP_3

production or change in IP$_3$R1 structure needed to induce channel gating might explain

it. However, the universal abrogation of Ca^{2+} oscillations by TPEN supports the view

drawn from cryo-EM-derived IP$_3$R1 models that signaling molecules can allosterically

induce channel gating from different starting positions in the channel by mechanically

coupling the binding effect to the ion-conducting pore in the C-terminal end of IP$_3$R (Fan

et al., 2015). The cytosolic C-terminal domain of each IP$_3$R1 subunit is alongside the IP$_3$-

binding domain of another subunit and, therefore, well positioned to sense IP_3 binding

and induce channel gating (Fan et al., 2015). Within each subunit, the LNK domain, which

contains a Zn^{2+}-finger motif (Fan et al., 2015), connects the opposite domains of the

molecule. Although there are no reports regarding the regulation of IP$_3$R1 sensitivity by

Zn^{2+}, such evidence exists for RyRs (Woodier et al., 2015), which also display a conserved

Zn^{2+}-finger motif (des Georges et al., 2016). Lastly, mutations of the two Cys or two His

residues of this motif, without exception, resulted in inhibition or inactivation of the IP$_3$R1

channel (Bhanumathy et al., 2012; Uchida et al., 2003). These results are consistent with

the view that the C-terminal end of IP$_3$Rs plays a dominant role in channel gating

(Bhanumathy et al., 2012; Uchida et al., 2003). We propose that TPEN inhibits Ca^{2+}

oscillations in mouse eggs because chelating Zn^{2+} interferes with the function of the LNK

domain and its Zn^{2+}-finger motif proposed role on the mechanical coupling induced by

agonist binding to the receptor that propagates to the pore-forming region and required to

gate the channel's ion-pore (Fan et al., 2022, 2015).

In support of this possibility, TPEN-induced Zn^{2+} deficient conditions altered the

Ca^{2+}-releasing kinetics in resting eggs or after fertilization. Tg increases intracellular Ca^{2+}

by inhibiting the SERCA pump (Thastrup et al., 1990) and preventing the reuptake into

the ER of the ebbing Ca^{2+} during the basal leak. Our previous studies showed that the

downregulation of IP$_3$R1 diminishes the leak, suggesting it occurs through IP$_3$R1 (Wakai

and Fissore, 2019). Consistent with this view, TPEN pre-treatment delayed the Ca^{2+}

response induced by Tg, implying that Zn^{2+} deficiency hinders Ca^{2+} release through IP$_3$R1.

An expected consequence would be increased Ca^{2+} content in the ER after Tg. Io that

mobilizes Ca^{2+} independently of IP$_3$Rs (Toeplitz et al., 1979) induced enhanced responses

in TPEN-treated eggs vs. controls, confirming the accumulation of Ca^{2+}- ER in Zn^{2+}

deficient conditions. We demonstrated that this accumulation is due to hindered emptying

of the Ca^{2+} ER evoked by agonists in Zn^{2+}-deficient environments resulting in reduced

cytosolic Ca^{2+} increases, as IP$_3$R1 is the pivotal intermediary channel between these

compartments. Noteworthy, the initial phase of the Tg-induced Ca^{2+} release out of the ER

did not appear modified by TPEN, as if it was mediated by a Zn^{2+}-insensitive Ca^{2+}

channel(s)/transporter, contrasting with Ach-induced emptying that was abrogated from

the outset. Remarkably, independently of Zn^{2+} chelators, emptying of Ca^{2+} ER was

modified in a genetic model of Zn^{2+}-deficient oocytes lacking two TRP channels,

confirming the impact of Zn^{2+} on Ca^{2+} release. It is worth noting that TPEN did not reduce

but increased the mass of IP$_3$R1, which might result in the inhibition of Zn^{2+}-dependent

ubiquitin ligase Ubc7 by the Zn-deficient conditions (Webster et al., 2003). We cannot

rule out that these conditions may undermine other conformational changes required to

trigger IP$_3$R1 degradation, thereby favoring the accumulation of IP$_3$R1.

Despite accruing Zn^{2+} during oocyte maturation, fertilization witnesses a

necessary Zn^{2+} release into the external milieu, known as "Zn^{2+} sparks" (Kim et al., 2011;

Que et al., 2019, 2015). This release of Zn^{2+} is a conserved event in fertilization across

species and is associated with several biological functions, including those related to

fending off polyspermy (Kim et al., 2011; Que et al., 2019; Wozniak et al., 2020). The

concomitant decrease in Zn^{2+} facilitates the resumption of the cell cycle and exit from the

MII stage (Kim et al., 2011). Congruent with this observation, artificial manipulation that

maintains high Zn^{2+} levels prevent egg activation (Kim et al., 2011), whereas lowering

Zn^{2+} with chelators leads to egg activation without Ca^{2+} mobilization (Suzuki et al.,

2010b). As posed by others, these results suggest that meiosis completion and the early

stages of fertilization unfold within a narrow window of permissible Zn^{2+} (Kim et al.,

2011, 2010). Here, we extend this concept and show that IP$_3$R1 function and the Ca^{2+}

oscillations in mouse eggs require this optimal level of labile Zn^{2+} because the Ca^{2+}
responses interrupted by TPEN-induced Zn^{2+}-insufficiency are rescued by restoring Zn^{2+}
levels with ZnPT. Furthermore, unopposed increases in Zn^{2+} by exposure to ZnPT
abrogated fertilization-initiated Ca^{2+} oscillations and prevented the expected egg
activation events. It is unclear how excess Zn^{2+} disturbs the function of IP$_3$R1.
Nevertheless, IP$_3$R1s have multiple cysteines whose oxidation enhances the receptor
sensitivity to IP$_3$ (Joseph et al., 2018), and it is possible that excessive Zn^{2+} aberrantly
modifies them, disturbing IP$_3$R1 structure and function. These results reveal a close
association between the Zn^{2+} levels controlling meiotic transitions and the Ca^{2+} release
necessary for egg activation, placing the IP$_3$R1 at the center of the crosstalk of these two
divalent cations.

Abrupt Zn^{2+} changes have emerged as critical signals for meiotic and mitotic
transitions in oocytes, eggs, embryos, and somatic cells (Kim et al., 2011, 2010; Lo et al.,
2020). Fertilization relies on prototypical Ca^{2+} rises and oscillations, and Zn^{2+} sparks are
an egg activation event downstream of this Ca^{2+} release, establishing a functional
association between these two divalent cations that continues to grow (Kim et al., 2011).
Here, we show that, in addition, these cations actively crosstalk during fertilization and
that the fertilization-induced Ca^{2+} oscillations rely on optimized IP$_3$R1 function
underpinned by ideal Zn^{2+} levels set during oocyte maturation. Future studies should
explore if artificial alteration of Zn^{2+} levels can extend the fertile lifespan of eggs,
improve developmental competence, or for developing non-hormonal methods of
contraception.

Materials and Methods

Key resources table

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Genetic reagent (*Mus musculus*)	CD1	Charles River	022	
Genetic reagent (*Mus musculus*)	C57BL/6J	JAX	JAX: 000664	
Genetic reagent (*Mus musculus*)	*Trpm7*-floxed	A generous gift from Dr. Carmen P. Williams (NIEHS) (PMID: 30322909)		C57BL6/J and 129s4/SvJae mixed background
Genetic reagent (*Mus musculus*)	*Gdf9-cre*	JAX	JAX: 011062	
Genetic reagent (*Mus musculus*)	*Trpv3⁻/⁻*	A generous gift from Dr H. Xu (PMID: 20403327)		C57BL/6J and 129/SvEv mixed background
Biological sample (mouse oocyte)	*Mus musculus*	this paper		Eggs at the metaphase of the second meiosis
Biological sample (mouse sperm)	*Mus musculus*	this paper		Matured sperm from cauda epididymis
Recombinant DNA reagent	pcDNA6-mouse *Plcz1-venus* (plasmid used as a template for mRNA synthesis)	Published in previous Fissore lab paper PMID: 34313315. Mouse *Plcz1*		mouse *Plcz1* mRNA was fused with Venus and

		sequence was a generous gift from Dr. Kiyoko Fukami (PMID:1802889 8)		inserted in pcDNA6 vector
Recombinant DNA reagent	pcDNA6-CALR-D1ER-KDEL (plasmid used as a template for mRNA synthesis)	Published in previous Fissore lab paper PMID: 24101727. Original D1ER vector was a generous gift from Dr. Roger Y Tsien (PMID: 15585581)		FRET construct D1ER was inserted between ER-targeting sequence of calreticuli n and KDEL ER retention signal in pcDNA6 vector
Recombinant DNA reagent	pcDNA6-human *ERp44-HA* (plasmid used as a template for mRNA synthesis)	This paper. Original human ERp44 sequence was a generous gift from Dr. Roberto Sitia (PMID: 11847130)		human *ERp44* mRNA fused with HA in pcDNA6/ Myc-His B vector
Antibody	Monoclonal HA (Mouse monoclonal)	Roche	115818160 01	Dilution: 1:200
Antibody	Polyclonal IP$_3$R1 (Rabbit polyclonal)	(Parys et al., 1995)		Dilution: 1:1000
Antibody	Monoclonal α-tubulin (Mouse monoclonal)	Sigma-Aldrich	T-9026	Dilution: 1:1000
Antibody	Alexa Fluor 488 (goat anti mouse)	Invitrogen	Invitrogen: A32723	Dilution: 1:400
Commercial assay or kit	T7 mMESSAGE mMACHINE Kit	Invitrogen	Invitrogen: AM1344	Used for *in vitro* mRNA synthesis
Commercial assay or kit	Poly(A) Tailing Kit	Invitrogen	Invitrogen: AM1350	Used for poly (A)

				tailing of synthesized mRNA
Chemical compound, drug	Hyaluronidase from bovine testes	Sigma-Aldrich	H3506	
Chemical compound, drug	3-Isobutyl-1-methylxanthine (IBMX)	Sigma-Aldrich	I5879	
Chemical compound, drug	Polyvinylpyrrolidone (PVP) (average molecular weight: 360,000)	Sigma-Aldrich	PVP360	Used for mRNA microinjection and ICSI
Chemical compound, drug	N,N, N′,N′-Tetrakis (2-pyridylmethyl) ethylenediamine (TPEN)	Sigma-Aldrich	P4413	Prepared in DMSO and kept at –20 °C until use
Chemical compound, drug	Zinc Pyrithione (ZnPT)	Sigma-Aldrich	PHR1401	Prepared in DMSO and kept at –20 °C until use
Chemical compound, drug	Strontium chloride hexahydrate ($SrCl_2$)	Sigma-Aldrich	255521	Freshly dissolved in water on the day of experiment
Chemical compound, drug	Calcium chloride dihydrate ($CaCl_2$)	Sigma-Aldrich	C3881	Freshly dissolved in water on the day of experiment
Chemical compound, drug	Magnesium chloride hexahydrate ($MgCl_2$)	Sigma-Aldrich	M2393	Freshly dissolved in water on the day of experiment
Chemical compound,	Zinc sulfate monohydrate ($ZnSO_4$)	Acros Organics	389802500	Freshly dissolved

drug				in water on the day of experiment
Chemical compound, drug	Ethylenediaminetetraacetic acid sodium dihydrate (EDTA)	LabChem	LC137501	Prepared as 0.5M aqueous solution with pH 8.0 adjusted by NaOH
Chemical compound, drug	Diethylenetriaminepentaacetic acid (DTPA)	Sigma-Aldrich	D6518	
Chemical compound, drug	Tris (2-pyridylmethyl) amine (TPA)	Santa Cruz	sc-477037	
Chemical compound, drug	Dimethyl sulphoxide (DMSO)	Sigma-Aldrich	D8418	Used as a solvent
Chemical compound, drug	Acetylcholine chloride	Sigma-Aldrich	A6625	
Chemical compound, drug	Thimerosal	Sigma-Aldrich	T5125	Freshly dissolved in water on the day of experiment and kept on ice until use
Chemical compound, drug	Ionomycin calcium salt	Tocris	1704	Working concentration: 2.5 μM
Chemical compound, drug	Thapsigargin	Calbiochem	#586500	Working concentration: 10 μM
Other	Pluronic F-127 (20% solution in DMSO) (Pluronic acid)	Invitrogen	P3000MP	
Other	Fura-2 AM	Invitrogen	F1221	Used at

				1.25 mM in TL-HEPES containing 0.02% Pluronic acid
Other	FluoZin-3 AM	Invitrogen	F24195	Used at 1.25 mM in TL-HEPES containing 0.02% Pluronic acid
Other	Fluo-4 AM	Invitrogen	F14201	Used at 1.25 mM in TL-HEPES containing 0.02% Pluronic acid
Other	Rhod2-AM	Invitrogen	R1244	Used at 2.2 mM in TL-HEPES containing 0.02% Pluronic acid.
Other	ci-IP3/ PM	Tocris	6210	Dissolved in DMSO and kept at –20 °C. Before use, the stock was diluted with water to make a final concentration of 0.25 mM.
Other	Pme1	New England	R0560S	Used to

		BioLabs	linearize pcDNA6 vectors for mRNA synthesis
Software, algorithm	Prism	GraphPad Software	Version 5.01

N,N,N′,N′-tetrakis (2-pyridinylmethyl)-1,2-ethylenediamine (TPEN) and Zinc pyrithione
(ZnPT) were dissolved in dimethyl sulfoxide (DMSO) at 10 mM and stored at -20°C until
use. SrCl$_2$, CaCl$_2$, ZnSO$_4$, and MgCl$_2$ were freshly dissolved with double-sterile water at
1M and diluted with the monitoring media just before use. Ethylenediaminetetraacetic
acid (EDTA) and diethylenetriaminepentaacetic acid (DTPA) were reconstituted with
double-sterile water at 0.5M and 10 mM, respectively, and the pH was adjusted to 8.0.
Tris(2-pyridylmethyl) amine (TPA) was diluted in DMSO at 100 mM and stored at -20°C
until use. Acetylcholine chloride and Thimerosal were dissolved in double-sterile water
at 550 mM and 100 mM, respectively. Acetylcholine was stored at -20°C until use,
whereas Thimerosal was made fresh in each experiment.

Mice
The University of Massachusetts Institutional Animal Care and Use Committee (IACUC)
approved all animal experiments and protocols. *Trpm7*-floxed (*Trpm7*$^{fl/fl}$) *Gdf9-Cre* and
Trpv3$^{-/-}$ mice were bred at our facility. *Trpm7*$^{fl/fl}$ mice were crossed with *Trpv3*$^{-/-}$ to
generate *Trpm7*$^{fl/fl}$; *Trpv3*$^{-/-}$ mouse line. Female *Trpm7*$^{fl/fl}$; *Trpv3*$^{-/-}$ mice were crossed
with *Trpm7*$^{fl/fl}$; *Trpv3*$^{-/-}$; *Gdf9-cre* male to generate females null for *Trpv3* and with
oocyte-specific deletion for *Trpm7*. Ear clips from offspring were collected prior to
weaning, and confirmation of genotype was performed after most experiments.

Egg Collection
All gamete handling procedures are as previously reported by us (Wakai and Fissore,
2019). MII eggs were collected from the ampulla of 6- to 8-week-old female mice.
Females were superovulated via intraperitoneal injections of 5 IU pregnant mare serum
gonadotropin (PMSG, Sigma, St. Louis, MO) and 5 IU human chorionic gonadotropin
(hCG, sigma) at 48hr. interval. Cumulus-oocyte-complexes (COCs) were obtained 13.5
hr. post-hCG injection by tearing the ampulla using forceps and needles in TL-HEPES
medium. COCs were treated with 0.26% (w/v) of hyaluronidase at room temperature (RT)
for 5 min to remove cumulus cells.

Intracytoplasmic sperm injection (ICSI)

ICSI was performed as previously reported by us (Kurokawa and Fissore, 2003) using described setup and micromanipulators (Narishige, Japan). Sperm from C57BL/6 or CD1 male mice (7-12 weeks old) were collected from the cauda epididymis in TL-HEPES medium, washed several times, heads separated from tails by sonication (XL2020; Heat Systems Inc., USA) for 5 s at 4°C. The sperm lysate was washed in TL-HEPES and diluted with 12% polyvinylpyrrolidone (PVP, MW = 360 kDa) to a final PVP concentration of 6%. A piezo micropipette-driving unit was used to deliver the sperm into the ooplasm (Primetech, Ibaraki, Japan); a few piezo-pulses were applied to puncture the eggs' plasma membrane following penetration of the zona pellucida. After ICSI, eggs were either used for Ca^{2+} monitoring or cultured in KSOM to evaluate activation and development at 36.5°C in a humidified atmosphere containing 5% CO_2.

Preparation and microinjection of mRNA

pcDNA6-m*Plcζ-mEGFP*, pcDNA6-CALR-D1ER-KDEL, and pcDNA6-*humanERp44-HA* were linearized with the restriction enzyme PmeI and *in vitro* transcribed using the T7 mMESSAGE mMACHINE Kit following procedures previously used in our laboratory (Ardestani et al., 2020). A poly(A) tail was added to the *in vitro* synthesized RNA (mRNA) using Tailing Kit followed by quantification and dilution to 0.5 μg/μL in nuclease-free water and stored at -80°C until use. Before microinjection, m*Plcζ*, D1ER, and *ERp44* mRNA were diluted to 0.01, 1.0, and 0.5 μg/μL, respectively, in nuclease-free water, heated at 95°C for 3 min followed by centrifugation at 13400×*g* for 10 min at 4°C. Cytoplasm injection of mRNA was performed under microscopy equipped with micromanipulators (Narishige, Japan). The zona pellucida and the plasma membrane of MII eggs were penetrated by applying small pulses generated by the piezo micromanipulator (Primetech, Ibaraki, Japan). The preparation of the injection pipette was as for ICSI (Kurokawa and Fissore, 2003), but the diameter of the tip was ~1 μm.

Ca^{2+} imaging

Before Ca^{2+} imaging, eggs were incubated in TL-HEPES containing 1.25 μM Fura2-AM, 1.25 μM FluoZin3-AM, or 2.2 μM Rhod2-AM and 0.02% Pluronic acid for 20 min at room temperature and then washed. The fluorescent probe-loaded eggs were allowed to attach to the bottom of the glass dish (Mat-Tek Corp., Ashland, MA). Eggs were monitored simultaneously using an inverted microscope (Nikon, Melville, NY) outfitted for fluorescence measurements. Fura-2 AM, FluoZin3-AM, and Rhod2-AM fluorescence

were excited with 340 nm and 380 nm, 480 nm, and 550 nm wavelengths, respectively,
every 20 sec, for different intervals according to the experimental design and as
previously performed in the laboratory. The illumination was provided by a 75-W Xenon
arc lamp and controlled by a filter wheel (Ludl Electronic Products Ltd., Hawthorne, NY).
The emitted light above 510 nm was collected by a cooled Photometrics SenSys CCD
camera (Roper Scientific, Tucson, AZ). Nikon Element software coordinated the filter
wheel and data acquisition. The acquired data were saved and analyzed using Microsoft
Excel and GraphPad using Prism software (Ardestani et al., 2020). For Figures 1A, 4A-
C, 5A, and 6H-I, values obtained from FluoZin3-AM, Fura2-AM, or Rhod2-AM
recordings were divided by the average of the first five recordings for each treatment that
was used as the F_0.
To estimate relative changes in Ca^{2+}-ER, emission ratio imaging of the D1ER (YFP/CFP)
was performed using a CFP excitation filter, dichroic beamsplitter, CFP and YFP emission
filters (Chroma technology, Rockingham, VT; ET436/20X, 89007bs, ET480/40m, and
ET535/30m). To measure Ca^{2+}-ER and cytosolic Ca^{2+} simultaneously, eggs that had been
injected with D1ER were loaded with Rhod-2AM, and CFP, YFP, and Rhod-2 intensities
were collected every 20 seconds.
526
Caged IP$_3$
Caged-IP$_3$/PM (cIP$_3$) was reconstituted in DMSO and stored at -20°C until use. Before
injection, cIP$_3$ stock was diluted to 0.25 mM with water and microinjected as above. After
incubation in KSOM media at 37°C for 1-hr., the injected eggs were loaded with the
fluorophore, 1.25 μM Fluo4-AM, and 0.02% Pluronic acid and handled as above for Fura-
2 AM. The release of cIP$_3$ was accomplished by photolysis using 0.5 to 5-sec pulses at
360 nm wavelengths. Ca^{2+} imaging was as above, but Fluo4 was excited at 488 nm
wavelength and emitted light above 510 nm collected as above.
535
Western blot analysis
Cell lysates from 20-50 mouse eggs were prepared by adding 2X- Laemmli sample buffer.
Proteins were separated on 5% SDS-PAGE gels and transferred to PVDF membranes
(Millipore, Bedford, MA). After blocking with 5% fat-free milk + TBS, membranes were
probed with the rabbit polyclonal antibody specific to IP$_3$R1 (1:1000; a generous gift from
Dr. Jan Parys, Katholieke Universiteit, Leuven, Belgium; Parys et al., 1995). Goat anti-
rabbit antibody conjugated to horseradish peroxidase (HRP) was used as a secondary
antibody (1:5000; Goat anti-Rabbit IgG (H+L) Cross-Adsorbed Secondary Antibody,
HRP; Invitrogen, Waltham, Ma). For detection of chemiluminescence, membranes were

developed using ECL Prime (Sigma) and exposed for 1–3 min to maximum sensitivity
film (VWR, Radnor, PA). Broad-range pre-stained SDS–PAGE molecular weight
markers (Bio-Rad, Hercules, CA) were run in parallel to estimate the molecular weight
of the immunoreactive bands. The same membranes were stripped at 50°C for 30 min
(62.5 mM Tris, 2% SDS, and 100 mM 2-beta mercaptoethanol) and re-probed with anti-
α-tubulin monoclonal antibody (1:1000).
551

Immunostaining and confocal microscopy
Immunostaining was performed according to our previous study (Akizawa et al., 2021).
After incubation with or without TPEN, MII eggs were fixed with 4% (w/v)
paraformaldehyde in house-made phosphate-buffered saline (PBS) for 20 min at room
temperature and then permeabilized for 60 min with 0.2% (v/v) Triton X-100 in PBS.
Next, the eggs were blocked for 45 min with a blocking buffer containing 0.2% (w/v)
skim milk, 2% (v/v) fetal bovine serum, 1% (w/v) bovine serum albumin, 0.1% (v/v)
TritonX-100, 0.75% (w/v) glycine in PBS. Eggs were incubated overnight at 4°C with
mouse anti-HA antibody (1:200) diluted in blocking buffer. Eggs were washed in blocking
buffer 3X for 10 min, followed by incubation at room temperature for 30 min with a
secondary antibody, Alexa Fluor 488 goat anti-mouse IgG (H + L) (1:400) diluted in
blocking buffer. Fluorescence signals were visualized using a laser-scanning confocal
microscope (Nikon A1 Resonant Confocal with six-color TIRF) fitted with a 63×, 1.4 NA
oil-immersion objective lens.
566

Statistical analysis
Comparisons for statistical significance of experimental values between treatments and
experiments were performed in three or more experiments performed on different batches
of eggs in most studies. Given the number of eggs needed, WB studies were repeated
twice. Prism-GraphPad software was used to perform the statistical comparisons that
include unpaired Student's t-tests, Fisher's exact test, and One-way ANOVA followed by
Tukey's multiple comparisons, as applicable, and the production of graphs to display the
data. All data are presented as mean±s.d. Differences were considered significant at $P <$
0.05.
576

577 **Acknowledgments**
578 We thank Ms. Changli He for technical support and Dr. James Chambers for support with
579 confocal microscopy support. We thank all members of the Fissore lab for useful
580 discussions and suggestions. We thank Jan B. Parys, K.U. Leuven, Belgium, for initial

581 discussions and advice.

Competing interests

584 The authors declare no competing or financial interests.

586 **Additional information**

587 **Funding Sources**

Funder	Grant reference number	Author
National Institute of Health	RO1 HD092499	Rafael A. Fissore
National Institute of Food and Agriculture	2021-06893	MG. Gervasi, R.A. Fissore, P.E. Visconti
Japan Society for the Promotion of Science Overseas Research Fellowship	Postdoctoral fellowship	Hiroki Akizawa
The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.		

Author contributions

Hiroki Akizawa, Data curation, Formal analysis, Validation, Investigation, Visualization,

Writing—original draft, Writing—review and editing; Emily Lopes, Data curation,

Formal analysis, Validation; Rafael A Fissore, Conceptualization, Formal analysis,

Supervision, Funding acquisition, Methodology, Writing—original draft, Project

administration, Writing—review and editing

References

Abbott AL, Xu Z, Kopf GS, Ducibella T, Schultz RM. 1998. In vitro culture retards

spontaneous activation of cell cycle progression and cortical granule exocytosis that

normally occur in vivo unfertilized mouse eggs. *Biol Reprod* **59**:1515–1521.

doi:10.1095/biolreprod59.6.1515

Ajduk A, Małagocki A, Maleszewski M. 2008. Cytoplasmic maturation of mammalian

oocytes: Development of a mechanism responsible for sperm-induced Ca2+

oscillations. *Reprod Biol* **8**:3–22. doi:10.1016/S1642-431X(12)60001-1

Akizawa H, Saito S, Kohri N, Furukawa E, Hayashi Y, Bai H, Nagano M, Yanagawa Y,

Tsukahara H, Takahashi M, Kagawa S, Kawahara-Miki R, Kobayashi H, Kono T,

Kawahara M. 2021. Deciphering two rounds of cell lineage segregations during

bovine preimplantation development. *FASEB J* **35**:1–14.

doi:10.1096/fj.202002762RR

Ardestani G, Mehregan A, Fleig A, Horgen FD, Carvacho I, Fissore RA. 2020. Divalent cation influx and calcium homeostasis in germinal vesicle mouse oocytes. *Cell Calcium* **87**:102181. doi:10.1016/j.ceca.2020.102181

Arslan P, Di Virgilio F, Beltrame M. 1985. Cytosolic Ca2+ homeostasis in Ehrlich and Yoshida carcinomas. A new, membrane-permeant chelator of heavy metals reveals that these ascites tumor cell lines have normal cytosolic free Ca2+. *J Biol Chem* **260**:2719–2727. doi:10.1016/s0021-9258(18)89421-2

Barnett B, Kretshmar H, Hartman F. 1977. Structural Characterization of Bis (jV-oxopyridine-2-thionato) zinc (II). *Inorg Chem* **16**:1834–1838.

Bernhardt ML, Kim AM, O'Halloran T V., Woodruff TK. 2011. Zinc requirement during meiosis I-meiosis II transition in mouse oocytes is independent of the MOS-MAPK pathway. *Biol Reprod* **84**:526–536. doi:10.1095/biolreprod.110.086488

Bernhardt ML, Kong BY, Kim AM, O'Halloran T V., Woodruff TK. 2012. A Zinc-Dependent Mechanism Regulates Meiotic Progression in Mammalian Oocytes. *Biol Reprod* **86**:1–10. doi:10.1095/biolreprod.111.097253

Bernhardt ML, Padilla-Banks E, Stein P, Zhang Y, Williams CJ. 2017. Store-operated Ca2+ entry is not required for fertilization-induced Ca2+ signaling in mouse eggs. *Cell Calcium* **65**:63–72. doi:10.1016/j.ceca.2017.02.004

Berridge MJ. 2016. The inositol trisphosphate/calcium signaling pathway in health and disease. *Physiol Rev* **96**:1261–1296. doi:10.1152/physrev.00006.2016

Bhanumathy C, Da Fonseca PCA, Morris EP, Joseph SK. 2012. Identification of functionally critical residues in the channel domain of inositol trisphosphate receptors. *J Biol Chem* **287**:43674–43684. doi:10.1074/jbc.M112.415786

Bootman MD, Taylor CW, Berridge MJ. 1992. The thiol reagent, thimerosal, evokes Ca2+ spikes in HeLa cells by sensitizing the inositol 1,4,5-trisphosphate receptor. *J Biol Chem* **267**:25113–25119. doi:10.1016/s0021-9258(19)74013-7

Brind S, Swann K, Carroll J. 2000. Inositol 1,4,5-trisphosphate receptors are downregulated in mouse oocytes in response to sperm or adenophostin A but not to increases in intracellular Ca2+ or egg activation. *Dev Biol* **223**:251–265. doi:10.1006/dbio.2000.9728

Broun ER, Greist A, Tricot G, Hoffman R. 1990. Excessive Zinc Ingestion: A Reversible Cause of Sideroblastic Anemia and Bone Marrow Depression. *JAMA J Am Med Assoc* **264**:1441–1443. doi:10.1001/jama.1990.03450110087033

Carraway RE, Dobner PR. 2012. Zinc pyrithione induces ERK- and PKC-dependent necrosis distinct from TPEN-induced apoptosis in prostate cancer cells. *Biochim Biophys Acta - Mol Cell Res* **1823**:544–557. doi:10.1016/j.bbamcr.2011.09.013

Carvacho I, Ardestani G, Lee HC, McGarvey K, Fissore RA, Lykke-Hartmann K. 2016.
TRPM7-like channels are functionally expressed in oocytes and modulate post-
fertilization embryo development in mouse. *Sci Rep* **6**:1–12. doi:10.1038/srep34236

Carvacho I, Lee HC, Fissore RA, Clapham DE. 2013. TRPV3 Channels mediate
strontium-induced mouse-egg activation. *Cell Rep* **5**:1375–1386.
doi:10.1016/j.celrep.2013.11.007

Deguchi R, Shirakawa H, Oda S, Mohri T, Miyazaki S. 2000. Spatiotemporal analysis of
Ca2+ waves in relation to the sperm entry site and animal-vegetal axis during Ca2+
oscillations in fertilized mouse eggs. *Dev Biol* **218**:299–313.
doi:10.1006/dbio.1999.9573

des Georges A, Clarke OB, Zalk R, Yuan Q, Condon KJ, Grassucci RA, Hendrickson WA,
Marks AR, Frank J. 2016. Structural Basis for Gating and Activation of RyR1. *Cell*
167:145-157.e17. doi:10.1016/j.cell.2016.08.075

Ducibella T, Huneau D, Angelichio E, Xu Z, Schultz RM, Kopf GS, Fissore R, Madoux
S, Ozil J-P. 2002. Egg-to-Embryo Transition Is Driven by Differential Responses to
Ca2+ Oscillation Number. *Dev Biol* **250**:280–291. doi:10.1006/dbio.2002.0788

Dupont G, McGuinness OM, Johnson MH, Berridge MJ, Borgese F. 1996. Phospholipase
C in mouse oocytes: Characterization of β and γ isoforms and their possible
involvement in sperm-induced Ca2+ spiking. *Biochem J* **316**:583–591.
doi:10.1042/bj3160583

Eppig JJ. 1996. Coordination of nuclear and cytoplasmic oocyte maturation in eutherian
mammals. *Reprod Fertil Dev* **8**:485–489. doi:10.1071/RD9960485

Evellin S, Nolte J, Tysack K, Dorp F Vom, Thiel M, Oude Weernink PA, Jakobs KH,
Webb EJ, Lomasney JW, Schmidt M. 2002. Stimulation of phospholipase C-ε by the
M3 muscarinic acetylcholine receptor mediated by cyclic AMP and the GTPase
Rap2B. *J Biol Chem* **277**:16805–16813. doi:10.1074/jbc.M112024200

Fan G, Baker ML, Wang Z, Baker MR, Sinyagovskiy PA, Chiu W, Ludtke SJ, Serysheva
II. 2015. Gating machinery of InsP3R channels revealed by electron cryomicroscopy.
Nature **527**:336–341. doi:10.1038/nature15249

Fan G, Baker MR, Terry LE, Arige V, Chen M, Seryshev AB, Baker ML, Ludtke SJ, Yule
DI, Serysheva II. 2022. Conformational motions and ligand-binding underlying
gating and regulation in IP3R channel. *Nat Commun* **13**:1–15. doi:10.1038/s41467-
022-34574-1

Fissore RA, Longo FJ, Anderson E, Parys JB, Ducibella T. 1999. Differential distribution
of inositol trisphosphate receptor isoforms in mouse oocytes. *Biol Reprod* **60**:49–57.
doi:10.1095/biolreprod60.1.49

Fujiwara T, Nakada K, Shirakawa H, Shunichi M. 1993. Development of inositol
trisphosphate-induced calcium release mechanism during maturation of hamster
oocytes. *Dev Biol*.

Hajnóczky G, Thomas AP. 1997. Minimal requirements for calcium oscillations driven
by the IP3 receptor. *EMBO J* **16**:3533–3543. doi:10.1093/emboj/16.12.3533

Hamada K, Terauchi A, Mikoshiba K. 2003. Three-dimensional Rearrangements within
Inositol 1,4,5-Trisphosphate Receptor by Calcium. *J Biol Chem* **278**:52881–52889.
doi:10.1074/jbc.M309743200

Hara T, Yoshigai E, Ohashi T, Fukada T. 2022. Zinc transporters as potential therapeutic
targets: An updated review. *J Pharmacol Sci* **148**:221–228.
doi:10.1016/j.jphs.2021.11.007

Heim A, Tischer T, Mayer TU. 2018. Calcineurin promotes APC/C activation at meiotic
exit by acting on both XErp1 and Cdc20. *EMBO Rep* **19**:1–10.
doi:10.15252/embr.201846433

Higo T, Hattori M, Nakamura T, Natsume T, Michikawa T, Mikoshiba K. 2005. Subtype-
specific and ER lumenal environment-dependent regulation of inositol 1,4,5-
trisphosphate receptor type 1 by ERp44. *Cell* **120**:85–98.
doi:10.1016/j.cell.2004.11.048

Iino M. 1990. Biphasic Ca2+ dependence of inositol 1,4,5-trisphosphate-induced Ca
release in smooth muscle cells of the guinea pig taenia caeci. *J Gen Physiol* **95**:1103–
1122. doi:10.1085/jgp.95.6.1103

Jean T, Klee CB. 1986. Calcium modulation of inositol 1,4,5-trisphosphate-induced
calcium release from neuroblastoma x glioma hybrid (NG108-15) microsomes. *J
Biol Chem* **261**:16414–16420. doi:10.1016/s0021-9258(18)66582-2

Jellerette T, He CL, Wu H, Parys JB, Fissore RA. 2000. Down-regulation of the Inositol
1,4,5-Triphosphate Receptor in Mouse Eggs Following Fertilization or
Parthenogenetic Activation. *Dev Biol* **223**:238–250. doi:10.1006/dbio.2000.9675

Jellerette T, Kurokawa M, Lee B, Malcuit C, Yoon SY, Smyth J, Vermassen E, De Smedt
H, Parys JB, Fissore RA. 2004. Cell cycle-coupled [Ca2+]i oscillations in mouse
zygotes and function of the inositol 1,4,5-trisphosphate receptor-1. *Dev Biol* **274**:94–
109. doi:10.1016/j.ydbio.2004.06.020

Joseph SK, Young MP, Alzayady K, Yule DI, Ali M, Booth DM, Hajnóczky G. 2018.
Redox regulation of type-I inositol trisphosphate receptors in intact mammalian cells.
J Biol Chem **293**:17464–17476. doi:10.1074/jbc.RA118.005624

Kang D, Park JY, Han J, Bae IH, Yoon SY, Kang SS, Choi WS, Hong SG. 2003.
Acetylcholine induces Ca2+ oscillations via m3/m4 muscarinic receptors in the

mouse oocyte. *Pflugers Arch Eur J Physiol* **447**:321–327. doi:10.1007/s00424-003-1184-y

Kaur K, Gupta R, Saraf SA, Saraf SK. 2014. Zinc: The metal of life. *Compr Rev Food Sci Food Saf* **13**:358–376. doi:10.1111/1541-4337.12067

Kim AM, Bernhardt ML, Kong BY, Ahn RW, Vogt S, Woodruff TK, O'Halloran T V. 2011. Zinc sparks are triggered by fertilization and facilitate cell cycle resumption in mammalian eggs. *ACS Chem Biol* **6**:716–723. doi:10.1021/cb200084y

Kim AM, Vogt S, O'Halloran T V., Woodruff TK. 2010. Zinc availability regulates exit from meiosis in maturing mammalian oocytes. *Nat Chem Biol* **6**:674–681. doi:10.1038/nchembio.419

Kurokawa M, Fissore RA. 2003. ICSI-generated mouse zygotes exhibit altered calcium oscillations, inositol 1,4,5-trisphosphate receptor-1 down-regulation, and embryo development. *Mol Hum Reprod* **9**:523–533. doi:10.1093/molehr/gag072

Lawrence Y, Ozil JP, Swann K. 1998. The effects of a Ca2+ chelator and heavy-metal-ion chelators upon Ca2+ oscillations and activation at fertilization in mouse eggs suggest a role for repetitive Ca2+ increases. *Biochem J* **335**:335–342. doi:10.1042/bj3350335

Lee B, Vermassen E, Yoon SY, Vanderheyden V, Ito J, Alfandari D, De Smedt H, Parys JB, Fissore RA. 2006. Phosphorylation of IP3R1 and the regulation of [Ca2+]i responses at fertilization: A role for the MAP kinase pathway. *Development* **133**:4355–4365. doi:10.1242/dev.02624

Lemos FO, Bultynck G, Parys JB. 2021. A comprehensive overview of the complex world of the endo- and sarcoplasmic reticulum Ca2+-leak channels. *Biochim Biophys Acta - Mol Cell Res* **1868**:119020. doi:10.1016/j.bbamcr.2021.119020

Liu M. 2011. The biology and dynamics of mammalian cortical granules. *Reprod Biol Endocrinol* **9**:1–17. doi:10.1186/1477-7827-9-149

Lo MN, Damon LJ, Tay JW, Jia S, Palmer AE. 2020. Single cell analysis reveals multiple requirements for zinc in the mammalian cell cycle. *Elife* **9**:1–24. doi:10.7554/eLife.51107

Lorca T, Cruzalegui FH, Fesquet D, Cavadore J, Méry J, Means A, Marcel D. 1993. Calmodulin-dependent protein kinase II mediates inactivation of MPF and CSF upon fertilization of Xenopus eggs. *Nature* **366**:270–273.

Maret W, Li Y. 2009. Coordination dynamics of zinc in proteins. *Chem Rev* **109**:4682–4707. doi:10.1021/cr800556u

Matsu-ura T, Shirakawa H, Suzuki KGN, Miyamoto A, Sugiura K, Michikawa T, Kusumi A, Mikoshiba K. 2019. Dual-FRET imaging of IP 3 and Ca 2+ revealed Ca 2+ -

753   induced IP 3 production maintains long lasting Ca 2+ oscillations in fertilized mouse
754   eggs. *Sci Rep* **9**:1–11. doi:10.1038/s41598-019-40931-w

755 Mendivil-Perez M, Velez-Pardo C, Jimenez-Del-Rio M. 2012. TPEN induces apoptosis
756   independently of zinc chelator activity in a model of acute lymphoblastic leukemia
757   and Ex vivo acute leukemia cells through oxidative stress and mitochondria caspase-
758   3- and AIF-dependent pathways. *Oxid Med Cell Longev* **2012**.
759   doi:10.1155/2012/313275

760 Miao YL, Stein P, Jefferson WN, Padilla-Banks E, Williams CJ. 2012. Calcium influx-
761   mediated signaling is required for complete mouse egg activation. *Proc Natl Acad*
762   *Sci U S A* **109**:4169–4174. doi:10.1073/pnas.1112333109

763 Miyazaki S. 1988. Inositol 1,4,5-trisphosphate-induced calcium release and guanine
764   nucleotide-binding protein-mediated periodic calcium rises in golden hamster eggs.
765   *J Cell Biol* **106**:345–353.

766 Miyazaki S, Ito M. 2006. Calcium signals for egg activation in mammals. *J Pharmacol*
767   *Sci* **100**:545–552. doi:10.1254/jphs.CPJ06003X

768 Miyazaki SI, Hashimoto N, Yoshimoto Y, Kishimoto T, Igusa Y, Hiramoto Y. 1986.
769   Temporal and spatial dynamics of the periodic increase in intracellular free calcium
770   at fertilization of golden hamster eggs. *Dev Biol* **118**:259–267. doi:10.1016/0012-
771   1606(86)90093-X

772 Miyazaki SI, Yuzaki M, Nakada K, Shirakawa H, Nakanishi S, Nakade S, Mikoshiba K.
773   1992. Block of Ca2+ wave and Ca2+ oscillation by antibody to the inositol 1,4,5-
774   trisphosphate receptor in fertilized hamster eggs. *Science (80-)* **257**:251–255.
775   doi:10.1126/science.1321497

776 Nomikos M, Elgmati K, Theodoridou M, Calver BL, Nounesis G, Swann K, Lai FA. 2011.
777   Phospholipase Cζ binding to PtdIns(4,5)P 2 requires the XY-linker region. *J Cell Sci*
778   **124**:2582–2590. doi:10.1242/jcs.083485

779 Nomikos M, Sanders JR, Parthimos D, Buntwal L, Calver BL, Stamatiadis P, Smith A,
780   Clue M, Sideratou Z, Swann K, Lai FA. 2015. Essential role of the EF-hand domain
781   in targeting sperm phospholipase Cζ to membrane phosphatidylinositol 4,5-
782   bisphosphate (PIP2). *J Biol Chem* **290**:29519–29530. doi:10.1074/jbc.M115.658443

783 Pace NJ, Weerapana E. 2014. A competitive chemical-proteomic platform to identify
784   zinc-binding cysteines. *ACS Chem Biol* **9**:258–265. doi:10.1021/cb400622

785 Paknejad N, Hite RK. 2018. Structural basis for the regulation of inositol trisphosphate
786   receptors by Ca2+ and IP3. *Nat Struct Mol Biol* **25**:660–668. doi:10.1038/s41594-
787   018-0089-6

Palmer AE, Jin C, Reed JC, Tsien RY. 2004. Bcl-2-mediated alterations in endoplasmic

reticulum Ca2+ analyzed with an improved genetically encoded fluorescent sensor.

Proc Natl Acad Sci U S A **101**:17404–17409. doi:10.1073/pnas.0408030101

Parrington J, Brind S, De Smedt H, Gangeswaran R, Anthony Lai F, Wojcikiewicz R,

Carroll J. 1998. Expression of inositol 1,4,5-trisphosphate receptors in mouse

oocytes and early embryos: The type I isoform is upregulated in oocytes and

downregulated after fertilization. *Dev Biol* **203**:451–461.

doi:10.1006/dbio.1998.9071

Parys JB, De Smedt H, Missiaen L, Bootman MD, Sienaert I, Casteels R. 1995. Rat

basophiliv leukemia cells as model system for inositol 1,4,5-trisphosphate receptor

IV, a receptor of the type II family: functional comparison and immunological

detection. *Cell Calcium* **17**:239–249. doi:10.1016/0143-4160(95)90070-5

Perreault SD, Barbee RR, Slott VL. 1988. Importance of glutathione in the acquisition

and maintenance of sperm nuclear decondensing activity in maturing hamster

oocytes. *Dev Biol* **125**:181–186. doi:10.1016/0012-1606(88)90070-X

Qin Y, Dittmer PJ, Park JG, Jansen KB, Palmer AE. 2011. Measuring steady-state and

dynamic endoplasmic reticulum and Golgi Zn 2+ with genetically encoded sensors.

Proc Natl Acad Sci U S A **108**:7351–7356. doi:10.1073/pnas.1015686108

Que EL, Bleher R, Duncan FE, Kong BY, Gleber SC, Vogt S, Chen S, Garwin SA, Bayer

AR, Dravid VP, Woodruff TK, O'Halloran T V. 2015. Quantitative mapping of zinc

fluxes in the mammalian egg reveals the origin of fertilization-induced zinc sparks.

Nat Chem **7**:130–139. doi:10.1038/nchem.2133

Que EL, Duncan FE, Lee HC, Hornick JE, Vogt S, Fissore RA, O'Halloran T V., Woodruff

TK. 2019. Bovine eggs release zinc in response to parthenogenetic and sperm-

induced egg activation. *Theriogenology* **127**:41–48.

doi:10.1016/j.theriogenology.2018.12.031

Richardson A, Taylor CW. 1993. Effects of Ca2+ chelators on purified inositol 1,4,5-

trisphosphate (InsP3) receptors and InsP3-stimulated Ca2+ mobilization. *J Biol*

Chem **268**:11528–11533. doi:10.1016/s0021-9258(19)50232-0

Ridgway EB, Gilkey JC, Jaffe LF. 1977. Free calcium increases explosively in activating

medaka eggs. *Proc Natl Acad Sci U S A* **74**:623–627. doi:10.1073/pnas.74.2.623

Robinson MA. 1964. Complexes of 1-hydroxy 2-pyridinethione. *J Inorg Nucl Chem*

26:1277–1281. doi:10.1016/0022-1902(64)80210-4

Sanders JR, Ashley B, Moon A, Woolley TE, Swann K. 2018. PLCζ induced Ca2+

oscillations in mouse eggs involve a positive feedback cycle of Ca2+ induced InsP3

formation from cytoplasmic PIP2. *Front Cell Dev Biol* **6**:1–14.

doi:10.3389/fcell.2018.00036

Saunders CM, Larman MG, Parrington J, Cox LJ, Royse J, Blayney LM, Swann K, Lai
FA. 2002. PLC zeta: a sperm-specific trigger of Ca(2+) oscillations in eggs and
embryo development. *Development* **129**:3533–44.

Schultz RM, Kopf GS. 1995. Molecular Basis of Mammalian Egg Activation. *Curr Top*
Dev Biol **30**:21–62. doi:10.1007/978-3-642-83139-3_7

Shoji S, Muto Y, Ikeda M, He F, Tsuda K, Ohsawa N, Akasaka R, Terada T, Wakiyama
M, Shirouzu M, Yokoyama S. 2014. The zinc-binding region (ZBR) fragment of
Emi2 can inhibit APC/C by targeting its association with the coactivator Cdc20 and
UBE2C-mediated   ubiquitylation.   *FEBS*   *Open*   *Bio*   **4**:689–703.
doi:10.1016/j.fob.2014.06.010

Shoji S, Yoshida N, Amanai M, Ohgishi M, Fukui T, Fujimoto S, Nakano Y, Kajikawa E,
Perry ACF. 2006. Mammalian Emi2 mediates cytostatic arrest and transduces the
signal   for   meiotic   exit   via   Cdc20.   *EMBO*   *J*   **25**:834–845.
doi:10.1038/sj.emboj.7600953

Sienaert I, Missiaen L, De Smedt H, Parys JB, Sipma H, Casteels R. 1997. Molecular and
functional evidence for multiple Ca2+-binding domains in the type 1 inositol 1,4,5-
trisphosphate   receptor.   *J*   *Biol*   *Chem*   **272**:25899–25906.
doi:10.1074/jbc.272.41.25899

Sikora J, Ouagazzal AM. 2021. Synaptic zinc: An emerging player in Parkinson's disease.
Int J Mol Sci **22**:1–16. doi:10.3390/ijms22094724

Sipma H, De Smet P, Sienaert I, Vanlingen S, Missiaen L, Parys JB, De Smedt H. 1999.
Modulation of inositol 1,4,5-trisphosphate binding to the recombinant ligand-
binding site of the type-1 inositol 1,4,5-trisphosphate receptor by Ca2+ and
calmodulin. *J Biol Chem* **274**:12157–12162. doi:10.1074/jbc.274.17.12157

Snitsarev VA, McNulty TJ, Taylor CW. 1996. Endogenous heavy metal ions perturb fura-
2 measurements of basal and hormone-evoked Ca2+ signals. *Biophys J* **71**:1048–
1056. doi:10.1016/S0006-3495(96)79305-0

Stein P, Savy V, Williams AM, Williams CJ. 2020. Modulators of calcium signalling at
fertilization. *Open Biol* **10**. doi:10.1098/rsob.200118

Stricker SA. 1999. Comparative biology of calcium signaling during fertilization and egg
activation in animals. *Dev Biol* **211**:157–176. doi:10.1006/dbio.1999.9340

Suzuki T, Suzuki E, Yoshida N, Kubo A, Li H, Okuda E, Amanai M, Perry ACF. 2010a.
Mouse Emi2 as a distinctive regulatory hub in second meiotic metaphase.
Development **137**:3281–3291. doi:10.1242/dev.052480

Suzuki T, Yoshida N, Suzuki E, Okuda E, Perry ACF. 2010b. Full-term mouse
development by abolishing Zn2+-dependent metaphase II arrest without Ca2+

release. *Development* **137**:2659–2669. doi:10.1242/dev.049791

Swann K, Parrington J. 1999. Mechanism of Ca2+ release at fertilization in mammals. *J*

Exp Zool **285**:267–275. doi:10.1002/(SICI)1097-010X(19991015)285:3<267::AID-

JEZ10>3.0.CO;2-P

Taylor CW, Tovey SC. 2010. IP3 Receptors: Toward Understanding Their Activation.

Cold Spring Harb Perspect Biol **2**:1–23.

Thastrup O, Cullen PJ, Drobak BK, Hanley MR, Dawson AP. 1990. Thapsigargin, a tumor

promoter, discharges intracellular Ca2+ stores by specific inhibition of the

endoplasmic reticulum Ca2+-ATPase. *Proc Natl Acad Sci U S A* **87**:2466–2470.

doi:10.1073/pnas.87.7.2466

Toeplitz BK, Cohen AI, Funke PT, Parker WL, Gougoutas JZ. 1979. Structure of

Ionomycin—A Novel Diacidic Polyether Antibiotic Having High Affinity for

Calcium Ions. *J Am Chem Soc* **101**:3344–3353. doi:10.1021/ja00506a035

Tokuhiro K, Dean J. 2018. Glycan-Independent Gamete Recognition Triggers Egg Zinc

Sparks and ZP2 Cleavage to Prevent Polyspermy. *Dev Cell* **46**:627-640.e5.

doi:10.1016/j.devcel.2018.07.020

Uchida K, Miyauchi H, Furuichi T, Michikawa T, Mikoshiba K. 2003. Critical regions

for activation gating of the inositol 1,4,5-trisphosphate receptor. *J Biol Chem*

278:16551–16560. doi:10.1074/jbc.M300646200

Wakai T, Fissore RA. 2019. Constitutive IP3R1-mediated Ca2+ release reduces Ca2+

store content and stimulates mitochondrial metabolism in mouse GV oocytes. *J Cell*

Sci **132**. doi:10.1242/jcs.225441

Wakai T, Fissore RA. 2013. Ca2+ homeostasis and regulation of ER Ca2+ in mammalian

oocytes/eggs. *Cell Calcium* **53**:63–67. doi:10.1016/j.ceca.2012.11.010

Wakai T, Harada Y, Miyado K, Kono T. 2014. Mitochondrial dynamics controlled by

mitofusins define organelle positioning and movement during mouse oocyte

maturation. *Mol Hum Reprod* **20**:1090–1100. doi:10.1093/molehr/gau064

Wakai T, Mehregan A, Fissore RA. 2019. Ca2+ signaling and homeostasis in mammalian

oocytes and eggs. *Cold Spring Harb Perspect Biol* **11**.

doi:10.1101/cshperspect.a035162

Wakai T, Vanderheyden V, Yoon SY, Cheon B, Zhang N, Parys JB, Fissore RA. 2012.

Regulation of inositol 1,4,5-trisphosphate receptor function during mouse oocyte

maturation. *J Cell Physiol* **227**:705–717. doi:10.1002/jcp.22778

Wakai T, Zhang N, Vangheluwe P, Fissore RA. 2013. Regulation of endoplasmic

reticulum Ca2+ oscillations in mammalian eggs. *J Cell Sci* **126**:5714–5724.

doi:10.1242/jcs.136549

Walker JW, Somlyo A V., Goldman YE, Somlyo AP, Trentham DR. 1987. Kinetics of
smooth and skeletal muscle activation by laser pulse photolysis of caged inositol
1,4,5-trisphosphate. *Nature* **327**:249–252.

Watanabe S, Amagai Y, Sannino S, Tempio T, Anelli T, Harayama M, Masui S, Sorrentino
I, Yamada M, Sitia R, Inaba K. 2019. Zinc regulates ERp44-dependent protein
quality control in the early secretory pathway. *Nat Commun* **10**:1–16.
doi:10.1038/s41467-019-08429-1

Webster JM, Tiwari S, Weissman AM, Wojcikiewicz RJH. 2003. Inositol 1,4,5-
trisphosphate receptor ubiquitination is mediated by mammalian Ubc7, a component
of the endoplasmic reticulum-associated degradation pathway, and is inhibited by
chelation of intracellular Zn2+. *J Biol Chem* **278**:38238–38246.
doi:10.1074/jbc.M305600200

Woodier J, Rainbow RD, Stewart AJ, Pitt SJ. 2015. Intracellular zinc modulates cardiac
ryanodine receptor-mediated calcium release. *J Biol Chem* **290**:17599–17610.
doi:10.1074/jbc.M115.661280

Wozniak KL, Bainbridge RE, Summerville DW, Tembo M, Phelps WA, Sauer ML,
Wisner BW, Czekalski ME, Pasumarthy S, Hanson ML, Linderman MB, Luu CH,
Boehm ME, Sanders SM, Buckley KM, Bain DJ, Nicotra ML, Lee MT, Carlson AE.
2020. Zinc protection of fertilized eggs is an ancient feature of sexual reproduction
in animals. *PLoS Biol* **18**:1–17. doi:10.1371/journal.pbio.3000811

Wu H, Smyth J, Luzzi V, Fukami K, Takenawa T, Black SL, Allbritton NL, Fissore RA.
2001. Sperm factor induces intracellular free calcium oscillations by stimulating the
phosphoinositide pathway. *Biol Reprod* **64**:1338–1349.
doi:10.1095/biolreprod64.5.1338

Wu L, Sweet T, Clapham DE. 2010. Current Progress in the Mammalian TRP Ion Channel
Family. *Pharmocol Rev* **62**:381–404. doi:10.1124/pr.110.002725.Abbreviations

Xu Z, Williams CJ, Kopf GS, Schultz RM. 2003. Maturation-associated increase in IP3
receptor type 1: Role in conferring increased IP3 sensitivity and Ca2+ oscillatory
behavior in mouse eggs. *Dev Biol* **254**:163–171. doi:10.1016/S0012-
1606(02)00049-0

Yamasaki S, Sakata-Sogawa K, Hasegawa A, Suzuki T, Kabu K, Sato E, Kurosaki T,
Yamashita S, Tokunaga M, Nishida K, Hirano T. 2007. Zinc is a novel intracellular
second messenger. *J Cell Biol* **177**:637–645. doi:10.1083/jcb.200702081

Zhang N, Duncan FE, Que EL, O'Halloran T V., Woodruff TK. 2016. The fertilization-
induced zinc spark is a novel biomarker of mouse embryo quality and early
development. *Sci Rep* **6**:1–9. doi:10.1038/srep22772

Table

Table 1. Addition of TPEN after ICSI does not prevent extrusion of the second polar

body but precludes pronuclear (PN) formation.

Group*	No. of zygotes	2nd polar body (2.5h)	PN	
			4h	7h
Untreated	26	25 (96.1%)	23 (88.5%)	23 (88.5%)
TPEN (10μM)	27	24 (88.9%)	1 (3.7%)***	2 (7.4%)***

937 ***$P < 0.001$

938 *Data from three different replicates for each group.

Figures and Legends

Figure 1. TPEN-induced Zn^{2+} deficiency inhibits fertilization-initiated [Ca^{2+}]$_i$ oscillations in a dose-dependent manner.

(**A**) (Left panel) Representative normalized Zn^{2+} recordings of MII eggs loaded with FluoZin-3AM following the addition of increasing concentrations of TPEN (0 μM, DMSO, black trace; 2.5 μM, sky blue; 5 μM, blue; 10 μM, navy). TPEN was directly added to the monitoring media. (Right panel) Representative fluorescent images of MII eggs loaded FluoZin-3AM supplemented with 0, 2.5, and 10 μM of TPEN. Scale bar: 10 μm. (**B-D**). (**B**) Representative Ca^{2+} oscillations following ICSI after the addition of 0, 5, 10, or 50 μM TPEN (arrowheads). Insets show representative traces for eggs that resumed Ca^{2+} oscillations after TPEN. (**C**) As above, but following the addition of 100 μM EDTA, 100 or 500 μM DTPA (time of addition denoted by arrowheads). (**D**) Ca^{2+} oscillations following ICSI after the addition of 50, 100, and 500 μM TPA (horizontal bars of increasing thickness). (**E**) Representative bright field images of ICSI fertilized eggs 2.5, 4, and 7 hr. after sperm injection. Arrows and arrowheads denote the second polar body and PN formation, respectively. Scale bar: 10 μm.

Figure 2. TPEN dose-dependently inhibits Ca²⁺ oscillations in eggs triggered by a broad-spectrum of agonists that stimulate the PI pathway or IP₃R1.

(**A-D**) Representative Ca²⁺ responses induced by (**A**) m*Plcζ* mRNA microinjection (0.01 µg/µl, black traces), (**B**) strontium chloride (10 mM, green), (**C**) acetylcholine chloride (50 µM, orange), and (**D**) thimerosal (25 µM, purple) in MII eggs. Increasing concentrations of TPEN were added to the monitoring media (arrowheads above traces denotes the time of adding). Insets in the upper row show representative traces of eggs that stop oscillating despite others continuing to oscillate. (**E**) Each bar graph summarizes the TPEN effect on Ca²⁺ oscillations at the selected concentrations for each of the agonists in A-D. (**F**) Western blot showing the intensities of IP₃R1 and alpha-tubulin bands in MII eggs or in eggs injected with m*Plcζ* mRNA and incubated or not with TPEN above (*P* < 0.01). Thirty eggs per lane in all cases. This experiment was repeated twice, and the mean relative intensity of each blot is shown in the bar graph below.

Figure 3. TPEN inhibition of cIP₃-induced Ca²⁺ release is precluded by ZnSO₄ supplementation before TPEN exposure.

(**A-G**) Representative Ca²⁺ responses in MII eggs triggered by the release of caged IP₃ (cIP₃) induced by UV light pulses of increasing duration (arrows). (**A**) A representative control trace without TPEN, and (**B**) following the addition of 10 µM TPEN between the third and the fourth pulses. The broken line rectangle is magnified in the inset, farthest right side of the panel displaying the near immediate termination of an ongoing rise. (**C, D**) Recordings started in the presence of 10 µM TPEN but in (**D**) 100 µM ZnSO₄ was added between the second and the third pulses. (**E**) Recording started in the presence of 100 µM ZnSO₄ followed by the addition of 10 µM TPEN between the third and the fourth pulses. (**F, G**) Recording started in the presence of 100 µM MgSO₄ (**F**) or 100 µM CaCl₂ (**G**) and 10 µM TPEN added as above. Arrowheads above the different panels indicate the time of TPEN or divalent cation addition. (**H**) Bar graphs summarizing the number and percentages of eggs that responded to a given duration of UV pulses under each of the TPEN±divalent ions.

Figure 4. Zn²⁺ depletion alters Ca²⁺ homeostasis and increases Ca²⁺ store content independent of IP₃R1 mass.

(**A, B**) Representative Ca²⁺ traces of MII eggs after the addition of Tg and Io in the presence or absence of TPEN. Blue trace recordings represent TPEN-treated eggs whereas gray traces represent control, untreated eggs. (**A**) Io was added to fresh MII eggs once Ca²⁺ returned to baseline after treatment with Tg. Comparisons of mean peak amplitudes after Tg and Io are shown in the bar graphs in the right panel ($P < 0.001$). (**B**) MII eggs were aged by 2-hr. incubation supplemented or not with TPEN followed by Io addition and Ca²⁺ monitoring ($P < 0.001$). (**C**) Western blot showing the intensities of IP₃R1 bands in MII eggs freshly collected, aged by 4-hr. incubation without TPEN, and with TPEN. Thirty eggs per lane in all cases. This experiment was repeated three times. (**D, E**) (Left panels) Representative traces of Ca²⁺ values in eggs loaded with the Ca²⁺-sensitive dye Rhod-2 AM and the ER Ca²⁺reporter, D1ER (1 μg/μl mRNA). TPEN was added into the media followed 10 min. later by (**D**) 10 μM Tg and (**E**) 50 μM Ach. (Right panel) Bars represent the difference of FRET value between at the time of Tg/ Ach addition and at 3 and 5 min. later of the addition ($P < 0.05$). Experiments were repeated two different times for each treatment. Black and green traces represent

cytosolic Ca^{2+} and Ca^{2+}-ER, respectively. Blue and black arrowheads indicate the time of addition of

TPEN and Tg/ Ach, respectively. (**F**) Basal Zn^{2+} level comparison in WT (open bar) and *Trpv3$^{-/-}$*

/Trpm7$^{-/-}$ (dKO, orange bar) MII eggs. Each plot represents the Fluozin3 measurement at 5 min. after

starting monitoring. (**G**) (Left panel) Representative Ca^{2+} traces of WT (black trace) and dKO

(orange trace) MII eggs after adding Tg. Insets represent the magnified traces at the peak of Ca^{2+}

spike from different sets of eggs. (Middle panel) Individual traces of WT and dKO eggs after Tg

addition. Dashed circles represent the flection point in dKO traces. (Right panel) Comparisons of

mean peak amplitudes after Tg and the time between Tg addition and the Ca^{2+} peak are shown in the

bar graphs in the right panel ($P < 0.001$)

1006

Figure 5. Restoring Zn²⁺ levels with ZnPT rescues oscillations interrupted by TPEN-induced Zn²⁺ deficiency.

(**A**) Representative traces of Zn²⁺ in MII eggs following the addition of 0.01 to 0.1 µM concentrations of ZnTP. The broken rectangular area is amplified in the next panel to appreciate the subtle increase in basal Zn²⁺ caused by the addition of ZnTP. (**B, C**) m*Plcζ* mRNA (0.01 µg/µl)-induced oscillations followed by the addition of TPEN (black arrowhead) (**B**), or after the addition of TPEN followed by ZnPT (red arrowhead) (**C**). (**D, E**) Thimerosal (25 µM) induced oscillations using the same sequence of TPEN (**D**) and ZnPT (**E**), but higher concentrations of ZnPT were required to rescue Thimerosal-initiated oscillations (**E**). These experiments were repeated at least two different times.

Figure 6. Excess Zn²⁺ hinders Ca²⁺ oscillations.

(**A-D**) ICSI-initiated Ca²⁺ response without (**A**) or following the addition of ZnPT (**B, C**) (the time of ZnPT addition and concentration are denoted above the tracing). (**E, F**) Representative Ca²⁺ responses induced by injection of 0.01 μg/μl m*Plcζ* mRNA in untreated eggs (**E**) or in eggs treated with 0.1 μM ZnPT followed by 10 μM TPEN first and then 50 μM (**F**). (**G**) cIP₃-induced Ca²⁺ release as expected when the UV pulses in the absence but not in the presence of 0.05 μM ZnPT (the time of addition is denoted by a bar above the tracing). (**H, I**) Representative traces of Ca²⁺ values in eggs loaded with the Ca²⁺-sensitive dye Rhod-2 AM and the ER Ca²⁺reporter, D1ER (1 μg/μl mRNA). Uninjected and 0.01 μg/μl m*Plcζ* mRNA-injected eggs were monitored. After initiation and establishment of the oscillations, 0.1 μM ZnPT was added into the media followed 30 min later by 2.5 μM Io. Experiments were repeated two different times. Red and black arrowheads indicate the time of addition of ZnPT and Io, respectively.

Figure 7. Schematic of proposed regulation of IP₃R1 function by Zn²⁺ in eggs and fertilized zygotes.

In MII eggs, left panel, IP₃R1s are in a Ca²⁺-release permissive state with optimal levels of cytoplasmic Ca²⁺ and Zn²⁺ and maximum ER content, but Ca²⁺ is maintained at resting levels by the combined actions of pumps, ER Ca²⁺ leak, and reduced influx. Once fertilization takes place, left center panel, robust IP₃ production induced by the sperm-borne PLCζ leads to Ca²⁺ release through ligand-induced gating of IP₃R1. Continuous IP₃ production and refilling of the stores via Ca²⁺ influx ensure the persistence of the oscillations. Zn²⁺ release occurs in association with first few Ca²⁺ rises and cortical granule exocytosis, Zn²⁺ sparks, lowering Zn²⁺ levels but not sufficiently to inhibit IP₃R1 function. Zn²⁺ deficiency caused by TPEN or other permeable Zn²⁺ chelators, right center panel, dose-dependently impairs IP₃R1 function and limits Ca²⁺ release. We propose this is accomplished by stripping the Zn²⁺ bound to the residues of the zinc-finger motif in the LNK domain of IP₃R1 that prevents the allosteric modulation of the gating process induced by IP₃ or other agonists. We propose that excess Zn²⁺, right panel, also inhibits IP₃R1-mediate Ca²⁺ release, possibly by non-specific binding of thiol groups present in cysteine residues throughout the receptor (denoted by ?). We submit that optimal Ca²⁺ oscillations in mouse eggs unfold in the presence of a permissive range of Zn²⁺ concentration.

Supplementary Figure 1. Overexpression of ER accessory protein ERp44 did not change the Ca²⁺ responses initiated by m*Plcζ* mRNA microinjection, Actylcholine, or SrCl₂.

(**A**) Representative immunofluorescent images of MII eggs with overexpression of ERp44. At 5 hr. post microinjection, eggs were treated with 10 or 50 μM of TPEN and incubated for 1 hr, after which they were fixed and stained. An anti-HA antibody was used. Scale bar: 10 μm. (**B**) Representative Ca²⁺ responses induced by m*Plcζ* mRNA microinjection (0.01 μg/ μl-left column), SrCl₂ (10 mM-center column), and acetylcholine (50 μM-right column) in eggs with (top panels) or without (bottom panels) ERp44 overexpression.

Supplementary Figure 2. Elevated Zn²⁺ impairs egg activation and the subsequent embryo

development.

(**A**) MII eggs were incubated in TL-HEPES containing 0, 0.1, or 1.0 μM ZnPT at room temperature

for 10 min and washed several times with fresh TL-HEPES and injected with m*Plcζ* mRNA. After it,

eggs and zygotes were cultured in KSOM for 24 hr. PN formation and 2-cell development were

checked at 7- and 24-hr post-microinjection. Bars represent the percentages of injected eggs that

reached the PN and the 2-cell stage. Scale bar: 50 μm. (**B**) MII eggs injected with m*Plcζ* mRNA were

incubated in KSOM without ZnPT for an hr. and then incubated in KSOM with 0 or 0.1 μM ZnPT for

24 hr. The second polar body extrusion, PN formation, and 2-cell development were checked at 2.5-,

7- and 24-hr. post-microinjection. Bars represent the percentages of injected eggs that reached the PN

and the 2-cell stage. Scale bar: 50 μm.