1 **FOXL2 interaction with different binding partners regulates the dynamics of granulosa**

2 **cell differentiation across ovarian development**

3 Roberta Migale[a,1], Michelle Neumann[a*], Richard Mitter[b*], Mahmoud-Reza Rafiee[c], Sophie

4 Wood[d], Jessica Olsen[d] & Robin Lovell-Badge[a,1]

Affiliations

[a]Laboratory of Stem Cell Biology and Developmental Genetics, The Francis Crick Institute,

London NW1 1AT, UK

[b]Bioinformatics core, The Francis Crick Institute, London NW1 1AT, UK

[c]RNA Networks Laboratory, The Francis Crick Institute, London NW1 1AT, UK

[d]Genetic Modification Service, The Francis Crick Institute, London NW1 1AT, UK

*contributed equally

[1]Correspondence to: roberta.migale@crick.ac.uk & robin.lovell-badge@crick.ac.uk

Abstract

FOXL2 is a transcription factor essential for female fertility, expressed in somatic cells of the
ovary, notably granulosa cells. In the mouse, *Foxl2* deletion leads to partial sex reversal
postnatally, with mutants developing dysgenic ovaries devoid of oocytes. However, deletion
of the gene in 8-week-old females leads to granulosa to Sertoli cell transdifferentiation and
gonadal sex reversal. We hypothesise that different outcomes of *Foxl2* deletion in embryonic
versus adult ovary may derive from a different role played by FOXL2 across ovarian
development. Therefore, in this study, we take a multi-omics approach to characterise the
dynamics of gene expression and chromatin accessibility changes in purified murine granulosa
cells across key developmental stages (E14.5, 1 and 8 weeks). We coupled these analyses with
genome wide identification of FOXL2 target genes and on-chromatin interacting partners by
ChIP-SICAP to reconstruct the gene regulatory networks underpinned by this essential
transcription factor. In the embryonic ovary, FOXL2 interacts with factors important for early
stages of gonadal development, such as GATA4 and WT1, whilst postnatally it interacts with
factors regulating primordial follicle activation, such as NR5A2, and with factors regulating
steroidogenesis including AR and ESR2. Integration of chromatin landscape dynamics with
gene expression changes and FOXL2 binding sites analysis revealed that its critical role in
ovarian cell fate maintenance goes beyond repression of the Sertoli-specific gene *Sox9*. Our
chromatome analysis revealed also that FOXL2 interacts with several proteins involved in
chromatin remodelling, DNA repair, splicing and gene repression. In summary, in this study
we identified target genes dynamically regulated by FOXL2 across ovarian development
including known and newly identified FOXL2 targets with a role in embryonic ovarian
development and folliculogenesis, as well as cofactors that point towards additional roles
played by FOXL2 besides transcriptional regulation. This work constitutes a comprehensive
resource for exploration of the molecular mechanisms of ovarian development and causes of
female infertility.

INTRODUCTION

The development of primary reproductive organs, ovaries in the female and testes in the male, starts from a pair of bipotential structures called genital ridges. These develop from E9.5 in the mouse and from the 5th week of gestation in humans [1]. Bipotential genital ridges contain undifferentiated cell types poised to develop into one of two sexes, depending on the genetic switch activated [2]. The time and robustness of this cell fate decision is critical for the healthy development of the gonads, successful reproduction and ultimately for the survival of a species. The presence or absence of the male sex determination gene *Sry*, usually located on the Y chromosome in mammals, determines which type of gonad it will become [3]. Its expression, starting at E10.75 and peaking at E11.5 in the mouse [4, 5], is sufficient to induce male differentiation in an XX female context[3].

This is also the case for its downstream target *Sox9*, which is able to induce male development in the absence of *Sry* [6]. Ovarian development was considered as a default pathway activated in the absence of *Sry* [7]. Recent work has challenged this idea showing that female sex determination is not a default pathway, but an active process involving the constant repression of the genetic program of the opposite sex [8-10]. A critical player in ovarian development and function is *Foxl2*, a gene encoding for the forkhead box L2 (FOXL2) protein, a transcription factor involved in regulating the differentiation and maintenance of granulosa cells in the ovary [11, 12]. Mutations in the FOXL2 gene in humans have been linked to premature ovarian insufficiency (POI), a condition in which the ovaries stop functioning before the age of 40, leading to infertility [13-15].

Null mutant ($Foxl2^{-/-}$) and conditional mutant (R26CreERT2;$Foxl2^{fl/fl}$) murine models aimed at studying the role of this factor in ovarian development provided intriguing results. $Foxl2^{-/-}$ ovaries displayed morphological anomalies only from 1 week postpartum, involving the lack of fragmentation of sex cords into individual follicles, failure to activate the primordial-to-primary transition of granulosa cells, thus resulting in impaired folliculogenesis and infertility. At the molecular level, mutant ovaries were characterised by downregulation of somatic sex markers of folliculogenesis and upregulation of the pro-testis factor *Sox9* as well as other male gonadal-specific markers including *Fgf9*, *Dhh*, *Dmrt1* and *Gata4* [16, 17].

These studies suggest that FOXL2 is not critical for initial ovarian specification because ovaries will differentiate correctly up to when folliculogenesis should have started, around one week postnatally [17]. In contrast, conditional deletion of *Foxl2* from adult ovaries resulted into a more dramatic phenotype, involving transdifferentiation of the supporting cells of the ovary, granulosa cells, into Sertoli cells, their opposite sex counterpart [9]. Crucially, the ectopic upregulation of SOX9 in this model occurred within three days following *Foxl2* deletion, indeed as soon as FOXL2 protein was lost, much earlier compared to the *Foxl2* null mutant model. This study demonstrated the plasticity of the adult ovary and the crucial role played by FOXL2 in this process. More recently, a similar role in ovarian sex maintenance was discovered for TRIM28, which we show to be recruited on chromatin together with FOXL2 and to regulate the sex-specific SUMOylation profile of ovarian-specific genes through its E3-SUMO ligase activity [8].

Collectively these data support a differential role played by FOXL2 depending on the
developmental stage of the ovary, as well as supporting the presence of additional factors
cooperating with FOXL2 in ensuring correct ovarian development.
We hypothesised that this differential function of FOXL2 during development may be
dependent on the availability of key interacting partners. Therefore, in this study, we combined
FOXL2 ChIP-SICAP with RNA-Seq and ATAC-Seq on purified $Foxl2^{EGFP}$ expressing cells to
address the differences in the role played by FOXL2 at embryonic versus adult stages and to
reconstruct the gene regulatory network underpinned by this transcription factor fundamental
to female fertility. We chose three timepoints for the analysis: E14.5 for embryonic stages, 1
week postnatally, because this is equivalent to prepubertal age and corresponds to the timepoint
at which SOX9 protein is first detected in $Foxl2^{-/-}$ ovaries, and 8 weeks when it is imperative
for ovary maintenance. We identified: i) the direct targets of FOXL2 at different stages of
ovarian development, ii) on-chromatin protein interactors, and iii) the complement of gene
expression changes and cis-regulatory element dynamics occurring throughout ovarian
development.
Our data support a more prominent role of FOX2 in postnatal stages compared to early
timepoints and provide a rich resource of novel protein interactors colocalising with FOXL2
on nearby genomic locations that could be used to generate new hypotheses on the molecular
mechanisms of ovarian development, POI and other causes of female infertility.
114
115
116   **RESULTS**
117
118   <u>ChIP-SICAP reveals differential genomic binding of FOXL2 throughout ovarian development.</u>
119
To profile the dynamic changes in FOXL2 DNA binding and to identify its on-chromatin
protein interactors, we performed a modified version of ChIP-SICAP (Methods & [18, 19]) on
whole mouse ovaries at three timepoints: E14.5, one week postnatally (1W) and at 8 weeks
(8W) (Fig.1A). ChIP-SICAP is a dual approach which allowed us to simultaneously identify
genome-wide FOXL2 binding sites, via ChIP-Sequencing, and proteins colocalising with
FOXL2 on chromatin, via mass spectrometry.
E14.5 was chosen because this embryonic stage is characterised by high FOXL2 expression in
the bipotential pre-granulosa cells of the ovarian medulla [20]; 1W coincides with the
activation of primordial follicles which start to develop into primary follicles, a process where
FOXL2 plays a central role by regulating the transition of the granulosa cells surrounding the
oocytes from squamous to cuboidal, and by inhibiting the growth of oocytes [12]; at 8W
FOXL2 is widely expressed in the follicles where it regulates folliculogenesis [21] and
maintains ovarian cell fate [9].
Principal-component analysis (PCA) of ChIP-Seq peaks highlighted the differences between
FOXL2 binding sites across the timecourse. PC1 (54% of variance, Figure 1B) showed a clear
separation of the E14.5 from both 1W an 8W. PC2 (27% of variance) further separated the 1W
and 8W samples. Embryonic and postnatal FOXL2 binding sites differed in their genomic
location. Annotation of peaks showed that most identified at E14.5 (50%) were mapped to
promoters, whilst about 30% of peaks identified at 1W and 8W were located within promoters

(Figure S1A). Indeed, most of the peaks from these two timepoints localised to either the gene
body or to distal intergenic regions. Motif enrichment analysis of consensus peaks across the
timecourse confirmed enrichment of FOXL2 and other forkhead factors motifs (Figure S1B).
Furthermore, we found enrichment of steroid hormone receptors amongst the top 10 predicted
motifs. These included estrogen receptors and androgen receptor factors known to regulate
hormone production as well as granulosa cell proliferation throughout ovarian development
[22, 23], as well as NR5A1, an important transcriptional regulator of genes involved in the
hypothalamic-pituitary-gonadal steroidogenic axis [24] and FOXL2 binding partner [25], as
well as it being essential for gonadal development in both sexes [26].
148

Figure 1

149
Figure 1: Timecourse ChIP-SICAP reveals differential FOXL2 binding across ovarian
development.

(A) Experimental design. FOXL2 ChIP-SICAP was performed on mouse ovaries collected at
three timepoints (E14.5, 1W and 8W, n=2). A FOXL2-specific antibody was used to capture
genomics fragments bound by FOXL2 as well as proteins colocalising with FOXL2 on
chromatin. DNA sequencing was used to identify FOXL2 target genes and mass-spectrometry
was used to find its on-chromatin interacting partners. (B) PCA of FOXL2 ChIP-Seq consensus
peaks identified with the nf-core MACS analysis (n=2). (C) Venn diagram depicting the
number of peaks identified at each timepoint and their overlap across time. Only peaks present
in both replicates (n=2) over the input were considered. (D-F) Integrative Genome Viewer
(IGV) tracks of representative FOXL2-bound regions. Main tracks show normalised read-
depth coverage (RPM). Grey bars and black arrow highlight significant peaks over input. (D)
Cited2, (E) *Esr2* [27], and (F) *Rspo1.* Star denotes proven functional *Esr2* enhancer (see [27]).

In total, we detected 725 peaks at E14.5, 4476 at 1W and 2654 at 8W (Figure 1C, Table S1).

Amongst these, we identified known FOXL2 targets bound throughout the timecourse, such as

Cited2 (Figure 1D), as well as targets dynamically bound including *Esr2* (Figure 1E) and a

novel target gene *Rspo1* (Figure 1F) encoding a positive effector of β-catenin signalling that is

required for the development of functional ovaries [28].

Differential binding analysis allowed the identification of FOXL2-binding events that were

statistically enriched in one timepoint or the other (Figure S1C-E & Table S1). We detected

the greatest number of differential binding (2503 differential peaks) when comparing E14.5

with 8W (Figure S1D), followed by the comparison between E14.5 and 1W (Figure S1C, 2098

differential peaks) and lastly by the comparison between 1W and 8W (Figure S1E, 634

differential peaks). Comparison of our FOXL2 ChIP-Seq dataset with a microarray analysis of

differentially expressed genes in mouse ovaries lacking *Foxl2* compared to wildtype controls

(*Foxl2*$^{-/-}$, Ortiz *et al.* [29], Table S1) allowed us to identify 770 genes more likely to be activated

by FOXL2 (because downregulated upon loss of *Foxl2* as well as bound by FOXL2, Figure

S2A). This group included, besides *Cited2* and *Esr2*, the cell-cycle repressor gene *Cdkn1b*

(Figure S2B), a known FOXL2 target that regulates granulosa cell proliferation [30], *Smad7*

which encodes for a regulator of primordial follicle activation together with SMAD3 [31]

(Figure S2C), and the follistatin gene (*Fst*) which encodes a secreted inhibitor of the TGFβ

pathway, essential for ovarian development [32] (Figure S2D).

Comparison of the FOXL2 targets identified, with genes upregulated in the absence of FOXL2

(Figure S2E) highlighted a total of 376 genes more likely to be repressed by FOXL2 (because

they are upregulated when *Foxl2* is deleted and are FOXL2 targets as assessed by ChIP-Seq).

This group included genes encoding for signalling molecules crucial for female sexual

development such as *Rspo1* [33] (Figure 1F), and *Wnt4* [34] (Figure S2F), as well as genes

important for testis development such as *Gadd45g* [35] (Figure S2G), and *Inhbb* (Figure S2H)

[36]. Lastly, we evaluated FOXL2 binding in the genomic region containing *Sox9*, a gene

critical for testis development [6, 37] (Figure S2I). We did not detect significant binding of

FOXL2 to the *Sox9* promoter at any timepoint. We then scanned the 1 Mb gene desert upstream

of *Sox9*, which was previously found to contain cis-regulatory elements essential for *Sox9*

expression in Sertoli cells [38, 39]. As previously shown [8, 27], neither TESCO nor enhancer

13, were significantly bound by FOXL2 at any point during ovarian development. Instead, we

observed a dynamic binding of FOXL2 in other known *Sox9* enhancers. For example, FOXL2

was bound to Enh33 at 1W and to Enh32 at E14.5. Significant binding events of FOXL2 to the

female-specific *Sox9* Enh8 [38], *Sox9* Enh2 and *Sox9* Enh6 were detected at 8W. The

remaining 5 FOXL2 binding events, 4 of which were detected at 8W and 1 at 1W, were

distributed in other regions, not previously identified as Sertoli-specific *Sox9* enhancers, and

therefore may constitute granulosa-specific enhancers of *Sox9* used by FOXL2 to repress it.

In total, by scanning the whole gene desert 1Mb upstream of *Sox9*, we detected one significant

peak at E14.5, two at 1W and eight at 8W (Figure S2J).

We then clustered the ChIP-Seq peaks into sets that showed similar patterns of variation in

FOXL2 occupancy across the timecourse (Figure 2A) and performed Gene Ontology (GO)

analysis on each cluster (Figure 2B, Table S1) to investigate which biological processes

FOXL2 is regulating across ovarian development.

Figure 2

Figure 2: The contribution of FOXL2 to the regulation of granulosa and Sertoli-enriched genes is greater at postnatal stages.

(A) ChIP peak occupancy scores (gene-level). The heatmap has been split based on ChIP peak presence/absence across the conditions. Rows are scaled by z-score to aid visualisation. Black/grey bars on the left indicate presence/absence of a peak in a specific cluster. Number of genes in each cluster is shown below the cluster name. (B) Bar chart showing representative GO biological processes enriched in each dynamic cluster of FOXL2 binding events. See also

Table S1. (C-D) Overview of datasets used for the comparison. Bar charts indicate number of
genes enriched in either granulosa or Sertoli cells in the embryonic stage (E13.5, C) and
postnatal stage (P7-P29, D). Microarray gene expression analysis performed at E13.5 by
Jameson *et al*. [40], RNA-Seq analysis by Lindeman *et al*. [37] of primary granulosa cells from
mice between 23-29 days of age compared to Sertoli cells isolated from *CAG-Stop^{flox-}*
^{tdTomato};DhhCre P7 XY pups [41]. Significant genes enriched in Sertoli cells are coloured blue,
granulosa-enriched genes are coloured purple. (E) Pie chart depicting the proportion of FOXL2
target genes, as identified at E14.5, that are classified as either granulosa- or Sertoli enriched
at E13.5 [40]. Grey denotes other FOXL2 target genes identified by our ChIP-SICAP and not
overlapping with the lists of granulosa/Sertoli-enriched genes as found in the other two studies.
(F) Proportion of FOXL2 target genes, as identified at 1W and (G) 8W, that are classified as
either granulosa- or Sertoli and enriched postnatally.

Known FOXL2 target genes were identified amongst those shared across the timecourse and
enriched for cell proliferation and ribosome biogenesis pathways, including the granulosa-
enriched genes *Cited2*, *Smad7*, *Fst*, *Thbs1*, and the Sertoli-specific genes *Inhbb* and *Fgf9*.
Rspo1, which encodes for a secreted signalling molecule that regulates WNT4 expression and
is essential for female sex determination [42], was a target uniquely regulated by FOXL2 at
E14.5, together with other genes including *Adora1*, *Pdgfc*, *Bcl9* and *Tert*, involved in cell
migration and proliferation, cell projection organisation and Wnt signalling. Amongst the
targets significantly bound by FOXL2 at 1W, we found genes involved in urogenital system
development including *Amh*, *Amhr2*, *Fgfr2*, *Ctnnb1* and *Igf2r*. Of those genes uniquely bound
by FOXL2 at 8W, we found an enrichment in those implicated in gland development and
steroidogenesis pathways. These include known FOXL2 targets such as *Cyp19a1*, the gene
encoding for aromatase [43], *Nr5a1*, known to physically interact with FOXL2 [25], and
Hsd17b7, encoding for a steroidogenic enzyme involved in the estradiol synthesis. Whilst the
timepoints E14.5-1W shared several targets involved in urogenital development, the group of
genes bound by FOXL2 between 1W and 8W featured genes encoding for members of the
steroidogenesis pathway, including the known FOXL2 targets *Esr2*, for which we detected
binding in the functional enhancer "Peak 3", located in one intronic region as previously
demonstrated [27] (Figure 1E), the human steroidogenic acute regulatory *Star* gene *[44]* and
Cyp17a1. Additionally, sex differentiation genes were stably bound by FOXL2 between 1W
and 8W, and these include granulosa-specific genes such as *Esr1* [45], *Irx3* [46] and *Wnt4* [34],
as well as Sertoli-specific genes, such as *Sox9* [37], *Inha* [47], *Fndc3a* [48] and bivalent factors
such as *Wt1* [49]. Additionally, we found binding within the cell-cycle repressor gene *Cdkn1b*,
a known FOXL2 target gene that regulates granulosa cell proliferation (Figure 2 and Figure
S2B) [30]. Lastly, targets shared between E14.5 and 8W were enriched for DNA repair
pathways, a recently discovered function of FOXL2 [50], and included genes such as *Ddx1*,
Ruvbl1, and *Exo5*.
Next, we assessed the relative contribution of FOXL2 to the regulation of granulosa *versus*
Sertoli-enriched genes at different developmental stages. To obtain genes enriched in granulosa
or Sertoli cells relevant for the embryonic and the postnatal stages of our timecourse, we used
datasets from two studies (Table S2): one by Jameson *et al.* reporting microarray analysis of

sorted XX (*Sry^EGFP*) and XY (*Sox9^ECFP*) supporting cells collected at E13.5 [40], and another
by Lindeman *et al.* reporting RNA-Seq analysis of primary granulosa cells from mice between
23-29 days of age compared to Sertoli cells isolated from *CAG-Stop^flox-tdTomato;DhhCre* P7 XY
pups [41], which constitute a reference for our postnatal timepoints.
We calculated the overlap between FOXL2 targets characterising each stage of ovarian
development, and the granulosa-, and Sertoli-enriched genes identified by comparing XX and
XY supporting cells transcriptomes at either E13.5, for the comparison with E14.5 FOXL2
targets (Figure 2C), or postnatally (Figure 2D), for the comparison with 1W and 8W FOXL2
target genes. We observed that during the embryonic stage E14.5 only 6% of FOXL2 targets
corresponded to either granulosa or Sertoli-enriched genes, whilst this percentage increased at
later timepoints, as 15% of FOXL2 target genes identified at 1W or 8W consisted of either
granulosa-enriched genes or Sertoli-enriched genes. These comparisons, together with the
greater number of targets identified at 1 and 8W compared to E14.5 in our timecourse analysis,
support a greater role played by FOXL2 in controlling supporting cell-specific genes
postnatally.
274
275
276 <u>Identification of FOXL2 on-chromatin protein interactors</u>
277
We next used ChIP-SICAP to identify on-chromatin protein interactors (chromatome) of
FOXL2 by applying mass-spectrometry on chromatin samples isolated at different stages of
ovarian development (Figure 1A). Proteins co-localising with FOXL2 on the same chromatin
fragments were pulled down using a specific antibody against FOXL2 followed by DNA
biotinylation to separate chromatin-bound proteins (see Methods). The two-step purification
strategy improves the purity of the samples and reduces the artificial interactions that are
generated during the immunoprecipitation procedure. We identified a total of 112 proteins at
E14.5, 430 at 1W, and 300 at 8W (>4-fold change FOXL2/no antibody control, n=2, adj-
pvalue<0.1, Fig. 3A and Table S3), thus reconstructing the FOXL2 chromatome across key
stages of ovarian development. Comparison of the relative protein intensities showed that the
most highly abundant proteins were known nuclear proteins, indicating a successful enrichment
for nuclear factors over potential cytoplasmic contaminants (Figure 3B). GO enrichment
analysis of FOXL2 interactors showed enrichment of proteins involved in mRNA splicing,
transcription, DNA damage and repair, and cell division (Figure 3C and Figure S3). FOXL2
was amongst the top 20 mostly enriched proteins identified (Figure 3D-F), together with
previously identified FOXL2 nuclear interactors namely GATAD2B, SUPT16H, ATRX,
SMARCE1, XRCC6 [51], and TRIM28 [8]. Amongst the FOXL2 protein partners present
across the timecourse we found members of the WNT signalling pathway including CTNN1B
(β-catenin), TCF4, and TCF21 (Figure S3, "transcription factors" and "transcriptional
regulators"), cell cycle regulators such as HMGA2, heterochromatin-interacting proteins such
as CBX3 and members of the splicing system including the DNA helicase DHX9, SRSF1, and
DDX17.
300

Figure 3

Figure 3: Identification of FOXL2 on-chromatin interactors by ChIP-SICAP

(A) Venn diagram representing the FOXL2 interactors identified at each timepoint by FOXL2 ChIP-SICAP. Total number of proteins identified at each stage is reported between round brackets. (B) Bar chart showing the relative intensities calculated using iBAQ (intensity Based Absolute Quantification) of the proteins identified in each timepoint and classified as either nuclear or cytoplasmatic only (proteins shuttling between nucleus and cytoplasm are classified as nuclear). (C) GO Biological processes enrichment analysis of the FOXL2 interactors. (D-F) Volcano plot showing enrichment (mean fold enrichment >4 and adj. p value <0.1, Benjamini-Hochberg adj. p values, FOXL2 immunoprecipitate *vs* no antibody control, n=2/timepoint) of proteins co-localising with FOLX2 on chromatin at E14.5 (D), 1W (E), and at 8W (F). Dotted lines separate significant proteins, also highlighted in yellow, orange and blue, depending on the timepoint. Black dots: selected FOXL2 interactors.

Importantly, we identified a series of FOXL2-interacting proteins with well-recognised roles in ovarian development and primordial follicle activation in human and mice, being differently present across the timecourse, as follows (Figure 3).

At E14.5 and 1W, factors essential for gonadal development such as GATA4 [52] and WT1 [53] were identified (Figure 3D). These were not detected at 8W (Figure 3F). Our analysis revealed the presence of NR5A2, a newly identified regulator of primordial follicle activation [54], co-localising with FOXL2 at 1W and 8W (Figure 3E-F). Additionally, we detected two

additional known regulators of the primordial-to-primary follicle transition, namely the
deacetylase HDAC6 [55] and the transcription factor STAT3 [56].
We found androgen receptor (AR), a factor that contributes greatly to normal follicle
development [22], enriched at 1W, but not later at 8W. Importantly, we identified estrogen
receptor 2 (ESR2) amongst the FOXL2 interactors at both postnatal timepoints, in agreement
with the previously postulated interplay between these two factors in orchestrating functions
key to ovarian physiology such as steroidogenesis [27] and cell fate maintenance [9].
We then compared our *in vivo* FOXL2 chromatome with a FOXL2 co-immunoprecipitation,
whole-cell proteomics dataset performed on the granulosa-like cell line AT29C [51] to assess
the level of overlap between our chromatin-specific dataset and one that would also include
cytoplasmic interactors. We found that 111 out of the 490 proteins we identified throughout
the timecourse were also found by this study (22%). Of these shared FOXL2 interacting
partners, 31 were found in our E14.5 timepoint, 95 at 1W and 72 at 8W (Figure S3A, Table
S3). These shared proteins included the known FOXL2 interactor involved in sex maintenance
TRIM28 [8], chromatin remodelling enzymes such as ATRX, which is involved in sexual
differentiation [57], regulators of the WNT signalling pathway such CXXC5, and DNA repair
proteins such as XRCC6, previously identified in a yeast-two-hybrid screen and co-
immunoprecipitation study [58].
In summary, these results represent the first description of the chromatin interactome of the
female-specific transcription factor FOXL2. We showed that this factor, essential for female
fertility, differentially interacts with proteins playing a role in a range of gonadal-essential
functions, from gonadogenesis, through primordial follicle activation, to steroidogenesis.
345
346
347 <u>Generation of a *Foxl2^EGFP^* mouse line to isolate granulosa cells.</u>
348
We next sought to analyse the transcriptomes and chromatin landscapes of granulosa cells
expressing FOXL2, to ultimately integrate these datasets with genomic binding locations of
this factor (ChIP-Seq), thus reconstructing the gene regulatory networks at play at different
developmental stages of the ovary. We therefore used CRISPR-Cas9 genome editing to
generate a *Foxl2^EGFP^* mouse line, allowing the purification of cells expressing FOXL2 (Figure
4A, D and Figure S4, see also Methods). *Foxl2^EGFP+/-^* ovaries imaged by fluorescence
microscopy showed a bright GFP signal specifically in the ovarian tissue at all stages examined
(E14.5, 1W, and 8W, Figure 4B). As expected, given the known absence of FOXL2 expression
in the testis, no GFP signal was detected in *Foxl2^EGFP+/-^* males, similarly to *Foxl2^EGFP-/-^*
wildtype testes and *Foxl2^EGFP-/-^* XX wildtype ovaries (Figure 4B and S5A). Colocalization of
GFP and endogenous FOXL2 protein in *Foxl2^EGF+/-^* ovaries and absence of this in *Foxl2^EGFP-/-^*
ovaries and *Foxl2^EGFP+/-^* testes was confirmed by immunofluorescence (Figure 4C). The line
was maintained as heterozygotes. Fertility performance of *Foxl2^EGFP+/-^* females was monitored
over 12 weeks and found to be normal compared to control wildtype females (C57BL/6J stock
mice, Figure S5B).
We dissected ovaries from *Foxl2^EGFP+/-^* mice at the following timepoints: E14.5, 1W and 8W
(Figure 4D). We used fluorescence-activated cell sorting (FACS) to isolate *Foxl2^EGFP^* positive
cells (Figure S6).

Figure 4

Figure 4: Generation of a *Foxl2^EGFP* mouse line to study granulosa cells across ovarian development

(A) Overview of the CRISPR/Cas9 mediated EGFP knock-in at the *Foxl2* locus. (B) Representative images of EGFP fluorescence in freshly collected gonads. Scale bars: 200 μm. Dotted lines outline the embryonic testis (left) and ovary (right). A minimum of three biological replicates were analysed. (C) Immunofluorescence staining of EGFP and endogenous FOXL2 in adult XX *Foxl2^EGFP-/-* ovaries (a-d), XX *Foxl2 ^EGFP+/-* ovaries (e-h), and of EGFP and endogenous SOX9 in XY *Foxl2 ^EGFP+/-* testes. Scale bars: 100 μm. (D) Overview of sample collection and downstream analyses performed. (E) PCA plot of RNA-Seq analysis of cells isolated from E14.5, 1W and 8W gonads (50k cells/replicate, n=3 biological replicates/timepoint). (F) Barplot summary illustrating the number of differentially expressed genes (DEG) identified in each pairwise comparison (DESeq2's Wald test, *pvalue*<0.01, fold-change of >2) (G) Heatmap of expression changes in genes validated as theca or granulosa cell markers.

A total of 50k and 100k of *Foxl2^EGFP* positive cells were used to perform bulk RNA-Seq and ATAC-Seq respectively, using three biological replicates for each timepoint (See Methods). PCA of RNA-Seq data (Figure 4E) based on the variance stabilised abundance estimates showed good separation between the timepoints indicating major transcriptional changes characterising granulosa *Foxl2^EGFP* positive cells throughout time. Most of the variance in the dataset was evident along the first principal component (57.3% variance), which separated

E14.5 samples from both W1 and W8. Poisson similarity metric indicated more similarity
between the transcriptome of postnatal samples (1W and 8W) compared to E14.5 (Figure S7A).
To examine the global transcriptome dynamics, we identify and selected genes significantly
changing across the timecourse in at least one pairwise comparison (Table S4 and Figure S7B).
Differential gene expression analysis comparing 8W and E14.5 revealed the greatest changes
with 1436 genes upregulated and 1134 downregulated; comparing 1W with E14.5 revealed
upregulation of 1307 genes and downregulation of 861 whilst the least number of changes were
detected between 8W and 1W, with 489 genes upregulated and 556 genes downregulated
(Figure 4F). Granulosa cells and the steroidogenic cell lineage including theca cells both
express FOXL2 [59]. We therefore checked expression of proven markers [60] expressed
exclusively in either theca or granulosa cells (Figure 4 G). This analysis indicated enrichment
of granulosa cells over theca cells in cells isolated using the $Foxl2^{EGFP}$ transgene.
We then used k-means clustering to analyse the dynamics of gene expression changes across
time (Figure S7B-C) and identified seven main patterns of gene expression. Clusters 7 and 6
consisted of genes highly expressed at E14.5 including markers of pre-granulosa cells such as
Rspo1, *Lgr5* and *Wnt4*, important for the initial proliferation of embryonic gonads [28]. GO
Biological processes analysis (GO BP, Figure S7D) of these clusters revealed enrichment of
biological processes such as Wnt signalling (*Wnt9a*, *Axin2*, *Fgfr2*, *Rspo1*) and urogenital
system development pathways (*Cdkn1b*, *Wt1*, *Wnt4*, *Cdkn1c*, *Rara*, *Emx2*, *Yap1*). Cluster 5
included genes moderately expressed at E14.5 and upregulated specifically at 1W and included
genes involved in kidney and urogenital system development (*Bmp7*, *Gli3*, *Smad3*, *Hey1*),
genetic imprinting (*Zfp57*, *Airn*, *Xist*), and extracellular matrix organisation (*Col18a1*,
Adamts9, *Eln*). Cluster 1 consisted of genes uniquely up regulated at 1W and participating in
extracellular matrix organisation, cell cycle phase transition, and DNA replication (*Pola2*,
Prim1, *Rad51*, *Cdc7*, *Dna2*). Cluster 4 grouped genes activated as the ovary develops from 1W
and remain upregulated in adulthood at 8W. GO analysis showed enrichment in nuclear
division, cell cycle (*Ube2c*, *Fbxo5*, *Top2a*), lipid transport, and synapses organisation (*Sema3e*,
Tanc2, *Ctnna2*). Cluster 3 included genes specifically upregulated at 8W and enriched in
steroid metabolic processes (*Star*, *Hsd17b1*, *Cyp19a1*) and angiogenesis (*Sema5a*, *Cd36*, *Jak1*,
Gadd45a). Lastly, Cluster 2 included genes expressed at the two divergent timepoints of E14.5
and 8W and characterised by high representation of lipid metabolism (*Apoe*, *Cyp17a1*, *Runx1*),
purine ribonucleotide biosynthesis (*Impdh1*, *Stat3*, *Bcl2l1*) and small molecule catabolic
processes (*Aldh1l1*, *Fah*, *Bcat2*). These data show the timely activation of genes important for
embryonic and postnatal ovarian development and function in our system.
424
425
426   <u>An integrative approach to reconstruct gene regulatory networks controlled by FOXL2.</u>
427
Having identified the gene expression changes characterising $Foxl2^{EGFP+/-}$ granulosa cells, we
then sought to reconstruct the gene regulatory networks (GRN) underpinned by this factor
(Figure 5 and Table S5).
We took the total number of FOXL2-bound genes identified by ChIP-Seq at any given
timepoint (Figure 1) and intersected it with the entire set of differentially expressed genes
(DEG) identified by RNA-Seq (Figure 4). Genes that changed in their expression value at any

point between E14.5, 1W and 8W (3333 genes) and that were assigned to a FOXL2 ChIP peak in at least one timepoint (4066 genes) were considered as "dynamic/FOXL2-bound" genes. The rest of the genes bound by FOXL2 but not changing their expression values were considered as "stable" (2966 in total) and divided between "stably repressed" and "stably activated", based on a variance stabilised (VST) cut-off of 8.3 (Figure S8 A-B). This threshold was chosen based on the expression values of known granulosa-enriched genes (*Foxl2*, *Esr2*, *Rspo1*), versus genes residing on the Y chromosome therefore not expected to be expressed at all in XX ovaries (*Sry*, *Usp9y* and *Uty*), as well as on the expression values of Sertoli-enriched genes (*Dmrt1*, *Sox9* and *Fgf9*) (Figure S8C).

Figure 5

Figure 5: Integrative approach combining RNA-Seq and ChIP-Seq to refine the gene regulatory networks controlled by FOXL2.

(A) Upset plot depicting the number of differentially expressed genes (DEG) identified by RNA-Seq, FOXL2 gene targets identified by ChIP-SICAP and the overlap between the two datasets. (B) Plot of variance stabilised (VST) gene expression values as detected by RNA-Seq on *Foxl2* $^{EGFP+/-}$ sorted ovarian cells (n=3). A threshold of VTS=8.3 was used to differentiate stably repressed from stably activated genes. (C) GO Biological processes enrichment analysis of the two groups of genes, stably repressed (left), and stably activated (right). (D) Hierarchical clustering of the 1100 genes bound by FOXL2 and differentially expressed across the timecourse. Gene examples from each cluster are shown on the right. (E) GO enrichment analysis of the clusters. Four representative pathways are depicted for each cluster.

Using this approach, we identified a total of 777 genes bound by FOXL2, but which are not expressed at any point during the timecourse and are therefore assumed to be repressed by FOXL2 (Figure 5B and Figure S8B). These were enriched in cell-cell adhesion, immune, and protein secretion pathways (Figure 5C). These included also Sertoli-enriched genes such as *Spz1* [61], *Cdh8*, *Rims1*, *Fgf9* [62], *Hsd17b3*, *Hopx* and *Tuba3* [41]. In contrast, a significant higher number of genes were bound by FOXL2 and were expressed throughout the timecourse but did not significantly change in levels across this (2189 genes). These were enriched in mRNA processing, ubiquitin-dependent catabolic processes, cell cycle and histone modification pathways (Figure 5C). In contrast, the "dynamic/FOXL2-bound" group of genes contained a total of 1100 genes (Figure 5D). We used unsupervised hierarchical clustering to analyse their pattern of expression throughout the timecourse and identified 7 main clusters (Figure 5D). GO enrichment analysis of these clusters showed that FOXL2 regulates a plethora of dynamically expressed genes enriched for functions critical for different stages of ovarian development. For example, we found enrichment in steroid metabolic and angiogenic processes, typical of folliculogenesis, in cluster 1 (C1), which included genes upregulated specifically at 8W. Of these, 53% were also a direct target of FOXL2 specifically at 8W (Figure 5D-E, Table S5). On the contrary, if considering C3, C4 and C6, clusters included genes upregulated at E14.5 compared to 8W, only 12% of the genes were significant targets of FOXL2 at E14.5. This confirmed the preponderant role of FOXL2 in regulating significantly changing genes at postnatal rather than during embryonic development. The high enrichment of Wnt signalling pathways found in C3, which contained a total of 263 genes upregulated at E14.5 only 36 of which were bound by FOXL2 at this timepoint (including 8 Wnt signalling pathway effectors such as *Rspo1*, *Zfp703* and *Bcl9*), was consistent with the established role of this pathway in ovarian development [34].

Foxl2^{EGFP} positive cells have a dynamic chromatin landscape across ovarian development

To explore the dynamics of chromatin remodelling underlying the gene expression changes throughout granulosa cells development, we performed ATAC-Seq on *Foxl2*^{EGFP} positive cells purified with FACS (Figure 6) at three timepoints (E14.5, 1W and 8W). PCA of genome-wide

chromatin accessibility on a total of 63831 consensus peaks (n=3) showed that most of the differences in chromatin landscape (57% variance) were found between the embryonic timepoint E14.5 and the late timepoints 1W and 8W (Figure 6A), in agreement with the transcriptional changes shown in Figure 4.

Figure 6

Figure 6: Granulosa cells have a dynamic chromatin landscape throughout development.

(A) PCA of ATAC-Seq performed on *Foxl2*^{EGFP+/-} sorted ovarian cells (n=3) at E14.5, 1W and 8W. (B) Venn diagram reporting the number of significant peaks identified at each stage. (C) Hierarchical clustering of chromatin landscape remodelling. (D-E) IGV snapshots of representative regions. *Sertoli H3K27Ac ChIP-Seq was re-analysed to match mm10 coordinates from Maatouk e*t al* [63]. (F) MonaLisa analysis of TF motifs enrichment analysis within the open chromatin regions as detected by ATAC-Seq. Top 6 motifs for each cluster are

reported (Table S6). (G) Heatmap displaying gene expression changes of representative TF, as
assessed by RNA-Seq (Table S4).

The second largest difference (21% variance) was observed between 1W and 8W, when
additional remodelling occurs in granulosa cells as folliculogenesis starts. Poisson metric also
confirmed a higher similarity between 1W and 8W samples, whilst E14.5 timepoint was
clustering separately (Figure S9A).
In total, we identified 45385 accessible regions at E14.5, 39708 at 1W and 44950 at 8W (Figure
6B, Table S6). Of these, 27079 were accessible throughout the timecourse, while 11130 were
unique to E14.5, 4333 unique to 1W, and 9233 unique to 8W. The genomic annotation of open
chromatin regions was similar across timepoints, with 32-35% mapping to promoters, 34-38%
to the gene body, 3% downstream of a gene, and 25-28 % mapping to distal intergenic regions
(Figure S9B). K-means clustering of consensus peaks revealed a total of 7 patterns representing
similar dynamic opening and closing of chromatin regions across ovarian development (Figure
6C). To assess the potential contribution of FOXL2 in binding to these open chromatin regions,
we specifically looked for enrichment values for the FOXL2 canonical motif (JASPAR id:
MA1607.1) in each ATAC-Seq cluster (Figure S9C-D). Motif analysis of these dynamic
regions revealed a greater enrichment for the canonical FOXL2 motif in clusters 6, 3, 4, 2 and
the least in cluster 7 (fold enrichment 1.4-1.2, adjPvalue>4). Amongst these, clusters 6 and 3,
which included peaks accessible preferentially at 1W and 8W and closed at E14.5, were the
two most significantly enriched with FOXL2 motifs (Figure S9C-D). In contrast, amongst
regions accessible at E14.5 (c1, c5 and c7) only c7 contained significant enrichment of FOXL2
binding sites, albeit with an enrichment score much lower than c6 and c3, containing elements
opened at 1 and 8W.
Differential accessibility analysis was then performed using DESeq2 [64] (Figure S10, Table
S6). This analysis revealed a total of 40165 differentially accessible regions (DARs) throughout
the timecourse, 32611 of which were changing between E14.5 and 8W, 24026 were changing
between E14.5 and 1W, and 16054 were changing between 1W and 8W (Figure S10A-D).
These chromatin landscape changes were consistent with the transcriptomic data, which
showed that most of the differences were detected comparing E14.5 and 8W samples (Figure
S7A).
Our profiling of chromatin accessibility dynamics contained novel potential cis-regulatory
regions, for example one within *Rspo1* gene, differentially bound by FOXL2 according to our
ChIP-Seq analysis (Figure 1F), and showing a similar change in its accessibility profile, as
indicated by ATAC-Seq (Figure S10E). This region was highly conserved and contained a
candidate cis-regulatory element sequence as predicted by the ENCODE Registry [65].
We also detected FOXL2 binding by ChIP-Seq at a known *Wt1* enhancer located 50 kb
upstream of its promoter [63] (Figure 6D). This binding event correlated with closing of this
region as ovarian development progressed, with no opening detected at 8W. In contrast, *Esr2*
was bound by FOXL2 at its known functional enhancer, located in the 8[th] intron of the gene,
only at postnatal stages (Figure 6E) [27]. This coincided with an opening of this region at 1W

and 8W. This enhancer was not marked by H3K27Ac (active enhancer mark) in Sertoli cells,
therefore can be considered as a granulosa-specific cis-regulatory region of *Esr2*.
K-means clustering of the changes identified seven clusters representative of the main
chromatin remodelling patterns (Figure 6C). Cluster 7 (c7) included peaks accessible at E14.5,
followed by c5, which included peaks significantly opened at E14.5 and starting to become
accessible at 1W. c1 included peaks closing down at E14.5 and opening up at 1W, whilst
remaining closed at 8W. c6 contained the majority of peaks accessible at 1W, which are also
starting to open up at 8W. c3 includes peaks progressively closing at 1W and opening at 8W,
while c4 included peaks that have gained accessibility at 8W. Lastly, c2 contained a group of
2713 peaks opened at the embryonic stage, E14.5, and in adulthood, at 8W, but not at 1W
(Figure 6C). Amongst the DARs we identified regions located nearby granulosa cell markers
such as *Fst* [66] and *Runx1* [10] which were highly conserved (Figure S10F). Additionally, we
identified open elements nearby markers of Sertoli cells, including structural markers such as
Cldn11 and *Itga6* (Figure S11A-B), hormonal markers such as *Amh* and *Hsd17b1* (Figure
S11C-D), and genes encoding for transcription factors such as *Fgf9* and *Nr5a1* (Figure S11E-
F). These were all bound by FOXL2 as shown by our ChIP-SICAP.
To investigate what transcription factors may bind to these dynamic regions that could
potentially drive the observed gene expression changes or that may direct changes in chromatin
organisation, we performed a transcription factor (TF) motif enrichment analysis of the DAR
dynamics (Figure 6F-G, Table S5). We selected only the significant DARs and used monaLisa
[67] to identify TF binding motifs enriched in accessible peaks. Regions that were more
accessible at E14.5, but lost accessibility at later stages (namely c7 and c5) were enriched with
nuclear factor family motifs (e.g. NFIC) and GATA motifs (GATA5, GATA3). NR5A1 motifs
were highly represented, as well as motifs from the RFX (RFX2, RFX5). Steroid receptor
motifs (ESR1 and ESRRB) were also enriched. Cluster 5 was also particularly enriched in
motifs typical of the transcriptional repressor CTCF. Regions accessible at 1W (c1 and c6)
were enriched in motifs for factors involved in Wnt signalling (TCFL5, TCF21) and Notch
signalling (HES1) and for ZBTB33 motifs, a transcriptional repressor also known as Kaiso,
which negatively regulates the Wnt signalling pathway [68]. Amongst these, TCF21, ATF1
were also found to localise with FOXL2 on chromatin, as we found by FOXL2 ChIP-SICAP
(Table S3 & Table S5). Clusters containing regions opened at 8W were highly enriched with
motifs for members of the steroid hormone receptor family, including ESRRB (containing a
half-site for the binding of ESR2) and AR, factors known to play critical roles in
folliculogenesis and ovulation [22, 69], which were also found in our proteomics analysis
(Figure 3). ESRRB motifs were also enriched at E14.5 although no expression of *Esr2* mRNA
was found at this timepoint by RNA-Seq (Figure 6G). Additionally, we noticed a remarkable
enrichment in motifs of members of the Jun and Fos family of transcription factors. Lastly, in
c2 we found enrichment for the TF NR5A2, known to regulate primordial follicle activation
[54], which was also detected with FOXL2 ChIP-SICAP at 1W and 8W timepoints (Figure 3).
RNA-Seq analysis revealed that some of these factors also show an expression pattern
matching the binding motif predictions (Figure 6G). For example, GATA4 and LHX9
expression was increased at E14.5, whilst factors predicted to bind at postnatal stages including
NR5A2, AR and ESR2, were upregulated from 1 week onwards.
586

DISCUSSION

Previous work has highlighted the importance of FOXL2 in many aspects of female sexual development[21] as well as in granulosa cell-related pathologies[70]. It is a sex-determining gene in goats [71], mutations of FOXL2 in humans lead to primary ovarian insufficiency [13], and it was shown to cooperate with other factors to establish ovarian sexual identity in the mouse embryonic gonad [72, 73]. Studies from our laboratory and that of Mathias Treier also demonstrated that FOXL2 becomes the sole gatekeeper of ovarian cell fate maintenance in adulthood [9].

In this study, we describe the set of FOXL2 genomic targets and on-chromatin protein interactors, as well as the gene expression changes and chromatin landscape dynamics characterising granulosa cells throughout mouse ovarian development. We found that FOXL2 binds to a greater number of genes at postnatal stages (1W and 8W) compared to embryonic (E14.5). The greatest number of targets overall was found at 1W, corresponding to the first timepoint at which ectopic SOX9 expression is observed in *Foxl2* null mutants [17]. These data support a more critical role for this factor during postnatal life compared to the embryo, at least in mice. GO analysis of FOXL2 targets at E14.5 showed enrichment of Wnt signalling, in keeping with the synergistic role played by these two pathways in reinforcing the female cell fate commitment in the embryonic gonad [72]. The Wnt pathway has recently been shown to be essential for pre-granulosa cell activation during primordial follicle activation [74]. This pathway was also highly featured amongst FOXL2 target genes identified at 1W, supporting the idea that a possible interplay between Wnt and FOXL2 around the time of primordial follicle activation may happen through regulation of similar target genes. Targets bound at 8W, once folliculogenesis is ongoing and mice have reached sexual maturity, are enriched for steroidogenesis and DNA repair pathways. Notably, we found FOXL2 to directly bind to a known ESR2 enhancer [27] specifically at 8W.

Indeed, in addition to binding sites for forkhead factors, FOXL2 peaks were highly enriched in sites for steroid receptors including ESR1 and AR. This supports previous studies performed in primary granulosa cell cultures which suggested a major role for FOXL2 on estradiol signalling through its direct regulation of *Esr2* [27].

One of the main roles of FOXL2 is direct repression of SOX9 [9]. Our data show that this repression may be time-specific, as we detected a greater number of FOXL2 peaks in the 'gene desert' regulatory region 5' to *Sox9* at 8W compared to earlier timepoints. Although this does not entirely exclude that the presence of a single peak at earlier timepoints may suffice and act as repressor site for *Sox9*. In agreement with other studies, we did not find binding of FOXL2 to TESCO [8, 21, 75], nor to the *Sox9*-specific Enh13 [38, 76], but we detected peaks in other regions, mostly not overlapping with Sertoli-specific *Sox9* enhancers [38]. The granulosa specific *Sox9* Enh8 [38] appeared significantly bound by FOXL2 only at 8W. Some binding seems to be present beforehand, however this was not considered significant because of our stringent selection criteria for ChIP binding events (that is, a peak was considered only if present above input in both replicates). FOXL2 appeared to bind to several Sertoli-enriched genes, various with structural roles (*Cldn11*, *Itga6*) or involved in hormone signalling (*Hsd17b1*, *Amh, Inhbb*). Comparison with published datasets of gene expression data obtained

from *Foxl2* null mutants ovaries [29], and from Sertoli and granulosa cells obtained at
embryonic and postnatal stages [41], also indicates a more prominent role of FOXL2
postnatally.
All together these data suggest that FOXL2 may repress testis development postnatally through
alternative *Sox9* enhancers, many of which are granulosa-specific, but also by directly
repressing genes important for Sertoli cell proliferation and function. The latter may as well be
the most important function of FOXL2 beyond repressing *Sox9*. This was suggested to be the
case in the conditional TRIM28 null mutant ovaries, where expression of Sertoli structural
genes precedes *Sox9* upregulation in an intermediate cell type along the female-to-male
transdifferentiation trajectory [8].
We then asked the question of which binding partners were co-localising on the same
chromatin regions with FOXL2 that could aid the multiple roles played by this factor along
ovarian development. Our ChIP-SICAP analysis represents the first *in vivo* analysis of
chromatin binding partners of FOXL2 and revealed that this factor interacts with proteins
crucial for each developmental milestone of ovarian development. For example, we found that
FOXL2 colocalises with the transcription coactivator of Wnt signalling Ctnnb1 (also known as
β-catenin) throughout the timecourse. It was previously shown that β-catenin stabilization and
activation in a normal XY testis leads to male-to-female sex-reversal [77] and is accompanied
by an increased expression of FOXL2 and reduced expression of SOX9 in the male gonad. The
co-localisation of β-catenin and FOXL2 on the same genomic locations observed in our study
may represent a mechanism to reinforce ovarian cell fate, suggesting that the functional
interplay between the two pathways may rely on their joint recruitment to DNA. This was also
supported by the identification of several other members of the pathway including TCF factors.
FOXL2 co-localised additionally with Wilms' tumour suppressor (WT1) at E14.5 and 1W.
WT1 is a transcriptional regulator essential for early development of the gonads [53] which is
found in two main splicing isoforms: +KTS, containing the amino acid trio KTS between the
last two zinc fingers, and -KTS, which lacks this. Importantly, only -KTS can act as a
transcriptional regulator because the addition of the +KTS portion abrogates WT1 ability to
bind to DNA, confining it to nuclear speckles [78, 79]. Given this knowledge and the fact that
our FOXL2 ChIP-SICAP experiments specifically identify only proteins bound to the
chromatin, which is typically absent from nuclear speckles [80], our data support an interplay
on the DNA of FOXL2 and the splicing isoform WT1 -KTS, which may aid the suggested role
of the latter in early ovarian development [81, 82]. We also identified interaction of FOXL2
with GATA4, which is essential for genital ridge formation in both sexes [52] and is known to
contribute to ovarian development and function [83, 84]. This is of interest because it was
previously shown that deficiency in GATA4-FOG2 interaction impaired ovarian development
through the alteration of the female gene expression program, including having a direct
negative effect on *Foxl2* expression [85]. Our data indicate that this novel FOXL2-GATA4
interaction may be at the base of GATA4's role in ovarian development.
FOXL2 is known to control the quiescence of primordial follicles by inhibiting the initiation
of the growth of oocytes. This is shown in *Foxl2* null mutants where the primordial follicles
form normally, but they do not develop into secondary follicles because pre-granulosa cells do
not undergo the necessary cuboidal transition [12]. FOXL2 was also shown to be differentially

expressed in subpopulations of pre-granulosa cells during primordial follicle formation being
downregulated in primary compared to primordial granulosa cells [20, 56].
In our study we identified four interactors known to have a role in primordial follicle activation
that had been presumed to be FOXL2-independent, namely NR5A2 [54], HDAC1 [86],
HDAC6 [55] and CSNK1A1[87]. This indicates that FOXL2 may create a chromatin hub
where factors regulating similar processes can interact. At 1 and 8W, we found FOXL2
interacting with ERS2, therefore confirming that the previously postulated interaction between
FOXL2 and estrogen signalling is mediated through the physical interaction of ESR2 and
FOXL2 at proximal genomic locations [27, 88]. In addition, we also detected AR at 1W in our
proteomics analysis, which also confirms the motif analysis of our ChIP-SICAP data showing
enrichment of AR motifs in the FOXL2 binding sites. Androgen receptor has a dual role in the
two sexes, regulating granulosa cell proliferation in the ovary [22], while it is critical to ensure
spermatogenesis is supported by Sertoli cells in the testis [89]. The co-localisation of AR and
FOXL2 may hint to a role for AR in the transdifferentiation phenotype observed in the
conditional *Foxl2* mutants [9]. Lastly, we also identified members of the DNA repair pathway,
such as XRCC6, which was also previously implicated in a newly discovered role played by
FOXL2 in DNA damage repair [50], alongside potential novel players such as PDS5B [90] and
DHX9 [91].
As part of the studies reported here, we generated a mouse line to allow the purification of
FOXL2-positive cells at all stages of ovarian development ($Foxl2^{EGFP}$). This valuable genetic
tool allowed us to generate age-matched transcriptome and chromatin landscape datasets to
integrate with our ChIP-SICAP data on the genomic binding sites of FOXL2 and be able to
draw conclusions as to which genes may be repressed or activated by this factor. FOXL2 is
also expressed in theca cells [59], therefore, we first assessed the expression values of known
theca and granulosa cell markers [60]. We found an enrichment of granulosa over theca
markers, confirming the suitability of this dataset for the exploration of granulosa-specific gene
regulatory networks. Our bulk RNA-Seq analysis shows expression of the pathways expected
to be activated at each developmental stage, notably an enrichment of WNT signalling and
urogenital development at E14.5, ECM remodelling and cell cycle at 1W, steroidogenesis and
angiogenesis at 8W.
By integrating the gene expression values with FOXL2 ChIP-SICAP we were able to identify
a high-confidence set of genes likely to be repressed, activated, or dynamically regulated by
FOXL2 across ovarian development. Our analysis showed that amongst all FOXL2 target
genes, the majority was either stably expressed throughout the timecourse, or dynamic. A lower
number of genes were bound by FOXL2 and stably repressed. Amongst the stably expressed
genes, we found genes with a role in mRNA processing, apoptosis and cell cycle, function
which have all been linked with FOXL2 in functional studies [92]. Analysis of those that are
dynamically expressed and controlled by FOXL2 provided an overview of the range of
functions critical for the ovary that this factor may regulate, from steroidogenesis and
extracellular matrix organisation to lipid catabolic processes. Genes involved in axon guidance
were also highlighted, which is of interest as the ovary is innervated, whereas the testis is not
[93].
A smaller number of genes was stably repressed, and this group included genes encoding for
Sertoli structural proteins suggesting that FOXL2 may inhibit the male pathway through the

regulation of genes critical for Sertoli cell function, beside regulating *Sox9* as previously
suggested by Uhlenhaut *et al.* [9]. Our findings are supported by another adult sex reversal
phenotype caused by gonadal deletion of TRIM28 in which authors found expression of
structural Sertoli cell markers preceding that of *Sox9* [8]. This evidence was further
corroborated by our ATAC-Seq analysis of the chromatin landscape of granulosa cells. Here,
we found that in agreement with the transcriptomic analysis, extensive chromatin remodelling
occurred across ovarian development. Importantly, we found that FOXL2 motifs were less
enriched in chromatin regions opened exclusively at E14.5, whilst their presence increased
significantly at 1 and 8W. This may suggest a more preponderant role of FOXL2 in regulating
genes postnatally, which agrees with the greater number of FOXL2 binding sites found at 1
and 8W compared to E14.5 This analysis led to the identification of several potential enhancers
of genes important for granulosa cell differentiation for example *Rspo1* and *Runx1*. Particularly
in the latter, we found several FOXL2 binding sites coinciding with open chromatin regions,
which is in agreement with the interplay between these two factors in regulating ovarian cell
fate maintenance [10]. Finally, the integration of ATAC-Seq with FOXL2 ChIP-SICAP and
comparison with ATAC-Seq datasets from sorted Sertoli cells [41, 94], also allowed us to
observe that genes important for Sertoli cells (*Amh*, *Cldn11*, *Itga6*, *Fgf9*, *Hsd17b1* amongst
others) contained regions opened in both Sertoli and granulosa cells and that these were bound
by FOXL2 in granulosa cells. This supports the concept that the role of FOXL2 in repressing
the male pathway may go beyond only repressing *Sox9* and may unfold additionally through
the direct repression of genes encoding proteins critical for the structure and function of Sertoli
cells.
In summary, our study represents the first description of the chromatin interactome of the
ovarian-specific transcription factor FOXL2 *in vivo*. We showed that this factor, essential for
female fertility, differentially interacts with proteins playing a role in a range of gonadal-
essential functions, from gonadogenesis, through primordial follicle activation, to
steroidogenesis and that this action is stage-specific. Further interrogation and mining of this
dataset may provide novel insights into the FOXL2-centric mechanisms at play in the ovary
during development and in the adult.

MATERIALS AND METHODS

Mice and timed mating

All animal experiments carried out were approved under the UK Animals (Scientific Procedures) Act 1986 and under the project licences n. 80/2405 and PP8826065. $Foxl2^{EGFP+/-}$ mice were maintained on a C57Bl6 background. $Foxl2^{EGFP+/-}$ embryos were produced by crossing heterozygotes with C57Bl6 wildtype animals and collected at E14.5 after timed matings. Noon of the day of plug was considered as E0.5. Litters of 1 week of age were sexed and females were culled to harvest the 1W timepoint. Fresh tissue was used for ATAC-Seq and RNA-Seq, whilst ovaries for ChIP-SICAP were snap-frozen and stored at -80°C until the day of the experiment.

Generation of $Foxl2^{EGFP}$ reporter strain

The $Foxl2^{EGFP}$ reporter strain was generated by the Genetic Modification Service at the Francis Crick Institute using CRISPR-Cas9 assisted targeting to insert a P2A and EGFP sequence after Exon 1 in frame of the $Foxl2$ sequence, followed by a stop codon. The donor vector contained a 783bp insert of P2A and eGFP, with 1kb homology arms either side, synthesised by ThermoFisher. The guide sequence used was 5'-TCTCTGAGTGCCAACGCGCG-3' and was cloned into the BbsI site of pX459 vector (Addgene #62988). The gene targeting was performed by co-transfection of pX459 and the synthetic donor vector into B6N 6.0 embryonic stem cells using Lipofectamine 2000. Clones were screened using an insertion PCR with primers within the EGFP (Table 1), followed by a long-range PCR with primers across the entire transgene (Table 1) and Sanger sequencing. Copy number was evaluated using digital droplet PCR (ddPCR) (Table 1 and Figure S4) and one correctly targeted homozygous ESC clone was microinjected into goGermline blastocysts [95], which were then transferred into pseudo pregnant females. The resulting chimeras were crossed to C57BL/6J albino (B6(Cg)-Tyr^{c-2J}/J) mice and offspring were validated for the correct integration of the transgene using the above screening PCRs followed by Sanger sequencing confirmation of the resulting amplicons. The ddPCR (Figure S4F) was performed for copy number evaluation of the F1 heterozygous mice and no evidence of random integrations of the transgene was found. Long read sequencing (Oxford Nanopore Technologies) was performed on two 3kb amplicons spanning the targeted insertion, generated by PCR and no aberrant on-target mutations or deletions were found 1.6kb upstream and 1.8kb downstream of the insertion.

Table 1. Primer Table

Primer/Probe	Sequence
Insert F	aggacgacggcaactacaag (within EGFP sequence)
Insert R	gtccatgccgagagtgatcc (within EGFP sequence)
LR-F	gagaagagagtgagagccgc (from sequence upstream of 5' homology arm)

LR-R	gggtatctttgccggtgtga (from sequence downstream of 3' homology arm)
eGFP ddPCR F	agcagaagaacggcatca
eGFP ddPCR R	gggtgttctgctggtagtg
eGFP ddPCR Probe	caagatccgccacaacatcgagga
ONT-F1	tgtccattcgtgactgggac (from sequence upstream of 5' homology arm)
ONT-R1	gccgtcgtccttgaagaaga (within EGFP sequence)
ONT-F2	aggacgacggcaactacaag (within EGFP sequence)
ONT-R2	gatgaagtgaggggtgcaca (from sequence downstream of 3' homology arm)

Immunofluorescence

Ovaries and testes were fixed after dissection by immersion in 4% PFA at 4ºC. Immunofluorescent stainings were performed on cryosections. The following primary antibodies were used: rabbit anti-SOX9 (Millipore, cat.#ab5535), rat anti-GFP (Nacalai Tesque 04404-84), rabbit anti-FOXL2 (a generous gift from Dr Dagmar Wilhelm). Slides were then incubated with Alexa Fluor secondary antibodies. Imaging of stained tissue sections was performed on a Leica SPE confocal microscope.

Tissue collection and chromatin preparation for ChIP-SICAP

Ovaries from E14.5, 1W and 8-weeks old C57BL/6 mice were dissected in PBS, snap-frozen, and stored at −80 °C. A pool of 140 embryonic E14.5 gonads, 30 1W ovaries, and 5 8W ovaries were used for each replicate. Chromatin was prepared using a modified version of the Active Motif High Sensitivity Chromatin Prep Kit protocol. Frozen tissue was pulverised under liquid nitrogen using mortar and pestle Bel-Art™ SP Scienceware™ Liquid Nitrogen-Cooled Mini Mortar. Ovaries were pooled to reach a minimum weight of 10 mg allowing a chromatin yield of at least 10 ug per replicate. Samples were fixed in 1 ml of Fixation Buffer (1.5 % methanol-free formaldehyde in 1% PBS) for 15 minutes on a roller at room temperature. Fixation was stopped using Stop buffer by Active Motif and incubating for 5 minutes. Chromatin preparation continued as per Active Motif protocol except for the sonication steps which were performed in a Bioruptor® Plus sonication device by Diagenode with the following settings: 40 cycles of 30"ON/30"OFF, power "high", constant temperature of 4ºC.

ChIP-SICAP

ChIP-SICAP experiments were performed in parallel using 2 biological replicates for each timepoint. ChIP-SICAP [18] allows the identification of chromatin-bound proteins that colocalise with a bait protein (FOXL2 in this study) on DNA. A total of 10 ug of sonicated chromatin was used for each immunoprecipitation using 3μl of FOXL2 antibody kindly donated by Dr Dagmar Wilhelm. The original ChIP-SICAP protocol [18, 19] was optimised to work on frozen tissue samples as follows. Chromatin was prepared with Active Motif

chromatin preparation kit as described above. After overnight immunoprecipitation at 4ºC with
a FOXL2-specific antibody, immune-protein complexes were captured on protein G
Dynabeads. DNA was biotinylated by TdT in the presence of biotin-11-ddUTP and biotin-11-
ddCTP as well as by Klenow3'exo in the presence of Biotin-7-dATP and eluted. Protein-DNA
complexes were captured with protease-resistant streptavidin beads [96], and proteins were
digested overnight at 37°C with 300 ng of LysC. Streptavidin beads carrying DNA were
separated using a magnet to be processed for ChIP-seq library preparation. The supernatant,
containing the digested peptides was collected and digested further by 8 hours of incubation
with 200 ng trypsin. Digested peptides were cleaned using the stage-tipping technique. Briefly,
digested samples were acidified by the addition of 2ul of TFA 10%. For each sample, 50 µL of
80% acetonitrile/0.1% formic acid was aliquoted in a LCMS Certified Clear Glass 12 x 32mm
Screw Neck Total Recovery Vial (Waters). ZipTip with 0.6 µL C18 resin was pre-treated by
pipetting 100% acetonitrile twice by aspirating, discarding and repeating. The ZipTip was
equilibrated by pipetting 0.1% TFA, three times by aspirating, discarding and repeating.
Acidified samples were then pipetted 10 times with ZipTip to allow peptides to bind to the
polymer within the tip. Following tip wash in 0.1% TFA, peptides were eluted in 80%
acetonitrile/0.1% formic acid in the glass vial and the eluent dried in a speed vac. Peptides were
reconstituted in 8 µL of 2% DMSO/0.1% formic acid. Peptides were separated on a 50 cm,
75 µm I.D. Pepmap column over a 70min gradient to be injected into the mass spectrometer
(Orbitrap Fusion Lumos) according to the universal Thermo Scientific HCD-IT method. The
instrument ran in data-dependent acquisition mode with the most abundant peptides selected
for MS/MS by HCD fragmentation. The raw data were analysed using Proteome Discoverer
2.1 (Thermo Scientific). Sequest HT node was used to search the spectra using the UniProt
Mus musculus database. The search parameters were as follows: the precursor mass tolerance
was 10 ppm, and MS/MS tolerance was 0.05 Da. Variable modification included Methionine
oxidation and N-terminal acetylation. Fixed modification included Cystenine
carbamidomethylation. Trypsin and LysC were chosen as the enzymes, and maximum 2 missed
cleavages were allowed. Perculator node was used to eliminate false identifications
(FDR < 0.01). Protein ratios in Foxl2 assays (n = 2) over no-antibody assays (n = 2) were
calculated in Proteome Discoverer. The quantification values were exported to R studio to
analyse significantly enriched proteins using t test limma package to determine Bayesian
moderated t-test p-values and Benjamini-Hochberg (BH) adjusted p values. We considered
proteins with mean fold enrichment >4 and adj. p value <0.1 as enriched proteins.

DNA purification for ChIP-SICAP-seq samples

Protease-resistant streptavidin beads carrying DNA samples from the above ChIP-SICAP were
processed for DNA purification. SDS buffer and proteinase K (20mg/ml) were added to
resuspend the beads followed by incubation for 30 minutes at 55°C. The beads were then heated
at 80°C for two hours to reverse the cross-linking. A magnetic rack was used to separate the
beads from the DNA, now in the supernatant. Supernatant was collected and 50 µL of Ampure
beads used to purify the DNA. Purified input (no SICAP) and ChIP-SICAP DNA samples were

used to generate libraries. NEB Ultra II DNA kit was used. Samples were sequenced on the
NovaSeq 6000 system. Paired-end sequencing was performed (30 million reads).

Ovary dissociation for fluorescence-activated cell sorting (FACS)

Ovaries were rinse in 1x PBS, then dissociated using add 450ul 0.05% trypsin and 50ul of 2.5%
collagenase. Tissue was incubated at 37°C for 6 minutes (for E14.5, up 20 minutes for 1W and
8 W ovaries after mincing). Enzymatic reaction was quenched by adding 200ul PBS/3% FBS.
Supernatant was removed without disrupting the gonads, then tissue clumps were resuspended
in 300ul PBS/3% FBS and gently pipetted up and down until gonads were disaggregated. Cell
mixture was passed through a 30 μm filter cap into a sorting tube. An extra 200ul of PBS/3%
FBS was used to rinse the tube and filtered. Samples were sorted with a FACS Aria III to
exclude debris (side scatter versus forward scatter) and doublets (area versus width) and to
select only live cells by excluding cells positive for DAPI. Cells were collected directly into a
low-retention 1.5 ml tube with 200ul PBS/3% FBS and kept on ice.

RNA extraction, library preparation and sequencing

A total of 50000 GFP positive cells were sorted, and RNA was extracted using the RNeasy
Plus Micro Kit (cat. no. 74034) by Qiagen. RNA quality was assessed by Bioanalyzer and
samples with a RIN value>8 were processed for library preparation. A total of 10ng of RNA
was used to prepare libraries using the NEBNext® Low Input RNA Library Prep Kit (E6420L).
Samples were subsequently sequenced on the NovaSeq 6000 system, at a dept of 25 million
reads (PE100) per sample.

ATAC-Seq, library preparation and sequencing

GFP positive cells isolated by FACS from gonads at E14.5, 1W and 8W (three biological
replicates per timepoint) were used for ATAC-Seq library preparation using Active Motif
ATAC-Seq Kit (cat. number 53150). In brief, 100 000 GFP positive sorted cells were washed
in PBS, pelleted, and lysed in ATAC Lysis Buffer to isolate intact nuclei. Tagmentation of
nuclei was performed for 30 minutes in a thermomixer at 37°C set at 800 rpm. Tagmented
DNA was purified using DNA purification columns and library amplification was performed
using Illumina's Nextera adapters. A unique combination of i7/i5 primers was used to allow
for multiplexing and sequencing on the same flow cell. PCR amplification included 5 minutes
incubation at 72°C, followed by 30 sec of DNA denaturation at 98°C and 10 cycles of: 98°C
for 10 sec, 63°C for 30 sec and 72°C for 1 minute. PCR products were cleaned up using SPRI
beads and size distribution of PCR amplified libraries was assessed on Tapestation. Samples
were subsequently sequenced on the NovaSeq 6000 system.

mRNA-Seq alignment and quantification

Reads were processed using the publicly available nf-core [97] rnseq pipeline v3.3 with the STAR [98]/RSEM [99] option against mouse genome assembly GRCm38 and Ensembl release 81 transcript annotations. Other options were left as default. Gene-level RSEM abundance estimates were imported into R using the Bioconductor package tximport's tximport function [100] for further analysis using DESeq2 [64]. Normalisation factors were calculated per gene using the default DESeq2 function, correcting for library size and feature length. Variance-stabilised abundance estimates were calculated using the VST function. PCA analysis was performed on the variance stabilised values using all available genes.

mRNA-Seq differential expression analysis and clustering

Differential expression between treatment groups was assessed using a DESeq2's Wald test in a pairwise manner, with false-discovery rate control based on an independent hypothesis weighting (IHW) (Ignatiadis et al 2016). Shrunken log2 fold changes were calculated using the lfcShrink function (type="ashr"). Significance was assessed based on a combined FDR<0.01, a shrunken log2FC>2 and a baseMean (i.e. mean abundance across all samples) >2. A combined list of significantly differentially expressed genes from all pairwise comparisons was created and used to generate a heatmap showing how patterns of expression changed across samples. Per-gene z-scores were calculated from the variance stabilised abundance estimates. K-means clustering was applied (km=7) to split the data into slices of similarly changing genes. Within each slice genes were clustered using a Euclidean distance metric and complete-linkage.

Gene Ontology enrichment analysis

Gene Ontology (GO) Biological Process (BP enrichment analysis of the individual gene lists and the kmeans clusters was calculated relative to a genomic background via the compareCluster function from the clusterProfiler Bioconductor package [101] (fun="enrichGO", OrgDb="org.Mm.eg.db", ont="BP", pAdjustMethod="BH", pvalueCutoff=0.05, qvalueCutoff=0.01, minGSSize=10, maxGSSize=500).

ChIP-SICAP

ChIP-Seq read alignment, peak calling and peak annotation

Reads were processed using the publicly available nf-core [97] chipseq pipeline v1.2.2 against the mouse genome assembly GRCm38 (--narrow_peak, --single_end, --deseq2_vst, min_reps_consensus 2, --macs_fdr 0.05), applying the Encode GRCm38 blacklist definitions to filter out problematic regions. A peak was only considered to be genuine if present in both biological replicates for each treatment group. A consensus peak set was then constructed by combining the peak calls from the individual treatment groups into a set of non-overlapping

intervals. Consensus peaks were annotated using the Bioconductor package ChIPpeakAnno's
annotatePeakInBatch function [102] against the protein-coding gene component of Ensembl
release 81. Gene assignments were made based on peak-centre minimal proximity to a gene's
TSS
(output="nearestLocation",multiple=FALSE,maxgap=1L,PeakLocForDistance="middle",Fea
tureLocForDistance="TSS",select="first",bindingType ="fullRange"). The distribution of
peaks mapping to genomic features was similarly assessed using ChIPpeakAnno's
"genomicElementDistribution" function, defining promoter regions as +/-2kb around a TSS
and downstream regions as the 2kb region downstream of genic boundaries.

DiffBind analysis

Consensus peak occupancy analysis was conducted using the Bioconductor package DiffBind
[103]. Briefly, reads were counted from the nfcore chipseq pipeline alignment files (BAM)
over a combined set of consensus peaks (intervals). Control (input) read counts were subtracted
from the targeted samples at each interval:

dba.count(samples.dba,peaks=consensus.gr,summits=FALSE,filter=0,bSubControl=TRUE,m
inCount=0,bUseSummarizeOverlaps=TRUE,mapQCth=0).

The resulting DiffBind object was normalised using a background approach, which calculates
scaled factors from read counts against a set of 15000bp genomic bins considered large enough
not to show differential enrichment across samples:

dba.normalize(samples.dba,background=TRUE, method=DBA_DESEQ2).

Normalised peak counts for all peaks were subsequently used for Pearson correlation analysis
and PCA analysis. Differential binding affinity analysis was conducted in a pairwise fashion
between treatment conditions using DESeq2 [64]. A FDR of <0.05 was used to threshold
significant changes in peak occupancy. Peak showing significant changes in occupancy were
submitted for read-depth profiling in the region +/-1.5kb from their centres. Plots are presented
at the merged replicate level, stratified based on gain/loss status (i.e. direction of fold change).

Clustering of ChIP-Seq peaks

The normalised peak occupancy estimates were used to cluster the consensus peaks into sets
that showed similar patterns of variation in occupancy across conditions. Peak estimates were
merged across replicates using a mean, a pseudo-count of 0.1 was added prior to a log2
transformation. Per-peak z-scores were calculated. K-means clustering was applied (km=7,
1000 repeats) to split the data into slices of similarly behaving peaks. Within each slice peaks
were clustered using a Euclidean distance metric and complete-linkage.

Motif enrichment analysis

Analysis was performed using the "calcBinnedMotifEnrR" from the monaLisa package [67], by scanning the sequence of consensus peaks +/-500bp from their centres against the PWMs from JASPAR 2020 [104]. Significance of enrichment was assessed against a background of equivalently sized sequences randomly sampled from the genome. Sequences were weighted to correct for GC and k-mer composition differences between fore- and background sets. Statistically significant enriched motifs within each cluster were determined using a one-tailed Fisher's exact test with Benjamini-Hochberg adjusted p-values pval<0.0001 and subsequently prioritised based on their log2 enrichment scores (descending).

ATAC-Seq

Read alignment and peak calling

Reads were processed using the publicly available nf-core [97] atacseq pipeline v1.2.1 against the mouse genome assembly GRCm38 (--deseq2_vst --min_reps_consensus), applying the Encode GRCm38 blacklist definitions to filter out problematic regions. A peak was only considered to be genuine if present in at least 2 biological replicates for each treatment group. A consensus peak set was then constructed by combining the peak calls from the individual treatment groups into a set of non-overlapping intervals.

Peak annotation

Consensus peaks were annotated using the Bioconductor package ChIPpeakAnno's annotatePeakInBatch function [102] against the protein-coding gene component of Ensembl release 81. Gene assignments were made based on peak-centre minimal proximity to a gene's TSS: (output="nearestLocation", multiple=FALSE, maxgap=-1L, PeakLocForDistance="middle", FeatureLocForDistance="TSS", select="first", bindingType="fullRange"). The distribution of peaks mapping to genomic features was similarly assessed using ChIPpeakAnno's genomicElementDistribution function, defining promoter regions as +/-2kb around a TSS and downstream regions as the 2kb region downstream of genic boundaries.

DiffBind analysis

Consensus peak occupancy analysis was conducted using the Bioconductor package DiffBind [103]. Briefly, reads were counted from the nfcore chipseq pipeline alignment files (BAM) over a combined set of consensus peaks (intervals) dba.count (samples.dba,peaks=consensus.gr,summits=FALSE,filter=0,bSubControl=FALSE,minCount =0,bUseSummarizeOverlaps=TRUE,mapQCth=0). The resulting DiffBind object was normalised DESeq2's RLE method [64] based on the peak counts dba.normalize(samples.dba,method=DBA_DESEQ2,normalize=DBA_NORM_RLE,library=

DBA_LIBSIZE_PEAKREADS). Normalised peak counts for all peaks were subsequently used
for Pearson correlation analysis and PCA analysis.

Differential binding affinity analysis was conducted in a pairwise fashion between treatment
conditions using DESeq2 [64]. A FDR of <0.05 was used to threshold significant changes in
peak occupancy. Peak showing significant changes in occupancy were submitted for read-
depth profiling in the region +/-1.5kb from their centres. Plots are presented at the merged
replicate level, stratified based on gain/loss status (i.e. direction of fold change).

1010

Clustering of peaks

The normalised peak occupancy estimates were used to cluster the consensus peaks into sets
that showed similar patterns of variation in occupancy across conditions. Peak estimates were
merged across replicates using a mean, a pseudo-count of 0.1 was added prior to a log2
transformation. Per-peak z-scores were calculated. K-means clustering was applied (km=7,
1000 repeats) to split the data into slices of similarly behaving peaks. Within each slice peaks
were clustered using a Euclidean distance metric and complete-linkage.

1018

Data integration

Data from the RNA, ATAC and ChIP experiments were integrated at the gene level. The
ATAC and ChIP data were assigned to protein-coding genes based on minimal peak proximity
to a TSS. Where this resulted in multiple peaks being assigned the same gene, the one closest
to the TSS was given precedence.

Identification of stably expressed and dynamically regulated genes: genes dynamically
regulated by FOXL22 were defined as the intersect of i) genes differentially expressed in at
least one of the pairwise comparisons from the RNA-seq analysis and ii) bound by Foxl2 based
on the presence of a ChIP-seq peak in at least 1 timepoint. Stably expressed genes were defined
as those bound by FOXL2 but now showing differential expression. Genes not bound by
FOXL2 but showing differential expression were not considered to be directly regulated by
FOXL2. Stably expressed genes were subsequently divided into "activated" and "repressed"
subsets based on the variance stabilised (VST) normalised counts from the RNA-seq analysis.
A VST score of >=8.3 in at least 1 sample was enough to quality a gene as "activated", while
all others were regarded as "repressed".

1034

1035 **Cytoscape visualisation of FOXL2 protein interactors**

1036 GO enrichment analysis of FOXL2 interactors identified by ChIP-SICAP and fold changes
1037 over no-antibody control in Table S3 were loaded onto Cytoscape v.3.9.1 [105]. Proteins were
1038 clustered based on their function. The cytoscape app enhancedGraphics [106] was used to
1039 overlay bar charts depicting the fold changes over control of each protein in each timepoint.

1040

Data and materials availability

All data are available in the main text or in the supplementary materials. Codes are available on request.

Aknowledgements

This work was supported by the Medical Research Council, U.K. (CC2116) and the Francis Crick Institute which receives its core funding from Cancer Research UK (CC2116), the UK Medical Research Council (CC2116), and the Wellcome Trust. RM holds a Postdoctoral Fellowship core-funded by the Francis Crick Institute (CC2116).

We are grateful for the support from present and past members of the Lovell-Badge lab. We are particularly grateful to Dr Nitzan Gonen (Bar-Ilan University), to Dr Francis Poulat (University of Montpellier), and to Dr Marie-Christine Chaboissier (Institut de Biologie Valrose) for scientific discussions and support.

We thank the Francis Crick institute platforms (GeMS, ASF, BABS, FLOW, Proteomics) for their excellent support and expertise. We thank Jacek Mor and all staff from the Biological Research facility (Francis Crick Institute) for taking excellent care of the animals. We thank Dr Dagmar Wilhelm for the generous gift of the anti-FOXL2 antibody.

Author contributions:

Conceptualization: RM

Visualization: RM

Investigation: RM, MN

Bioinformatic analyses: RM, RM, MRR

Methodology: RM, MN, MRR, SW, JO

Supervision: RM, RLB

Funding acquisition: RLB

Writing-original draft: RM

Writing-review&editing: RM and all authors

Competing interests

The authors declare no competing interests.

 SUPPLEMENTARY FIGURE LEGENDS

1077

Figure S1. FOXL2 genome-wide occupancy across ovarian development, Related to Figure
1.

(A) Annotation of consensus peaks by ChIPpeakAnno against the default iGenomes Ensembl
GTF file. (B) Enrichment analysis of transcription factor binding sites associated with
consensus peaks significantly changing across the timecourse. Analysis performed using the
calcBinnedMotifEnrR from the monaLisa package. Top 10 representative TF are shown (*p-*
value<0.0001). (C-E) Heatmaps of differentially bound peaks in each pairwise comparison.

1085

Figure S2. Comparison of FOXL2 target genes with DEGs identified in a *Foxl2$^{-/-}$* null
model (Ortiz *et al.*), Related to Figure 1.

(A) Venn diagram showing the overlap between FOXL2 target genes identified at any point
throughout ovarian development (E14.5, 1W and 8W) and genes downregulated in *Foxl2$^{-/-}$* null
mutant ovaries compared to wildtype controls (Ortiz *et al.*[29] and Table S2). Genomic
overviews of FOXL2 peaks in representative genes likely to be activated by FOXL2 including
Cdkn1b (B), *Smad7* (C), and *Fst* (D). (E) Venn diagram showing overlap between FOXL2
target genes and genes upregulated in in FOXL2^{KO} ovaries. Representative genes within this
overlap are shown and include *Wnt4* (F), *Gadd45g* (G), *Inhbb* (H) and *Sox9* (I). (J) Bar chart
displaying the total number of significant peaks identified in each timepoint in the 1Mb gene
desert upstream of *Sox9*.

1097

Figure S3 Cytoscape network of FOXL2 interactome across ovarian development,
Related to Figure 3.

(A) Proteins interacting with FOXL2 on-chromatin were clustered by GO Processes and
visualised with Cytoscape [107]. Enrichment values over no-antibody control were depicted as
barcharts (yellow=E14.5, orange=1W, blue=8W, n=2, fold change over no antibody control >4
in at least one timepoint, adj-*pvalue*<0.1) and visualised within the network using
enhancedGraphics Cytoscape plugin [106]. (B) Overlap of protein interactors found in our
study compared to a cell line (AT29c) whole proteome study by Penrad-Mobayed *et al.* [51].

1106

Figure S4. Design of *Foxl2^{EGFP}* construct and genotyping strategy to ensure correct
insertion of EGFP into *Foxl2* locus, Related to Figure 4.

(A) Design of gRNA (yellow box) and associated silent mutation (SM) in PAM sequence as
well as 783bp P2A-EGFP construct (green boxes). (B) Location of primers for insert, junction,
long-range and digital droplet PCR. (C) ES cell screening identified clone E11 (green arrows),
which showed successful EGFP integration depicted by a single band in insert, junction, and
long-range PCR. (D) Correct integration of EGFP into Foxl2 locus of clone E11 was confirmed
via sanger sequencing. (E) Long-range PCR of chimeric offspring as well as clone E11
confirmed heterozygous and homozygous EGFP insertion, respectively. No evidence of

backbone integration was found. BB pos= backbone positive; BB nrg= backbone negative. (F)
Copy number evaluation of chimeric offspring as well as ES cell clone E11 by ddPCR
confirmed absence of random transgene integration.

1119

Figure S5. Characterisation of the _Foxl2^{EGFP}_ mouse line, Related to Figure 4.

(A) Representative images of ovaries and testes dissected from adult mice (8 weeks) and
imaged with a fluorescence microscope. Upper row: GFP channel, lower row: brightfield
channel. Adult mouse ovaries wildtype (a-b) and heterozygous (c-d) as well as testes wildtype
(e-f) and heterozygous (g-h) are shown. Scale bars=100 µM. (B) Fertility assessment of
Foxl2$^{EGFP+/-}$ females compared to wildtype controls (C57BL/6J stock mice).

1126

Figure S6. Fluorescence-activated cell sorting of EGFP positive cells throughout ovarian
development, Related to Figure 4.

Gating strategy used to isolate live cells (A), singlets (B), and positive for GFP. An example
of FACS profile of a negative samples is shown in (C). (D-F) Dissociated cells were gated and
purified by FACS at E14.5 (D), 1W (E), and 8W (F). A total of 50k cells at each timepoint was
sorted for RNA-Seq and 100k for ATAC-Seq.

1133

Figure S7. Overview of RNA-Seq analysis of _Foxl2$^{EGFP+/-}$_ cells isolated throughout
ovarian development, Related to Figure 4 and Methods

(A) Poisson dissimilarity metric to assess RNA-Seq sample similarity, n=3 biological
replicates. (B) Boxplot illustrating the dynamics of gene expression changes granulosa
Foxl2$^{EGFP+/-}$ cells. (C) K-means clustering of the gene expression dynamics and representative
genes on the right-hand side. (D) Top terms from the GO enrichment test (Cluster Profiler)
showing the processes associated with DEGs from each of the seven clusters.

1141

Figure S8. Integration of FOXL2 ChIP-Seq data with RNA-Seq of _Foxl2$^{EGFP+/-}$_ cells,
Related to Figure 5.

(A) Plot of gene density and VST cut-off =8.3 used to distinguish genes stably repressed from
those stably expressed. (B) Barplot depicting the number of genes either stably repressed or
activated and bound by FOXL2. (C) Example of gene expression changes used to choose the
VST cut-off. The male marker _Fgf9_ was used as reference for the cut-off as known to not be t
expressed in the ovary.

1149

Figure S9. ATAC-Seq analysis of chromatin accessibility of _Foxl2^{EGFP}_ positive cells
collected across ovarian development, Related to Figure 6.

(A) Poisson metric showing peak occupancy correlation between timepoints (E14.5, 1W and
8W). (B) Genomic annotation of consensus peaks obtained with ChIPpeakAnno [102] against

the default iGenomes Ensembl GTF file. (C) Boxplot of z-scores representing overall trends of
chromatin opening dynamics over time within each cluster derived from the clustering of
normalised abundancies of peaks. (D) Heatmap of enrichment scores for the FOXL2 canonical
motif in each cluster, ranked on enrichment score and filtered by -log10pval. Cut-off: -
log10pval > 4 (i.e. pval<0.0001).

1159

Figure S10. Overview of differential accessibility analysis of ATAC-Seq data, Related to
Figure 6.

(A-C) Density plots of peaks differentially opened in each pairwise comparison. (D) Summary
of shared and unique open chromatin regions across the timecourse. (E-F) IGV snapshots
representative of several putative enhancers identified within *Rspo1*, *Fst*, *Runx1* and *Sox9*
genes, as assessed by ATAC-Seq and FOXL2 ChIP-SICAP. UCSC Genome Brower
conservation analysis in pop-up squares. Black tracks show E13.5 Sertoli ATAC-Seq data re-
analysed from Garcia-Moreno *et al*. [94], and P7 Sertoli ATAC-Seq from Lindeman *et al.* [41].

1168

Figure S11. Overview of differential accessibility analysis of ATAC-Seq data, Sertoli-
specific genes, Related to Figure 6.

IGV snapshots representative of several putative enhancers identified within Sertoli-enriched
genes. (A) *Cldn11*, (B) *Itga6*, (C) *Amh*, (D) *Hsd17b1*, (E) *Nr5a1*, (F) *Fgf9*. Black tracks show
E13.5 Sertoli ATAC-Seq data re-analysed from Garcia-Moreno *et al*.[94], and P7 Sertoli
ATAC-Seq from Lindeman *et al.* [41].

1175

Sup. Table 1: FOXL2 ChIP-SICAP analysis of genomic targets, GO of clusters and diffbind
analysis, relative to Figure 1 and 2.

Sup. Table 2: Comparison of FOXL2 ChIP-SICAP with other datasets from Jameson *et*
al.[40], Lindeman *et al.[41]*, and Ortiz *et al.[29]*, relative to Figure 2.

Sup. Table 3: FOXL2 ChIP-SICAP, proteomics dataset of FOXL2 interactors, relative to
Figure 3.

Sup. Table 4: RNA-Seq of *Foxl2$^{EGFP+/-}$* cells isolated throughout ovarian development**,**
relative to Figure 4.

Sup. Table 5: Data integration of ChIP-Seq, ATAC and RNA-Seq and gene-level dynamics,
relative to Figure 5.

Sup. Table 6: ATAC-Seq normalised peak counts, diffbind analysis**,** clustering analysis.

References

1. Bunce, C., J. McKey, and B. Capel, *Concerted morphogenesis of genital ridges and nephric ducts in the mouse captured through whole-embryo imaging.* Development, 2021. **148**(18).
2. Migale, R., M. Neumann, and R. Lovell-Badge, *Long-Range Regulation of Key Sex Determination Genes.* Sex Dev, 2021. **15**(5-6): p. 360-380.
3. Koopman, P., et al., *Male development of chromosomally female mice transgenic for Sry.* Nature, 1991. **351**(6322): p. 117-21.
4. Hacker, A., et al., *Expression of Sry, the mouse sex determining gene.* Development, 1995. **121**(6): p. 1603-14.
5. Bullejos, M. and P. Koopman, *Spatially dynamic expression of Sry in mouse genital ridges.* Dev Dyn, 2001. **221**(2): p. 201-5.
6. Vidal, V.P., et al., *Sox9 induces testis development in XX transgenic mice.* Nat Genet, 2001. **28**(3): p. 216-7.
7. Yao, H.H., *The pathway to femaleness: current knowledge on embryonic development of the ovary.* Mol Cell Endocrinol, 2005. **230**(1-2): p. 87-93.
8. Rossitto, M., et al., *TRIM28-dependent SUMOylation protects the adult ovary from activation of the testicular pathway.* Nat Commun, 2022. **13**(1): p. 4412.
9. Uhlenhaut, N.H., et al., *Somatic sex reprogramming of adult ovaries to testes by FOXL2 ablation.* Cell, 2009. **139**(6): p. 1130-42.
10. Nicol, B., et al., *RUNX1 maintains the identity of the fetal ovary through an interplay with FOXL2.* Nat Commun, 2019. **10**(1): p. 5116.
11. Tucker, E.J., *The Genetics and Biology of FOXL2.* Sex Dev, 2022. **16**(2-3): p. 184-193.
12. Schmidt, D., et al., *The murine winged-helix transcription factor Foxl2 is required for granulosa cell differentiation and ovary maintenance.* Development, 2004. **131**(4): p. 933-42.
13. Crisponi, L., et al., *The putative forkhead transcription factor FOXL2 is mutated in blepharophimosis/ptosis/epicanthus inversus syndrome.* Nat Genet, 2001. **27**(2): p. 159-66.
14. Harris, S.E., et al., *Identification of novel mutations in FOXL2 associated with premature ovarian failure.* Mol Hum Reprod, 2002. **8**(8): p. 729-33.
15. De Baere, E., et al., *Spectrum of FOXL2 gene mutations in blepharophimosis-ptosis-epicanthus inversus (BPES) families demonstrates a genotype--phenotype correlation.* Hum Mol Genet, 2001. **10**(15): p. 1591-600.
16. Uda, M., et al., *Foxl2 disruption causes mouse ovarian failure by pervasive blockage of follicle development.* Hum Mol Genet, 2004. **13**(11): p. 1171-81.
17. Ottolenghi, C., et al., *Foxl2 is required for commitment to ovary differentiation.* Hum Mol Genet, 2005. **14**(14): p. 2053-62.
18. Rafiee, M.R., et al., *Expanding the Circuitry of Pluripotency by Selective Isolation of Chromatin-Associated Proteins.* Mol Cell, 2016. **64**(3): p. 624-635.
19. Rafiee, M.R. and J. Krijgsveld, *Using ChIP-SICAP to Identify Proteins That Co-localize in Chromatin.* Methods Mol Biol, 2021. **2351**: p. 275-288.
20. Niu, W. and A.C. Spradling, *Two distinct pathways of pregranulosa cell differentiation support follicle formation in the mouse ovary.* Proc Natl Acad Sci U S A, 2020. **117**(33): p. 20015-20026.
21. Georges, A., et al., *FOXL2: a central transcription factor of the ovary.* J Mol Endocrinol, 2014. **52**(1): p. R17-33.
22. Franks, S. and K. Hardy, *Androgen Action in the Ovary.* Front Endocrinol (Lausanne), 2018. **9**: p. 452.
23. Drummond, A.E. and P.J. Fuller, *The importance of ERbeta signalling in the ovary.* J Endocrinol, 2010. **205**(1): p. 15-23.
24. Bashamboo, A. and K. McElreavey, *NR5A1/SF-1 and development and function of the ovary.* Ann Endocrinol (Paris), 2010. **71**(3): p. 177-82.

25. Park, M., et al., *FOXL2 interacts with steroidogenic factor-1 (SF-1) and represses SF-1-induced*
CYP17 transcription in granulosa cells. Mol Endocrinol, 2010. **24**(5): p. 1024-36.
26. Shen, W.H., et al., *Nuclear receptor steroidogenic factor 1 regulates the mullerian inhibiting*
substance gene: a link to the sex determination cascade. Cell, 1994. **77**(5): p. 651-61.
27. Georges, A., et al., *The transcription factor FOXL2 mobilizes estrogen signaling to maintain the*
identity of ovarian granulosa cells. Elife, 2014. **3**.
28. Chassot, A.A., et al., *WNT4 and RSPO1 together are required for cell proliferation in the early*
mouse gonad. Development, 2012. **139**(23): p. 4461-72.
29. Garcia-Ortiz, J.E., et al., *Foxl2 functions in sex determination and histogenesis throughout*
mouse ovary development. BMC Dev Biol, 2009. **9**: p. 36.
30. Gustin, S.E., et al., *WNT/beta-catenin and p27/FOXL2 differentially regulate supporting cell*
proliferation in the developing ovary. Dev Biol, 2016. **412**(2): p. 250-60.
31. Granados-Aparici, S., et al., *SMAD3 directly regulates cell cycle genes to maintain arrest in*
granulosa cells of mouse primordial follicles. Sci Rep, 2019. **9**(1): p. 6513.
32. Yao, H.H., W. Whoriskey, and B. Capel, *Desert Hedgehog/Patched 1 signaling specifies fetal*
Leydig cell fate in testis organogenesis. Genes Dev, 2002. **16**(11): p. 1433-40.
33. Parma, P., et al., *R-spondin1 is essential in sex determination, skin differentiation and*
malignancy. Nat Genet, 2006. **38**(11): p. 1304-9.
34. Vainio, S., et al., *Female development in mammals is regulated by Wnt-4 signalling.* Nature,
1999. **397**(6718): p. 405-9.
35. Johnen, H., et al., *Gadd45g is essential for primary sex determination, male fertility and testis*
development. PLoS One, 2013. **8**(3): p. e58751.
36. Rodriguez, K.F., et al., *Somatic cell fate maintenance in mouse fetal testes via*
autocrine/paracrine action of AMH and activin B. Nat Commun, 2022. **13**(1): p. 4130.
37. Sekido, R., et al., *SOX9 is up-regulated by the transient expression of SRY specifically in Sertoli*
cell precursors. Dev Biol, 2004. **274**(2): p. 271-9.
38. Gonen, N., et al., *Sex reversal following deletion of a single distal enhancer of Sox9.* Science,
2018. **360**(6396): p. 1469-1473.
39. Sekido, R. and R. Lovell-Badge, *Sex determination involves synergistic action of SRY and SF1 on*
a specific Sox9 enhancer. Nature, 2008. **453**(7197): p. 930-4.
40. Jameson, S.A., et al., *Temporal transcriptional profiling of somatic and germ cells reveals*
biased lineage priming of sexual fate in the fetal mouse gonad. PLoS Genet, 2012. **8**(3): p.
e1002575.
41. Lindeman, R.E., et al., *The conserved sex regulator DMRT1 recruits SOX9 in sexual cell fate*
reprogramming. Nucleic Acids Res, 2021. **49**(11): p. 6144-6164.
42. Chassot, A.A., et al., *Activation of beta-catenin signaling by Rspo1 controls differentiation of*
the mammalian ovary. Hum Mol Genet, 2008. **17**(9): p. 1264-77.
43. Wang, D.S., et al., *Foxl2 up-regulates aromatase gene transcription in a female-specific*
manner by binding to the promoter as well as interacting with ad4 binding
protein/steroidogenic factor 1. Mol Endocrinol, 2007. **21**(3): p. 712-25.
44. Pisarska, M.D., et al., *Forkhead l2 is expressed in the ovary and represses the promoter activity*
of the steroidogenic acute regulatory gene. Endocrinology, 2004. **145**(7): p. 3424-33.
45. Dupont, S., et al., *Effect of single and compound knockouts of estrogen receptors alpha*
(ERalpha) and beta (ERbeta) on mouse reproductive phenotypes. Development, 2000. **127**(19):
p. 4277-91.
46. Fu, A., et al., *IRX3 and IRX5 collaborate during ovary development and follicle formation to*
establish responsive granulosa cells in the adult mousedagger. Biol Reprod, 2020. **103**(3): p.
620-629.
47. Cai, K., et al., *Action mechanism of inhibin alpha-subunit on the development of Sertoli cells*
and first wave of spermatogenesis in mice. PLoS One, 2011. **6**(10): p. e25585.

48.   Obholz, K.L., et al., *FNDC3A is required for adhesion between spermatids and Sertoli cells.* Dev
Biol, 2006. **298**(2): p. 498-513.
49.   Gao, F., et al., *The Wilms tumor gene, Wt1, is required for Sox9 expression and maintenance*
of tubular architecture in the developing testis. Proc Natl Acad Sci U S A, 2006. **103**(32): p.
11987-92.
50.   Jin, H., et al., *FOXL2 directs DNA double-strand break repair pathways by differentially*
interacting with Ku. Nat Commun, 2020. **11**(1): p. 2010.
51.   Penrad-Mobayed, M., et al., *Conventional and unconventional interactions of the transcription*
factor FOXL2 uncovered by a proteome-wide analysis. FASEB J, 2020. **34**(1): p. 571-587.
52.   Hu, Y.C., L.M. Okumura, and D.C. Page, *Gata4 is required for formation of the genital ridge in*
mice. PLoS Genet, 2013. **9**(7): p. e1003629.
53.   Kreidberg, J.A., et al., *WT-1 is required for early kidney development.* Cell, 1993. **74**(4): p. 679-
91.
54.   Meinsohn, M.C., et al., *A role for orphan nuclear receptor liver receptor homolog-1 (LRH-1,*
NR5A2) in primordial follicle activation. Sci Rep, 2021. **11**(1): p. 1079.
55.   Zhang, T., et al., *HDAC6 regulates primordial follicle activation through mTOR signaling*
pathway. Cell Death Dis, 2021. **12**(6): p. 559.
56.   Ernst, E.H., et al., *Granulosa cells from human primordial and primary follicles show differential*
global gene expression profiles. Hum Reprod, 2018. **33**(4): p. 666-679.
57.   Leon, N.Y. and V.R. Harley, *ATR-X syndrome: genetics, clinical spectrum, and management.*
Hum Genet, 2021. **140**(12): p. 1625-1634.
58.   L'Hote, D., et al., *Discovery of novel protein partners of the transcription factor FOXL2 provides*
insights into its physiopathological roles. Hum Mol Genet, 2012. **21**(14): p. 3264-74.
59.   Zhou, J., et al., *Dissecting the fate of Foxl2-expressing cells in fetal ovary using lineage tracing*
and single-cell transcriptomics. Cell Discov, 2022. **8**(1): p. 139.
60.   Morris, M.E., et al., *A single-cell atlas of the cycling murine ovary.* Elife, 2022. **11**.
61.   Hsu, S.H., et al., *Spz1, a novel bHLH-Zip protein, is specifically expressed in testis.* Mech Dev,
2001. **100**(2): p. 177-87.
62.   Stevant, I., et al., *Deciphering Cell Lineage Specification during Male Sex Determination with*
Single-Cell RNA Sequencing. Cell Rep, 2018. **22**(6): p. 1589-1599.
63.   Maatouk, D.M., et al., *Genome-wide identification of regulatory elements in Sertoli cells.*
Development, 2017. **144**(4): p. 720-730.
64.   Love, M.I., W. Huber, and S. Anders, *Moderated estimation of fold change and dispersion for*
RNA-seq data with DESeq2. Genome Biol, 2014. **15**(12): p. 550.
65.   Consortium, E.P., et al., *Expanded encyclopaedias of DNA elements in the human and mouse*
genomes. Nature, 2020. **583**(7818): p. 699-710.
66.   Jorgez, C.J., et al., *Granulosa cell-specific inactivation of follistatin causes female fertility*
defects. Mol Endocrinol, 2004. **18**(4): p. 953-67.
67.   Machlab, D., et al., *monaLisa: an R/Bioconductor package for identifying regulatory motifs.*
Bioinformatics, 2022. **38**(9): p. 2624-2625.
68.   Kim, S.W., et al., *Non-canonical Wnt signals are modulated by the Kaiso transcriptional*
repressor and p120-catenin. Nat Cell Biol, 2004. **6**(12): p. 1212-20.
69.   Couse, J.F., et al., *Estrogen receptor-beta is critical to granulosa cell differentiation and the*
ovulatory response to gonadotropins. Endocrinology, 2005. **146**(8): p. 3247-62.
70.   Shah, S.P., et al., *Mutation of FOXL2 in granulosa-cell tumors of the ovary.* N Engl J Med, 2009.
360(26): p. 2719-29.
71.   Boulanger, L., et al., *FOXL2 is a female sex-determining gene in the goat.* Curr Biol, 2014. **24**(4):
p. 404-8.
72.   Ottolenghi, C., et al., *Loss of Wnt4 and Foxl2 leads to female-to-male sex reversal extending*
to germ cells. Hum Mol Genet, 2007. **16**(23): p. 2795-804.

73. Auguste, A., et al., *Loss of R-spondin1 and Foxl2 amplifies female-to-male sex reversal in XX*
mice. Sex Dev, 2011. **5**(6): p. 304-17.
74. Habara, O., et al., *WNT signaling in pre-granulosa cells is required for ovarian folliculogenesis*
and female fertility. Development, 2021. **148**(9).
75. Nicol, B., et al., *Genome-wide identification of FOXL2 binding and characterization of FOXL2*
feminizing action in the fetal gonads. Hum Mol Genet, 2018. **27**(24): p. 4273-4287.
76. Croft, B., et al., *Human sex reversal is caused by duplication or deletion of core enhancers*
upstream of SOX9. Nat Commun, 2018. **9**(1): p. 5319.
77. Maatouk, D.M., et al., *Stabilization of beta-catenin in XY gonads causes male-to-female sex-*
reversal. Hum Mol Genet, 2008. **17**(19): p. 2949-55.
78. Englert, C., et al., *Truncated WT1 mutants alter the subnuclear localization of the wild-type*
protein. Proc Natl Acad Sci U S A, 1995. **92**(26): p. 11960-4.
79. Larsson, S.H., et al., *Subnuclear localization of WT1 in splicing or transcription factor domains*
is regulated by alternative splicing. Cell, 1995. **81**(3): p. 391-401.
80. Spector, D.L. and A.I. Lamond, *Nuclear speckles.* Cold Spring Harb Perspect Biol, 2011. **3**(2).
81. Barbaux, S., et al., *Donor splice-site mutations in WT1 are responsible for Frasier syndrome.*
Nat Genet, 1997. **17**(4): p. 467-70.
82. Hammes, A., et al., *Two splice variants of the Wilms' tumor 1 gene have distinct functions*
during sex determination and nephron formation. Cell, 2001. **106**(3): p. 319-29.
83. Heikinheimo, M., et al., *Expression and hormonal regulation of transcription factors GATA-4*
and GATA-6 in the mouse ovary. Endocrinology, 1997. **138**(8): p. 3505-14.
84. Efimenko, E., et al., *The transcription factor GATA4 is required for follicular development and*
normal ovarian function. Dev Biol, 2013. **381**(1): p. 144-58.
85. Manuylov, N.L., et al., *Ovarian development in mice requires the GATA4-FOG2 transcription*
complex. Development, 2008. **135**(22): p. 3731-43.
86. Ma, P., et al., *Compensatory functions of histone deacetylase 1 (HDAC1) and HDAC2 regulate*
transcription and apoptosis during mouse oocyte development. Proc Natl Acad Sci U S A, 2012.
109(8): p. E481-9.
87. Zhang, D., et al., *Oocyte Casein kinase 1alpha deletion causes defects in primordial follicle*
formation and oocyte loss by impairing oocyte meiosis and enhancing autophagy in developing
mouse ovary. Cell Death Discov, 2022. **8**(1): p. 388.
88. Herman, L., et al., *Genomic exploration of the targets of FOXL2 and ESR2 unveils their*
implication in cell migration, invasion, and adhesion. FASEB J, 2021. **35**(4): p. e21355.
89. De Gendt, K., et al., *A Sertoli cell-selective knockout of the androgen receptor causes*
spermatogenic arrest in meiosis. Proc Natl Acad Sci U S A, 2004. **101**(5): p. 1327-32.
90. Morales, C., et al., *PDS5 proteins are required for proper cohesin dynamics and participate in*
replication fork protection. J Biol Chem, 2020. **295**(1): p. 146-157.
91. Chakraborty, P. and K. Hiom, *DHX9-dependent recruitment of BRCA1 to RNA promotes DNA*
end resection in homologous recombination. Nat Commun, 2021. **12**(1): p. 4126.
92. Benayoun, B.A., et al., *Transcription factor FOXL2 protects granulosa cells from stress and*
delays cell cycle: role of its regulation by the SIRT1 deacetylase. Hum Mol Genet, 2011. **20**(9):
p. 1673-86.
93. McKey, J., et al., *Neural crest-derived neurons invade the ovary but not the testis during mouse*
gonad development. Proc Natl Acad Sci U S A, 2019. **116**(12): p. 5570-5575.
94. Garcia-Moreno, S.A., et al., *Gonadal supporting cells acquire sex-specific chromatin landscapes*
during mammalian sex determination. Dev Biol, 2019. **446**(2): p. 168-179.
95. Koentgen, F., et al., *Exclusive transmission of the embryonic stem cell-derived genome through*
the mouse germline. Genesis, 2016. **54**(6): p. 326-33.
96. Rafiee, M.R., et al., *Protease-resistant streptavidin for interaction proteomics.* Mol Syst Biol,
2020. **16**(5): p. e9370.

97.   Ewels, P.A., et al., *The nf-core framework for community-curated bioinformatics pipelines.* Nat
1388   Biotechnol, 2020. **38**(3): p. 276-278.
1389   98.   Dobin, A., et al., *STAR: ultrafast universal RNA-seq aligner.* Bioinformatics, 2013. **29**(1): p. 15-
1390   21.
1391   99.   Li, B. and C.N. Dewey, *RSEM: accurate transcript quantification from RNA-Seq data with or
1392   without a reference genome.* BMC Bioinformatics, 2011. **12**: p. 323.
1393   100.   Soneson, C., M.I. Love, and M.D. Robinson, *Differential analyses for RNA-seq: transcript-level
1394   estimates improve gene-level inferences.* F1000Res, 2015. **4**: p. 1521.
1395   101.   Wu, T., et al., *clusterProfiler 4.0: A universal enrichment tool for interpreting omics data.*
1396   Innovation (Camb), 2021. **2**(3): p. 100141.
1397   102.   Zhu, L.J., et al., *ChIPpeakAnno: a Bioconductor package to annotate ChIP-seq and ChIP-chip
1398   data.* BMC Bioinformatics, 2010. **11**: p. 237.
1399   103.   Ross-Innes, C.S., et al., *Differential oestrogen receptor binding is associated with clinical
1400   outcome in breast cancer.* Nature, 2012. **481**(7381): p. 389-93.
1401   104.   Fornes, O., et al., *JASPAR 2020: update of the open-access database of transcription factor
1402   binding profiles.* Nucleic Acids Res, 2020. **48**(D1): p. D87-D92.
1403   105.   Shannon, P., et al., *Cytoscape: a software environment for integrated models of biomolecular
1404   interaction networks.* Genome Res, 2003. **13**(11): p. 2498-504.
1405   106.   Morris, J.H., et al., *enhancedGraphics: a Cytoscape app for enhanced node graphics.* F1000Res,
1406   2014. **3**: p. 147.
1407   107.   Cline, M.S., et al., *Integration of biological networks and gene expression data using
1408   Cytoscape.* Nat Protoc, 2007. **2**(10): p. 2366-82.

1409