<u>**Marine transmissible cancer navigates urbanised waters, threatening to spillover**</u>

Hammel M. [1,2*], Touchard F. [1], Burioli E. A. V [1,2], Paradis L.[1], Cerqueira F[1]., Chailler E.[1],

Bernard I.[3], Cochet H.[4], Simon A.[1], Thomas F.[5], Destoumieux-Garzón D.[2], Charrière G.

M.[2], Bierne N.[1]

[1] ISEM, Univ Montpellier, CNRS, IRD, Montpellier, France

[2] IHPE, Univ Montpellier, CNRS, Ifremer, Univ Perpignan Via Domitia, Montpellier, France

[3] Eurêka Mer, Lézardrieux, France

[4] Cochet Environnement, 56550 Locoal, France

[5] CREEC/CANECEV (CREES), MIVEGEC, Unité Mixte de Recherches, IRD 224–CNRS

5290–Université de Montpellier, Montpellier, France

*Corresponding author address: maurine.hammel@umontpellier.fr

Abstract

Inter-individual transmission of cancer cells represents a unique form of microparasites

increasingly reported in marine bivalves. In this study, we sought to understand the ecology of

the propagation of *Mytilus trossulus* Bivalve Transmissible Neoplasia 2 (MtrBTN2), a

transmissible cancer affecting four *Mytilus* mussel species worldwide. We investigated the

prevalence of MtrBTN2 in the mosaic hybrid zone of *M. edulis* and *M. galloprovincialis* along

the French Atlantic coast, sampling contrasting natural and anthropogenic habitats. We observed

a similar prevalence in both species, likely due to the spatial proximity of the two species in this

region. Our results showed that ports had higher prevalence of MtrBTN2, with a possible hotspot

observed at a shuttle landing dock. No cancer was found in natural beds except for two sites

close to the hotspot, suggesting spillover. Ports may provide favourable conditions for the

transmission of MtrBTN2, such as high mussel density, stressful conditions, sheltered and

confined shores, or buffered temperatures. Ships may also spread the disease through biofouling. Our results suggest ports may serve as epidemiological hubs, with maritime routes providing artificial gateways for MtrBTN2 propagation. This highlights the importance of preventing biofouling on docks and ship hulls to limit the spread of marine pathogens hosted by fouling species.

Keywords: Transmissible cancer, Bivalve Transmissible Neoplasia, epidemiology, biofouling, spillover, *Mytilus* mussels.

1. Introduction

Transmissible cancers are caused by malignant cell lineages that have acquired the ability to infect new hosts through the transmission of living cancer cells. Fourteen transmissible cancer lineages have been described to date: one in dogs (canine transmissible venereal tumour, CTVT [1, 2]), two in Tasmanian devils (devil facial tumour, DFT1 and DFT2 [3, 4]), and eleven in different marine bivalve species (bivalve transmissible neoplasia, BTNs [5–10]). While direct contact is necessary for CTVT and DFTs transmission, via coitus for the former and biting for the latter, the transmission of BTNs between shellfish is assumed to occur through the seawater column [11, 12]. *Mytilus* spp. mussels are affected by two BTN lineages known as MtrBTN1 and MtrBTN2 that originate from two different *M. trossulus* founder hosts. MtrBTN2 is distributed worldwide and has crossed the species barrier several times to infect four *Mytilus* species: *M. trossulus* in East Asia and Northern Europe [13–15], *M. chilensis* in South America [9], and *M. edulis* and *M. galloprovincialis* in Europe [9, 16]. Cancerous process in mussels was firstly described fifty years ago as disseminated neoplasia (DN, [17]) but the demonstration that some DNs are transmissible is recent [6, 7]. As non-transmissible neoplasia also occurs in mussels [9, 16], a genetic diagnosis is required to ensure the identification of MtrBTNs. The oldest known genetically validated MtrBTN2 was sampled in 2009 [9]. Genetic data suggest that it could be at least one or two orders of magnitude more ancient, although dating is difficult due to an apparent acceleration of the mitochondrial clock [15, 16].

The global distribution of the MtrBTN2 lineage remains enigmatic [9, 14]. The other known
transmissible cancer with such a worldwide distribution is CTVT in dogs. CTVT emerged 4,000
to 8,000 years ago, probably in Eastern Asia, and has expanded rapidly worldwide over the past
500 years, probably facilitated by the development of maritime transportation [18]. Transmission
of MtrBTN2 within mussel populations occurs presumably through filter-feeding (discussed in
[11, 19]). MtrBTN2 cells can survive for several days in seawater [11], which greatly increases
their chances of infecting new hosts. However, it is unlikely that transport of cancer cells by
marine currents alone explains such a global distribution. Indeed, the dilution of cancer cells in
the marine environment may strongly limit the success of transmission over long distances.
Shipping traffic has been proposed as the most likely explanation for the global distribution of
MtrBTN2 -i.e., the transport of disease-carrying mussels on ship hulls [9, 14, 15]. To our
knowledge, these claims have remained speculative and no formal evidence of the effect of
shipping traffic and port habitat on the epidemiology of BTN has been provided to date.
69
European mussels are affected by MtrBTN2, with a much higher prevalence in *M. edulis* (of the
order of 1/100) than in *M. galloprovincialis* (of the order of 1/1000) [16]. These two host species
are parapatric in Europe and can coexist in contact zones where hybridisation is taking place
[20–22]. Associations between genetic backgrounds and environmental variables indicate that
the environment partly influences the structure and maintenance of the hybrid zones: in the study
area, *M. edulis* genotypes are more frequent in sheltered habitats under freshwater influence,
while *M. galloprovincialis* genotypes are more frequent in habitats exposed to wave action and
ocean salinities [23, 24]. In this context, the distribution and propagation of MtrBTN2 is likely to
be influenced by host genetic background and population composition [16], as well as by other
environmental factors such as population density, reduced hydrodynamics, and stressful
environmental factors (e.g., heat, salinity or pollution stresses). MtrBTN2 global distribution
indicates its resilience across a broad range of temperature with varying seasonal patterns [14],
reflecting host species distribution range. Prior to evidence of horizontal transmission, pollutants
have been suggested as a potential cause of DN; however, the association with disease
prevalence remained inconclusive [25–27], and as we now know that some neoplasias are
transmissible, it is no longer the initial carcinogenesis but the transmission that needs to be
explained. Seasonal influence on DN prevalence was observed at a local scale (higher during

winter in *M. trossulus*, [28, 29], and *M. edulis*, [30]), whereas no significant influence of

seasonality was noted at the European scale [31]. Unfortunately, in the absence of genetic

diagnostics, we cannot know how many of these DNs were indeed BTNs (i.e. transmissible) or

not, and rigorous evaluation of MtrBTN2 prevalence seasonality is still lacking. In brief, little is

known about the ecology of MtrBTN2 in relation to its host and the possible environmental

factors influencing the epidemiology of this transmissible cancer.

In the present study, we investigated the occurrence of MtrBTN2 in a survey area in Southern

Brittany (France) which has contrasting natural and anthropogenic habitats and a mosaic

distribution of *M. edulis* and *M. galloprovincialis*. We sampled 40 natural populations as well as

9 mussel farms, 7 floating buoys, and 20 ports. To detect MtrBTN2-infected mussels, we used

digital PCR and real-time PCR, amplifying one nuclear and two mitochondrial genes with

published and newly designed primers targeting specific variants. We found a low prevalence of

MtrBTN2 (23/1516), equally affecting the two coexisting *Mytilus* species. Interestingly, most

affected sites (7/9) were ports, suggesting that ports could be favourable habitats for MtrBTN2

transmission and/or that maritime transport could play a role as a vector for the spread of the

disease. This highlights the possible role of ports as epidemiological hubs for MtrBTN2

propagation.

2.　Material and methods

2.1. Sampling

We collected 1516 *Mytilus* spp. mussels from 76 sites of the French Atlantic coast, from the Bay

of Quiberon to Pornic, between January and March 2020. Approximately 20 mussels were

sampled at each site (Table S1), allowing us to have a large number of closely located sites. We

collected hemolymph from adductor muscle (with a 1 ml syringe, 26G needle) and a piece of

mantle; both samples were fixed with 96% ethanol as described in [16]. DNA extraction was

performed with the NucleoMag 96 Tissue kit (Macherey-Nagel) using a Kingfisher Flex

extraction robot (serial number 711-920, ThermoFisher Scientific). We followed the kit protocol

with a modified lysis time of 1 hour at 56°C and modified the volumes of some reagents: beads

were diluted twice in water, 200 µl of MB3 and MB4, 300 µl of MB5 and 100 µl of MB6. DNA

concentration (ng/µl) was quantified using the Nanodrop 8000 spectrophotometer (ThermoFisher

Scientific). DNA from hemolymph was used for MtrBTN2 detection and DNA from mantle was

used for mussel genotyping (Figure S1).

2.2. MtrBTN2 detection

As the prevalence of MtrBTN2 is expected to be low [16, 32] and the number of samples is high

(n = 1516), we chose to pre-screen by pooling 12 samples (pooled screening step; Figure S1) and

then demultiplexing positive pools to specifically target cancerous samples (simplex screening

step; Figure S1). Prior to pooling, DNA concentrations were adjusted to 10 ng/µl to obtain an

unbiased representation of each sample in the pool. This allowed us to considerably reduce the

cost and time of detection, but we acknowledge that this protocol could result in some very early

stages of the disease being missed. To partly circumvent this problem, MtrBTN2 detection in

pools was performed by digital PCR (ddPCR) targeting one nuclear (Elongation Factor, EF) and

one mitochondrial (Control region, described in [9]) marker (Table S2). ddPCR is a sensitive

method, based on the Taqman method which requires two primers and one probe, and which

directly estimates the copy number of the targeted sequence. In addition, the use of a

mitochondrial marker should provide a more sensitive detection as the mitochondrial genome has

more copies than the nuclear genome. ddPCR primers design and analysis was subcontracted to

the company IAGE using the QIAcuity™ system (QIAGEN). Positive pools were then

demultiplexed (simplex screening step; Figure S1) and MtrBTN2 detection was performed by

real-time PCR targeting one nuclear (Elongation Factor, EF1α-i3, described in [11]) and one

mitochondrial (cytochrome c oxidase I, mtCOI-sub) marker (Table S2). To design the mtCOI-

sub primers, we used sequences of *M. edulis*, *M. galloprovincialis*, *M. trossulus*, and MtrBTN2

available from the National Center for Biotechnology Information (NCBI). Please note that

mtCOI-sub primers were initially designed to amplify a polymorphism that discriminates

between two MtrBTN2 sub-lineages "a" and "b" [16]. However, attempts to identify sub-

lineages based on melting temperature (Tm) did not yield conclusive results, most likely due to

additional polymorphism. Amplification of EF1α-i3 and mtCOI-sub markers was performed

using a three-step cycling protocol (95°C for 10 s, 58°C for 20 s, 72°C for 25 s) for 40 cycles.

We carried out real-time PCRs using the SensiFAST™ SYBR No-ROX Kit (Bioline) and the

LightCycler 480 Real-Time PCR System (Roche Diagnostics). We also confirmed the positive

samples diagnosed by real-time PCR using ddPCR (simplex screening step; Figure S1). The

specificity of both methods and markers was tested using various control samples of *M. edulis*,

M. galloprovincialis, *M. trossulus*, and MtrBTN2 samples of early, moderate, or late stages

(based on cytological observations, following [32], Tableau S3). To test the sensitivity, we used

positive controls with MtrBTN2 DNA diluted 10-fold (x10) and 100-fold (x100) in *M. edulis*

DNA.

Real-time PCR results were analysed using the LightCycler480 software. Two parameters were

extracted: threshold cycle (CT), obtained by Absolute Quantification analysis using the Second

Derivative Maximum method, and Tm, obtained by Tm calling analysis. Positive samples were

defined as those with a CT of less than 35 cycles and a Tm between 77.46 to 78.71 (based on Tm

values of the positive control samples; see result, Figure S2). ddPCR results were analysed with

the QuantaSoft™ Analysis Pro software (Bio-Rad), which provides the copy number for each

sample. Samples were considered positive when there were more than two positive droplets.

2.3. Mussel genotyping

We used 10 biallelic SNPs known to discriminate between *M. edulis* and *M. galloprovincialis*

mussel species (Table S4). These markers were identified as being ancestry-informative (fixed

for alternative alleles in *M. edulis* and *M. galloprovincialis*) in the [33] dataset and were

subsequently confirmed as near diagnostic by analysis of larger datasets [22, 34]. Genotyping

was performed using the Kompetitive Allele Specific PCR (KASP) method [35]. As we wanted

to know the mean ancestry of each sample (G ancestry) and reduce the cost of genotyping, we

developed a multiSNPs marker by multiplexing the 10 SNPs. Rapidly, 1µL of assay mix (KASP-

TF V4.0 2X Master Mix, 1X target concentration; primers, 1µM target concentration;

HyClone™ HyPure water) and 0.5 µL of DNA at 10 ng/µL were mixed in qPCR 384-well plates

using Labcyte Echo525 (BECKMAN COULTER). KASP analysis was performed on the

LightCycler 480 Instrument (Roche Diagnostics) with the following thermal cycling conditions:

initialisation 15 min at 95°C, first amplification 20 s at 94°C and 1 min at 61°C to 55°C with

steps of 0.6°C for 10 cycles, second amplification 20 s 94°C and 1 min at 55°C for 29 cycles,

read 1 min at 37°C and 1 s at 37°C.

KASP results are a combination of two fluorescence values, one for allele X, and another for

allele Y. Here, we oriented all SNPs so that allele X corresponds to *M. edulis* and allele Y to *M.*

galloprovincialis. Following [36], we transformed the data to obtain a single measure of the

relative fluorescence of the two alleles, using the following formula: $y_{i,j} = Y_{i,j}/(X_{i,j} + Y_{i,j})$;

where $X_{i,j}$ and $Y_{i,j}$ are the fluorescence values of allele X and allele Y, respectively for, i[th] SNP of

the j[th] sample. To scale $y_{i,j}$ value from 0 when *M. edulis* allele fluorescence dominates at the 10

SNPs to 1 when *M. galloprovincialis* allele fluorescence dominates, we used the following

formula: $y'_{i,j} = (y_{i,j} - \min(y_i))/(\max(y_i) - \min(y_i))$; where $y_{i,j}$ is the relative fluorescence

value for the i[th] SNP of the j[th] sample. In the results section and figures, $y'_{i,j}$ corresponds to "G

ancestry".

2.4. Environmental data

Source of environmental data

The 76 sampling sites were selected to cover four different habitats (ports, natural beds, farms

and floating buoys) presenting contrasted environments and genetic backgrounds (*M. edulis*, *M.*

galloprovincialis and hybrids). The description of environmental data considered in this study is

reported in Table S5 and all site information is available in Table S1. Note that natural beds are

intertidal sites, while the other habitats are subtidal. During sampling, we collected two habitat

discrete descriptive variables: population density (1: isolated, 2: patchy beds, 3: continuous beds)

and wave action (1: exposed, 2: sheltered). We used the E.U. Copernicus Marine Service

Information (doi: 10.48670/moi-00027) to access temperature (°C), salinity (PSU 1e-3) and

surface current velocity (m.s-1) data for each site (monthly recorded from 2021, resolution of

0.028°x0.028°). The velocity variable was formed by combining its Northward (N) and Eastward

(E) components, as follow: $\sqrt{E^2 + N^2}$. We also used EMODnet Human Activities (source:

EMSA Route Density Map) to retrieve the maritime traffic density of fishing, passenger, and

recreational boats in the study area (seasonal record from 2018, 2019 and 2020, resolution 1x1

km).

Preparation of environmental data

Among the 76 sampled sites, 13 fell outside the coverage scope of the Copernicus model (i.e

situated beyond the model's pixel boundaries). Consequently, for these sites (see Table S1), we

were unable to extract temperature, salinity, or velocity data. Certain sites recorded a velocity of

0 for all month (Table S1), raising concerns about the model resolution for this variable. To

validate the accuracy of Copernicus data, we computed the mean values of neighbouring pixels

(MNP, within a 0.05° buffer zone) and compared them to the pixel values at the respective sites

of interest (SP, referred to as the 'central' pixel, Table S6). We note that, due to the proximity of

our sampled sites to the coastline, the number of neighbouring pixels varied within a certain

range (Table S6). Notably, there was no correlation between these variations and the absolute

difference between SP and MNP values (Figure S3C). The comparison between SP and MNP

values was carried out for all sites, encompassing temperature, salinity, and surface current

velocity, and the results are presented in Figure S3. We observed a robust positive correlation for

temperature and salinity variables, providing confidence in the reliability of the extracted data for

our small-scale analysis. However, as expected, the inconclusive correlation for surface current

velocity led us to exclude this environmental variable from the analyses.

In the statistical analysis (see section 2.5) we used the annual average of salinity and boat

density (maritime traffic). For temperature, we used the annual variance as we identified strong

seasonal variations in temperature that would be misinterpreted if only the annual average

temperature were used (Figure S4).

2.5. Statistical analysis: relationship between MtrBTN2 prevalence, host genotype and

environmental variables

To ascertain whether cancer prevalence varied among the four habitats sampled (ports, natural

beds, buoys and farms), we performed Fisher's exact test on the contingency table (*fisher.test()*

function of the *stats* package *v4.1.3* [37]). Subsequently, we aimed to determine whether specific

environmental variables could provide a more comprehensive explanation of MtrBTN2

prevalence across these habitats. To do so, we pursued the following steps: an exploratory

analysis with a Principal Component Analysis (PCA), an examination of data dependencies

(Spearman correlation test and Moran's I test), and lastly, a regression model adapted to our

dataset (spatial logistic regression model). All statistical analyses were performed in R *v4.1.3*

[37].

Principal component analysis

As an exploratory analysis, we firstly performed a PCA to investigate the relationship between the quantitative environmental variables. Missing values were estimated using *imputePCA()* (*missMDA* package *v1.18* [38]), prior to performing the PCA using *PCA()* (*FactoMineR* package *v2.6* [39]).

Spearman correlation test

To test for correlation between environmental variables, we performed a non-parametric test of Spearman using *rcorr()* function from *Hmisc* package [40], that automatically excludes missing data.

Moran I test for spatial autocorrelation

Given the proximity of the sampled sites, which constitutes a fine spatial matrix, we used the Moran's I statistic to test for spatial autocorrelation among the environmental variables. After converting longitude and latitude data into UTM coordinates (*spTransform()* function of *rgdal* package v1.6-7 [41]), this analysis was performed using the moran.test() function (*spdep* package version 1.2-8 [42]) with the k-nearest neighbour classification.

Spatial logistic regression model

The association of MtrBTN2 prevalence and environmental variables was tested at the site level using a multivariate spatial logistic regression (*fitme()* function, *spaMM* package *v4.3.0* [43]). Spatial autocorrelation was accounted for by including a Matern covariance function as a random effect (i.e. *Matern(1 | Longitude +Latitude)*). MtrBTN2 prevalence was represented as a binary matrix (N, N_{site}-N) with N and N_{site} indicating site-specific MtrBTN2 case counts and total number of sampled mussels, respectively. The following explanatory variables were included into the full model: temperature variance, salinity mean, population abundance, wave action, average *M. galloprovincialis* ancestry of the mussel host population (G ancestry), maritime traffic (as described above, section 2.4). We also included a binary variable "port" into the model, where 1 denotes port habitat and 0 represents other habitats. Model suitability was evaluated using the residual diagnostics tool implemented in the "DHARMa" package [44]. To assess the significance of fixed effects (environmental variables), we performed Bartlett-

272 corrected likelihood ratio tests (LR tests) by comparing the full model with a model that
273 excluded the variable of interest (*fixedLRT()* function of *spaMM* package *v4.3.0* [45]). To
274 account for bias associated with rare events and collinearity of environmental variables (see
275 results), LR tests were performed using 999 bootstraps.
276
277 3. Results
278 3.1. Validation of the screening method
In order to detect MtrBTN2 in thousands of individuals, we developed dedicated ddPCR and
qPCR screening tools. The design of real-time PCR and ddPCR primers and probes allowed us to
amplify nuclear and mitochondrial markers specific to *M. trossulus* mussels and MtrBTN2
(Figure S5, TableS7). However, for the mtCOI-sub locus, the melting temperature (Tm) range of
MtrBTN2 was different from *M. trossulus* controls and allowed us to specifically detect
MtrBTN2 based on Tm values (Figure S2). With ddPCR, all MtrBTN2 control samples were
detected, regardless of marker or dilution, except for one MtrBTN2 sample diluted at x100 which
was negative for the EF locus (Figure S5). Copy numbers appeared higher for the mtCR marker,
showing a higher sensitivity for this mitochondrial marker. The same pattern was observed for
EF1α-i3 and mtCOI-sub in real-time PCR, with a higher sensitivity of the mitochondrial marker
based on the lower CT values observed (Figure S5). However, two MtrBTN2 samples diluted at
x100 were negative for mtCOI-sub and positive for EF1α-i3. Even though the mitochondrial
marker appears to be more sensitive in both methods, we chose to screen for MtrBTN2 using
both markers in order to have comparisons and additional validation.
293
294 3.2. MtrBTN2 screening reveals a low prevalence
We amplified nuclear and mitochondrial markers using ddPCR (EF, mtCR) and real-time PCR
(EF1α-i3, mtCOI-sub) to detect the presence of MtrBTN2 in the 1516 mussels sampled from the
76 sites of the study area (Figure S1). All results of the pooled and simplex screening are
presented in TableS8.
The pooled screening step using ddPCR revealed 25/127 positive pools: 9/127 with EF, 14/127
with mtCR, and 2/127 were positive for both markers (Figure S5). As the 14 mtCR-positive
pools showed low copy numbers (median = 7.59), we assume that EF did not amplify because of
a detection threshold. Although the 9 EF-positive pools were not expected, they were included

for further investigation by simplex screening (see below). For pools positive for both markers,

the EF copy numbers appear relatively low compared to mtCR copy numbers. This is due to the

high mitochondrial fluorescence masking the nuclear fluorescence and underestimating EF copy

numbers. To eliminate this bias in the simplex ddPCR step (Figure S1), we amplified the two

markers separately. In order to maximise cancer detection in the simplex screening step, we

chose to keep the pools positive for at least one gene, corresponding to 25 positive pools of 12

samples (n = 300).

We performed the simplex screening step using real-time PCR and then confirmed positive

samples with ddPCR (Figure S1). In total, 44/300 samples were positive for at least one real-time

PCR marker: 16 with EF1α-i3 only, 23 with mtCOI-sub only, and 5 with both markers (Figure

S5, Table S9). Pooled and simplex screening results are mostly consistent, as 20 of the 25

positive pools had at least one individual detected by real-time PCR amplification (Table S10).

The 5 pools without positive samples had low ddPCR copy numbers (less than 3.37 for EF and

less than 6.65 for mtCR). These false positives were to be expected as we intentionally applied

low threshold values in the pooled screening step. Two samples from mtCR-positive pools were

found to be positive for both markers at the simplex level, which was expected as the

mitochondrial marker is likely to be more sensitive than the nuclear marker. On the contrary, we

did not expect to observe samples positive for EF1α-i3/EF and negative for mtCOI-sub/mtCR.

However, all but one of the individuals from the EF-positive pools were positive only for the

EF1α-i3 marker (Table S10). The absence of mitochondrial genes in these samples does not

seem to be related to a detection threshold issue, especially as ddPCR and real-time PCR target

two different genes (mtCR and mtCOI-sub). We suspect that some host alleles that were present

at low frequencies in the 'dock mussels' populations (the hybrid lineage found in the port of

Saint-Nazaire [34]) may sometimes be amplified with EF1α-i3. This could be explained by

shared polymorphism between host species rather than the presence of MtrBTN2. Therefore, the

16 samples positive only for EF1α-i3 were conservatively considered as false positives (note

they were mostly sampled in the port of Saint-Nazaire, which does not affect our main result).

For the remaining 28 samples, we performed ddPCR with both markers to confirm the real-time

mitochondrial PCR diagnosis (Figure S1, TableS9). For 23/28 samples, mtCR ddPCR confirmed

the mtCOI-sub real-time PCR diagnosis. For 10/23 samples positive only for mtCOI-sub, ddPCR

revealed few EF copies. Finally, we considered all samples positive for mitochondrial markers

with both methods to be cancerous samples. We found 23/1516 positive samples from 9/76 sites

in which the number of cases varied from 1 to 8 (Figure 1). The highest proportion of cancerous

mussels was found at site P20 (Croisic Pontoon), and the next highest in the vicinity of this site

(P19, P21, and P23; Figure 1).

Figure 1: Location of sampling sites. Zoomed-in sites correspond to sites where mussels are

affected by MtrBTN2. Pie graphs and numbers in italics correspond to the proportion of

cancerous samples in each site. Coloured points represent the type of site, as indicated in the

figure's legend. Map origin: Ersi gray (light).

3.3. *M. edulis* **and** *M. galloprovincialis* **are equally affected by MtrBTN2**

To determine whether host genetic background or population composition influences cancer
prevalence and distribution, mussels were genotyped using 10 SNPs known to be diagnostic
between *M. edulis* and *M. galloprovincialis* (the two species coexisting in the studied area).
To reduce the cost of genotyping, we developed a multiplex assay, mixing 10 SNPs that directly
give the average ancestry of the sample (G ancestry, Figure S1). This approach was validated by
comparing the multiplex fluorescence and average simplex fluorescence of SNPs from a set of
samples (Figure S6a). We genotyped each individual from the 9 sites affected by MtrBTN2 (180
mussels) and estimated the average ancestry of each site by genotyping pools of individuals (76
pools). The positive correlation between the average individual fluorescence of a site and the
fluorescence of the corresponding pool validated our approach (Figure S6b). Mussels with G
ancestry values below 0.3 were assigned to *M. edulis,* those with values above 0.59 to *M.
galloprovincialis*, and those in between were assigned as hybrids (Figure 2, FigureS7). Of the
mussels affected by MtrBTN2, 15/23 were assigned to *M. edulis*, 7/23 to *M. galloprovincialis*
and 1/23 was a hybrid (Table S11). Considering the sites affected by MtrBTN2, MtrBTN2
prevalence did not appear to be significantly different between species with 15/105 (14%) in *M.
edulis*, 7/60 (11%) in *M. galloprovincialis* and 1/15 (6%) in hybrids (Fisher's exact test, p-value
= 0.83). Most *M. edulis* mussels affected by MtrBTN2 were found in populations with a majority
of *M. edulis* (P44, P23, P20)*,* only two were found in populations with some hybrids and *M.
galloprovincialis* mussels (P25, P71) and only one in a population with a majority of *M.
galloprovincialis* (P11, Figure 2, Figure S7). Interestingly, all *M. galloprovincialis* mussels
affected by MtrBTN2 were found in *M. galloprovincialis* populations (P19, P21, Figure 2,
FigureS7), which are close to the site with the highest prevalence (Croisic pontoon, Figure 1).
Finally, the hybrid individual affected by MtrBTN2 was found in a population where the two
species coexist (P22, Figure 2, FigureS7).
371

Figure 2: Individual and population ancestry. G ancestry of healthy (empty grey circle) and
MtrBTN2 affected (dark red) hosts is shown for the 9 sites with cancer (bold labels). Population
average ancestry for all sampled sites is represented by black crosses. Blue, white and orange
rectangles correspond to the G ancestry range of *M. edulis*, hybrids and *M. galloprovincialis*
respectively.

3.4. *MtrBTN2 infected mussels are mostly found in ports*
After the MtrBTN2 detection step, which revealed 23/1516 individuals infected with this
transmissible cancer, we investigated associations with habitat types. Cancerous samples were
found in 7 ports (16/23 infected mussels) and 2 natural beds (7/23) near the Croisic pontoon
(P20, Figure 1). MtrBTN2 was not detected in the 180 individuals from the 9 mussel farms or in
the 140 individuals from the 7 floating buoys. Mussels affected by MtrBTN2 were significantly
more frequent in ports than in other habitats (Fisher's exact test, p-value = 0.0001, Table S12). In
addition, the sampled ports are not significantly spatially autocorrelated (Moran I test, p-value =
0.25, Table S13).

3.5. *Exploring the effect of some environmental variables, none other than the port habitat*
**   *explains MtrBTN2 prevalence after accounting for spatial autocorrelation***

Given that MtrBTN2 was mostly found in ports, we aimed to gain a deeper understanding of whether environmental variables (including population ancestry) could explain this pattern. Univariate correlation between MtrBTN2 prevalence and all environmental variables across habitats are presented in Figure S8. The two best correlations are found with population density and maritime traffic, but we also have strong multicollinearity among variables that needs to be accounted for (see Table S14). The PCA reveals an association between environmental variables and habitat types rather than the presence of cancer itself (Figure 3). Indeed, anthropogenic habitats (ports, floating buoys and farms) with high population density and sheltered from wave action are well separated from natural beds on the second axis of the PCA. Interestingly, maritime traffic (i.e., annual mean of passenger, fishing, and recreational boat density) explains a direction of variation towards human-altered habitats, but also towards sites affected by MtrBTN2. MtrBTN2 seems to be found in sites with buffered temperature (low annual variance) and higher mean annual salinity, environmental variables that are associated with ports in our study. Water temperature indeed tends to be buffered in ports, being warmer in winter and cooler in summer (Figure S4). However, these variables are difficult to interpret as they also explain well the inertia of axis 1 of the PCA which differentiates the sites located near estuaries (La Loire or La Vilaine) from the others. To investigate more rigorously whether some environmental variables could explain prevalence of MtrBTN2, we used a spatial logistic regression model.

Figure 3: Principal Component Analysis on environmental variables. Axis 1 mainly captures the variance associated with annual temperature variance and annual salinity mean and Axis 2 mainly captures the variance associated with wave action, population density and maritime traffic.

As MtrBTN2 was mostly found in ports (see section 3.4), we included an environmental variable "port" (1 for port habitat, 0 for others) in the regression model analysis. To test for spatial structure, we conducted Moran's I tests on all environmental variables. The results indicate significant spatial autocorrelation in most environmental variables (Table S13), which was expected as we selected this area due to its mosaic of contrasted environments. However, "population density" and "port" variables were found randomly distributed in space (Table S13). To overcome spatial autocorrelation bias we performed a logistic regression model that corrects for spatial effect (*spaMM* package, [43]) and assessed environmental variables significance via bootstrapped Likelihood Ratio (LR) tests. Bootstrapped LRTs are expected to handle the problem of variables multicollinearity. MtrBTN2 prevalence is significantly explained by the

port habitat variable (LR test: $\chi^2 = 7.21$, df $= 1$, p-value $= 0.0072$; Table S15), and none of

the other environmental variables significantly explain MtrBTN2 prevalence after accounting for

spatial effects (Table S15). Despite the residuals following the distributional assumptions of the

model (Figure S9) and the significance of the 'Port' variable (also supported by the Fisher's

exact test, see 3.4), we acknowledge that quantifying the magnitude of this trend remains

challenging and requires further investigations.

4.   **Discussion**

Our results show that MtrBTN2 is present at low prevalence along the French Atlantic coast

(23/1516 = 1.5%). Strikingly, it is mostly found in ports and is equally distributed in *M. edulis*

and *M. galloprovincialis*. Our analysis shows that ports are likely epidemiological hubs for

MtrBTN2 propagation. However, the precise environmental factors influencing MtrBTN2

epidemiology have yet to be fully characterised. In this context, we discuss various

environmental variables and assess their relevance to the port environment and marine parasite

epidemiology.

Most MtrBTN2-infected mussels were found in ports and a prevalence hotspot was found around

the Croisic pontoon. Given the low prevalence of MtrBTN2 previously described [16, 32] and

confirmed in this study, we acknowledge that our sample sizes per site may be too small (n = 20)

to assess MtrBTN2 prevalence with confidence at a site level. However, this was not the scale of

resolution we were interested in. Instead, our overall sampling design, capturing replications of

habitat features, was chosen to evaluate the effect of habitats and genetic composition of host

populations. This allowed us to reveal similar MtrBTN2 prevalence in both species of the mosaic

hybrid zone, and a higher prevalence in ports.

The highest prevalence of MtrBTN2 in ports could be explained by two non-exclusive

hypotheses. Firstly, ports could offer a favourable environment as it is conducive to disease

development and propagation. Indeed, ports are often polluted, confined and permanently

immersed habitats (without tidal constraints) with high density of mussels. It is important to note

here that our objective is not to explain the emergence of a transmissible cancer, i.e. the initial

carcinogenesis in the founder host, but its maintenance and spread. The emergence probably took
place long before the industrial period and, despite studying a cancer, the mutagenic effect of
pollutants does not have to be put forward to explain our results, but rather their effect on host
resistance to infection by transmissible cancer cells. Secondly, there is increased connectivity
between ports due to maritime traffic and ports are connected to other sites (other ports or wild
sites) through biofouling from vessels.

Ports are often confined areas with longer particle residence times which could increase the
contact time between parasites and hosts [46, 47], thereby increasing the probability of
transmission and favouring the persistence of a passively dispersing parasite. However, the
possible hotspot on Croisic Pontoon, which is affected by strong tidal currents, provides
evidence against this hypothesis given that it is not a confined area.

Host population density is an important epidemiological parameter, as high density is known to
favour parasite persistence in host populations [48, 49]. Our results suggest that ports are prone
to harbouring high population density (Figure 3). However, we found no evidence of MtrBTN2
cases in mussel farms or floating buoys despite high mussel population density in these sites as
well. Although population density could favour MtrBTN2 persistence in an affected population,
this parameter alone cannot explain the higher prevalence of MtrBTN2 in ports.

Port-based shipping activities are major contributors to water pollution, including issues such as
oil spills, plastic contamination, wastewater discharge, and the use of antifouling paint [50, 51].
Pollution is a major environmental stressor that could affect host physiology and increase
susceptibility to parasites, particularly by altering its immune response (reviewed in [52]). As
filter-feeders, species of bivalve molluscs are particularly concerned [53–55]. However, in our
study, most of the affected individuals (15/23) were found in three open sea sites known to be
healthy and protected ecosystems with abundant resources (Croisic Pontoon, Natural beds 19 and
21 part of Natura 2000 area; Figure 1), suggesting that pollution alone cannot explain the higher
prevalence of MtrBTN2 cancers in ports. Conversely, it can be hypothesised that a healthy, less
stressful environment may increase the host permissiveness and allow MtrBTN2 (and possibly
other BTNs) to persist at a higher detectable prevalence. Host nutrition can affect disease

outcome by driving host immunity and parasite resource availability [56–58]. In invertebrates,

increased host nutrition can lead to increased parasite load [59] and improved immunity against

parasites [57, 58]. This balance could apply to mussel-MtrBTN2 interaction, and a healthy

environment with abundant resources could increase host carrying capacity and/or slow down

disease progression. A polluted environment could also affect the survival of MtrBTN2 cells in

seawater and reduce their chances of infecting a new host. However, the impact of pollution on

mussel hosts and cancer cells needs to be further investigated to clearly establish its influence on

the transmission and persistence of such transmissible cancers.

Connectivity through maritime traffic appears to be the main common feature between affected

ports, although this explanatory variable does not withstand spatial logistic regression analysis

any better than others. Vessels could facilitate MtrBTN2 propagation through transport of

affected mussels fixed on the hull. Maritime traffic has been identified as a vector of marine

mollusk pathogens (for review see [60]), microorganisms [61] and non-indigenous species [62,

63], through ballast water and/or hull biofouling. As such, ports are considered hotspots of

marine non-indigenous species introductions [62, 64, 65] and of marine parasites [66]. *Mytilus*

mussels are commonly reported as principal taxa composing hull biofouling (fixed adult mussels;

[67–69]). If a port is affected by a local outbreak, it could become a source of contamination to

other ports through biofouling of vessels, while natural beds will probably remain isolated by

density troughs that could halt the propagation (Figure 4). It is also conceivable that prevalence

hotspots exist at other unsampled locations along the coast, either due to environmental factors

that have yet to be characterised, or simply due to stochastic local outbreaks. These local

outbreaks can be the source of contamination to the anthropogenic connectivity network

providing the disease reaches a port. While this does not disqualify the factors discussed above –

e.g., pollution, population density and confinement- as drivers of contamination within ports, our

study suggests that maritime routes between ports could be anthropogenic gateways to MtrBTN2

propagation. Furthermore, despite a high density of maritime traffic around the floating buoys

sampled, these sites remained free of MtrBTN2. Long mooring times in ports increase the

probability of hull surface colonisation [69–71] as well as the risk of pathogen transmission [60].

This suggests that, in addition to cruising, ships may have to dock for the spread of MtrBTN2 to

be effective, either through transmission of cancer cells or detachment of infected mussels

(Figure 4). This could provide an explanation as to why maritime traffic is not a sufficient
explanatory variable for MtrBTN2 prevalence. Further investigations would be necessary to
ascertain the requisite propagule density (comprising both numerical and time-scale components)
for effective transmission, as well as to evaluate possible disruption of byssus quality in affected
hosts.

Figure 4: Diagram illustrating the discussion on the propagation of MtrBTN2 across
Mytilus **sp. populations.** Arrows' widths correspond to the probability of MtrBTN2 propagation.
Blue and orange mussels correspond to *M. edulis* and *M. galloprovincialis*, respectively.
Diagram created with BioRender.

A possible hotspot of MtrBTN2 prevalence was found in the area around the Croisic pontoon.
Even though this possible hotspot will need to be confirmed, this site could represent a source for
other ports and surrounding natural beds. Indeed, this pontoon is frequently used by maritime
shuttles, as well as by recreational and fishing boats, and is connected by maritime traffic to
many ports in the studied area. In particular, this pontoon is the departure site of boats travelling
to Hoedic (site P44 in Figure 1), an isolated island for which the presence of cancer is most
likely explained by anthropogenic transport given the distance from the mainland. MtrBTN2 is
also found in foreshores near Croisic. This reveals a possible spillover from ports to adjacent
wild populations (Figure 4). Maritime traffic has been identified as a vector for non-indigenous
species from commercial ports to natural populations via marinas [72, 73]. In our study, when
sites outside standard enclosed ports (e.g., enclosed marinas) were sampled, MtrBTN2 was never
detected, suggesting that contagion often remains confined to the port area. Spillover may
therefore depend on the degree of confinement of a port.

Mytilus galloprovincialis was thought to be more resistant to MtrBTN2 because of lower prevalence in this species [16], and population patches of *M. galloprovincialis* could also have acted as barriers to disease propagation between patches of *M. edulis*. However, our results suggest that *M. galloprovincialis* population patches close to *M. edulis* populations may also show enhanced prevalence. This does not contradict the idea that *M. galloprovincialis* population patches could impose a slightly higher resistance to MtrBTN2 propagation. Under the hypothesis that natural propagation is hindered by density troughs between population patches, or by population patches of a more resistant species, ports and ship traffic would create anthropogenic gateways that favour connections between patches of host populations (Figure 4). This hypothesis could also suggest that BTNs spread naturally and progressively from one host to another by close contact rather than drifting with the currents, making enhanced R0 by anthropogenic dispersal a threat to host populations.

Our results represent a significant advance in our understanding of MtrBTN2 epidemiology, both by proposing ports as epidemiological hubs and by suggesting that the natural propagation of this transmissible cancer could be naturally halted by density troughs in the distribution of host mussel populations. Detection of MtrBTN2 presence on boat biofouling would provide more direct evidence of its role as a vector. It would also be interesting to investigate whether prevalence hotspots are more commonly located in areas with particular environmental conditions, or whether they are stochastic local outbreaks. Sampling vessel biofouling in international ports linking known MtrBTN2 sites (East and North Europe, North West and South Pacific) and nearby marinas, along with genetic comparison of cancer cells, would provide a better understanding of the global and local transmission routes of this transmissible cancer since its emergence. While MtrBTN2 is not associated with extensive mortality events [30], other pathogens could be translocated and affect wild and farmed mussel populations. Various measures are employed to mitigate biofouling, including antifouling painting, dry docking, in-water cleaning with capture, and ballast water filtration (as discussed in [60]). While biofouling management for international vessel exchanges is established [74–76], regulations for biofouling control in local recreational boats remain to be improved [67, 77]. This is a significant concern, as the involvement of recreational boats in the spread of invasive species and possible associated

pathogens is increasingly documented [78–80]. Prioritising the survey of vessels that stay in port

for long periods [81] or identifying nodes of busy maritime route [72] could contribute

considerably to local and international biofouling management.

5.   Conclusion

Our study reveals that the prevalence of MtrBTN2 along the South Brittany French coast is low,

but that the most affected sites are ports (marinas, commercial ports, and docks). This result

suggests that ports are epidemiological hubs and that maritime routes are anthropogenic

gateways for the propagation of MtrBTN2. Investigations into the presence of other BTNs in

boat biofouling would provide more direct evidence of its role as vector, although mussels are

the most common species of bivalve mollusc in biofouling. The spread of marine

(micro)parasites and pathogens through vessel biofouling is a legitimate concern, and results

such as ours -i.e., on a smaller scale than translocations identified in the literature on marine

bioinvasion or by [9]- highlight the critical need for politicy regulation to limit the effects of

biofouling, both on ship hulls and port docks, in addition to ballast water control.

Data accessibility

All data used in this study are available in the supplementary material.

Author's contribution

M.H., Conceptualization, Data curation, Formal analysis, Investigation, Methodology,

Supervision, Visualization, Writing – original draft, Writing – review and editing. F.To.,

Investigation, Data curation, Writing – review and editing. E.A.V.B., Formal analysis,

Investigation, Writing – review and editing. L.P., Data curation, Investigation, Writing – review

and editing. F.C., Investigation, Writing – review and editing. E.G., Investigation, Writing –

review and editing. I.B., Resources, Writing – review and editing. H.C., Resources, Writing –

review and editing. A.S., Formal analysis, Writing – review and editing. F.Th.,

Conceptualization, Funding acquisition, Writing – review and editing. D.D-G.,

Conceptualization, Funding acquisition, Writing – review and editing. G.M.C.,

Conceptualization, Project administration, Funding acquisition, Writing – review and editing.

N.B., Conceptualization, Funding acquisition, Formal analysis, Investigation, Methodology,
Project administration, Supervision, Writing – original draft, Writing – review and editing.
608

609 **Conflict of interest declaration**
610 We declare we have no competing interests.
611

Fundings
This work was supported by Montpellier Université d'Excellence (BLUECANCER project) and
Agence Nationale de la Recherche (TRANSCAN project, ANR-18-CE35-0009 and DockEvol
project, ANR-23-CE02-0020-01). This study falls within the framework of the "Laboratoires
d'Excellence (LABEX)" CEMEB (10-LABX-0004) and Tulip (ANR-10-LABX-41). FT was
supported by the EVOSEXCAN project (ANR-23-CE13-0007), the MAVA Foundation and the
HOFFMANN Family.
619

Acknowledgement
We are grateful to the IAGE company for carrying out the ddPCR assay of this study. We thank
the GENSEQ and QPCR HAUT-DEBIT platforms for access to equipments and their expertise.
Credit is owed to Olivier Gimenez, François Rousset and an anonymous reviewer for their
valuable advice on statistical analysis. Finally, we thank Cécile Perrin for her valuable reading of
the manuscript.
626

627 **References**
628

[1]   Murgia C, Pritchard JK, Kim SY, et al. Clonal Origin and Evolution of a Transmissible
Cancer. *Cell* 2006; 126: 477–487.
[2]   Rebbeck CA, Thomas R, Breen M, et al. Origins and evolution of a transmissible cancer.
Evolution (N Y) 2009; 63: 2340–2349.
[3]   Pearse AM, Swift K. Allograft theory: Transmission of devil facial-tumour disease. *Nature*
2006; 439: 549.
[4]   Pye RJ, Pemberton D, Tovar C, et al. A second transmissible cancer in Tasmanian devils.
Proceedings of the National Academy of Sciences 2016; 113: 374–379.

[5] Garcia-Souto D, Bruzos AL, Diaz S, et al. Mitochondrial genome sequencing of marine
leukaemias reveals cancer contagion between clam species in the Seas of Southern
Europe. *Elife*; 11. Epub ahead of print 1 January 2022. DOI: 10.7554/ELIFE.66946.

[6] Metzger MJ, Reinisch C, Sherry J, et al. Horizontal transmission of clonal cancer cells
causes leukemia in soft-shell clams. *Cell* 2015; 161: 255–263.

[7] Metzger MJ, Villalba A, Carballal MJ, et al. Widespread transmission of independent
cancer lineages within multiple bivalve species. *Nature* 2016; 534: 705–709.

[8] Michnowska A, Hart SFM, Smolarz K, et al. Horizontal transmission of disseminated
neoplasia in the widespread clam Macoma balthica from the Southern Baltic Sea. *Mol
Ecol* 2022; 31: 3128–3136.

[9] Yonemitsu MA, Giersch RM, Polo-Prieto M, et al. A single clonal lineage of transmissible
cancer identified in two marine mussel species in South America and Europe. *Elife*; 8.
Epub ahead of print 1 November 2019. DOI: 10.7554/eLife.47788.

[10] Yonemitsu MA, Sevigny JK, Vandepas LE, et al. Multiple lineages of transmissible
neoplasia in the basket cockle (Clinocardium nuttallii) with repeated horizontal transfer of
mitochondrial DNA. *bioRxiv* 2023; 2023.10.11.561945.

[11] Burioli EAV, Hammel M, Bierne N, et al. Traits of a mussel transmissible cancer
are reminiscent of a parasitic life style. *Scientific Reports 2021 11:1* 2021; 11: 1–11.

[12] Giersch RM, Hart SFM, Reddy SG, et al. Survival and Detection of Bivalve Transmissible
Neoplasia from the Soft-Shell Clam Mya arenaria (MarBTN) in Seawater. Epub ahead of
print 2022. DOI: 10.3390/pathogens11030283.

[13] Skazina M, Odintsova N, Maiorova M, et al. Two lineages of bivalve transmissible
neoplasia affect the blue mussel Mytilus trossulus Gould in the subarctic Sea of Okhotsk.
Curr Zool 2022; 00: 1–12.

[14] Skazina M, Ponomartsev N, Maiorova M, et al. Genetic features of bivalve transmissible
neoplasia in blue mussels from the Kola Bay (Barents Sea) suggest a recent trans-Arctic
migration of the cancer lineages. *Mol Ecol*. Epub ahead of print 5 October 2023. DOI:
10.1111/MEC.17157.

[15] Skazina M, Odintsova N, Maiorova M, et al. First description of a widespread Mytilus
trossulus-derived bivalve transmissible cancer lineage in M. trossulus itself. *Sci Rep*; 11.
Epub ahead of print 2021. DOI: 10.1038/s41598-021-85098-5.

[16] Hammel M, Simon A, Arbiol C, et al. Prevalence and polymorphism of a mussel transmissible cancer in Europe. *Mol Ecol* 2022; 31: 736–751.

[17] Farley CA. Sarcomatoid proliferative disease in a wild population of blue mussels (mytilus edulis). *J Natl Cancer Inst* 1969; 43: 509–516.

[18] Baez-Ortega A, Gori K, Strakova A, et al. Somatic evolution and global expansion of an ancient transmissible cancer lineage. *Science (1979)*; 365. Epub ahead of print 2 August 2019. DOI: 10.1126/science.aau9923.

[19] Caza F, E B, FJ V, et al. Hemocytes released in seawater act as Trojan horses for spreading of bacterial infections in mussels. *Sci Rep* 2020; 10: 19696–19696.

[20] Skibinski DOF, Beardmore JA, Cross TF. Aspects of the population genetics of Mytilus (Mytilidae; Mollusca) in the British Isles. *Biological Journal of the Linnean Society* 1983; 19: 137–183.

[21] Coustau C, Renaud F, Delay B. Genetic characterization of the hybridization between Mytilus edulis and M. galloprovincialis on the Atlantic coast of France. *Mar Biol* 1991; 111: 87–93.

[22] Simon A, Fraïsse C, El Ayari T, et al. How do species barriers decay? Concordance and local introgression in mosaic hybrid zones of mussels. *J Evol Biol* 2021; 34: 208–223.

[23] Bierne N, Bonhomme F, David P. Habitat preference and the marine-speciation paradox. *Proc R Soc Lond B Biol Sci* 2003; 270: 1399–1406.

[24] Bierne N, David P, Langlade A, et al. Can habitat specialisation maintain a mosaic hybrid zone in marine bivalves? *Mar Ecol Prog Ser* 2002; 245: 157–170.

[25] Lowe DM, Moore MN. Cytology and Quantitative Cytochemistry of a Proliferative Atypical Hemocytic Condition in Mytilus edulis (Bivalvia, Mollusca). *J Natl Cancer Inst* 1978; 60: 1455–1459.

[26] Landsberg JH. Neoplasia and biotoxins in bivalves: Is there a connection? *J Shellfish Res* 1996; 15: 203–230.

[27] Green M, Alderman DJ. Neoplasia in Mytilus edulis L. from United Kingdom waters. *Aquaculture* 1983; 30: 1–10.

[28] Mix MC. Haemic neoplasms of bay mussels, Mytilus edulis L., from Oregon: occurrence, prevalence, seasonality and histopathological progression. *J Fish Dis* 1983; 6: 239–248.

698   [29]   Rasmussen LPD. Occurrence, prevalence and seasonality of neoplasia in the marine
699   mussel Mytilus edulis from three sites in Denmark. *Marine Biology: International Journal*
700   *on Life in Oceans and Coastal Waters* 1986; 92: 59–64.

701   [30]   Charles M. Etude des pathogènes, des conditions physiologiques et pathologiques
702   impliqués dans les mortalités anormales de moules (Mytilus sp.), https://tel.archives-
703   ouvertes.fr/tel-03053331 (2019, accessed 13 October 2021).

704   [31]   Bramwell G, Schultz AG, Sherman CDH, et al. A review of the potential effects of climate
705   change on disseminated neoplasia with an emphasis on efficient detection in marine
706   bivalve populations. *Science of The Total Environment* 2021; 775: 145134.

707   [32]   Burioli EAV, Trancart S, Simon A, et al. Implementation of various approaches to study
708   the prevalence, incidence and progression of disseminated neoplasia in mussel stocks. *J*
709   *Invertebr Pathol* 2019; 168: 107271.

710   [33]   Fraïsse C, Belkhir K, Welch JJ, et al. Local interspecies introgression is the main cause of
711   extreme levels of intraspecific differentiation in mussels. *Mol Ecol* 2016; 25: 269–286.

712   [34]   Simon A, Arbiol C, Nielsen EE, et al. Replicated anthropogenic hybridisations reveal
713   parallel patterns of admixture in marine mussels. *Evol Appl* 2020; 13: 575–599.

714   [35]   Semagn K, Babu R, Hearne S, et al. Single nucleotide polymorphism genotyping using
715   Kompetitive Allele Specific PCR (KASP): Overview of the technology and its application
716   in crop improvement. *Molecular Breeding* 2014; 33: 1–14.

717   [36]   Cuenca J, Aleza P, Navarro L, et al. Assignment of SNP allelic configuration in polyploids
718   using competitive allele-specific PCR: application to citrus triploid progeny. *Ann Bot*
719   2013; 111: 731–742.

720   [37]   R Core Team. R: A language and environment for statistical computing, http://www.r-
721   project.org/index.html (2020).

722   [38]   Josse J, Husson F. missMDA: A Package for Handling Missing Values in Multivariate
723   Data Analysis. *J Stat Softw* 2016; 70: 1–31.

724   [39]   Lê S, Josse J, Husson F. FactoMineR: An R Package for Multivariate Analysis. *J Stat*
725   *Softw* 2008; 25: 1–18.

726   [40]   Harrell JF. _Hmisc: Harrell Miscellaneous_. *R package version 51-0* .

727   [41]   Bivand R, Keitt T, Rowlingson B. rgdal: Bindings for the 'Geospatial' Data Abstraction
728   Library.

729   [42]   Bivand R. R Packages for Analyzing Spatial Data: A Comparative Case Study with Areal
730   Data. *Geogr Anal* 2022; 54: 488–518.

731   [43]   Rousset F, Ferdy J-B. Testing environmental and genetic effects in the presence of spatial
732   autocorrelation. *Ecography* 2014; 37: 781–790.

733   [44]   Hartig F. _DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed)
734   Regression Models_, https://CRAN.R-project.org/package=DHARMa (2022, accessed 19
735   December 2023).

736   [45]   Thomas F, Fisher D, Fort P, et al. Applying ecological and evolutionary theory to cancer: a
737   long and winding road. *Evol Appl* 2013; 6: 1–10.

738   [46]   Morrisey D, Gadd J, Page M, et al. *In-water cleaning of vessels: biosecurity and chemical*
739   *contamination risks. New Zealand Ministry for Primary Industries Technical Paper No*,
740   https://www.researchgate.net/publication/276920418 (November 2013, accessed 3
741   November 2023).

742   [47]   Gadd JB, Lewis JA. In-water cleaning of vessels: biosecurity and chemical contamination
743   risks, https://www.researchgate.net/publication/276920418 (2011, accessed 23 October
744   2023).

745   [48]   Bommarito C, Thieltges DW, Pansch C, et al. Effects of first intermediate host density,
746   host size and salinity on trematode infections in mussels of the south-western Baltic Sea.
747   *Parasitology* 2021; 148: 486–494.

748   [49]   Hopkins SR, Fleming-Davies AE, Belden LK, et al. Systematic review of modelling
749   assumptions and empirical evidence: Does parasite transmission increase nonlinearly with
750   host density? *Methods Ecol Evol* 2020; 11: 476–486.

751   [50]   Široka M, Piličić S, Milošević T, et al. A novel approach for assessing the ports'
752   environmental impacts in real time – The IoT based port environmental index. *Ecol Indic*
753   2021; 120: 106949.

754   [51]   Roberts T, Williams I, Preston J, et al. Ports in a Storm: Port-City Environmental
755   Challenges and Solutions. *Sustainability 2023, Vol 15, Page 9722* 2023; 15: 9722.

756   [52]   Destoumieux-Garzón D, Mavingui P, Boetsch G, et al. The one health concept: 10 years
757   old and a long road ahead. *Front Vet Sci* 2018; 5: 14.

[53] Canesi L, Ciacci C, Bergami E, et al. Evidence for immunomodulation and apoptotic
processes induced by cationic polystyrene nanoparticles in the hemocytes of the marine
bivalve Mytilus. *Mar Environ Res* 2015; 111: 34–40.

[54] Girón-Pérez MI. Relationships between innate immunity in bivalve molluscs and
environmental pollution. *ISJ* 2010; 7: 149–156.

[55] Renault T. Immunotoxicological effects of environmental contaminants on marine
bivalves. *Fish Shellfish Immunol* 2015; 46: 88–93.

[56] Chandra RK. Nutrition, immunity and infection: From basic knowledge of dietary
manipulation of immune responses to practical application of ameliorating suffering and
improving survival. *Proc Natl Acad Sci U S A* 1996; 93: 14304–14307.

[57] Tseng M, Myers JH. The Relationship between Parasite Fitness and Host Condition in an
Insect - Virus System. *PLoS One* 2014; 9: e106401.

[58] Pike VL, Lythgoe KA, King KC. On the diverse and opposing effects of nutrition on
pathogen virulence. *Proceedings of the Royal Society B*; 286. Epub ahead of print 10 July
2019. DOI: 10.1098/RSPB.2019.1220.

[59] Cressler CE, Nelson WA, Day T, et al. Disentangling the interaction among host resources,
the immune system and pathogens. *Ecol Lett* 2014; 17: 284–293.

[60] Georgiades E, Scianni C, Davidson I, et al. The Role of Vessel Biofouling in the
Translocation of Marine Pathogens: Management Considerations and Challenges .
Frontiers in Marine Sciences; 8. Epub ahead of print 2021. DOI:
10.3389/fmars.2021.660125.

[61] Ruiz GM, Rawlings TK, Dobbs FC, et al. Global spread of microorganisms by ships.
Nature 2000 408:6808 2000; 408: 49–50.

[62] Massé C, Viard F, Humbert S, et al. An Overview of Marine Non-Indigenous Species
Found in Three Contrasting Biogeographic Metropolitan French Regions: Insights on
Distribution, Origins and Pathways of Introduction. *Diversity (Basel)* 2023; 15: 161.

[63] Molnar JL, Gamboa RL, Revenga C, et al. Assessing the global threat of invasive species
to marine biodiversity. *Front Ecol Environ* 2008; 6: 485–492.

[64] Bishop JDD, Wood CA, Yunnie ALE, et al. Unheralded arrivals: non-native sessile
invertebrates in marinas on the English coast. *Aquat Invasions* 2015; 10: 249–264.

[65] Couton M, Lévêque L, Daguin-Thiébaut C, et al. Water eDNA metabarcoding is effective in detecting non-native species in marinas, but detection errors still hinder its use for passive monitoring. *Biofouling* 2022; 38: 367–383.

[66] Pagenkopp Lohan KM, Darling JA, Ruiz GM. International shipping as a potent vector for spreading marine parasites. *Divers Distrib* 2022; 28: 1922–1933.

[67] Clarke Murray C, Pakhomov EA, Therriault TW. Recreational boating: a large unregulated vector transporting marine invasive species. *Divers Distrib* 2011; 17: 1161–1172.

[68] Salta M, Chambers L, Wharton J, et al. Marine fouling organisms and their use in antifouling bioassays.

[69] Davidson IC, Brown CW, Sytsma MD, et al. The role of containerships as transfer mechanisms of marine biofouling species. *Biofouling* 2009; 25: 645–655.

[70] Bouda A, Bachari NI, Nacef L, et al. Risk Analysis of Invasive Species Introduction in the Port of Arzew, by Calculation of Biofouling Surface on Ships' Hulls. *Environmental Modeling and Assessment* 2018; 23: 185–192.

[71] Sylvester F, Kalaci O, Leung B, et al. Hull fouling as an invasion vector: can simple models explain a complex problem? *Journal of Applied Ecology* 2011; 48: 415–423.

[72] Floerl O, Inglis GJ, Dey K, et al. The importance of transport hubs in stepping-stone invasions. *Journal of Applied Ecology* 2009; 46: 37–45.

[73] Guzinski J, Ballenghien M, Daguin-Thiébaut C, et al. Population genomics of the introduced and cultivated Pacific kelp Undaria pinnatifida: Marinas—not farms—drive regional connectivity and establishment in natural rocky reefs. *Evol Appl* 2018; 11: 1582–1597.

[74] International Maritime Organization [IMO]. *International Convention for the Control and Management of Ships' Ballast Water and Sediments.* Lodon, 2004.

[75] International Maritime Organization [IMO]. *Guidelines for the Control and Management of Ships' Biofouling to Minimize the Transfer of Invasive Aquatic Species*. London, 2011.

[76] McClay T, Zabin C, Davidson I, et al. *Vessel biofouling prevention and management options report.* 2015.

[77] GEF-UNDP-IMO GloFouling Partnerships Project 2022. Biofouling Management for Recreational Boating: Recommendations to Prevent the Introduction and Spread of

Invasive Aquatic Species. *GEF-UNDP-IMO GloFouling Partnerships Project, 2022.*
Biofouling Management for Recreational Boating: Recommendations to Prevent the
Introduction and Spread of Invasive Aquatic Species.

[78] Briski E, Wiley CJ, Bailey SA. Role of domestic shipping in the introduction or secondary
spread of nonindigenous species: biological invasions within the Laurentian Great Lakes.
Journal of Applied Ecology 2012; 49: 1124–1130.

[79] Ulman A, Ferrario J, Forcada A, et al. Alien species spreading via biofouling on
recreational vessels in the Mediterranean Sea. *Journal of Applied Ecology* 2019; 56:
2620–2629.

[80] Ashton G V., Zabin CJ, Davidson IC, et al. Recreational boats routinely transfer organisms
and promote marine bioinvasions. *Biol Invasions* 2022; 24: 1083–1096.

[81] Lim CS, Leong YL, Tan KS. Managing the risk of non-indigenous marine species transfer
in Singapore using a study of vessel movement. *Mar Pollut Bull* 2017; 115: 332–344.

Supplementary material

Figure S1: Diagram of screening and genotyping experimental steps.

Figure S2: TM threshold determination for EF1α-i3 (left) and mtCOI-sub (right) real-time PCR markers. Grey rectangles correspond to TM threshold based on MtrBTN2 positive control samples (77.46<TM<78.71). Sample type colour and diagnostic based on TM values are indicated in the legend. *M. edulis* and *M. galloprovincialis* correspond to negative controls, *M. trossulus* are *M. trossulus* controls, and MtrBTN2 samples are positive controls (ND: cancer stage not defined, initial: early stage, moderate: moderate stage, heavy: late stage, see TableS3)

Figure S3: Evaluation of Copernicus model resolution for salinity, temperature and velocity data. Correlation between the value of the site pixel (SP) and the mean value of neighbouring pixels (MNP) for each variable. **(A)** Correlation including all months of year 2021 and **(B)** each month separately. No correlation observed between the absolute difference of SP and MNP values and the number of neighbouring pixels **(C)**.

Figure S4: Sampled sites temperature, recorded monthly in 2021. Points correspond to a temperature measurement, solid lines refer to MtrBTN2-free sites (grey) and MtrBTN2 affected sites (darkred).

Figure S5: Real-time PCR and digital PCR results. (A) and (C) represent ddPCR results of EF

and mtCR, and (B) and (D) represent real-time PCR results of EF1α-i3 and mtCOI-sub,

respectively. Dotted lines correspond to the negative thresholds, 0 for no DNA copy detected by

ddPCR and 40 for no amplification in real-time PCR. Solid lines in each graph refer to the same

sample diluted to different concentrations (pure, x10, x100). *M. edulis* and *M. galloprovincialis*

correspond to negative controls, *M. trossulus* are *M. trossulus* controls, and MtrBTN2 samples

are positive controls (ND: cancer stage not defined, Initial: early stage, moderate: moderate

stage, heavy: late stage, see TableS2). Screening results using both method and markers are

shown for Pool and Simplex. Sample type colours and diagnostic status are indicated in the

figure legend.

Figure S6: Genotyping method validation. (A) Positive correlation between multiSNPs G

ancestry and the mean of the 10 single SNPs from a subset of Individual and Pool samples. **(B)**

Positive correlation between population DNA pool G ancestry and the mean individual G

ancestry of individuals from the same population.

Figure S7: Population and individual ancestry composition across the sampled area. Zoomed-in sites correspond to sites with mussels affected by MtrBTN2. Barplots represent the estimated ancestry of individuals based on the 10SNPs multiplex tool. MtrBTN2 individuals are indicated by stars under each barplot. Coloured points represent the population average ancestry, as indicated in the figure's legend. Map origin: Ersi gray (light).

894 **Figure S8: Univariate correlations between the prevalence of MtrBTN2 and environmental**

895 **variables.** (A) MtrBTN2 prevalence among habitats. (B) Correlations for the six environmental

896 variables studied. Coloured points correspond to the prevalence of MtrBTN2 as indicated in the

897 legend.

Figure S9: Residual diagnostics of the logistic regression model accounting for spatial autocorrelation. Model results are presented in Table S15. Plot generated using *simulateResiduals()* function of DHARMa package in R.

Supplementary Tables (➔ <u>See excel TableS1-15</u>)

TableS1 : Description of sampled sites. Sources of environmental variables are provided in TableS5. Site: site name; N: number of sampled mussels; MtrBTN2 prevalence: number of MtrBTN2 affected mussels; Temperature: annual variance; Salinity: annual mean; Maritime traffic: annual mean in log; temp: temperature; sal: salinity; vel_E and vel_N: Eastward and Northward current velocity, respectively; [fish-pass-other][19-21]_[1-4] fish: fishing boat density, pass: passenger boat density, other: recreational boat density, 19 to 21 correspond to the year of record, and 1 to 4 correspond to spring, summer, autumn, winter, respectively.

Temperature and Salinity missing data (blank) correspond to sites outside of Copernicus model pixels (n = 13).

Table S2: Primer and probe sequences used in real-time PCR and digital PCR.

Table S3: Control samples used to validate real-time PCR and ddPCR screening tools. All samples are from our laboratory sample collection. *M. edulis* and *M. galloprovincialis* correspond to negative controls, *M. trossulus* are *M. trossulus* controls, and MtrBTN2 samples are positive controls. MtrBTN2 stages were defined by cytology of the hemolymph, according to the following classification: early (<15% cancer cells), moderate (15-75% cancer cells), late (>75% cancer cells), ND when cancer stage was not defined.

Table S4: Bi-allelic nuclear SNPs used for the multiSNPs genotyping.

TableS5: Source and description of environmental variables collected for each habitat (ports, natural beds, farms, buoys).

TableS6: Data for the evaluation of Copernicus model resolution. Sources of environmental variables are provided in TableS5. Site: site name; N_pixels: number of neighbouring pixels within a buffer zone of 0.05°; SP: value of the site pixel; MNP: mean value of the neighbouring pixels; SP_MNP: absolute difference between SP and MNP values; .s, .t and .v refer to salinity, temperature and velocity values, respectively. Velocity correspond to the combination of squared Northward and Eastward velocity values. Missing data (blank) correspond to sites outside of Copernicus model pixels (n = 13).

Table S7: Control sample results validate real-time PCR and ddPCR screening tools. EF_CopyNb: copy number for EF ddPCR marker; mtCR_CopyNb: copy number for mtCR ddPCR marker; CT: threshold cycle, TM: melting temperature for EF1α-i3 and mtCOI-sub real-time PCR markers. NP: not performed.

TableS8: Pooled and simplex screening results. Pool_CopyNb_EF: copy number for EF
ddPCR marker (pooled screening); Pool_CopyNb_mtCR: copy number for mtCR ddPCR marker
(pooled screening); CT: threshold cycle, TM: melting temperature for EF1α-i3 and mtCOI-sub
real-time PCR markers; Simplex_CopyNb_EF: copy number for EF ddPCR marker (simplex
screening); Simplex_CopyNb_mtCR: copy number for mtCR ddPCR marker (simplex
screening). NP: not performed.

TableS9: Simplex screening results for positive samples. CT: threshold cycle, TM: melting
temperature for EF1α-i3 and mtCOI-sub real-time PCR markers; PCR_result: marker name
positive in real-time PCR; EF_CopyNb: copy number for EF ddPCR marker; mtCR_CopyNb:
copy number for mtCR ddPCR marker; ddPCR_result: marker name positive in ddPCR;
MtrBTN2: final diagnostic status. NP: not performed.

Table S10: Correspondence between pool and simplex screening. Pool_CopyNb_EF : copy
number for EF ddPCR marker (pooled screening); Pool_CopyNb_mtCR : copy number for
mtCR ddPCR marker (pooled screening); Pool_ddPCR_result: marker name positive in ddPCR
(pool); Nb_ind_positive: number of individuals in the pool found positive for one or two marker
in simplex screening; Simplex_PCR_result: marker name of the positive individual in real-time
PCR; Simplex_ddPCR_result: marker name of the positive individual in ddPCR, negative if the
sample did not amplify. NP: not performed.

Table S11: MultiSNPs genotyping results. G ancestry: fluorescence value; Ancestry: ancestry
defined based on G ancestry values.

Table S12: Contingency table used for the Fisher exact test.

Table S13: Moran I test on environmental variables. Results were obtained using moran.test()
function in R.

Table S14: Correlation coefficient and Spearman correlation test results for each
environmental data pairs.

Table S15: Spatial logistic regression model testing the effect of environmental variables on MtrBTN2 prevalence. Model results were obtained using fitme() from spaMM package in R. Matern covariance function was included to account for spatial autocorrelation (i.e. Matern(1 | Longitude + Latitude). Effects of environmental variables were tested using Likelihood ratio (LR) tests using fixedLRT() function from the same package. Temperature: annual variance; Salinity: annual mean; Maritime traffic: annual mean.