Migratory herbivorous waterfowl track multiple resource waves during spring migration

Fei Xu[1,2*], Wei Wu[3], Jie Wei[2*], Qinchuan Xin[3], Ben Wielstra[4,5], Frank A. La Sorte[6], Zhijun Ma[7], Guangchun Lei[8], Jialin Lei[8], Wenzhao Wu[2], Yongchuan Yang[1], Peng Gong[2,9], Bing Xu[2], Yali Si[2,10*]

[1] Key Laboratory of the Three Gorges Reservoir Region's Eco-Environment, Ministry of Education, Chongqing University, Chongqing 400045, China;

[2] Ministry of Education Field Research Station for East Asian Migratory Birds, Department of Earth System Science, Tsinghua University, Beijing 100086, China;

[3] School of Geography and Planning, Sun Yat-sen University, Guangzhou 510275, China;

[4] Institute of Biology Leiden, Leiden University, Leiden, the Netherlands;

[5] Naturalis Biodiversity Center, Leiden, the Netherlands;

[6] Cornell Lab of Ornithology, Cornell University, Ithaca, NY, USA;

[7] Ministry of Education Key Laboratory for Biodiversity Science and Ecological Engineering, Coastal Ecosystems Research Station of the Yangtze River Estuary, and Shanghai Institute of Eco-Chongming (SIEC), Fudan University, Shanghai, 200433, China;

[8] School of Ecology and Nature Conservation, Beijing Forestry University, Beijing 100083, China;

[9] Department of Geography, Department of Earth Sciences, Institute for Climate and Carbon Neutrality, The University of Hong Kong, Hong Kong, China;

[10] Institute of Environmental Sciences, Leiden University, 2333 CC, Leiden, Netherlands

* Correspondence to: feixu@cqu.edu.cn; weij16@tsinghua.org.cn; y.si@cml.leidenuniv.nl

Abstract

East Asian herbivorous waterfowl intensively use farmland in spring, next to their natural habitat. Accordingly, they might have expanded their migration strategy from merely tracking the green wave of newly emerging vegetation to also incorporating the availability of post-harvest agricultural seeds (here dubbed the seed wave). However, if and how waterfowl use multiple food resources to time their seasonal migration is still unknown. We test this migration strategy using 167 spring migration tracks of five East Asian herbivorous waterfowl species and mixed-effect resource selection function models. We find all study species arrive at their core stopover sites in the Northeast China Plain after agricultural seeds become available, extend their stay after spring vegetation emerges, and arrive at breeding sites around the emergence of vegetation. Moreover, the cue used in exploiting the availability of both food resources varies in different regions and among species. At the core stopover sites, larger-bodied species use the seed wave and smaller-bodied ones adopt the green wave. At the breeding sites, the former follow the green wave, and the latter track the snow-free time. Our findings suggest that waterfowl track multiple resource waves to finetune their migration, highlighting new opportunities for conservation.

Keywords

Resource tracking, seed wave, onset of vegetation green-up; agricultural land; migration phenology; seasonal bird migration

Introduction

Migratory species move between their breeding and non-breeding grounds and benefit from

the exploitation of seasonal food resources across the annual cycle [1]. Foraging

opportunities occur in a series of pulses, at different times and places, which generate

"resource waves" across the landscape [1-5]. The green wave hypothesis posits that the

migration progress of herbivores is driven by the seasonal growth of new vegetation at

stopover sites along the migration route [6, 7]. Studies have verified this hypothesis for a

variety of herbivorous animal groups, such as geese [4, 6, 8-10] and ungulates [11-13]. Not

only individual-level information compiled using tracking technology [6] but also population-

level information compiled by large-scale community science initiatives [14] support such

hypothesis. During vernal migration, birds mainly arrive at their stopover sites after the

emergence of spring vegetation, but can reach their breeding grounds earlier, thus overtaking

the green wave, and allowing their offspring to benefit from the peaks of food resources [4,

9]. While herbivorous species have been documented to track the green wave in Northern

Europe [6, 10, 15] and North America [14], this has not been the case for many herbivorous

waterfowl populations worldwide, especially within East Asia [15].

Due to extensive land cover change worldwide, herbivorous waterfowl have expanded

from using natural ecosystems to exploiting food resources in agricultural landscapes [16,

17]. For example, during the spring-staging period, herbivorous waterfowl have been

observed to consume new sown cereals in Europe [18], post-harvest corn remaining in

cornfields, and spilled grain in rice paddies in North America [19-21] and in Asia [22-27].

This way they benefit from the post-harvest management of agricultural lands [28, 29]. This

is particularly the case for East Asian waterfowl populations, where in China large areas of

natural wetland have been converted into farmland since the 1950s [30-33]. As a

consequence, East Asian waterfowl might take the availability of both food resources into

account during spring migration. However, this possibility of tracking multiple resource

waves during migration has not yet been investigated.

Migratory herbivorous waterfowl use the natural habitat (e.g., wetlands, grasslands,

shrublands) and farmland habitat differently. Since many waterbodies and grasslands in East

Asia are semi-natural, we use the term 'natural habitat' hereafter to represent areas where

birds feed on spring sprouts of vegetation (representing the green wave). Within farmlands,

East Asian waterfowl primarily consume seeds, rather than crop vegetation, e.g., spilled

grains and leftover maize that remained after the harvest [22] in the previous fall and so have

been carried over into spring. These seeds have been found to offer a higher food quality

compared to natural vegetation [16]. Spilled seeds are only available after the harvest and

during the period that the ground is not frozen, resulting in two waves in autumn and in

spring, that we here dub the seed wave (Figure S1). In winter at the core stopover area

Northeast China Plain, the daily minimum temperature can drop below -25 °C [34, 35]. It

often starts snowing from November and ends around the end of March [34, 36]. This makes

the post-harvest seeds not accessible in winter but these seeds can be used by waterfowl after

the ground thaws around the end of March [36]. The seed wave then ends at the onset of

sowing in early May [36], when the new seeds that are sowed covered with pesticides (Figure

S2*C*). Thus, the seed wave in spring in the core stopover area starts around the end of March

(with the end of snowmelt as a surrogate) and ends in early May with the start of sowing

(Figures S1 and S2) [37]. The timing of seed availability starts before, and partially overlaps

with, the emergence of vegetation [15]. How would the herbivorous waterfowl time their

migration based on these two resource waves is still unclear.

We aim to investigate if and how herbivorous waterfowl track multiple resource waves

during spring migration. We analyze 167 spring migration tracks from 2015 to 2021, derived

from satellite tracking of 99 individuals for five herbivorous waterfowl species (swan goose

Anser cygnoides, tundra bean goose *A. serrirostris*, greater white-fronted goose *A. albifrons*,

lesser white-fronted goose *A. erythropus*, and tundra swan *Cygnus columbianus*) in East Asia,

using a mixed-effect resource selection function model. We hypothesize that migratory

herbivorous waterfowl select resources upon arrival at stopover and breeding sites based on

the timing of food resource availability in both natural habitat and farmland (Figure 1). We

predict that: 1) All study species time their migration to use both resources (seeds and natural

vegetation) at their core stopover sites. They use farmland more frequently upon arrival at the

core stopover sites, and shift to natural habitat after the vegetation emerges. All study species

arrive at their natural habitat in the breeding sites around the time of vegetation emergence; 2)

Waterfowl tend to track the start of seed availability (i.e., the seed wave) at their core

stopover sites and then follow the newly emerged plants (i.e., the green wave) as a cue to

arrive at their breeding sites (Figure 1). Understanding how herbivorous waterfowl use the

spatiotemporal availability of multiple food resources during migration is fundamental to

predict species' responses to land cover and climate change. The wider implications of our

findings are new opportunities for effective management of multiple types of habitats to

conserve migratory waterfowl.

Figure 1. Schematic diagram of tracking multiple resource waves by East Asian

migratory herbivorous waterfowl. (A) Core stopover sites (S1) and breeding sites (S2)

along the spring migration route. **(B)** During spring migration, birds arrive on farmlands at

the core stopover sites after the start of seed availability (seed wave; orange line), extend their

stay after spring vegetation in natural habitats emerges (green wave; green lines) (S1) and

reach the breeding grounds after the emergence of vegetation (green wave; green lines) (S2).

(C) Photo of greater white-fronted geese (*Anser albifrons*) and tundra bean geese (*A.*

serrirostris) foraging in farmlands (taken 19 March 2021) on Yuquan island, Hunchun,

Yanbian Korean Nationality Autonomous Prefecture, Jilin, China (photo credit: Haixiang

Zhou). Inset photo shows leftover maize (taken 2 May 2016) on farmlands in Fuyu Village,

Nehe, Qiqihar, China (photo credit: Jie Wei). **(D)** Photo of new spring vegetation (taken 1

May 2016) embedded in old vegetation in Baishan town, Longjiang, Qiqihar, China (photo

credit: Jie Wei).

Materials and Methods

Analysing bird migration patterns

We analyzed the satellite tracking data (i.e., 167 spring migration tracks) of five East Asian

herbivorous waterfowl species from 2015 to 2021, covering 22 tracks of swan geese from 19

individuals, 8 of tundra swans from 4 individuals, 54 of tundra bean geese from 28

individuals, 63 of greater white-fronted geese from 34 individuals, and 20 of lesser white-

fronted geese from 14 individuals. Lesser white-fronted geese (*Anser erythropus*), greater

white-fronted geese (*A. albifrons*), tundra bean geese (*A. serrirostris*), and tundra swans

(*Cygnus columbianus*) were captured and tagged at their East Dongting Lake (29°N, 113°E)

and Poyang Lake (29°N, 116°E) wintering sites in the Yangtze River Floodplain, China

during the 2014 to 2018 wintering seasons. Swan geese (*A. cygnoides*) were captured at the

breeding site Hulun Lake, Hulun Buir, China (48.3°N, 117.4°E) during the 2017 molting

season. All birds were equipped with GPS - GSM (Global Positioning System – Global

System for Mobile Communications), solar-powered loggers. GPS locations were recorded

every 2 h. Information used in the analysis on migration patterns include Bird ID, track year,

longitude, latitude, and time of the GPS location. Detailed methods on birds capture and

deployment of the satellite transmitters can be found in Si, Xu [22] and Lei, Jia [38].

We quantified the migration parameters (Table S1) in three steps. First, we subset all

GPS locations between January 1st to July 31st, thereby including months of the whole

spring migration period (February-June) and the months in which all five herbivorous

waterfowl have arrived at breeding sites (June-July). The definition of the spring migration

period is that GPS locations recorded from the day that birds left their wintering sites to the

day that they reached their breeding sites [8]. Second, we labeled daytime and nighttime

locations, by identifying the sunrise and sunset time for each location based on algorithms

provided by the National Oceanic & Atmospheric Administration (NOAA)

(https://www.esrl.noaa.gov/) [39]. Third, we identified stopover sites and the length of stay in

each site through the space-time permutation model (Supplementary Text) in the SaTScan

statistics (http://www.satscan.org) [22, 40]. We defined the stopover sites as those sites where

birds stayed for at least two days [41] and moved within a radius of less than 50 km [22, 41-

43]. The arrival/departure time of the wintering/stopover/breeding sites for each individual in

each track year were defined as the first day it arrived/left a specific site in which it showed a

continuous presence/absence [23]. Cumulative length of stay in the Northeast China Plain

and Russian stopover sites was defined as the sum of all days spent at stopover sites by an

individual in its respective region. Migration distance was defined as the cumulative distance

travelled between the wintering site and breeding site [22].

We calculated mean dates of migration timing upon arrival at or departure from

wintering sites, core stopover sites, and breeding sites for all tracks of each species, i.e., 1)

mean departure date from the wintering grounds, 2) mean arrival date at core stopover sites,

3) mean departure date from core stopover sites, and 4) mean arrival date at breeding sites.

For each of the four types of migration timing, we compared the dates among five

herbivorous waterfowl species in relation to their body size, which was obtained from

EltonTraits 1.0 [44]. Differences on the migration timing across species were assessed using

one-way ANOVA followed by Tukey's honestly significant difference (HSD) multiple

comparison.

Investigating resource utilization in migration

To examine the habitat use of tracked birds along the spring migration route, we used two

land cover products. The 30 m spatial resolution spring land cover product [45] from 2015

was selected to capture the natural habitats and farmlands within China. The European Space

Agency Climate Change Initiative (ESA-CCI) 300 m spatial resolution land cover product

from 2015 (http://maps.elie.ucl.ac.be/CCI/viewer/index.php) was used to capture the

relatively homogeneous landscape within Russia. Two classification systems from the two

products were aggregated into seven land cover types: grassland, farmland, water, wetland,

forest, shrubland, and 'other' (i.e., barren land, lichens and mosses, snow and ice, or

impervious). Habitat use at each stopover site was estimated using the percentage of bird

GPS locations on each land cover type [22]. Habitat use at each latitude was estimated using

the percentage of bird GPS locations on each land cover type across all stopover sites (their

rounded centroid coordinates) located in the same latitude. The proportion of land cover types

in use at each latitude was then calculated as the percentage of bird GPS locations on each

land cover type divided by the sum of the percentage of bird GPS locations across all land

cover types at the corresponding latitude.

As waterfowl mostly forage during the day, we also calculated the habitat use at daytime

(i.e., the proportion of land cover in use during the day) across the whole spring migration

period, to better interpret how waterfowl use the food resources across time. To better

investigate the multiple resource use across time within the core stopover sites in the

Northeast China Plain, we divide the length of stay (Table S1 and Figure S3) of herbivorous

waterfowl into two periods, i.e., 1) from the arrival date at each stopover site in the Northeast

China Plain to the local emergence of vegetation, and 2) from the local emergence of

vegetation till they departure from this region. We then calculate the foraging distribution

(i.e., the proportion of GPS locations during the day on farmlands and natural habitats) for all

study species during these two different time windows (Figure 3 and Figure S7).

Calculating the timing of the seed wave and the green wave

For each track, the spring vegetation emergence time in natural habitats and the start of seed

availability time on farmlands were calculated. We used the second-order derivative [46, 47]

of the smoothed time series of Enhanced Vegetation Index (EVI) derived from the Moderate

Resolution Imaging Spectroradiometer (MODIS) Terra 8-day 250 m spatial resolution surface

reflectance product MOD09Q1 to extract the spring vegetation emergence time (details are

provided in Supplementary Text). Only pure pixels [48] where the proportion of herbaceous

area (i.e., wetlands, grasslands, and shrublands) was larger than 0.5 within a 250 m grid cell

were used (details are provided in Supplementary Text and Figure S9).

The start of the seed availability time was determined by the end of snowmelt in spring

because after the snow melt seeds (remaining from previous year's harvest) could be easily

dug out from the ground. Following the method of Rickbeil, Coops [49] we used the MODIS

Terra Snow Cover daily 500 m spatial resolution product MOD10A1 to measure the different

degree of snowmelt (i.e., the onset and the end of snowmelt). We first created a time-series to

capture the daily percent of snow cover (from day 1 to day 210) for each stopover and

breeding site, for each track. We then applied three-piece segmented regression to extract the

onset and the end of snowmelt for each pixel (Figure S10). To facilitate interpreting the

resource use, in addition to the vegetation availability date and the seed availability date, we

also calculated the onset of snowmelt and the end of snowmelt at each site, for each

individual bird, in each track year.

Determining the resource availability – bird arrival relationship

We used a resource selection function (RSF) framework [50, 51] to investigate to what extent

the start of resource availability can be matched with the arrival date at the stopover or

breeding sites for migratory herbivorous waterfowl during spring migration. Using a logistic

regression of presence/absence points, RSFs can be used to define the selection of resources

on the day when waterfowl arrive at a specific site [23]. The presence points were defined as

all GPS locations recorded on the day of arrival. The absence points were drawn randomly on

the date before and after arrival at a specific site during the spring migration period. For each

pixel in the specific site, we calculated the number of days between each of the four indices

(i.e., the spring vegetation emergence, the start of seed availability, the onset of snowmelt,

and the end of snowmelt) and the bird arrival date. Migratory waterfowl might not track the

exact day that a specific resource becomes available but may display a lag effect, where birds

arrive before (positive lag value) or after (negative lag value) the resource becomes available

[15]. In order to obtain the focal lag (i.e., the number of days before or after the resource

available or snow melt dates), we tested the predictive ability of the models across a series of

temporal lags, by adding or subtracting days to the resource becoming available or the snow

melt dates, following Laforge, Bonar [51]. For example, if a bird arrives at the stopover site

two days after seed availability, the model with a lag of two (by subtracting two from the day

to the seed availability date) would have the highest likelihood.

We executed RSFs in two regions for each species: the core stopover area Northeast

China Plain and the breeding sites. We ran two models in the Northeast China Plain stopover

sites (35°-55°N latitude) using vegetation and seed availability as the predictors, and three

both at the mid-latitude (around 46°- 49°N latitude) and arctic breeding sites (around 70°-

75°N latitude) using vegetation availability, the onset of snowmelt, and the end of snowmelt

as predictors. The response variable was the probability of bird arrival. The presence was set

to 1 and absence was set to 0. For each of the five waterfowl species, within each region, we

fit a mixed-effects conditional logistic regression model using a binomial distribution as the

link function. The fixed effect for each model was a specific type of temporal lag, i.e., lag

days to 1) the spring vegetation emergence, 2) the start of seed availability, 3) the onset of

snowmelt, or 4) the end of snowmelt. For each individual tracked in each year, we obtained

the identity of the track (track ID), named as individual (bird ID) and year (ID-year). The

track ID was included in the model as a random effect. We used the R package glmmTMB

[52] to run the analyses.

We modelled the lags in days to food availability or the snow melt conditions that fell

between the 10th and 90th percentile of all locations within each region used by herbivorous

waterfowl. We selected the top model for the focal lag based on the lowest Akaike

Information Criterion (AIC) [51]. The range around the optimal lag was also estimated by

predicting the day at which interpolated log-likelihood values have a difference of 1 (ΔAIC =

2). The difference in range was then calculated by the number of days between the minimum

and maximum estimate. The predictive performance of the top models with the optimal

estimated lag was evaluated by k-fold cross-validation [51, 53], in according to previous

studies [54, 55]. This k-fold cross-validation procedure withheld a fraction of the data using a

k-fold partitioning [54], where k represented the number of partitions (i.e., test-training sets),

ranging from two to N-1 (number of observations minus one) [54, 55]. We split our origin

samples by the identity of the track (track ID) into k (k = 3) training-testing sets and 10 bins.

We trained our model iteratively on two out of the three datasets using a logistic regression.

The remaining dataset was used for validation. To examine model performance, the pattern of

predicted RSF scores for partitioned test-training sets is investigated against categories of

RSF scores (bins) [55]. We calculated Spearman rank correlations (r_s) between 10 RSF bin

ranks and three test-training sets (k = 3). A model with good predictive performance would be

expected to be one with a strong positive correlation (i.e., a high r_s value) [55]. For each

region and each species, we therefore considered the predictor used in the model with the

highest r_s value as the cue used by herbivorous waterfowl in timing their migration [56].

The calculation of the green wave was performed in MATLAB R2020a [57]. Calculating

the timing of seed wave, investigating resource utilization, and analysing the relationship

between resource availability and bird arrival were performed in R 4.1.0 [58].

Results

Resource utilization along the migration routes

We analyze the relationship between spatiotemporal bird migration patterns (Table S1 and Figure S3) and the availability of two food resources (seeds and newly emerged vegetation). Spring migration tracks (n = 167) from 2015 to 2021 of five waterfowl species within the East Asian – Australasian Flyway were used: 22 tracks of swan geese, 8 of tundra swans, 54 of tundra bean geese, 63 of greater white-fronted geese, and 20 of lesser white-fronted geese. Swan geese migrate around 2500 km from their wintering grounds in Poyang, China to reach their mid-latitude breeding sites in Inner Mongolia and Mongolia (Table S1 and Figure S3*C*). The other four species are arctic breeders, with a migration distance over 5000 km (Table S1 and Figure S3*A, B, D, E*). Seed resources at farmlands are all distributed below 52°N latitude (Figure S4). Tracked birds utilize both natural habitats (water and surrounding wetlands, grassland, and shrublands) and farmlands (especially in the core stopover sites in the Northeast China Plain). In Russia (above 52°N latitude), waterfowl mainly use water, shrublands, grasslands, and wetlands. They briefly stay at wetlands or rivers that occur within forests, which explains the high proportion of forest habitats used in this region (Figure S4*F-H*). At breeding sites, tracked birds use grasslands, wetlands, and shrublands as their main habitats (Figure S4).

Herbivorous waterfowl track multiple resource waves during spring migration

Herbivorous waterfowl in East Asia adopt a strategy in which they use both the seed wave

and the green wave during spring migration (Figure 2). All study species arrive after seeds

become available (using the end of snowmelt as a surrogate) and depart after the new

vegetation become available in their core stopover sites in the Northeast China Plain (Figure

2 and Figure S5). The period when seeds become available in this region is at the end of

March (86 ± 31 Julian days, Figure 2), whereas the emergence of new vegetation occurs

around early May (114 ± 30 Julian days, Figure 2). Birds arrive at the core stopover sites in

early April (94 ± 14 Julian days, Figure 2). All arctic-breeding species depart from the

Northeast China Plain region around mid-May (131 ± 10 Julian days, Figure 2*E, H, K, N*). All

study species arrive at their breeding sites around the onset of vegetation growth (Figure 2).

In the Russian stopover sites where they briefly pause, the observed arrival timings primarily

match the end of snowmelt (around the end of May and the beginning of June). Birds reach

their arctic breeding sites around early June (Table S1, Figure 2*F, I, L, O*, and Figure S5).

Figure 2. Tracking multiple resource waves during spring migration. All study species

(D-R) arrive (vertical red line with error bar) at the Northeast China Plain after the start of

seed availability (vertical orange line with error bar) and departed (vertical black line with

error bar) from this region after the emergence of vegetation (vertical green line with error

bar), thereby use both food resources. Greater white-fronted geese and lesser white-fronted

geese depart from the wintering grounds **(J, M)** and arrive at the Northeast China Plain **(K,**

N) later than larger-bodied species **(D, E, G, H, P, Q)** to better use the newly emerged

vegetation (vertical green line with error bar). All study species arrive at the breeding sites

around the time vegetation emerges (**F, I, L, O, R**). Grey lines show annual food availability

indicated by enhanced vegetation index (EVI) time series. Dots show the centroid locations

of wintering grounds (green dots in **A**), core stopover sites in the Northeast China Plain

(orange dots in **B**), breeding sites in the mid-latitude (dark blue dots in **C**) and arctic (light

blue dots in **C**) used by five herbivorous waterfowl.

Herbivorous waterfowl indeed spend relatively more time on farmland when they arrive

at their core stopover sites, and then shift to spend more time on natural habitat after newly

emerged vegetation becomes available (Figure 3, Figures S6 and S7). For all study species,

the proportion of habitat use on farmlands at daytime is 42% before the emergence of

vegetation, whereas this decreases to 32% after the emergence of vegetation, resulting in a

10% shift in foraging distribution from farmlands towards natural habitats (i.e., grasslands,

shrublands, wetlands, and water areas) (Figure 3). This habitat shift pattern is more obviously

in the two larger-bodied species, with a 28% shift for tundra swans and 16% for swan geese

(Figure 3).

Figure 3. Foraging distribution before and after the emergence of vegetation. The

proportion of habitat use showing foraging distribution at core stopover sites in the Northeast

China Plain was calculated using the proportion of GPS locations during the daytime on each

land cover type. The proportion of GPS locations during the daytime on farmlands decrease,

shifting on average 10% towards natural habitats after the emergence of vegetation for all

study species. Larger-bodied herbivorous waterfowl (i.e., tundra swan *Cygnus columbianus*

and swan geese *Anser cygnoides*) have relatively more shifts from farmlands to natural

habitats after the emergence of vegetation than relatively smaller-bodied species (i.e., tundra

bean geese *A. serrirostris*, greater white-fronted geese *A. albifrons*, and lesser white-fronted

geese *A. erythropus*).

Variation in tracking multiple resource waves among species

Although all study species track multiple resource waves during spring migration, the specific spring migration timing varies slightly among species (Figure 2 and Figure S8). Species with a relatively larger body size (tundra swans and tundra bean geese) show a relatively earlier departure from the wintering sites ($F = 9.98$, $df = 4$, $P < 0.001$, Figure 2*D, G,* Figure S8*A,* and Table S1), and a relatively earlier arrival on the Northeast China Plain ($F = 20.08$, $df = 4$, $P < 0.001$, Figure 2*E, H,* Figure S8*C* and Table S1), in comparison to smaller-bodied species (greater white-fronted geese and lesser white-fronted geese; Figure 2*J, K, M, N*). The two smaller-bodied geese stay in the wintering grounds for an extended period of time after the emergence of vegetation: 10 days in greater white-fronted geese and 28 days in lesser white-fronted geese (Figure 2*J, M*). The departure time from the core stopover sites is significantly earlier for mid-latitude breeding swan geese than the other four arctic-breeding species ($F = 10.45$, $df = 4$, $P < 0.001$, Figure 2*,* Figure S8*D* and Table S1). The smallest-bodied lesser white-fronted geese arrive at the arctic breeding sites the latest among all five herbivorous waterfowl species ($F = 40.67$, $df = 4$, $P < 0.001$, Figure 2, Figure S8*B* and Table S1).

The arrival probability at the core stopover sites and breeding sites in relation to the availability of food waves and snow conditions shows that optimal lags to the availability of multiple resources also vary slightly among species (Figure 4 and Table 1). For arctic-breeding species, the highest arrival probability occurs after the start of seed availability and before the emergence of vegetation in the core stopover sites (Figure 4*A, C, E, G*). Tundra

swans arrive at the core stopover sites soon (only one day) after seeds become accessible

(Table 1 and Figure 4*A*), whereas smaller-bodied goose species arrive later, after seeds have

already been available for about one week (for tundra bean geese; Figure 4*C*) to up to three

weeks (for greater white-fronted geese and lesser white-fronted geese; Figure 4*E and G*). On

the arctic breeding grounds, three goose species (tundra bean geese, greater white-fronted

geese, and lesser white-fronted geese) arrive at their breeding sites after the plant emergence

(Table 1, Figure 4*D*, *F*, and *H*), whereas tundra swans arrive at the breeding sites relatively

earlier, slightly before the emergence of vegetation (Table 1, Figure 4*B*). Mid-latitude

breeding swan geese arrive at the core stopover sites and the breeding sites around three

weeks after the local onset of vegetation growth (Table 1, Figure 4*I* and *J*).

Figure 4. Top resource selection models showing the species variation in the use of

multiple resources. Arrival probability in relation to the availability of seeds and newly

emerged vegetation for East Asian migratory herbivorous waterfowl during spring migration.

The highest arrival probability for arctic-breeding birds occurred after the start of seed

availability (SSA; orange lines) and before the emergence of vegetation (EOV; green lines) in

the Northeast China Plain (NCP) stopover sites **(A, C, E, G)**. All arctic-breeding geese

reached their breeding sites after the emergence of vegetation (EOV in **D, F, H**; green lines)

and close to the end of snowmelt (EOS in **D, F, H**; blue lines). Tundra swans arrived at the

arctic breeding ground earlier (after the onset of snowmelt; OOS in B; grey lines). Mid-

latitude breeding swan geese **(I, J)** reached the Northeast China Plain and the breeding sites

after the emergence of vegetation (EOV; green lines). Colored dashed lines represent 95%

confidence intervals. Vertical dashed lines represent the bird arrival date. Lags (x axis)

indicate the lag days in comparison to the timing of resource availability or the timing of

snowmelt.

All study species use multiple resource waves during spring migration, but variation

among species is observed, as indicated by the best resource selection models evaluated by k-

fold cross validation (higher score, stronger effect; Table 1). At the core stopover sites, the

models for larger-bodied species (tundra swans and swan geese) using the start of seed

availability as a predictor have relatively higher k-fold cross-validation scores, indicating the

use of the seed wave as a cue. For smaller-bodied arctic-breeding geese, the models using the

emergence of plants as a predictor have a stronger effect, indicating the use of the green wave

as a cue. At the breeding sites, the models for larger-bodied species using the emergence of

vegetation have the highest k-fold cross-validation scores, suggesting these waterfowl species

track the green wave. For the relatively smaller-bodied species (tundra bean geese, greater

white-fronted geese, and lesser white-fronted geese), the models that use the end of snowmelt

as predictors show the strongest effect, indicating the use of the snow-free time as a cue.

Table 1. Top resource selection models showing that species using multiple resource waves in migration but relying on specific cues in specific region. The estimated optimal lags (days) on resource availability and snowmelt conditions shows the variation in tracking multiple resource waves at core stopover sites the Northeast China Plain and the breeding sites for five East Asian herbivorous waterfowl species. The estimated range represents the day at which interpolated log-likelihood values have a difference of 1 (ΔAIC = 2). The difference is the number of days between the minimum and maximum estimate of the range. Bold font represents strongest models evaluated by the relatively stronger k-fold cross-validation score (r_s) in each region and for each species.

Region	Species	Models	Estimated days with range	Difference	Log-likelihood	K-fold scores (r_s) Mean	SE
Northeast China Plain	Tundra swan	Plant [*]	4 (-3.4, 13.7)	17.1	-213.57	0.427	0.450
		Seed [†]	**-1 (-5.5, 4.8)**	**10.3**	**-24.10**	**0.571**	**0.169**
	Tundra bean goose	**Plant**	**26 (24.3, 27.8)**	**3.5**	**-101.14**	**0.873**	**0.125**
		Seed	-9 (-11.1, -5.9)	5.2	-42.04	0.503	0.143
	Greater white-fronted goose	**Plant**	**19 (17.7, 21.7)**	**4**	**-1522.14**	**0.935**	**0.025**
		Seed	-20 (-20.7, -18.9)	1.8	-4067.08	0.394	0.278
	Lesser white-fronted goose	**Plant**	**22 (20.2, 24.2)**	**4**	**-466.73**	**0.914**	**0.014**
		Seed	-21 (-22.2, -19.9)	2.3	-1398.77	0.358	0.053
	Swan goose	Plant	-23 (-25.1, -19.5)	5.6	-1107.01	0.403	0.397
		Seed	**-30 (-35.7, -20.6)**	**15.1**	**-1380.99**	**0.487**	**0.546**
Breeding sites	Tundra swan	**Plant**	**14 (12.9, 16)**	**3.1**	**-6198.94**	**0.530**	**0.130**
		Onset of snowmelt	-5 (-6.6, -2.3)	4.3	-5916.80	0.417	0.451
		End of snowmelt	17 (16.1, 17.4)	1.3	-6789.48	0.523	0.108
	Tundra bean goose	Plant	-25 (-26, -23.5)	2.5	-336.80	0.301	0.164
		Onset of snowmelt	-23 (-23.4, -21.6)	1.8	-484.11	0.253	0.165
		End of snowmelt	**-7 (-9.5, -4.7)**	**4.8**	**-514.02**	**0.572**	**0.152**
	Greater white-fronted goose	Plant	-12 (-12.2, -11)	1.2	-6332.97	0.628	0.182
		Onset of snowmelt	-16 (-17.1, -15.6)	1.5	-7455.17	0.434	0.081
		End of snowmelt	**2 (0.5, 3.1)**	**2.6**	**-7569.63**	**0.859**	**0.093**
	Lesser white-fronted goose	Plant	-17 (-17.8, -16.9)	0.9	-2464.9	0.406	0.140
		Onset of snowmelt	-16 (-16.9, -15.9)	1	-1999.08	0.384	0.082
		End of snowmelt	**-10 (-10.1, -9.9)**	**0.2**	**-2171.43**	**0.648**	**0.128**
	Swan goose	**Plant**	**-26 (-26.3, -25.8)**	**0.5**	**-3562.83**	**0.595**	**0.340**
		Onset of snowmelt	-39 (-39.7, -37.7)	2	-3449.32	0.312	0.730
		End of snowmelt	-15 (-16, -14.4)	1.6	-3785.92	0.466	0.242

[*] The model using the spring vegetation emergence as a predictor

[†] The model using the start of seed availability as a predictor

Discussion

We test a new migration strategy in which East Asian herbivorous waterfowl use multiple

resource waves to time their spring migration. The variation in tracking multiple resource

waves for each species and in each region are identified regarding the use of the seed wave

(agricultural seeds in farmland) and the green wave (newly emerging vegetation in natural

habitat). Larger-bodied species follow the seed wave and smaller-bodied ones adopt the green

wave as the cue at the core stopover sites. When arriving at the breeding sites, larger-bodied

species track the green wave whereas smaller-bodied ones track the snow-free time. These

strategies in which migration is timed to be able to take advantage of both natural and

anthropogenic food resources allows East Asian herbivorous waterfowl to exploit the vast and

variable landscape along their migration routes.

Consistent with our first prediction, East Asian herbivorous waterfowl time their staging

in the core stopover sites Northeast China Plain to make use of both anthropogenic and

natural food resources. Seeds left over from previous year's harvest on the unfrozen ground

(from the beginning of April) and newly emerged plants (in early May) allow migratory birds

to replenish their energy for an extended period of time during stopover, which facilitates

them arriving at the breeding grounds in an optimal condition. The Northeast China Plain

contains 15% of the total farming area in China, with maize, soybean, and rice being the three

major crop types [59]. Crop seeds on farmlands provide a high-nutrient food source for

herbivorous waterfowl [28], especially during early spring, when snow just starts melting and

food in the natural habitat is insufficient. Mechanical ploughing (Figure S2*A*) at the end of

April predicts sowing (Figure S2*B* and *C*). In early May, seeds covered with pesticides are

sowed (Figure S2*C*), which means crop seeds on farmlands are no longer useable for

waterfowl (personal field observations and personal communications with local farmers;

Figure S2). When seeds are not available anymore, all study species show a pattern in which

they shift towards the use of the natural habitat.

The exact migration timing differs among species relating to their body size. One

explanation could be that larger-bodied species do not rely on the emergence of natural food

resources as much as the smaller-bodied goose species. Larger-bodied arctic-breeders

generally invest more time on breeding activities and face a more time-constrained window

in which they have to lay eggs, molt, and raise chicks, than smaller-bodied species [4, 60,

61]. The slower pace in their life-histories (i.e., laying larger eggs with a longer incubation

and gosling-rearing period) drives birds to reach the breeding grounds and lay eggs earlier

[61]. This is particularly the case for tundra swans, with the largest body size among the five

herbivorous waterfowl studied, which arrive at the core stopover sites and the arctic breeding

sites the earliest and overtake the green wave, but which can use underwater and underground

food resources once the ice melts [62]. Smaller-bodied grazing species, namely the greater

and lesser white-fronted geese, depart later from the wintering site (Figure 2*J, M* and Table

S1), so that they can forage there on newly emerging grasses before moving north [63-65].

They thereby arrive much later in the core stopover sites and the breeding sites. This strategy

enables them to maximize their time foraging on the newly emergent shoots in their wintering

and stopover areas [8]. Tundra bean geese, slightly larger than greater and lesser white-

fronted geese, arrive at the core stopover sites only one week after seeds have become

available, spend longer stopovers to make use of both resources (Table S1), and arrive at the

breeding sites after the emergence of vegetation. These patterns showing species-dependent

resource utilization at different times likely explain how the species can coexist in a shared

landscape.

In line with our second prediction, larger-bodied herbivorous waterfowl, namely tundra

swans and swan geese, use the seed wave to time their arrival at the core stopover sites. They

generally are less sensitive to forage quality, and therefore can utilize a relatively more

diverse assortment of food resources [39, 64]. They prefer open water where underwater food

resources can be easily accessed (e.g., tubers and algae) that, just like seeds, only become

accessible after snowmelt [62]. This is in line with the habitat use after they reach the

stopover sites, that they spend relatively longer time in water than other smaller-bodied

species (Figure 3). Tundra swan then follow the green wave to arrive at their arctic breeding

sites the earliest. This provides direct benefits for larger-bodied waterfowl, such as

prioritizing resource consumption, and having sufficient time to raise their offspring [61].

Different from our prediction, we found that three smaller-bodied arctic-breeding geese

rely relatively more on natural food resources, and accordingly follow the green wave at core

stopover sites. They arrive in the arctic after vegetation has newly emerged, using the snow-

free time as a cue. While newly emergent vegetation provides a nutritional resource, melting

snow facilitates the emergence of vegetation and provide a cue for animals to exploit

resources at the correct time, which helps to reduce movement and foraging costs related to

snow cover [51]. This finding is in line with a previous study reporting that East Asian geese

arrive at their arctic breeding sites prior to the end of snowmelt [66]. We highlight that

smaller-bodied goose species use the newly emerged vegetation as a primary cue to enable

accumulating sufficient fuel during migration, and avoiding that they arrive too early and

deplete their body stores while awaiting the local thaw, and at the meantime ensuring that

their goslings benefit from abundant food resources later in the season [4, 6, 66].

Our study provides the first evidence for multiple resource tracking in waterfowl spring

migration. The central tenet of resource tracking theory is that mobile consumers can benefit

by moving to exploit phenological variation in resources across space [5]. Tracking multiple

resource waves allows migratory herbivorous waterfowl to prolong the overall resource

availability of natural and anthropogenic resources. Similar to our results, previous study

found that forest birds and stream fishes increase foraging opportunities by feeding on

emerging stream insects early in the year and then switching to terrestrial arthropods later on,

in response to the asynchronous occurrence of fluxes of invertebrates between terrestrial and

freshwater environments [67]. Recent empirical work shows that elephants track both

vegetation green-up in natural habitat and farmland brown down in a coupled human-nature

system [68], maximizing their energy intake by integrating agricultural food resources when

natural food resources is limited. To further our understanding on the response of species to

environmental change, the capability of tracking multiple resource waves should be further

tested in additional study systems and different geographical regions.

Conservation implications

Climate change might induce shifts in vegetation phenology in the spring [69-71], thereby

changing food availability for herbivorous waterfowl at a different pace at different sites [72].

A reduced predictability of food supply due to phenological asynchrony would be

exacerbated when different food resources are exploited. As a consequence, species may

struggle to advance their migration timing to match food availability [73]. Even though some

goose species can accelerate their migration speed under global warming, they might still

suffer population declines if they cannot store sufficient nutrients for egg-laying [60]. For

species taking advantages of the seed wave, migration timing and pace could be influenced

by changes in cultivation practices [36]. For example, changes in farming calendars might

affect seed availability [17, 23, 36]. Different responses to environmental change between the

timing of cultivation and bird migration phenology might cause ecological traps and

population declines of migratory birds [74, 75]. Therefore, changes in habitat management

should consider conservation needs, especially for species that have come to depend on

anthropogenic food resources [76].

Study limitations

We use two different seasonal metrics derived from satellite imagery (i.e., timing of

snowmelt and the second derivative of EVI) to measure two different resource waves (i.e., the

start of seed availability and the emergence of vegetation). Given that data that directly

measure food availability site-by-site in the field are costly to obtain, satellite-derived indices

provide an effective way to measure the seed wave and green wave. They cover more

detailed pixel-by-pixel information in space and time, and better match the GPS locations for

tracked birds. However, satellite-derived indices necessarily contain uncertainties. Firstly, our

estimated plant emergence time (the green wave) based on EVI time-series data reflects the

time that the greening increases at the fastest rate. This time might be later than the shoot

emergence time. Secondly, since seeds are generally spilled and available during the previous

fall and carried over into spring after soil thawing, it is crucial to investigate how many seeds

are actually available in spring, and how much of those seeds are eaten by waterfowl moving

through. Ground-truth data can further optimize the derived plant and seed availability.

Conclusions

We test and demonstrate how migratory herbivorous waterfowl time their spring migration

using both the post-harvest seed availability in cultivated landscapes and the emergence of

vegetation in more natural landscapes. Specifically, we document how the dynamics of

anthropogenic and natural resources act in combination to shape the spring migration

phenology of herbivorous waterfowl. To what extent populations of migratory herbivorous

waterfowl benefit from or are threatened by this migration strategy of tracking multiple

resource waves needs to be thoroughly investigated. For example, how the seed wave itself is

affected by climate and agricultural practices, is still unknown. A proper understanding of the

impact of global environmental change and different human cultivating activities on

migratory bird species will guide effective habitat management for migratory bird

conservation worldwide.

Acknowledgments

We thank Kun Xiao (Sun Yat-sen University, China) for helping processing the remote

sensing data. We are grateful for John Takekawa, Sivananintha Balachandran, Guanhua Liu,

Hao Luo, Duanji Tao, Saras Kumar Behera, Murugesan Ponnaian, Wenyuan Zhang, Yanjie

Xu, Boyu Gao, Zhiyuan Lv, Lingtong Yang, Yongzhi Xiao, and Yunrui Liu who joined the

field work. This research was supported by the National Natural Science Foundation of China

(Grant No. 42301055, No. 41471347, and No. U22A20563).

Author contributions

Conceptualization, F.X., P.G., and Y.S.; Data curation, F.X., J.W., G.L., J.L., B.W., W.Z.W.,

and Y.S.; Methodology, F.X., W.W., J.W., Q.X., W.Z.W., and Y.S.; Software, F.X., W.W., and

J.W.; Investigation, F.X., J.W., and Y.S.; Formal Analysis, F.X., and W.W.; Visualization,

F.X., and Y.S., Writing—Original Draft: F.X., and Y.S.; Writing—Review & Editing: F.X.,

B.W., F.A.L., Z.M., Y.Y., P.G., B.X., and Y.S.; Supervision, B.W., F.A.L., Z.M., Y.Y., and

Y.S.; Project Administration, F.X., and Y.S.; Funding Acquisition, F.X., G.L., and Y.S.

Declaration of interests

564 The authors declare that they have no competing interests.

Data and materials availability

567 Movement tracks of five herbivorous waterfowl species are stored in Movebank

568 (www.movebank.org), under ID 731644623 for lesser white-fronted geese, 1614083125 for

569 greater white-fronted geese, 1614098614 for tundra bean geese, 1614096661 for swan geese,

570 and 1614102131 for tundra swans. The authors declare that all other data supporting the

571 findings of this study are available in the main text or the supplementary materials.

References

1. Thorup K., Tottrup A.P., Willemoes M., Klaassen R.H.G., Strandberg R., Vega M.L., Dasari H.P., Araujo M.B., Wikelski M., Rahbek C. 2017 Resource tracking within and across continents in long-distance bird migrants. *Science Advances* **3**(1). (doi:10.1126/sciadv.1601360).

2. Armstrong J.B., Takimoto G., Schindler D.E., Hayes M.M., Kauffman M.J. 2016 Resource waves: phenological diversity enhances foraging opportunities for mobile consumers. *Ecology* **97**(5), 1099-1112. (doi:10.1890/15-0554.1).

3. Aikens E.O., Kauffman M.J., Merkle J.A., Dwinnell S.P.H., Fralick G.L., Monteith K.L. 2017 The greenscape shapes surfing of resource waves in a large migratory herbivore. *Ecology Letters* **20**(6), 741-750. (doi:10.1111/ele.12772).

4. Si Y., Xin Q., de Boer W.F., Gong P., Ydenberg R.C., Prins H.H.T. 2015 Do Arctic breeding geese track or overtake a green wave during spring migration? *Scientific Reports* **5**(1), 8749. (doi:10.1038/srep08749).

5. Abrahms B., Aikens E.O., Armstrong J.B., Deacy W.W., Kauffman M.J., Merkle J.A. 2021 Emerging Perspectives on Resource Tracking and Animal Movement Ecology. *Trends in Ecology & Evolution* **36**(4), 308-320. (doi:10.1016/j.tree.2020.10.018).

6. van der Graaf S., Stahl J., Klimkowska A., Bakker J.P., Drent R.H. 2006 Surfing on a green wave - how plant growth drives spring migration in the Barnacle Goose *Branta leucopsis*. *Ardea* **94**(3), 567-577.

7. Drent R.H. 1980 Balancing the energy budgets of arctic-breeding geese throughout the annual cycle : a progress report. *Verh Orn Ges Bayern* **23**, 239-264.

8. Kölzsch A., Müskens G.J.D.M., Kruckenberg H., Glazov P., Weinzierl R., Nolet B.A., Wikelski M. 2016 Towards a new understanding of migration timing: slower spring than autumn migration in geese reflects different decision rules for stopover use and departure. *Oikos* **125**(10), 1496-1507. (doi:10.1111/oik.03121).

9. Kölzsch A., Bauer S., de Boer R., Griffin L., Cabot D., Exo K.M., van der Jeugd H.P., Nolet B.A. 2015 Forecasting spring from afar? Timing of migration and predictability of phenology along different migration routes of an avian herbivore. *Journal of Animal Ecology* **84**(1), 272-283. (doi:10.1111/1365-2656.12281).

10. Shariati Najafabadi M., Wang T., Skidmore A.K., Toxopeus A.G., Koelzsch A., Nolet B.A., Exo K.-M., Griffin L., Stahl J., Cabot D. 2014 Migratory Herbivorous Waterfowl Track Satellite-Derived Green Wave Index. *Plos One* **9**(9), e108331. (doi:10.1371/journal.pone.0108331).

11. Rivrud I.M., Heurich M., Krupczynski P., Müller J., Mysterud A. 2016 Green wave tracking by large herbivores: an experimental approach. *Ecology* **97**(12), 3547-3553. (doi:10.1002/ecy.1596).

12. Bischof R., Loe L.E., Meisingset E.L., Zimmermann B., Moorter B.V., Mysterud A. 2012 A Migratory Northern Ungulate in the Pursuit of Spring: Jumping or Surfing the Green Wave? *The American Naturalist* **180**(4), 407-424. (doi:10.1086/667590).

13. Merkle J.A., Monteith K.L., Aikens E.O., Hayes M.M., Hersey K.R., Middleton A.D., Oates B.A., Sawyer H., Scurlock B.M., Kauffman M.J. 2016 Large herbivores surf waves of green-up during spring. *Proceedings Biological sciences* **283**(1833), 20160456. (doi:10.1098/rspb.2016.0456).

14. La Sorte F.A., Graham C.H. 2021 Phenological synchronization of seasonal bird

migration with vegetation greenness across dietary guilds. *Journal of Animal Ecology* **90**(2),

343-355. (doi:10.1111/1365-2656.13345).

15. Wang X., Cao L., Fox A.D., Fuller R., Griffin L., Mitchell C., Zhao Y., Moon O.K.,

Cabot D., Xu Z., et al. 2019 Stochastic simulations reveal few green wave surfing

populations among spring migrating herbivorous waterfowl. *Nat Commun* **10**(1), 2187.

(doi:10.1038/s41467-019-09971-8).

16. Fox A.D., Elmberg J., Tombre I.M., Hessel R. 2017 Agriculture and herbivorous

waterfowl: a review of the scientific basis for improved management. *Biological Reviews*

92(2), 854-877. (doi:10.1111/brv.12258).

17. Blount J.D., Horns J.J., Kittelberger K.D., Neate-Clegg M.H.C., Şekercioğlu Ç.H. 2021

Avian Use of Agricultural Areas as Migration Stopover Sites: A Review of Crop Management

Practices and Ecological Correlates. *Frontiers in Ecology and Evolution* **9**.

(doi:10.3389/fevo.2021.650641).

18. Fox A.D., Madsen J., Boyd H., Kuijken E., Norriss D.W., Tombre I.M., Stroud D.A.

2005 Effects of agricultural change on abundance, fitness components and distribution of two

arctic-nesting goose populations. *Global Change Biology* **11**(6), 881-893.

(doi:10.1111/j.1365-2486.2005.00941.x).

19. Sherfy M.H., Anteau M.J., Bishop A.A. 2011 Agricultural practices and residual corn

during spring crane and waterfowl migration in Nebraska. *The Journal of Wildlife*

Management **75**(5), 995-1003. (doi:10.1002/jwmg.157).

20. Sesser K.A., Iglecia M., Reiter M.E., Strum K.M., Hickey C.M., Kelsey R., Skalos D.A.

2018 Waterbird response to variable-timing of drawdown in rice fields after winter-flooding. *PLoS One* **13**(10), e0204800. (doi:10.1371/journal.pone.0204800).

21. Reynolds M.D., Sullivan B.L., Hallstein E., Matsumoto S., Kelling S., Merrifield M., Fink D., Johnston A., Hochachka W.M., Bruns N.E., et al. 2017 Dynamic conservation for migratory species. *Science Advances* **3**(8), e1700707. (doi:10.1126/sciadv.1700707).

22. Si Y., Xu Y., Xu F., Li X., Zhang W., Wielstra B., Wei J., Liu G., Luo H., Takekawa J., et al. 2018 Spring migration patterns, habitat use, and stopover site protection status for two declining waterfowl species wintering in China as revealed by satellite tracking. *Ecol Evol* **8**(12), 6280-6289. (doi:10.1002/ece3.4174).

23. Si Y., Wei J., Wu W., et al. . 2020 Reducing human pressure on farmland could rescue China's declining wintering geese. *Movement Ecology* **8**(1), 35. (doi:10.1186/s40462-020-00220-y).

24. Amano T. 2009 Conserving bird species in Japanese farmland: Past achievements and future challenges. *Biological Conservation* **142**(9), 1913-1921. (doi:10.1016/j.biocon.2008.12.025).

25. Amano T., Ushiyama K., Fujita G., Higuchi H. 2007 Predicting grazing damage by white-fronted geese under different regimes of agricultural management and the physiological consequences for the geese. *Journal of Applied Ecology* **44**(3), 506-515. (doi:10.1111/j.1365-2664.2007.01314.x).

26. Lane S.J., Azuma A., Higuchi H. 1998 Wildfowl damage to agriculture in Japan. *Agriculture Ecosystems & Environment* **70**(1), 69-77. (doi:10.1016/s0167-8809(98)00114-5).

27. Shimada T. 2002 Daily activity pattern and habitat use of Greater White-fronted Geese

661 wintering in Japan: Factors of the population increase. *Waterbirds* **25**(3), 371-377.

662 (doi:10.1675/1524-4695(2002)025[0371:Dapahu]2.0.Co;2).

663 28. Fox A.D., Abraham K.F. 2017 Why geese benefit from the transition from natural

664 vegetation to agriculture. *Ambio* **46**(2), 188-197. (doi:10.1007/s13280-016-0879-1).

665 29. Donnelly J.P., Moore J.N., Casazza M.L., Coons S.P. 2022 Functional Wetland Loss

666 Drives Emerging Risks to Waterbird Migration Networks. *Frontiers in Ecology and*

667 *Evolution* **10**. (doi:10.3389/fevo.2022.844278).

668 30. Xu W., Fan X., Ma J., Pimm S.L., Kong L., Zeng Y., Li X., Xiao Y., Zheng H., Liu J., et

669 al. 2019 Hidden Loss of Wetlands in China. *Current Biology* **29**(18), 3065-3071.

670 (doi:10.1016/j.cub.2019.07.053).

671 31. Niu Z., Zhang H., Wang X., Yao W., Zhou D., Zhao K., Zhao H., Li N., Huang H., Li C.,

672 et al. 2012 Mapping wetland changes in China between 1978 and 2008. *Chinese Science*

673 *Bulletin* **57**(22), 2813-2823. (doi:10.1007/s11434-012-5093-3).

674 32. Mao D.H., Luo L., Wang Z.M., Wilson M.C., Zeng Y., Wu B.F., Wu J.G. 2018

675 Conversions between natural wetlands and farmland in China: A multiscale geospatial

676 analysis. *Science of the Total Environment* **634**, 550-560.

677 (doi:10.1016/j.scitotenv.2018.04.009).

678 33. Yan F., Zhang S., Liu X., Yu L., Chen D., Yang J., Yang C., Bu K., Chang L. 2017

679 Monitoring spatiotemporal changes of marshes in the Sanjiang Plain, China. *Ecological*

680 *Engineering* **104**, 184-194. (doi:10.1016/j.ecoleng.2017.04.032).

681 34. Miao S., Qiao Y., Han X., Brancher Franco R., Burger M. 2014 Frozen Cropland Soil in

682 Northeast China as Source of N2O and CO2 Emissions. *PLOS ONE* **9**(12), e115761.

(doi:10.1371/journal.pone.0115761).

35. Mwagona P.C., Yao Y., Shan Y., Yu H., Zhang Y. 2018 Trend and Abrupt Regime Shift of

Temperature Extreme in Northeast China, 1957–2015. *Advances in Meteorology* **2018**,

2315372. (doi:10.1155/2018/2315372).

36. Fu W., Zhang X., Zhao J., Du S., Horton R., Hou M. 2018 Artificial warming-mediated

soil freezing and thawing processes can regulate soybean production in Northeast China.

Agricultural and Forest Meteorology **262**, 249-257. (doi:10.1016/j.agrformet.2018.07.015).

37. Zhang Q., Hu Z. 2018 Assessment of drought during corn growing season in Northeast

China. *Theoretical and Applied Climatology* **133**(3), 1315-1321. (doi:10.1007/s00704-018-

2469-6).

38. Lei J., Jia Y., Zuo A., Zeng Q., Shi L., Zhou Y., Zhang H., Lu C., Lei G., Wen L. 2019

Bird Satellite Tracking Revealed Critical Protection Gaps in East Asian⁻Australasian Flyway.

International journal of environmental research and public health **16**(7), 1147.

(doi:10.3390/ijerph16071147).

39. Zhang W., Li X., Yu L., Si Y. 2018 Multi-scale habitat selection by two declining East

Asian waterfowl species at their core spring stopover area. *Ecological Indicators* **87**, 127-

135. (doi:10.1016/j.ecolind.2017.12.035).

40. Kulldorff M., Heffernan R., Hartman J., Assunção R., Mostashari F. 2005 A Space–Time

Permutation Scan Statistic for Disease Outbreak Detection. *PLOS Medicine* **2**(3), e59.

(doi:10.1371/journal.pmed.0020059).

41. Drent R.J., Fox A.D., Stahl J. 2006 Travelling to breed. *Journal of Ornithology* **147**(2),

122-134. (doi:10.1007/s10336-006-0066-4).

42. Ackerman J.T., Takekawa J.Y., Orthmeyer D.L., Fleskes J.P., Yee J.L., Kruse K.L. 2006

Spatial use by wintering Greater White-fronted Geese relative to a decade of habitat change

in California's Central Valley. *The Journal of Wildlife Management* **70**(4), 965-976.

(doi:10.2193/0022-541X(2006)70[965:SUBWGW]2.0.CO;2).

43. Johnson W.P., Schmidt P.M., Taylor D.P. 2014 Foraging flight distances of wintering

ducks and geese: a review. *Avian Conservation and Ecology* **9**, 2. (doi:10.5751/ACE-00683-

090202).

44. Wilman H., Belmaker J., Simpson J., de la Rosa C., Rivadeneira M.M., Jetz W. 2014

EltonTraits 1.0: Species-level foraging attributes of the world's birds and mammals. *Ecology*

95(7), 2027-2027. (doi:10.1890/13-1917.1).

45. Liu H., Gong P., Wang J., Wang X., Ning G., Xu B. 2021 Production of global daily

seamless data cubes and quantification of global land cover change from 1985 to 2020 - iMap

World 1.0. *Remote Sensing of Environment* **258**, 112364. (doi:10.1016/j.rse.2021.112364).

46. Wu W., Sun Y., Xiao K., Xin Q. 2021 Development of a global annual land surface

phenology dataset for 1982–2018 from the AVHRR data by implementing multiple

phenology retrieving methods. *International Journal of Applied Earth Observation and*

Geoinformation **103**, 102487. (doi:10.1016/j.jag.2021.102487).

47. Xin Q., Li J., Li Z., Li Y., Zhou X. 2020 Evaluations and comparisons of rule-based and

machine-learning-based methods to retrieve satellite-based vegetation phenology using

MODIS and USA National Phenology Network data. *International Journal of Applied Earth*

Observation and Geoinformation **93**, 102189. (doi:10.1016/j.jag.2020.102189).

48. Chen X., Wang D., Chen J., Wang C., Shen M. 2018 The mixed pixel effect in land

surface phenology: A simulation study. *Remote Sensing of Environment* **211**, 338-344.

(doi:10.1016/j.rse.2018.04.030).

49. Rickbeil G.J.M., Coops N.C., Berman E.E., McClelland C.J.R., Bolton D.K., Stenhouse

G.B. 2020 Changing spring snow cover dynamics and early season forage availability affect

the behavior of a large carnivore. *Global Change Biology* **26**, 6266-6275.

(doi:10.1111/gcb.15295).

50. Muff S., Signer J., Fieberg J. 2020 Accounting for individual-specific variation in

habitat-selection studies: Efficient estimation of mixed-effects models using Bayesian or

frequentist computation. *Journal of Animal Ecology* **89**(1), 80-92. (doi:10.1111/1365-

2656.13087).

51. Laforge M.P., Bonar M., Vander Wal E. 2021 Tracking snowmelt to jump the green

wave: phenological drivers of migration in a northern ungulate. *Ecology* **102**(3), e03268.

(doi:10.1002/ecy.3268).

52. Brooks M.E., Kristensen K., van Benthem K.J., Magnusson A., Berg C.W., Nielsen A.,

Skaug H.J., Machler M., Bolker B.M. 2017 The R Journal: glmmTMB Balances Speed and

Flexibility Among Packages for Zero-inflated Generalized Linear Mixed Modeling. *The R*

Journal **9**(2), 378-400. (doi:10.32614/rj-2017-066).

53. Roberts D.R., Bahn V., Ciuti S., Boyce M.S., Elith J., Guillera-Arroita G., Hauenstein S.,

Lahoz-Monfort J.J., Schröder B., Thuiller W., et al. 2017 Cross-validation strategies for data

with temporal, spatial, hierarchical, or phylogenetic structure. *Ecography* **40**(8), 913-929.

(doi:10.1111/ecog.02881).

54. Fielding A.H., Bell J.F. 1997 A review of methods for the assessment of prediction errors

in conservation presence/absence models. *Environmental Conservation* **24**(1), 38-49.

(doi:10.1017/S0376892997000088).

55. Boyce M.S., Vernier P.R., Nielsen S.E., Schmiegelow F.K.A. 2002 Evaluating resource

selection functions. *Ecological Modelling* **157**(2), 281-300. (doi:10.1016/S0304-

3800(02)00200-4).

56. Chudzińska M.E., van Beest F.M., Madsen J., Nabe-Nielsen J. 2015 Using habitat

selection theories to predict the spatiotemporal distribution of migratory birds during

stopover – a case study of pink-footed geese *Anser brachyrhynchus*. *Oikos* **124**(7), 851-860.

(doi:10.1111/oik.01881).

57. The MathWorks Inc. 2022 MATLAB version: 9.12.0 (R2022a), Natick, Massachusetts:

The MathWorks Inc. (https://www.mathworks.com)

58. R Core Team. 2021 R: A language and environment for statistical computing. In *R*

Foundation for Statistical Computing (Vienna, Austria).

59. Wang C., Linderholm H.W., Song Y., Wang F., Liu Y., Tian J., Xu J., Song Y., Ren G.

2020 Impacts of Drought on Maize and Soybean Production in Northeast China During the

Past Five Decades. *International journal of environmental research and public health* **17**(7),

2459. (doi:10.3390/ijerph17072459).

60. Lameris T.K., van der Jeugd H.P., Eichhorn G., Dokter A.M., Bouten W., Boom M.P.,

Litvin K.E., Ens B.J., Nolet B.A. 2018 Arctic Geese Tune Migration to a Warming Climate

but Still Suffer from a Phenological Mismatch. *Current Biology* **28**(15), 2467-2473.

(doi:10.1016/j.cub.2018.05.077).

61. Klaassen M. 2003 Relationships Between Migration and Breeding Strategies in Arctic

771 Breeding Birds. *Avian Migration*, 237-249. (doi:10.1007/978-3-662-05957-9_16).

772 62. Nuijten R.J.M., Kölzsch A., van Gils J.A., Hoye B.J., Oosterbeek K., de Vries P.P.,

773 Klaassen M., Nolet B.A. 2014 The exception to the rule: retreating ice front makes Bewick's

774 swans *Cygnus columbianus bewickii* migrate slower in spring than in autumn. *Journal of*

775 *Avian Biology* **45**(2), 113-122. (doi:10.1111/j.1600-048X.2013.00287.x).

776 63. Richman S.E., Leafloor J.O., Karasov W.H., McWilliams S.R. 2015 Ecological

777 implications of reduced forage quality on growth and survival of sympatric geese. *Journal of*

778 *Animal Ecology* **84**(1), 284-298. (doi:10.1111/1365-2656.12270).

779 64. Heuermann N., van Langevelde F., van Wieren S.E., Prins H.H.T. 2011 Increased

780 searching and handling effort in tall swards lead to a Type IV functional response in small

781 grazing herbivores. *Oecologia* **166**(3), 659-669. (doi:10.1007/s00442-010-1894-8).

782 65. Zhang Y., Prins H.H.T., Cao L., Zhao M., Boer W.F.d. 2016 Variation in Elevation and

783 Sward Height Facilitate Coexistence of Goose Species through Allometric Responses in

784 Wetlands. *Waterbirds* **39**(1), 34-44. (doi:10.1675/063.039.0105).

785 66. Li H., Fang L., Wang X., Yi K., Cao L., Fox A.D. 2020 Does snowmelt constrain spring

786 migration progression in sympatric wintering Arctic-nesting geese? Results from a Far East

787 Asia telemetry study. *Ibis* **162**(2), 548-555. (doi:10.1111/ibi.12767).

788 67. Nakano S., Murakami M. 2001 Reciprocal subsidies: Dynamic interdependence between

789 terrestrial and aquatic food webs. *Proceedings of the National Academy of Sciences* **98**(1),

790 166-170. (doi:10.1073/pnas.98.1.166).

791 68. Branco P.S., Merkle J.A., Pringle R.M., Pansu J., Potter A.B., Reynolds A., Stalmans M.,

792 Long R.A. 2019 Determinants of elephant foraging behaviour in a coupled human-natural

system: Is brown the new green? *Journal of Animal Ecology* **88**(5), 780-792.

(doi:10.1111/1365-2656.12971).

69. Socolar J.B., Epanchin P.N., Beissinger S.R., Tingley M.W. 2017 Phenological shifts

conserve thermal niches in North American birds and reshape expectations for climate-driven

range shifts. *Proceedings of the National Academy of Sciences of the United States of*

America **114**(49), 12976-12981. (doi:10.1073/pnas.1705897114).

70. Elsen P.R., Tingley M.W. 2015 Global mountain topography and the fate of montane

species under climate change. *Nature Climate Change* **5**(8), 772-776.

(doi:10.1038/nclimate2656).

71. Tingley M.W., Koo M.S., Moritz C., Rush A.C., Beissinger S.R. 2012 The push and pull

of climate change causes heterogeneous shifts in avian elevational ranges. *Global Change*

Biology **18**(11), 3279-3290. (doi:10.1111/j.1365-2486.2012.02784.x).

72. Mayor S.J., Guralnick R.P., Tingley M.W., Otegui J., Withey J.C., Elmendorf S.C.,

Andrew M.E., Leyk S., Pearse I.S., Schneider D.C. 2017 Increasing phenological asynchrony

between spring green-up and arrival of migratory birds. *Scientific Reports* **7**, 1902.

(doi:10.1038/s41598-017-02045-z).

73. Clausen K.K., Clausen P. 2013 Earlier Arctic springs cause phenological mismatch in

long-distance migrants. *Oecologia* **173**(3), 1101-1112. (doi:10.1007/s00442-013-2681-0).

74. Santangeli A., Lehikoinen A., Bock A., Peltonen-Sainio P., Jauhiainen L., Girardello M.,

Valkama J. 2018 Stronger response of farmland birds than farmers to climate change leads to

the emergence of an ecological trap. *Biological Conservation* **217**, 166-172.

(doi:10.1016/j.biocon.2017.11.002).

75. Outhwaite C.L., McCann P., Newbold T. 2022 Agriculture and climate change are

reshaping insect biodiversity worldwide. *Nature* **605**(7908), 97-102. (doi:10.1038/s41586-

022-04644-x).

76. Li L., Hu R., Huang J., Bürgi M., Zhu Z., Zhong J., Lü Z. 2020 A farmland biodiversity

strategy is needed for China. *Nature Ecology & Evolution* **4**(6), 772-774.

(doi:10.1038/s41559-020-1161-2).

Supporting information

Supplementary information is available for this paper.