Migratory herbivorous waterfowl couple the seed wave and the green

wave during spring migration

Fei Xu[a,b*], Wei Wu[c], Jie Wei[b*], Qinchuan Xin[c], Ben Wielstra[d,e], Frank A. La Sorte[f], Zhijun Ma[g],

Guangchun Lei[h], Jialin Lei[h], Wenzhao Wu[b], Yongchuan Yang[a], Peng Gong[b,i], Bing Xu[b], Yali Si[b,j*]

[a]Key Laboratory of the Three Gorges Reservoir Region's Eco-Environment, Ministry of Education,

Chongqing University, Chongqing 400045, China; [b]Ministry of Education Field Research Station

for East Asian Migratory Birds, Department of Earth System Science, Tsinghua University, Beijing

100086, China; [c]School of Geography and Planning, Sun Yat-sen University, Guangzhou

510275, China; [d]Institute of Biology Leiden, Leiden University, Leiden, the Netherlands; [e]Naturalis

Biodiversity Center, Leiden, the Netherlands; [f]Cornell Lab of Ornithology, Cornell University,

Ithaca, NY, USA; [g]Ministry of Education Key Laboratory for Biodiversity Science and Ecological

Engineering, Coastal Ecosystems Research Station of the Yangtze River Estuary, and Shanghai

Institute of Eco-Chongming (SIEC), Fudan University, Shanghai, 200433, China; [h]School of

Ecology and Nature Conservation, Beijing Forestry University, Beijing 100083, China;

[i]Department of Geography, Department of Earth Sciences, Institute for Climate and Carbon

Neutrality, The University of Hong Kong, Hong Kong, China; and [j]Institute of Environmental

Sciences, Leiden University, 2333 CC, Leiden, Netherlands

*Fei Xu, Jie Wei, Yali Si

Email: feixu@cqu.edu.cn; jwei1029@163.com; y.si@cml.leidenuniv.nl

Author Contributions: F.X., P.G., and Y.S. designed research; F.X., J.W., and Y.S. performed

research; F.X., J.W., G.L., J.L., B.W., W.Z.W., and Y.S. performed data collection; F.X., W.W.,

J.W., Q.X., W.Z.W., and Y.S. analyzed the data; and F.X., B.W., F.A.L., Z.M., Y. Y., P.G., B.X.,

and Y.S. wrote the paper.

Competing Interest Statement: The authors declare no competing interest.

Classification: Biological sciences; Ecology

Keywords: onset of vegetation green-up; agricultural land; migration phenology; resource

tracking; seasonal bird migration

Abstract

In 1980s, many wetlands in China have been transformed into farmlands. In response, East Asian waterfowl populations have taken advantage of spilled seeds left after harvest and now intensively use farmlands during migration. Accordingly, waterfowl might have expanded their migration strategy from merely surfing the green wave of newly emerging vegetation to also incorporating post-harvest agricultural seeds for refueling en route. However, how migratory birds use the phenology of multiple food resources to time their seasonal migration is yet to be explored. Here, we use 166 spring migration tracks of five East Asian herbivorous waterfowl species with a mixed-effect resource selection function model to test how waterfowl use the newly emerging vegetation (the green wave) and the post-harvest agricultural seeds (the seed wave) during spring migration. We find waterfowl arrive at their core stopover sites in the Northeast China Plain after agricultural seeds become available and extend their stay after spring vegetation emerges. Smaller-bodied species tend to arrive at this area later to better use the newly emerged vegetation. Regarding the breeding site, four geese species arrive after the emergence of vegetation, while tundra swans are slightly earlier. Our findings suggest that a strategy of resource coupling has allowed waterfowl to cope with the environmental change that has occurred along their migration routes: they time their migration to exploit both natural and anthropogenic food resources. Management initiatives that combine the preservation of natural habitats with adaptive agricultural practices could safeguard these threatened, long-distance migratory bird species.

Significance Statement

East Asia is experiencing human-induced climate and habitat change at an alarming rate, impacting migratory birds. Conservation challenges largely depend on the cues used by migratory birds to track resources. For the first time, we test a migration strategy of coupling the phenology of natural and anthropogenic food resources, which act in combination to shape the spring migration phenology of herbivorous waterfowl. Waterfowl use both resources by arriving at core stopover sites after seeds become available on farmland and extend their stay after spring vegetation emerges. The exact timing varies among species, with smaller-bodied species, preferring for newly emerging vegetation, arriving later. These findings advance our understanding of migration mechanisms, which will inform and guide effective management and conservation efforts.

Main Text

Introduction

Migratory species move between their breeding and non-breeding grounds and benefit from the exploitation of seasonal food resources across the annual cycle. Foraging opportunities occur in a

series of pulses, at different times and places, which generate "resource waves" across the landscape (1-4). The green wave hypothesis posits that the migration progress of herbivores is driven by the seasonal growth of new vegetation at stopover sites along the migration route (5, 6). This hypothesis has been verified in a variety of herbivorous animal groups, such as geese (3, 5, 7-9) and ungulates (10-12), according to individual-level information compiled using tracking technology (5) to population-level information compiled by large-scale community science initiatives (13). During northward migration, birds mainly arrive at their stopover sites after the emergence of spring vegetation, but can reach their breeding grounds earlier, thus overtaking the green wave, and allowing their offspring to benefit from the peaks of food resources (3, 8). While herbivorous species have been documented to track the green wave in Northern Europe (5, 9, 14) and North America (13), this has not been the case for many herbivorous waterfowl populations worldwide, especially within East Asia (14).

Due to extensive land cover change worldwide, herbivorous waterfowl have expanded from using natural ecosystems to exploiting food resources in agricultural landscapes (15, 16). For example, during the spring-staging period, migratory herbivorous waterfowl have been observed to consume new sown cereals in Europe (17) and post-harvest corn remaining in cornfields (18) and spilled grain in rice paddies (19, 20) in North America. This way they benefit from the spread and post-harvest management of agricultural lands (21, 22). In 1980s, large areas of wetland in China have been converted into farmlands (23-25). East Asian waterfowl populations have started to take advantage of the spilled seeds left after harvest and now intensively use farmlands (26), especially in their core stopover area, the Northeast China Plain (27, 28). As a consequence, the green wave hypothesis (3, 5), generated mainly based on the spatio-temporal dynamics of the natural habitats, does not fully encapsulate the migration strategy adopted by East Asian waterfowl.

Migratory herbivorous waterfowl use the natural habitat (e.g., wetlands, grasslands, shrublands) and farmland habitat differently. Since many waterbodies and grasslands in East Asia are semi-natural, we use the term 'natural habitat' hereafter to represent areas where birds feed on spring sprouts of vegetation. Within farmlands, East Asian waterfowl primarily consume seeds, rather than crop vegetation, e.g., spilled grains and leftover maize (Fig. 1*B*) that remain after harvest (28). These items have been found to offer a higher food quality compared to natural vegetation (15). Spilled seeds on farmland are not available all year round, but only after the harvest during the previous fall and during the time that the ground is not frozen (i.e., during the winter this resource is not accessible). However, how herbivorous waterfowl exploit seeds when they become available (the seed wave) in relation to how they take advantage of the emergence of new vegetation (the green wave) during migration is yet to be explored.

We hypothesize that migratory herbivorous waterfowl could take advantage of the phenology of both seed wave (i.e., seed availability in farmlands using the end of snowmelt as a surrogate; Fig.

1*B*) and green wave (i.e., newly emerged vegetation in natural habitats; Fig. 1*C*) during spring
migration (Fig. 1*A*). We present 166 spring migration tracks from 2015 to 2021, derived from
satellite tracking of 99 individuals for five herbivorous waterfowl species (swan goose *Anser*
cygnoides, tundra bean goose *A. serrirostris*, greater white-fronted goose *A. albifrons*, lesser white-
fronted goose *A. erythropus*, and tundra swan *Cygnus columbianus*) in East Asia. Using a mixed-
effect resource selection function model, we analyze whether birds time their migration to couple
the seed wave and the green wave during spring migration. Investigating how herbivorous
waterfowl track the spatiotemporal availability of these widely different food resources allows
furthering our understanding of migration mechanisms, predicting the impact of land cover and
climate change on migratory birds, and guiding effective habitat management for migratory bird
conservation.

Results

We analyze the relationship between spatiotemporal bird migration patterns (*SI Appendix,* Table
S1 and Fig. S1) and the availability of two food resources (i.e., seeds and newly emerged
vegetation). Spring migration tracks (*n* = 166) from 2015 to 2021 of five waterfowl species within
the East Asian – Australasian Flyway are used: 22 tracks of swan geese, 8 of tundra swans, 54 of
tundra bean geese, 62 of greater white-fronted geese, and 20 of lesser white-fronted geese. Seed
resources at farmlands are all distributed below 55°N latitude (*SI Appendix,* Fig. S2). Tracked birds
utilize both natural habitats (water and surrounding wetlands, grassland, and shrublands) and
farmlands (especially in the Northeast China Plain). In Russia (above 55°N latitude), waterfowl
mainly use water, shrublands, grasslands, and wetlands. They briefly stay at wetlands or rivers that
occurred within forests, which explains the high proportion of forest habitats used in this region (*SI*
Appendix, Fig. S2*F-H*). At breeding sites, tracked birds use grasslands, wetlands, and shrublands
as major habitats (*SI Appendix,* Fig. S2).
Herbivorous waterfowl in East Asia adopt a strategy in which they couple the seed wave with the
green wave during spring migration (Fig. 2 and Fig. 3). Observed arrival timings of all study species
match the period that seeds become available (using the end of snowmelt as a surrogate) within
their core stopover area, the Northeast China Plain, before new vegetation become available (Fig.
2, *SI Appendix,* Fig. S3 and Fig. S4). Across all study species, birds arrive at the core stopover
sites in the Northeast China Plain in early April (96 ± 14 Julian days, Fig. 2). The period when seeds
become available in this region is at the end of March, whereas the emergence of new vegetation
occurs around early May. Waterfowl depart from the Northeast China Plain region in mid-May (131
± 12 Julian days, Fig. 2). In the Russian stopover sites, the observed arrival timings primarily match

the end of snowmelt (around the end of May and the beginning of June). Birds reach their breeding
sites after onset of vegetation growth (around early June) (*SI Appendix,* Fig. S4 and Table S1).
The strategy of coupling the two types of food resources, seeds and newly emerged plants,
varies among species (Fig. 2 and Fig. 3). For arctic-breeding species, those with a relatively larger
body size (i.e., tundra swans and tundra bean geese) show a relatively earlier departure from the
wintering sites ($F = 16.14$, $P < 0.001$, *SI Appendix,* Table S1), and a relatively earlier arrival on the
Northeast China Plain ($F = 40.04$, $P < 0.001$, Fig. 3*A, C, E, and G* and *SI Appendix,* Table S1) and
breeding sites ($F = 7.79$, $P < 0.001$, Fig. 3*B, D, F, and H* and *SI Appendix,* Table S1), in comparison
to those smaller-bodied species (i.e., greater white-fronted geese and lesser white-fronted geese).
Tundra swans arrive at the Northeast China Plain soon (i.e., only one day) after the snow melts
and seeds become accessible (Table 1; Fig. 3*A*), whereas smaller-bodied goose species arrive
later, after seeds have already become available for almost one week (for tundra bean geese; Fig.
3*C*) or for three weeks (for greater white-fronted geese and lesser white-fronted geese; Fig. 3*E and*
G). All arctic-breeding species extend their stay at the Northeast China Plain to be able to forage
on newly emerged plants (Fig. 2).
On the breeding grounds, three arctic-breeding goose species (i.e., tundra bean geese, greater
white-fronted geese, and lesser white-fronted geese) arrive at their breeding sites after the plant
emergence (Table 1, Fig. 3*D*, *F*, and *H*), whereas tundra swans arrive at the breeding sites relatively
earlier, soon after the onset of snowmelt (Table 1, Fig. 3*B*). Mid-latitude breeding swan geese
depart from the wintering site during mid-March (*SI Appendix,* Table S1) and arrive at the Northeast
China Plain and the breeding sites around three weeks after the local onset of vegetation growth
(Table 1, Fig. 3*I* and *J*). They spend a larger percentage of time on farmlands before mid-April, and
then mainly use natural food resource later in the season (*SI Appendix,* Fig. S3).
159
160 **Discussion**
161
We discover a spring migration strategy used by East Asian herbivorous waterfowl in which they
couple the 'seed wave' of seed accessibility on farmland with the 'green wave' of newly emerging
vegetation in natural habitat. All study species arrive at their core stopover sites in the Northeast
China Plain after the start of seed availability on farmlands, extend their stay to use newly emerged
vegetation, and arrive at the breeding sites after the emergence of spring vegetation. Only tundra
swans arrive at the breeding grounds earlier, after the onset of snowmelt. This strategy in which
migration is timed to be able to take advantage of both natural and anthropogenic food resources
allows East Asian herbivorous waterfowl to exploit the vast and variable landscapes along their
migration routes.

East Asian herbivorous waterfowl time their staging in the core stopover sites Northeast China
Plain to make use of both anthropogenic and natural food resources. Seeds leftover from previous
year's harvest on the unfrozen ground (i.e., the beginning of April) and newly emerged plants (i.e.,
in early May) allow migratory birds to replenish their energy for an extended period of time during
stopover, which facilitates arriving at the breeding grounds in an optimal condition. The Northeast
China Plain contains 15% of the total farming area in China, with maize, soybean, and rice being
the three major crop types (29). Crop seeds on farmlands could provide a high-nutrient food source
for herbivorous waterfowl, especially during early spring, when snow just starts melting and food in
their natural habitats is insufficient. Management of agricultural lands has had a great impact on
how birds use seeds. The seed-availability period in the Northeast China Plain ends with the start
of sowing, at the end of April (30) (*SI Appendix,* Fig. S5). Mechanical ploughing (*SI Appendix,* Fig.
S5*A*) at the end of April predicts sowing (*SI Appendix,* Fig. S5*B* and *C*). In early May, seeds covered
with pesticides are sowed (*SI Appendix,* Fig. S5*C*), which means crop seeds on farmlands are no
longer useable for waterfowl (field observations and personal communications with local farmers;
SI Appendix, Fig. S5). When seeds are not available anymore, all study species are able to forage
on newly emerged plants in natural habitats, and depart from the Northeast China Plain in mid-
May.
Although all study species arrive at the core stopover sites after the seed wave and depart after
the spring vegetation emergence, their exact migration timing differs. Arctic breeders (i.e., tundra
swans, tundra bean geese, greater white-fronted geese, and lesser white-fronted geese) arrive at
the Northeast China Plain stopover sites after the start of seed availability and before new
vegetation emerges. The departure date from the wintering sites is later for smaller-bodied species.
The largest-bodied tundra swans leave the earliest and the smallest-bodied lesser white-fronted
geese leave the latest (*SI Appendix,* Table S1), suggesting body size is of importance for migration
strategy. One explanation could be that smaller-bodied grazing species, namely the greater and
lesser white-fronted geese, depart later from the wintering site (*SI Appendix,* Table S1), so that
they can forage on newly emerging grasses in their wintering ground (31-33). By arriving later in
the stopover areas, they could also make use of the new spring vegetation for a longer period.
Larger-bodied herbivorous waterfowl, namely tundra swans and tundra bean geese, generally are
less sensitive to forage quality, and therefore can utilize relatively more diverse assortment of food
resources (32, 34).
Tundra swans prefer open water where underwater food resources can be easily accessed (e.g.,
tubers and algae) that, just like seeds, only become accessible after snowmelt (35). They arrive
the earliest at their core stopover sites (soon after seeds and underwater food become available),
and also move on towards their breeding sites the earliest. Tundra bean geese arrive at the
Northeast China Plain only one week after seeds have become available. In contrast, mid-latitude

breeding swan geese only arrive at the core stopover sites after the emergence of vegetation. This
probably reflects that swan geese could easily get access to nutrient-rich food and reach their
destination under relatively warm weather conditions (36). According to our findings, East Asian
herbivorous waterfowl species we studied take species-dependent utilization of food resources
temporally, though they share similar spatial distributions in the Northeast China Plain. This likely
explains how the species can coexist in the same landscape.

All study species arrive at their breeding sites after the onset of vegetation growth and close to
the end of snowmelt, except for tundra swans, which arrive relatively earlier. This is probably
because larger-bodied arctic-breeders generally invest more time on breeding activities and face a
more time-constrained window in which they have to lay eggs, molt, and raise chicks, than smaller-
bodied species and mid-latitude breeders (3, 37, 38). The slower pace in their life-histories (i.e.,
laying larger eggs with a longer incubation and gosling-rearing period) drives birds to reach the
breeding grounds earlier and lay eggs earlier (38). This is particularly the case for tundra swans,
with the largest body size among the five herbivorous waterfowl, which overtake the green wave,
but can use underwater food resources once the ice melts (35). The three smaller-bodied species
arrive at the arctic later, after vegetation has newly emerged, and close to the end of snowmelt.
This is in line with a previous study reporting that East Asian geese follow the retreating snow
northwards and arrive at their arctic breeding sites prior to the end of snowmelt (39). We find geese
couple the use of seeds and newly emerged vegetation as they move north to enable accumulating
sufficient fuel during migration, and avoiding to arrive so early that they deplete their body stores
prior to the local thaw, therewith ensuring that their goslings benefit from abundant food resources
later in the season (3, 5, 39).

Mid-latitude breeding swan geese reach their breeding grounds after spring vegetation
emergence and around two weeks after the end of snowmelt (i.e., around late April and early May).
The growing season in Inner Mongolia (i.e., breeding grounds for swan geese) typically lasts six
months from May to October (40). This long breeding window makes swan geese less time-
constrained than arctic-breeding species. Swan geese specialize on submerged tubers, which are
only available when lakes have melted, although both field observations and our tracking data
suggest swan geese can graze on sedges and grass (41, 42). A former study observed that swan
geese breeding in Eastern Mongolia (i.e., the same population as our study species) take
significantly longer periods to arrive at their breeding grounds than those breeding in Central or
Western Mongolian, due to a prolonged staging at the core stopover sites in the Northeast China
Plain (43). Their arrival on the breeding grounds coincides with the date that minimum temperatures
are above the freezing point of water (43).

Climate change might induce shifts in vegetation phenology in the spring (44-46), thereby
changing food availability for herbivorous waterfowl at a different pace at different sites (47). A

reduced predictability of food supply due to phenological asynchrony would be exacerbated when
different food resources are coupled. As a consequence, species may struggle to advance their
migration timing to match food availability (48). Even though some goose species can accelerate
their migration speed under global warming, they might still suffer population declines if they cannot
store sufficient nutrients for egg-laying (37). For species taking advantages of the seed wave,
migration timing and pace could be influenced by changes in cultivation practices. For example,
changes in farming calendars might affect seed availability (16, 27). Different responses to
environmental change between the timing of cultivation and bird migration phenology might cause
ecological traps and population declines of migratory birds (49, 50). Therefore, changes in habitat
management should consider conservation needs, especially for species that have been forced to
depend on human-dominated landscapes (51).
Our study offers the first empirical evidence that birds couple both phenology of anthropogenic
and natural food resources in migration. Specifically, we document that anthropogenic and natural
factors act in combination to shape the spring migration phenology of herbivorous waterfowl. We
recommend the resource coupling strategy is further tested in additional species and within different
flyways. To what extent populations of migratory herbivorous waterfowl benefit or suffer from this
coupling strategy needs to be thoroughly investigated. For example, how the seed wave itself is
affected by climate and habitat change, is still unknown. A proper understanding of the impact of
global environmental change and different human cultivating activities on migratory bird species
will inform and guide effective habitat management for migratory bird conservation worldwide.
263
264
265   **Materials and Methods**
266
Analyzing bird migration patterns
We analyzed the satellite tracking data of five East Asian herbivorous waterfowl species. Lesser
white-fronted geese (*Anser erythropus*), greater white-fronted geese (*A. albifrons*), tundra bean
geese (*A. serrirostris*), and tundra swans (*Cygnus columbianus*) were captured and tagged at their
East Dongting Lake (29°N, 113°E) and Poyang Lake (29°N, 116°E) wintering sites in the Yangtze
River Floodplain, China during the 2014 to 2018 wintering seasons. Swan geese (*A. cygnoides*)
were captured at the breeding site Hulun Lake, Hulun Buir, China (48.3°N, 117.4°E) during the
2017 molting season. All birds were equipped with GPS - GSM (Global Positioning System – Global
System for Mobile Communications), solar-powered loggers. GPS locations were recorded every
2 h. Information used in the analysis on migration patterns include Bird ID, track year, longitude,
latitude, and time of the GPS location. Detailed methods on birds capture and deployment of the
satellite transmitters can be found in Si*, et al.* (28) and Lei*, et al.* (26).

279  Migration parameters (*SI Appendix,* Table S1) were derived from 166 spring migration tracks of

280 five species from 2015 to 2021, covering 22 tracks of swan geese from 19 individuals, 8 of tundra

281 swans from 4 individuals, 54 of tundra bean geese from 28 individuals, 62 of greater white-fronted

282 geese from 34 individuals, and 20 of lesser white-fronted geese from 14 individuals. All tracking

283 data of five species are stored in Movebank (www.movebank.org), under ID 731644623 for lesser

284 white-fronted geese, 1614083125 for greater white-fronted geese, 1614098614 for tundra bean

285 geese, 1614096661 for swan geese, and 1614102131 for tundra swans. For each track, we labeled

286 daytime and nighttime locations, by identifying the sunrise and sunset time for each location based

287 on algorithms provided by the National Oceanic & Atmospheric Administration (NOAA)

288 (https://www.esrl.noaa.gov/) (34). The spring migration period, defined as GPS locations recorded

289 from the day that birds left their wintering sites to the day that they reached their breeding sites (7),

290 was included in the analysis. Stopover sites and the length of stay in each site were identified

291 through the space-time permutation model in the SaTScan statistics (http://www.satscan.org) (28).

292 We defined the stopover sites as those sites where birds stayed for at least two days and moved

293 within a radius of less than 50 km (28, 52). The arrival/departure time of the

294 wintering/stopover/breeding sites for each individual in each track year were defined as the first

295 day it arrived/left a specific site in which it showed a continuous presence/absence for at least three

296 days (27). Differences on the migration timing across species were assessed using one-way

297 ANOVA followed by Tukey's honestly significant difference (HSD) multiple comparison. Cumulative

298 length of stay in the Northeast China Plain and Russian stopover sites was defined as the sum of

299 all days spent at stopover sites by an individual in its respective region. Migration distance was

300 defined as the cumulative distance travelled between the wintering site and breeding site (28).

302 **Investigating habitat use along the spring migration route**

303 To examine the habitat use of tracked birds along the migration route, we used two land cover

304 products. The 30 m spatial resolution spring land cover product from 2015 (53) was selected to

305 capture the natural habitats and farmlands within China. The European Space Agency Climate

306 Change Initiative (ESA-CCI) 300 m spatial resolution land cover product from 2015

307 (http://maps.elie.ucl.ac.be/CCI/viewer/index.php) was used to capture the relatively homogeneous

308 landscape within Russia. Two classification systems from the two products were aggregated into

309 seven land cover types: grassland, farmland, water, wetland, forest, shrubland, and 'other' (i.e.,

310 barren land, lichens and mosses, snow and ice, or impervious). Habitat use at each stopover site

311 was estimated using the percentage of bird GPS locations on each land cover type (28). The

312 relative proportion of land cover types in use at each latitude was then calculated as the sum of the

313 percentage of bird GPS locations on each land cover type across all stopover sites (their rounded

314 centroid coordinates) located within the same latitudinal range. As waterfowl mostly forage during

the day, we also calculated the habitat use at daytime (i.e., the proportion of land cover in use

during the day) across the whole spring migration period, to better interpret how waterfowl use the

food resources across time.

Calculating the timing of the seed wave and the green wave

For each track, the spring vegetation emergence time in natural habitats and the start of seed

availability time on farmlands were calculated. We used the second-order derivative (54, 55) of the

smoothed time series of Enhanced Vegetation Index (EVI) derived from the Moderate Resolution

Imaging Spectroradiometer (MODIS) Terra 8-day 250 m spatial resolution surface reflectance

product MOD09Q1 to extract the spring vegetation emergence time (details are provided in *SI*

Appendix, SI methods S1). Only pure pixels where the proportion of herbaceous area (i.e.,

wetlands, grasslands, and shrublands) was larger than 0.5 within a 250 m grid cell were used

(details are provided in *SI Appendix*, SI methods S2 and Fig. S6).

The start of the seed availability time was determined by the end of snowmelt in spring because

after the snow melt seeds (remaining from previous year's harvest) could be easily dug out from

the ground. Following the method of Rickbeil *, et al.* (56), we used the MODIS Terra Snow Cover

daily 500 m spatial resolution product MOD10A1 to measure the different degree of snowmelt (i.e.,

the onset and the end of snowmelt). We first created a time-series to capture the daily percent of

snow cover (from day 1 to day 210) for each stopover and breeding site, for each track. We then

applied three-piece segmented regression to extract the onset and the end of snowmelt for each

pixel (*SI Appendix,* Fig. S7). To facilitate interpreting the resource use, in addition to the vegetation

availability date and the seed availability date, we also calculated the onset of snowmelt and the

end of snowmelt at each site, for each individual bird, in each track year.

Determining the resource availability – bird arrival relationship

We used a resource selection function (RSF) framework (57, 58) to investigate to what extent the

start of resource availability can be matched with the arrival date at the stopover or breeding sites

for migratory herbivorous waterfowl during spring migration. Using a logistic regression of

presence/absence points, RSFs can be used to define the selection of resources on the day when

waterfowl arrive at a specific site (27). The presence points were defined as all GPS locations

recorded on the day of arrival. The absence points were drawn randomly on the date before and

after arrival at a specific site during the spring migration period. For each pixel in the specific site,

we calculated the number of days between each of the four indices (i.e., the spring vegetation

emergence, the start of seed availability, the onset of snowmelt, and the end of snowmelt) and the

bird arrival date. Migratory waterfowl might not track the exact day that a specific resource becomes

available but may display a lag effect, where birds arrive before (positive lag value) or after

(negative lag value) the resource becomes available (14). In order to obtain the focal lag (i.e., the

number of days before or after the resource available or snow melt dates), we tested the predictive

ability of the models across a series of temporal lags, by adding or subtracting days to the resource

becoming available or the snow melt dates, following (58). For example, if a bird arrives at the

stopover site two days after seed availability, the model with a lag of two (by subtracting two from

the day to the seed availability date) would have the highest likelihood.

We executed RSFs in two regions for each species: the core stopover area Northeast China

Plain and the breeding sites. We ran two models in the Northeast China Plain stopover sites (35°-

55°N latitude) using vegetation and seed availability as the predictors, and three both at the mid-

latitude (around 46°-49°N latitude) and arctic breeding sites (around 70°-75°N latitude) using

vegetation availability, the onset of snowmelt, and the end of snowmelt as predictors. For each of

the five waterfowl species, within each region, we fit a mixed effects conditional logistic regression

model using a binomial distribution as the link function. The response variable was the probability

of bird arrival. The presence was set to 1 and absence was set to 0. The fixed effect for each model

was a specific type of temporal lag, i.e., lag days to 1) the spring vegetation emergence, 2) the start

of seed availability, 3) the onset of snowmelt, or 4) the end of snowmelt. The random effect was

the identity of the individual track (track ID). We used the R package glmmTMB to run the analysis

(59).

We modelled the lags in days to food availability or the snow melt conditions that fell between

the 10th and 90th percentile of all locations within each region used by herbivorous waterfowl. We

selected the top model for the focal lag based on the lowest Akaike Information Criterion (AIC) (58).

The range around the optimal lag was also estimated by predicting the day at which interpolated

log-likelihood values have a difference of 1 (ΔAIC = 2). The difference in range was then calculated

by the number of days between the minimum and maximum estimate. The predictive performance

of the top models with the optimal estimated lag was evaluated by k-fold cross-validation (58, 60).

Acknowledgments

We thank Kun Xiao (Sun Yat-sen University, China) for helping processing the remote sensing

data. We are grateful for John Takekawa, Sivananintha Balachandran, Guanhua Liu, Hao Luo,

Duanji Tao, Saras Kumar Behera, Murugesan Ponnaian, Wenyuan Zhang, Yanjie Xu, Boyu Gao,

Zhiyuan Lv, Lingtong Yang, Yongzhi Xiao, and Yunrui Liu who joined the field work. This research

was supported by the National Natural Science Foundation of China (No. 41471347 and No.

U22A20563).

References

1. J. B. Armstrong, G. Takimoto, D. E. Schindler, M. M. Hayes, M. J. Kauffman, Resource
waves: phenological diversity enhances foraging opportunities for mobile consumers.
Ecology **97**, 1099-1112 (2016).
2. E. O. Aikens *et al.*, The greenscape shapes surfing of resource waves in a large migratory
herbivore. *Ecology Letters* **20**, 741-750 (2017).
3. Y. Si *et al.*, Do Arctic breeding geese track or overtake a green wave during spring
migration? *Scientific Reports* **5**, 8749 (2015).
4. B. Abrahms *et al.*, Emerging Perspectives on Resource Tracking and Animal Movement
Ecology. *Trends in Ecology & Evolution* **36**, 308-320 (2021).
5. S. van der Graaf, J. Stahl, A. Klimkowska, J. P. Bakker, R. H. Drent, Surfing on a green
wave - how plant growth drives spring migration in the Barnacle Goose *Branta leucopsis*.
Ardea **94**, 567-577 (2006).
6. R. H. Drent, Balancing the energy budgets of arctic-breeding geese throughout the
annual cycle : a progress report. *Verh Orn Ges Bayern* **23**, 239-264 (1980).
7. A. Kölzsch *et al.*, Towards a new understanding of migration timing: slower spring than
autumn migration in geese reflects different decision rules for stopover use and
departure. *Oikos* **125**, 1496-1507 (2016).
8. A. Kölzsch *et al.*, Forecasting spring from afar? Timing of migration and predictability of
phenology along different migration routes of an avian herbivore. *The Journal of animal
ecology* **84**, 272-283 (2015).
9. M. Shariati Najafabadi *et al.*, Migratory Herbivorous Waterfowl Track Satellite-Derived
Green Wave Index. *Plos One* **9**, e108331 (2014).
10. I. M. Rivrud, M. Heurich, P. Krupczynski, J. Müller, A. Mysterud, Green wave tracking by
large herbivores: an experimental approach. *Ecology* **97**, 3547-3553 (2016).
11. R. Bischof *et al.*, A Migratory Northern Ungulate in the Pursuit of Spring: Jumping or
Surfing the Green Wave? *The American Naturalist* **180**, 407-424 (2012).
12. J. A. Merkle *et al.*, Large herbivores surf waves of green-up during spring. *Proceedings.
Biological sciences* **283**, 20160456 (2016).
13. F. A. La Sorte, C. H. Graham, Phenological synchronization of seasonal bird migration
with vegetation greenness across dietary guilds. *Journal of Animal Ecology* **90**, 343-355
(2021).
14. X. Wang *et al.*, Stochastic simulations reveal few green wave surfing populations among
spring migrating herbivorous waterfowl. *Nat Commun* **10**, 2187 (2019).
15. A. D. Fox, J. Elmberg, I. M. Tombre, R. Hessel, Agriculture and herbivorous waterfowl: a
review of the scientific basis for improved management. *Biological Reviews* **92**, 854-877
(2017).
16. J. D. Blount, J. J. Horns, K. D. Kittelberger, M. H. C. Neate-Clegg, Ç. H. Şekercioğlu, Avian
Use of Agricultural Areas as Migration Stopover Sites: A Review of Crop Management
Practices and Ecological Correlates. *Frontiers in Ecology and Evolution* **9** (2021).
17. A. D. Fox *et al.*, Effects of agricultural change on abundance, fitness components and
distribution of two arctic-nesting goose populations. *Global Change Biology* **11**, 881-893
(2005).
18. M. H. Sherfy, M. J. Anteau, A. A. Bishop, Agricultural practices and residual corn during
spring crane and waterfowl migration in Nebraska. *The Journal of Wildlife Management*
75, 995-1003 (2011).

19. K. A. Sesser *et al.*, Waterbird response to variable-timing of drawdown in rice fields after
winter-flooding. *PLoS One* **13**, e0204800 (2018).
20. M. D. Reynolds *et al.*, Dynamic conservation for migratory species. *Science Advances* **3**,
e1700707 (2017).
21. A. D. Fox, K. F. Abraham, Why geese benefit from the transition from natural vegetation
to agriculture. *Ambio* **46**, 188-197 (2017).
22. J. P. Donnelly, J. N. Moore, M. L. Casazza, S. P. Coons, Functional Wetland Loss Drives
Emerging Risks to Waterbird Migration Networks. *Frontiers in Ecology and Evolution* **10**
(2022).
23. W. Xu *et al.*, Hidden Loss of Wetlands in China. *Current Biology* **29**, 3065-3071 (2019).
24. Z. Niu *et al.*, Mapping wetland changes in China between 1978 and 2008. *Chinese
Science Bulletin* **57**, 2813-2823 (2012).
25. D. H. Mao *et al.*, Conversions between natural wetlands and farmland in China: A
multiscale geospatial analysis. *Science of the Total Environment* **634**, 550-560 (2018).
26. J. Lei *et al.*, Bird Satellite Tracking Revealed Critical Protection Gaps in East
Asian‾Australasian Flyway. *International journal of environmental research and public
health* **16**, 1147 (2019).
27. Y. Si, J. Wei, W. Wu, et al. , Reducing human pressure on farmland could rescue China's
declining wintering geese. *Movement Ecology* **8**, 35 (2020).
28. Y. Si *et al.*, Spring migration patterns, habitat use, and stopover site protection status for
two declining waterfowl species wintering in China as revealed by satellite tracking. *Ecol
Evol* **8**, 6280-6289 (2018).
29. C. Wang *et al.*, Impacts of Drought on Maize and Soybean Production in Northeast China
During the Past Five Decades. *International journal of environmental research and public
health* **17**, 2459 (2020).
30. Q. Zhang, Z. Hu, Assessment of drought during corn growing season in Northeast China.
Theoretical and Applied Climatology **133**, 1315-1321 (2018).
31. S. E. Richman, J. O. Leafloor, W. H. Karasov, S. R. McWilliams, Ecological implications of
reduced forage quality on growth and survival of sympatric geese. *Journal of Animal
Ecology* **84**, 284-298 (2015).
32. N. Heuermann, F. van Langevelde, S. E. van Wieren, H. H. T. Prins, Increased searching
and handling effort in tall swards lead to a Type IV functional response in small grazing
herbivores. *Oecologia* **166**, 659-669 (2011).
33. Y. Zhang, H. H. T. Prins, L. Cao, M. Zhao, W. F. d. Boer, Variation in Elevation and Sward
Height Facilitate Coexistence of Goose Species through Allometric Responses in
Wetlands. *Waterbirds* **39**, 34-44 (2016).
34. W. Zhang, X. Li, L. Yu, Y. Si, Multi-scale habitat selection by two declining East Asian
waterfowl species at their core spring stopover area. *Ecological Indicators* **87**, 127-135
(2018).
35. R. J. M. Nuijten *et al.*, The exception to the rule: retreating ice front makes Bewick's
swans *Cygnus columbianus bewickii* migrate slower in spring than in autumn. *Journal of
Avian Biology* **45**, 113-122 (2014).
36. N. Batbayar *et al.*, Migration strategies of Swan Geese *Anser cygnoides* from northeast
Mongolia. *Wildfowl* **61**, 90-109 (2011).
37. T. K. Lameris *et al.*, Arctic Geese Tune Migration to a Warming Climate but Still Suffer
from a Phenological Mismatch. *Current Biology* **28**, 2467-2473 (2018).

38. M. Klaassen, Relationships Between Migration and Breeding Strategies in Arctic
Breeding Birds. in *Avian Migration*, eds P. Berthold, E. Gwinner, E. Sonnenschein
(Springer Berlin Heidelberg, Berlin, Heidelberg, 2003), pp 237-249.
39. H. Li *et al.*, Does snowmelt constrain spring migration progression in sympatric wintering
Arctic-nesting geese? Results from a Far East Asia telemetry study. *Ibis* **162**, 548-555
(2020).
40. Z. Jin, Y. Qi, D. Yunshe, M. Domroes, Seasonal patterns of soil respiration in three types
of communities along grass-desert shrub transition in Inner Mongolia, China. *Advances*
in Atmospheric Sciences **26**, 503-512 (2009).
41. A. D. Fox *et al.*, Preliminary observations of diurnal feeding patterns of Swan Geese
Anser cygnoides using two different habitats at Shengjin Lake, Anhui Province, China.
Wildfowl, 20-30 (2013).
42. Y. Chen, Y. Zhang, L. Cao, W. F. de Boer, A. D. Fox, Wintering Swan Geese maximize
energy intake through substrate foraging depth when feeding on buried *Vallisneria*
natans tubers. *Avian Research* **10**, 6 (2019).
43. I. Damba *et al.*, Seasonal and regional differences in migration patterns and
conservation status of Swan Geese (*Anser cygnoides*) in the East Asian Flyway. *Avian*
Research **12**, 73 (2021).
44. J. B. Socolar, P. N. Epanchin, S. R. Beissinger, M. W. Tingley, Phenological shifts conserve
thermal niches in North American birds and reshape expectations for climate-driven
range shifts. *Proceedings of the National Academy of Sciences of the United States of*
America **114**, 12976-12981 (2017).
45. P. R. Elsen, M. W. Tingley, Global mountain topography and the fate of montane species
under climate change. *Nature Climate Change* **5**, 772-U192 (2015).
46. M. W. Tingley, M. S. Koo, C. Moritz, A. C. Rush, S. R. Beissinger, The push and pull of
climate change causes heterogeneous shifts in avian elevational ranges. *Global Change*
Biology **18**, 3279-3290 (2012).
47. S. J. Mayor *et al.*, Increasing phenological asynchrony between spring green-up and
arrival of migratory birds. *Scientific Reports* **7**, 1902 (2017).
48. K. K. Clausen, P. Clausen, Earlier Arctic springs cause phenological mismatch in long-
distance migrants. *Oecologia* **173**, 1101-1112 (2013).
49. A. Santangeli *et al.*, Stronger response of farmland birds than farmers to climate change
leads to the emergence of an ecological trap. *Biological Conservation* **217**, 166-172
(2018).
50. C. L. Outhwaite, P. McCann, T. Newbold, Agriculture and climate change are reshaping
insect biodiversity worldwide. *Nature* **605**, 97-102 (2022).
51. L. Li *et al.*, A farmland biodiversity strategy is needed for China. *Nature Ecology &*
Evolution **4**, 772-774 (2020).
52. R. J. Drent, A. D. Fox, J. Stahl, Travelling to breed. *Journal of Ornithology* **147**, 122-134
(2006).
53. H. Liu *et al.*, Production of global daily seamless data cubes and quantification of global
land cover change from 1985 to 2020 - iMap World 1.0. *Remote Sensing of Environment*
258, 112364 (2021).
54. W. Wu, Y. Sun, K. Xiao, Q. Xin, Development of a global annual land surface phenology
dataset for 1982–2018 from the AVHRR data by implementing multiple phenology

retrieving methods. *International Journal of Applied Earth Observation and*
Geoinformation **103**, 102487 (2021).
55.   Q. Xin, J. Li, Z. Li, Y. Li, X. Zhou, Evaluations and comparisons of rule-based and machine-
learning-based methods to retrieve satellite-based vegetation phenology using MODIS
and USA National Phenology Network data. *International Journal of Applied Earth*
Observation and Geoinformation **93**, 102189 (2020).
56.   G. J. M. Rickbeil *et al.*, Changing spring snow cover dynamics and early season forage
availability affect the behavior of a large carnivore. *Global Change Biology* **26**, 6266-
6275 (2020).
57.   S. Muff, J. Signer, J. Fieberg, Accounting for individual-specific variation in habitat-
selection studies: Efficient estimation of mixed-effects models using Bayesian or
frequentist computation. *Journal of Animal Ecology* **89**, 80-92 (2020).
58.   M. P. Laforge, M. Bonar, E. Vander Wal, Tracking snowmelt to jump the green wave:
phenological drivers of migration in a northern ungulate. *Ecology* **102**, e03268 (2021).
59.   M. E. Brooks *et al.*, glmmTMB Balances Speed and Flexibility Among Packages for Zero-
inflated Generalized Linear Mixed Modeling. *R Journal* **9**, 378-400 (2017).
60.   D. R. Roberts *et al.*, Cross-validation strategies for data with temporal, spatial,
hierarchical, or phylogenetic structure. *Ecography* **40**, 913-929 (2017).

Figures and Tables

543
Figure 1. Schematic diagram of the newly discovered spring migration strategy adopted by East
Asian migratory herbivorous waterfowl in which they couple two food resources. (A) Core stopover
sites (S1) and breeding sites (S2) along the spring migration route. (B) During spring migration,
birds arrive on farmlands at the core stopover sites after the start of seed availability (seed wave;
orange line), extend their stay after spring vegetation in natural habitats emerges (green wave;
green lines) (S1) and reach the breeding grounds after the emergence of vegetation (green wave;
green lines) (S2). (C) Photo of greater white-fronted geese (*Anser albifrons*) and tundra bean geese
(*A. serrirostris*) foraging in farmlands (taken 19 March 2021) on Yuquan island, Hunchun, Yanbian
Korean Nationality Autonomous Prefecture, Jilin, China (photo credit: Haixiang Zhou). Inset photo
shows leftover maize (taken 2 May 2016) on farmlands in Fuyu Village, Nehe, Qiqihar, China (photo
credit: Jie Wei). (D) Photo of new spring vegetation (taken 1 May 2016) embedded in old vegetation
in Baishan town, Longjiang, Qiqihar, China (photo credit: Jie Wei).
556

557
Figure 2. Coupling the seed and green wave at core stopover sites in the Northeast China Plain.
All study species (A-E) arrive (vertical red line with error bar) at the Northeast China Plain after the
start of seed availability (vertical orange line with error bar) and departed (vertical black line with
error bar) from this region after the emergence of vegetation (vertical green line with error bar),
thereby use both food resources. Grey lines show annual food availability indicated by enhanced
vegetation index (EVI) time series.
564

565
566
Figure 3. Arrival probability in relation to the availability of seeds and newly emerged vegetation
for East Asian migratory herbivorous waterfowl during spring migration. The highest arrival
probability for arctic-breeding birds occurred after the start of seed availability (SSA; orange lines)
and before the emergence of vegetation (EOV; green lines) in the Northeast China Plain (NCP)
stopover sites (A, C, E, G). All arctic-breeding geese reached their breeding sites after the
emergence of vegetation (EOV in D, F, H; green lines) and close to the end of snowmelt (EOS in
D, F, H; blue lines). Tundra swans arrived at the arctic breeding ground earlier (after the onset of
snowmelt; OOS in B; grey lines). Mid-latitude breeding swan geese (I, J) reached the Northeast
China Plain and the breeding sites after the emergence of vegetation (EOV; green lines). Colored
dashed lines represent 95% confidence intervals. Vertical dashed lines represent the bird arrival
date. Lags (x axis) indicate the lag days in comparison to the timing of resource availability or the
timing of snowmelt.

Table 1. The top resource selection models showing the estimated optimal lags (days) on
resource availability and snowmelt conditions for five East Asian herbivorous waterfowl species at
core stopover sites the Northeast China Plain and the breeding sites.

Region	Species	Models	Estimated days with range	Difference	Log-likelihood	K-fold scores Mean	SE
Northeast China Plain	Tundra swan	Plant *	4 (-3.4, 13.7)	17.1	-213.57	0.427	0.450
		Seed †	-1 (-5.5, 4.8)	10.3	-24.10	0.571	0.169
	Tundra bean goose	Plant	26 (24.3, 27.8)	3.5	-101.14	0.873	0.125
		Seed	-9 (-11.1, -5.9)	5.2	-42.04	0.503	0.143
	Greater white-fronted goose	Plant	19 (17.7, 21.7)	4	-1522.14	0.935	0.025
		Seed	-20 (-20.7, -18.9)	1.8	-4067.08	0.394	0.278
	Lesser white-fronted goose	Plant	22 (20.2, 24.2)	4	-466.73	0.914	0.014
		Seed	-21 (-22.2, -19.9)	2.3	-1398.77	0.358	0.053
	Swan goose	Plant	-23 (-25.1, -19.5)	5.6	-1107.01	0.403	0.397
		Seed	-30 (-35.7, -20.6)	15.1	-1380.99	0.487	0.546
Breeding sites	Tundra swan	Plant	14 (12.9, 16)	3.1	-6198.94	0.530	0.130
		Onset of snowmelt	-5 (-6.6, -2.3)	4.3	-5916.80	0.417	0.451
		End of snowmelt	17 (16.1, 17.4)	1.3	-6789.48	0.523	0.108
	Tundra bean goose	Plant	-25 (-26, -23.5)	2.5	-336.80	0.301	0.164
		Onset of snowmelt	-23 (-23.4, -21.6)	1.8	-484.11	0.253	0.165
		End of snowmelt	-7 (-9.5, -4.7)	4.8	-514.02	0.572	0.152
	Greater white-fronted goose	Plant	-12 (-12.2, -11)	1.2	-6332.97	0.628	0.182
		Onset of snowmelt	-16 (-17.1, -15.6)	1.5	-7455.17	0.434	0.081
		End of snowmelt	2 (0.5, 3.1)	2.6	-7569.63	0.859	0.093
	Lesser white-fronted goose	Plant	-17 (-17.8, -16.9)	0.9	-2464.9	0.406	0.140
		Onset of snowmelt	-16 (-16.9, -15.9)	1	-1999.08	0.384	0.082
		End of snowmelt	-10 (-10.1, -9.9)	0.2	-2171.43	0.648	0.128
	Swan goose	Plant	-26 (-26.3, -25.8)	0.5	-3562.83	0.595	0.340
		Onset of snowmelt	-39 (-39.7, -37.7)	2	-3449.32	0.312	0.730
		End of snowmelt	-15 (-16, -14.4)	1.6	-3785.92	0.466	0.242

583 * The model using the spring vegetation emergence as a predictor

584 † The model using the start of seed availability as a predictor