What AlphaFold tells us about cohesin's retention on

and release from chromosomes

Kim A. Nasmyth[1], Byung-Gil Lee[2, 3], Maurici B. Roig[1] and Jan Löwe[2]

[1] Department of Biochemistry, University of Oxford, Oxford OX1 3QU, UK;

[2] MRC Laboratory of Molecular Biology, Cambridge CB2 0QH, UK

[3]Current address: Department of Biochemistry, Lee Gil Ya Cancer and Diabetes Institute,

Gachon University College of Medicine, Incheon 21999, Republic of Korea

ABSTRACT

Cohesin is a trimeric complex containing a pair of SMC proteins (Smc1 and Smc3) whose

ATPase domains at the end of long coiled coils (CC) are interconnected by Scc1. During

interphase, it organizes chromosomal DNA topology by extruding loops in a manner dependent

on Scc1's association with two large hook shaped proteins called SA (yeast: Scc3) and Nipbl

(Scc2). The latter's replacement by Pds5 recruits Wapl, which induces release from chromatin

via a process requiring dissociation of Scc1's N-terminal domain (NTD) from Smc3. If blocked

by Esco (Eco)-mediated Smc3 acetylation, cohesin containing Pds5 merely maintains pre-

existing loops, but a third fate occurs during DNA replication, when Pds5-containing cohesin

associates with Sororin and forms structures that hold sister DNAs together. How Wapl induces

and Sororin blocks release has hitherto remained mysterious. In the twenty years since their

discovery, not a single testable hypothesis has been proposed as to their role. Here, AlphaFold

2 (AF) three-dimensional protein structure predictions lead us to propose formation of a

quarternary complex between Wapl, SA, Pds5, and Scc1's NTD, in which the latter is

juxtaposed with (and subsequently sequestered by) a highly conserved cleft within Wapl's C-

terminal domain (CTD). AF also reveals how Scc1's dissociation from Smc3 arises from a

distortion of Smc3's CC induced by engagement of SMC ATPase domains, how Esco acetyl

transferases are recruited to Smc3 by Pds5, and how Sororin prevents release by binding to the

Smc3/Scc1 interface. Our hypotheses explain the phenotypes of numerous existing mutations

and are highly testable.

INTRODUCTION

Cohesin is a multi-subunit complex responsible for determining much of the topology of chromosomal DNA during interphase as well as sister chromatid cohesion (Yatskevich et al., 2019). At its heart are a pair of rod-shaped Structural Maintenance of Chromosomes (SMC) proteins, Smc1 and Smc3, that have a dimerization domain called the hinge at one end and an ABC-like ATPase domain at the other, separated by 50 nm long intra-molecular anti-parallel coiled coils. These are interrupted in two places, at an "elbow", 16 nm from the hinge, around which the coiled coils fold onto each other (Bürmann et al., 2019), and a disruption of the coiled coil called the joint (Diebold-Durand et al., 2017), about 6.5 nm from the ATPase domains (Fig. 1).

In the presence of ATP, the Smc1 and Smc3 ATPase domains associate with each other, thereby sandwiching a pair of ATP molecules between them (Shi et al., 2020). Crucially, they are also inter-connected by a kleisin subunit (Schleiffer et al., 2003), called Scc1, whose N- and C-terminal domains bind to the "base" and "neck" of the Smc1 and Smc3 ATPases respectively, creating a huge tripartite SMC-kleisin ring, even in the absence of ATP (Chapard et al., 2019; Haering et al., 2002) (Fig. 1). Cohesin's various activities depend on the association with the largely unstructured sections of the kleisin polypeptide connecting its N- and C-terminal domains, of three large hook-shaped HEAT repeat containing regulatory subunits called HAWKs (HEAT repeat proteins Associated With Kleisins). Only two of these are thought to be bound at any one time, Scc3 (human: SA) with either Pds5 or Scc2 (human: Nipbl). Complexes containing Scc2/Nipbl are active as DNA dependent ATPases, while those containing Pds5 are not (Petela et al., 2018).

In addition to holding sister chromatids together in post-replicative cells, cohesin is capable of extruding long loops of DNA (Davidson et al., 2019; Kim et al., 2019). This loop extruding (LE) activity occurs throughout interphase in mitotic cells and during prophase in meiosis when it is essential for formation of the synaptonemal complex (Klein et al., 1999). Importantly, LE and sister chromatid cohesion are conferred by distinct populations of cohesin, as LE requires the continuous presence of Scc2/Nipbl (Mitter et al., 2020), whereas cohesion does not (Ciosk et al., 2000; Srinivasan et al., 2019). LE facilitates VDJ recombination (Zhang et al., 2022), as well as the interaction of enhancers with distant promoters and is regulated throughout the genome by the site-specific DNA binding protein CTCF (Fudenberg et al., 2016; Li et al., 2020).

Cohesin's initial association with chromosomes as well as active loop extrusion (LE) are
mediated by Smc-kleisin rings occupied by Scc2/Nipbl and Scc3/SA (Fig. 2). LE is halted at
asymmetric CTCF sites, but only when they face in one direction (Fudenberg et al., 2016). This
arrest depends on modification by yeast Eco1 (Bastie et al., 2022) or human Esco1 and 2 (van
Ruiten et al., 2022; Wutz et al., 2020) acetyl-transferases of a pair of lysine residues (yeast
Smc3K112 and K113; human Smc3K105 and K106) on the ATPase domain of Smc3 (Unal et
al., 2008), an event that depends on replacement of Scc2/Nipbl by Pds5 (Chan et al., 2013)
(Fig. 2). Unlike Scc2/Nipbl, which is required for LE and for building but not maintaining
cohesion, SA/Scc3 is essential for all of cohesin's activities, namely loading onto chromatin
and loop extrusion, as well as the establishment and maintenance of cohesion (Davidson et al.,
2019; Roig et al., 2014; Toth et al., 1999). Unrestricted LE between convergent CTCF sites,
but not past them, ensures that any sequence between convergent sites is more likely to be
associated with other sequences within this interval than with those beyond these sites, giving
rise to the phenomenon of Topologically Associated Domains (TADs) in HiC genome contact
maps (Rao et al., 2017). The inhibition of cohesin's LE by CTCF is presumed to depend on
Nipbl's replacement by Pds5, as well as acetylation of Smc3 by Esco1 and is accompanied by
cohesin's association with CTCF sites (Parelho et al., 2008; Wendt et al., 2008).

Figure 1. *Overview of cohesin, its subunits and regulatory proteins. Top left: cohesin's*

trimeric Smc1/Smc3/Scc1 ring. The approximate HAWK binding sites on the kleisin Scc1 are

indicated. Top right: cohesin in the DNA-clamped and folded state (hybrid model combining

PDBs 6WG3, 6ZZ6 and 7OGT). Bottom: cohesin's HAWKs and regulatory proteins

(AlphaFold 2 [AF] predictions).

Figure 2. *Cohesin's trimeric ring (top left), represented throughout the rest of the figure as a*

triangle, associates with DNA loops when occupied by SA/Scc3 and Nipbl/Scc2. Nipbl/Scc2

stimulates ATP hydrolysis, leading to DNA translocation and loop extrusion. This is halted by

Nipbl's replacement by Pds5, creating a complex that can bind Wapl and detach from

chromatin, in a process that depends on a pair of lysine residues (yeast: K112 and K113;

humans: K105 and K106) on Smc3's ATPase head, whose acetylation by Eco1 (yeast) or

Esco1/2 in vertebrates blocks release. Acetylated complexes containing Pds5 either form stable

loops between CTCF sites or, during DNA replication, generate cohesion between sister DNAs,

whose maintenance (in animal and plant cells) requires Sororin (as well as continued

acetylation) to prevent Wapl-mediated release.

Post-replicative cells have three major types of cohesin complexes: those holding sisters
together (sister chromatid cohesion), those actively extruding loops, and those whose LE has
been arrested by CTCF and which hold together convergent CTCF sites at TAD boundaries.
Crucially, all three are susceptible to the action of a fourth regulatory subunit called Wapl
(Chan et al., 2012; Gandhi et al., 2006; Kueng et al., 2006), which induces cohesin's release
from chromatin at a certain frequency, albeit only when Nipbl/Scc2 has been replaced by Pds5
and when Smc3(K105) and K106 are unmodified (Fig. 2). The increased processivity of
cohesin's LE upon Wapl's artificial depletion induces formation of chromatid-like structures
during interphase (Tedeschi et al., 2013), a process that occurs naturally during meiosis (Klein
et al., 1999). Moreover, Wapl's downregulation during B cell development is essential for
recombination between DJ junctions and distal V genes (Hill et al., 2020), which requires
highly processive LE, mediated by cohesin to bring them together in the correct orientation so
that they can be joined by the RAG recombinase.

Because its maintenance does not require Scc2/Nipbl, sister chromatid cohesion is thought to
be a passive structure that does not require continual ATP hydrolysis. Once generated during
replication, cohesion must also be shielded from Wapl. This protection depends on Smc3
acetylation, and in animal and plant cells on cohesin's association during S phase with a protein
called Sororin (Nishiyama et al., 2010; Rankin et al., 2005). When cells enter M phase,
Sororin's phosphorylation by Cdk1 abrogates its cohesion protection and Wapl thereupon
destroys most cohesion along chromosome arms but not at centromeres, where PP2A
associated with Shugoshins counteracts Cdk1 (Liu et al., 2013). Wapl's destruction of all
cohesion along chromosome arms when cells are arrested in mitosis by spindle poisons is
responsible for the canonical image of mitotic chromosomes, with cohesion persisting solely
at centromeres.

Cohesion is thought to be mediated through the topological entrapment of sister DNAs by
individual tripartite Smc-kleisin rings (Haering et al., 2008; Srinivasan et al., 2018), either
through co-entrapment of sister DNAs inside a single ring, or alternatively through formation
of a Borromean ring that links sister DNAs (best thought of as infinitively large rings) and
cohesin rings in a topology that does not involve catenation of any one by the other (Collier et
al., 2020). In either case, ring opening through Scc1's cleavage by Separase destroys the
topological entanglement, thereby destroying cohesion and triggering the metaphase to
anaphase transition (Oliveira et al., 2010; Uhlmann et al., 2000). Unlike Separase, Wapl
removes cohesin and destroys cohesion by opening the cohesin ring in a manner that does not

involve proteolysis. Importantly, the finding that Wapl-mediated release is blocked by fusing

Smc3's C-terminus to Scc1's N-terminus suggests that release depends on dissociation of

Scc1's NTD from Smc3 (Beckouet et al., 2016; Chan et al., 2012), thereby creating a gate

through which DNAs may pass. If sisters are held together by their co-entrapment within a

cohesin ring, then the latter must also possess an entry gate that differs from the exit gate and

probably resides at the hinge interface (Collier and Nasmyth, 2022), because, unlike release,

fusion of Smc3 to Scc1 does not block cohesion establishment (Gruber et al., 2006; Srinivasan

et al., 2018). Less is known about the topology of cohesin's association with DNA during LE.

The observation that it is unaffected, at least in vitro, by fusing all three of the ring's interfaces

has led to the suggestion that LE does not require (single strand) passage of DNA through any

of the ring's interfaces (Davidson et al., 2019), but this does not exclude the possibility of

DNAs passing through cohesin rings in two directions (two strand passage).

X-ray crystallography and electron cryomicroscopy (cryo-EM) have revealed many details of

the topology of Smc1/Smc3 heterodimers (Fig. 1), how they dimerize via hinge domains

(Haering et al., 2002), how their coiled coils are folded around an elbow (Bürmann et al., 2019),

how their ATPase domains associate in the presence of ATP and bind the N- and C-terminal

domains of Scc1 (Gligoris et al., 2014; Haering et al., 2004), the structure of cohesin's HAWK

subunits (Hara et al., 2014; Kikuchi et al., 2016; Lee et al., 2016; Ouyang et al., 2016; Roig et

al., 2014) and how they bind DNA (Li et al., 2018), how Scc2/Nipbl clamps DNA on top of

engaged ATPase heads (Collier et al., 2020; Higashi et al., 2020; Shi et al., 2020), and lastly

the structure of Wapl (Chatterjee et al., 2013; Ouyang et al., 2013). Despite this progress, many

molecular aspects of cohesin's activities remain mysterious. These include how cohesin

actually extrudes DNA loops, how Scc2/Nipbl and Pds5 exchange with each other, how

cohesin's exit and entry gates are opened, how Wapl facilitates exit gate opening and thereby

induces cohesin's dissociation from chromosomes, how Esco/Eco acetyl transferases are

recruited to cohesin, how they recognize Smc3, and how this is facilitated by Pds5 and the

Mcm helicase, and how Sororin hinders exit gate opening and thereby thwarts Wapl.

We report here a complementary approach to in vitro biochemistry and experimental structural

biology, namely in silico 3D protein structure prediction by AlphaFold 2 (AF) (Evans et al.,

2021; Jumper et al., 2021), based on amino acid co-evolution analyses and deep learning

approaches to sequence/structure relationships, to predict structurally hitherto unreported

interactions between cohesin subunits. When performed with subunits from multiple

eukaryotic lineages and critically judged in the light of amino acid conservation (beyond that

used by AF), these predictions have furnished insights into several of the outstanding questions

posed above. It has revealed a possible role in opening cohesin's exit gate of interactions

between the Smc1 and Smc3 coiled coils in the vicinity of their joints when ATPases engage

in the absence of Scc2/Nipbl. AF also predicts how Eco/Esco acetyl transferases are recruited

to cohesin and thereby bind to K112 and K113 (human: K105 and K106) within Smc3's

ATPase domain. Crucially, it predicts that Wapl induces formation of a tripartite complex

between itself, SA/Scc3, and Pds5. If this occurs after Scc1's NTD has already dissociated

from Smc3, then the manner in which Scc1 associates with Pds5 places the kleisin's NTD close

to a highly conserved cleft within Wapl's CTD, raising the possibility that the latter acts by

sequestering Scc1's N-terminal 20 amino acids. Importantly, the tripartite SA/Wapl/Pds5

complex is predicted to form not only in animal but also in plant and fungal lineages, suggesting

it has deep evolutionary origins. Likewise, the prediction that Sororin's essential C-terminal

domain binds to the interface between Smc3 and Scc1's NTD provides a mechanism by which

it protects the latter from Wapl.

RESULTS

Closing the Smc1 and Smc3 coiled coil joints plays a part in the dissociation of Scc1's

NTD from Smc3's neck.

Wapl is thought to induce cohesin's release from chromosomes by facilitating the passage of

DNAs through the Smc3/Scc1 interface. It could do this either by inducing dissociation of the

interface, by holding the gate open once Scc1 has already dissociated, or by a combination of

both mechanisms. If it merely held the gate open, then there must exist a mechanism capable

of opening it that is Wapl-independent. Evidence for this comes from the observation that

inactivation of Scc2 during late G1 triggers cohesin's dissociation from chromosomes via a

mechanism that is blocked by fusion of Smc3 to Scc1 or by mutations such as *SMC1(L1129V)*

or *SMC1(D1164E)* that suppress *eco1Δ* lethality (Srinivasan et al., 2019). Crucially, this

release is independent of Wapl.

A clue as to a mechanism for Wapl-independent release comes from observations on the related

condensin complex. In the absence of DNA, ATP induces dissociation from Smc2 (equivalent

to Smc3) of Brn1's NTD (equivalent to Scc1). Because it also occurs in the presence of ADP

and BeF$_3$, Brn1's dissociation is presumably induced by the engagement of Smc2 and Smc4

ATPase heads (Hassler et al., 2019; Lee et al., 2022). This raises two related questions. Does a

similar process occur in cohesin and if so, how might head engagement induce dissociation?

An important clue as to the possible mechanism stems from cryo-EM structures of both cohesin

and condensin in the presence of DNA and non-hydrolyzable ATP analogues, which reveal

that DNA is "clamped" on top of engaged ATPase heads by Scc2 and Ycs4 respectively.

Importantly, in both cases, Smc coiled coils are splayed open and the N-terminal domains of

Brn1 and Scc1 remain associated with the necks of Smc2 and Smc3 respectively (Collier et al.,

2020; Lee et al., 2022; Shaltiel et al., 2022). This raises the question as to why DNA's presence

leads to retention of the Smc-kleisin interaction. Is it due to DNA binding per se or to an event

that accompanies that binding, namely opening of the Smc coiled coils. Clearly, head

engagement does not per se induce dissociation, at least when DNA is bound on top of them,

as in the cryo-EM clamped structures.

To address these questions, we explored the potential of structural predictions using AlphaFold

2 multimer (AF) (Evans et al., 2021; Jumper et al., 2021). Importantly, presumably due to the

co-evolution of amino acids within the interface of engaged Smc ATPase heads, AF routinely

attempts to engage them even though ATP is absent, which means that it is possible, to some

extent, to observe the effects of head engagement on the structure of cohesin's coiled coils

using AF (Fig. 3A, f1). As with all further figures, the data files (f1- f98) can be found at

https://doi.org/10.6084/m9.figshare.22567318.v1. To address whether head engagement is

compatible with association of the N- and C-terminal domains of Scc1 with Smc3 and Smc1

ATPase heads respectively, we asked AF to predict the structures of Smc1 and Smc3 heads

associated with coiled coils extending up to but not including their joints in the presence of a

truncated Scc1 polypeptide in which its conserved NTD and CTDs were joined by a GS

(glycine, serine) linker (Fig. 3B, f2). AF predicted with high confidence in five out of five

models heads whose engagement was very similar to that of the clamped cryo-EM structures

(Fig. 3C), as was the orientation of their coiled coils. Importantly, the N- and C-terminal

domains of Scc1 were invariably bound to the neck of Smc3 and the base of Smc1 respectively,

as observed in the clamped structures. As shown by cryo-EM and predicted by AF, head

engagement is therefore not per se incompatible with Scc1's association with Smc3.

Figure 3. A, *AlphaFold 2 multimer (AF) prediction of yeast Smc1 and Smc3 ATPase heads*

with associated coiled coils up to, but not including their joints. The actual predictions and

sequences used are available in folder f1 at https://doi.org/10.6084/m9.figshare.22567318.v1.

***B**, As **A**, but including Scc1's N- and CTD connected by a GS linker (f2). **C**, Cryo-EM structure*

of DNA clamped on top of engaged Smc1 and Smc3 heads by Scc2 (PDB 6ZZ6).

A very different arrangement of Smc1 and Smc3 coiled coils has been observed by cryo-EM

in the absence of ATP. The coiled coils are juxtaposed from their emergence from the hinge

all the way to their joints close to the ATPase domains, which is accompanied by folding of

both coiled coils around their elbows, thereby locating the hinge domain close to a segment of

Smc3's coiled coil just above its joint (Petela et al., 2021) (PDB 7OGT). To address whether

AF can reproduce this state, it was used to predict the structure of a yeast Smc1/Smc3

heterodimer containing all hinge and coiled coil sequences, but lacking both ATPase domains

(Fig. 4A, f3). This revealed, with high confidence in five out of five models, a structure

resembling that seen by cryo-EM (Fig. 4C), namely, with the Smc1 and Smc3 coiled coils

juxtaposed in parallel throughout their length and folded around their elbows placing the hinge

asymmetrically on top of the Smc3's coiled coil, just above its joint. Noticeably, in the AF

prediction, the latter was intimately inter-digitated with Smc1's joint, creating a compact

junction between the two (Fig. 4A). AF predicted an almost identical structure for the

equivalent parts of human Smc1/Smc3 heterodimers (Fig. 4B, f4). Importantly, the asymmetric

manner in which the Smc1 and Smc3 coiled coils emerge from the hinge and thereby associate

with each other, as soon as they do so, resembles that seen by cryo-EM as well as a crystal

structure of the condensin hinge associated with stretches of coiled coil (Lee et al., 2020; Soh

et al., 2015) (PDB 4RSI, 6YVU). This relationship between the hinge and its associated coils

has the appearance of a wrist, folded back on its associated arm. Interestingly, according to AF,

the folded wrist conformation is not predicted when the coiled coils associated with the hinge

were cut short at the elbow (Fig. 5A, f5). The shortened Smc1 and Smc3 coiled coils instead

emerge symmetrically with opened arms at an angle of 120° from each other, in a manner

resembling the crystal structure of the *T. maritima* hinge associated with short coiled coils (Fig.

5B) (Haering et al., 2002) (PDB 1GXL). This suggests that folding at the elbow as well as the

consequent packing of the hinge on top of the Smc3 coiled coil may have an important role in

organizing how the coiled coils emerge from the hinge. Crucially, the association of full-length

elbow-containing coiled coils and especially their joint junction depends on the hinge, as Smc1

and Smc3 coiled coils in which the hinge is replaced by either a GSx4 (Fig. 6A, f6) or GSx15

(Fig. 6B, f7) linker fail to associate, even at their joints. Though the elbow appears to be a

feature that is autonomous, at least according to AF, coiled coil association, including joint

junction, is affected by a distant "organizing locus", namely the hinge in its wrist conformation.

***Figure 4. A**, AF prediction for yeast Smc1 and Smc3 lacking their ATPase domains (f3).* ***B**, As*

A, but human Smc1 and Smc3 (f4). ***C**, Cryo-EM structure of yeast Smc1 and Smc3 hinge and*

associated coiled coils, PDB 7OGT.

Figure 5. **A**, *AF prediction for yeast Smc1 and Smc3 hinge domains with short associated coiled coils (f5).* **B**, *Crystal structure of T. maritima hinge associated with short coiled coils, PDB 1GXL.*

Figure 6. *AF predictions for yeast Smc1 and Smc3 coiled coils including their elbow and joints, but joined by GSx4 (**A**, f6) and GSx15 (**B**, f7) linkers instead of their hinges.*

Though influenced by elbow and hinge, the joints nevertheless have an affinity for each other, as they form compact junctions when the coiled coils distal to the elbow are excluded and do so despite the lack of any extensive association between coiled coils away from the joint (Fig. 7A, f8). Indeed, AF predicted with high confidence a compact heterodimer formed between the joints of Smc1 and Smc3, associated with only short sections of coiled coils (Fig. 7B, f9). Though AF predicted heterodimers for isolated Smc1 and Smc3 joints from yeast, it failed to do so for joints from humans (Fig. 7C, f10), despite the fact that their joints are also joined when part of hinge-associated coiled coils. A comparison of the yeast and human joints revealed the probable reason for this difference, namely that the human Smc1 joint contains an alpha helical insertion that hinders the intimate joint inter-digitation observed in yeast.

Consistent with this theory, human Smc3 joints interdigitate with yeast Smc1 (Fig. 7D, f11)

but not vice versa. Though it hinders autonomous joint junction, the human Smc1 joint can

nevertheless be induced to interdigitate with its Smc3 partner when organized by intact folded

coiled coils organized by their hinge (Fig. 4B). Though joint junction may be a universal feature

of Smc1/Smc3 heterodimers, the propensity to form this structure may vary between species

and such differences that may be of physiological importance. The differences seen may also

reflect the varying degrees of prediction accuracy achieved by AF with the joint sequences.

Figure 7. *AF predictions of Smc1 and Smc3 joints with short associated coiled coils.* ***A*** *(f8)*

and ***B*** *(f9), yeast.* ***C*** *(f10), human.* ***D*** *(f11), human Smc3 and yeast Smc1 chimera.*

It is important to note that the junction between Smc1 and Smc3 joints was fully compatible

with the simultaneous association of Scc2 with the Smc3 joint. This was demonstrated by

running an AF prediction of Scc2's NTD either with an isolated Smc3 joint segment or with

an Smc1/Smc3 joint dimer (Fig. 8A, f12). Aligning the Smc3 moiety of the AF prediction with

the equivalent residues in the cryo-EM structure of DNAs clamped by Scc2 on top of engaged

Smc1/3 heads (Fig. 8B) (PDB 6ZZ6) shows the prediction to recapitulate the same Scc2:Smc3

binding mode. Because of this compatibility, competition with Scc2 is unlikely therefore to

have a role in joint disjunction (and thereby retention of Scc1's association with Smc3) upon

clamping.

***Figure 8. A**, AF prediction for yeast Smc1 and Smc3 joints with associated coiled coils together*

with Scc1 and Scc2's NTDs (f12). ***B**, Cryo-EM structure of DNA clamped by Scc2 on top of*

engaged Smc1 and Smc3 ATPase, PDB 6ZZ6.

These predictions imply that dimerization at opposite ends of the Smc1 and Smc3 coiled coils

have opposing influences on their organization. Engagement of the ATPase heads opens their

associated coiled coils, while hinge dimerization has the opposite effect, forcing them together,

including junction of their joints proximal to the ATPase heads. What happens then when both

types of dimerization co-exist? Due to their size, AF predictions of full length Smc1/Smc3

heterodimers are close to limits of what is possible for current implementations of AlphaFold

2 multimer (Evans et al., 2021). Despite this, we obtained four models for yeast and five for

human heterodimers. Four out of four models for the yeast complex predicted heterodimers

whose coiled coils were folded and their joints interdigitated despite heads being fully engaged

(Fig. 9A and B, f13). In the human case, three out of five models (ranks 1, 2, and 5) had a

similar configuration but in one model (rank 3) joint junction was dominant over head

engagement (and the latter were disengaged), while in another head engagement was dominant

over joint junction and the latter were disjoined (Fig. 9C, f14). As one might expect,

coexistence of joint junction and head engagement is accompanied by distortions of the coiled

coils connecting joints and heads, and in some predictions this is associated with extreme

bending and/or actual unfolding of the alpha helices. The more extreme coiled coil distortions

observed in some models of the full length Smc1/Smc3 heterodimers models are absent when

the hinge and distal coiled coils are removed from the yeast complex. Thus, four out of five

(ranks 1, 2, 3, and 4) models of Smc1/3 ATPase heads associated with coiled coils extending

to but not beyond the elbow predict with high confidence junction of their joints co-existing

with fully engaged heads (Fig. 10A, f15).

Figure 9. *AF predictions for full length yeast (**A** and **B**, f13) and human (**C**, f14) Smc1/Smc3 heterodimers.*

Though some of the more baroque AF models may reflect unrealistic scenarios, the emerging

principle that head engagement and joint junction can co-exist but only do so if the intervening

coiled coils are distorted is consistent with a similar configuration in low resolution cryo-EM

structures of Smc1/Smc3 heterodimers in the presence of ATP (Petela et al., 2021). Indeed,

there is evidence that the state also occurs in vivo. By introducing cysteine pairs in the interface

between engaged Smc1/3 ATPases, it is possible to use the bifunctional thiol-specific

crosslinking reagent BMOE (bis-maleimidoethane) to detect their proximity in vivo. A

different pair of cysteines can be used to detect association between Smc1 and Smc3 coiled

coils in the vicinity of their joints. In an observation surprising at the time, it was observed that

BMOE induces appreciable double crosslinking (head and joint crosslinks) with a frequency

not dissimilar to the product of the frequency of the individual pairs, which indicates

simultaneous crosslinking (Chapard et al., 2019).

These observations lead us to propose that head engagement under conditions in which the

hinge reinforces a junction of Smc1/Smc3 joints, namely in the absence of DNA or Scc2, will

create distortions of the coiled coil (Smc3's neck) to which Scc1's NTD is normally associated

and that this distortion may cause its dissociation. A corollary is that maintaining the

association of Scc1's NTD with the Smc3 neck, should prevent either joint junction or even

head engagement. Consistent with this hypothesis, the addition of Scc1's NTD (Fig. 10B, f16)

but not its CTD (Fig. 10C, f17) to the Smc1 and Smc3 ATPase heads associated with coiled

coils extending to the elbow eliminated the joint junction previously observed in four out of

five models. In this case, where joint junction is not reinforced by the hinge, AF clearly gave

preference to the Smc3-kleisin interaction over joint junction. Likewise, the top three ranked

models obtained with Scc1's NTD together with full length Smc1/Smc3 heterodimers

predicted engaged heads with Scc1's NTD associated with Smc3's neck but joints fully

disjoined (f18). Though the in silico prediction that association between Scc1's NTD and the

Smc3 neck is dominant over joint junction may not be an accurate reflection of nature, the

notion that joint junction while heads are engaged may be the driving force for opening

cohesin's exit gate nevertheless provides a plausible mechanism for this key event.

Importantly, the hypothesis explains why non-lethal mutations that compromise head

engagement (Elbatsh et al., 2016) abrogate releasing activity (RA). Crucially, it is eminently

testable, as it predicts that dissociation of Scc1's NTD from the Smc3 neck depends on joints

and probably also on distal coiled coils and their associated hinge.

Figure 10. *AF predictions of yeast Smc1 and Smc3 ATPases with coiled coils extending to the elbow but not beyond, without Scc1 (**A**, f15), with Scc1's NTD (**B**, f16) and with Scc1's CTD (**C**, f17).*

How does Wapl enhance exit gate opening?

If ATP-dependent dissociation of Scc1's NTD from the Smc3 neck is a feature intrinsic to Smc-kleisin trimers, then how might Wapl enhance this process? To address this, we used AF to explore how Wapl is first recruited to cohesin and second what effect it has on the configuration of cohesin's HAWK regulatory proteins and /or on the Smc3/Scc1 interface. Our starting point was existing knowledge based on biochemical and genetic assays. In vitro, Wapl is known to bind SA/Scc3 and this binding is abrogated by mutation of a highly conserved lysine residue (K404 in yeast; K330 in humans) within a highly conserved surface (known as the "CES", Conserved and Essential Surface) that sits underneath the pronounced "nose" of Scc3/SA. This binding must be of physiological importance as K to E mutations abrogate Wapl-dependent releasing activity in vivo, both in yeast (Beckouet et al., 2016) and in cultured human cells (Hara et al., 2014). Exactly which part of Wapl binds to the CES has never been fully established. It has been proposed that sequences within Wapl's unstructured N-terminal domain are responsible, either one or other of its FGF motifs (Shintomi and Hirano, 2009) or a different segment between residues 510 and 570, known as Wapl-M (Hara et al., 2014; Ouyang et al., 2013). As will become clear below, AF predictions imply that neither proposal

is correct. Meanwhile, a short YSR motif at Wapl's N-terminus binds a highly conserved loop

(consensus sequence AP[D/E]AP) between helices 2 and 3 within Pds5's NTD, which is also

bound by a YSR-like motif within the kinase Haspin (Ouyang et al., 2016). This locus within

Pds5's NTD was in fact first identified as being crucial for Wapl-mediated release through

mutations clustering at this location that suppress the lethality of *eco1* mutants in yeast

(Rowland et al., 2009). However, Wapl's N-terminal YSR has never been rigorously

demonstrated to be necessary for its releasing activity (RA). Most fungal Wapls also possess

at or near their N-termini a similar motif whose consensus is [M/Φ]xxYG[K/R]. Importantly,

its deletion in *S. cerevisiae* does not suppress the lethality of *eco1Δ* mutants (Fig. 11A),

implying that unlike Pds5's AP[D/E]AP loop to which it is thought to bind, the

[M/Φ]xxYG[K/R] motif is not essential for Wapl-mediated releasing activity. Some other

sequence must therefore bind the AP[D/E]AP loop in vivo. Interestingly, the

[M/Φ]xxYG[K/R] motif is lacking in *S. pombe*, though a different one (TYxxxR[T/S]ΦL),

which is much closer to Wapl's conserved CTD, is present and highly conserved in that

location among all fungi (Fig. 11B). Unlike the N-terminal motif, the more C-terminal

TYxxxR[T/S]ΦL motif is essential for releasing activity, as its deletion supresses the lethality

of *eco1Δ* mutants in *S. cerevisiae* (Fig. 11A). Thus, the picture that emerges is primarily one

of a loose association between Wapl and SA/Scc3 or Pds5 via short motifs within Wapl's

unstructured N-terminal domain. Crucially, none of these interactions help explain how Wapl's

more conserved CTD composed of eight HEAT repeats actually promotes release.

Figure 11. **A**, *Deletion of Wapl's N-terminal [M/Φ]xxYG[K/R] motif from S. cerevisiae (waplΔ4-9) does not suppress the lethality of eco1 mutations while deletion of its more C-terminal TYxxxR[T/S]ΦL motif (waplΔ208-215) does so. Both deletions were initially created within a version of the WAPL gene tagged with EGFP (KN18714) and their ability to suppress the lethality of eco1Δ determined by crossing to a waplΔ eco1Δ strain (strain KN16432) followed by tetrad analysis.* **B**, *A multiple amino acid sequence alignment of fungal Wapl sequences, showing the TYxxxR[T/S]ΦL motif on the left and the first highly conserved alpha helix within Wapl's CTD on the right. Yeast Wapl is the top sequence (W7PW2).*

We first used AF to explore the association between Wapl and Pds5. For yeast, this predicted

a single point of contact between a short motif ([F/W]xFLD), which is conserved in numerous

ascomycetes including *S. cerevisiae* (FNFLD- see Fig. 11A) and *S. pombe* (WNFLD), and a

trough situated between a pair of HEAT repeats located within Pds5's C-terminal half, at the

point where its longitudinal axis starts to curve (Fig. 12A, f19). Because the very same trough

is predicted also to bind FGF motifs from human Wapl (Fig. 13B, f20) and Sororin (Fig. 13B,

f21) as well as a separate motif within Esco1 (Fig.13B, f22), we will refer to it as the Wapl

Esco Sororin Trough or "WEST". Intriguingly, AF models of individual proteins in the

AlphaFold Protein Structure Database (Varadi et al., 2022) predict that the *S. cerevisiae* and *S.*

pombe FNFLD motifs of Wapl also bind a highly conserved cleft within Wapl itself, at the C-

terminal end of its CTD (Fig. 12B). However, given the choice between binding Pds5 and

itself, AF chooses the former. In humans, as already alluded to, AF predicted in five out of five

models an interaction between the central of three FGF motifs within Wapl's N-terminal

extension and the equivalent C-terminal trough within Pds5A and B (Fig. 13B, f20). Various

other sequences within this unstructured part of Wapl were predicted to bind Pds5's N-terminal

APDAP loop, including a TKWGE motif (2/5 models) (Fig. 14A, f20) and in only one out five

models and with low confidence also its YSR motif (Fig. 14A, f20). In the case of human (Fig.

14B, f23), but not yeast Wapl (Fig. 12A, f19), AF predicted with high confidence in five out

of five models a rigid body association between its HEAT-repeat containing CTD and HEAT

repeats at Pds5's N-terminus, close to but independent of Pds5's APDAP loop, an interaction

that proved pivotal for our subsequent analyses (see below).

Though AF did not predict interaction of the *S. cerevisiae* TYxxxR[T/S]ΦL motif with full

length Pds5 (f19A), it predicted, in five out of five models, interaction of the equivalent motif

from *Neurospora crassa* with Pds5's APDAP loop (Fig. 12C, f19B). Subsequent predictions

using truncated versions of Pds5 and Wapl revealed that the *S. cerevisiae* TYxxxR[T/S]ΦL

motif also reproducibly bound to this site (f19C), as did (using a different Wapl truncation) its

[M/Φ]xxYG[K/R] motif (f19D), in this case in a manner resembling a crystal structure of the

equivalent complex from humans (PDB: 5HDT) (Ouyang et al., 2016). Given that both Pds5's

APDAP loop and Wapl's TYxxxR[T/S]ΦL motif are essential for releasing activity in yeast,

we presume that the interaction predicted by AF is crucial for their function, a feature whose

importance will become apparent later. It is reasonable to suppose that the Pds5 APDAP loop

also binds Wapl's N-terminal [M/Φ]xxYG[K/R] motif, albeit not at the stage when the

releasing reaction requires the APDAP loop's interaction with the TYxxxR[T/S]ΦL motif.

Figure 12. *A, AF prediction for yeast Wapl's association with Pds5 (f19A).* ***B****, AF prediction*
of yeast Wapl alone (AF-Q99359-F1, AlphaFold Protein Structure Database). ***C****, AF predicts*
interaction between Wapl's TYxxxR[T/S]ΦL motif and Pds5's APDAP loop in Neurospora
crassa (f19B).

Figure 13. Interaction between a trough in Pds5's CTD called "WEST" and an FNFLD motif

*within the N-terminal part of yeast Wapl (**A left and right**, f19), FGF motifs within human*

*Wapl (**B upper left and right**, f20) and D. rerio Sororin (**B lower right**, f21), and a highly*

*conserved, partly helical, ILxLCEEIAGEIESD motif within human Esco1 (**B lower left**, f22).*

Figure 14. *AF predictions for the interaction between human Pds5B and Wapl's unstructured N-terminal extension (**A**, f20), and between Pds5A and the N-terminal half of Wapl's highly conserved CTD (**B**, f23). Both types of interaction were observed with both Pds5A and B.*

In contrast to Pds5, AF did predict with high confidence in five out of five models a rigid body contact between a complex between yeast Scc3 and Scc1, and the first (α1) and second (α2) α helices within Wapl's CTD (Fig. 15A, f24), which are highly conserved (Fig. 11). The first of these binds across Scc3's HEAT repeats, including the helices at the base of its nose and those associated with the CES. Though the Wapl helix runs parallel to, and contacts, the Scc1 polypeptide, exclusion of the latter made little or no difference to AF's predictions. Since the axis of Wapl's first helix is nearly orthogonal to those of Scc3's HEAT repeats, so too are most of the HEAT repeat helices within Wapl's CTD and these therefore rise above Scc3 like a

"tower" that follows the line of Scc3's nose, albeit leaning away from the latter's axis by 30°.

Towards the top of this tower is a hydrophobic cleft (Chatterjee et al., 2013), lined by the most

conserved α helix within fungal Wapls. This is the same cleft that in the absence of other

proteins is occupied by Wapl's own N-terminal F/WxFLD motif (Fig. 12B), not only in fungi

but also in a wide variety of eukaryotes (see below).

A very similar if not identical interaction was predicted between the human SA2:Scc1 complex

and a fragment of human Wapl containing its entire N-terminal extension as well as half of its

CTD (a prediction with full length SA2 and human Wapl would have been close or over AF's

size limit) (Fig. 15B, f25). In addition, in two out five models (ranked 1 and 3) a long motif

(IFSGPKRSPTKAVYNARHWN) within Wapl's unstructured N-terminal extension

associated with a quite separate segment of SA2, in this case binding parallel to one of its

central HEAT repeats (Fig. 15B). Contact with the SA2:Scc1 complex is extensive, running

along the entire length of the relevant SA2 HEAT repeat, crossing and in so doing, contacting

the Scc1 polypeptide and finally interacting via its conserved tryptophan residue C-terminal

HEAT repeats on the other side of SA2's hook. This part of Wapl corresponds to the Wapl-M

sequences (510-570) important for its association with SA2:Scc1 and previously thought to

bind the CES (Hara et al., 2014; Ouyang et al., 2013). AF predicts a very different interaction,

and besides which, it suggests that the CES is occupied instead by Wapl's CTD (Fig. 15B).

The AVYNARHWN part of the motif is conserved in animals (consensus

A[V/L]Yxx[R/K][H/-]W) and its mode of binding may explain why Wapl binds to Scc1/Rad21

alone in vitro (Shintomi and Hirano, 2009). Significantly, AF failed to predict any interaction

between SA2:Scc1 and Wapl's FGF motifs. However, because the tryptophan-containing

SA2:Scc1-binding motif is separated from Wapl's middle FGF motif by a long polypeptide,

there is no reason why the former and latter could not be bound simultaneously to SA2:Scc1and

Pds5's WEST respectively.

The first alpha helix of Wapl's CTD (α1) is highly conserved (Fig. 17A, right), especially, but

not exclusively (for reasons that will become apparent) residues facing SA2. Crucially, its

interaction with SA/Scc3 appears to be of physiological importance. Scc3(K404) likely forms

a conserved salt bridge with yeast Wapl's D272, while the equivalent residue in SA2, K330

does likewise with human Wapl's D657. Remarkably, mutation of all four residues

(individually) has been shown to either abrogate or to greatly reduce RA mediated by Wapl

(Beckouet et al., 2016; Ouyang et al., 2013; Rowland et al., 2009). *S. cerevisiae* Wapl(D272G)

was in fact isolated as a spontaneous suppressor of *eco1* lethality in yeast. The AF prediction

for the human proteins explains the greater conservation of the surface of Wapl's α1 facing

SA2 as well as the more N-terminal V639, which also contacts SA2. Crucially, the AF model

explains why individual mutation of E653, D656, D657, or Y660 within the CTD's α1 helix

abrogates RA, as does mutation of L676 or K684 in its α2 helix (Ouyang et al., 2013). All

these residues are predicted by AF to contact SA2 (Fig. 15B, residues marked in yellow).

***Figure 15. A**, AF prediction for the interaction between Scc3/Scc1 and the CTD of yeast Wapl*

*(f24). **B**, AF prediction for the interaction between SA2/Scc1 and a fragment of human Wapl*

containing its N-terminal sequences as well as the N-terminal half of its helical CTD (f25).

Mutations that abolish RA are marked in yellow: E653, D656, D657, L676 and K684.

The prediction that in humans Wapl's CTD forms a complex with Pds5 as well as with

SA2:Scc1 raises the possibility that both interactions could occur simultaneously and that Wapl

would therefore glue all three proteins together in a ternary complex. This seems indeed to be

the case, as AF failed to predict any direct interaction between full length SA1 and Pds5A

(f26), but did so with high confidence in the top two ranked models created in the presence of

the part of Wapl's CTD that it predicts to bind both SA and Pds5 (Fig. 16, f27). The prediction

statistics indicated that the most accurate images of the ternary complex were obtained when

only the N-terminal domains of SA1 and Pds5A were included. In this case, AF predicted a

ternary complex with high confidence in five out of five models (Fig. 17A, f28). Using this

template, we screened for the formation of ternary SA:Wapl:Pds5 complexes in a variety of

eukaryotic lineages. Identical ternary complexes were predicted with high confidence in plants

(*Arabidopsis thaliana*), in insects (*Drosophila melanogaster*) and in a basal fungus, namely

the chytrid *Spizellomyces punctatus* (Fig. 17B, f29, f30, and f31). Crucially, formation of the

ternary complex explains the conservation and functional importance of E647 (Ouyang et al.,

2013) within human Wapl's α1 helix despite not facing SA2, as it contacts Pds5's NTD.

Likewise, contact with Pds5's NTD in the AF model explains abrogation of Wapl-mediated

RA by R695E as well as conservation of M692, R695, R731, and L732 in animals (Figure 17A,

right). We will refer to this as the canonical Pds5:SA/Scc3:Wapl rigid body ternary complex.

Figure 16. *AF prediction of a complex between human Pds5A, SA1, and a fragment of Wapl's CTD (f27). Pds5 and SA1 were not predicted to interact in the absence of Wapl (f26).*

Figure 17. *AF predictions for complexes between N-terminal fragments of Pds5 and SA/Scc3*

*and the N-terminal half of Wapl's CTD for humans (**A**, f28), A. thaliana (**B**, f29) D.*

melanogaster (f30), and S. punctatus (f31). Also shown is Wapl's conservation mapped on the

*human complex, **A**, right.*

In contrast, despite the robust formation of binary complexes between Scc3 and Wapl

orthologs, AF failed to predict the same type of ternary complexes with the equivalent protein

fragments from *S. cerevisiae* (f32A). Further analysis of the way Wapl interacts with Pds5 in

S. punctatus revealed a possible explanation. When the former's CTD was attached to N-

terminal residues containing its TYxxxR[T/S]ΦL motif, AF predicted that Pds5's NTD formed

a rigid body complex with the CTD, whose conformation was identical to that observed in the

canonical Pds5:SA/Scc3:Wapl rigid body ternary complex (Fig. 17B), while its APDAP loop

simultaneously bound Wapl's TYxxxR[T/S]ΦL motif (Fig. 18, f19E), in the manner observed

in *S. cerevisiae* (f19C) and *N. crassa* (f19B). This raised the possibility that an interaction

between Wapl's TYxxxR[T/S]ΦL motif and Pds5's APDAP loop might actually facilitate

formation of ternary complexes and that this might explain why the TYxxxR[T/S]ΦL motif is

essential for releasing activity in *S. cerevisiae*. We therefore repeated the AF predictions for

the previously described *S. cerevisiae* Pds5, Scc3, and Wapl fragments (f32A) but in this case

included sequences containing Wapl's TYxxxR[T/S]ΦL motif. Remarkably, the top three

ranked models now predicted with high confidence a rigid body ternary complex (Fig.19A,

f32B), whose conformation closely resembles the canonical Pds5:SA/Scc3:Wapl rigid body

ternary complex observed in animals, plants, and basal fungi. Moreover, this was accompanied

by interaction between the TYxxxR[T/S]ΦL motif and Pds5's APDAP loop, as observed in *S.*

punctatus. The dramatic effect on including the TYxxxR[T/S]ΦL motif is best visualized by

comparing the PAEs of the top three ranked models with and without the motif (Fig. 19B).

AF also predicted with high confidence in two out of five models a very similar type of

canonical ternary complex in *N. crassa* (f33A) and the basidiomycete *Ustilago Maydis* (f33B).

As in *S. cerevisiae*, formation of the canonical Pds5:Scc3:Wapl ternary complex in *Neurospora*

was also strictly dependent on inclusion of its TYxxxR[T/S]ΦL motif (compare f33A & f33C).

These observations not only suggest that formation of a canonical Pds5:SA/Scc3:Wapl rigid

body ternary complex is conserved in fungi as well as animals and plants, but also that its

formation may be an essential feature of releasing activity, because deleting the

TYxxxR[T/S]ΦL motif necessary for its in silico prediction in *S. cerevisiae* (f32) abolishes

releasing activity in vivo (Fig. 11B). It is remarkable that the TYxxxR[T/S]ΦL motif is not

only highly conserved among fungi but that it is connected to α1 in Wapl's CTD by a short

polypeptide linker whose length (25 to 50 residues) does not vary greatly. In other words, the

shortness of this tether may be important for ensuring the particular type of interaction between

Wapl's CTD and Pds5's NTD observed in canonical Pds5:SA/Scc3:Wapl rigid body ternary

complexes.

Figure 18. *AF prediction for the interaction between the N-terminal half of S. punctatus Pds5*

and a fragment of Wapl containing its TYxxxRT/SΦL motif connected to its CTD (f19E). Both

the motif and the CTD bind to Pds5

Fig. 19. A, AF prediction for a S. cerevisiae Pds5:Scc3:Wapl rigid body ternary complex accompanied by interaction between Wapl's TYxxxR[T/S]ΦL motif and Pds5's APDAP loop (f32B). B, PAEs for the top three ranked models for S. cerevisiae's Pds5:Scc3:Wapl ternary complex with (top row, f32B)) and without (bottom row, f32A) Wapl's TYxxxR[T/S]ΦL motif. Note that the lower confidence complexes predicted by AF for those lacking Wapl's TYxxxR[T/S]ΦL motif are not of the canonical variety.

Though the structure of the ternary Pds5:SA/Scc3:Wapl RA complex in animals (Fig. 20, f35),

plants, and fungi is clearly intriguing and presumably important for RA, it does not per se offer

insight into the actual mechanism by which these three proteins facilitate cohesin's release.

This is especially true for the conserved cleft at the top (C-terminal end) of Wapl's CTD.

Despite differences in the precise patterns of conservation in animals, plants, and fungi, it is

remarkable that in all three lineages, motifs within Wapl's unstructured N-terminal extension

are predicted by the AlphaFold structure database to bind this cleft (yeast shown in Fig. 12B).

As will become apparent later, AF also predicts that Sororin's conserved FGF motif (which is

related to the yeast Wapl's FNF) also binds the cleft. The fact that Sororin is an antagonist of

Wapl suggests that the binding by N-terminal Wapl sequences represents a form of auto-

inhibition.

Figure 20. *Composite model of the human Pds5:SA1:Wapl:Scc1 quarternary complex (f35)*

created using by superimposing rank 1 models from f23, f25, and f28.

Potential insight into the function of the highly conserved C-terminal Wapl cleft, which is

predicted to be bound by motifs within Wapl's N-terminal sequences was revealed by a

separate set of predictions, namely investigation of how Scc1 binds to Pds5, either in the

presence or absence of Smc3's neck. Functional studies in yeast combined with X-ray

crystallography of a Pds5:Scc1 complex from *Lanchea thermotolerans* revealed association of

an extended Scc1 peptide, from *Lt* S125 to V141, along the axis of Pds5's spine, half way down

from its N terminus (Lee et al., 2016). This mode of interaction is essential as mutations

affecting the interaction, namely Scc1(V137K) and Pds5(Y458K) abrogate binding of the two

proteins in vivo and are lethal (Lee et al., 2016). Pds5 is required to maintain cohesion in post-

replicative cells (Hartman et al., 2000; Panizza et al., 2000), when Scc1's NTD is bound to

Smc3's neck (Gligoris et al., 2014), as well as being involved in RA (Rowland et al., 2009),

when Scc1's NTD must dissociate from Smc3, at least transiently (Beckouet et al., 2016; Chan

et al., 2012). With the initial aim of revealing the orientation of Pds5 connected to the Smc3

neck via Scc1, we used AF to predict how the N-terminal half of Pds5 interacts with the N-

terminal 160 amino acids of Scc1 in the presence of Smc3's ATPase and neck. Not surprisingly,

all five models predicted Scc1's simultaneous association with both the Smc3 neck and with

Pds5, in both cases in the manner expected from crystallographic studies (PDB 5F0N & 4UX3)

(Fig. 21, f36). More unexpected were the predictions for the sequences that connected the Scc1

S125-V141 interval to the NTD bound to Smc3. This revealed that Scc1's association with

Pds5's spine extended, in an N-terminal direction on both proteins, from the previously

characterized site of interaction almost but not quite to the C-terminal end of the long α3 helix

that forms a three-helical bundle with Smc3's coiled coil (Fig. 21). In all five models, the short

peptide linking α3 with Scc1 associated with Pds5's spine adopts the same conformation,

which ensures that Pds5's NTD points towards the Smc3 ATPase head. Though we have

previously suggested that such an orientation might help to direct Eco1 to its K112 and K113

targets on the Smc3 ATPase (Lee et al., 2016), subsequent AF predictions (see below) imply

that this orientation would not in fact facilitate acetylation and the orientation presumably has

some other function. Given that the sequences linking Scc1$_N$'s α3 with its canonical Pds5

binding motif are poorly conserved between fungi and animals, it was remarkable that AF

predicted a very similar structure with the orthologous human proteins (Fig. 21, f37).

Figure 21. *AF predictions for the interaction of Scc1's N-terminal 160 residues with Smc3 and with an N-terminal fragment of Pds5 for both S. cerevisiae (f36) and human (f37).*

Because RA depends on Wapl's association with Pds5, Scc1$_N$ is presumably bound to Pds5 upon its dissociation from Smc3 during a release reaction. We therefore used AF to predict how Scc1$_N$ alone binds to Pds5. In the absence of Smc3, Smc3's coiled coil no longer poses a constraint on how Scc1$_N$ binds with Pds5, and AF predicts that yeast Scc1's association now extends to the C-terminal end of α3, whose interaction with the ends of a pair of Pds5's HEAT repeats leads to a pronounced rotation, as well as levering backwards of the Scc1 NTD (Fig. 22, f38). This change is not accompanied by any alteration of its actual structure or in the manner in which more C-terminal Scc1 sequences bind the Pds5 spine. In humans, α3's association with the Pds5 spine was less affected by Smc3's removal but it was nevertheless levered further backwards, becoming more orthogonal to the spine's axis (Fig. 22, f39). Crucially, in both cases, as well as in *A. thaliana* (Fig. 22, f40), the C-terminal end of Scc1$_N$'s α3 is anchored on the Pds5 spine and juts out from it at a defined angle. The change in the

manner of Scc1$_N$'s association with Pds5 upon dissociation from Smc3 might account for the

observation that Pds5 inhibits Scc1$_N$'s association with Smc3 in vitro, but does not affect its

association once bound (Ouyang et al., 2016). Given that Pds5 is intimately involved in RA

and that the latter requires dissociation of Scc1's NTD from Smc3, it is plausible to imagine

that upon its dissociation (presumably in response to head engagement and joint junction),

Scc1's NTD remains associated with Pds5 in the conformations observed in the above AF

models. Crucially, AF predicts that under these conditions, Scc1's NTD adopts a defined

position relative to Pds5's NTD that is similar in fungi, animals, and plants.

S. cerevisiae   Human   *A. thaliana*

Figure 22*. AF predictions for the interaction between Scc1's N-terminal 160 residues and*

Pds5 in S. cerevisiae (f38), humans (f39), and A. thaliana (f40).

Having realized that Pds5 organizes the spatial position of Scc1's NTD, we superimposed the

human Pds5:Scc1$_N$ model (Fig. 22 middle, f39) onto the canonical Pds5:SA/Scc3:Wapl rigid

body ternary complex (Fig. 23, f41). Remarkably, this revealed that the N-terminal end of

Scc1's NTD is placed, through its association with the spine of Pds5, in the vicinity of the

aforementioned cleft at the C-terminal end of Wapl's CTD. Supposition of the canonical

Pds5:Scc3:Wapl rigid body ternary complex from yeast (f32B) with AF predictions for Scc3's

association with Scc1 and full length Wapl (f24) and full length Pds5 associated with $Scc1_N$

(f38B) revealed a very similar looking quarternary complex (Fig. 31B).

As mentioned, in humans and most other metazoa, as well as in many fungi (Fig. 12B) and

plants, this cleft is predicted to be occupied by a motif (PSCLSVCNVT) within Wapl's N-

terminal sequences (AlphaFold Protein Structure Database: AF-Q7Z5K2-F1). The revelation

that Scc1's NTD is positioned close to this cleft in a putative Pds5:Sacc3:Wapl:Scc1 complex

suggested that the role of this site might be to sequester Scc1's NTD following its dissociation

from Smc3's neck. Testing this hypothesis was not possible using a full quarternary complex,

due to its size. However, predictions of Wapl's position relative to Pds5's NTD are in fact

similar whether or not Wapl's CTD is also bound to the SA subunit and we therefore used AF

to predict the structure of a complex containing Wapl's CTD, the first 150 residues of

$Scc1(Scc1_N)$, and the N-terminal straight section of Pds5 whose spine binds Scc1. All five

models predicted with high confidence a ternary complex between Wapl, Pds5, and $Scc1_N$ (Fig.

24, f42). In three of the five, including ranks 1, 2, and 5, Scc1's NTD was placed in close

vicinity of the Wapl cleft. The most N-terminal (α-1) of its four short helices (α-1, α0, α1, and

α2), normally packed on top of α3 N-terminal helix, dissociates from α3 and binds in the cleft

(f42). However, these peripheral associations between $Scc1_N$'s α-1 and Wapl were not

predicted with high confidence (f42), did not look convincing, and more seriously differed in

their details between models.

Figure 23. *A putative quartenary complex between human Scc1$_N$ Pds5, SA1, and Wapl, created*

by super-imposing f23, f25, f28, and f39. The Scc1's NTD points towards, but clashes with the

highly conserved cleft in Wapl's CTD that is otherwise predicted to bind a motif PSCLSVCNVT

(conserved among metazoa) located in Wapl's N-terminal sequences (see Fig. 26B).

Figure 24. *AF's predictions for ternary complexes (models ranks1 and 4) containing Scc1$_N$,*
Pds5's NTD, and Wapl's CTD (f42).

To test whether the positioning and hence orientation of Scc1's NTD through its association

with Pds5 hinders its association with Wapl's CTD, we asked how the latter interacts with

Scc1's NTD alone, i.e when not constrained in space through its association with Pds5 within

the rigid body ternary complex. In this case, AF predicts with high confidence and in four out

of five models (Fig. 25, f34A) that Scc1's N-terminal helix (α-1) binds the cleft in Wapl's CTD

that otherwise binds Wapl's own SCLSVCNVT peptide (Fig. 26B), while its second helix (α0),

lying at 90° to α-1, is sandwiched between two long antiparallel helices that form a pronounced

nose at the C-terminal end of Wapl and a second pair that cross over at 45°. AF predicts similar

interactions for *D. melanogaster* (f34B), *C. elegans* (f34C), and for *A. thaliana* (f34D).

Moreover, a slightly different but clearly related interaction was also predicted for *N. crassa*

(f34E). Remarkably, only modest changes in the conformation of Scc1's NTD are required to

accommodate binding of its α-1 and α0 helices to Wapl's CTD in this manner, principally a

slight bending of α3 to avoid a clash with the pair of Wapl CTD helices that cross its nose. As

a consequence, α3 lies on top of and almost parallel with Wapl's nose and this kind of

alignment is presumably why AF failed to predict Wapl's interaction with Scc1's α-1 and α0

when Scc1's NTD is connected to Pds5 when the latter is part of a ternary complex with Wapl

(Fig. 24, f42). In the latter case, Scc1's NTD approaches Wapl's CTD from an angle

incompatible with the stereo-specific engagement depicted in Figure 25. Additional AF

predictions confirmed that Scc1's first three mini-helices (α-1, α0, and α1) are sufficient for

this type of interaction not only for humans (Fig. 26A, f43) but also for *D. melanogaster* (Fig.

26C, f44), for *C. elegans* (Fig. 26D, f45), for the sponge *A. queenslandica* (Fig. 26E, f46), for

the green alga *P. salinarum* (Fig. 26F, f47), and for the vascular plant *A. thaliana* (Fig. 26G,

f48). Likewise, despite considerable sequence variation, AF predicted a very similar interaction

between Wapl and the equivalent helices of a fragment of human Rec8 containing its first 36

amino acids (Fig. 26H, f49).

Figure 25. A*, AlphaFold's prediction for the interaction of human Scc1's NTD with Wapl's*

CTD (f34A).

These results suggest that there may indeed be an intrinsic affinity between Wapl and Scc1's

α-1 and α0 helices but the interaction is not possible when the helices are packed onto α3 while

the latter is simultaneously connected to Pds5 bound to Wapl's CTD. If positioning Scc1's

NTD in the immediate vicinity of Wapl's CTD, through formation of a stereo-specific ternary

complex between Pds5, Wapl, and Scc1$_N$ is to promote sequestration of Scc1's α-1 α0 helices

by Wapl's CTD, then these helices would have to dissociate from α3. Crucially, AF would not

predict the unfolding necessary if it gave higher priority to packing of Scc1's N-terminal

helices with α3 than to partial unfolding and association with Wapl. That this appears to be the

case should not be surprising, given that Scc1's NTD spends most of its life associated with

Smc3 during which the α0 helix is packed onto top of α3. Even when not associated with

Smc3, Scc1 may not actually be engaged in a release reaction, which after all is probably a rare

event in the life of cohesin, only occurring once every 20 minutes in G1 cells (Gerlich et al.,

2006; Wutz et al., 2020).

There are two possible explanations for the conundrum that Scc1's NTD appears to approach

Wapl's CTD from the wrong direction: either the hypothesis that Wapl sequesters Scc1's NTD

is wrong or Wapl does indeed sequester Scc1's NTD when delivered precisely to the right

location by the Pds5 moiety of the Pds5:Scc3:Wapl:Scc1 quarternary complex, but to do so in

the manner depicted in Figure 26 requires a partial unfolding of Scc1$_N$, in particular

dissociation of Scc1$_N$'s α-1, α0 and α1 helices from α3, a feature that AF may not on its own

be able to predict. We therefore explored whether the packing of the three most N-terminal

helices around α3 could be weakened by changing key residues within α3, without altering the

latter's structure. After a series of trials testing the effects of α3 mutations on the structure

predictions of Scc1's NTD alone, we settled on a pentuple α3 mutant Scc1$_N$(H58A, L59S,

G62D, R65A, R69A) that, according to the PAE and pLDDT plots (Fig. 27, f50, f51) weakens

the association between Scc1$_N$'s three most N-terminal helices and α3, without greatly

affecting the folding of α3 itself. Importantly, AF's prediction for Scc1's interaction with the

Smc3 neck was also unaltered by the pentuple mutant, not only with regard to α3's interaction

with the Smc3 coiled coil, but also with regard to the packing on and around α3 of Scc1's more

N-terminal helices. The latter may be surprising given the extensive changes but AF predictions

are remarkable resilient to mutations and do not accurately reflect the underlying physico-

chemical changes. We then tested the effect of these mutations on AF's predictions for the

complex containing Wapl's CTD, the first 150 residues of Scc1, and the N-terminal straight

section of Pds5. Remarkably, the mutations led to AF predicting with high confidence and in

five out five models the unpacking of Scc1$_N$'s first two helices and their docking with Wapl's

CTD (Fig. 28, f52A) in a manner very similar if not identical to that observed when the first

two or three helices are run alone with Wapl's CTD (Fig. 26A) or in AF predictions for Wapl's

association with isolated Scc1 NTDs (Fig. 25, f34). AF even managed to predict a similar

outcome when the appropriate fragment of SA was included, thereby creating a quarternary

complex whose Scc1$_N$ is sequestered by Wapl (f52B). It is important to note that the mutations

did not affect any of the residues that actually contact Wapl in these models. Being in α3 itself,

which does not associate with Wapl, the mutations merely affect how the two N-terminal

helices associate with α3 (Fig. 27).

Figure 26. *AF predictions for the interactions between Wapl's CTD and the N-terminal 35*

*amino acids of Scc1 for humans (**A**, f43), D. melanogaster (**C**, f44), C. elegans (**D**, f45), the*

*sponge A. queenslandica (**E**, f46), the green alga P. salinarum (**F**, f47), and A. thaliana (**G**,*

*f48). **B**, AF's prediction for Human Wapl alone (AlphaFold Protein Structure Database AF-*

*Q7Z5K2-F1). **H**, AF prediction for the interaction between Wapl's CTD and the N-terminal*

*36 amino acids of human Rec8 (f49). **I**, interaction between Wapl's CTD and the N-terminal*

35 amino acids of Scc1 in a ternary complex containing Scc1$_N$(H58A, L59S, G62D, R65A,

R69A), Pds5's NTD, and Wapl's CTD (f52A). See Fig. 28 for the same structure in the complete

ternary complex.

Figure 27. ***A***, *PAEs for WT Scc1$_N$ (top, f50) and mutant Scc1$_N$(H58A, L59S, G62D, R65A,*

*R69A) (bottom, f51). **B**, pLDDT plots for WT Scc1$_N$ (left, f50) and mutant Scc1$_N$(H58A, L59S,*

G62D, R65A, R69A) (right, f51).

Figure 28. *AF's prediction of a ternary complex containing Scc1$_N$(H58A, L59S, G62D, R65A, R69A), Pds5's NTD, and Wapl's CTD (f52A). The location of α3 Scc1$_N$ mutations that distinguish it from f42 shown in Fig. 24 are marked in red.*

If sequestration of Scc1's N-terminal (α-1 & α0) helices by Wapl's CTD is of physiological importance, then the Wapl residues involved in sequestration should be conserved. This consideration cannot be applied to Scc1's NTD as it has multiple other functions. Highly conserved (amongst animals) human Wapl surface residues involved in the interaction include N939 and N990, D979 and E1117, which could interact with Scc1(R11), L983 (which could interact with Scc1[L8]), A1112 and H1115 (which could interact with Scc1[W23]), M1116 (whose sulphur atom could interact simultaneously with Wapl[F1165] and Scc1[W18]), and lastly M1166 (which could interact with Scc1[Y3]). Remarkably, several of these residues have been mutated and shown to abrogate Wapl's RA activity in vivo, namely Wapl(D979K), M1116A, E1117K, and F1165A (Ouyang et al., 2013). In conclusion, sequestration not only helps explain the pattern of conservation but also the physiological consequences of mutating key residues. Crucially, AF predicts that sequestration can also occur when Scc1$_N$ and Wapl's CTD are juxtaposed within a quarternary complex with SA and Pds5 (f52B). Because very

similar rigid body ternary complexes are formed between Pds5, Scc3, and Wapl in plants as

well as fungi (Fig. 23B), the N-terminal helices of their Scc1$_{N}$s will be presented to Wapl's

CTD in a similar manner to that observed in humans. It is therefore entirely plausible that the

proposed sequestration mechanism applies to cohesin releasing activity in all three kingdoms.

The cleft within Wapl's CTD putatively bound by Scc1$_{N}$'s α-1 (from animals, plants, and even

certain fungi such as the *Sordariale*s) has previously been highlighted by Chatterjee and co-

workers (Chatterjee et al., 2013), who described a crystal structure (PDB 3ZIL) in which the

equivalent cleft from the yeast *Ashbya gossypii* was occupied by a peptide from Smc3's

ATPase domain, a phenomenon that seemed implausible from a structural point of view and

has subsequently not been confirmed. *A. gossypii,* like *S .cerevisiae* belongs to the

Saccharomycetacae, a group of yeasts in which α-1 is missing, a feature unique amongst

eukaryotes, and replaced by a short string of non-conserved residues. In *Ashbya*'s case, AF

predicts that the cleft is bound by α0 instead of α-1 (f58), a feature that might also apply to *S.*

cerevisiae.

835   Yeast genetics implies that formation of the canonical Pds5:Scc3:Wapl ternary complex

836   involves multiple steps

Our AF predictions suggested that formation of canonical rigid body Pds5:Scc3:Wapl ternary

complexes in ascomycetes depends on a juxtaposition of Wapl's CTD and Pds5's NTD

facilitated by Wapl's TYxxxR[T/S]ΦL motif (Fig. 19B), which is essential for RA in vivo. It

is however curious that despite its clear effect on canonical ternary complex formation and

despite highly reproducible AF predictions for the interaction of this motif with Pds5's APDAP

loop using versions of Wapl that lack sequences N-terminal to the motif, AF fails to predict its

association with the APDAP loop of full-length yeast Pds5 when Wapl's entire unstructured

N-terminal sequences are present (f19A, f19D). A possible reason for this is that AF robustly

predicts that the FNFLD motif within yeast Wapl's N-terminal sequences binds to the WEST

locus situated at the opposite end of Pds5 and the limited distance between the FNFLD and

TYxxxR[T/S]ΦL motifs reduces the likelihood of the latter's interaction with the APDAP loop.

Under these circumstances, AF in fact prefers to dock the N-terminal [M/Φ]xxYG[K/R] motif

to the APDAP (f19D), possibly because is it further from the FNFLD motif and therefore less

restricted by the latter's association with WEST. How then does association between the

APDAP loop and the TYxxxR[T/S]ΦL motif come about? A clue emerges from AF predictions

for the interaction between Scc3 and Pds5 in yeast. Unlike humans where there is no direct

interaction between SA and Pds5, AF predicts with high confidence in five out of five models

an interaction between Scc3's C-terminal face (Scc3$_C$) and the pronounced protrusion within

the middle of Pds5 (Pds5$_P$) (Fig. 29A, f54A). Crucially, both surfaces involved are the sites of

mutations (Fig. 29B) that abrogate releasing activity in vivo, namely R996S and R1043L

within Scc3's C-terminal face and R578I, C599F, and E602K within the face of Pds5$_P$

predicted to interact with Scc3$_C$ (Rowland et al., 2009). The remarkable congruence between

the AF prediction and mutations isolated as spontaneous suppressors of *eco1Δ* is unlikely to be

a co-incidence and therefore implies that the interaction must be of physiological importance

and necessary for releasing activity, at least in yeast. There is a simple explanation why no

such interaction was detected in humans, namely that SA proteins have seven extra HEAT

repeats at their C-termini, which means that they lack the Scc3$_C$ face that interacts with Pds5$_P$

in yeast. These extra repeats are shared by Scc3 orthologs in plants, basal fungi, and

basidiomycetes, but lacking in *S. pombe* and *N. crassa* as well as *S. cerevisiae*. AF also

predicted, albeit with low confidence a similar association between Scc3$_C$ and Pds5$_P$ in *N.*

crassa (f54E).

Figure 29. *A, AF prediction for the interaction between Scc3 and Pds5 in S. cerevisiae (f54A).*

B, *close up showing residues whose mutation abrogates RA and suppresses eco1Δ.*

Though the Scc3$_C$ face involved in the interaction appears unique to ascomycetes, it has a

potentially important implication for how canonical rigid body Pds5:Scc3:Wapl ternary

complexes are generated in *S. cerevisiae*. This emerges from the fact that the Scc3$_C$:Pds5$_P$

interaction aligns the axes of the two proteins in parallel and thereby juxtaposes their N-termini,

albeit modestly spaced apart (Fig. 29A). As a consequence, any Wapl CTD bound to Scc3's

CES will now be placed in the vicinity of Pds5's NTD, which may favour the binding to the

latter's APDAP loop of the TYxxxR[T/S]ΦL motif (associated with Wapl's CTD) over the

more distant N-terminal [M/Φ]xxYG[K/R] motif.

To test this, we used AF to predict the structure created by a fragment of Wapl containing its

TYxxxR[T/S]ΦL motif together with the part of its CTD that binds Scc3's CES together with

full length Scc3 and a long fragment of Pds5 that contains its entire N-terminal part as well as

its Pds5$_P$ protrusion predicted to bind Scc3$_C$. Remarkably, AF predicted with high confidence

in five out of five models (f54B) a complex with Scc3$_C$ bound to Pds5$_P$, Wapl's CTD bound to

Scc3's CES, a close juxtaposition of the latter with Pds5's NTD, and interaction of Wapl's

TYxxxR[T/S]ΦL with Pds5's APDAP loop (Fig. 30). A similar result was obtained when all

of Wapl's N-terminal sequences were included and crucially under these circumstances Wapl's

TYxxxR[T/S]ΦL and not its N-terminal [M/Φ]xxYG[K/R] motif bound to Pds5's APDAP

loop (f54C).

Figure 30. AF prediction for a complex between S. cerevisiae Scc3, Pds5, and Wapl's CTD

with its associated TYxxxR[T/S]ΦL motif (f54B).

This finding suggests that the Scc3$_C$:Pds5$_P$ interaction facilitates juxtaposition of Wapl's CTD

with Pds5's NTD in manner that would place them close to but not identical to that adopted in

896 the canonical configuration. It also promotes the interaction of Pds5's APDAP loop with

897 Wapl's TYxxxR[T/S]ΦL motif that is ultimately necessary for formation of the canonical rigid

898 body Pds5:Scc3:Wapl ternary complex in yeast. However, the latter is not in fact generated

899 because its precise adoption is incompatible with the manner in which the axes of Scc3 and

900 Pds5 are oriented due to their Scc3$_C$:Pds5$_P$ interaction. As a consequence, in this alternative

901 state, Scc1$_N$ associated with Pds5 is not placed in the position necessary for its sequestration

902 by the cleft in Wapl's CTD (Fig. 31A). For this to occur, the Scc3$_C$:Pds5$_P$ interaction must

903 subsequently be broken, which allows Pds5's axis to rotate in a manner that slots it into the

904 canonical configuration, which then delivers Scc1$_N$ to the cleft in Wapl's CTD (Fig. 31B).

905 Though speculative, the scheme has the merit of explaining why Wapl's TYxxxR[T/S]ΦL

906 motif as well as the Scc3$_C$:Pds5$_P$ interaction are essential for RA in yeast but not in animals,

907 and yet RA in both types of organism appears to share the fundamental mechanism, namely

908 delivery of Scc1N to Wapl's CTD when part of the canonical Pds5:Scc3/SA:Wapl ternary

909 complex. If this mechanism is indeed correct and universal, then we must conclude that not all

910 of the interactions revealed by AF and by yeast genetics can occur simultaneously. The

911 Scc3$_C$:Pds5$_P$ interaction is clearly incompatible with formation of the canonical ternary

912 complex and yet the former is essential for RA in yeast. In other words, the proteins involved

913 in RA in yeast must exist in multiple states and the process of release must involve transitions

914 between these that presumably depend on a specific combination of on and off rates for the

915 various interactions involved.

Figure 31. Putative quarternary complexes between yeast $Scc1_N$, Pds5, Scc3, and Wapl. A, when formed with $Scc3_C$ interacting with $Pds5_P$ (f55), created by superimposing f24 and f38B onto f54B. B, when formed without any interaction between $Scc3_C$ and $Pds5_P$ (f53), which involves a canonical rigid body Pds5:Scc3:Wapl ternary complex. Model created by superimposing f24 and f38B onto f32B. A movie showing the transition between A and B at https://doi.org/10.6084/m9.figshare.22567525.v1.

Yeast genetics and AF suggest a novel function for Smc3 K112 and K113

Though our current AF predictions explain many of the existing mutations known to abrogate releasing activity in yeast as well as providing the first plausible explanation for how Wapl

facilitates release, they do not explain how acetylation of Smc3(K112) and K113 by Eco1 in

yeast blocks release in a manner that does not require Sororin, nor why release is abrogated by

mutations of Smc3's KKD loop, Smc3(K113T, K113N, K112Q K113Q and D114Q/Y) and its

surroundings Smc3(S75R, R107I/S, G110R/V/W) or by D189E and E202K situated on the N-

terminal face of Scc3 (Scc3$_N$). Because it was unclear whether or not Smc3's KKD loop binds

another protein or merely binds DNA, we focused on what proteins might bind Scc3$_N$. Because

the predictions described above excluded both Wapl (f24) and Pds5 (f54A), we tested whether

Scc3$_N$ might bind Smc3's ATPase domain. Remarkably, AF predicted with high confidence,

in the top four ranked models, an interaction between the face of Scc3$_N$ containing D189 and

E202 and the part of Smc3's ATPase containing K112 and K113 (Fig. 32A, f56A). Not only

do both Smc3(K112) and Smc3(K113) participate in the interaction but so does Smc3S75,

whose substitution by R abrogates RA in yeast (Rowland et al., 2009). Again, the striking

congruence between the predicted interaction and residues known to be essential for RA

implies that the interaction is of physiological importance. We therefore presume that

Scc3(D189E) and E202K abrogate RA because they compromise the interaction between

Scc3$_N$ and Smc3's KKD loop. Because Scc3(E202K) suppresses the lethality caused by

unregulated RA caused by Smc3(K112R, K113R) double mutants as well as *eco1Δ* (Rowland

et al., 2009), we can also be confident that the AF prediction (f56A) recognizes K112 and K113

in their unmodified state. Acetylation of K112 and K113 would presumably also abrogate the

interaction in vivo and given that the phenotype of Scc3(D189E) and E202K mutations imply

that the interaction is necessary for RA, the AF model finally provides an explanation for how

acetylation would be sufficient to block RA in yeast, without any need for Sororin.

The face of Scc3$_N$, including D189 and E202, is highly conserved in fungi and AF predicted a

similar interaction between Scc3$_N$ and Smc3's ATPase domain in *Candida albicans* (f57A) but

not, strangely, in *N. crassa* (f57B). D189 and E202 are not conserved in metazoa and AF also

failed to detect any interaction between SA/Scc3$_N$ and Smc3's ATPase domain in humans

(f57C), where the physiological role of K105 and K106 acetylation has never been rigorously

investigated. What is known is that in G2 cells Sororin is necessary to protect cohesion from

Wapl and if one assumes that K105 and K106 acetylation still takes place when Sororin is

depleted (which is not in fact known), then unlike yeast their acetylation cannot be sufficient

to abrogate a form of RA capable of destroying cohesion.

Finally, AF predicted a very similar interaction between Scc3$_N$ and Smc3's ATPase when the

former was bound by Wapl's CTD (Fig. 32B, f56B) and even when it was part of a canonical

ternary complex with Wapl and Pds5 (Fig. 32C, f56C), implying that the sequestration of

Scc1's NTD by Wapl's CTD could in principle occur when Scc3$_N$ is bound to Smc3's KKD

loop. If so, Scc3$_N$ would be held at 110° from Smc3's neck, which may help prevent their re-

association during the release reaction (Fig. 33). However, it is also possible that the interaction

between Scc3$_N$ and Smc3 has a role at some other stage of the release process. Furthermore, it

is presently unclear how the complex portrayed in Fig. 31 facilitates cohesin's release from

DNA. If the latter is entrapped within the Smc-kleisin ring, then it must pass through the opened

Scc1$_N$:Smc3 interface, which is hard to envisage if Scc3 remains bound at this stage to Smc3's

KKD loop. Though essential, the binding of Scc3$_N$ to Smc3's KKD loop is not necessarily the

latter's sole function during release. For instance, K112 and K113 might at some other stage

during the reaction facilitate the binding of DNA destined to be released. Interestingly, there

would be sufficient space between Smc1 and Smc3 coiled coils for DNA to bind to engaged

heads even when this is accompanied by junction of Smc1 and Smc3 joints, which we suggest

here may trigger dissociation of Scc1's NTD from Smc3. Importantly, Smc3(K112) and K113

would be in an ideal position to contribute to such DNA binding.

Figure 32. *Interaction between Smc3's ATPase and Scc3's N-terminal domain.* ***A*** *Left, Detail*

of the interaction between Smc3 and Scc3 (f56A) showing Scc3(D189) and Scc3(E202)

(marked green) and Smc3(S75, K112, K113) (marked red). ***A*** *Right, PAEs for rank 1 model in*

A. ***B***, *Interaction between Smc3's ATPase, Scc3's N-terminal domain, and Wapl's CTD (f56B).*

C, *Interaction of the canonical rigid body Pds5:Scc3:Wapl ternary complex with Smc3's*

ATPase domain (f56C).

Figure 33. *A model for the yeast quarternary Pds5:Scc3:Wapl:Scc1 complex in the canonical*

configuration when its Scc3 moiety is bound to the Smc3 ATPase head as predicted by AF,

created by super-imposing f56A with the model in Fig. 31B. There is a modest clash between

Scc1 and the Smc3 coiled coil that would presumably have been avoided if AF were able to

predict the entire complex, which is beyond its capability.

Regulation of release by Smc3 acetylation

Acetylation of Smc3(K112) and K113 by Eco1 is essential for blocking releasing activity in

G2 cells, a process dependent on Pds5 (Chan et al., 2013). How does Eco1 recognize Smc3's

KKD loop and how might Pds5 facilitate this? We therefore asked whether AF can predict how

Eco1 binds to Smc3, Pds5, or other cohesin subunits. In two (ranks 1 and 2) out of five models

with *S. cerevisiae* (f67) and in five out of five models with *Candida glabrata* (f68), AF predicts

with high confidence an interaction between yeast Eco1 and Smc3's ATPase with Scc1's NTD

bound to the coiled coils emerging from it (Fig. 34A). Importantly, Eco1 binds in a manner in

which K113 would enter the enzyme's catalytic site as observed in a crystal structure of a

K106-CoA conjugate bound to xEco2 from *Xenopus laevis* (PDB: 5N22) (Chao et al., 2017)

(f69) (Fig. 34B, left). The prediction envisions a very different type of association to that

previously proposed on the basis of attempts to dock the acetyl transferase onto Smc3 heads

(Chao et al., 2017). AF predicts, again with high confidence, a very similar mode of association

of Esco1's catalytic domain with the Smc3 ATPase head, albeit in only one out five models, in

the absence (f70) but not in the presence (f71) of Scc1's NTD. No interaction of any nature

was predicted for Esco2, either with or without Scc1. The reason for these differences lie in

the zinc fingers that adjoin the acetyl transferase. In the yeast model, the α helix underlying

the zinc finger almost clashes with the N-terminal helices from Scc1 bound to Smc3, a feature

clearly ignored by the top two AF models (Fig. 34A and Fig. 34B right). It turns out that the

zinc fingers of Esco1 and Esco2 differ from Eco1 with respect to their orientation relative to

the catalytic domain and this difference creates a clash with Scc1 that AF cannot avoid, which

explains why Esco1 is predicted to bind only when Scc1 is not bound to the Smc3 neck. Given

the plausibility of the yeast model, we suggest that binding in vivo is normally accompanied

by an allosteric change in the way the zinc fingers adjoin the catalytic domain, a change that

AF does not predict, presumably because it is not as present in the evolutionary record

imprinted in the sequences used.

Passage through S phase is necessary for acetylation by Eco1 (Beckouet et al., 2010; Rolef

Ben-Shahar et al., 2008) and Esco2 (Ivanov et al., 2018) and likely also stimulates that by

Esco1. S phase-specific acetylation by all three enzymes requires a small motif (PIP box)

within unstructured sequences immediately N-terminal to their zinc fingers that bind to PCNA

(Moldovan et al., 2006) and this association was confirmed by AF (f72, f73, & f74). In the case

of Eco1, association between Eco1 and PCNA was clearly compatible with its association with

the Smc3 ATPase head. The recent finding that Smc3's acetylation by Eco1 in vitro depends

on PCNA having been loaded on DNA (Minamino et al., 2023) raises the possibility that the

latter induces a change in the Zinc finger's conformation that is necessary for docking the

human Smc3 ATPase in the manner depicted in Fig. 34.

1026

Figure 34. A, *AF predictions for Smc3's interaction with Eco1 from S. cerevisiae (f67) and C.*
glabrata (f68) with the Scc1 NTD present. **B**, *Left, similarity of this interaction with a crystal*
structure (PDB: 5N22) of Esco1 with a K106-CoA conjugate. A superimposition of f67 and
5N22 is shown in f69. Right: as in A, left, without Scc1 NTD.

In addition to predicting binding of Smc3 to the catalytic cleft on the front face of Eco1's

catalytic domain, AF predicts with high confidence, in five out of five models, binding of three

loops on the back side of the enzyme to a conserved loop associated with *S. cerevisiae* Pds5's

most C-terminal HEAT repeats (Fig. 35A, f75). A similar but not identical type of association

is predicted for Esco1 (Fig. 35B, f22) and Esco2 (Fig. 35C, f76), both to Pds5A (Fig. 35) and

Pds5B (not shown). In these cases, an additional contact is made by a pair of anti-parallel β

sheets within the catalytic domain with the α helices that protrude from the centre of Pds5A &

B, which due to Pds5's hook shape are in close proximity to its C-terminal helices. Given that

the association between *S. cerevisiae* Pds5's CTD with the back side of Eco1 is their sole form

of interaction, it is reasonable to assume that this interaction alone recruits Eco1 to cohesin and

thereby facilitates its association with the Smc3 ATPase. This has an implication. If Smc3 is

acetylated by Eco1 when bound simultaneously to Pds5's CTD as well as to Scc1's NTD, then

Pds5 must adopt a very different orientation, relative to the Smc3 ATPase, to that predicted by

AF in the context of releasing activity, where its NTD and not its CTD points towards the

ATPase (Fig. 36). Either the segment of Scc1 linking Scc1's NTD to Smc3 is sufficiently

flexible for Pds5 to adopt more than one orientation relative to the Smc3 ATPase or acetylation

promoted by Pds5 would have to take place after Scc1's NTD has dissociated from the Smc3

neck.

1051

Figure 35. *AF predictions for interactions between Pds5 and S. cerevisiae Eco1 (**A**, f75), human Esco1 (**B**, f22), and human Esco2 (**C**, f76).*

1054

1055

1056   ***Figure 36.*** *Projected position of Eco1 bound to Pds5 if Scc1$_N$ bound to the latter adopts the*

1057   *orientation predicted by AF (f36). The model was created by superimposing f36 and f75 via*

1058   *their Pds5 moieties. For Eco1 to bind Smc3 K112 and K113, while still bound to Pds5, the*

1059   *Scc1 sequences between Scc1$_N$ α3 and the part bound to Pds5 must bend by 180^0, or Scc1's*

1060   *NTD must dissociate from Smc3.*

1061

Unlike yeast Eco1, Esco1 and Esco2 have long unstructured N-terminal extensions and AF

predicts numerous interactions involving specific motifs within these, not only with various

cohesin subunits (Figs. 13 & 37), but also with the MCM helicase (Fig. 38). Thus, in addition

to the interaction between Esco1's catalytic domain and Pds5's CTD, AF predicts with high

confidence and in five out five models, interactions of a highly conserved, partly helical, motif

(ILxLCEEIAGEIESD) within Esco1 with the WEST site on both Pds5A (f22) and Pds5B,

namely the very same site bound by Wapl's central FGF motif (Fig. 13). Wapl and Esco1 have

opposing effects on release and presumably compete for binding. In two out of five models,

AF predicts an additional interaction between a conserved RRSQRLQQL motif and an adjacent

(more C-terminal) groove within Pds5's CTD. AF detected no such interactions between Pds5

and Esco2's N-terminal sequences (f76).

The theme of competition between Wapl and Esco1/2 is also observed with SA1:Scc1, where

AF predicts, albeit with less confidence, that the aforementioned ILxLCEEIAGEIESD helical

motif from Esco1 predicted to bind Pds5's WEST has an affinity also to SA1's CES, binding

orthogonal to SA1's HEAT repeats, in a manner similar to the first helix from Wapl's CTD

(Fig. 37A, f77). More importantly, AF predicts that yet another conserved motif

(RFFPSRKxxP[V/L]KC) binds the same groove within the central section of SA1 bound by

Wapl (Fig. 37A, f77). Like the latter, binding extends across the Scc1 polypeptide. Despite

binding to the same locus, the motif has only modest resemblance to the one used by Wapl

Interestingly, a largely unrelated but broadly conserved motif within Esco2's N-terminal

domain is predicted to bind in a similar fashion to the equivalent locus on a SA2:Scc1 complex

(Fig. 37B, f78). This motif was originally identified as the PBM-B motif required for the

recruitment of Xenopus Eco2 (xEco2) to chromatin in a pre-RC dependent manner (Higashi et

al., 2012) and subsequently postulated to bind cohesin (Ivanov et al., 2018). A second

conserved motif (PBM-A) is required for xEco2's recruitment to chromatin and for its ability

to acetylate Smc3 (Higashi et al., 2012). This was subsequently shown by co-

immunoprecipitation to bind the MCM helicase. Because analysis of amine-reactive chemical

crosslinking using mass spectrometry indicated Mcm4 or Mcm7 as the most likely partners

(Ivanov et al., 2018), we used AF to predict Esco2's interaction with each of these subunits No

interaction was observed with Mcm4 (f79), but three out of five AF models predicted with high

confidence an interaction between PBM-A and Mcm7 (Fig. 38, f80). The motif is predicted to

form two short helices connected by a conserved loop. The first of these binds to the surface

of Mcm7's CTD while part of the loop together with the second helix is sandwiched between

the CTD and a C-terminal winged helical domain that binds the surface of ORC1 and Cdc6

within the pre-RC (Zhai et al., 2017). This interaction together with its PIP box may impart at

least some of the S phase specificity of Smc3 acetylation by Esco2.

1098

1099

1100   ***Figure 37. A**, AF predictions for the interactions between SA1:Scc1 and the N-terminal*

1101   *extension of Esco1 (left, f77) and between SA2:Scc1 and Wapl-M (right, f25). **B**, AF prediction*

1102   *for the interaction between SA2:Scc1 and Esco2's PBM-B (f78).*

1103

1104

1105   ***Figure 38.*** *AF prediction for the interaction of Esco2's PBM-A and Mcm7 (f80).*

1106

Regulation of release by Sororin

Though Smc3 acetylation may be sufficient to protect sister chromatid cohesion from Wapl-
mediated release in yeast, it is necessary (Nishiyama et al., 2010) but not sufficient in
vertebrate, insect, and plant cells, where proteins called Sororins (vertebrates and plants) or
Dalmatians (insects) (Mota et al., 2022; Rankin et al., 2005; Yamada et al., 2017) are also
required. Sororin is absent from G1 mammalian cells because it is degraded at the hands of
APC[Cdh1] and therefore only accumulates shortly before S phase. However, accumulation alone
is insufficient for its interaction with chromosomal cohesin as its association also depends on
DNA replication in Xenopus extracts that lack Cdh1 (Nishiyama et al., 2010). Association with
cohesin only takes place when Smc3 is acetylated in an S phase-specific manner, largely by
Esco2, and on DNA that has actually been replicated, whereupon it is essential for maintaining
cohesion between sisters (Ladurner et al., 2016). Sororin is thought to bind to and protect from
Wapl only cohesin engaged in cohesion, but whether this is really the case is not known.
Importantly, though Smc3 acetylation is necessary for Sororin's association, it is insufficient
and some other change in the state of cohesin that occurs during replication may be necessary.
Sororins are largely unstructured and for the most part poorly conserved. Common to all

members of the family are so called C-terminal Sororin boxes predicted to contain a short (six

turns)   α   helix   and   whose   consensus   in   animals   is

[W/F/Y]xxxΦNxxFxE[A/Φ][E/D]x[F/Y]ELxΦE.

Three other features are common to vertebrate Sororins, namely a short stretch of basic amino

acids immediately N-terminal to the Sororin box, which is preceded by a FGF motif (consensus

[S/T]ΦFGFExL), separated by ten residues from a so called YSR motif (consensus

[M/W]S[K/Q]KVRRSYSRL). Functional studies indicate that the Sororin box is essential,

while the YSR and FGF motifs are not, at least when the protein is over-expressed (Wu et al.,

2011). Because the cohesion defects associated with sororin inactivation are largely suppressed

by Wapl mutants (Ladurner et al., 2016; Nishiyama et al., 2010), Sororin is believed to be

required merely to suppress Wapl-mediated release, a function it fulfils despite a residence

time on cohesin of about 60 seconds (Ladurner et al., 2016). It could perform this function

either by inhibiting Wapl's sequestration of Scc1's NTD or by preventing the latter's

dissociation from Smc3 in the first place. Though AF cannot address why Sororin only

associates with replicated DNA, it has the potential to reveal which cohesin subunits interact

with its essential Sororin box and/or FGF and YSR motifs. Accordingly, we used AF to predict

the interaction of full-length mouse Sororin with cohesin subunits known to be involved in

Wapl-mediated RA. AF predicted a potential interaction between Sororin's FGF motif and the

CES of an SA:Scc1 complex, albeit in only one out five models (f81) and not in the same

manner as CTCF's YDF motif (Li et al., 2020). In contrast, it predicted with high confidence

an interaction between its YSR and FGF motifs and Pds5B's C-terminal domain (Fig. 39), both

in mice (f82) and in zebra fish (f21). Like Wapl, Sororin's FGF was predicted to bind to Pds5's

WEST locus while the YSR bound to an adjacent (more C-terminal) pair of helices. In both

cases, the highly conserved aromatic residues in each motif were inserted into grooves between

adjacent HEAT repeats and, notably, all five models predicted simultaneous association.

Crucially, none of them predicted binding of the YSR to Pds5's N-terminal APDAP loop,

which was initially surprising as it has been shown to compete with the binding of Wapl's YSR

to this site (Ouyang et al., 2016). The reason for this becomes apparent when one takes into

account that Sororin's YSR and FGF motifs are separated by no more than 15 amino acids,

which is too short to permit simultaneous binding of the YSR and FGF motifs to Pds5's N-

terminal APDAP loop and C-terminal WEST respectively.

1154

1155

1156

1157 **Figure 39**. *AF predictions for Sororin's interaction with Pds5B in mouse (f82) and Zebra fish*
1158 *(f21).*

1159

Given that AF failed to predict any association between the C-terminal Sororin box and either
SA2, Pds5B, or indeed Wapl itself (f83), we considered whether it might instead bind to the
Smc3:Scc1 interface. Remarkably, AF predicted, with high confidence in two out of five
human models, interaction with a Smc3 ATPase domain whose coiled coils were bound by
Scc1's NTD (Fig. 40, f84). A very similar interaction was observed with high confidence in
five out of five models using orthologous proteins from *Xenopus laevis* (f85), *Danio rerio*
(zebra fish) (f86), *D. melanogaster* (f87), *Tribolium casaneum* (red flour beetle) (f88), *A.*
thaliana (f89), and even the ascomycete *Aspergillus fumigatus* (f90). The similarity of the
predictions is remarkable given that the Sororin orthologs of vertebrates, insects, plants, and
fungi have very little sequence homology and even exhibit considerable variation between their
Sororin boxes (Mota et al., 2022). In contrast, no interaction was predicted between Sororin
and a complex between Smc1 and Scc1's CTD in *D. melanogaster* (Fig. 41, f91), *A. thaliana*
(Fig. 41, f92), and *A. fumigata* (Fig. 41, f93). Intriguingly, AF suggests that Sororin's C-
terminal helix might interact with the longer of the two helices within Scc1's CTD in mammals
(Fig. 41, f94). Thus, AF's prediction that the Smc3:Scc1$_N$ interface is a highly conserved
partner of Sororin's CTD is not shared by Smc1:Scc1$_C$.

1176

1177

1178   ***Figure 40.*** *AF predictions for Sororin's interaction with Smc3 bound to Scc1's NTD in humans*

1179   *(f84), X. laevis (f85), D. rerio (f86), D. melanogaster (f87), T. casaneum (f88), A. thaliana (89),*

1180   *and A. fumigatus (f90).*

1181

1182

1183   ***Figure 41***. *AF predictions for the Sororin box's interaction with a complex between Smc1 and*

1184   *Scc1's CTD in D. melanogaster (f91), in A. thaliana (f92), in A. fumigatus (f93), and Mouse*

1185   *(f94).*

1186

According to all of the AF models, Sororin's C-terminal α helix straddles Scc1's most N-

terminal helix and the loop connecting its fourth helix to the long α helix that forms a three-

helical bundle with the Smc3 coiled coil (Fig. 40). Meanwhile, the C-terminal end of the

Sororin helix contacts the C-terminal α helix emerging from the Smc3 ATPase just before it

forms a coiled coil with its partner from the N-terminal part of the protein, while the terminal

4-5 amino acids straddle a pair of adjacent β sheets. Aromatic residues play a key part in the

interface. In vertebrates, a conserved tryptophan (W233 in humans) sits on top of Scc1's most

N-terminal helix, while a pair of highly conserved phenylalanines (F241 and F247) interact

with the aforementioned Smc3 α helix respectively (Fig. 40, human rank 1, f84). The nature

of Sororin's interaction with the Smc3:Scc1 interface is not only universal across eukaryotic

kingdoms but also helps explain how Sororin hinders release. By binding Scc1's N-terminal

helix, Sororin would prevent its unfolding from Scc1$_N$'s long α3 helix, thereby preventing

sequestration occurring even before the NTD has dissociated from Smc3. Perhaps more

importantly, by binding over Scc1's N-terminal helices and simultaneously binding Smc3,

Sororin could actually stabilize the interaction between Scc1's NTD and the Smc3 neck,

thereby thwarting any subsequent sequestration of its N-terminal helices by Wapl. Such

stabilization might also prevent the distortions in Smc3's neck produced by simultaneous head

engagement and joint junction, thereby preventing such an event. Despite these insights, AF

predictions provided no clues with regards to the role of Smc3 acetylation.

Finally, we tested whether Sororin might also bind Wapl's CTD. Remarkably, AF predicted

with high confidence in five out five models that Sororin's [S/T]ΦFGFExL motif would bind

to exactly the same cleft within Wapl's CTD that sequesters the most N-terminal α helix from

Scc1 and does so in a manner that would be mutually exclusive (Fig. 42, f83). As already

mentioned, this motif is also predicted to bind Pds5's WEST (Fig. 13B, f21 & Fig. 39, f82) and

by doing so ensures that Sororin's [M/W]S[K/Q]KVRRSYSRL motif binds to an adjacent

groove in Pds5's CTD (Fig. 39, f82) and not to its N-terminal APDAP loop. However, if

Sororin's [S/T]ΦFGFExL motif was instead bound to Wapl's CTD (via the latter's putative

Scc1$_N$ sequestration cleft), when part of the canonical ternary RA complex with SA and Pds5,

then the [K/Q]KVRRSYSRL motif would be free to bind to Pds5's APDAP, which is brought

into close proximity to the aforementioned cleft in Wapl's CTD by ternary complex formation.

In other words, if Sororin's [S/T]ΦFGFExL and [M/W]S[K/Q]KVRRSYSRL motifs are to

bind cohesin simultaneously, the short linker connecting them will constrain the distance

allowed between their binding sites. If the [S/T]ΦFGFExL motif binds to Pds5, its partner must

also bind to an adjacent site on Pds5 (Fig. 39) but if [S/T]ΦFGFExL instead binds to Wapl's

CTD, its [M/W]S[K/Q]KVRRSYSRL partner could in principle bind simultaneously to Pds5's

APDAP loop.

To test this, we asked AF to predict the interaction between Sororin and an N-terminal fragment

of Pds5 that lacked its WEST [S/T]ΦFGFExL binding site. In this case, AF predicted with high

confidence and in five out of five models an interaction exclusively between Sororin's

Dxx[M/W]S[K/Q]KVRRSYSRL motif and Pds5's APDAP (Fig. 43 left, f95). In this case, the

interaction involved conserved aspartic acid (D135) and methionine (M138) residues at the N-

terminal end of the motif, which are predicted to be part of a short helix (marked in grey in Fig.

43), as well as the YSR sub-motif at its C-terminal end, whose interaction with Pds5's APDAP

is predicted by AF to be identical to that observed in a crystal structure containing Wapl's N-

terminal YSR motif bound to Pds5 (Ouyang et al., 2016) (PDB 5HDT) (Fig. 43, right). These

predictions therefore raise the possibility that Sororin may have a second line of defence against

Wapl in case dissociation of Scc1's NTD from Smc3 is accompanied by formation of the

canonical ternary (SA:Wapl:Pds5) RA complex, namely occupation of the key cleft in Wapl's

CTD by Sororin's [S/T]ΦFGFExL motif, which would exclude sequestration of Scc1$_N$ (Fig.

42), and be facilitated by the simultaneous binding of Sororin's [M/W]S[K/Q]KVRRSYSRL

motif to the neighbouring N-terminal APDAP loop of Pds5 (Fig. 43).

1238

1239

1240 ***Figure 42.*** *AF's prediction for the interaction between Sororin and Wapl's CTD (f83).*

1241

1242

Figure 43. AF predicts that Sororin's [M/W]S[K/Q]KVRRSYSRL (YSR) motif binds to Pds5's
APDAP loop (f95) (left) in a manner similar to Wapl's YSR peptide (right, PDB 5HDT), but
only if binding of Sororin's [S/T]ΦFGFExL motif is prevented from binding Pds5's WEST site
by deleting the C-terminal part of Pds5 containing this WEST site.

Interactions between Scc2/Nipbl and Scc3/SA

One of the key insights revealed here by AF is the canonical rigid body association between
SA and Pds5 induced by their simultaneous association with Wapl's CTD (Fig. 16). According
to AF, SA and Pds5 do not otherwise interact directly and only do so in the presence of Wapl.
Note however, that the yeast orthologs do so via their Scc3$_C$:Pds5$_P$ interaction (Fig. 29). In the
course of investigating their association, we used AF to undertake a more systematic
investigation of how cohesin's three HAWK subunits interact. Similar to SA/Scc3, AF also
predicted no interaction between Nipbl/Scc2 and Pds5. In contrast, it predicted with high
confidence and in five out five models a pair of simultaneous interactions between full length
versions of SA and Nipbl (Fig. 44A, f96). One of these interactions is identical to that observed
in a cryo-EM structure of the clamped state (Fig. 44C, PDB 6WG3) (Shi et al., 2020) and
involves interactions between Nipbl helices next to its N-terminal protrusion and the outside
surface of SA where there is a twist in the alignment of its anti-parallel HEAT repeats at the
point where the protein starts to bend. The second region of interactions has not hitherto been
observed and involves an interaction between helices at the point where Nipbl starts to bend
(on its outer surface) and the antiparallel helices that make up SA's pronounced "nose". An
identical pair of interactions were predicted for the orthologous proteins from *A. thaliana* (Fig.
44B, f97).

1266

1267

1268   **Figure 44.** *AF predictions for interactions between Nipbl/Scc2 and SA/Scc3 for humans (**A**,*
1269   *f96), A. thaliana (**B**, f97), and S. cerevisiae (**D**, f98). **C**, cryo-EM structure PDB 6WG3.*

1270

In contrast, AF predicted only the nose interaction for yeast Scc2 and Scc3 (Fig. 44D, f98). In
this case, the residues involved are highly conserved on both the Scc3/SA (Scc3: Q323, E327
and SA1: Q165, E169) and Scc2/Nipbl (Scc2: P958, Y959 and Nipbl: P2029, Y2030) sides of
the interface and the interaction is therefore in all probability of physiological importance.
Apart from microsporidians (Roig et al., 2014), where it is absent, the nose is a highly
conserved feature of Scc3/SA orthologs. Its deletion is lethal in yeast (Roig et al., 2014), but
its function has hitherto remained obscure. It is interesting that the interaction between SA's
nose and Nipbl was not observed in the PDB 6WG3 cryo-EM structure (Shi et al., 2020). This

is probably because it would have been incompatible with DNA binding to Nipbl, which was

an important feature of the clamped structure. Now that AF has revealed the details, it will be

possible to create specific mutations that affect it and thereby address whether the interaction

is necessary either for LE or for DNA entrapment within S-K rings, both in vitro and in vivo.

1283

DISCUSSION

Our use of AF to predict interactions between cohesin subunits has led to numerous hypotheses

concerning how cohesin's association with chromatin is regulated. Can one be sufficiently

confident to undertake the biochemical and genetic experiments needed to test them? And if

so, what will be the most effective techniques to employ? Aside from AF's excellent track

record, it so happens that in several instances experimental data, particularly the phenotype of

mutations, was already in the public domain but had hitherto not been interpretable. Thus, AF's

predictions about the interaction between Wapl's CTD and Scc3/SA explains why mutation of

both Wapl and Scc3/SA residues predicted by AF to affect their interaction are defective in RA

activity in vivo (Beckouet et al., 2016; Hara et al., 2014; Ouyang et al., 2013; Rowland et al.,

2009). Likewise, the phenotypes of an extensive set of mutations affecting highly conserved

surface residues within the C-terminal half of Wapl's CTD (Ouyang et al., 2013) are fully

consistent with AF's prediction that its role is to sequester Scc1's N-terminal helices (α-1 and

α0). Even more impressive is the remarkable congruence between cohesin subunit interfaces

revealed by AF and the locations of mutations that abrogate RA in yeast and were isolated

without any prior hypothesis, as spontaneous suppressors of *eco1* mutant lethality (Rowland et

al., 2009). Yet further support comes from the congruence between AF predictions and

sequence conservation (beyond that used by AF), which reflects natural selection and hence is

also a robust measure of function. As a consequence, our analyses have been guided not only

by the degree of AF's confidence but also by a requirement that predictions be broadly based

from a taxonomic viewpoint, namely similar in multiple and as distant as possible eukaryotic

lineages, and concern sequences and domains that are evolutionarily conserved. Despite this,

AF revealed instances of physiologically important interactions specific to particular

eukaryotic lineages, for example the interactions between the N- and C-terminal faces of Scc3

with Smc3's ATPase and Pds5's protrusion respectively, which were only detected in yeast

and related fungi. Last but not least, a key guiding principle has been whether the models

appear plausible and above all explanatory.

In so far that AF largely functions by correlating the query sequences with known structures
and their sequences, and not by calculating chemical or physical properties from first
principles, there is reason to be optimistic that the results will be in many cases comparable in
robustness to those obtained by experimental structural biology, which often uses or even
requires unwarranted conditions (protein concentration, salt concentration, divalent cations,
pH, temperature, etc) and usually evaluates interactions either pairwise or among a very limited
number of constituents and not as in vivo where proteins must compete with a large fraction of
the proteome, as well as with other macromolecules. We contend that many if not most of the
hypotheses described in this paper are plausible, novel, in some cases provocative, and above
all explanatory, and therefore deserve airing in public before validation using orthogonal
approaches.

What then might these be? At the top of the list must be renewed attempts to visualize the
predicted complexes by cryo-EM alongside the use of bi-functional thiol specific reagents to
crosslink both in vitro and in vivo cysteine pairs inserted into the predicted interfaces (Chapard
et al., 2019; Srinivasan et al., 2018). Mutation of key residues will also have an important role,
but it is important to stress that mutation alone rarely addresses whether the resulting
physiological defects are due to the loss of a specific interaction. They merely reveal whether
or not one or other interaction surface is important. To achieve this goal using mutations alone,
it would be necessary to generate not just one or two mutations but a large series and
demonstrate a robust structure-function relationship, or better still demonstrate that mutations
in both sides of an interface produce identical phenotypes. A good example of the latter is our
finding of RA-abrogating mutations identified through classical (i.e. hypothesis free) forward
genetics mapping to both sides of interfaces identified by AF, for example that between $Scc3_C$
and $Pds5_P$ and between $Scc3_N$ and Smc3's KKD loop. Combining cross-linking with a more
limited mutational analysis is probably therefore the most economical way of proceeding, but
even this is will be a considerable undertaking as all substitutions should be made within
endogenous genes.

Though our experience is anecdotal, the modelling that we undertook for this analysis was
sufficiently extensive that it is worth sharing the impressions gained into the strengths and
weaknesses of AF as well as what may transpire to be best practice. First and foremost, is the
issue of the optimum scale for embarking on a prediction. On the one hand, AF's ability to
detect patterns between specific domains clearly diminishes as the size and number of
components increase. The combinatorial space increases with the square of the number of

residues provided in the query sequence(s). On the other hand, predicted interactions clearly

have greater value if AF manages to predict them within the context of intact and therefore

possibly large proteins or protein assemblies, whose size approaches the limits of what is

currently practical or even possible given GPU memory limitations. It is our experience that

initial forays should be performed while making as few assumptions as possible as to the nature

of the interactions and therefore giving AF as large a landscape as is practically possible.

Nevertheless, following up interactions revealed in this manner also has value. It is also clear

that there is probably no relationship between the confidence of AF predictions and any actual

affinity between proteins, as would be expected given the AF algorithm. If biology demands a

weak interaction and natural selection has delivered this, AF clearly excels in picking up the

resulting patterns and in this regard it provides an approach that is complementary to

conventional biochemistry and structural biology with which weak interactions are sometimes

difficult to investigate. As a consequence, many of the interactions detected by AF may be

challenging to pursue using these techniques and combining experimental structural work with

the sort of cross-linking techniques described above may prove useful in this regard. Another

important lesson, which may prove to be of general validity, is that complexes such as cohesin

should not be regarded as conventional solids, which imply a lack of constituent mobility or as

liquids, which would require massively larger stoichiometries. They should instead be

considered as promiscuous molecular communities made up by a large set of potential

interactions with a great heterogeneity of on and off rates and extensive competition between

different constituents, many of which have multiple possible partners and multiple binding

interfaces. Some of the interactions are between relatively rigid domains but many are between

short motifs embedded in very long intrinsically disordered loops and extensions.

Understanding the physical dynamics of such communities as well as the mechanisms

underlying the switches between states will be a formidable challenge. In the case of cohesin,

both loop extruding and cohesive forms must be capable of switching between forms that

associate stably with chromatin fibres to ones that dissociate, and a competition for shared

binding sites between factors promoting release or its inhibition presumably create bi-stable

switches.

In summary, our AF predictions have yielded several hypotheses for how cohesin's association

with chromatin is regulated. These include:

1) The role of cohesin's hinge in effecting a junction between the joints within Smc1 and

Smc3's coiled coils. When combined with ATP-driven head engagement in the absence of

Scc2/Nipbl, but in the presence of Pds5, this junction is predicted to create a distortion of the

coiled coil to which the Scc1's NTD normally binds and thereby triggers its dissociation.

2) The proposal that Wapl's CTD forms a ternary complex with the pair of cohesin's HAWK

subunits previously implicated in release, namely SA/Scc3 and Pds5. Upon dissociation from

Smc3, Scc1 is predicted to bind to Pds5 in a manner that places Scc1's short N-terminal helices

in close proximity to a highly conserved site within Wapl's CTD when part of the ternary

complex with SA and Pds5. The subsequent unfolding of Scc1's α-1 and α0 helices from the

long α3 helix that normally associates with Smc3's neck then leads to their sequestration by

this part of Wapl. We propose that sequestration in this manner is the chief function of Wapl

and that it serves to keep the exit gate open long enough for DNA to pass out of the cohesin

ring.

3) The finding that in yeast formation of the canonical ternary complex needed to deliver Scc1's

N-terminal helices to Wapl may be a multi-stage process during which an alternative ternary

complex involving the $Scc3_C$:$Pds5_P$ interaction is required for subsequent formation of the

canonical one.

4) The first insight into how Smc3's KKD loop facilitates release, at least in *S. cerevisiae* where

it binds the N-terminal face of Scc3.

5) An understanding how the acetyl transferases that block release and compete with Wapl for

binding cohesin's HAWK subunits bind to their substrate and how this is facilitated during S

phase by simultaneous binding to Pds5 and to the Mcm helicase.

6) The first insight into how Sororin is recruited to cohesin and how it hinders Wapl-mediated

release, through the association of a conserved C-terminal α helix with cohesin's exit gate,

namely the junction of Scc1's NTD with Smc3's neck.

7) The discovery that Scc2/Nipbl and SA/Scc3 interact with each other via an association

involving the latter's highly conserved nose, for which there has hitherto been no explanation.

Whether this association has a role during LE or in the entrapment of DNA inside the cohesin

ring remains to be addressed.

Of these, the notion that a key function of Wapl's CTD is to bind α-1 and α0 from $Scc1_N$ is

probably the most remarkable as it is based not on a single AF prediction but on an elaborate

concatenation of numerous models, producing a quarternary complex involving Wapl, Pds5,

SA/Scc3, and Scc1, in which Scc1's N-terminal α helices are positioned next to a pair of crucial

sites within Wapl's CTD. Key observations underpinning this notion include the conservation

of Wapl's putative $Scc1_N$ binding site, the existence of mutations within it that abrogate release

in vivo both in yeast and in mammalian cells, the prediction by AF that Pds5, SA/Scc3, and

Wapl form a highly specific ternary complex in animal, plant, and fungal lineages, that $Scc1_N$'s

association with the spine of Pds5's NTD places the formers' $\alpha-1$ and $\alpha0$ helices next to the

putative binding site within Wapl's CTD, that formation of the ternary complex in fungi

depends on a motif within Wapl's unstructured N-terminal sequences which binds Pds5's

APDAP loop and is essential for RA in *S. cerevisiae*, AF predictions for the interaction between

$Scc1_N$ and Wapl that appear conserved among a very wide variety of eukaryotes, and lastly on

the fact that dissociation of $Scc1_N$ from Smc3 depends on Wapl in vivo in *S. cerevisiae* and

that RA is abrogated by their co-translation. Whether like a jigsaw puzzle, our hypothesis

proves to be the sole solution to these explicanda will nevertheless require numerous types of

wet experiments.

Last but not least, it should be pointed out that due to the nature of AF, none of our models

incorporate DNA and therefore shed no direct insight into how sequestration of Scc1N by Wapl

actually facilitates cohesin's dissociation from chromosomes in vivo.

METHODS

AF predictions prior to February 2023 were performed via Google CoLab using Alphafold 2,

while those performed thereafter used Colabfold v1.5.2 AF2 with MMSeqs2. Supposition of

different models were performed using UCSF Chimera while all figures in this manuscript and

Movie 1 were created using PyMOL v2.5 (Schrödinger, LLC). Each subunit was drawn in all

figures using the following colours: Smc1=red; Smc3=blue; Scc1=green; Scc2/Nipbl=yellow;

Scc3/SA1/SA2=orange; Pds5=violet; Wapl=cyan; Eco1/Esco1/Esco2=ivory; Sororin=grey.

The *WAPL* deletions were introduced into the genome of *S. cerevisiae* using CRISPR-Cas9.

Cells of strain KN18714 (a W303 related strain whose endogenous *WAPL* gene was tagged

with EGFP) were transformed with a 738 nt (for Δ4-9) or a 138 nt (for Δ208-215) PCR "healing

fragment" along with a 2 micron *LEU2* plasmid, constitutively expressing Cas9 endonuclease

and transcribing the structural RNA that contains the Cas9 binding site and the guide RNA that

will target the double strand break formation by Cas9 to the *WAPL* (*RAD61*) gene. The guide

RNA was designed 20 nt upstream of the NGG (E228, V229) using the annealed oligos

MB526F/MB527R (MB526F: 5'-ATCAAATGAAATAATGGGAGAAG-3', MB527R: 5'-

AACCTTCTCCCATTATTTCATTT-3') and fused afterwards at the 5' end of the structural

RNA with Sap1 cloning. The PCR "healing fragment" was created using the oligos

MB543F/MB525R for Δ4-9 and MB534F/MB525R for Δ208-215 (MB543F: 5'-

TCGCAAAACGAAACCATCTTCTTACCCTAAAGCATCCTGTTTCTGAAAAAATGAG

AGCAGTTCTCAGAACTCCTTTTAGATCTAATAAAGGCTTGCCATC-3', MB525R: 5'-

TTACTGTTGTCAGCGTTATTTAGGGACAAGGTGTTAGCTTTCTGATCTACTTCTTC

TCCCATTATTTCATTTTC-3',   MB534F: 5'-

GCATACGCACATCCAAAGACAAAAAATAAACAAAATAAAGAAAATG-3')   and

contained the deletion, a mutagenized NGG Cas9 sequence target (E228, V229: GAGGTA to

GAAGTA) and 5'-, 3'- homology arms needed for repair of the double strand break created by

Cas9. As a control, no colonies were obtained after yeast transformation of the 2 micron *LEU2*

plasmid alone without the "healing fragment". All mutations were confirmed by sequencing

the genomic locus.

1452

The ability of the EGFP-tagged *WAPL* mutations to suppress *eco1Δ* lethality was tested by

crossing them with strain KN16432, which carries deletions of both *ECO1* and *WAPL*.

Dissection of 18 tetrads from crosses involving *WAPLΔ208-215* yielded 60 viable spores, of

which 27 carried *eco1Δ*. Of these 15 carried the *waplΔ* and 12 the *WAPLΔ208-215* mutation.

Thus, *WAPL Δ208-215* was as effective as *waplΔ* in suppressing *eco1Δ* lethality and this allele

is therefore defective in releasing activity. In contrast, dissection of 18 tetrads from crosses

involving *WAPLΔ4-9* yielded 49 viable spores of which 16 carried *eco1Δ.* Of these, all 16

carried *waplΔ.* Thus, *WAPLΔ4-9* cannot suppress *eco1Δ* lethality and is therefore proficient for

releasing activity. Separate experiments demonstrated that *WAPLΔ208-215* suppressed *eco1Δ*

when untagged or tagged with PK9 while *WAPLΔ4-9* never did so. Mutation of both motifs

(*Δ4-9 Δ208-215*) also suppressed *eco1Δ* lethality as did an allele in which the TYxxxR[T/S]ΦL

motif (YGKKRTIL) was inserted at *Δ4−9* (creating the sequence MRAYGKKRTILVLR)

within *Δ4-9 Δ208-215*. In other words, the TYxxxR[T/S]ΦL motif does not appear to function

when placed at Wapl's N-terminus, namely when further from its CTD.

1467

1468   **YEAST STRAINS**

1469

1470

Strain number	Genotype
KN18714	*Mat a, Wapl-EGFP::His*
KN24262	*Mat a, wapl(Δ4-9)-EGFP::His*
KN24060	*Mat a, wapl(Δ208-215)-EGFP::His*
KN24224	*Mat a, wapl(Δ208-215)-EGFP::His, eco1::NatMx*
KN16432	*Mat alpha, eco1::NatMx, wapl::HphMx*
KN26163	*Mat a, wapl(Δ208-215), eco1::NatMx*
KN25817	*Mat a, wapl(Δ208-215)-PK9::NatMX, eco1::HphMx*
KN25168	*Mat alpha, eco1::HphMx, leu2::Eco1p-Eco1::LEU2*

1471

1472   **DATA AVAILABILIY**

1473   All data files (f1-f98) referred to in the main text and figure legends are available at

1474   https://doi.org/10.6084/m9.figshare.22567318.v1. The actual amino acid sequences used for

1475   each AF prediction are to be found embedded in the predicted structure files (in PDB format).

1476

1477   **FUNDING**

1478   This work was supported by the Medical Research Council (U105184326 to J.L.) and by

1479   Cancer Research UK (26747 to K.A.N).

1480

1481   **ACKNOWELDGEMENTS**

1482   K.A.N would like to thank Madhusudhan Srinivasan (Oxford, UK) for numerous discussions

1483   that stimulated and informed this work.

REFERENCES

Bastie N, Chapard C, Dauban L, Gadal O, Beckouet F, Koszul R. 2022. Smc3 acetylation, Pds5 and Scc2 control the translocase activity that establishes cohesin-dependent chromatin loops. *Nat Struct Mol Biol* **29**:575–585. doi:10.1038/s41594-022-00780-0

Beckouet F, Hu B, Roig MB, Sutani T, Komata M, Uluocak P, Katis VL, Shirahige K, Nasmyth K. 2010. An Smc3 acetylation cycle is essential for establishment of sister chromatid cohesion. *Mol Cell* **39**:689–99. doi:10.1016/j.molcel.2010.08.008

Beckouet F, Srinivasan M, Roig MB, Chan KL, Scheinost JC, Batty P, Hu B, Petela N, Gligoris T, Smith AC, Strmecki L, Rowland BD, Nasmyth K. 2016. Releasing Activity Disengages Cohesin's Smc3/Scc1 Interface in a Process Blocked by Acetylation. *Mol Cell* **61**:563–574. doi:10.1016/j.molcel.2016.01.026

Bürmann F, Lee B-G, Than T, Sinn L, O'Reilly FJ, Yatskevich S, Rappsilber J, Hu B, Nasmyth K, Löwe J. 2019. A folded conformation of MukBEF and cohesin. *Nat Struct Mol Biol* **26**:227–236. doi:10.1038/s41594-019-0196-z

Chan KL, Gligoris T, Upcher W, Kato Y, Shirahige K, Nasmyth K, Beckouet F. 2013. Pds5 promotes and protects cohesin acetylation. *Proc Natl Acad Sci U S A* **110**:13020–5. doi:10.1073/pnas.1306900110

Chan K-L, Roig MB, Hu B, Beckouët F, Metson J, Nasmyth K. 2012. Cohesin's DNA exit gate is distinct from its entrance gate and is regulated by acetylation. *Cell* **150**:961–974. doi:10.1016/j.cell.2012.07.028

Chao WCH, Wade BO, Bouchoux C, Jones AW, Purkiss AG, Federico S, O'Reilly N, Snijders AP, Uhlmann F, Singleton MR. 2017. Structural Basis of Eco1-Mediated Cohesin Acetylation. *Sci Rep* **7**:44313. doi:10.1038/srep44313

Chapard C, Jones R, van Oepen T, Scheinost JC, Nasmyth K. 2019. Sister DNA Entrapment between Juxtaposed Smc Heads and Kleisin of the Cohesin Complex. *Mol Cell* **75**:224-237.e5. doi:10.1016/j.molcel.2019.05.023

Chatterjee A, Zakian S, Hu XW, Singleton MR. 2013. Structural insights into the regulation of cohesion establishment by Wpl1. *EMBO J* **32**:677–87. doi:10.1038/emboj.2013.16

Ciosk R, Shirayama M, Shevchenko A, Tanaka T, Toth A, Shevchenko A, Nasmyth K. 2000. Cohesin's binding to chromosomes depends on a separate complex consisting of Scc2 and Scc4 proteins. *Mol Cell* **5**:243–54. doi:10.1016/s1097-2765(00)80420-7

Collier JE, Lee B-G, Roig MB, Yatskevich S, Petela NJ, Metson J, Voulgaris M, Gonzalez Llamazares A, Löwe J, Nasmyth KA. 2020. Transport of DNA within cohesin involves clamping on top of engaged heads by Scc2 and entrapment within the ring by Scc3. *Elife* **9**:e59560. doi:10.7554/eLife.59560

Collier JE, Nasmyth KA. 2022. DNA passes through cohesin's hinge as well as its Smc3-kleisin interface. *Elife* **11**. doi:10.7554/eLife.80310

Davidson IF, Bauer B, Goetz D, Tang W, Wutz G, Peters JM. 2019. DNA loop extrusion by human cohesin. *Science* **366**:1338–1345. doi:10.1126/science.aaz3418

Diebold-Durand M-L, Lee H, Ruiz Avila LB, Noh H, Shin H-C, Im H, Bock FP, Bürmann F, Durand A, Basfeld A, Ham S, Basquin J, Oh B-H, Gruber S. 2017. Structure of Full-Length SMC and Rearrangements Required for Chromosome Organization. *Mol Cell* **67**:334-347.e5. doi:10.1016/j.molcel.2017.06.010

Elbatsh AMO, Haarhuis JHI, Petela N, Chapard C, Fish A, Celie PH, Stadnik M, Ristic D, Wyman C, Medema RH, Nasmyth K, Rowland BD. 2016. Cohesin Releases DNA through Asymmetric ATPase-Driven Ring Opening. *Mol Cell* **61**:575–588. doi:10.1016/j.molcel.2016.01.025

Evans R, O'Neill M, Pritzel A, Antropova N, Senior A, Green T, Žídek A, Bates R, Blackwell S, Yim J, Ronneberger O, Bodenstein S, Zielinski M, Bridgland A,

Potapenko A, Cowie A, Tunyasuvunakool K, Jain R, Clancy E, Kohli P, Jumper J,
Hassabis D. 2021. Protein complex prediction with AlphaFold-Multimer.
doi:10.1101/2021.10.04.463034

Fudenberg G, Imakaev M, Lu C, Goloborodko A, Abdennur N, Mirny LA. 2016. Formation
of Chromosomal Domains by Loop Extrusion. *Cell Rep* **15**:2038–49.
doi:10.1016/j.celrep.2016.04.085

Gandhi R, Gillespie PJ, Hirano T. 2006. Human Wapl is a cohesin-binding protein that
promotes sister-chromatid resolution in mitotic prophase. *Curr Biol* **16**:2406–17.
doi:10.1016/j.cub.2006.10.061

Gerlich D, Koch B, Dupeux F, Peters JM, Ellenberg J. 2006. Live-cell imaging reveals a
stable cohesin-chromatin interaction after but not before DNA replication. *Curr Biol*
16:1571–8. doi:10.1016/j.cub.2006.06.068

Gligoris TG, Scheinost JC, Burmann F, Petela N, Chan KL, Uluocak P, Beckouet F, Gruber
S, Nasmyth K, Lowe J. 2014. Closing the cohesin ring: structure and function of its
Smc3-kleisin interface. *Science* **346**:963–7. doi:10.1126/science.1256917

Gruber S, Arumugam P, Katou Y, Kuglitsch D, Helmhart W, Shirahige K, Nasmyth K. 2006.
Evidence that loading of cohesin onto chromosomes involves opening of its SMC
hinge. *Cell* **127**:523–537. doi:10.1016/j.cell.2006.08.048

Haering CH, Farcas A-M, Arumugam P, Metson J, Nasmyth K. 2008. The cohesin ring
concatenates sister DNA molecules. *Nature* **454**:297–301. doi:10.1038/nature07098

Haering CH, Löwe J, Hochwagen A, Nasmyth K. 2002. Molecular architecture of SMC
proteins and the yeast cohesin complex. *Mol Cell* **9**:773–788. doi:10.1016/s1097-
2765(02)00515-4

Haering CH, Schoffnegger D, Nishino T, Helmhart W, Nasmyth K, Löwe J. 2004. Structure
and stability of cohesin's Smc1-kleisin interaction. *Mol Cell* **15**:951–964.
doi:10.1016/j.molcel.2004.08.030

Hara K, Zheng G, Qu Q, Liu H, Ouyang Z, Chen Z, Tomchick DR, Yu H. 2014. Structure of
cohesin subcomplex pinpoints direct shugoshin-Wapl antagonism in centromeric
cohesion. *Nat Struct Mol Biol* **21**:864–70. doi:10.1038/nsmb.2880

Hartman T, Stead K, Koshland D, Guacci V. 2000. Pds5p is an essential chromosomal
protein required for both sister chromatid cohesion and condensation in
Saccharomyces cerevisiae. *J Cell Biol* **151**:613–26. doi:10.1083/jcb.151.3.613

Hassler M, Shaltiel IA, Kschonsak M, Simon B, Merkel F, Thärichen L, Bailey HJ, Macošek
J, Bravo S, Metz J, Hennig J, Haering CH. 2019. Structural Basis of an Asymmetric
Condensin ATPase Cycle. *Mol Cell* **74**:1175-1188.e9.
doi:10.1016/j.molcel.2019.03.037

Higashi TL, Eickhoff P, Sousa JS, Locke J, Nans A, Flynn HR, Snijders AP, Papageorgiou
G, O'Reilly N, Chen ZA, O'Reilly FJ, Rappsilber J, Costa A, Uhlmann F. 2020. A
Structure-Based Mechanism for DNA Entry into the Cohesin Ring. *Mol Cell* **79**:917-
933.e9. doi:10.1016/j.molcel.2020.07.013

Higashi TL, Ikeda M, Tanaka H, Nakagawa T, Bando M, Shirahige K, Kubota Y, Takisawa
H, Masukata H, Takahashi TS. 2012. The prereplication complex recruits XEco2 to
chromatin to promote cohesin acetylation in Xenopus egg extracts. *Curr Biol* **22**:977–
88. doi:10.1016/j.cub.2012.04.013

Hill L, Ebert A, Jaritz M, Wutz G, Nagasaka K, Tagoh H, Kostanova-Poliakova D, Schindler
K, Sun Q, Bonelt P, Fischer M, Peters JM, Busslinger M. 2020. Wapl repression by
Pax5 promotes V gene recombination by Igh loop extrusion. *Nature* **584**:142–147.
doi:10.1038/s41586-020-2454-y

Ivanov MP, Ladurner R, Poser I, Beveridge R, Rampler E, Hudecz O, Novatchkova M,
Heriche JK, Wutz G, van der Lelij P, Kreidl E, Hutchins JR, Axelsson-Ekker H,

Ellenberg J, Hyman AA, Mechtler K, Peters JM. 2018. The replicative helicase MCM
recruits cohesin acetyltransferase ESCO2 to mediate centromeric sister chromatid
cohesion. *EMBO J* **37**. doi:10.15252/embj.201797150
Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K,
Bates R, Žídek A, Potapenko A, Bridgland A, Meyer C, Kohl SAA, Ballard AJ,
Cowie A, Romera-Paredes B, Nikolov S, Jain R, Adler J, Back T, Petersen S, Reiman
D, Clancy E, Zielinski M, Steinegger M, Pacholska M, Berghammer T, Bodenstein S,
Silver D, Vinyals O, Senior AW, Kavukcuoglu K, Kohli P, Hassabis D. 2021. Highly
accurate protein structure prediction with AlphaFold. *Nature* **596**:583–589.
doi:10.1038/s41586-021-03819-2
Kikuchi S, Borek DM, Otwinowski Z, Tomchick DR, Yu H. 2016. Crystal structure of the
cohesin loader Scc2 and insight into cohesinopathy. *Proc Natl Acad Sci U S A*
113:12444–12449. doi:10.1073/pnas.1611333113
Kim Y, Shi Z, Zhang H, Finkelstein IJ, Yu H. 2019. Human cohesin compacts DNA by loop
extrusion. *Science* **366**:1345–1349. doi:10.1126/science.aaz4475
Klein F, Mahr P, Galova M, Buonomo SB, Michaelis C, Nairz K, Nasmyth K. 1999. A
central role for cohesins in sister chromatid cohesion, formation of axial elements,
and recombination during yeast meiosis. *Cell* **98**:91–103. doi:10.1016/S0092-
8674(00)80609-1
Kueng S, Hegemann B, Peters BH, Lipp JJ, Schleiffer A, Mechtler K, Peters JM. 2006. Wapl
controls the dynamic association of cohesin with chromatin. *Cell* **127**:955–67.
doi:10.1016/j.cell.2006.09.040
Ladurner R, Kreidl E, Ivanov MP, Ekker H, Idarraga-Amado MH, Busslinger GA, Wutz G,
Cisneros DA, Peters JM. 2016. Sororin actively maintains sister chromatid cohesion.
EMBO J **35**:635–53. doi:10.15252/embj.201592532
Lee B-G, Merkel F, Allegretti M, Hassler M, Cawood C, Lecomte L, O'Reilly FJ, Sinn LR,
Gutierrez-Escribano P, Kschonsak M, Bravo S, Nakane T, Rappsilber J, Aragon L,
Beck M, Löwe J, Haering CH. 2020. Cryo-EM structures of holo condensin reveal a
subunit flip-flop mechanism. *Nat Struct Mol Biol* **27**:743–751. doi:10.1038/s41594-
020-0457-x
Lee B-G, Rhodes J, Löwe J. 2022. Clamping of DNA shuts the condensin neck gate. *Proc
Natl Acad Sci U S A* **119**:e2120006119. doi:10.1073/pnas.2120006119
Lee BG, Roig MB, Jansma M, Petela N, Metson J, Nasmyth K, Lowe J. 2016. Crystal
Structure of the Cohesin Gatekeeper Pds5 and in Complex with Kleisin Scc1. *Cell
Rep* **14**:2108–2115. doi:10.1016/j.celrep.2016.02.020
Li Y, Haarhuis JHI, Sedeño Cacciatore Á, Oldenkamp R, van Ruiten MS, Willems L,
Teunissen H, Muir KW, de Wit E, Rowland BD, Panne D. 2020. The structural basis
for cohesin-CTCF-anchored loops. *Nature* **578**:472–476. doi:10.1038/s41586-019-
1910-z
Li Y, Muir KW, Bowler MW, Metz J, Haering CH, Panne D. 2018. Structural basis for Scc3-
dependent cohesin recruitment to chromatin. *Elife* **7**. doi:10.7554/eLife.38356
Liu H, Rankin S, Yu H. 2013. Phosphorylation-enabled binding of SGO1-PP2A to cohesin
protects sororin and centromeric cohesion during mitosis. *Nat Cell Biol* **15**:40–9.
doi:10.1038/ncb2637
Minamino M, Bouchoux C, Canal B, Diffley JFX, Uhlmann F. 2023. A replication fork
determinant for the establishment of sister chromatid cohesion. *Cell* **186**:837-849 e11.
doi:10.1016/j.cell.2022.12.044
Mitter M, Gasser C, Takacs Z, Langer CCH, Tang W, Jessberger G, Beales CT, Neuner E,
Ameres SL, Peters JM, Goloborodko A, Micura R, Gerlich DW. 2020. Conformation

of sister chromatids in the replicated human genome. *Nature* **586**:139–144.
doi:10.1038/s41586-020-2744-4

Moldovan GL, Pfander B, Jentsch S. 2006. PCNA controls establishment of sister chromatid
cohesion during S phase. *Mol Cell* **23**:723–32. doi:10.1016/j.molcel.2006.07.007

Mota IP, Galova M, Schleiffer A, Nguyen T-T, Kovacikova I, Nishiyama T, Gregan J, Peters
J-M, Schlögelhofer P. 2022. SORORIN is an evolutionary conserved antagonist of
WAPL. doi:10.1101/2022.10.24.513534

Nishiyama T, Ladurner R, Schmitz J, Kreidl E, Schleiffer A, Bhaskara V, Bando M,
Shirahige K, Hyman AA, Mechtler K, Peters JM. 2010. Sororin mediates sister
chromatid cohesion by antagonizing Wapl. *Cell* **143**:737–49.
doi:10.1016/j.cell.2010.10.031

Oliveira RA, Hamilton RS, Pauli A, Davis I, Nasmyth K. 2010. Cohesin cleavage and Cdk
inhibition trigger formation of daughter nuclei. *Nat Cell Biol* **12**:185–92.
doi:10.1038/ncb2018

Ouyang Z, Zheng G, Song J, Borek DM, Otwinowski Z, Brautigam CA, Tomchick DR,
Rankin S, Yu H. 2013. Structure of the human cohesin inhibitor Wapl. *Proc Natl*
Acad Sci U S A **110**:11355–60. doi:10.1073/pnas.1304594110

Ouyang Z, Zheng G, Tomchick DR, Luo X, Yu H. 2016. Structural Basis and IP6
Requirement for Pds5-Dependent Cohesin Dynamics. *Mol Cell* **62**:248–259.
doi:10.1016/j.molcel.2016.02.033

Panizza S, Tanaka T, Hochwagen A, Eisenhaber F, Nasmyth K. 2000. Pds5 cooperates with
cohesin in maintaining sister chromatid cohesion. *Curr Biol* **10**:1557–64.
doi:10.1016/s0960-9822(00)00854-x

Parelho V, Hadjur S, Spivakov M, Leleu M, Sauer S, Gregson HC, Jarmuz A, Canzonetta C,
Webster Z, Nesterova T, Cobb BS, Yokomori K, Dillon N, Aragon L, Fisher AG,
Merkenschlager M. 2008. Cohesins functionally associate with CTCF on mammalian
chromosome arms. *Cell* **132**:422–433. doi:10.1016/j.cell.2008.01.011

Petela NJ, Gligoris TG, Metson J, Lee B-G, Voulgaris M, Hu B, Kikuchi S, Chapard C, Chen
W, Rajendra E, Srinivisan M, Yu H, Löwe J, Nasmyth KA. 2018. Scc2 Is a Potent
Activator of Cohesin's ATPase that Promotes Loading by Binding Scc1 without Pds5.
Mol Cell **70**:1134-1148.e7. doi:10.1016/j.molcel.2018.05.022

Petela NJ, Gonzalez Llamazares A, Dixon S, Hu B, Lee BG, Metson J, Seo H, Ferrer-
Harding A, Voulgaris M, Gligoris T, Collier J, Oh BH, Lowe J, Nasmyth KA. 2021.
Folding of cohesin's coiled coil is important for Scc2/4-induced association with
chromosomes. *Elife* **10**. doi:10.7554/eLife.67268

Rankin S, Ayad NG, Kirschner MW. 2005. Sororin, a substrate of the anaphase-promoting
complex, is required for sister chromatid cohesion in vertebrates. *Mol Cell* **18**:185–
200. doi:10.1016/j.molcel.2005.03.017

Rao SSP, Huang SC, Glenn St Hilaire B, Engreitz JM, Perez EM, Kieffer-Kwon KR,
Sanborn AL, Johnstone SE, Bascom GD, Bochkov ID, Huang X, Shamim MS, Shin J,
Turner D, Ye Z, Omer AD, Robinson JT, Schlick T, Bernstein BE, Casellas R, Lander
ES, Aiden EL. 2017. Cohesin Loss Eliminates All Loop Domains. *Cell* **171**:305-320
e24. doi:10.1016/j.cell.2017.09.026

Roig MB, Lowe J, Chan KL, Beckouet F, Metson J, Nasmyth K. 2014. Structure and
function of cohesin's Scc3/SA regulatory subunit. *FEBS Lett* **588**:3692–702.
doi:10.1016/j.febslet.2014.08.015

Rolef Ben-Shahar T, Heeger S, Lehane C, East P, Flynn H, Skehel M, Uhlmann F. 2008.
Eco1-dependent cohesin acetylation during establishment of sister chromatid
cohesion. *Science* **321**:563–6. doi:10.1126/science.1157774

Rowland BD, Roig MB, Nishino T, Kurze A, Uluocak P, Mishra A, Beckouet F, Underwood
P, Metson J, Imre R, Mechtler K, Katis VL, Nasmyth K. 2009. Building sister
chromatid cohesion: smc3 acetylation counteracts an antiestablishment activity. *Mol*
Cell **33**:763–74. doi:10.1016/j.molcel.2009.02.028
Schleiffer A, Kaitna S, Maurer-Stroh S, Glotzer M, Nasmyth K, Eisenhaber F. 2003.
Kleisins: a superfamily of bacterial and eukaryotic SMC protein partners. *Mol Cell*
11:571–575. doi:10.1016/s1097-2765(03)00108-4
Shaltiel IA, Datta S, Lecomte L, Hassler M, Kschonsak M, Bravo S, Stober C, Ormanns J,
Eustermann S, Haering CH. 2022. A hold-and-feed mechanism drives directional
DNA loop extrusion by condensin. *Science* **376**:1087–1094.
doi:10.1126/science.abm4012
Shi Z, Gao H, Bai XC, Yu H. 2020. Cryo-EM structure of the human cohesin-NIPBL-DNA
complex. *Science* **368**:1454–1459. doi:10.1126/science.abb0981
Shintomi K, Hirano T. 2009. Releasing cohesin from chromosome arms in early mitosis:
opposing actions of Wapl-Pds5 and Sgo1. *Genes Dev* **23**:2224–36.
doi:10.1101/gad.1844309
Soh Y-M, Bürmann F, Shin H-C, Oda T, Jin KS, Toseland CP, Kim C, Lee H, Kim SJ, Kong
M-S, Durand-Diebold M-L, Kim Y-G, Kim HM, Lee NK, Sato M, Oh B-H, Gruber S.
2015. Molecular basis for SMC rod formation and its dissolution upon DNA binding.
Mol Cell **57**:290–303. doi:10.1016/j.molcel.2014.11.023
Srinivasan M, Petela NJ, Scheinost JC, Collier J, Voulgaris M, M BR, Beckouet F, Hu B,
Nasmyth KA. 2019. Scc2 counteracts a Wapl-independent mechanism that releases
cohesin from chromosomes during G1. *Elife* **8**. doi:10.7554/eLife.44736
Srinivasan M, Scheinost JC, Petela NJ, Gligoris TG, Wissler M, Ogushi S, Collier JE,
Voulgaris M, Kurze A, Chan K-L, Hu B, Costanzo V, Nasmyth KA. 2018. The
Cohesin Ring Uses Its Hinge to Organize DNA Using Non-topological as well as
Topological Mechanisms. *Cell* **173**:1508-1519.e18. doi:10.1016/j.cell.2018.04.015
Tedeschi A, Wutz G, Huet S, Jaritz M, Wuensche A, Schirghuber E, Davidson IF, Tang W,
Cisneros DA, Bhaskara V, Nishiyama T, Vaziri A, Wutz A, Ellenberg J, Peters JM.
2013. Wapl is an essential regulator of chromatin structure and chromosome
segregation. *Nature* **501**:564–8. doi:10.1038/nature12471
Toth A, Ciosk R, Uhlmann F, Galova M, Schleiffer A, Nasmyth K. 1999. Yeast cohesin
complex requires a conserved protein, Eco1p(Ctf7), to establish cohesion between
sister chromatids during DNA replication. *Genes Dev* **13**:320–33.
doi:10.1101/gad.13.3.320
Uhlmann F, Wernic D, Poupart MA, Koonin EV, Nasmyth K. 2000. Cleavage of cohesin by
the CD clan protease separin triggers anaphase in yeast. *Cell* **103**:375–86.
doi:10.1016/s0092-8674(00)00130-6
Unal E, Heidinger-Pauli JM, Kim W, Guacci V, Onn I, Gygi SP, Koshland DE. 2008. A
molecular determinant for the establishment of sister chromatid cohesion. *Science*
321:566–9. doi:10.1126/science.1157880
van Ruiten MS, van Gent D, Sedeno Cacciatore A, Fauster A, Willems L, Hekkelman ML,
Hoekman L, Altelaar M, Haarhuis JHI, Brummelkamp TR, de Wit E, Rowland BD.
2022. The cohesin acetylation cycle controls chromatin loop length through a PDS5A
brake mechanism. *Nat Struct Mol Biol* **29**:586–591. doi:10.1038/s41594-022-00773-z
Varadi M, Anyango S, Deshpande M, Nair S, Natassia C, Yordanova G, Yuan D, Stroe O,
Wood G, Laydon A, Žídek A, Green T, Tunyasuvunakool K, Petersen S, Jumper J,
Clancy E, Green R, Vora A, Lutfi M, Figurnov M, Cowie A, Hobbs N, Kohli P,
Kleywegt G, Birney E, Hassabis D, Velankar S. 2022. AlphaFold Protein Structure
Database: massively expanding the structural coverage of protein-sequence space

with high-accuracy models. *Nucleic Acids Res* **50**:D439–D444.
doi:10.1093/nar/gkab1061
Wendt KS, Yoshida K, Itoh T, Bando M, Koch B, Schirghuber E, Tsutsumi S, Nagae G,
Ishihara K, Mishiro T, Yahata K, Imamoto F, Aburatani H, Nakao M, Imamoto N,
Maeshima K, Shirahige K, Peters J-M. 2008. Cohesin mediates transcriptional
insulation by CCCTC-binding factor. *Nature* **451**:796–801. doi:10.1038/nature06634
Wu FM, Nguyen JV, Rankin S. 2011. A conserved motif at the C terminus of sororin is
required for sister chromatid cohesion. *J Biol Chem* **286**:3579–86.
doi:10.1074/jbc.M110.196758
Wutz G, Ladurner R, St Hilaire BG, Stocsits RR, Nagasaka K, Pignard B, Sanborn A, Tang
W, Varnai C, Ivanov MP, Schoenfelder S, van der Lelij P, Huang X, Durnberger G,
Roitinger E, Mechtler K, Davidson IF, Fraser P, Lieberman-Aiden E, Peters JM.
2020. ESCO1 and CTCF enable formation of long chromatin loops by protecting
cohesin(STAG1) from WAPL. *Elife* **9**. doi:10.7554/eLife.52091
Yamada T, Tahara E, Kanke M, Kuwata K, Nishiyama T. 2017. Drosophila Dalmatian
combines sororin and shugoshin roles in establishment and protection of cohesion.
EMBO J **36**:1513–1527. doi:10.15252/embj.201695607
Yatskevich S, Rhodes J, Nasmyth K. 2019. Organization of Chromosomal DNA by SMC
Complexes. *Annu Rev Genet* **53**:445–482. doi:10.1146/annurev-genet-112618-043633
Zhai Y, Cheng E, Wu H, Li N, Yung PY, Gao N, Tye BK. 2017. Open-ringed structure of the
Cdt1-Mcm2-7 complex as a precursor of the MCM double hexamer. *Nat Struct Mol*
Biol **24**:300–308. doi:10.1038/nsmb.3374
Zhang Y, Zhang X, Dai HQ, Hu H, Alt FW. 2022. The role of chromatin loop extrusion in
antibody diversification. *Nat Rev Immunol* **22**:550–566. doi:10.1038/s41577-022-
00679-3
1756