What AlphaFold tells us about cohesin's retention on
and release from chromosomes

Kim A. Nasmyth[1], Byung-Gil Lee[2, 3], Maurici B. Roig[1] and Jan Löwe[2]

[1] Department of Biochemistry, University of Oxford, Oxford OX1 3QU, UK;

[2] MRC Laboratory of Molecular Biology, Cambridge CB2 0QH, UK

[3]Current address: Department of Biochemistry, Lee Gil Ya Cancer and Diabetes Institute, Gachon University College of Medicine, Incheon 21999, Republic of Korea

ABSTRACT

Cohesin is a trimeric complex containing a pair of SMC proteins (Smc1 and Smc3) whose

ATPase domains at the end of long coiled coils (CC) are interconnected by Scc1. During

interphase, it organizes chromosomal DNA topology by extruding loops in a manner dependent

on Scc1's association with two large hook shaped proteins called SA (yeast: Scc3) and Nipbl

(Scc2). The latter's replacement by Pds5 recruits Wapl, which induces release from chromatin

via a process requiring dissociation of Scc1's N-terminal domain (NTD) from Smc3. If blocked

by Esco (Eco)-mediated Smc3 acetylation, cohesin containing Pds5 merely maintains pre-

existing loops, but a third fate occurs during DNA replication, when Pds5-containing cohesin

associates with Sororin and forms structures that hold sister DNAs together. How Wapl induces

and Sororin blocks release has hitherto remained mysterious. In the twenty years since their

discovery, not a single testable hypothesis has been proposed as to their role. Here, AlphaFold

2 (AF) three-dimensional protein structure predictions lead us to propose formation of a

quarternary complex between Wapl, SA, Pds5, and Scc1's NTD, in which the latter is

juxtaposed with (and subsequently sequestered by) a highly conserved cleft within Wapl's C-

terminal domain (CTD). AF also reveals how Scc1's dissociation from Smc3 arises from a

distortion of Smc3's CC induced by engagement of SMC ATPase domains, how Esco acetyl

transferases are recruited to Smc3 by Pds5, and how Sororin prevents release by binding to the

Smc3/Scc1 interface. Our hypotheses explain the phenotypes of numerous existing mutations

and are highly testable.

INTRODUCTION

Cohesin is a multi-subunit complex responsible for determining much of the topology of chromosomal DNA during interphase as well as sister chromatid cohesion (Yatskevich et al., 2019). At its heart are a pair of rod-shaped Structural Maintenance of Chromosomes (SMC) proteins, Smc1 and Smc3, that have a dimerization domain called the "hinge" at one end and an ABC-like ATPase domain at the other, separated by 50 nm long intra-molecular anti-parallel coiled coils. These are interrupted in two places, at an "elbow", 16 nm from the hinge, around which the coiled coils fold onto each other (Bürmann et al., 2019), and a disruption of the coiled coil called the "joint" (Diebold-Durand et al., 2017), about 6.5 nm from the ATPase domains (Fig. 1).

In the presence of ATP, the Smc1 and Smc3 ATPase domains associate with each other, thereby sandwiching a pair of ATP molecules between them (Shi et al., 2020). Crucially, they are also inter-connected by a kleisin subunit (Schleiffer et al., 2003), called Scc1, whose N- and C-terminal domains bind to the "base" and "neck" of the Smc1 and Smc3 ATPases respectively, creating a huge tripartite SMC-kleisin ring, even in the absence of ATP (Chapard et al., 2019; Haering et al., 2002) (Fig. 1). Cohesin's various activities depend on the association with the largely unstructured sections of the kleisin polypeptide connecting its N- and C-terminal domains, of three large hook-shaped HEAT repeat containing regulatory subunits called HAWKs (HEAT repeat proteins Associated With Kleisins). Only two of these are thought to be bound at any one time, Scc3 (human: SA) with either Pds5 or Scc2 (human: Nipbl). Complexes containing Scc2/Nipbl are active as DNA dependent ATPases, while those containing Pds5 are not (Petela et al., 2018).

In addition to holding sister chromatids together in post-replicative cells, cohesin is capable of extruding long loops of DNA (Davidson et al., 2019; Kim et al., 2019). This loop extruding (LE) activity occurs throughout interphase in mitotic cells and during prophase in meiosis when it is essential for formation of the synaptonemal complex (Klein et al., 1999). Importantly, LE and sister chromatid cohesion are conferred by distinct populations of cohesin, as LE requires the continuous presence of Scc2/Nipbl (Mitter et al., 2020), whereas cohesion does not (Ciosk et al., 2000; Srinivasan et al., 2019). LE facilitates VDJ recombination (Zhang et al., 2022), as well as the interaction of enhancers with distant promoters and is regulated throughout the genome by the site-specific DNA binding protein CTCF (Fudenberg et al., 2016; Li et al., 2020).

Cohesin's initial association with chromosomes as well as active loop extrusion (LE) are

mediated by Smc-kleisin rings occupied by Scc2/Nipbl and Scc3/SA (Fig. 2). LE is halted at

asymmetric CTCF sites, but only when they face in one direction (Fudenberg et al., 2016). This

arrest depends on modification by yeast Eco1 (Bastie et al., 2022) or human Esco1 and 2 (van

Ruiten et al., 2022; Wutz et al., 2020) acetyl-transferases of a pair of lysine residues (yeast

Smc3K112 and K113; human Smc3K105 and K106) on the ATPase domain of Smc3 (Unal et

al., 2008), an event that depends on replacement of Scc2/Nipbl by Pds5 (Chan et al., 2013)

(Fig. 2). Unlike Scc2/Nipbl, which is required for LE and for building but not maintaining

cohesion, SA/Scc3 is essential for all of cohesin's activities, namely loading onto chromatin

and loop extrusion, as well as the establishment and maintenance of cohesion (Davidson et al.,

2019; Roig et al., 2014; Toth et al., 1999). Unrestricted LE between convergent CTCF sites,

but not past them, ensures that any sequence between convergent sites is more likely to be

associated with other sequences within this interval than with those beyond these sites, giving

rise to the phenomenon of Topologically Associated Domains (TADs) in HiC genome contact

maps (Rao et al., 2017). The inhibition of cohesin's LE by CTCF is presumed to depend on

Nipbl's replacement by Pds5, as well as acetylation of Smc3 by Esco1 and is accompanied by

cohesin's association with CTCF sites (Parelho et al., 2008; Wendt et al., 2008).

Figure 1. *Overview of cohesin, its subunits and regulatory proteins. Top left: cohesin's*

trimeric Smc1/Smc3/Scc1 ring. The approximate HAWK binding sites on the kleisin Scc1 are

indicated. Top right: cohesin in the DNA-clamped and folded state (hybrid model combining

PDBs 6WG3, 6ZZ6 and 7OGT). Bottom: cohesin's HAWKs and regulatory proteins

(AlphaFold 2 [AF] predictions).

Figure 2. *Cohesin's trimeric ring (top left), represented throughout the rest of the figure as a*

triangle, associates with DNA loops when occupied by SA/Scc3 and Nipbl/Scc2. Nipbl/Scc2

stimulates ATP hydrolysis, leading to DNA translocation and loop extrusion. This is halted by

Nipbl's replacement by Pds5, creating a complex that can bind Wapl and detach from

chromatin, in a process that depends on a pair of lysine residues (yeast: K112 and K113;

humans: K105 and K106) on Smc3's ATPase head, whose acetylation by Eco1 (yeast) or

Esco1/2 in vertebrates blocks release. Acetylated complexes containing Pds5 either form stable

loops between CTCF sites or, during DNA replication, generate cohesion between sister DNAs,

whose maintenance (in animal and plant cells) requires Sororin (as well as continued

acetylation) to prevent Wapl-mediated release.

Post-replicative cells have three major types of cohesin complexes: those holding sisters
together (sister chromatid cohesion), those actively extruding loops, and those whose LE has
been arrested by CTCF and which hold together convergent CTCF sites at TAD boundaries.
Crucially, all three are susceptible to the action of a fourth regulatory subunit called Wapl
(Chan et al., 2012; Gandhi et al., 2006; Kueng et al., 2006), which induces cohesin's release
from chromatin at a certain frequency, albeit only when Nipbl/Scc2 has been replaced by Pds5
and when Smc3(K105) and K106 are unmodified (Fig. 2). The increased processivity of
cohesin's LE upon Wapl's artificial depletion induces formation of chromatid-like structures
during interphase (Tedeschi et al., 2013), a process that occurs naturally during meiosis (Klein
et al., 1999). Moreover, Wapl's downregulation during B cell development is essential for
recombination between DJ junctions and distal V genes (Hill et al., 2020), which requires
highly processive LE, mediated by cohesin to bring them together in the correct orientation so
that they can be joined by the RAG recombinase.

Because its maintenance does not require Scc2/Nipbl, sister chromatid cohesion is thought to
be a passive structure that does not require continual ATP hydrolysis. Once generated during
replication, cohesion must also be shielded from Wapl. This protection depends on Smc3
acetylation, and in animal and plant cells on cohesin's association during S phase with a protein
called Sororin (Nishiyama et al., 2010; Rankin et al., 2005). When cells enter M phase,
Sororin's phosphorylation by Cdk1 abrogates its cohesion protection and Wapl thereupon
destroys most cohesion along chromosome arms but not at centromeres, where PP2A
associated with Shugoshins counteracts Cdk1 (Liu et al., 2013). Wapl's destruction of all
cohesion along chromosome arms when cells are arrested in mitosis by spindle poisons is
responsible for the canonical image of mitotic chromosomes, with cohesion persisting solely
at centromeres.

Cohesion is thought to be mediated through the topological entrapment of sister DNAs by
individual tripartite Smc-kleisin rings (Haering et al., 2008; Srinivasan et al., 2018), either
through co-entrapment of sister DNAs inside a single ring, or alternatively through formation
of a Borromean ring that links sister DNAs (best thought of as infinitively large rings) and
cohesin rings in a topology that does not involve catenation of any one by the other (Collier et
al., 2020). In either case, ring opening through Scc1's cleavage by Separase destroys the
topological entanglement, thereby destroying cohesion and triggering the metaphase to
anaphase transition (Oliveira et al., 2010; Uhlmann et al., 2000). Unlike Separase, Wapl
removes cohesin and destroys cohesion by opening the cohesin ring in a manner that does not

involve proteolysis. Importantly, the finding that Wapl-mediated release is blocked by fusing

Smc3's C-terminus to Scc1's N-terminus suggests that release depends on dissociation of

Scc1's NTD from Smc3 (Beckouet et al., 2016; Chan et al., 2012), thereby creating a gate

through which DNAs may pass. If sisters are held together by their co-entrapment within a

cohesin ring, then the latter must also possess an entry gate that differs from the exit gate and

probably resides at the hinge interface (Collier and Nasmyth, 2022), because, unlike release,

fusion of Smc3 to Scc1 does not block cohesion establishment (Gruber et al., 2006; Srinivasan

et al., 2018). Less is known about the topology of cohesin's association with DNA during LE.

The observation that it is unaffected, at least in vitro, by fusing all three of the ring's interfaces

has led to the suggestion that LE does not require (single) passage of DNA through any of the

ring's interfaces (Davidson et al., 2019), but this does not exclude the possibility of DNAs

passing through cohesin rings in two directions (double passage).

X-ray crystallography and electron cryomicroscopy (cryo-EM) have revealed many details of

the topology of Smc1/Smc3 heterodimers (Fig. 1), how they dimerize via hinge domains

(Haering et al., 2002), how their coiled coils are folded around an elbow (Bürmann et al., 2019),

how their ATPase domains associate in the presence of ATP and bind the N- and C-terminal

domains of Scc1 (Gligoris et al., 2014; Haering et al., 2004), the structure of cohesin's HAWK

subunits (Hara et al., 2014; Kikuchi et al., 2016; Lee et al., 2016; Ouyang et al., 2016; Roig et

al., 2014) and how they bind DNA (Li et al., 2018), how Scc2/Nipbl clamps DNA on top of

engaged ATPase heads (Collier et al., 2020; Higashi et al., 2020; Shi et al., 2020), and lastly

the structure of Wapl (Chatterjee et al., 2013; Ouyang et al., 2013). Despite this progress, many

molecular aspects of cohesin's activities remain mysterious. These include how cohesin

actually extrudes DNA loops, how Scc2/Nipbl and Pds5 exchange with each other, how

cohesin's exit and entry gates are opened, how Wapl facilitates exit gate opening and thereby

induces cohesin's dissociation from chromosomes, how Esco/Eco acetyl transferases are

recruited to cohesin, how they recognize Smc3, and how this is facilitated by Pds5 and the

Mcm helicase, and how Sororin hinders exit gate opening and thereby thwarts Wapl.

We report here a complementary approach to in vitro biochemistry and experimental structural

biology, namely in silico 3D protein structure prediction by AlphaFold 2 (AF) (Evans et al.,

2021; Jumper et al., 2021), based on amino acid co-evolution analyses and deep learning

approaches to sequence/structure relationships, to predict structurally hitherto unreported

interactions between cohesin subunits. When performed with subunits from multiple

eukaryotic lineages and critically judged in the light of amino acid conservation (beyond that

used by AF), these predictions have furnished insights into several of the outstanding questions
posed above. It has revealed a possible role in opening cohesin's exit gate of interactions
between the Smc1 and Smc3 coiled coils in the vicinity of their joints when ATPases engage
in the absence of Scc2/Nipbl. AF also predicts how Eco/Esco acetyl transferases are recruited
to cohesin and thereby bind to K112 and K113 (human: K105 and K106) within Smc3's
ATPase domain. Crucially, it predicts that Wapl induces formation of a tripartite complex
between itself, SA/Scc3, and Pds5. If this occurs after Scc1's NTD has already dissociated
from Smc3, then the manner in which Scc1 associates with Pds5 places the kleisin's NTD close
to a highly conserved cleft within Wapl's CTD, raising the possibility that the latter acts by
sequestering Scc1's N-terminal 20 amino acids. Importantly, the tripartite SA/Wapl/Pds5
complex is predicted to form not only in animal but also in plant and fungal lineages, suggesting
it has deep evolutionary origins. Likewise, the prediction that Sororin's essential C-terminal
domain binds to the interface between Smc3 and Scc1's NTD provides a mechanism by which
it protects the latter from Wapl.

RESULTS

Closing the Smc1 and Smc3 coiled coil joints plays a part in the dissociation of Scc1's

NTD from Smc3's neck.

Wapl is thought to induce cohesin's release from chromosomes by facilitating the passage of

DNAs through the Smc3/Scc1 interface. It could do this either by inducing dissociation of the

interface, by holding the gate open once Scc1 has already dissociated, or by a combination of

both mechanisms. If it merely held the gate open, then there must exist a mechanism capable

of opening it that is Wapl-independent. Evidence for this comes from the observation that

inactivation of Scc2 during late G1 triggers cohesin's dissociation from chromosomes via a

mechanism that is blocked by fusion of Smc3 to Scc1 or by mutations such as *SMC1(L1129V)*

or *SMC1(D1164E)* that suppress *eco1Δ* lethality (Srinivasan et al., 2019). Crucially, this

release is independent of Wapl.

A clue as to a mechanism for Wapl-independent release comes from observations on the related

condensin complex. In the absence of DNA, ATP induces dissociation from Smc2 (equivalent

to Smc3) of Brn1's NTD (equivalent to Scc1). Because it also occurs in the presence of ADP

and BeF$_3$, Brn1's dissociation is presumably induced by the engagement of Smc2 and Smc4

ATPase heads (Hassler et al., 2019; Lee et al., 2022). This raises two related questions. Does a

similar process occur in cohesin and if so, how might head engagement induce dissociation?

An important clue as to the possible mechanism stems from cryo-EM structures of both cohesin

and condensin in the presence of DNA and non-hydrolyzable ATP analogues, which reveal

that DNA is "clamped" on top of engaged ATPase heads by Scc2 and Ycs4, respectively.

Importantly, in both cases, Smc coiled coils are splayed open and the N-terminal domains of

Brn1 and Scc1 remain associated with the necks of Smc2 and Smc3 respectively (Collier et al.,

2020; Lee et al., 2022; Shaltiel et al., 2022). This raises the question as to why DNA's presence

leads to retention of the Smc-kleisin interaction. Is it due to DNA binding per se or to an event

that accompanies that binding, namely opening of the Smc coiled coils. Clearly, head

engagement does not per se induce dissociation, at least when DNA is bound on top of them,

as in the clamped cryo-EM structures.

To address these questions, we explored the potential of structural predictions using AlphaFold

2 multimer (AF) (Evans et al., 2021; Jumper et al., 2021). Importantly, presumably due to the

co-evolution of amino acids within the interface of engaged Smc ATPase heads, AF routinely

attempts to engage them even though ATP is absent, which means that it is possible, to some

extent, to observe the effects of head engagement on the structure of cohesin's coiled coils

using AF (Fig. 3A, f1). As with all further figures, the data files (f1- f98), including atomic

coordinates   and   AF   output   files   can   be   found   at

https://doi.org/10.6084/m9.figshare.22567318.v2. To address whether head engagement

is compatible with association of the N- and C-terminal domains of Scc1 with Smc3 and Smc1

ATPase heads respectively, we asked AF to predict the structures of Smc1 and Smc3 heads

associated with coiled coils extending up to but not including their joints in the presence of a

truncated Scc1 polypeptide in which its conserved NTD and CTDs were joined by a GS

(glycine, serine) linker (Fig. 3B, f2). AF predicted with high confidence in five out of five

models heads whose engagement was very similar to that of the clamped cryo-EM structures

(Fig. 3C), as was the orientation of their coiled coils. Importantly, the N- and C-terminal

domains of Scc1 were invariably bound to the neck of Smc3 and the base of Smc1 respectively,

as observed in the clamped structures. As shown by cryo-EM and predicted by AF, head

engagement is therefore not per se incompatible with Scc1's association with Smc3.

Figure 3. A*, AlphaFold 2 multimer (AF) prediction of yeast Smc1 and Smc3 ATPase heads*
with associated coiled coils up to, but not including their joints. The actual predictions and
sequences   used   are   available   in   folder   f1   at
https://doi.org/10.6084/m9.figshare.22567318.v2*. **B**, As **A***, but including Scc1's N- and*
CTD connected by a GS linker (f2). ***C****, Cryo-EM structure of DNA clamped on top of engaged*
Smc1 and Smc3 heads by Scc2 (PDB 6ZZ6).

A very different arrangement of Smc1 and Smc3 coiled coils has been observed by cryo-EM
in the absence of ATP. The coiled coils are juxtaposed from their emergence from the hinge
all the way to their joints close to the ATPase domains, which is accompanied by folding of
both coiled coils around their elbows, thereby locating the hinge domain close to a segment of
Smc3's coiled coil just above its joint (Petela et al., 2021) (PDB 7OGT). To address whether
AF can reproduce this state, it was used to predict the structure of a yeast Smc1/Smc3
heterodimer containing all hinge and coiled coil sequences, but lacking both ATPase domains
(Fig. 4A, f3). This revealed, with high confidence in five out of five models, a structure
resembling that seen by cryo-EM (Fig. 4C), namely, with the Smc1 and Smc3 coiled coils
juxtaposed in parallel throughout their length and folded around their elbows placing the hinge
asymmetrically on top of the Smc3's coiled coil, just above its joint. Noticeably, in the AF
prediction, the latter was intimately inter-digitated with Smc1's joint, creating a compact
junction between the two (Fig. 4A). AF predicted an almost identical structure for the
equivalent parts of human Smc1/Smc3 heterodimers (Fig. 4B, f4). Importantly, the asymmetric
manner in which the Smc1 and Smc3 coiled coils emerge from the hinge and thereby associate
with each other, as soon as they do so, resembles that seen by cryo-EM as well as a crystal
structure of the condensin hinge associated with stretches of coiled coil (Lee et al., 2020; Soh
et al., 2015) (PDB 4RSI, 6YVU). This relationship between the hinge and its associated coils
has the appearance of a wrist, folded back on its associated arm. Interestingly, according to AF,
the folded wrist conformation is not predicted when the coiled coils associated with the hinge
were cut short at the elbow (Fig. 5A, f5). The shortened Smc1 and Smc3 coiled coils instead
emerge symmetrically with opened arms at an angle of 120° from each other, in a manner
resembling the crystal structure of the *T. maritima* hinge associated with short coiled coils (Fig.
5B) (Haering et al., 2002) (PDB 1GXL). This suggests that folding at the elbow as well as the
consequent packing of the hinge on top of the Smc3 coiled coil may have an important role in
organizing how the coiled coils emerge from the hinge. Crucially, the association of full-length

elbow-containing coiled coils and especially their joint junction depends on the hinge, as Smc1

and Smc3 coiled coils in which the hinge is replaced by either a GSx4 (Fig. 6A, f6) or GSx15

(Fig. 6B, f7) linker fail to associate in the predictions, even at their joints. Though the elbow

appears to be a feature that is autonomous, at least according to AF, coiled coil association,

including joint junction, is affected by a distant "organizing locus", namely the hinge in its

wrist conformation.

Figure 4. A*, AF prediction for yeast Smc1 and Smc3 lacking their ATPase domains (f3).* ***B****, As*

A, but human Smc1 and Smc3 (f4). ***C****, Cryo-EM structure of yeast Smc1 and Smc3 hinge and*

associated coiled coils, PDB 7OGT.

Figure 5*. **A**, AF prediction for yeast Smc1 and Smc3 hinge domains with short associated*
*coiled coils (f5). **B**, Crystal structure of T. maritima hinge associated with short coiled coils,*
PDB 1GXL.

Figure 6. *AF predictions for yeast Smc1 and Smc3 coiled coils including their elbow and joints, but joined by GSx4 (**A**, f6) and GSx15 (**B**, f7) linkers instead of their hinges.*

Though influenced by elbow and hinge, the joints nevertheless have an affinity for each other, as they form compact junctions when the coiled coils distal to the elbow are excluded and do so despite the lack of any extensive association between coiled coils away from the joint (Fig. 7A, f8). Indeed, AF predicted with high confidence a compact heterodimer formed between the joints of yeast Smc1 and Smc3, associated with only short sections of coiled coils (Fig. 7B, f9). A very similar prediction, albeit in only the top ranked model, was made for the equivalent human proteins (f10). In both yeast and humans, the predicted interactions between isolated joints were very similar to those within Smc1/Smc3 heterodimers lacking merely their ATPase domains (Fig. 4).

Figure 7. *AF predictions of Smc1 and Smc3 joints with short associated coiled coils.* ***A*** *(f8)*
and ***B*** *(f9), yeast.*

It is important to note that the junction between Smc1 and Smc3 joints was fully compatible
with the simultaneous association of Scc2 with the Smc3 joint. This was demonstrated by
running an AF prediction of Scc2's NTD either with an isolated Smc3 joint segment or with
an Smc1/Smc3 joint dimer (Fig. 8A, f12). Aligning the Smc3 moiety of the AF prediction with
the equivalent residues in the cryo-EM structure of DNAs clamped by Scc2 on top of engaged
Smc1/3 heads (Fig. 8B) (PDB 6ZZ6) shows the prediction to recapitulate the same Scc2:Smc3
binding mode. Because of this compatibility, competition with Scc2 is unlikely therefore to
have a role in joint disjunction (and thereby retention of Scc1's association with Smc3) upon
clamping.

Figure 8. A, AF prediction for yeast Smc1 and Smc3 joints with associated coiled coils together with Scc1 and Scc2's NTDs (f12). B, Cryo-EM structure of DNA clamped by Scc2 on top of engaged Smc1 and Smc3 ATPase, PDB 6ZZ6.

These predictions imply that dimerization at opposite ends of the Smc1 and Smc3 coiled coils have opposing influences on their organization. Engagement of the ATPase heads opens their associated coiled coils, while hinge dimerization has the opposite effect, forcing them together, including junction of their joints proximal to the ATPase heads. What happens then when both types of dimerization co-exist? Due to their size, AF predictions of full length Smc1/Smc3 heterodimers are close to limits of what is possible for current implementations of AlphaFold 2 multimer (Evans et al., 2021). Despite this, we obtained four models for yeast and five for human heterodimers. Four out of four models for the yeast complex predicted heterodimers whose coiled coils were folded and their joints interdigitated despite heads being fully engaged

(Fig. 9A and B, f13). In the human case, three out of five models (ranks 1, 2, and 5) had a

similar configuration but in one model (rank 3) joint junction was dominant over head

engagement (and the latter were disengaged), while in another head engagement was dominant

over joint junction and the latter were disjoined (Fig. 9C, f14). However, as one might expect,

coexistence of joint junction and head engagement is accompanied by distortions of the coiled

coils connecting joints and heads, and in some predictions this is associated with extreme

bending and/or actual unfolding of the alpha helices. The more extreme coiled coil distortions

observed in some models of the full length Smc1/Smc3 heterodimers models are absent when

the hinge and distal coiled coils are removed from the yeast complex. Thus, four out of five

(ranks 1, 2, 3, and 4) models of Smc1/3 ATPase heads associated with coiled coils extending

to but not beyond the elbow predict with high confidence junction of their joints co-existing

with fully engaged heads (Fig. 10A, f15).

Figure 9. *AF predictions for full length yeast (**A** and **B**, f13) and human (**C**, f14) Smc1/Smc3*

heterodimers.

Though some of the more baroque AF models may reflect unrealistic scenarios, the emerging

principle that head engagement and joint junction can co-exist but only do so if the intervening

coiled coils are distorted is consistent with a similar configuration in low resolution cryo-EM

structures of Smc1/Smc3 heterodimers in the presence of ATP (Petela et al., 2021). Indeed,

there is evidence that the state also occurs in vivo. By introducing cysteine pairs in the interface

between engaged Smc1/3 ATPases, it is possible to use the bifunctional thiol-specific

crosslinking reagent BMOE (bis-maleimidoethane) to detect their proximity in vivo. A

different pair of cysteines can be used to detect association between Smc1 and Smc3 coiled

coils in the vicinity of their joints. In an observation surprising at the time, it was observed that

BMOE induces appreciable double crosslinking (head and joint crosslinks) with a frequency

not dissimilar to the product of the frequency of the individual pairs, which indicates

simultaneous crosslinking (Chapard et al., 2019).

These observations lead us to propose that head engagement under conditions in which the

hinge reinforces a junction of Smc1/Smc3 joints, namely in the absence of DNA or Scc2, will

create distortions of the coiled coil (Smc3's neck) to which Scc1's NTD is normally associated

and that this distortion may cause its dissociation. A corollary is that maintaining the

association of Scc1's NTD with the Smc3 neck should prevent either joint junction or even

head engagement. Consistent with this hypothesis, the addition of Scc1's NTD (Fig. 10B, f16)

but not its CTD (Fig. 10C, f17) to the Smc1 and Smc3 ATPase heads associated with coiled

coils extending to the elbow eliminated the joint junction previously observed in four out of

five models. In this case, where joint junction is not reinforced by the hinge, AF clearly gave

preference to the Smc3-kleisin interaction over joint junction. Likewise, the top three ranked

models obtained with Scc1's NTD together with full length Smc1/Smc3 heterodimers

predicted engaged heads with Scc1's NTD associated with Smc3's neck but joints fully

disjoined (f18). Though the in silico prediction that association between Scc1's NTD and the

Smc3 neck is dominant over joint junction may not be an accurate reflection of nature, the

notion that joint junction while heads are engaged may be the driving force for opening

cohesin's exit gate nevertheless provides a plausible mechanism for this key event.

Importantly, the hypothesis explains why non-lethal mutations that compromise head

engagement (Elbatsh et al., 2016) abrogate releasing activity (RA). Crucially, it is eminently

testable, as it predicts that dissociation of Scc1's NTD from the Smc3 neck depends on joints

and probably also on distal coiled coils and their associated hinge.

Figure 10. *AF predictions of yeast Smc1 and Smc3 ATPases with coiled coils extending to the elbow but not beyond, without Scc1 (**A**, f15), with Scc1's NTD (**B**, f16) and with Scc1's CTD (**C**, f17).*

How does Wapl enhance exit gate opening?

If ATP-dependent dissociation of Scc1's NTD from the Smc3 neck is a feature intrinsic to Smc-kleisin trimers, then how might Wapl enhance this process? To address this, we used AF to explore how Wapl is first recruited to cohesin and second what effect it has on the configuration of cohesin's HAWK regulatory proteins and /or on the Smc3/Scc1 interface. Our starting point was existing knowledge based on biochemical and genetic assays. In vitro, Wapl is known to bind SA/Scc3 and this binding is abrogated by mutation of a highly conserved lysine residue (K404 in yeast; K330 in humans) within a highly conserved surface (known as the "CES", Conserved and Essential Surface) that sits underneath the pronounced "nose" of Scc3/SA. This binding must be of physiological importance as K to E mutations abrogate Wapl-dependent releasing activity in vivo, both in yeast (Beckouet et al., 2016) and in cultured human cells (Hara et al., 2014). Exactly which part of Wapl binds to the CES has never been fully established. It has been proposed that sequences within Wapl's unstructured N-terminal domain are responsible, either one or other of its FGF motifs (Shintomi and Hirano, 2009) or a different segment between residues 510 and 570, known as Wapl-M (Hara et al., 2014; Ouyang et al., 2013). As will become clear below, AF predictions imply that neither proposal

is correct. Meanwhile, a short YSR motif at Wapl's N-terminus binds a highly conserved loop

(consensus sequence AP[D/E]AP) between helices 2 and 3 within Pds5's NTD, which is also

bound by a YSR-like motif within the kinase Haspin (Ouyang et al., 2016). This locus within

Pds5's NTD was in fact first identified as being crucial for Wapl-mediated release through

mutations clustering at this location that suppress the lethality of *eco1* mutants in yeast

(Rowland et al., 2009). However, Wapl's N-terminal YSR has never been rigorously

demonstrated to be necessary for its releasing activity (RA). Most fungal Wapls also possess

at or near their N-termini a similar motif whose consensus is [M/Φ]xxYG[K/R]. Importantly,

its deletion in *S. cerevisiae* does not suppress the lethality of *eco1Δ* mutants (Fig. 11A),

implying that unlike Pds5's AP[D/E]AP loop to which it is thought to bind, the

[M/Φ]xxYG[K/R] motif is not essential for Wapl-mediated releasing activity. Some other

sequence must therefore bind the AP[D/E]AP loop in vivo. Interestingly, the

[M/Φ]xxYG[K/R] motif is lacking in *S. pombe*, though a different one (TYxxxR[T/S]ΦL),

which is much closer to Wapl's conserved CTD, is present and highly conserved in that

location among all fungi (Fig. 11B). Unlike the N-terminal motif, the more C-terminal

TYxxxR[T/S]ΦL motif is essential for releasing activity, as its deletion supresses the lethality

of *eco1Δ* mutants in *S. cerevisiae* (Fig. 11A). Thus, the picture that emerges is primarily one

of a loose association between Wapl and SA/Scc3 or Pds5 via short motifs within Wapl's

unstructured N-terminal domain. Crucially, none of these interactions help explain how Wapl's

more conserved CTD composed of eight HEAT repeats actually promotes release.

Figure 11. *A, Deletion of Wapl's N-terminal [M/Φ]xxYG[K/R] motif from S. cerevisiae (waplΔ4-9) does not suppress the lethality of eco1 mutations while deletion of its more C-terminal TYxxxR[T/S]ΦL motif (waplΔ208-215) does so. Both deletions were initially created within a version of the WAPL gene tagged with EGFP (KN18714) and their ability to suppress the lethality of eco1Δ determined by crossing to a waplΔ eco1Δ strain (strain KN16432) followed by tetrad analysis. B, A multiple amino acid sequence alignment of fungal Wapl sequences, showing the TYxxxR[T/S]ΦL motif on the left and the first highly conserved alpha helix within Wapl's CTD on the right. Yeast Wapl is the top sequence (W7PW2).*

We first used AF to explore the association between Wapl and Pds5. For yeast, this predicted

a single point of contact between a short Wapl motif ([F/W]xFLD), which is conserved in

numerous ascomycetes including *S. cerevisiae* (FNFLD- see Fig. 11A) and *S. pombe*

(WNFLD), and a trough situated between a pair of HEAT repeats located within Pds5's C-

terminal half, at the point where its longitudinal axis starts to curve (Fig. 12A, f19). Because

the very same trough is predicted also to bind FGF motifs from human Wapl (Fig. 13B, f20)

and Sororin (Fig. 13B, f21) as well as a separate motif within Esco1 (Fig.13B, f22), we will

refer to it as the Wapl Esco Sororin Trough or "WEST". Intriguingly, AF models of individual

proteins in the AlphaFold Protein Structure Database (Varadi et al., 2022) predict that the *S.*

cerevisiae and *S. pombe* FNFLD motifs of Wapl also bind a highly conserved cleft within Wapl

itself, at the C-terminal end of its CTD (Fig. 12B). However, given the choice between binding

Pds5 and itself, AF chooses the former. In humans, as already alluded to, AF predicted in five

out of five models an interaction between the central of three FGF motifs within Wapl's N-

terminal extension and the equivalent C-terminal trough within Pds5A and B (Fig. 13B, f20).

In contrast, AF predicted association between Wapl's N-terminal YSR motif and Pds5's N-

terminal APDAP loop in just one out of five models. In the case of human (Fig. 14, f23), but

not yeast Wapl (Fig. 12A, f19), AF predicted with high confidence in five out of five models a

rigid body association between its HEAT-repeat containing CTD and HEAT repeats at Pds5's

N-terminus, close to but independent of Pds5's APDAP loop, an interaction that proved pivotal

for our subsequent analyses (see below).

Though AF did not predict interaction of the *S. cerevisiae* TYxxxR[T/S]ΦL motif with full

length Pds5 (f19A), it predicted, in five out of five models, interaction of the equivalent motif

from *Neurospora crassa* with Pds5's APDAP loop (Fig. 12C, f19B). Subsequent predictions

using truncated versions of Pds5 and Wapl revealed that the *S. cerevisiae* TYxxxR[T/S]ΦL

motif also reproducibly bound to this site (f19C), as did (using a different Wapl truncation) its

[M/Φ]xxYG[K/R] motif (f19D), in this case in a manner resembling a crystal structure of the

equivalent complex from humans (PDB: 5HDT) (Ouyang et al., 2016). Given that both Pds5's

APDAP loop and Wapl's TYxxxR[T/S]ΦL motif are essential for releasing activity in yeast,

we presume that the interaction predicted by AF is crucial for their function, a feature whose

importance will become apparent later. It is reasonable to suppose that the Pds5 APDAP loop

also binds Wapl's N-terminal [M/Φ]xxYG[K/R] motif, albeit not at the stage when the

releasing reaction requires the APDAP loop's interaction with the TYxxxR[T/S]ΦL motif.

Figure 12. **A**, *AF prediction for yeast Wapl's association with Pds5 (f19A).* **B**, *AF prediction of yeast Wapl alone (AF-Q99359-F1, AlphaFold Protein Structure Database).* **C**, *AF predicts interaction between Wapl's TYxxxR[T/S]ΦL motif and Pds5's APDAP loop in Neurospora crassa (f19B).*

Figure 13. Interaction between a trough in Pds5's CTD called "WEST" and an FNFLD motif within the N-terminal part of yeast Wapl (**A left and right**, f19), FGF motifs within human Wapl (**B upper left and right**, f20) and D. rerio Sororin (**B lower right**, f21), and a highly conserved, partly helical, ILxLCEEIAGEIESD motif within human Esco1 (**B lower left**, f22).

Figure 14. *AF prediction for the interaction between Pds5A and the N-terminal half of Wapl's highly conserved CTD (f23). The same interaction was observed with both Pds5A and B.*

In contrast to yeast Pds5, AF did predict with high confidence in five out of five models a rigid body contact between a complex between yeast Scc3 and Scc1, and the first (α1) and second (α2) α helices within Wapl's CTD (Fig. 15A, f24), which are highly conserved (Fig. 11). The first of these binds across Scc3's HEAT repeats, including the helices at the base of its nose and those associated with the CES. Though the Wapl helix runs parallel to, and contacts, the Scc1 polypeptide, exclusion of the latter made little or no difference to AF's predictions. Since the axis of Wapl's first helix is nearly orthogonal to those of Scc3's HEAT repeats, so too are most of the HEAT repeat helices within Wapl's CTD and these therefore rise above Scc3 like a "tower" that follows the line of Scc3's nose, albeit leaning away from the latter's axis by 30°. Towards the top of this tower is a hydrophobic cleft (Chatterjee et al., 2013), lined by the most conserved α helix within fungal Wapls. This is the same cleft that in the absence of other proteins is occupied by Wapl's own N-terminal F/WxFLD motif (Fig. 12B), not only in fungi but also in a wide variety of eukaryotes (see below).

A very similar if not identical interaction was predicted between the human SA2:Scc1 complex

and a fragment of human Wapl containing its entire N-terminal extension as well as half of its

CTD (a prediction with full length SA2 and human Wapl would have been close or over AF's

size limit) (Fig. 15B, f25). In addition, a long motif (IFSGPKRSPTKAVYNARHWN) within

Wapl's unstructured N-terminal extension associated with a quite separate segment of SA2, in

this case binding parallel to one of its central HEAT repeats (Fig. 15B). Contact of this motif

with the SA2:Scc1 complex is extensive, running along the entire length of the relevant SA2

HEAT repeat, crossing and in so doing, contacting the Scc1 polypeptide and finally interacting

via its conserved tryptophan residue C-terminal HEAT repeats on the other side of SA2's hook.

This part of Wapl corresponds to the Wapl-M sequences (510-570) important for its association

with SA2:Scc1 and previously thought to bind the CES (Hara et al., 2014; Ouyang et al., 2013).

AF predicts a very different interaction, and besides which, it suggests that the CES is occupied

instead by Wapl's CTD (Fig. 15B). The AVYNARHWN part of the motif is conserved in

animals (consensus A[V/L]Yxx[R/K][H/-]W) and its mode of binding may explain why Wapl

binds to Scc1/Rad21 alone in vitro (Shintomi and Hirano, 2009). Significantly, AF failed to

predict any interaction between SA2:Scc1 and Wapl's FGF motifs, at least when the N-terminal

helices within Wapl's CTD are present and predicted to bind to SA2's CES.

The first alpha helix of Wapl's CTD (α1) is highly conserved (Fig. 17A, right; see also Fig.

11B), especially, but not exclusively (for reasons that will become apparent) residues facing

SA2. Crucially, its interaction with SA/Scc3 appears to be of physiological importance.

Scc3(K404) likely forms a conserved salt bridge with yeast Wapl's D272, while the equivalent

residue in SA2, K330 does likewise with human Wapl's D657. Remarkably, mutation of all

four residues (individually) has been shown to either abrogate or to greatly reduce RA mediated

by Wapl (Beckouet et al., 2016; Ouyang et al., 2013; Rowland et al., 2009). *S. cerevisiae*

Wapl(D272G) was in fact isolated as a spontaneous suppressor of *eco1* lethality in yeast. The

AF prediction for the human proteins explains the greater conservation of the surface of Wapl's

α1 facing SA2 as well as the more N-terminal V639, which also contacts SA2. Crucially, the

AF model explains why individual mutation of E653, D656, D657, or Y660 within the CTD's

α1 helix abrogates RA, as does mutation of L676 or K684 in its α2 helix (Ouyang et al., 2013).

All these residues are predicted by AF to contact SA2 (Fig. 15B, residues marked in yellow).

Figure 15. A, *AF prediction for the interaction between Scc3/Scc1 and the CTD of yeast Wapl*

(f24). ***B***, *AF prediction for the interaction between SA2/Scc1 and a fragment of human Wapl*

containing its N-terminal sequences as well as the N-terminal half of its helical CTD (f25).

Mutations that abolish RA are marked in yellow: E653, D656, D657, L676 and K684.

The prediction that in humans Wapl's CTD forms a complex with Pds5 as well as with

SA2:Scc1 raises the possibility that both interactions might occur simultaneously and that Wapl

517 would therefore glue all three proteins together in a ternary complex. This seems indeed to be

518 the case, as AF failed to predict any direct interaction between full length SA1 and Pds5A

519 (f26), but did so with high confidence in the top two ranked models created in the presence of

520 the part of Wapl's CTD that it predicts to bind both SA and Pds5 (Fig. 16, f27). The prediction

521 statistics indicated that the most accurate images of the ternary complex were obtained when

522 only the N-terminal domains of SA1 and Pds5A were included. In this case, AF predicted a

523 ternary complex with high confidence in five out of five models (Fig. 17A, f28). Using this

524 template, we screened for the formation of ternary SA:Wapl:Pds5 complexes in a variety of

525 eukaryotic lineages. Identical ternary complexes were predicted with high confidence in plants

526 (*Arabidopsis thaliana*), in insects (*Drosophila melanogaster*) and in a basal fungus, namely

527 the chytrid *Spizellomyces punctatus* (Fig. 17B, f29, f30, and f31). Crucially, formation of the

528 ternary complex explains the conservation and functional importance of E647 (Ouyang et al.,

529 2013) within human Wapl's α1 helix despite not facing SA2, as it contacts Pds5's NTD.

530 Likewise, contact with Pds5's NTD in the AF model explains abrogation of Wapl-mediated

531 RA by R695E as well as conservation of M692, R695, R731, and L732 in animals (Figure 17A,

532 right). We will refer to this as the canonical Pds5:SA/Scc3:Wapl rigid body ternary complex.

Figure 16. *AF prediction of a complex between human Pds5A, SA1, and a fragment of*

Wapl's CTD (f27). Pds5 and SA1 were not predicted to interact in the absence of Wapl (f26).

Figure 17. AF predictions for complexes between N-terminal fragments of Pds5 and SA/Scc3 and the N-terminal half of Wapl's CTD for humans (**A**, f28), A. thaliana (**B**, f29) D. melanogaster (f30), and S. punctatus (f31). Also shown is Wapl's conservation mapped on the human complex, **A**, right.

In contrast, despite the robust formation of binary complexes between Scc3 and Wapl orthologs, AF failed to predict the same type of ternary complexes with the equivalent protein fragments from *S. cerevisiae* (f32A). Further analysis of the way Wapl interacts with Pds5 in *S. punctatus* revealed a possible explanation. When the former's CTD was attached to N-terminal residues containing its TYxxxR[T/S]ΦL motif, AF predicted that Pds5's NTD formed a rigid body complex with the CTD, whose conformation was identical to that observed in the canonical Pds5:SA/Scc3:Wapl rigid body ternary complex (Fig. 17B), while its APDAP loop simultaneously bound Wapl's TYxxxR[T/S]ΦL motif (Fig. 18, f19E), in the manner observed in *S. cerevisiae* (f19C) and *N. crassa* (f19B). This raised the possibility that an interaction between Wapl's TYxxxR[T/S]ΦL motif and Pds5's APDAP loop might actually facilitate formation of ternary complexes and that this might explain why the TYxxxR[T/S]ΦL motif is

essential for releasing activity in *S. cerevisiae*. We therefore repeated the AF predictions for

the previously described *S. cerevisiae* Pds5, Scc3, and Wapl fragments (f32A) but in this case

included sequences containing Wapl's TYxxxR[T/S]ΦL motif. Remarkably, the top three

ranked models now predicted with high confidence a rigid body ternary complex (Fig.19A,

f32B), whose conformation closely resembles the canonical Pds5:SA/Scc3:Wapl rigid body

ternary complex observed in animals, plants, and basal fungi. Moreover, this was accompanied

by interaction between the TYxxxR[T/S]ΦL motif and Pds5's APDAP loop, as observed in *S.*

punctatus. The dramatic effect of including the TYxxxR[T/S]ΦL motif is best visualized by

comparing the PAEs of the top three ranked models with and without the motif (Fig. 19B).

AF also predicted with high confidence in two out of five models a very similar type of

canonical ternary complex in *N. crassa* (f33A) and the basidiomycete *Ustilago Maydis* (f33B).

As in *S. cerevisiae*, formation of the canonical Pds5:Scc3:Wapl ternary complex in *Neurospora*

was also strictly dependent on inclusion of its TYxxxR[T/S]ΦL motif (compare f33A & f33C).

These observations not only suggest that formation of a canonical Pds5:SA/Scc3:Wapl rigid

body ternary complex is conserved in fungi as well as animals and plants, but also that its

formation may be an essential feature of releasing activity, because deleting the

TYxxxR[T/S]ΦL motif necessary for its in silico prediction in *S. cerevisiae* (f32) abolishes

releasing activity in vivo (Fig. 11B). It is remarkable that the TYxxxR[T/S]ΦL motif is not

only highly conserved among fungi but that it is connected to α1 in Wapl's CTD by a short

polypeptide linker whose length (25 to 50 residues) does not vary greatly. In other words, the

shortness of this tether may be important for ensuring the particular type of interaction between

Wapl's CTD and Pds5's NTD observed in canonical Pds5:SA/Scc3:Wapl rigid body ternary

complexes.

Figure 18. AF prediction for the interaction between the N-terminal half of S. punctatus Pds5 and a fragment of Wapl containing its TYxxxRT/SΦL motif connected to its CTD (f19E). Both the motif and the CTD bind to Pds5

Fig. 19. A*, AF prediction for a S. cerevisiae Pds5:Scc3:Wapl rigid body ternary complex accompanied by interaction between Wapl's TYxxxR[T/S]ΦL motif and Pds5's APDAP loop (f32B). **B**, PAEs for the top three ranked models for S. cerevisiae's Pds5:Scc3:Wapl ternary complex with (top row, f32B)) and without (bottom row, f32A) Wapl's TYxxxR[T/S]ΦL motif. Note that the lower confidence complexes predicted by AF for those lacking Wapl's TYxxxR[T/S]ΦL motif are not of the canonical variety. In both **A** and **B**, chains A, B, and C correspond to Wapl, Pds5, and Scc3 respectively. Interaction between Wapl's TYxxxR[T/S]ΦL*

motif and Pds5 is represented by the thin blue lines on the left and upper squares of the top

*three maps in **B**.*

Though the structure of the ternary Pds5:SA/Scc3:Wapl RA complex in animals (Fig. 20, f35),

plants, and fungi is clearly intriguing and presumably important for RA, it does not per se offer

insight into the actual mechanism by which these three proteins facilitate cohesin's release.

This is especially true for the conserved cleft at the top (C-terminal end) of Wapl's CTD.

Despite differences in the precise patterns of conservation in animals, plants, and fungi, it is

remarkable that in all three lineages, motifs within Wapl's unstructured N-terminal extension

are predicted by the AlphaFold structure database to bind their own clefts (yeast shown in Fig.

12B). As will become apparent later, AF also predicts that Sororin's conserved FGF motif

(which is related to the yeast Wapl's FNF) also binds the cleft. The fact that Sororin is an

antagonist of Wapl suggests that the binding by N-terminal Wapl sequences represents a form

of auto-inhibition.

Figure 20. *Composite model of the human Pds5:SA1:Wapl:Scc1 quarternary complex (f35)*

created using by superimposing rank 1 models from f23, f25, and f28.

Potential insight into the function of the highly conserved C-terminal Wapl cleft, which is

predicted to be bound by motifs within Wapl's N-terminal sequences was revealed by a

separate set of predictions, namely investigation of how Scc1 binds to Pds5, either in the

presence or absence of Smc3's neck. Functional studies in yeast combined with X-ray

crystallography of a Pds5:Scc1 complex from *Lanchea thermotolerans* revealed association of

an extended Scc1 peptide, from *Lt* S125 to V141, along the axis of Pds5's spine, half way down

from its N terminus (Lee et al., 2016). This mode of interaction is essential as mutations

affecting the interaction, namely Scc1(V137K) and Pds5(Y458K) abrogate binding of the two

proteins in vivo and are lethal (Lee et al., 2016). Pds5 is required to maintain cohesion in post-

replicative cells (Hartman et al., 2000; Panizza et al., 2000), when Scc1's NTD is bound to

Smc3's neck (Gligoris et al., 2014), as well as being involved in RA (Rowland et al., 2009),

when Scc1's NTD must dissociate from Smc3, at least transiently (Beckouet et al., 2016; Chan

et al., 2012). With the initial aim of revealing the orientation of Pds5 connected to the Smc3

neck via Scc1, we used AF to predict how the N-terminal half of Pds5 interacts with the N-

terminal 160 amino acids of Scc1 in the presence of Smc3's ATPase and neck. Not surprisingly,

all five models predicted Scc1's simultaneous association with both the Smc3 neck and with

Pds5, in both cases in the manner expected from crystallographic studies (PDB 5F0N & 4UX3)

(Fig. 21, f36). More unexpected were the predictions for the sequences that connected the Scc1

S125-V141 interval to the NTD bound to Smc3. This revealed that Scc1's association with

Pds5's spine extended, in an N-terminal direction on both proteins, from the previously

characterized site of interaction almost but not quite to the C-terminal end of the long α3 helix

that forms a three-helical bundle with Smc3's coiled coil (Fig. 21). In all five models, the short

peptide linking α3 with the part of Scc1 that is associated with Pds5's spine adopts the same

conformation, which ensures that Pds5's NTD points towards the Smc3 ATPase head. Though

we have previously suggested that such an orientation might help to direct Eco1 to its K112

and K113 targets on the Smc3 ATPase (Lee et al., 2016), subsequent AF predictions (see

below) imply that this orientation would not in fact facilitate acetylation and the orientation

presumably has some other function. Given that the sequences linking Scc1$_N$'s α3 with its

canonical Pds5 binding motif are poorly conserved between fungi and animals, it was

remarkable that AF predicted a very similar structure with the orthologous human proteins

(Fig. 21, f37).

S. cerevisiae   Human

Figure 21. *AF predictions for the interaction of Scc1's N-terminal 160 residues with Smc3 and*

with an N-terminal fragment of Pds5 for both S. cerevisiae (f36) and human (f37).

Because RA depends on Wapl's association with Pds5, Scc1$_N$ is presumably bound to Pds5

upon its dissociation from Smc3 during a release reaction. We therefore used AF to predict

how Scc1$_N$ alone binds to Pds5. In the absence of Smc3, Smc3's coiled coil no longer poses a

constraint on how Scc1$_N$ binds with Pds5, and AF predicts that yeast Scc1's association now

extends to the C-terminal end of α3, whose interaction with the ends of a pair of Pds5's HEAT

repeats leads to a pronounced rotation, as well as levering backwards of the Scc1 NTD (Fig.

22, f38). This change is not accompanied by any alteration of its actual structure or in the

manner in which more C-terminal Scc1 sequences bind the Pds5 spine. In humans, α3's

association with the Pds5 spine was less affected by Smc3's removal but it was nevertheless

levered further backwards, becoming more orthogonal to the spine's axis (Fig. 22, f39).

Crucially, in both cases, as well as in *A. thaliana* (Fig. 22, f40), the C-terminal end of Scc1$_N$'s

α3 is anchored on the Pds5 spine and juts out from it at a defined angle. The change in the

manner of Scc1$_N$'s association with Pds5 upon dissociation from Smc3 might account for the

observation that Pds5 inhibits Scc1$_N$'s association with Smc3 in vitro, but does not affect its

association once bound (Ouyang et al., 2016). Given that Pds5 is intimately involved in RA

and that the latter requires dissociation of Scc1's NTD from Smc3, it is plausible to imagine

that upon its dissociation (presumably in response to head engagement and joint junction),

Scc1's NTD remains associated with Pds5 in the conformations observed in the above AF

models. Crucially, AF predicts that under these conditions, Scc1's NTD adopts a defined

position relative to Pds5's NTD that is similar in fungi, animals, and plants.

667   *S. cerevisiae*   Human   *A. thaliana*

Figure 22*. AF predictions for the interaction between Scc1's N-terminal 160 residues and*

Pds5 in S. cerevisiae (f38), humans (f39), and A. thaliana (f40).

Having realized that Pds5 organizes the spatial position of Scc1's NTD, we superimposed the

human Pds5:Scc1$_N$ model (Fig. 22 middle, f39) onto the canonical Pds5:SA/Scc3:Wapl rigid

body ternary complex (Fig. 23, f41). Remarkably, this revealed that the N-terminal end of

Scc1's NTD is placed, through its association with the spine of Pds5, in the vicinity of the

aforementioned cleft at the C-terminal end of Wapl's CTD. Supposition of the canonical

Pds5:Scc3:Wapl rigid body ternary complex from yeast (f32B) with AF predictions for Scc3's

association with Scc1 and full length Wapl (f24) and full length Pds5 associated with Scc1$_N$

(f38B) revealed a very similar looking quarternary complex (Fig. 31B).

As mentioned, in humans and most other metazoa, as well as in many fungi (Fig. 12B) and

plants, this cleft is predicted to be occupied by a motif (PSCLSVCNVT) within Wapl's N-

terminal sequences (AlphaFold Protein Structure Database: AF-Q7Z5K2-F1). The revelation

that Scc1's NTD is positioned close to this cleft in a putative Pds5:Scc3:Wapl:Scc1 complex

suggested that the role of this site might be to sequester Scc1's NTD following its dissociation

from Smc3's neck. Testing this hypothesis was not possible using a full quarternary complex,

due to its size. However, predictions of Wapl's position relative to Pds5's NTD are in fact

similar whether or not Wapl's CTD is also bound to the SA subunit and we therefore used AF

to predict the structure of a complex containing Wapl's CTD, the first 150 residues of

Scc1(Scc1$_N$), and the N-terminal straight section of Pds5 whose spine binds Scc1. All five

models predicted with high confidence a ternary complex between Wapl, Pds5, and Scc1$_N$ (Fig.

24, f42). In three of the five, including ranks 1, 2, and 5, Scc1's NTD was placed in close

vicinity of the Wapl cleft. The most N-terminal (α-1) of its four short helices (α-1, α0, α1, and

α2), normally packed on top of the α3 N-terminal helix, dissociates from α3 and binds in the

cleft (f42). However, these peripheral associations between Scc1$_N$'s α-1 and Wapl were not

predicted with high confidence (f42), did not look convincing, and more seriously, differed in

their details between models.

Figure 23. *A putative quarternary complex between human Scc1ₙ Pds5, SA1, and Wapl,*
created by super-imposing f23, f25, f28, and f39. The Scc1's NTD points towards, but clashes
with the highly conserved cleft in Wapl's CTD that is otherwise predicted to bind a motif
PSCLSVCNVT (conserved among metazoa) located in Wapl's N-terminal sequences (see Fig.
26B).

Figure 24. *AF's predictions for ternary complexes (models ranks1 and 4) containing Scc1$_N$, Pds5's NTD, and Wapl's CTD (f42).*

To test whether the positioning and hence orientation of Scc1's NTD through its association with Pds5 hinders its association with Wapl's CTD, we asked how the latter interacts with Scc1's NTD alone, i.e. when not constrained in space through its association with Pds5 within the rigid body ternary complex. In this case, AF predicts with high confidence and in four out of five models (Fig. 25, f34A) that Scc1's N-terminal helix (α-1) binds the cleft in Wapl's CTD that otherwise binds Wapl's own SCLSVCNVT peptide (Fig. 26B), while its second helix (α0), lying at 90° to α-1, is sandwiched between two long antiparallel helices that form a pronounced nose at the C-terminal end of Wapl and a second pair that cross over at 45°. AF predicts similar interactions for *D. melanogaster* (f34B), *C. elegans* (f34C), and for *A. thaliana* (f34D). Moreover, a slightly different but clearly related interaction was also predicted for *N. crassa* (f34E). Remarkably, only modest changes in the conformation of Scc1's NTD are required to accommodate binding of its α-1 and α0 helices to Wapl's CTD in this manner, principally a slight bending of α3 to avoid a clash with the pair of Wapl CTD helices that cross its nose. As a consequence, α3 lies on top of and almost parallel with Wapl's nose and this kind of

alignment is presumably why AF failed to predict Wapl's interaction with Scc1's α-1 and α0

when Scc1's NTD is connected to Pds5 when the latter is part of a ternary complex with Wapl

(Fig. 24, f42). In the latter case, Scc1's NTD approaches Wapl's CTD from an angle

incompatible with the stereo-specific engagement depicted in Figure 25. Additional AF

predictions confirmed that Scc1's first three mini-helices (α-1, α0, and α1) are sufficient for

this type of interaction not only for humans (Fig. 26A, f43) but also for *D. melanogaster* (Fig.

26C, f44), for *C. elegans* (Fig. 26D, f45), for the sponge *A. queenslandica* (Fig. 26E, f46), for

the green alga *P. salinarum* (Fig. 26F, f47), and for the vascular plant *A. thaliana* (Fig. 26G,

f48). Likewise, despite considerable sequence variation, AF predicted a very similar interaction

between Wapl and the equivalent helices of a fragment of human Rec8 containing its first 36

amino acids (Fig. 26H, f49).

Figure 25. A *, AlphaFold's prediction for the interaction of human Scc1's NTD with Wapl's*

CTD (f34A).

These results suggest that there may indeed be an intrinsic affinity between Wapl and Scc1's

α-1 and α0 helices but the interaction is not possible when the helices are packed onto α3 while

the latter is simultaneously connected to Pds5 bound to Wapl's CTD. If positioning Scc1's

NTD in the immediate vicinity of Wapl's CTD, through formation of a stereo-specific ternary

complex between Pds5, Wapl, and Scc1$_N$ is to promote sequestration of Scc1's α-1 α0 helices

by Wapl's CTD, then these helices would have to dissociate from α3. Crucially, AF would not

predict the unfolding necessary if it gave higher priority to packing of Scc1's N-terminal

helices with α3 than to partial unfolding and association with Wapl. That this appears to be the

case should not be surprising, given that Scc1's NTD spends most of its life associated with

Smc3 during which the α0 helix is packed onto top of α3. Even when not associated with

Smc3, Scc1 may not actually be engaged in a release reaction, which after all is probably a rare

event in the life of cohesin, only occurring once every 20 minutes in G1 cells (Gerlich et al.,

2006; Wutz et al., 2020).

There are two possible explanations for the conundrum that Scc1's NTD appears to approach

Wapl's CTD from the wrong direction: either the hypothesis that Wapl sequesters Scc1's NTD

is wrong or Wapl does indeed sequester Scc1's NTD when delivered precisely to the right

location by the Pds5 moiety of the Pds5:Scc3:Wapl:Scc1 quarternary complex, but to do so in

the manner depicted in Figure 26 requires a partial unfolding of Scc1$_N$, in particular

dissociation of Scc1$_N$'s α-1, α0 and α1 helices from α3, a feature that AF may not on its own

be able to predict. We therefore explored whether the packing of the three most N-terminal

helices around α3 could be weakened by changing key residues within α3, without altering the

latter's structure. After a series of trials testing the effects of α3 mutations on the structure

predictions of Scc1's NTD alone, we settled on a pentuple α3 mutant Scc1$_N$(H58A, L59S,

G62D, R65A, R69A) that, according to the PAE and pLDDT plots (Fig. 27, f50, f51) weakens

the association between Scc1$_N$'s three most N-terminal helices and α3, without greatly

affecting the folding of α3 itself. Importantly, AF's prediction for Scc1's interaction with the

Smc3 neck was also unaltered by the pentuple mutant, not only with regard to α3's interaction

with the Smc3 coiled coil, but also with regard to the packing on and around α3 of Scc1's more

N-terminal helices. The latter may be surprising given the extensive changes but AF predictions

are remarkable resilient to mutations and do not reflect the underlying physico-chemical

changes. We then tested the effect of these mutations on AF's predictions for the complex

containing Wapl's CTD, the first 150 residues of Scc1, and the N-terminal straight section of

Pds5. Remarkably, the mutations led to AF predicting with high confidence and in five out five

models the unpacking of Scc1$_N$'s first two helices and their docking with Wapl's CTD (Fig.

28, f52A) in a manner very similar, if not identical, to that observed when the first two or three

helices are run alone with Wapl's CTD (Fig. 26A) or in AF predictions for Wapl's association

with isolated Scc1 NTDs (Fig. 25, f34). AF even managed to predict a similar outcome when

the appropriate fragment of SA was included, thereby creating a quarternary complex whose

Scc1$_N$ is sequestered by Wapl (f52B). It is important to note that the mutations did not affect

any of the residues that actually contact Wapl in these models. Being in α3 itself, which does

not associate with Wapl, the mutations merely affect how the two N-terminal helices associate

with α3 (Fig. 27).

Figure 26. AF predictions for the interactions between Wapl's CTD and the N-terminal 35 amino acids of Scc1 for humans (A, f43), D. melanogaster (C, f44), C. elegans (D, f45), the sponge A. queenslandica (E, f46), the green alga P. salinarum (F, f47), and A. thaliana (G, f48). B, AF's prediction for human Wapl alone (AlphaFold Protein Structure Database AF-Q7Z5K2-F1). H, AF prediction for the interaction between Wapl's CTD and the N-terminal 36 amino acids of human Rec8 (f49). I, interaction between Wapl's CTD and the N-terminal 35 amino acids of Scc1 in a ternary complex containing Scc1$_N$(H58A, L59S, G62D, R65A, R69A), Pds5's NTD, and Wapl's CTD (f52A). See Fig. 28 for the same structure in the complete ternary complex.

Figure 27. A, PAEs for WT Scc1$_N$ (top, f50) and mutant Scc1$_N$(H58A, L59S, G62D, R65A, R69A) (bottom, f51). B, pLDDT plots for WT Scc1$_N$ (left, f50) and mutant Scc1$_N$(H58A, L59S, G62D, R65A, R69A) (right, f51).

Figure 28. *AF's prediction of a ternary complex containing Scc1$_N$(H58A, L59S, G62D, R65A, R69A), Pds5's NTD, and Wapl's CTD (f52A). The location of α3 Scc1$_N$ mutations that distinguish it from f42 shown in Fig. 24 are marked in red.*

If sequestration of Scc1's N-terminal (α-1 & α0) helices by Wapl's CTD is of physiological importance, then the Wapl residues involved in sequestration should be conserved. This consideration cannot be applied to Scc1's NTD as it has multiple other functions. Highly conserved (amongst animals) human Wapl surface residues involved in the interaction include N939 and N990, D979 and E1117, which could interact with Scc1[R11], L983 (which could interact with Scc1[L8]), A1112 and H1115 (which could interact with Scc1[W23]), M1116 (whose sulphur atom could interact simultaneously with Wapl[F1165] and Scc1[W18]), and lastly M1166 (which could interact with Scc1[Y3]). Remarkably, several of these residues have been mutated and shown to abrogate Wapl's RA activity in vivo, namely Wapl(D979K), M1116A, E1117K, and F1165A (Ouyang et al., 2013). In conclusion, sequestration not only helps explain the pattern of conservation but also the physiological consequences of mutating key residues. Crucially, AF predicts that sequestration can also occur when Scc1$_N$ and Wapl's CTD are juxtaposed within a quarternary complex with SA and Pds5 (f52B). Because very

similar rigid body ternary complexes are formed between Pds5, Scc3, and Wapl in plants as

well as fungi (Fig. 23B), the N-terminal helices of their Scc1$_N$s will be presented to Wapl's

CTD in a similar manner to that observed in humans. It is therefore entirely plausible that the

proposed sequestration mechanism applies to cohesin releasing activity in all three kingdoms.

The cleft within Wapl's CTD putatively bound by Scc1$_N$'s α-1 (from animals, plants, and even

certain fungi such as the *Sordariale*s) has previously been highlighted by Chatterjee and co-

workers (Chatterjee et al., 2013), who described a crystal structure (PDB 3ZIL) in which the

equivalent cleft from the yeast *Ashbya gossypii* was occupied by a peptide from Smc3's

ATPase domain, a phenomenon that seemed implausible from a structural point of view and

has subsequently not been confirmed. *A. gossypii,* like *S. cerevisiae* belongs to the

Saccharomycetacae, a group of yeasts in which α-1 is missing, a feature unique amongst

eukaryotes, and replaced by a short string of non-conserved residues. In *Ashbya*'s case, AF

predicts that the cleft is bound by α0 instead of α-1 (f58), a feature that might also apply to *S.*

cerevisiae.

Yeast genetics implies that formation of the canonical Pds5:Scc3:Wapl ternary complex

involves multiple steps

Our AF predictions suggested that formation of canonical rigid body Pds5:Scc3:Wapl ternary

complexes in ascomycetes depends on a juxtaposition of Wapl's CTD and Pds5's NTD

facilitated by Wapl's TYxxxR[T/S]ΦL motif (Fig. 19B), which is essential for RA in vivo. It

is however curious that despite its clear effect on canonical ternary complex formation and

despite highly reproducible AF predictions for the interaction of this motif with Pds5's APDAP

loop using versions of Wapl that lack sequences N-terminal to the motif, AF fails to predict its

association with the APDAP loop of full-length yeast Pds5 when Wapl's entire unstructured

N-terminal sequences are present (f19A, f19D). A possible reason for this is that AF robustly

predicts that the FNFLD motif within yeast Wapl's N-terminal sequences binds to the WEST

locus situated at the opposite end of Pds5 and the limited distance between the FNFLD and

TYxxxR[T/S]ΦL motifs reduces the likelihood of the latter's interaction with the APDAP loop.

Under these circumstances, AF in fact prefers to dock the N-terminal [M/Φ]xxYG[K/R] motif

to the APDAP (f19D), possibly because is it further from the FNFLD motif and therefore less

restricted by the latter's association with WEST. How then does association between the

APDAP loop and the TYxxxR[T/S]ΦL motif come about? A clue emerges from AF predictions

for the interaction between Scc3 and Pds5 in yeast. Unlike humans where there is no direct

interaction between SA and Pds5, AF predicts with high confidence in five out of five models

an interaction between Scc3's C-terminal face (Scc3$_C$) and the pronounced protrusion within

the middle of Pds5 (Pds5$_P$) (Fig. 29A, f54A). Crucially, both surfaces involved are the sites of

mutations (Fig. 29B) that abrogate releasing activity in vivo, namely R996S and R1043L

within Scc3's C-terminal face and R578I, C599F, and E602K within the face of Pds5$_P$

predicted to interact with Scc3$_C$ (Rowland et al., 2009). The remarkable congruence between

the AF prediction and mutations isolated as spontaneous suppressors of *eco1Δ* is unlikely to be

a co-incidence and therefore implies that the interaction must be of physiological importance

and necessary for releasing activity, at least in yeast. There is a simple explanation why no

such interaction was detected in humans, namely that SA proteins have seven extra HEAT

repeats at their C-termini, which means that they lack the Scc3$_C$ face that interacts with Pds5$_P$

in yeast. These extra repeats are shared by Scc3 orthologs in plants, basal fungi, and

basidiomycetes, but lacking in *S. pombe* and *N. crassa* as well as *S. cerevisiae*. AF also

predicted, albeit with low confidence a similar association between Scc3$_C$ and Pds5$_P$ in *N.*

crassa (f54E).

Figure 29. **A**, *AF prediction for the interaction between Scc3 and Pds5 in S. cerevisiae (f54A).*

B, *close up showing residues whose mutation abrogates RA and suppresses eco1Δ.*

Though the Scc3$_C$ face involved in the interaction appears unique to ascomycetes, it has a

potentially important implication for how canonical rigid body Pds5:Scc3:Wapl ternary

complexes are generated in *S. cerevisiae*. This emerges from the fact that the Scc3$_C$:Pds5$_P$

interaction aligns the axes of the two proteins in parallel and thereby juxtaposes their N-termini,

albeit modestly spaced apart (Fig. 29A). As a consequence, any Wapl CTD bound to Scc3's

CES will now be placed in the vicinity of Pds5's NTD, which may favour the binding to the

latter's APDAP loop of the TYxxxR[T/S]ΦL motif (associated with Wapl's CTD) over the

more distant N-terminal [M/Φ]xxYG[K/R] motif.

To test this, we used AF to predict the structure created by a fragment of Wapl containing its

TYxxxR[T/S]ΦL motif together with the part of its CTD that binds Scc3's CES together with

full length Scc3 and a long fragment of Pds5 that contains its entire N-terminal part as well as

its Pds5$_P$ protrusion predicted to bind Scc3$_C$. Remarkably, AF predicted with high confidence

in five out of five models (f54B) a complex with Scc3$_C$ bound to Pds5$_P$, Wapl's CTD bound to

Scc3's CES, a close juxtaposition of the latter with Pds5's NTD, and interaction of Wapl's

TYxxxR[T/S]ΦL with Pds5's APDAP loop (Fig. 30). A similar result was obtained when all

of Wapl's N-terminal sequences were included and crucially under these circumstances Wapl's

TYxxxR[T/S]ΦL and not its N-terminal [M/Φ]xxYG[K/R] motif bound to Pds5's APDAP

loop (f54C).

Figure 30. *AF prediction for a complex between S. cerevisiae Scc3, Pds5, and Wapl's CTD*

with its associated TYxxxRⁱ[T/S]ΦL motif (f54B).

This finding suggests that the $Scc3_C$:$Pds5_P$ interaction facilitates juxtaposition of Wapl's CTD

with Pds5's NTD in manner that would place them close to but not identical to that adopted in

the canonical configuration. It also promotes the interaction of Pds5's APDAP loop with

Wapl's TYxxxR[T/S]ΦL motif that is ultimately necessary for formation of the canonical rigid

body Pds5:Scc3:Wapl ternary complex in yeast. However, the latter is not in fact generated

because its precise adoption is incompatible with the manner in which the axes of Scc3 and

Pds5 are oriented due to their $Scc3_C$:$Pds5_P$ interaction. As a consequence, in this alternative

state, $Scc1_N$ associated with Pds5 is not placed in the position necessary for its sequestration

by the cleft in Wapl's CTD (Fig. 31A). For this to occur, the $Scc3_C$:$Pds5_P$ interaction must

subsequently be broken, which allows Pds5's axis to rotate in a manner that slots it into the

canonical configuration, which then delivers $Scc1_N$ to the cleft in Wapl's CTD (Fig. 31B).

Though speculative, the scheme has the merit of explaining why Wapl's TYxxxR[T/S]ΦL

motif as well as the $Scc3_C$:$Pds5_P$ interaction are essential for RA in yeast but not in animals,

and yet RA in both types of organism appears to share the fundamental mechanism, namely

delivery of Scc1N to Wapl's CTD when part of the canonical Pds5:Scc3/SA:Wapl ternary

complex. If this mechanism is indeed correct and universal, then we must conclude that not all

of the interactions revealed by AF and by yeast genetics can occur simultaneously. The

$Scc3_C$:$Pds5_P$ interaction is clearly incompatible with formation of the canonical ternary

complex and yet the former is essential for RA in yeast. In other words, the proteins involved

in RA in yeast must exist in multiple states and the process of release must involve transitions

between these that presumably depend on a specific combination of on and off rates for the

various interactions involved.

Figure 31. *Putative quarternary complexes between yeast Scc1ₙ, Pds5, Scc3, and Wapl. **A,***
when formed with Scc3ᴄ interacting with Pds5ₚ (f55), created by superimposing f24 and f38B
*onto f54B. **B,** when formed without any interaction between Scc3ᴄ and Pds5ₚ (f53), which*
involves a canonical rigid body Pds5:Scc3:Wapl ternary complex. Model created by super-
*imposing f24 and f38B onto f32B. A movie showing the transition between **A** and **B** at*
https://doi.org/10.6084/m9.figshare.22567525.v1.

Yeast genetics and AF suggest a novel function for Smc3(K112) and K113

Though our current AF predictions explain many of the existing mutations known to abrogate

releasing activity in yeast, as well as providing the first plausible explanation for how Wapl

facilitates release, they do not explain how acetylation of Smc3(K112) and K113 by Eco1 in

yeast blocks release in a manner that does not require Sororin, nor why release is abrogated by

mutations of Smc3's KKD loop, Smc3(K113T, K113N, K112Q K113Q and D114Q/Y) and its

surroundings Smc3(S75R, R107I/S, G110R/V/W) or by Scc3(D189E) and E202K (Rowland

et al., 2009), situated on the N-terminal face of Scc3 (Scc3$_N$). Because it was unclear whether

or not Smc3's KKD loop binds another protein or merely binds DNA, we focused on what

proteins might bind Scc3$_N$. Because the predictions described above excluded both Wapl (f24)

and Pds5 (f54A), we tested whether Scc3$_N$ might bind Smc3's ATPase domain. Remarkably,

AF predicted with high confidence, in the top four ranked models, an interaction between the

face of Scc3$_N$ containing D189 and E202 and the part of Smc3's ATPase containing K112 and

K113 (Fig. 32A, f56A). Not only do both Smc3(K112) and Smc3(K113) participate in the

interaction but so does Smc3(S75), whose substitution by R abrogates RA in yeast (Rowland

et al., 2009). Again, the striking congruence between the predicted interaction and residues

known to be essential for RA implies that the interaction is of physiological importance. We

therefore presume that Scc3(D189E) and E202K abrogate RA because they compromise the

interaction between Scc3$_N$ and Smc3's KKD loop. Because Scc3(E202K) suppresses the

lethality caused by unregulated RA caused by Smc3(K112R, K113R) double mutants as well

as *eco1Δ* (Rowland et al., 2009), we can also be confident that the AF prediction (f56A)

recognizes K112 and K113 in their unmodified state. Acetylation of K112 and K113 would

presumably also abrogate the interaction in vivo and given that the phenotype of Scc3(D189E)

and E202K mutations imply that the interaction is necessary for RA, the AF model finally

provides an explanation for how acetylation would be sufficient to block RA in yeast, without

any need for Sororin.

The face of Scc3$_N$, including D189 and E202, is highly conserved in fungi and AF predicted a

similar interaction between Scc3$_N$ and Smc3's ATPase domain in *Candida albicans* (f57A) but

not, strangely, in *N. crassa* (f57B). D189 and E202 are not conserved in metazoa and AF also

failed to detect any interaction between SA/Scc3$_N$ and Smc3's ATPase domain in humans

(f57C), where the physiological role of Smc3(K105) and K106 acetylation has never been

rigorously investigated.

Finally, AF predicted a very similar interaction between Scc3$_N$ and Smc3's ATPase when the

former was bound by Wapl's CTD (Fig. 32B, f56B) and even when it was part of a canonical

ternary complex with Wapl and Pds5 (Fig. 32C, f56C), implying that the sequestration of

Scc1's NTD by Wapl's CTD could in principle occur when Scc3$_N$ is bound to Smc3's KKD

loop. If so, Scc3N would be held at 110° from Smc3's neck, which may help prevent their re-association during the release reaction (Fig. 33). However, it is also possible that the interaction between Scc3N and Smc3 has a role at some other stage of the release process. Furthermore, it is presently unclear how the complex portrayed in Fig. 31 facilitates cohesin's release from DNA. If the latter is entrapped within the Smc-kleisin ring, then it must pass through the opened Scc1N:Smc3 interface, which is hard to envisage if Scc3 remains bound at this stage to Smc3's KKD loop. Though essential, the binding of Scc3N to Smc3's KKD loop is not necessarily the latter's sole function during release. For instance, K112 and K113 might at some other stage during the reaction facilitate the binding of DNA destined to be released. Interestingly, there would be sufficient space between Smc1 and Smc3 coiled coils for DNA to bind to engaged heads even when this is accompanied by junction of Smc1 and Smc3 joints, which we suggest here may trigger dissociation of Scc1's NTD from Smc3. Importantly, Smc3(K112) and K113 would be in an ideal position to contribute to such DNA binding.

Figure 32. *Interaction between Smc3's ATPase and Scc3's N-terminal domain.* **A** *Left, Detail of the interaction between Smc3 and Scc3 (f56A) showing Scc3(D189) and Scc3(E202)*

*(marked green) and Smc3(S75, K112, K113) (marked red). **A** Right, PAEs for rank 1 model in*

*A. Chains A and B correspond to Smc3 while C to Scc3.   **B**, Interaction between Smc3's*

*ATPase, Scc3's N-terminal domain, and Wapl's CTD (f56B). **C**, Interaction of the canonical*

rigid body Pds5:Scc3:Wapl ternary complex with Smc3's ATPase domain (f56C).

***Figure 33.** A model for the yeast quarternary Pds5:Scc3:Wapl:Scc1 complex in the canonical*

configuration when its Scc3 moiety is bound to the Smc3 ATPase head as predicted by AF,

created by super-imposing f56A with the model in Fig. 31B. There is a modest clash between

Scc1 and the Smc3 coiled coil that would presumably have been avoided if AF were able to

predict the entire complex, which is beyond its capability.

Regulation of release by Smc3 acetylation

Acetylation of Smc3(K112) and K113 by Eco1 is essential for blocking releasing activity in

G2 cells, a process dependent on Pds5 (Chan et al., 2013). How does Eco1 recognize Smc3's

KKD loop and how might Pds5 facilitate this? We therefore asked whether AF can predict how

Eco1 binds to Smc3, Pds5, or other cohesin subunits. In two (ranks 1 and 2) out of five models

with *S. cerevisiae* (f67) and in five out of five models with *Candida glabrata* (f68), AF predicts

with high confidence an interaction between yeast Eco1 and Smc3's ATPase with Scc1's NTD

bound to the coiled coils emerging from it (Fig. 34A). Importantly, Eco1 binds in a manner in

which Smc3(K113) would enter the enzyme's catalytic site as observed in a crystal structure

of a K106-CoA conjugate bound to xEco2 from *Xenopus laevis* (PDB: 5N22) (Chao et al.,

2017) (f69) (Fig. 34B, left). The prediction envisions a very different type of association to that

previously proposed on the basis of attempts to dock the acetyl transferase onto Smc3 heads

(Chao et al., 2017). AF predicts, again with high confidence, a very similar mode of association

between Smc3 and Esco1's catalytic domain (f70), either in the absence or presence of Scc1's

NTD, and a similar mode of interaction was predicted in the top ranked model for Esco2 (f71).

Passage through S phase is necessary for acetylation by Eco1 (Beckouet et al., 2010; Rolef

Ben-Shahar et al., 2008) and Esco2 (Ivanov et al., 2018) and likely also stimulates that by

Esco1. S phase-specific acetylation by all three enzymes requires a small motif (PIP box)

within unstructured sequences immediately N-terminal to their zinc fingers that bind to PCNA

(Moldovan et al., 2006) and this association was confirmed by AF (f72, f73, & f74).

Figure 34. A*, AF predictions for Smc3's interaction with Eco1 from S. cerevisiae (f67) and C. glabrata (f68) with the Scc1 NTD present.* **B***, Left, similarity of this interaction with a crystal structure (PDB: 5N22) of Esco1 with a K106-CoA conjugate. A superimposition of f67 and 5N22 is shown in f69. Right: as in A, left, without Scc1 NTD.*

In addition to predicting binding of Smc3 to the catalytic cleft on the front face of Eco1's

catalytic domain, AF predicts with high confidence, in five out of five models, binding of three

loops on the back side of the enzyme to a conserved loop associated with *S. cerevisiae* Pds5's

most C-terminal HEAT repeats (Fig. 35A, f75). A similar but not identical type of association

is predicted for Esco1 (Fig. 35B, f22) and Esco2 (Fig. 35C, f76), both to Pds5A (Fig. 35) and

Pds5B (not shown). In these cases, an additional contact is made by a pair of anti-parallel β

sheets within the catalytic domain with the α helices that protrude from the centre of Pds5A &

B, which due to Pds5's hook shape are in close proximity to its C-terminal helices. Given that

the association between *S. cerevisiae* Pds5's CTD with the back side of Eco1 is their sole form

of interaction, it is reasonable to assume that this interaction alone recruits Eco1 to cohesin and

thereby facilitates its association with the Smc3 ATPase. This has an implication. If Smc3 is

acetylated by Eco1 when bound simultaneously to Pds5's CTD as well as to Scc1's NTD, then

Pds5 must adopt a very different orientation, relative to the Smc3 ATPase, to that predicted by

AF in the context of releasing activity, where its NTD and not its CTD points towards the

ATPase (Fig. 36). Either the segment of Scc1 linking Scc1's NTD to Smc3 is sufficiently

flexible for Pds5 to adopt more than one orientation relative to the Smc3 ATPase or acetylation

promoted by Pds5 would have to take place after Scc1's NTD has dissociated from the Smc3

neck.

1024

1025   ***Figure 35.*** *AF predictions for interactions between Pds5 and S. cerevisiae Eco1 (**A**, f75),*

1026   *human Esco1 (**B**, f22), and human Esco2 (**C**, f76).*

1027

1028

1029   ***Figure 36.*** *Projected position of Eco1 bound to Pds5 if Scc1$_N$ bound to the latter adopts the*

1030   *orientation predicted by AF (f36). The model was created by superimposing f36 and f75 via*

1031   *their Pds5 moieties. For Eco1 to bind Smc3 K112 and K113, while still bound to Pds5, the*

1032   *Scc1 sequences between Scc1$_N$ α3 and the part bound to Pds5 must bend by 180^0, or Scc1's*

1033   *NTD must dissociate from Smc3.*

1034

Unlike yeast Eco1, Esco1 and Esco2 have long unstructured N-terminal extensions and AF

predicts numerous interactions involving specific motifs within these, not only with various

cohesin subunits (Figs. 13 & 37), but also with the MCM helicase (Fig. 38). Thus, in addition

to the interaction between Esco1's catalytic domain and Pds5's CTD, AF predicts with high

confidence and in five out five models, interactions of a highly conserved, partly helical, motif

(ILxLCEEIAGEIESD) within Esco1 with the WEST site on both Pds5A (f22) and Pds5B,

namely the very same site bound by Wapl's central FGF motif (Fig. 13). Wapl and Esco1 have

opposing effects on release and presumably compete for binding.

The theme of competition between Wapl and Esco1/2 is also observed with SA1:Scc1, where

AF predicts, albeit with less confidence, that the aforementioned ILxLCEEIAGEIESD helical

motif from Esco1 predicted to bind Pds5's WEST has an affinity also to SA1's CES, binding

orthogonal to SA1's HEAT repeats, in a manner similar to the first helix from Wapl's CTD

(Fig. 37A, f77). More importantly, AF predicts that yet another conserved motif

(RFFPSRKxxP[V/L]KC) binds the same groove within the central section of SA1 bound by

Wapl (Fig. 37A, f77). Like the latter, binding extends across the Scc1 polypeptide. Despite

binding to the same locus, the motif has only modest resemblance to the one used by Wapl.

Interestingly, a largely unrelated but broadly conserved motif within Esco2's N-terminal

domain is predicted to bind in a similar fashion to the equivalent locus on a SA2:Scc1 complex

(Fig. 37B, f78). This is adjacent to an even more conserved sequence, the PBM-B motif

(GAAFF) required for the recruitment of *Xenopus* Eco2 (xEco2) to chromatin in a pre-RC

dependent manner (Higashi et al., 2012), subsequently postulated to bind cohesin (Ivanov et

al., 2018), and predicted by AF to bind as a very short helix to SA2's C-terminal HEAT repeats

(Fig. 37B). A second conserved motif (PBM-A) is required for xEco2's recruitment to

chromatin and for its ability to acetylate Smc3 (Higashi et al., 2012). This was subsequently

shown by co-immunoprecipitation to bind the MCM helicase. Because analysis of amine-

reactive chemical crosslinking using mass spectrometry indicated Mcm4 or Mcm7 as the most

likely partners (Ivanov et al., 2018), we used AF to investigate Esco2's interaction with each

of these subunits. No interaction was observed with Mcm4 (f79), but three out of five AF

models predicted with high confidence an interaction between PBM-A and Mcm7 (Fig. 38,

f80). The motif is predicted to form two short helices connected by a conserved loop. The first

of these binds to the surface of Mcm7's CTD while part of the loop together with the second

helix is sandwiched between the CTD and a C-terminal winged helical domain that binds the

surface of ORC1 and Cdc6 within the pre-RC (Zhai et al., 2017). This interaction together with

its PIP box may impart at least some of the S phase specificity of Smc3 acetylation by Esco2.

1069

Figure 37. A, *AF predictions for the interactions between SA1:Scc1 and the N-terminal extension of Esco1 (left, f77) and between SA2:Scc1 and Wapl-M (right, f25).* **B**, *AF prediction for the interaction between SA2:Scc1 and Esco2's PBM-B (f78).*

1075

1076   ***Figure 38.*** *AF prediction for the interaction of Esco2's PBM-A and Mcm7 (f80).*

1077

Regulation of release by Sororin

Though Smc3 acetylation may be sufficient to protect sister chromatid cohesion from Wapl-
mediated release in yeast, it is necessary (Nishiyama et al., 2010) but not sufficient in
vertebrate, insect, and plant cells, where proteins called Sororins (vertebrates and plants) or
Dalmatians (insects) (Mota et al., 2022; Rankin et al., 2005; Yamada et al., 2017) are also
required. Sororin is absent from G1 mammalian cells because it is degraded at the hands of
APC^{Cdh1} and therefore only accumulates shortly before S phase. However, accumulation alone
is insufficient for its interaction with chromosomal cohesin as its association also depends on
DNA replication in *Xenopus* extracts that lack Cdh1 (Nishiyama et al., 2010). Association with
cohesin only takes place when Smc3 is acetylated in an S phase-specific manner, largely by
Esco2, and on DNA that has actually been replicated, whereupon it is essential for maintaining
cohesion between sisters (Ladurner et al., 2016). Sororin is thought to bind to and protect from
Wapl only cohesin engaged in cohesion, but whether this is really the case is not known.
Importantly, though Smc3 acetylation is necessary for Sororin's association, it is insufficient
and some other change in the state of cohesin that occurs during replication may be necessary.
Sororins are largely unstructured and for the most part poorly conserved. Common to all
members of the family are C-terminal so called Sororin boxes predicted to contain a short (six
turns)   α   helix   and   whose   consensus   in   animals   is
[W/F/Y]xxxΦNxxFxE[A/Φ][E/D]x[F/Y]ELxΦE.

Three other features are common to vertebrate Sororins, namely a short stretch of basic amino

acids immediately N-terminal to the Sororin box, which is preceded by a FGF motif (consensus

[S/T]ΦFGFExL), separated by ten residues from a so called YSR motif (consensus

[M/W]S[K/Q]KVRRSYSRL). Functional studies indicate that the Sororin box is essential,

while the YSR and FGF motifs are not, at least when the protein is over-expressed (Wu et al.,

2011). Because the cohesion defects associated with sororin inactivation are largely suppressed

by Wapl mutants (Ladurner et al., 2016; Nishiyama et al., 2010), Sororin is believed to be

required merely to suppress Wapl-mediated release, a function it fulfils despite a residence

time on cohesin of about 60 seconds (Ladurner et al., 2016). It could perform this function

either by inhibiting Wapl's sequestration of Scc1's NTD or by preventing the latter's

dissociation from Smc3 in the first place. Though AF cannot address why Sororin only

associates with replicated DNA, it has the potential to reveal which cohesin subunits interact

with its essential Sororin box and/or FGF and YSR motifs. Accordingly, we used AF to predict

the interaction of full-length mouse Sororin with cohesin subunits known to be involved in

Wapl-mediated RA. AF predicted a potential interaction between Sororin's FGF motif and the

CES of an SA:Scc1 complex, albeit in only one out five models (f81) and not in the same

manner as CTCF's YDF motif (Li et al., 2020). In contrast, it predicted with high confidence

an interaction between its YSR and FGF motifs and Pds5B's C-terminal domain (Fig. 39), both

in mice (f82) and in zebrafish (f21). Like Wapl, Sororin's FGF was predicted to bind to Pds5's

WEST locus while the YSR bound to an adjacent (more C-terminal) pair of helices. In both

cases, the highly conserved aromatic residues in each motif were inserted into grooves between

adjacent HEAT repeats and, notably, all five models predicted simultaneous association.

Crucially, none of them predicted binding of the YSR to Pds5's N-terminal APDAP loop,

which was initially surprising as it has been shown to compete with the binding of Wapl's YSR

to this site (Ouyang et al., 2016). The reason for this becomes apparent when one takes into

account that Sororin's YSR and FGF motifs are separated by no more than 15 amino acids,

which is too short to permit simultaneous binding of the YSR and FGF motifs to Pds5's N-

terminal APDAP loop and C-terminal WEST respectively.

1125

1126

1127

1128   ***Figure 39**. AF predictions for Sororin's interaction with Pds5B in mouse (f82) and zebrafish*

1129   *(f21).*

1130

Given that AF failed to predict any association between the C-terminal Sororin box and either

SA2, Pds5B, or indeed Wapl itself (f83), we considered whether it might instead bind to the

Smc3:Scc1 interface. Remarkably, AF predicted, with high confidence in two out of five

human models, interaction with an Smc3 ATPase domain whose coiled coils were bound by

Scc1's NTD (Fig. 40, f84). A very similar interaction was observed with high confidence in

five out of five models using orthologous proteins from *Xenopus laevis* (f85), *Danio rerio*

(zebrafish) (f86), *D. melanogaster* (f87), *Tribolium casaneum* (red flour beetle) (f88), *A.*

thaliana (f89), and even the ascomycete *Aspergillus fumigatus* (f90). The similarity of the

predictions is remarkable given that the Sororin orthologs of vertebrates, insects, plants, and

fungi have very little sequence homology and even exhibit considerable variation between their

Sororin boxes (Mota et al., 2022). In contrast, no interaction was predicted between Sororin

and a complex between Smc1 and Scc1's CTD in *D. melanogaster* (Fig. 41, f91), *A. thaliana*

(Fig. 41, f92), and *A. fumigata* (Fig. 41, f93). Intriguingly, AF suggests that Sororin's C-

terminal helix might interact with the longer of the two helices within Scc1's CTD in mammals

(Fig. 41, f94). Thus, AF's prediction that the Smc3:Scc1$_N$ interface is a highly conserved

partner of Sororin's CTD is not shared by Smc1:Scc1$_C$.

1147

Figure 40. *AF predictions for Sororin's interaction with Smc3 bound to Scc1's NTD in humans (f84), X. laevis (f85), D. rerio (f86), D. melanogaster (f87), T. casaneum (f88), A. thaliana (89), and A. fumigatus (f90).*

1153

1154   ***Figure 41***. *AF predictions for the Sororin box's interaction with a complex between Smc1 and*

1155   *Scc1's CTD in D. melanogaster (f91), in A. thaliana (f92), in A. fumigatus (f93), and Mouse*

1156   *(f94).*

1157

According to all of the AF models, Sororin's C-terminal α helix straddles Scc1's most N-

terminal helix and the loop connecting its fourth helix to the long α helix that forms a three-

helical bundle with the Smc3 coiled coil (Fig. 40). Meanwhile, the C-terminal end of the

Sororin helix contacts the C-terminal α helix emerging from the Smc3 ATPase just before it

forms a coiled coil with its partner from the N-terminal part of the protein, while the terminal

4-5 amino acids straddle a pair of adjacent β sheets. Aromatic residues play a key part in the

interface. In vertebrates, a conserved tryptophan (W233 in humans) sits on top of Scc1's most

N-terminal helix, while a pair of highly conserved phenylalanines (F241 and F247) interact

with the aforementioned Smc3 α helix respectively (Fig. 40, human rank 1, f84). The nature

of Sororin's interaction with the Smc3:Scc1 interface is not only universal across eukaryotic

kingdoms but also helps explain how Sororin hinders release. By binding Scc1's N-terminal

helix, Sororin would prevent its unfolding from $Scc1_N$'s long α3 helix, thereby preventing

sequestration occurring even before the NTD has dissociated from Smc3. Perhaps more

importantly, by binding over Scc1's N-terminal helices and simultaneously binding Smc3,

Sororin could actually stabilize the interaction between Scc1's NTD and the Smc3 neck,

thereby thwarting any subsequent sequestration of its N-terminal helices by Wapl. Such

stabilization might also prevent the distortions in Smc3's neck produced by simultaneous head

engagement and joint junction, thereby preventing such an event. Despite these insights, AF

predictions provided no clues with regards to the role of Smc3 acetylation.

Finally, we tested whether Sororin might also bind Wapl's CTD. Remarkably, AF predicted

with high confidence in five out five models that Sororin's [S/T]ΦFGFExL motif would bind

to exactly the same cleft within Wapl's CTD that sequesters the most N-terminal α helix from

Scc1 and does so in a manner that would be mutually exclusive (Fig. 42, f83). As already

mentioned, this motif is also predicted to bind Pds5's WEST (Fig. 13B, f21 & Fig. 39, f82) and

by doing so ensures that Sororin's [M/W]S[K/Q]KVRRSYSRL motif binds to an adjacent

groove in Pds5's CTD (Fig. 39, f82) and not to its N-terminal APDAP loop. However, if

Sororin's [S/T]ΦFGFExL motif was instead bound to Wapl's CTD (via the latter's proposed

$Scc1_N$ sequestration cleft), when part of the canonical ternary RA complex with SA and Pds5,

then the [K/Q]KVRRSYSRL motif would be free to bind to Pds5's APDAP, which is brought

into close proximity to the aforementioned cleft in Wapl's CTD by ternary complex formation.

In other words, if Sororin's [S/T]ΦFGFExL and [M/W]S[K/Q]KVRRSYSRL motifs are to

bind cohesin simultaneously, the short linker connecting them will constrain the distance

allowed between their binding sites. If the [S/T]ΦFGFExL motif binds to Pds5, its partner must

also bind to an adjacent site on Pds5 (Fig. 39) but if [S/T]ΦFGFExL instead binds to Wapl's

CTD, its [M/W]S[K/Q]KVRRSYSRL partner could in principle bind simultaneously to Pds5's

APDAP loop.

To test this, we asked AF to predict the interaction between Sororin and an N-terminal fragment

of Pds5 that lacked its WEST [S/T]ΦFGFExL binding site. In this case, AF predicted with high

confidence and in five out of five models an interaction exclusively between Sororin's

Dxx[M/W]S[K/Q]KVRRSYSRL motif and Pds5's APDAP (Fig. 43 left, f95). In this case, the

interaction involved conserved aspartic acid (D135) and methionine (M138) residues at the N-

terminal end of the motif, which are predicted to be part of a short helix (marked in grey in Fig.

43), as well as the YSR sub-motif at its C-terminal end, whose interaction with Pds5's APDAP

is predicted by AF to be identical to that observed in a crystal structure containing Wapl's N-

terminal YSR motif bound to Pds5 (Ouyang et al., 2016) (PDB 5HDT) (Fig. 43, right). These

predictions therefore raise the possibility that Sororin may have a second line of defence against

Wapl in case dissociation of Scc1's NTD from Smc3 is accompanied by formation of the

canonical ternary (SA:Wapl:Pds5) RA complex, namely occupation of the key cleft in Wapl's

CTD by Sororin's [S/T]ΦFGFExL motif, which would exclude sequestration of Scc1$_N$ (Fig.

42), and be facilitated by the simultaneous binding of Sororin's [M/W]S[K/Q]KVRRSYSRL

motif to the neighbouring N-terminal APDAP loop of Pds5 (Fig. 43).

1209

1210

1211   ***Figure 42.*** *AF's prediction for the interaction between Sororin and Wapl's CTD (f83).*

1212

1213

Figure 43. AF predicts that Sororin's [M/W]S[K/Q]KVRRSYSRL (YSR) motif binds to Pds5's
APDAP loop (f95) (left) in a manner similar to Wapl's YSR peptide (right, PDB 5HDT), but
only if binding of Sororin's [S/T]ΦFGFExL motif is prevented from binding Pds5's WEST site
by deleting the C-terminal part of Pds5 containing this WEST site.

Interactions between Scc2/Nipbl and Scc3/SA

One of the key insights revealed here by AF is the canonical rigid body association between
SA and Pds5 induced by their simultaneous association with Wapl's CTD (Fig. 16). According
to AF, SA and Pds5 do not otherwise interact directly and only do so in the presence of Wapl.
We note however, that the yeast orthologs do so via their Scc3$_C$:Pds5$_P$ interaction (Fig. 29). In
the course of investigating their association, we used AF to undertake a more systematic
investigation of how cohesin's three HAWK subunits interact. Similar to SA/Scc3, AF also
predicted no interaction between Nipbl/Scc2 and Pds5. In contrast, it predicted with high
confidence and in five out five models a pair of simultaneous interactions between full length
versions of SA and Nipbl (Fig. 44A, f96). One of these interactions is identical to that observed
in a cryo-EM structure of the clamped state (Fig. 44C, PDB 6WG3) (Shi et al., 2020) and
involves interactions between Nipbl helices next to its N-terminal protrusion and the outside
surface of SA where there is a twist in the alignment of its anti-parallel HEAT repeats at the
point where the protein starts to bend. The second region of interactions has not hitherto been
observed and involves an interaction between helices at the point where Nipbl starts to bend
(on its outer surface) and the antiparallel helices that make up SA's pronounced "nose". An
identical pair of interactions were predicted for the orthologous proteins from *A. thaliana* (Fig.
44B, f97).

1237

1238

Figure 44. *AF predictions for interactions between Nipbl/Scc2 and SA/Scc3 for humans (**A**, f96), A. thaliana (**B**, f97), and S. cerevisiae (**D**, f98). **C**, cryo-EM structure PDB 6WG3.*

1241

In contrast, AF predicted only the nose interaction for yeast Scc2 and Scc3 (Fig. 44D, f98). In this case, the residues involved are highly conserved on both the Scc3/SA (Scc3: Q323, E327 and SA1: Q165, E169) and Scc2/Nipbl (Scc2: P958, Y959 and Nipbl: P2029, Y2030) sides of the interface and the interaction is therefore in all probability of physiological importance. Apart from microsporidians (Roig et al., 2014), where it is absent, the nose is a highly conserved feature of Scc3/SA orthologs. Its deletion is lethal in yeast (Roig et al., 2014), but its function has hitherto remained obscure. It is interesting that the interaction between SA's nose and Nipbl was not observed in the PDB 6WG3 cryo-EM structure (Shi et al., 2020). This

1250   is probably because it would have been incompatible with DNA binding to Nipbl. Now that

1251   AF has revealed the details of the interaction, it will be possible to create specific mutations

1252   that affect it and thereby address whether the interaction is necessary either for LE or for DNA

1253   entrapment within S-K rings, both in vitro and in vivo.

1254

DISCUSSION

Our use of AF to predict interactions between cohesin subunits has led to numerous hypotheses

concerning how cohesin's association with chromatin is regulated. Can one be sufficiently

confident to undertake the biochemical and genetic experiments needed to test them? And if

so, what will be the most effective techniques to employ? Aside from AF's excellent track

record, it so happens that in several instances experimental data, particularly the phenotypes of

mutations, was already in the public domain but had hitherto not been interpretable. Thus, AF's

predictions about the interaction between Wapl's CTD and Scc3/SA explains why mutation of

both Wapl and Scc3/SA residues predicted by AF to affect their interaction are defective in RA

activity in vivo (Beckouet et al., 2016; Hara et al., 2014; Ouyang et al., 2013; Rowland et al.,

2009). Likewise, the phenotypes of an extensive set of mutations affecting highly conserved

surface residues within the C-terminal half of Wapl's CTD (Ouyang et al., 2013) are fully

consistent with AF's prediction that its role is to sequester Scc1's N-terminal helices (α-1 and

α0). Even more impressive is the remarkable congruence between cohesin subunit interfaces

revealed by AF and the locations of mutations that abrogate RA in yeast and were isolated

without any prior hypothesis, as spontaneous suppressors of *eco1* mutant lethality (Rowland et

al., 2009). Yet further support comes from the congruence between AF predictions and

sequence conservation (beyond that used by AF), which reflects natural selection and hence is

also a robust measure of function. As a consequence, our analyses have been guided not only

by the degree of AF's confidence but also by a requirement that predictions be broadly based

from a taxonomic viewpoint, namely similar in multiple and as distant as possible eukaryotic

lineages, and concern sequences and domains that are evolutionarily conserved. Despite this,

AF revealed instances of physiologically important interactions specific to particular

eukaryotic lineages, for example the interactions between the N- and C-terminal faces of Scc3

with Smc3's ATPase and Pds5's protrusion, respectively, which were only detected in yeast

and related fungi. Last but not least, a key guiding principle has been whether the models

appear plausible and above all explanatory.

In so far that AF largely functions by correlating the query sequences with known structures
and their sequences, and not by calculating chemical or physical properties from first
principles, there is reason to be optimistic that the results will be in many cases comparable in
robustness to those obtained by experimental structural biology, which often uses or even
requires unwarranted conditions (protein concentration, salt concentration, divalent cations,
pH, temperature, etc) and usually evaluates interactions either pairwise or among a very limited
number of constituents and not as in vivo where proteins must compete with a large fraction of
the proteome, as well as with other macromolecules. We contend that many if not most of the
hypotheses described in this paper are plausible, novel, in some cases provocative, and above
all explanatory, and therefore deserve communication before validation using orthogonal
approaches.

What then might these be? At the top of the list must be renewed attempts to visualize the
predicted complexes by cryo-EM alongside the use of bi-functional thiol specific reagents to
crosslink both in vitro and in vivo cysteine pairs inserted into the predicted interfaces (Chapard
et al., 2019; Srinivasan et al., 2018). Mutation of key residues will also have an important role,
but it is important to stress that mutations alone rarely address whether the resulting
physiological defects are due to the loss of a specific interaction. They merely reveal whether
or not one or other interaction surface is important. To achieve this goal using mutations alone,
it would be necessary to generate not just one or two mutations but a large series and
demonstrate a robust structure-function relationship, or better still demonstrate that mutations
in both sides of an interface produce identical phenotypes. A good example of the latter is our
finding of RA-abrogating mutations identified through classical (i.e. hypothesis free) forward
genetics mapping to both sides of interfaces identified by AF, for example that between $Scc3_C$
and $Pds5_P$ and between $Scc3_N$ and Smc3's KKD loop. Combining cross-linking with a more
limited mutational analysis is probably therefore the most economical way of proceeding, but
even this is will be a considerable undertaking as all substitutions should be made within
endogenous genes.

Though our experience is anecdotal, the modelling that we undertook for this analysis was
sufficiently extensive that it is worth sharing the impressions gained into the strengths and
weaknesses of AF as well as what may transpire to be best practice. First and foremost, is the
issue of the optimum scale for embarking on a prediction. On the one hand, AF's ability to
detect patterns between specific domains clearly diminishes as the size and number of
components increase. The combinatorial space increases with the square of the number of

residues provided in the query sequence(s). On the other hand, predicted interactions clearly
have greater value if AF manages to predict them within the context of intact and therefore
possibly large proteins or protein assemblies, whose size approaches the limits of what is
currently practical or even possible given GPU memory limitations. It is our experience that
initial forays should be performed while making as few assumptions as possible as to the nature
of the interactions and therefore giving AF as large a landscape as is practically possible.
Nevertheless, following up interactions revealed in this manner also has value. It is also clear
that there is probably no relationship between the confidence of AF predictions and any actual
affinity between proteins, as would be expected given the AF algorithm. If biology demands a
weak interaction and natural selection has delivered this, AF clearly excels in picking up the
resulting patterns and in this regard it provides an approach that is complementary to
conventional biochemistry and structural biology with which weak interactions are sometimes
difficult to investigate. As a consequence, many of the interactions detected by AF may be
challenging to pursue using these techniques and combining experimental structural work with
the sort of cross-linking techniques described above may prove useful in this regard. Another
important lesson, which may prove to be of general validity, is that complexes such as cohesin
should not be regarded as conventional solids, which imply a lack of constituent mobility or as
liquids, which would require massively larger stoichiometries. They should instead be
considered as promiscuous molecular communities made up by a large set of potential
interactions with a great heterogeneity of on and off rates and extensive competition between
different constituents, many of which have multiple possible partners and multiple binding
interfaces. Some of the interactions are between relatively rigid domains but many are between
short motifs embedded in very long intrinsically disordered loops and extensions.
Understanding the physical dynamics of such communities as well as the mechanisms
underlying the switches between states will be a formidable challenge. In the case of cohesin,
both loop extruding and cohesive forms must be capable of switching between forms that
associate stably with chromatin fibres to ones that dissociate, and a competition for shared
binding sites between factors promoting release or its inhibition presumably create bi-stable
switches.

In summary, our AF predictions have yielded several hypotheses for how cohesin's association
with chromatin is regulated. These include:

1) The role of cohesin's hinge in effecting a junction between the joints within Smc1 and
Smc3's coiled coils. When combined with ATP-driven head engagement in the absence of

Scc2/Nipbl, but in the presence of Pds5, this junction is predicted to create a distortion of the

coiled coil to which the Scc1's NTD normally binds and thereby triggers its dissociation.

2) The proposal that Wapl's CTD forms a ternary complex with the pair of cohesin's HAWK

subunits previously implicated in release, namely SA/Scc3 and Pds5. Upon dissociation from

Smc3, Scc1 is predicted to bind to Pds5 in a manner that places Scc1's short N-terminal helices

in close proximity to a highly conserved site within Wapl's CTD when part of the ternary

complex with SA and Pds5. The subsequent unfolding of Scc1's α-1 and α0 helices from the

long α3 helix that normally associates with Smc3's neck then leads to their sequestration by

this part of Wapl. We propose that sequestration in this manner is the chief function of Wapl

and that it serves to keep the exit gate open long enough for DNA to pass out of the cohesin

ring.

3) The finding that in yeast formation of the canonical ternary complex needed to deliver Scc1's

N-terminal helices to Wapl may be a multi-stage process during which an alternative ternary

complex involving the $Scc3_C$:$Pds5_P$ interaction is required for subsequent formation of the

canonical one.

4) The first insight into how Smc3's KKD loop facilitates release, at least in *S. cerevisiae* where

it binds the N-terminal face of Scc3.

5) An understanding how the acetyl transferases that block release and compete with Wapl for

binding cohesin's HAWK subunits bind to their substrate and how this is facilitated during S

phase by simultaneous binding to Pds5 and to the Mcm helicase.

6) The first insight into how Sororin is recruited to cohesin and how it hinders Wapl-mediated

release, through the association of a conserved C-terminal α helix with cohesin's exit gate,

namely the junction of Scc1's NTD with Smc3's neck.

7) The discovery that Scc2/Nipbl and SA/Scc3 interact with each other via an association

involving the latter's highly conserved nose, for which there has hitherto been no explanation.

Whether this association has a role during LE or in the entrapment of DNA inside the cohesin

ring remains to be addressed.

Of these, the notion that a key function of Wapl's CTD is to bind α-1 and α0 from $Scc1_N$ is

probably the most remarkable as it is based not on a single AF prediction but on an elaborate

concatenation of numerous models, producing a quarternary complex involving Wapl, Pds5,

SA/Scc3, and Scc1, in which Scc1's N-terminal α helices are positioned next to a pair of crucial

sites within Wapl's CTD. Key observations underpinning this notion include the conservation

of Wapl's putative $Scc1_N$ binding site, the existence of mutations within it that abrogate release

in vivo both in yeast and in mammalian cells, the prediction by AF that Pds5, SA/Scc3, and

Wapl form a highly specific ternary complex in animal, plant, and fungal lineages, that $Scc1_N$'s

association with the spine of Pds5's NTD places the formers' $\alpha-1$ and $\alpha0$ helices next to the

putative binding site within Wapl's CTD, that formation of the ternary complex in fungi

depends on a motif within Wapl's unstructured N-terminal sequences which binds Pds5's

APDAP loop and is essential for RA in *S. cerevisiae*, AF predictions for the interaction between

$Scc1_N$ and Wapl that appear conserved among a very wide variety of eukaryotes, and lastly on

the fact that dissociation of $Scc1_N$ from Smc3 depends on Wapl in vivo in *S. cerevisiae* and

that RA is abrogated by their co-translation. Whether like a jigsaw puzzle, our hypothesis

proves to be the sole solution to these explicanda will nevertheless require numerous types of

wet experiments.

Last but not least, it should be pointed out that due to the nature of AF, none of our models

incorporate DNA and therefore shed no direct insight into how sequestration of Scc1N by Wapl

actually facilitates cohesin's dissociation from chromosomes in vivo.

METHODS

AF predictions prior to February 2023 were performed via Google CoLab using Alphafold 2, while those performed thereafter used Colabfold v1.5.2 AF2 with MMSeqs2. Supposition of different models were performed using UCSF Chimera while all figures in this manuscript and Movie 1 were created using PyMOL v2.5 (Schrödinger, LLC). Each subunit was drawn in all figures using the following colours: Smc1=red; Smc3=blue; Scc1=green; Scc2/Nipbl=yellow; Scc3/SA1/SA2=orange; Pds5=violet; Wapl=cyan; Eco1/Esco1/Esco2=ivory; Sororin=grey.

The *WAPL* deletions were introduced into the genome of *S. cerevisiae* using CRISPR-Cas9. Cells of strain KN18714 (a W303 related strain whose endogenous *WAPL* gene was tagged with EGFP) were transformed with a 738 nt (for Δ4-9) or a 138 nt (for Δ208-215) PCR "healing fragment" along with a 2 micron *LEU2* plasmid, constitutively expressing Cas9 endonuclease and transcribing the structural RNA that contains the Cas9 binding site and the guide RNA that will target the double strand break formation by Cas9 to the *WAPL* (*RAD61*) gene. The guide RNA was designed 20 nt upstream of the NGG (E228, V229) using the annealed oligos MB526F/MB527R (MB526F: 5'-ATCAAATGAAATAATGGGAGAAG-3', MB527R: 5'-AACCTTCTCCCATTATTTCATTT-3') and fused afterwards at the 5' end of the structural RNA with Sap1 cloning. The PCR "healing fragment" was created using the oligos MB543F/MB525R for Δ4-9 and MB534F/MB525R for Δ208-215 (MB543F: 5'-TCGCAAAACGAAACCATCTTCTTACCCTAAAGCATCCTGTTTCTGAAAAAATGAG AGCAGTTCTCAGAACTCCTTTTAGATCTAATAAAGGCTTGCCATC-3', MB525R: 5'-TTACTGTTGTCAGCGTTATTTAGGGACAAGGTGTTAGCTTTCTGATCTACTTCTTC TCCCATTATTTCATTTTC-3', MB534F: 5'-GCATACGCACATCCAAAGACAAAAAATAAACAAAATAAAGAAAATG-3') and contained the deletion, a mutagenized NGG Cas9 sequence target (E228, V229: GAGGTA to GAAGTA) and 5'-, 3'- homology arms needed for repair of the double strand break created by Cas9. As a control, no colonies were obtained after yeast transformation of the 2 micron *LEU2* plasmid alone without the "healing fragment". All mutations were confirmed by sequencing the genomic locus.

The ability of the EGFP-tagged *WAPL* mutations to suppress *eco1Δ* lethality was tested by crossing them with strain KN16432, which carries deletions of both *ECO1* and *WAPL*. Dissection of 18 tetrads from crosses involving *WAPLΔ208-215* yielded 60 viable spores, of which 27 carried *eco1Δ*. Of these 15 carried the *waplΔ* and 12 the *WAPLΔ208-215* mutation.

Thus, *WAPL Δ208-215* was as effective as *waplΔ* in suppressing *eco1Δ* lethality and this allele

is therefore defective in releasing activity. In contrast, dissection of 18 tetrads from crosses

involving *WAPLΔ4-9* yielded 49 viable spores of which 16 carried *eco1Δ*. Of these, all 16

carried *waplΔ*. Thus, *WAPLΔ4-9* cannot suppress *eco1Δ* lethality and is therefore proficient for

releasing activity. Separate experiments demonstrated that *WAPLΔ208-215* suppressed *eco1Δ*

when untagged or tagged with PK9 while *WAPLΔ4-9* never did so. Mutation of both motifs

(*Δ4-9 Δ208-215*) also suppressed *eco1Δ* lethality as did an allele in which the TYxxxR[T/S]ΦL

motif (YGKKRTIL) was inserted at *Δ4−9* (creating the sequence MRAYGKKRTILVLR)

within *Δ4-9 Δ208-215*. In other words, the TYxxxR[T/S]ΦL motif does not appear to function

when placed at Wapl's N-terminus, namely when further from its CTD.

1438

1439   **YEAST STRAINS**

1440

1441

Strain number	Genotype
KN18714	*Mat a, Wapl-EGFP::His*
KN24262	*Mat a, wapl(Δ4-9)-EGFP::His*
KN24060	*Mat a, wapl(Δ208-215)-EGFP::His*
KN24224	*Mat a, wapl(Δ208-215)-EGFP::His, eco1::NatMx*
KN16432	*Mat alpha, eco1::NatMx, wapl::HphMx*
KN26163	*Mat a, wapl(Δ208-215), eco1::NatMx*
KN25817	*Mat a, wapl(Δ208-215)-PK9::NatMX, eco1::HphMx*
KN25168	*Mat alpha, eco1::HphMx, leu2::Eco1p-Eco1::LEU2*

1442

1443   **DATA AVAILABILIY**

1444   All data files (f1-f98) referred to in the main text and figure legends are available at

1445   https://doi.org/10.6084/m9.figshare.22567318.v2. The actual amino acid sequences used

1446   for each AF prediction are to be found embedded in the predicted structure files (in PDB

1447   format).

1448

1449   **FUNDING**

1450   This work was supported by the Medical Research Council (U105184326 to J.L.) and by

1451   Cancer Research UK (26747 to K.A.N).

1452

1453   **ACKNOWELDGEMENTS**

1454   K.A.N would like to thank Madhusudhan Srinivasan (Oxford, UK) for numerous discussions

1455   that stimulated and informed this work.

REFERENCES

Bastie N, Chapard C, Dauban L, Gadal O, Beckouet F, Koszul R. 2022. Smc3 acetylation, Pds5 and Scc2 control the translocase activity that establishes cohesin-dependent chromatin loops. *Nat Struct Mol Biol* **29**:575–585. doi:10.1038/s41594-022-00780-0

Beckouet F, Hu B, Roig MB, Sutani T, Komata M, Uluocak P, Katis VL, Shirahige K, Nasmyth K. 2010. An Smc3 acetylation cycle is essential for establishment of sister chromatid cohesion. *Mol Cell* **39**:689–99. doi:10.1016/j.molcel.2010.08.008

Beckouet F, Srinivasan M, Roig MB, Chan KL, Scheinost JC, Batty P, Hu B, Petela N, Gligoris T, Smith AC, Strmecki L, Rowland BD, Nasmyth K. 2016. Releasing Activity Disengages Cohesin's Smc3/Scc1 Interface in a Process Blocked by Acetylation. *Mol Cell* **61**:563–574. doi:10.1016/j.molcel.2016.01.026

Bürmann F, Lee B-G, Than T, Sinn L, O'Reilly FJ, Yatskevich S, Rappsilber J, Hu B, Nasmyth K, Löwe J. 2019. A folded conformation of MukBEF and cohesin. *Nat Struct Mol Biol* **26**:227–236. doi:10.1038/s41594-019-0196-z

Chan KL, Gligoris T, Upcher W, Kato Y, Shirahige K, Nasmyth K, Beckouet F. 2013. Pds5 promotes and protects cohesin acetylation. *Proc Natl Acad Sci U S A* **110**:13020–5. doi:10.1073/pnas.1306900110

Chan K-L, Roig MB, Hu B, Beckouët F, Metson J, Nasmyth K. 2012. Cohesin's DNA exit gate is distinct from its entrance gate and is regulated by acetylation. *Cell* **150**:961–974. doi:10.1016/j.cell.2012.07.028

Chao WCH, Wade BO, Bouchoux C, Jones AW, Purkiss AG, Federico S, O'Reilly N, Snijders AP, Uhlmann F, Singleton MR. 2017. Structural Basis of Eco1-Mediated Cohesin Acetylation. *Sci Rep* **7**:44313. doi:10.1038/srep44313

Chapard C, Jones R, van Oepen T, Scheinost JC, Nasmyth K. 2019. Sister DNA Entrapment between Juxtaposed Smc Heads and Kleisin of the Cohesin Complex. *Mol Cell* **75**:224-237.e5. doi:10.1016/j.molcel.2019.05.023

Chatterjee A, Zakian S, Hu XW, Singleton MR. 2013. Structural insights into the regulation of cohesion establishment by Wpl1. *EMBO J* **32**:677–87. doi:10.1038/emboj.2013.16

Ciosk R, Shirayama M, Shevchenko A, Tanaka T, Toth A, Shevchenko A, Nasmyth K. 2000. Cohesin's binding to chromosomes depends on a separate complex consisting of Scc2 and Scc4 proteins. *Mol Cell* **5**:243–54. doi:10.1016/s1097-2765(00)80420-7

Collier JE, Lee B-G, Roig MB, Yatskevich S, Petela NJ, Metson J, Voulgaris M, Gonzalez Llamazares A, Löwe J, Nasmyth KA. 2020. Transport of DNA within cohesin involves clamping on top of engaged heads by Scc2 and entrapment within the ring by Scc3. *Elife* **9**:e59560. doi:10.7554/eLife.59560

Collier JE, Nasmyth KA. 2022. DNA passes through cohesin's hinge as well as its Smc3-kleisin interface. *Elife* **11**. doi:10.7554/eLife.80310

Davidson IF, Bauer B, Goetz D, Tang W, Wutz G, Peters JM. 2019. DNA loop extrusion by human cohesin. *Science* **366**:1338–1345. doi:10.1126/science.aaz3418

Diebold-Durand M-L, Lee H, Ruiz Avila LB, Noh H, Shin H-C, Im H, Bock FP, Bürmann F, Durand A, Basfeld A, Ham S, Basquin J, Oh B-H, Gruber S. 2017. Structure of Full-Length SMC and Rearrangements Required for Chromosome Organization. *Mol Cell* **67**:334-347.e5. doi:10.1016/j.molcel.2017.06.010

Elbatsh AMO, Haarhuis JHI, Petela N, Chapard C, Fish A, Celie PH, Stadnik M, Ristic D, Wyman C, Medema RH, Nasmyth K, Rowland BD. 2016. Cohesin Releases DNA through Asymmetric ATPase-Driven Ring Opening. *Mol Cell* **61**:575–588. doi:10.1016/j.molcel.2016.01.025

Evans R, O'Neill M, Pritzel A, Antropova N, Senior A, Green T, Žídek A, Bates R, Blackwell S, Yim J, Ronneberger O, Bodenstein S, Zielinski M, Bridgland A,

Potapenko A, Cowie A, Tunyasuvunakool K, Jain R, Clancy E, Kohli P, Jumper J,
Hassabis D. 2021. Protein complex prediction with AlphaFold-Multimer.
doi:10.1101/2021.10.04.463034

Fudenberg G, Imakaev M, Lu C, Goloborodko A, Abdennur N, Mirny LA. 2016. Formation
of Chromosomal Domains by Loop Extrusion. *Cell Rep* **15**:2038–49.
doi:10.1016/j.celrep.2016.04.085

Gandhi R, Gillespie PJ, Hirano T. 2006. Human Wapl is a cohesin-binding protein that
promotes sister-chromatid resolution in mitotic prophase. *Curr Biol* **16**:2406–17.
doi:10.1016/j.cub.2006.10.061

Gerlich D, Koch B, Dupeux F, Peters JM, Ellenberg J. 2006. Live-cell imaging reveals a
stable cohesin-chromatin interaction after but not before DNA replication. *Curr Biol*
16:1571–8. doi:10.1016/j.cub.2006.06.068

Gligoris TG, Scheinost JC, Burmann F, Petela N, Chan KL, Uluocak P, Beckouet F, Gruber
S, Nasmyth K, Lowe J. 2014. Closing the cohesin ring: structure and function of its
Smc3-kleisin interface. *Science* **346**:963–7. doi:10.1126/science.1256917

Gruber S, Arumugam P, Katou Y, Kuglitsch D, Helmhart W, Shirahige K, Nasmyth K. 2006.
Evidence that loading of cohesin onto chromosomes involves opening of its SMC
hinge. *Cell* **127**:523–537. doi:10.1016/j.cell.2006.08.048

Haering CH, Farcas A-M, Arumugam P, Metson J, Nasmyth K. 2008. The cohesin ring
concatenates sister DNA molecules. *Nature* **454**:297–301. doi:10.1038/nature07098

Haering CH, Löwe J, Hochwagen A, Nasmyth K. 2002. Molecular architecture of SMC
proteins and the yeast cohesin complex. *Mol Cell* **9**:773–788. doi:10.1016/s1097-
2765(02)00515-4

Haering CH, Schoffnegger D, Nishino T, Helmhart W, Nasmyth K, Löwe J. 2004. Structure
and stability of cohesin's Smc1-kleisin interaction. *Mol Cell* **15**:951–964.
doi:10.1016/j.molcel.2004.08.030

Hara K, Zheng G, Qu Q, Liu H, Ouyang Z, Chen Z, Tomchick DR, Yu H. 2014. Structure of
cohesin subcomplex pinpoints direct shugoshin-Wapl antagonism in centromeric
cohesion. *Nat Struct Mol Biol* **21**:864–70. doi:10.1038/nsmb.2880

Hartman T, Stead K, Koshland D, Guacci V. 2000. Pds5p is an essential chromosomal
protein required for both sister chromatid cohesion and condensation in
Saccharomyces cerevisiae. *J Cell Biol* **151**:613–26. doi:10.1083/jcb.151.3.613

Hassler M, Shaltiel IA, Kschonsak M, Simon B, Merkel F, Thärichen L, Bailey HJ, Macošek
J, Bravo S, Metz J, Hennig J, Haering CH. 2019. Structural Basis of an Asymmetric
Condensin ATPase Cycle. *Mol Cell* **74**:1175-1188.e9.
doi:10.1016/j.molcel.2019.03.037

Higashi TL, Eickhoff P, Sousa JS, Locke J, Nans A, Flynn HR, Snijders AP, Papageorgiou
G, O'Reilly N, Chen ZA, O'Reilly FJ, Rappsilber J, Costa A, Uhlmann F. 2020. A
Structure-Based Mechanism for DNA Entry into the Cohesin Ring. *Mol Cell* **79**:917-
933.e9. doi:10.1016/j.molcel.2020.07.013

Higashi TL, Ikeda M, Tanaka H, Nakagawa T, Bando M, Shirahige K, Kubota Y, Takisawa
H, Masukata H, Takahashi TS. 2012. The prereplication complex recruits XEco2 to
chromatin to promote cohesin acetylation in Xenopus egg extracts. *Curr Biol* **22**:977–
88. doi:10.1016/j.cub.2012.04.013

Hill L, Ebert A, Jaritz M, Wutz G, Nagasaka K, Tagoh H, Kostanova-Poliakova D, Schindler
K, Sun Q, Bonelt P, Fischer M, Peters JM, Busslinger M. 2020. Wapl repression by
Pax5 promotes V gene recombination by Igh loop extrusion. *Nature* **584**:142–147.
doi:10.1038/s41586-020-2454-y

Ivanov MP, Ladurner R, Poser I, Beveridge R, Rampler E, Hudecz O, Novatchkova M,
Heriche JK, Wutz G, van der Lelij P, Kreidl E, Hutchins JR, Axelsson-Ekker H,

Ellenberg J, Hyman AA, Mechtler K, Peters JM. 2018. The replicative helicase MCM
recruits cohesin acetyltransferase ESCO2 to mediate centromeric sister chromatid
cohesion. *EMBO J* **37**. doi:10.15252/embj.201797150
Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K,
Bates R, Žídek A, Potapenko A, Bridgland A, Meyer C, Kohl SAA, Ballard AJ,
Cowie A, Romera-Paredes B, Nikolov S, Jain R, Adler J, Back T, Petersen S, Reiman
D, Clancy E, Zielinski M, Steinegger M, Pacholska M, Berghammer T, Bodenstein S,
Silver D, Vinyals O, Senior AW, Kavukcuoglu K, Kohli P, Hassabis D. 2021. Highly
accurate protein structure prediction with AlphaFold. *Nature* **596**:583–589.
doi:10.1038/s41586-021-03819-2
Kikuchi S, Borek DM, Otwinowski Z, Tomchick DR, Yu H. 2016. Crystal structure of the
cohesin loader Scc2 and insight into cohesinopathy. *Proc Natl Acad Sci U S A*
113:12444–12449. doi:10.1073/pnas.1611333113
Kim Y, Shi Z, Zhang H, Finkelstein IJ, Yu H. 2019. Human cohesin compacts DNA by loop
extrusion. *Science* **366**:1345–1349. doi:10.1126/science.aaz4475
Klein F, Mahr P, Galova M, Buonomo SB, Michaelis C, Nairz K, Nasmyth K. 1999. A
central role for cohesins in sister chromatid cohesion, formation of axial elements,
and recombination during yeast meiosis. *Cell* **98**:91–103. doi:10.1016/S0092-
8674(00)80609-1
Kueng S, Hegemann B, Peters BH, Lipp JJ, Schleiffer A, Mechtler K, Peters JM. 2006. Wapl
controls the dynamic association of cohesin with chromatin. *Cell* **127**:955–67.
doi:10.1016/j.cell.2006.09.040
Ladurner R, Kreidl E, Ivanov MP, Ekker H, Idarraga-Amado MH, Busslinger GA, Wutz G,
Cisneros DA, Peters JM. 2016. Sororin actively maintains sister chromatid cohesion.
EMBO J **35**:635–53. doi:10.15252/embj.201592532
Lee B-G, Merkel F, Allegretti M, Hassler M, Cawood C, Lecomte L, O'Reilly FJ, Sinn LR,
Gutierrez-Escribano P, Kschonsak M, Bravo S, Nakane T, Rappsilber J, Aragon L,
Beck M, Löwe J, Haering CH. 2020. Cryo-EM structures of holo condensin reveal a
subunit flip-flop mechanism. *Nat Struct Mol Biol* **27**:743–751. doi:10.1038/s41594-
020-0457-x
Lee B-G, Rhodes J, Löwe J. 2022. Clamping of DNA shuts the condensin neck gate. *Proc
Natl Acad Sci U S A* **119**:e2120006119. doi:10.1073/pnas.2120006119
Lee BG, Roig MB, Jansma M, Petela N, Metson J, Nasmyth K, Lowe J. 2016. Crystal
Structure of the Cohesin Gatekeeper Pds5 and in Complex with Kleisin Scc1. *Cell
Rep* **14**:2108–2115. doi:10.1016/j.celrep.2016.02.020
Li Y, Haarhuis JHI, Sedeño Cacciatore Á, Oldenkamp R, van Ruiten MS, Willems L,
Teunissen H, Muir KW, de Wit E, Rowland BD, Panne D. 2020. The structural basis
for cohesin-CTCF-anchored loops. *Nature* **578**:472–476. doi:10.1038/s41586-019-
1910-z
Li Y, Muir KW, Bowler MW, Metz J, Haering CH, Panne D. 2018. Structural basis for Scc3-
dependent cohesin recruitment to chromatin. *Elife* **7**. doi:10.7554/eLife.38356
Liu H, Rankin S, Yu H. 2013. Phosphorylation-enabled binding of SGO1-PP2A to cohesin
protects sororin and centromeric cohesion during mitosis. *Nat Cell Biol* **15**:40–9.
doi:10.1038/ncb2637
Minamino M, Bouchoux C, Canal B, Diffley JFX, Uhlmann F. 2023. A replication fork
determinant for the establishment of sister chromatid cohesion. *Cell* **186**:837-849 e11.
doi:10.1016/j.cell.2022.12.044
Mitter M, Gasser C, Takacs Z, Langer CCH, Tang W, Jessberger G, Beales CT, Neuner E,
Ameres SL, Peters JM, Goloborodko A, Micura R, Gerlich DW. 2020. Conformation

of sister chromatids in the replicated human genome. *Nature* **586**:139–144.
doi:10.1038/s41586-020-2744-4
Moldovan GL, Pfander B, Jentsch S. 2006. PCNA controls establishment of sister chromatid
cohesion during S phase. *Mol Cell* **23**:723–32. doi:10.1016/j.molcel.2006.07.007
Mota IP, Galova M, Schleiffer A, Nguyen T-T, Kovacikova I, Nishiyama T, Gregan J, Peters
J-M, Schlögelhofer P. 2022. SORORIN is an evolutionary conserved antagonist of
WAPL. doi:10.1101/2022.10.24.513534
Nishiyama T, Ladurner R, Schmitz J, Kreidl E, Schleiffer A, Bhaskara V, Bando M,
Shirahige K, Hyman AA, Mechtler K, Peters JM. 2010. Sororin mediates sister
chromatid cohesion by antagonizing Wapl. *Cell* **143**:737–49.
doi:10.1016/j.cell.2010.10.031
Oliveira RA, Hamilton RS, Pauli A, Davis I, Nasmyth K. 2010. Cohesin cleavage and Cdk
inhibition trigger formation of daughter nuclei. *Nat Cell Biol* **12**:185–92.
doi:10.1038/ncb2018
Ouyang Z, Zheng G, Song J, Borek DM, Otwinowski Z, Brautigam CA, Tomchick DR,
Rankin S, Yu H. 2013. Structure of the human cohesin inhibitor Wapl. *Proc Natl*
Acad Sci U S A **110**:11355–60. doi:10.1073/pnas.1304594110
Ouyang Z, Zheng G, Tomchick DR, Luo X, Yu H. 2016. Structural Basis and IP6
Requirement for Pds5-Dependent Cohesin Dynamics. *Mol Cell* **62**:248–259.
doi:10.1016/j.molcel.2016.02.033
Panizza S, Tanaka T, Hochwagen A, Eisenhaber F, Nasmyth K. 2000. Pds5 cooperates with
cohesin in maintaining sister chromatid cohesion. *Curr Biol* **10**:1557–64.
doi:10.1016/s0960-9822(00)00854-x
Parelho V, Hadjur S, Spivakov M, Leleu M, Sauer S, Gregson HC, Jarmuz A, Canzonetta C,
Webster Z, Nesterova T, Cobb BS, Yokomori K, Dillon N, Aragon L, Fisher AG,
Merkenschlager M. 2008. Cohesins functionally associate with CTCF on mammalian
chromosome arms. *Cell* **132**:422–433. doi:10.1016/j.cell.2008.01.011
Petela NJ, Gligoris TG, Metson J, Lee B-G, Voulgaris M, Hu B, Kikuchi S, Chapard C, Chen
W, Rajendra E, Srinivisan M, Yu H, Löwe J, Nasmyth KA. 2018. Scc2 Is a Potent
Activator of Cohesin's ATPase that Promotes Loading by Binding Scc1 without Pds5.
Mol Cell **70**:1134-1148.e7. doi:10.1016/j.molcel.2018.05.022
Petela NJ, Gonzalez Llamazares A, Dixon S, Hu B, Lee BG, Metson J, Seo H, Ferrer-
Harding A, Voulgaris M, Gligoris T, Collier J, Oh BH, Lowe J, Nasmyth KA. 2021.
Folding of cohesin's coiled coil is important for Scc2/4-induced association with
chromosomes. *Elife* **10**. doi:10.7554/eLife.67268
Rankin S, Ayad NG, Kirschner MW. 2005. Sororin, a substrate of the anaphase-promoting
complex, is required for sister chromatid cohesion in vertebrates. *Mol Cell* **18**:185–
200. doi:10.1016/j.molcel.2005.03.017
Rao SSP, Huang SC, Glenn St Hilaire B, Engreitz JM, Perez EM, Kieffer-Kwon KR,
Sanborn AL, Johnstone SE, Bascom GD, Bochkov ID, Huang X, Shamim MS, Shin J,
Turner D, Ye Z, Omer AD, Robinson JT, Schlick T, Bernstein BE, Casellas R, Lander
ES, Aiden EL. 2017. Cohesin Loss Eliminates All Loop Domains. *Cell* **171**:305-320
e24. doi:10.1016/j.cell.2017.09.026
Roig MB, Lowe J, Chan KL, Beckouet F, Metson J, Nasmyth K. 2014. Structure and
function of cohesin's Scc3/SA regulatory subunit. *FEBS Lett* **588**:3692–702.
doi:10.1016/j.febslet.2014.08.015
Rolef Ben-Shahar T, Heeger S, Lehane C, East P, Flynn H, Skehel M, Uhlmann F. 2008.
Eco1-dependent cohesin acetylation during establishment of sister chromatid
cohesion. *Science* **321**:563–6. doi:10.1126/science.1157774

Rowland BD, Roig MB, Nishino T, Kurze A, Uluocak P, Mishra A, Beckouet F, Underwood P, Metson J, Imre R, Mechtler K, Katis VL, Nasmyth K. 2009. Building sister chromatid cohesion: smc3 acetylation counteracts an antiestablishment activity. *Mol Cell* **33**:763–74. doi:10.1016/j.molcel.2009.02.028

Schleiffer A, Kaitna S, Maurer-Stroh S, Glotzer M, Nasmyth K, Eisenhaber F. 2003. Kleisins: a superfamily of bacterial and eukaryotic SMC protein partners. *Mol Cell* **11**:571–575. doi:10.1016/s1097-2765(03)00108-4

Shaltiel IA, Datta S, Lecomte L, Hassler M, Kschonsak M, Bravo S, Stober C, Ormanns J, Eustermann S, Haering CH. 2022. A hold-and-feed mechanism drives directional DNA loop extrusion by condensin. *Science* **376**:1087–1094. doi:10.1126/science.abm4012

Shi Z, Gao H, Bai XC, Yu H. 2020. Cryo-EM structure of the human cohesin-NIPBL-DNA complex. *Science* **368**:1454–1459. doi:10.1126/science.abb0981

Shintomi K, Hirano T. 2009. Releasing cohesin from chromosome arms in early mitosis: opposing actions of Wapl-Pds5 and Sgo1. *Genes Dev* **23**:2224–36. doi:10.1101/gad.1844309

Soh Y-M, Bürmann F, Shin H-C, Oda T, Jin KS, Toseland CP, Kim C, Lee H, Kim SJ, Kong M-S, Durand-Diebold M-L, Kim Y-G, Kim HM, Lee NK, Sato M, Oh B-H, Gruber S. 2015. Molecular basis for SMC rod formation and its dissolution upon DNA binding. *Mol Cell* **57**:290–303. doi:10.1016/j.molcel.2014.11.023

Srinivasan M, Petela NJ, Scheinost JC, Collier J, Voulgaris M, M BR, Beckouet F, Hu B, Nasmyth KA. 2019. Scc2 counteracts a Wapl-independent mechanism that releases cohesin from chromosomes during G1. *Elife* **8**. doi:10.7554/eLife.44736

Srinivasan M, Scheinost JC, Petela NJ, Gligoris TG, Wissler M, Ogushi S, Collier JE, Voulgaris M, Kurze A, Chan K-L, Hu B, Costanzo V, Nasmyth KA. 2018. The Cohesin Ring Uses Its Hinge to Organize DNA Using Non-topological as well as Topological Mechanisms. *Cell* **173**:1508-1519.e18. doi:10.1016/j.cell.2018.04.015

Tedeschi A, Wutz G, Huet S, Jaritz M, Wuensche A, Schirghuber E, Davidson IF, Tang W, Cisneros DA, Bhaskara V, Nishiyama T, Vaziri A, Wutz A, Ellenberg J, Peters JM. 2013. Wapl is an essential regulator of chromatin structure and chromosome segregation. *Nature* **501**:564–8. doi:10.1038/nature12471

Toth A, Ciosk R, Uhlmann F, Galova M, Schleiffer A, Nasmyth K. 1999. Yeast cohesin complex requires a conserved protein, Eco1p(Ctf7), to establish cohesion between sister chromatids during DNA replication. *Genes Dev* **13**:320–33. doi:10.1101/gad.13.3.320

Uhlmann F, Wernic D, Poupart MA, Koonin EV, Nasmyth K. 2000. Cleavage of cohesin by the CD clan protease separin triggers anaphase in yeast. *Cell* **103**:375–86. doi:10.1016/s0092-8674(00)00130-6

Unal E, Heidinger-Pauli JM, Kim W, Guacci V, Onn I, Gygi SP, Koshland DE. 2008. A molecular determinant for the establishment of sister chromatid cohesion. *Science* **321**:566–9. doi:10.1126/science.1157880

van Ruiten MS, van Gent D, Sedeno Cacciatore A, Fauster A, Willems L, Hekkelman ML, Hoekman L, Altelaar M, Haarhuis JHI, Brummelkamp TR, de Wit E, Rowland BD. 2022. The cohesin acetylation cycle controls chromatin loop length through a PDS5A brake mechanism. *Nat Struct Mol Biol* **29**:586–591. doi:10.1038/s41594-022-00773-z

Varadi M, Anyango S, Deshpande M, Nair S, Natassia C, Yordanova G, Yuan D, Stroe O, Wood G, Laydon A, Žídek A, Green T, Tunyasuvunakool K, Petersen S, Jumper J, Clancy E, Green R, Vora A, Lutfi M, Figurnov M, Cowie A, Hobbs N, Kohli P, Kleywegt G, Birney E, Hassabis D, Velankar S. 2022. AlphaFold Protein Structure Database: massively expanding the structural coverage of protein-sequence space

with high-accuracy models. *Nucleic Acids Res* **50**:D439–D444.
doi:10.1093/nar/gkab1061
Wendt KS, Yoshida K, Itoh T, Bando M, Koch B, Schirghuber E, Tsutsumi S, Nagae G,
Ishihara K, Mishiro T, Yahata K, Imamoto F, Aburatani H, Nakao M, Imamoto N,
Maeshima K, Shirahige K, Peters J-M. 2008. Cohesin mediates transcriptional
insulation by CCCTC-binding factor. *Nature* **451**:796–801. doi:10.1038/nature06634
Wu FM, Nguyen JV, Rankin S. 2011. A conserved motif at the C terminus of sororin is
required for sister chromatid cohesion. *J Biol Chem* **286**:3579–86.
doi:10.1074/jbc.M110.196758
Wutz G, Ladurner R, St Hilaire BG, Stocsits RR, Nagasaka K, Pignard B, Sanborn A, Tang
W, Varnai C, Ivanov MP, Schoenfelder S, van der Lelij P, Huang X, Durnberger G,
Roitinger E, Mechtler K, Davidson IF, Fraser P, Lieberman-Aiden E, Peters JM.
2020. ESCO1 and CTCF enable formation of long chromatin loops by protecting
cohesin(STAG1) from WAPL. *Elife* **9**. doi:10.7554/eLife.52091
Yamada T, Tahara E, Kanke M, Kuwata K, Nishiyama T. 2017. Drosophila Dalmatian
combines sororin and shugoshin roles in establishment and protection of cohesion.
EMBO J **36**:1513–1527. doi:10.15252/embj.201695607
Yatskevich S, Rhodes J, Nasmyth K. 2019. Organization of Chromosomal DNA by SMC
Complexes. *Annu Rev Genet* **53**:445–482. doi:10.1146/annurev-genet-112618-043633
Zhai Y, Cheng E, Wu H, Li N, Yung PY, Gao N, Tye BK. 2017. Open-ringed structure of the
Cdt1-Mcm2-7 complex as a precursor of the MCM double hexamer. *Nat Struct Mol*
Biol **24**:300–308. doi:10.1038/nsmb.3374
Zhang Y, Zhang X, Dai HQ, Hu H, Alt FW. 2022. The role of chromatin loop extrusion in
antibody diversification. *Nat Rev Immunol* **22**:550–566. doi:10.1038/s41577-022-
00679-3
1728