Multiple behavioural mechanisms shape development in a highly social cichlid fish

Isabela P. Harmon[a], Emily A. McCabe[a], Madeleine R. Vergun[a], Julia Weinstein[b], Hannah L.

Graves[b], Deijah D. Bradley[c], Clare M. Boldt[a], June Lee[b], Jessica M. Maurice[a], Tessa K.

Solomon-Lane[a-c]*

[a]Scripps College, Claremont, CA

[b]Claremont McKenna College, Claremont, CA

[c]Pitzer College, Claremont, CA

*Corresponding author

Tessa Solomon-Lane

W.M. Keck Science Department

Scripps, Pitzer, and Claremont McKenna Colleges

925 N. Mills Ave, Claremont, CA 91711, USA

TSolomonLane@kecksci.claremont.edu

+1 (909) 607-7857

ABSTRACT

Early-life social experiences shape adult phenotype, yet the underlying behavioural mechanisms remain poorly understood. We manipulated early-life social experience in the highly social African cichlid fish *Astatotilapia burtoni* to investigate the effects on behaviour and neuroendocrine stress axis function. Juveniles experienced different numbers of early-life social partners in stable pairs (1 partner), stable groups (6 fish; 5 partners), and socialized pairs (a novel fish was exchanged every 5 days; 5 partners). Treatments differed in group size (groups vs. pairs) and stability (stable vs. socialized). We then measured behaviour in multiple contexts and collected water-borne cortisol. We found effects of treatment on behaviour across all assays: open field exploration, social cue investigation, dominant behaviour, and subordinate behaviour. Cortisol did not differ across treatments. Principal components (PC) analysis revealed robust co-variation of behaviour across contexts, including with cortisol, to form behavioural syndromes sensitive to early-life social experience. PC1 (25.1%) differed by numbers of social partners: juveniles with more social partners were more active during the social cue investigation, spent less time in the territory, and were more interactive as dominants. Differences in PC5 (8.5%) were based on stability: socialized pairs were more dominant, spent less time in and around the territory, were more socially investigative, and had lower cortisol than stable groups or pairs. Behaviour observations in the home tanks provided further insights into the behavioural mechanisms underlying these effects. These results contribute to our understanding of how early-life social experiences are accrued and exert strong, lasting effects on adult phenotype.

INTRODUCTION

Early-life social environments and experiences are potent drivers of developmental plasticity for social species and, as a result, can have strong, long-term effects on organismal phenotype (Bateson, 2001; Bateson et al., 2004; Kuijper & Johnstone, 2019; Taborsky, 2017; Weaver, 2009). Early-life social effects have been documented across vertebrate taxa (e.g., Arnold & Taborsky, 2010; Bölting & von Engelhardt, 2017; Champagne & Curley, 2005; Moretz et al., 2007; Perkeybile & Bales, 2017; White et al., 2010), yet the specific attributes of early social environments and experiences that cause phenotypic changes are often unidentified (Kasumovic, 2013; Taborsky, 2016). Social interactions and stimuli can make up a substantial part of juvenile experience (Kohn, 2019). Depending on the species and group structure, early social interactions can involve parents (maternal, paternal, or biparental care) (Champagne & Curley, 2005; McClelland et al., 2011; Perkeybile et al., 2013), parental helpers (Arnold & Taborsky, 2010; Taborsky et al., 2012), siblings (Branchi et al., 2013; Buist et al., 2013; D'Andrea et al., 2007; Monclús et al., 2012), peers (Ahloy Dallaire & Mason, 2017; Bölting & von Engelhardt, 2017; Förster & Cords, 2005; Moretz et al., 2007; Weinstein et al., 2014), and other members of the group (Bray, Murray, et al., 2021; Förster & Cords, 2005; Jin et al., 2015), as well as observations of others interacting (Clay & de Waal, 2013; Desjardins et al., 2012; Oliveira et al., 1998). Identifying the specific, proximate causes—the behavioural mechanisms—is critical to understanding how gene-by-environment interactions shape processes of developmental plasticity and behavioural developmental trajectories. Given the fitness and health consequences of social behaviour (Bennett et al., 2006; Meyer-Lindenberg & Tost, 2012; Silk, 2007; Solomon-Lane et al., 2015; Wilson, 1980), the developmental origins of adult behavioural phenotype are particularly important to understand, including as a target for natural selection.

Manipulating the early social environment is a common approach to studying early-life

social effects. For example, rearing animals in groups of different sizes and/or complexities often

has significant effects on development and phenotype (reviewed in Taborsky, 2016). In ravens

(*Corvus corax*), family size affected social attentiveness (Gallego-Abenza et al., 2022); in zebra

finches (*Taeniopygia guttata*), rearing group size and age diversity of early-life flocks affected

courtship and aggressive behaviour, as well as plumage development (Bölting & von Engelhardt,

2017); in mice, communal nesting affected social skills and neuroendocrine function, with

separate effects of maternal care and peer interactions (Branchi et al., 2009, 2013); and in

Daffodil cichlids (*Neolamprologus pulcher*), the presence of parents and brood helpers increased

social competence (Arnold & Taborsky, 2010; Taborsky et al., 2012). In general, larger social

groups are more complex than smaller groups, and early exposure to social complexity tends to

benefit social skills and competence (e.g., Branchi et al., 2009, 2013; Fischer et al., 2015; White

et al., 2010). However, individuals accrue different social experiences, even within shared

environments. For example, mouse pups in mixed-age, communal nests interact with siblings at

varying rates and receive different levels of maternal care (Branchi et al., 2013); infant blue

monkeys (*Cercopithecus mitis stuhlmanni*) receive varying rates of allomaternal care, and from

different non-mothers in the group (Förster & Cords, 2005); immature male chimpanzees

socially associate with adult males at different rates (Bray, Feldblum, et al., 2021); and young

male long-tailed manakins occupy varied positions in the social network (McDonald, 2007). This

variation can have long-term effects on social decision-making and behaviour (e.g., Branchi et

al., 2013; Bray, Murray, et al., 2021; McDonald, 2007). Therefore, to identify the behavioural

mechanisms underlying behavioural development, it is necessary to observe individuals in the

rearing environment and test mechanistic hypotheses directly by manipulating the quality and/or

quantity of social experiences (Kasumovic, 2013; Taborsky, 2016).

Early social experiences exert long-term effects on organismal phenotype through

persistent changes in underlying neural and neuroendocrine mechanisms (e.g., Antunes et al.,

2021; Branchi et al., 2013; Champagne & Curley, 2005; McClelland et al., 2011). The

neuroendocrine stress axis, or hypothalamic-pituitary-adrenal (interrenal in fish; HPA/I) axis, is a

highly-conserved mechanism underlying early-life effects, including social effects (Banerjee et

al., 2012; Champagne & Curley, 2005; Crespi & Denver, 2005; Ensminger et al., 2018; Francis

et al., 1999; Jonsson & Jonsson, 2014; McClelland et al., 2011; Taborsky, 2016). The HPA/I axis

translates environmental conditions and experiences into coordinated physiological and

behavioural responses through a neuroendocrine cascade that initiates in response to an

environmental stressor. Detection of a stressor, which can include any external condition that

disrupts or threatens to disrupt homeostasis, leads to the release of corticotropin-releasing factor

(CRF) from the hypothalamus. The pituitary responds to CRF with the release of

adrenocorticotropic (ACTH) hormone, which signals for the adrenal (or interrenal) glands to

release glucocorticoids (cortisol in fishes) into circulation (Denver, 2009; Lowry & Moore, 2006;

Wendelaar Bonga, 1997). Early-life social experiences shape HPA/I axis function in multiple

ways (Champagne & Curley, 2005; Francis et al., 1999; McClelland et al., 2011; Spencer, 2017;

Taborsky, 2016; Turecki & Meaney, 2016). For example, peer-reared rhesus macaques (*Macaca*

mulatta) had higher baseline and stress-induced levels of ACTH and cortisol, as well as a more

reactive HPA axis, compared to maternal-reared juveniles (Stevens et al., 2009). Similarly, zebra

finch chicks reared with only their fathers had hyperresponsive HPA axes, along with altered

levels of neural glucocorticoid receptor (GR) and mineralocorticoid receptor (MR) expression,

compared to biparental rearing (Banerjee et al., 2012). In *N. pulcher* cichlids, rearing with or

without parents affected *gr1* expression in the telencephalon (Nyman et al., 2018), and rearing

with or without brood helpers altered neural *crf* and *gr* expression, as well as the ratio of *mr* to

gr1 (Taborsky et al., 2013). The HPA/I axis is also an important source of individual variation in

social behaviour (Boogert et al., 2014; Dettmer et al., 2017; Farine et al., 2015; Freeman et al.,

2021; Pryce et al., 2011; Reyes-Contreras et al., 2019; Sih, 2011) and health (e.g., Turecki &

Meaney, 2016).

In this study, we used Burton's Mouthbrooder (*Astatotilapia burtoni*), a highly social

cichlid fish and model system in social neuroscience (Fernald & Maruska, 2012; Hofmann,

2003; Stevenson et al., 2018), to investigate the behavioural mechanisms through which early-

life social experiences affect behaviour and HPI axis function. The vast majority of research on

this species has focused on adults, which form mixed-sex, hierarchical social communities.

Dominant males are colourful, territorial, and reproductively active. In contrast, subordinate

males are drab in coloration, shoal with females, and are reproductively suppressed. Social status

is socially-regulated, and males regularly transition between dominant and subordinate positions

(Fernald & Maruska, 2012; Hofmann, 2003). Juveniles also form status relationships (Solomon-

Lane et al., 2022), and both juveniles and adults express a suite of highly-conserved social

behaviours (Fernald & Hirata, 1979; Fernald & Maruska, 2012; Solomon-Lane et al., 2022;

Weitekamp & Hofmann, 2017). Although sex and social status have strong effects on adult

behaviour, individual variation persists, including in the frequency and quality of behavioural

expression, such as aggression, territoriality, courtship, cooperation, reproductive behaviour, and

maternal behaviour (Alward et al., 2021; Friesen et al., 2022; Fulmer et al., 2017; Kidd et al.,

2013; Maruska, Becker, et al., 2013; Renn et al., 2009; Weitekamp & Hofmann, 2017); tenure in

a dominant vs. subordinate role (Hofmann et al., 1999); female mate choice (Kidd et al., 2013);

social learning (Rodriguez-Santiago et al., 2020); and cognition (Wallace & Hofmann, 2021).

HPI axis function also varies among adults (Alcazar et al., 2016; Chen & Fernald, 2008; Clement

et al., 2005; Dijkstra et al., 2017; Greenwood et al., 2003; Korzan et al., 2014; Maruska et al.,

2022) and is highly responsive to social context, such as social status and changes in status

(Carpenter et al., 2014; Chen & Fernald, 2008; Clement et al., 2005; Fox et al., 1997; Huffman et

al., 2015; Korzan et al., 2014; Maruska et al., 2022; Maruska, Zhang, et al., 2013; Parikh et al.,

2006), group social dynamics (Maguire et al., 2021), social habituation (Weitekamp & Hofmann,

2017), and in response to an intruder (Alcazar et al., 2016; Weitekamp et al., 2017).

Developmental plasticity may be an important source of individual variation in behaviour

and HPI axis function among adults (Fernald & Hirata, 1979; Fraley & Fernald, 1982; Solomon-

Lane & Hofmann, 2019). We have previously demonstrated early-life social effects in juvenile

A. burtoni in response to early-life social group size. Juveniles reared in social groups (16 fish)

developed to be more active, more socially interactive, and less submissive in a subordinate role

compared to pair-reared juveniles. Whole brain gene expression related to HPI axis function was

also altered, including elevated GR1a expression in group-reared animals and tight co-expression

of candidate genes (including GR1a, GR1b, GR2, MR, and CRF) in pair-reared, but not group-

reared or isolated, juveniles (Solomon-Lane & Hofmann, 2019). Juveniles in the previous study

were not observed in the rearing environment, and there are multiple, possible behavioural

mechanisms responsible for these effects (Solomon-Lane & Hofmann, 2019; Taborsky, 2016).

Individuals reared in larger and/or more complex groups may interact socially at higher rates,

experience a greater diversity of types of social interaction (e.g., affiliative, aggressive,

cooperative, etc.), gain experience in multiple social roles (e.g., dominant and subordinate,

younger and older) (Chase et al., 2022; Solomon-Lane et al., 2022; Williamson et al., 2016),

have more social partners, have social partners of varied identities / traits (e.g., age, body size,

life history stage diversity) (Arnold & Taborsky, 2010; Bölting & von Engelhardt, 2017; Branchi

& Alleva, 2006; Taborsky et al., 2012; White et al., 2010), experience interactions involving

more actors simultaneously (i.e., 3-way, or more, interactions), and learn indirectly by watching

others (e.g., Desjardins et al., 2012; Oliveira et al., 1998)). For juvenile *A. burtoni*, group size

and the relative body sizes of individuals affects social behaviour and group structure. For

example, pairs and triads containing same-sized fish are less hierarchical (Solomon-Lane et al.,

2022).

170   In this study, we tested the hypothesis that the number of social partners experienced by

juvenile *A. burtoni* during early-life is a behavioural mechanism that shapes behavioural

development and HPI axis signalling. The number of social partners is one clear way that groups

and pairs differ (Solomon-Lane & Hofmann, 2019). There is also evidence from other species

that this may be a meaningful attribute of social experience that varies across individuals.

Assortative social interactions have been observed in a variety of species. For example, more

socially interactive juvenile yellow-bellied marmots (*Marmota flaviventris*) had more novel

social partners (Monclús et al., 2012); juvenile male geladas (*Theropithecus gelada*) had more

novel play partners than juvenile females (Barale et al., 2015); juvenile rhesus macaques exhibit

variation in the number of friendships initiated and reciprocated (Weinstein et al., 2014); and

bold three-spined sticklebacks (*Gasterosteus aculeatus*) had a larger number of social contacts

compared to shy fish (Pike et al., 2008). For juvenile *A. burtoni* in pairs and triads, individuals

are not equally likely to initiate interactions (Solomon-Lane et al., 2022), suggesting that in

larger, more naturalistic groups, social network connectivity may vary considerably, as it does in

adults (Maguire et al., 2021). Direct manipulations of social partner numbers can also affect

social behavioural development. For example, juvenile male brown-headed cowbirds reared in

dynamic flocks outcompeted males reared in stable flocks of the same size (White et al., 2010).

To manipulate the number of social partners, we reared juveniles in stable pairs (2 fish, 1

partner each), in stable groups (6 fish, 5 partners each), or in socialized pairs, in which one

member of the pair was changed 5 times over the course of the ~1 month experiment (2 fish at a

time in the pair, 5 total partners each) (Figure 1). This design also allowed us to compare the

effects of group size (pair vs. group) and social stability (stable vs. socialized). We observed

social interactions in the different rearing environments to quantify social experience and then

tested the effects on individual social behaviour through a series of behaviour assays. We

hypothesized that the number of social partners would be the strongest influence on behavioural

phenotype. If the number of early-life social partners was an important behavioural mechanism

underlying group- vs. pair-reared development in Solomon-Lane & Hofmann (2019), then in this

study, we predict that juveniles from the stable groups and socialized pairs will be more active

and socially interactive in novel social context, as well as less submissive in a subordinate role.

To test the hypothesis that early-life social experience affects HPI axis signalling, we collected

water-borne cortisol following the individual behaviour assays. By investigating specific

behavioural mechanisms underlying behavioural development and neuroendocrine function, this

research can provide insights into developmental plasticity processes and uncover the origins of

adult phenotype.

METHODS

Animals

The juvenile *A. burtoni* used in this experiment were bred in the laboratory. The breeding

adults are a laboratory population that descended ~65-70 generations from a wild caught stock

from Lake Tanganyika (Fernald & Hirata, 1977). The adults are housed in naturalistic, mixed-

sex breeding communities. Dominant males court females to lay eggs in his territory, after which

she immediately scoops up the eggs into her mouth, where the male fertilizes them. The

developing larvae are brooded in the female's buccal cavity for ~14 days before being released.

In the wild, and under some laboratory conditions, females display maternal behaviour by

guarding their offspring for 10 days or more after initially releasing the free-swimming fry from

her mouth (Renn et al., 2009).

In this study, 86 fry were removed from the buccal cavities of 7 mothers approximately

5-7 days after fertilization. We selected fry at this early developmental stage before overt social

interactions occurred (Fraley & Fernald, 1982) to ensure that the scope of our experiment

captured meaningful social experience as early as possible during development. Individual

broods were then placed in shallow water in a petri dish, and a digital image was taken with a

ruler for scale. ImageJ was used to measure standard length (SL, mm) (Schneider et al., 2012),

from the tip of the jaw to the caudal peduncle. Broods ranged in size from 4-21 fry (mean: 12.43

$\pm$ 2.84 fry, median: 15), and fry size ranged from 4.59-8.13 mm SL (mean: 5.96 ± 0.095 mm,

median: 5.81 mm). See Supplemental Table 1 for mean, median, minimum, and maximum SL of

fry per brood. All 7 broods were then placed briefly (~30 min) in a common bucket to intermix.

Given their developmental stage, social interactions were highly unlikely to occur, but visual,

chemosensory, and tactile social sensory cues were likely exchanged. We used a hand net to

remove individuals from the bucket and haphazardly assigned them to different home tanks and

treatment groups (stable groups, socialized pairs, or stable pairs; see below for descriptions).

Methylene blue was added to the water to reduce the incidence of fungal infection during this

time. After 7 days, we confirmed visually that the fry were all mobile and their yolks had been

fully absorbed. The sex ratio of fish in this experiment is not known because sex cannot be

determined anatomically until reproductive maturation. The sex ratio of *A. burtoni* broods (a

laboratory population descended from wild-caught fish) is approximately 1:1 (Heule et al.,

2014).

Early-life social experiences

Fry were reared for 32 or 35 days (depending on the behavioural test day, see below) in

one of three conditions to manipulate early-life social experience: stable groups of 6 fish (n=6

groups), stable pairs of 2 fish (n=7 pairs), and socialized pairs of 2 fish (n=8 pairs) (Figure 1).

These treatment groups were designed to manipulate the number of early-life social partners. In

the stable groups, every individual had 5 social partners. In the stable pairs, every individual had

1 social partner. In the socialized pairs, 1 individual was removed from the tank using a hand net

every 5 days and was replaced with a novel partner. The fish that was removed was transferred to

another socialized pair tank with a novel partner. The social exchanges began on experimental

day 12 (all fry were sufficiently mature on day 7, followed by 5 days with the first social

partner). After a total of 4 exchanges, each of the fish in the socialized pairs were exposed to a

total of 5 social partners, the same number of partners that each individual experienced in the

stable groups. However, the size of the social group for socialized pairs was always 2 fish, the

same as in the stable pairs. As a control, on the day that fish in the socialized pairs were

transferred, we used a hand net to remove one fish from each of the stable pairs and stable

groups. That fish was immediately returned to the same home tank so that group membership in

the stable pairs and stable groups was consistent.

254   Stable and socialized pairs were housed in small acrylic tanks ($23 \times 15 \times 15$ cm), while

the stable groups were housed in larger acrylic tanks (30 x 20 x 17 cm). One small terracotta pot

(5 cm at the widest diameter) was placed in each tank, and an air stone was used to keep the

water oxygenated. There was no substrate on the bottom of the tank. White plastic dividers were

placed outside of the tanks to prevent visual contact between tanks. The fish were maintained on

a 12:12 light:dark schedule at 24°C, and they were fed Hikari Middle Larval Stage Plankton

(Pentair Aquatic Eco-Systems, Cary, NC) once a day. We mixed 0.015 g of fish food in 200 mL

of water. The mixture was actively stirred, and a transfer pipette was used to add 1 mL to the

pairs and 3 mL to the groups.

Tagging individual fish

265   To track individuals as they developed over the course of the experiment, we tagged

every fish following the first rotation in the socialized pairs. We waited until this time so the fish

would be slightly bigger, as their very small size is a challenge for existing tagging methods

(Lotrich & Meredith, 1974; Solomon-Lane & Hofmann, 2018; Thresher & Gronell, 1978). We

first lightly anesthetized each fry in 0.03 g MS-222 / L aquarium water (buffered with sodium

bicarbonate to pH 7.0 –7.5). They were then removed from the water and placed on a wet paper

towel. We placed a small amount of Alcian Blue (Fisher Scientific, Pittsburgh, PA) powder onto

the dorsal muscle of the fish and perforated the surface of the fish using a pulled capillary tube

micropipette. This method of tattooing allowed the ink to seep into the tissue of the fish. We

used different tag locations (left or right; anterior, middle, or posterior regions) on the dorsal

muscle to differentiate individuals in the same tank. In the socialized pairs, tagging was

sufficient in most tanks to identify and move the same individual each time. The persistence of

the tag varied, and in the best cases, it remained visible for 3 weeks. If the tags were not visible

in the socialized pair, the smaller individual was moved.

Survival and number of early-life social partners

The tagging procedure had an 85% survival rate. Fish that did not survive tagging or that

died at another point in the experiment were replaced with a fish of similar size and age, selected

from a community tank. The replacement fish were not included in the individual behaviour

assays conducted at the end of the experimental rearing period (see below). We tracked the exact

number of social partners for each experimental tank. All fish in the socialized pairs had 5 social

partners. Stable group fish had an average of 6.43 ± 0.14 social partners. Stable pair fish had an

average of 1.2 ± 0.13 social partners. In our analyses, we consider the treatment groups as

categorical, although there is limited variation in the number of social partners (Supplemental

Figure 1).

Behavioural Observations

<u>Home tanks</u>: We recorded social behaviour in the home tanks twice during the rearing period, on

days 20 and 25 of the experiment, at the same time of day. Cameras (Warrior 4.0, Security

Camera Warehouse, Asheville, NC) were positioned above the tanks, and with the tank lids

removed, all areas of the tank were visible except under the terracotta pot. Using Solomon Coder

(www.solomoncoder.com), 10 minutes of video was scored for approaches and displacements.

Approaches were counted anytime one fish swam directly towards any part of another fish,

within 3 body lengths. Approaches can range from affiliative to aggressive, and the response of

the approached fish provides insight into the degree of agonism. If the approached fish swam

away in any direction, it was scored as a displacement, which is an indication that the approach

was aggressive. Because the identification tags were not visible on the video, we counted the

total number of social interactions (approaches, displacements) for the group or pair. We also

quantified the number of times a fish entered the terracotta pot territory, defined as at least 50%

of a fish's body crossing into / under the pot. Exiting the territory was highly correlated with

entering the territory ($p<0.0001$, $r^2=0.97$); therefore, we only present data for entering. To

measure social grouping, we captured screenshots from the video every 30 s for a total of 20 time

points. Distance was measured between every dyad, from a focal point in the centre of the head,

using ImageJ (Schneider et al., 2012). The researchers scoring these behaviours were unaware of

the treatment of the pairs (socialized vs. stable), but the stable groups were easily recognizable.

For behaviour and distance, we analysed the raw values (behaviours per minute and distance), as

well as scaled these values to account for differences in group size (behaviours per minute

divided by 2 for the pairs and divided by 6 for the groups) and tank size (distance divided by the

hypotenuse of the tank) among treatments.

Individual Behaviour Assays: After the 32 or 35-day rearing period, we measured individual

behaviour in a series of assays, including open field exploration, social cue investigation,

dominant behaviour, and subordinate behaviour. Every fish in the socialized (final sample size:

n=9) and stable pairs (final sample size: n=10) was included in the individual assays. Three of

the 6 fish from each stable group were selected randomly to be included in the individual assays

using a random number generator (final sample size: n=14). The procedures have been described

previously in detail (Solomon-Lane & Hofmann, 2019). Briefly, these assays (or similar) are

used across species to assess different elements of behavioural phenotype, including behaviours

affected by early-life social experience (Sih, 2011; Sih et al., 2015; Solomon-Lane & Hofmann,

2019; Taborsky, 2016). The open field exploration is used to assess locomotion and anxiety-like

behaviours (e.g., Cachat et al., 2010; Prut & Belzung, 2003). The social cue investigation is used

as a measure of social motivation or preference (e.g., Bonuti & Morato, 2018; Moy et al., 2004).

The dominant and subordinate behaviour assays are used to assess how the focal fish behaves

when in the position to be dominant or subordinate. In social groups, including for juvenile *A.*

burtoni (Solomon-Lane et al., 2022), most individuals will be subordinate to some group

members and dominant over others at a given time (e.g., Chase et al., 2022). Social status also

changes over a lifetime (Fernald & Maruska, 2012; Hofmann, 2003).

The behavioural assays took place in novel experimental aquaria, which were the same

size as the pairs' home tanks (3.27 L, 23 × 15 × 15 cm). Cameras recorded behaviour from

above, and all areas of the tank were visible except under the terracotta pot. A permanent marker

was used to delineate different zones (Figure 1B): a territory zone that contained a small

terracotta pot, a close zone, a far zone, and an investigate zone where the social cue was placed

for the social cue investigation assay. Each assay lasted for 30 min. The focal fish was alone in

the tank for the open field exploration (minutes 20-30 were analysed). For the social cue

investigation, a live, novel cue fish inside of a 20 mL glass vial was placed in the investigate

zone (see placement in Figure 1B, minutes 2-12 were analysed). In the open field and social cue

assays, we recorded the number of times a fish entered a zone and how long it spent there. After

removing the vial and cue fish, a free-swimming, novel fish that was smaller than the focal fish

was added to the tank to assess dominant behaviour (minutes 2-12 were analysed). After

removing the small fish, a free-swimming novel fish larger than the focal was added to assess

subordinate behaviour (minutes 2-12 were analysed). In the dominant and subordinate behaviour

assays, we scored approaches and displacements. The researchers scoring these behaviours were

unaware of the treatment of the focal fish.

From the approaches and displacements in the dominant and subordinate behaviour

assays, we calculated two additional social measures using the *compete* package in R (Curley,

2016). First, we calculated an adjusted version of David's Score, which is a dominance index.

Rather than categorizing all social interactions as a one-dimensional "win" or "loss," we

incorporated both approaches and displacements, such that the agonistic outcome of individual i

in interaction with another individual j is the number of times that i displaces j, divided by the

total number of interactions (approaches) between i and j (i.e., i displaces j / (i approaches j + j

approaches i)). The rest of the calculations were done as described in Gammell et al. (2003). See

the Supplemental Information in Solomon-Lane et al. (2022) for the R code. Second, we

calculated directional consistency index (DCI) to assess whether patterns of approaching in the

dyads are reciprocal, from perfectly reciprocal (0) to unidirectional (1). We also used the

compete package to run a randomization test to determine if the directionality was significantly

more asymmetrical (unidirectional) than expected by chance (Leiva et al., 2008).

Water-borne cortisol

Following the individual behaviour assays, we collected water-borne hormone samples.

This is a non-invasive method of quantifying steroid hormones, and values strongly correlate

with circulating hormone levels (Kidd et al., 2010). We chose cortisol, as HPI axis function is

often affected by early-life experience (e.g., Champagne & Curley, 2005; Taborsky et al., 2013)

and cortisol can correlate with behaviour and behavioural syndromes (e.g., Koolhaas et al.,

1999). Focal fish were placed individually in clean glass vials filled with 15 mL of clean

aquarium water for 2 hrs. Visual barriers were placed around the vials to limit disturbances to the

fish during the collection. After two hours, fish were removed from the vial using a hand net

rinsed in clean aquarium water, and samples were frozen at 4°C until processing. Hormones were

extracted from the water samples using 3 cc Sep-Pak Vac C18 columns (Water Associates,

Milford, MA). Columns were first primed by passing 2 mL methanol through the columns twice,

being careful not to let the columns run dry. This was followed by running 2 mL ultrapure water

through the columns twice. The thawed samples were then passed through the columns from the

sample vials using the vacuum pump. Ultrapure water (2 mL twice) was then passed through the

columns again, followed by 5 min of the vacuum to dry the columns. Finally, the hormone

samples were eluted into 13 x 100 borosilicate test tubes by passing 2 mL methanol through the

columns twice. The vacuum was used again to dry the columns. The sample in methanol was

then evaporated under a gentle stream of nitrogen at 37°C and resuspended to a volume of 200

µL per sample (5% EtOH and 95% ELISA buffer). Resuspended samples were shaken on a

multitube vortex for 1 hr before being stored at -20°C. Before measuring cortisol, samples were

thawed and shaken again on the multitube vortex for 1 hr. ELISAs were completed according to

the supplier's instructions (Cayman Chemical, Ann Arbor, Michigan).

Cortisol data are presented here as pg hormone per mL sample volume per hr sample

collection per g body weight. During hormone collection, fish excrete hormones into the water

through their gills, urine, and faeces. The gills of larger fish have a larger surface area compared

to smaller fish, thus body size can influence water-borne hormone levels. Fish size was measured

following the collection of water-borne hormones. Mass was measured on an analytical balance,

and SL was measured using ImageJ (Schneider et al., 2012) from digital images with a rule for

scale. We calculated the condition index as Fulton's K, which estimates expected mass as SL^3.

We used the mass and SL of 400 laboratory *A. burtoni* across a range of ages and sizes to

empirically calculate expected mass as: mass$=0.00002*SL^{3.02}$ (Stevenson & Woods, 2006).

Ethical Note

All research was done in compliance with the Institutional Animal Care and use Committee

(IACUC Protocol # 19-001). We used non-invasive approaches when possible, such as collecting

water-borne hormones, and we used a tested and effective anaesthetic (MS-222) for tagging to

minimize stress and pain. We took steps to minimize stress throughout the experiment, including

limiting handling time to less than 2 min when measuring size or transferring between tanks. A

total of 76 juvenile *A. burtoni* were included in this study, and fish were returned to community

tanks at the end of the experiment.

Statistical Analyses

All statistical analyses were conducted using R Studio (R version 4.2.1) (RStudio Team,

2022). Results were considered significant at the $p<0.05$ level, and averages ± standard error of

the mean are included in the text. The boxes of the box and whisker plots show the median and

the first and third quartiles. The whiskers extend to the largest and smallest observations within

or equal to 1.5 times the interquartile range. To test for the effects of early-life social experience,

we compared body size and condition index; behaviour in the home tank rearing environments

(total behaviours per minute and behaviours per fish per minute); spatial distancing in the home

tank rearing environments (raw distance and scaled to account for the different sized aquaria for

pairs vs. groups); behaviour in the open field exploration and social cue investigation (frequency

entering and time in the territory, close, far, and investigate zones); social behaviour

(approaches, displacements, approaches received, submissions) and status (David's score, DCI)

in the dominant and subordinate assays; cortisol; and principal components (see below, PCs 1-5)

across fish reared in the stable groups, socialized pairs, or stable pairs. For data that met the

assumptions of parametric statistics, we used one-way ANOVAs. The cortisol data were log

transformed first to meet the assumption of normal distribution. *Post hoc* analysis of significant

ANOVA results was calculated using Tukey HSD tests. For data that did not meet the

assumptions of parametric statistics, we used non-parametric Kruskal-Wallis tests, and Dunn's

tests with Bonferroni correction were used for *post hoc* analysis of significant results. Eta

squared is reported for the effect size of both the one-way ANVOAs and Kruskal-Wallis tests

(small effect: $0 < \eta2 < 0.01$; moderate: $0.01 < \eta2 < 0.06$; large: $0.06 < \eta2$). We used Principal

Components Analysis (PCA) to identify how behaviour in the open field exploration, social cue

investigation, dominance behaviour assay, subordinate behaviour assay, and cortisol clustered.

RESULTS

Body size and condition

To determine whether early-life experience in a stable group, socialized pair, or stable

pair affected body size and condition, we compared SL, mass, and condition index across

treatment groups. We found that mass differed significantly across treatment groups ($\chi^2_2=6.39$,

p=0.041, η^2=0.15; Supplemental Figure 2A). *Post hoc* analysis showed juveniles reared in stable

pairs were significantly heavier than those from stable groups (p=0.036). There were no

significant differences in mass between stable groups and socialized pairs (p=0.63) or between

stable pairs and socialized pairs (p=0.82). There were no significant treatment differences in SL

($F_{2,30}$=2.15, p=0.14; Supplemental Figure 2B). Because we found an effect of rearing experience

on mass, but not SL, we also asked whether condition index was affected by treatment. We

found no significant treatment differences in condition index ($F_{2,30}$=0.45, p=0.62, Supplemental

Figure 2C). Mass was also positively correlated with water-borne cortisol levels (p=0.017,

r^2=0.15, Supplemental Figure 2D); therefore, we corrected for body size in the cortisol analyses

below.

Home tank social behaviour

We observed juveniles in the stable groups, socialized pairs, and stable pairs to

understand the differences in social experience and environment among the rearing treatments.

Unsurprisingly, there were significantly more approaches and displacements in groups compared

to pairs (both stable and socialized), and there were significantly more entrances to the territory

in groups compared to stable pairs (Figure 2A). There were no differences in efficiency (Figure

2B). There were also no differences in rates of behavioural interactions per fish. Mean dyad

distance was significantly larger in groups than pairs (Figure 2C), but interestingly, the minimum

dyad distance was significantly smaller in groups than socialized pairs (Figure 2D). There were

no treatment differences in mean dyad distance when scaled for tank size. Statistics are reported

in Table 1 and Supplemental Table 2.

Open field exploration and social cue investigation

For the open field exploration and social cue investigation, we compared across treatment

groups for the frequency of entering each zone of the tank, as well as time spent in each zone.

Statistics are reported in Table 2 (Supplemental Figure 3). Overall, we found for the open field

exploration that fish from the socialized pairs entered the territory zone significantly more

frequently than stable pair fish, and fish from the stable groups spent significantly more time in

the far zone than stable pair fish. In the social cue investigation, we found that social group fish

entered the far zone significantly more frequently and spent significantly more time in the

territory and far zones, than fish reared in stable pairs.

Dominance and Subordinate Assays

To determine whether rearing experience affected social behaviour and status, we

quantified patterns of social interaction between the focal fish and a novel social partner (cue

fish). Because relative physical size often affects dominant and subordinate social dynamics for

juveniles (Solomon-Lane et al., 2022), pairing the focal fish with a smaller fish (dominance

assay) presented the focal fish with the opportunity to express social behaviours as the dominant.

In pairing the focal fish with a larger fish (subordinate assay), the focal fish can express social

behaviours as the subordinate. We found significant differences in social behaviour and status in

both the dominance and subordinate assays (statistics reported in Table 3, Figure 3). In the

dominance assay, socialized pair fish approached and displaced significantly more than stable

pair fish, and their David's Scores were significantly higher than both stable pair and stable

group fish. In the subordinate assay, socialized pair fish received significantly more approaches

than stable pair fish from the larger cue fish with which they were paired. Socialized pairs with

their cue fish also had significantly lower directional consistency than stable group or stable pair

fish.

 Cortisol

Following the behaviour assays, we collected water-borne cortisol to determine whether

hormone levels were affected by rearing experience and/or associated with behavioural

phenotype (see PCA below). We found there were no significant differences in cortisol across

treatments ($F_{2,29}$=2.30, p=0.12, Figure 4).

Integrative analysis of behaviour, social status, and cortisol

Given the treatment differences we identified across the open field exploration, social cue

investigation, dominance assay, and subordinate assay, we next used PCA to integrate across

assays. This allowed us to identify subsets of factors that together contribute to juvenile

phenotype and test if suites of factors—principal components (PCs)—differed significantly

across treatment groups. To determine which variables to include from the open field exploration

and social cue investigation (time and frequency in the territory, close, far, and investigate

zones), we ran an initial PCA with just these variables. Examining the vector plot for PC1 with

PC2 (Supplemental Figure 4), we found that nearly all of the open field exploration measures

were strongly aligned with the same measure in the social cue investigation (i.e., the vectors for

open field and social cue time in the territory zone are identical). The exceptions were for time

spent in the far and investigate zones. Therefore, to avoid unnecessary replication, we chose to

include all of the variables from the social cue investigation, in addition to time spent in the far

and investigate zones during the open field exploration. From the dominant and subordinate

assays, we included approaches, displacements, approaches received, submissions, and David's

score. Finally, we included cortisol.

Here, we focused on the first 5 PCs, which together explain 76.6% of the variation in the

data (Figure 5A). We found significant effects of rearing experience on PC1 (25.1%) and PC5

(8.48%). For PC1, stable pairs were significantly higher than stable groups, and there was a trend

for stable pairs to be higher than socialized pairs (p=0.063) (Figure 5B). Behaviours from the

dominant behaviour assay and the social cue investigation assay loaded strongly on PC1.

Approaches and displacements when in a dominant social role; time in and frequency entering

the far and investigate zones during the social cue investigation; and frequency entering the close

zone during the social cue investigation loaded strongly in the same direction. Time spent in the

territory zone during the social cue investigation loaded strongly in the opposite direction. For

PC5, socialized pairs were significantly higher than both stable groups and stable pairs (Figure

5C). David's Score from the dominance assay and time in the investigate zone during the social

cue investigation loaded strongly together in the same direction. In the opposite direction,

submissions as a dominant; cortisol; time and frequency in the close zone during the social cue

investigation; and the frequency entering the territory zone during the social cue investigation

loaded strongly together. Statistics for the PC1-PC5 treatment comparisons are in Table 4 (also

see Supplemental Figure 5).

DISCUSSION

We investigated the effects of early-life social experience, in the first month of life, on

social behaviour and neuroendocrine stress axis function in juvenile *A. burtoni*. We tested the

hypothesis that the number of novel social partners experienced during early-life is a behavioural

mechanism driving variation in development. In manipulating the number of social partners, our

experimental treatment groups—stable groups, socialized pairs, and stable pairs—also varied in

social group size (pairs vs. group) and group stability (socialized vs. stable). We present strong

evidence for early-life social effects on juvenile social behaviour, consistent with previous

studies in this species (Fernald & Hirata, 1979; Fraley & Fernald, 1982; Solomon-Lane &

Hofmann, 2019). In particular, our results support two behavioural mechanisms: the number of

social partners and social stability. Although early-life social effects are widely documented

across social species, the specific attributes of social experience that affect the mechanisms and

trajectory of developmental plasticity are not typically identified (Kasumovic, 2013; Taborsky,

2016). The social interactions and social spacing we observed in the home tanks also provide

insights into how social environments vary within and across treatments and may shape the

experiences individuals accrue during development. Causation at this level is needed because it

is these environmental elements that interact with genes dynamically over development to

influence plasticity and the emergence of adult phenotype (Kasumovic, 2013; Taborsky, 2016,

2017). Overall, it is likely that multiple behavioural mechanisms contribute and interact to shape

development (e.g., Branchi et al., 2013) and adult phenotype.

We found that manipulating early-life social experience affected behaviour across

multiple contexts, including the open field exploration, social cue investigation, dominant

behaviour assay, and subordinate behaviour assay. Principal components analysis revealed that

these effects were correlated across contexts. Principal component 1 (25.1%) and PC5 (8.5%),

which both differed significantly across treatment groups, had behaviours from multiple assays

that loaded strongly. These results support our previous finding that juvenile *A. burtoni*

behaviour can form a syndrome, and an individual's position along the syndrome continuum is

sensitive to early-life social experience (Solomon-Lane & Hofmann, 2019). A syndrome is a

population-level measure in which rank-order differences between individuals are correlated

550 across contexts and/or over time (Bell, 2007). Behavioural syndromes have been identified

551 across species and can indicate consistency in individual behaviour across contexts and/or over

552 time (Bell, 2007; Sih, Bell, & Johnson, 2004; Sih, Bell, Johnson, et al., 2004). The PC1

553 syndrome included approaches and displacements in the dominance assay loading strongly in the

554 same direction with time in and frequency entering the investigate and far zones, and frequency

555 entering the close zone, during the social cue investigation. Time in the territory zone during the

556 social cue investigation loaded in the opposite direction. This syndrome is highly similar to the

557 one we identified after rearing juvenile *A. burtoni* in pairs or social groups of 16 fish (Solomon-

558 Lane & Hofmann, 2019).

559   The treatment differences in PC1 are most consistent with an effect of the number of

560 social partners. Stable pairs had significantly higher PC1 values compared to stable groups, and

561 there was a strong trend to be higher than socialized pairs (p=0.063). There were no differences

562 between stable groups and socialized pairs. Juveniles that experienced more social partners

563 during early-life were more active during the social cue investigation, spent less time in the

564 territory in the presence of a social cue, and were more socially interactive in a dominant social

565 role. The direction of this effect is also the same as in our previous study: group-reared juveniles

566 were more active and socially interactive than pair-reared juveniles (Solomon-Lane & Hofmann,

567 2019). This suggests that social experiences resulting from more novel partners may be an

568 important behavioural mechanism underlying the effect of group size. Syndromes involving

569 activity and social interaction are common across species (e.g., Conrad et al., 2011; Näslund &

570 Johnsson, 2016) and may also be related to bold-shy and proactive-reactive behaviours (Bell,

571 2007; Conrad et al., 2011; Groothuis & Carere, 2005; Koolhaas et al., 1999; Sih, Bell, Johnson,

572 et al., 2004).

It is not yet known how variable the number of social partners is for juvenile *A. burtoni* in

the wild or in larger, more naturalistic laboratory social groups. Evidence from a diversity of

other species suggests there can be considerable individual variation. In many species, siblings

(and parents) are the most proximate—and sometimes the only—early-life social contacts. The

number of offspring produced can vary both among individuals and within individuals across

reproductive bouts / over time. Within a group or population, individuals can also vary in the size

and makeup of their social network or niche (Barale et al., 2015; Beirão-Campos et al., 2016;

Branchi et al., 2013; Croft et al., 2005; Förster & Cords, 2005; Maguire et al., 2021; Monclús et

al., 2012; Pike et al., 2008; Saltz et al., 2016; Weinstein et al., 2014; Weinstein & Capitanio,

2008, 2008). For juveniles, this can have long-term, phenotypic effects (Branchi et al., 2013;

Monclús et al., 2012; Weinstein et al., 2014; Weinstein & Capitanio, 2008). Socialization

strategies are also used by humans with animals, such as working dogs (Gfrerer et al., 2018),

family dogs (Howell et al., 2015), and livestock like piglets (Morgan et al., 2014; Salazar et al.,

2018). Although many studies have manipulated the number of early-life social partners as a

consequence of group size, group size could exert independent or interacting effects on

phenotype. In studies that also controlled for group size, brown-headed cowbirds (*Molothrus*

ater) were housed in stable or dynamic flocks, in which flock members were exchanged multiple

times with novel birds. Dynamic flock males had more variable social networks over time, larger

signing networks, and outcompeted stable flock males in mating opportunities (White et al.,

2010). When housing conditions were later reversed, the new dynamic flock males still had

higher reproductive success, which was achieved via changes in social strategy (Gersick et al.,

2012). For juvenile *A. burtoni* in pairs and triads, individuals are not equally likely to initiate

interactions. Both group size and relative body size influenced social group structure (Solomon-

Lane et al., 2022). This suggests that, like adults (Maguire et al., 2021), social network position

and connectivity may vary considerably across individuals, with consequences for

developmental plasticity and behavioural development.

The treatment differences in PC5 (8.5%) support an effect of early-life social stability.

Socialized pairs had significantly higher PC5 values than stable groups and stable pairs, which

were not different from each other. David's score in the dominance behaviour assay loaded in the

same direction as time in the investigate zone during the social cue investigation. Submissions in

the dominance behaviour assay, frequency entering the territory zone in the social cue

investigation, time in and frequency entering the close zone in the social cue investigation, and

water-borne cortisol loaded together in the opposite direction. Socialized pairs were more

dominant in the dominance behaviour assay, spent less time in and near the territory zone, spent

more time in the investigate zone, and had lower cortisol levels than stable groups or pairs. This

suite of behaviours for PC5 shares multiple similarities with the PC1 syndrome, suggesting these

syndromes may not be independent of each other. Dominance behaviour is represented as high

rates of approaching and displacing on PC1 and as the opposing loadings of David's score

(indication of dominant status) and submissions (indication of subordinate status) on PC5.

Dominant juvenile *A. burtoni* approach and displace at significantly higher rates than

subordinates (Solomon-Lane et al., 2022). For PC5, being in or near the territory zone, with less

time in the investigate zone, was associated with low status when given the opportunity to be

dominant. This is mirrored on PC1 by the negative association between being in the territory

zone and approaching and displacing in the dominance behaviour assay. The relative dominance

of juveniles from the socialized pairs is also consistent with having significantly lower

directional consistency (i.e., more agonistically symmetrical) in the subordinate behaviour assay.

In our previous study, group-reared juveniles were less submissive in a subordinate role than

pair-reared juveniles (Solomon-Lane & Hofmann, 2019). It is possible this was not the case for

stable group juveniles in this study due to group size differences (6 fish vs. 16 fish).

The most striking difference for PC5 is the involvement of cortisol. We found no

treatment differences in water-borne cortisol levels, and PC5 was the only PC on which cortisol

loaded strongly. Higher cortisol was associated with lower social status, less exploratory

behaviour, and more time in the territory away from the novel cue fish. This suite of behaviours

with cortisol resembles previously identified syndromes (Réale et al., 2007), such as the pace-of-

life syndrome (Careau & Garland, 2012) and coping styles (Koolhaas et al., 1999). A "fast"

pace-of-life is associated with increased activity, exploration, boldness, and aggressiveness,

along with one or more traits from the slow-fast metabolic continuum (Careau & Garland, 2012).

Coping styles are an integrative phenotype in which a behavioural syndrome aligns with stress

physiology (Koolhaas et al., 1999). Coping styles have been observed across species (Alfonso et

al., 2019; Conrad et al., 2011; Øverli et al., 2007) and are sensitive to early-life effects (Sih,

2011). Behaviourally, proactive copers are more active, aggressive, and bold compared to

reactive copers. In response to stress, proactive copers have higher sympathetic reactivity and

lower HPA/I activity (Koolhaas et al., 1999). Whether socialized pairs (this study) or group-

reared juveniles (Solomon-Lane & Hofmann, 2019) exhibit a fast-pace-life or are proactive

copers are hypotheses that should be tested directly. The association between high cortisol and

low status is consistent with previous studies of juvenile *A. burtoni*, which showed higher whole

brain GR1a and GR1b expression, and lower GR2 and MR expression, in fish with higher

dominance scores (Solomon-Lane et al., 2022). Efficient negative feedback may be mediated, in

part, by GR1 expression, leading to lower cortisol levels (Solomon-Lane & Hofmann 2019). The

relationship between adult *A. burtoni* status and cortisol varies across studies and can be elevated

in subordinates (Maruska et al., 2022). Adult *A. burtoni* do not appear to form coping styles

(Butler et al., 2018), and this suite of behaviour is not always correlated with stress

responsiveness across species (e.g., Thomson et al., 2011). Overall, early-life exposure to social

complexity tends to benefit social competence and skills, whereas social instability tends to have

lasting, negative phenotypic effects, such as elevated HPA/I axis activity, weight loss, elevated

aggression, and decreased activity (Kohn, 2019). This suggests that exchanging members of the

pair in the socialized treatment may not have been perceived as instability, potentially because of

the predictable schedule (Kohn, 2019). An alternative explanation is that familiarity played a role

in the treatment differences between socialized pairs vs. stable groups and pairs. The dominant

and subordinate assays were highly similar to the way we socialized the pairs, and familiarity

with the assay could lead to appearing more active, interactive, and dominant. Future studies can

test these potential behavioural mechanisms directly.

We observed fish in their rearing environments to gain insights into the specific social

experiences and social sensory cues, or proximate behavioural mechanisms, responsible for

early-life social effects (Taborsky, 2016). Unsurprisingly rates of behaviour were higher in the

stable groups compared to the pairs. Rates of behaviour per fish did not differ across treatment

groups, and there were no differences in total agonistic efficiency. These data confirm that a

group (compared to pair) social environment has more opportunities for social experiences, such

as direct involvement in an interaction and observations of others interacting (Desjardins et al.,

2012; Oliveira et al., 1998). The mean distance between dyads was also significantly larger in

stable groups than the pairs, and although the tank was larger for groups, the smallest dyad

distance was significantly smaller in groups than social pairs. There were no differences across

treatments when scaled for the size of the tank. We were unable to identify and track individuals

because the ink tags were not visible on video. As a result, we could not quantify individual

social experience, social status, social network position, or spatial position, which we

hypothesize are causally related to behavioural development and phenotype (Kasumovic, 2013;

Taborsky, 2016). The evidence that social interactions are assortative within groups is

overwhelming across species (e.g., Chase et al., 2022; Croft et al., 2005, 2009; Pike et al., 2008;

Williamson et al., 2016), and juvenile *A. burtoni* form social status relationships and nuanced

social structures (Solomon-Lane et al., 2022). Therefore, we expect social status experience and

the degree of agonistic asymmetry across social partners to be particularly important for

behavioural development. In addition to its role as an indicator of social relationships, spatial

proximity may also affect social sensory cue perception and communication, for example, via

mechanosensory and chemosensory cues (e.g., in adult *A. burtoni*, Butler & Maruska, 2016;

Nikonov et al., 2017) that could be stronger for fish in closer proximity.

Overall, our work demonstrates that multiple behavioural mechanisms—the number of

early-life social partners and social stability or familiarity—affect juvenile *A. burtoni*

development and phenotype, including integrated behavioural and neuroendocrine traits. In

addition to manipulating social experience directly, we observed juveniles in their rearing

environments, which are necessary steps towards understanding the social experiences accrued

during development and the mechanisms by which early experiences exert long-term and

disproportionately strong effects on adult phenotype (Buist et al., 2013; Jonsson & Jonsson,

2014; Kasumovic, 2013; Taborsky, 2016, 2017). Although the simplistic social contexts we used

in this study are unlikely to reflect the dynamics of f larger, more complex groups found in

nature (Chase et al., 2003), we expect that these behavioural mechanisms, and potentially

others—acting additively or synergistically—will also be influential in more naturalistic

contexts. Testing the hypotheses we generated here will be key to uncovering the behavioural

mechanisms of developmental plasticity and phenotype, as well as the role of neuroendocrine

stress axis function, in *A. burtoni* and other social species.

ACKNOWLEDGEMENTS

This work was supported by start-up funds from the W.M. Keck Science Department to TKSL,

W.M. Keck Science Department Summer Fellowships to DDB, HLG, IPH, and EAM; Claremont

McKenna College Sponsored Internships & Experiences to JL, and the Claremont Colleges

Intercollegiate Neuroscience Summer Research Fellowship to JW. We thank Maya Shah for

contributing to scoring behaviour.

DECLARATION OF INTEREST: none.

DATA AVAILABILITY: Data will be made available in an online repository (Dryad).

REFERENCES

Ahloy Dallaire, J., & Mason, G. J. (2017). Juvenile rough-and-tumble play predicts adult sexual

behaviour in American mink. *Animal Behaviour*, *123*, 81–89.

https://doi.org/10.1016/j.anbehav.2016.10.023

Alcazar, R. M., Becker, L., Hilliard, A. T., Kent, K. R., & Fernald, R. D. (2016). Two types of

dominant male cichlid fish: Behavioural and hormonal characteristics. *Biology Open*,

5(8), 1061–1071. https://doi.org/10.1242/bio.017640

Alfonso, S., Sadoul, B., Gesto, M., Joassard, L., Chatain, B., Geffroy, B., & Bégout, M.-L.

(2019). Coping styles in European sea bass: The link between boldness, stress response

and neurogenesis. *Physiology & Behaviour*, *207*, 76–85.

https://doi.org/10.1016/j.physbeh.2019.04.020

Alward, B. A., Cathers, P. H., Blakkan, D. M., Hoadley, A. P., & Fernald, R. D. (2021). A

behavioural logic underlying aggression in an African cichlid fish. *Ethology*, *127*(7),

572–581. https://doi.org/10.1111/eth.13164

Antunes, D. F., Teles, M. C., Zuelling, M., Friesen, C. N., Oliveira, R. F., Aubin-Horth, N., &

Taborsky, B. (2021). Early social deprivation shapes neuronal programming of the social

decision-making network in a cooperatively breeding fish. *Molecular Ecology*, *30*(16),

4118–4132. https://doi.org/10.1111/mec.16019

Arnold, C., & Taborsky, B. (2010). Social experience in early ontogeny has lasting effects on

social skills in cooperatively breeding cichlids. *Animal Behaviour*, *79*(3), 621–630.

https://doi.org/10.1016/j.anbehav.2009.12.008

Banerjee, S. B., Arterbery, A. S., Fergus, D. J., & Adkins-Regan, E. (2012). Deprivation of

maternal care has long-lasting consequences for the hypothalamic–pituitary–adrenal axis

726  of zebra finches. *Proceedings of the Royal Society B: Biological Sciences*, *279*(1729),

727  759–766. https://doi.org/10.1098/rspb.2011.1265

728 Barale, C. L., Rubenstein, D. I., & Beehner, J. C. (2015). Juvenile social relationships reflect

729  adult patterns of behaviour in wild geladas. *American Journal of Primatology*, *77*(10),

730  1086–1096. https://doi.org/10.1002/ajp.22443

731 Bateson, P. (2001). Fetal experience and good adult design. *International Journal of*

732  *Epidemiology*, *30*(5), 928–934. https://doi.org/10.1093/ije/30.5.928

733 Bateson, P., Barker, D., Clutton-Brock, T., Deb, D., D'Udine, B., Foley, R. A., Gluckman, P.,

734  Godfrey, K., Kirkwood, T., Lahr, M. M., McNamara, J., Metcalfe, N. B., Monaghan, P.,

735  Spencer, H. G., & Sultan, S. E. (2004). Developmental plasticity and human health.

736  *Nature*, *430*(6998), 419–421. https://doi.org/10.1038/nature02725

737 Beirão-Campos, L., Cantor, M., Flach, L., & Simões-Lopes, P. C. (2016). Guiana dolphins form

738  social modules in a large population with high ranging overlap and small demographic

739  changes. *Behavioural Ecology and Sociobiology*, *70*(11), 1821–1830.

740  https://doi.org/10.1007/s00265-016-2188-x

741 Bell, A. M. (2007). Future directions in behavioural syndromes research. *Proceedings of the*

742  *Royal Society B: Biological Sciences*, *274*(1611), 755–761.

743  https://doi.org/10.1098/rspb.2006.0199

744 Bennett, D., Schneider, J., Tang, Y., Arnold, S., & Wilson, R. S. (2006). The effect of social

745  networks on the relation between Alzheimer's disease pathology and level of cognitive

746  function in old people: A longitudinal cohort study. *The Lancet Neurology*, *5*(5), 406–

747  412. https://doi.org/10.1016/S1474-4422(06)70417-3

Bölting, S., & von Engelhardt, N. (2017). Effects of the social environment during adolescence

on the development of social behaviour, hormones and morphology in male zebra finches

(Taeniopygia guttata). *Frontiers in Zoology*, *14*(1), 5. https://doi.org/10.1186/s12983-

017-0190-4

Bonuti, R., & Morato, S. (2018). Proximity as a predictor of social behaviour in rats. *Journal of*

Neuroscience Methods, *293*, 37–44. https://doi.org/10.1016/j.jneumeth.2017.08.027

Boogert, N. J., Farine, D. R., & Spencer, K. A. (2014). Developmental stress predicts social

network position. *Biology Letters*, *10*, 20140561. https://doi.org/10.1098/rsbl.2014.0561

Branchi, I., & Alleva, E. (2006). Communal nesting, an early social enrichment, increases the

adult anxiety-like response and shapes the role of social context in modulating the

emotional behaviour. *Behavioural Brain Research*, *172*(2), 299–306.

https://doi.org/10.1016/j.bbr.2006.05.019

Branchi, I., Curley, J. P., D'Andrea, I., Cirulli, F., Champagne, F. A., & Alleva, E. (2013). Early

interactions with mother and peers independently build adult social skills and shape

BDNF and oxytocin receptor brain levels. *Psychoneuroendocrinology*, *38*(4), 522–532.

https://doi.org/10.1016/j.psyneuen.2012.07.010

Branchi, I., D'Andrea, I., Gracci, F., Santucci, D., & Alleva, E. (2009). *Birth spacing in the*

mouse communal nest shapes adult emotional and social behaviour.

Bray, J., Feldblum, J. T., & Gilby, I. C. (2021). Social bonds predict dominance trajectories in

adult male chimpanzees. *Animal Behaviour*, *179*, 339–354.

https://doi.org/10.1016/j.anbehav.2021.06.031

Bray, J., Murray, C. M., Gilby, I. C., & Stanton, M. A. (2021). Immature Male Chimpanzees'

(Pan troglodytes schweinfurthii) Social Relationships with Adult Males, but Not Peers,

Persist into Adulthood. *International Journal of Primatology*, *42*(5), 701–721.

https://doi.org/10.1007/s10764-021-00235-1

Buist, K. L., Deković, M., & Prinzie, P. (2013). Sibling relationship quality and psychopathology

of children and adolescents: A meta-analysis. *Clinical Psychology Review*, *33*(1), 97–

106. https://doi.org/10.1016/j.cpr.2012.10.007

Butler, J. M., & Maruska, K. P. (2016). Mechanosensory signaling as a potential mode of

communication during social interactions in fishes. *Journal of Experimental Biology*,

219(18), 2781–2789. https://doi.org/10.1242/jeb.133801

Butler, J. M., Whitlow, S. M., Roberts, D. A., & Maruska, K. P. (2018). Neural and behavioural

correlates of repeated social defeat. *Scientific Reports*, *8*(1), 6818.

https://doi.org/10.1038/s41598-018-25160-x

Cachat, J., Stewart, A., Grossman, L., Gaikwad, S., Kadri, F., Chung, K. M., Wu, N., Wong, K.,

Roy, S., Suciu, C., Goodspeed, J., Elegante, M., Bartels, B., Elkhayat, S., Tien, D., Tan,

J., Denmark, A., Gilder, T., Kyzar, E., … Kalueff, A. V. (2010). Measuring behavioural

and endocrine responses to novelty stress in adult zebrafish. *Nature Protocols*, *5*(11),

1786–1799. https://doi.org/10.1038/nprot.2010.140

Careau, V., & Garland, T. (2012). Performance, Personality, and Energetics: Correlation,

Causation, and Mechanism. *Physiological and Biochemical Zoology*, *85*(6), 543–571.

https://doi.org/10.1086/666970

Carpenter, R. E., Maruska, K. P., Becker, L., & Fernald, R. D. (2014). Social Opportunity

Rapidly Regulates Expression of CRF and CRF Receptors in the Brain during Social

Ascent of a Teleost Fish, Astatotilapia burtoni. *PLoS ONE*, *9*(5), e96632.

https://doi.org/10.1371/journal.pone.0096632

Champagne, F. A., & Curley, J. P. (2005). How social experiences influence the brain. *Current*

Opinion in Neurobiology, *15*(6), 704–709. https://doi.org/10.1016/j.conb.2005.10.001

Chase, I. D., Coelho, D., Lee, W., Mueller, K., & Curley, J. P. (2022). Networks never rest: An

investigation of network evolution in three species of animals. *Social Networks*, *68*, 356–

373. https://doi.org/10.1016/j.socnet.2021.09.002

Chase, I. D., Tovey, C., & Murch, P. (2003). Two's Company, Three's a Crowd: Differences in

Dominance Relationships in Isolated versus Socially Embedded Pairs of Fish. *Behaviour*,

140(10,), 1193–1217.

Chen, C.-C., & Fernald, R. D. (2008). Sequences, expression patterns and regulation of the

corticotropin-releasing factor system in a teleost. *General and Comparative*

Endocrinology, *157*(2), 148–155. https://doi.org/10.1016/j.ygcen.2008.04.003

Clay, Z., & de Waal, F. B. M. (2013). Bonobos Respond to Distress in Others: Consolation

across the Age Spectrum. *PLoS ONE*, *8*(1), e55206.

https://doi.org/10.1371/journal.pone.0055206

Clement, T. S., Parikh, V., Schrumpf, M., & Fernald, R. D. (2005). Behavioural coping strategies

in a cichlid fish: The role of social status and acute stress response in direct and displaced

aggression. *Hormones and Behaviour*, *47*(3), 336–342.

https://doi.org/10.1016/j.yhbeh.2004.11.014

Conrad, J. L., Weinersmith, K. L., Brodin, T., Saltz, J. B., & Sih, A. (2011). Behavioural

syndromes in fishes: A review with implications for ecology and fisheries management.

Journal of Fish Biology, *78*(2), 395–435. https://doi.org/10.1111/j.1095-

8649.2010.02874.x

Crespi, E. J., & Denver, R. J. (2005). Ancient origins of human developmental plasticity.

American Journal of Human Biology, *17*(1), 44–54. https://doi.org/10.1002/ajhb.20098

Croft, D. P., James, R., Ward, A. J. W., Botham, M. S., Mawdsley, D., & Krause, J. (2005).

Assortative interactions and social networks in fish. *Oecologia*, *143*(2), 211–219.

https://doi.org/10.1007/s00442-004-1796-8

Croft, D. P., Krause, J., Darden, S. K., Ramnarine, I. W., Faria, J. J., & James, R. (2009).

Behavioural trait assortment in a social network: Patterns and implications. *Behavioural*

Ecology and Sociobiology, *63*(10), 1495–1503. https://doi.org/10.1007/s00265-009-0802-

x

Curley, J. P. (2016). *Compete: Analyzing competitive interaction data: R package version 0.1.*

D'Andrea, I., Alleva, E., & Branchi, I. (2007). Communal nesting, an early social enrichment,

affects social competences but not learning and memory abilities at adulthood.

Behavioural Brain Research, *183*(1), 60–66. https://doi.org/10.1016/j.bbr.2007.05.029

Denver, R. J. (2009). Stress hormones mediate environment-genotype interactions during

amphibian development. *General and Comparative Endocrinology*, *164*(1), 20–31.

https://doi.org/10.1016/j.ygcen.2009.04.016

Desjardins, J. K., Hofmann, H. A., & Fernald, R. D. (2012). Social Context Influences

Aggressive and Courtship Behaviour in a Cichlid Fish. *PLoS ONE*, *7*(7), e32781.

https://doi.org/10.1371/journal.pone.0032781

Dettmer, A. M., Wooddell, L. J., Rosenberg, K. L., Kaburu, S. S. K., Novak, M. A., Meyer, J. S.,

& Suomi, S. J. (2017). Associations between early life experience, chronic HPA axis

activity, and adult social rank in rhesus monkeys. *Social Neuroscience*, *12*(1), 92–101.

https://doi.org/10.1080/17470919.2016.1176952

Dijkstra, P. D., Maguire, S. M., Harris, R. M., Rodriguez, A. A., DeAngelis, R. S., Flores, S. A.,

& Hofmann, H. A. (2017). The melanocortin system regulates body pigmentation and

social behaviour in a colour polymorphic cichlid fish <sup/>. *Proceedings of the Royal*

Society B: Biological Sciences, *284*(1851), 20162838.

https://doi.org/10.1098/rspb.2016.2838

Ensminger, D. C., Langkilde, T., Owen, D. A. S., MacLeod, K. J., & Sheriff, M. J. (2018).

Maternal stress alters the phenotype of the mother, her eggs and her offspring in a wild-

caught lizard. *Journal of Animal Ecology*, *87*(6), 1685–1697.

https://doi.org/10.1111/1365-2656.12891

Farine, D. R., Spencer, K. A., & Boogert, N. J. (2015). Early-Life Stress Triggers Juvenile Zebra

Finches to Switch Social Learning Strategies. *Current Biology*, *25*(16), 2184–2188.

https://doi.org/10.1016/j.cub.2015.06.071

Fernald, R. D., & Hirata, N. R. (1977). Field study of Haplochromis burtoni: Habitats and co-

habitant. *Environmental Biology Of Fishes*, *2*(3), 299–308.

Fernald, R. D., & Hirata, N. R. (1979). The Ontogeny of Social behaviour and body coloration in

the african cichlid fish Haplochromis burtoni. *Zeitschrif Für Tierpsychologie*, *50*, 180–

187.

Fernald, R. D., & Maruska, K. P. (2012). Social information changes the brain. *Proceedings of*

the National Academy of Sciences, *109*(supplement_2), 17194–17199.

https://doi.org/10.1073/pnas.1202552109

Fischer, S., Bessert-Nettelbeck, M., Kotrschal, A., & Taborsky, B. (2015). Rearing-Group Size

Determines Social Competence and Brain Structure in a Cooperatively Breeding Cichlid.

The American Naturalist, *186*(1). https://doi.org/10.1086/681636

Förster, S., & Cords, M. (2005). Socialization of infant blue monkeys (Cercopithecus mitis

stuhlmanni): Allomaternal interactions and sex differences. *Behaviour*, *142*(7), 869–896.

https://doi.org/10.1163/1568539055010138

Fox, H. E., White, S. A., Kao, M. H. F., & Fernald, R. D. (1997). Stress and Dominance in a

Social Fish. *The Journal of Neuroscience*, *17*(16), 6463–6469.

https://doi.org/10.1523/JNEUROSCI.17-16-06463.1997

Fraley, N. B., & Fernald, R. D. (1982). Social control of developmental rate in the African

cichlid, Haplochromis burtoni. *Zeitschrif Für Tierpsychologie*, *60*(1), 66–82.

Francis, D. D., Caldji, C., Champagne, F., Plotsky, P. M., & Meaney, M. J. (1999). The role of

corticotropin-releasing factor–norepinephrine systems in mediating the effects of early

experience on the development of behavioural and endocrine responses to stress.

Biological Psychiatry, *46*(9), 1153–1166. https://doi.org/10.1016/S0006-3223(99)00237-

1

Freeman, N. E., Norris, D. R., Sutton, A. O., Strickland, D., Kyser, T. K., & Newman, A. E. M.

(2021). Early-Life Corticosterone Body Condition Influence Social Status and Survival in

a Food-Caching Passerine. *Integrative and Comparative Biology*, *61*(1), 9–19.

https://doi.org/10.1093/icb/icab038

Friesen, C. N., Maclaine, K. D., & Hofmann, H. A. (2022). Social status mediates behavioural,

endocrine, and neural responses to an intruder challenge in a social cichlid, Astatotilapia

burtoni. *Hormones and Behaviour*, *145*, 105241.

https://doi.org/10.1016/j.yhbeh.2022.105241

Fulmer, A. G., Neumeister, H., & Preuss, T. (2017). Social plasticity in non-territorial male

African cichlid fish Astatotilapia burtoni. *Journal of Ethology*, *35*(1), 109–119.

https://doi.org/10.1007/s10164-016-0498-0

Gallego-Abenza, M., Boucherie, P. H., & Bugnyar, T. (2022). Early social environment affects

attention to social cues in juvenile common ravens, Corvus corax. *Royal Society Open*

Science, *9*(6), 220132. https://doi.org/10.1098/rsos.220132

Gammell, M. P., de Vries, H., Jennings, D. J., Carlin, C. M., & Hayden, T. J. (2003). David's

score: A more appropriate dominance ranking method than Clutton-Brock et al.'s index.

Animal Behaviour, *66*(3), 601–605. https://doi.org/10.1006/anbe.2003.2226

Gersick, A. S., Snyder-Mackler, N., & White, D. J. (2012). Ontogeny of social skills: Social

complexity improves mating and competitive strategies in male brown-headed cowbirds.

Animal Behaviour, *83*(5), 1171–1177. https://doi.org/10.1016/j.anbehav.2012.02.005

Gfrerer, N., Taborsky, M., & Würbel, H. (2018). Benefits of intraspecific social exposure in

adult Swiss military dogs. *Applied Animal Behaviour Science*, *201*, 54–60.

https://doi.org/10.1016/j.applanim.2017.12.016

Greenwood, A. K., Butler, P. C., White, R. B., DeMarco, U., Pearce, D., & Fernald, R. D.

(2003). Multiple Corticosteroid Receptors in a Teleost Fish: Distinct Sequences,

Expression Patterns, and Transcriptional Activities. *Endocrinology*, *144*(10), 4226–4236.

https://doi.org/10.1210/en.2003-0566

Groothuis, T. G. G., & Carere, C. (2005). Avian personalities: Characterization and epigenesis.

Neuroscience & Biobehavioural Reviews, *29*(1), 137–150.

https://doi.org/10.1016/j.neubiorev.2004.06.010

Heule, C., Göppert, C., Salzburger, W., & Böhne, A. (2014). Genetics and timing of sex

determination in the East African cichlid fish Astatotilapia burtoni. *BMC Genetics*, *15*,

140. https://doi.org/10.1186/s12863-014-0140-5

Hofmann, H. A. (2003). Functional genomics of neural and behavioural plasticity. *Journal of*

Neurobiology, *54*(1), 272–282. https://doi.org/10.1002/neu.10172

Hofmann, H. A., Benson, M. E., & Fernald, R. D. (1999). Social status regulates growth rate:

Consequences for life-history strategies. *Proceedings of the National Academy of*

Sciences of the United States of America, *96*(24), 14171–14176.

Howell, T., King, T., & Bennett, P. (2015). Puppy parties and beyond: The role of early age

socialization practices on adult dog behaviour. *Veterinary Medicine: Research and*

Reports, 143. https://doi.org/10.2147/VMRR.S62081

Huffman, L. S., Hinz, F. I., Wojcik, S., Aubin-Horth, N., & Hofmann, H. A. (2015). Arginine

vasotocin regulates social ascent in the African cichlid fish Astatotilapia burtoni. *General*

and Comparative Endocrinology, *212*, 106–113.

https://doi.org/10.1016/j.ygcen.2014.03.004

Jin, T., Wang, D., Pan, W., & Yao, M. (2015). Nonmaternal Infant Handling in Wild White-

Headed Langurs (Trachypithecus leucocephalus). *International Journal of Primatology*,

36(2), 269–287. https://doi.org/10.1007/s10764-015-9823-7

Jonsson, B., & Jonsson, N. (2014). Early environment influences later performance in fishes:

Effects of early experiences. *Journal of Fish Biology*, *85*(2), 151–188.

https://doi.org/10.1111/jfb.12432

Kasumovic, M. M. (2013). The multidimensional consequences of the juvenile environment:

Towards an integrative view of the adult phenotype. *Animal Behaviour*, *85*(5), 1049–

1059. https://doi.org/10.1016/j.anbehav.2013.02.009

Kidd, C. E., Kidd, M. R., & Hofmann, H. A. (2010). Measuring multiple hormones from a single

water sample using enzyme immunoassays. *General and Comparative Endocrinology*,

165(2), 277–285. https://doi.org/10.1016/j.ygcen.2009.07.008

Kidd, M. R., O'Connell, L. A., Kidd, C. E., Chen, C. W., Fontenot, M. R., Williams, S. J., &

Hofmann, H. A. (2013). Female preference for males depends on reproductive

physiology in the African cichlid fish Astatotilapia burtoni. *General and Comparative*

Endocrinology, *180*, 56–63. https://doi.org/10.1016/j.ygcen.2012.10.014

Kohn, G. M. (2019). How social systems persist: Learning to build a social network in an

uncertain world. *Animal Behaviour*, *154*, 1–6.

https://doi.org/10.1016/j.anbehav.2019.06.006

Koolhaas, J. M., Korte, S. M., De Boer, S. F., Van Der Vegt, B. J., Van Reenen, C. G., Hopster,

H., De Jong, I. C., Ruis, M. A. W., & Blokhuis, H. J. (1999). Coping styles in animals:

Current status in behaviour and stress-physiology. *Neuroscience & Biobehavioural*

Reviews, *23*(7), 925–935. https://doi.org/10.1016/S0149-7634(99)00026-3

Korzan, W. J., Fernald, R. D., & Grone, B. P. (2014). Social regulation of cortisol receptor gene

expression. *Journal of Experimental Biology*, jeb.104430.

https://doi.org/10.1242/jeb.104430

Kuijper, B., & Johnstone, R. A. (2019). The evolution of early-life effects on social behaviour—

Why should social adversity carry over to the future? *Philosophical Transactions of the*

Royal Society B: Biological Sciences, *374*, 20180111.

https://doi.org/10.1098/rstb.2018.0111

Leiva, D., Solanas, A., & Salafranca, L. (2008). Testing reciprocity in social interactions: A

comparison between the directional consistency and skew-symmetry statistics. *Behaviour*

Research Methods, *40*(2), 626–634. https://doi.org/10.3758/BRM.40.2.626

Lotrich, V. A., & Meredith, W. H. (1974). A Technique and the Effectiveness of Various Acrylic

Colors for Subcutaneous Marking of Fish. *Transactions of the American Fisheries*

Society, *1*, 3.

Lowry, C. A., & Moore, F. L. (2006). Regulation of behavioural responses by corticotropin-

releasing factor. *General and Comparative Endocrinology*, *146*(1), 19–27.

https://doi.org/10.1016/j.ygcen.2005.12.006

Maguire, S. M., DeAngelis, R., Dijkstra, P. D., Jordan, A., & Hofmann, H. A. (2021). Social

network dynamics predict hormone levels and behaviour in a highly social cichlid fish.

Hormones and Behaviour, *132*, 104994. https://doi.org/10.1016/j.yhbeh.2021.104994

Maruska, K. P., Anselmo, C. M., King, T., Mobley, R. B., Ray, E. J., & Wayne, R. (2022).

Endocrine and neuroendocrine regulation of social status in cichlid fishes. *Hormones and*

Behaviour, *139*, 105110. https://doi.org/10.1016/j.yhbeh.2022.105110

Maruska, K. P., Becker, L., Neboori, A., & Fernald, R. D. (2013). Social descent with territory

loss causes rapid behavioural, endocrine, and transcriptional changes in the brain. *Journal*

of Experimental Biology, jeb.088617. https://doi.org/10.1242/jeb.088617

Maruska, K. P., Zhang, A., Neboori, A., & Fernald, R. D. (2013). Social Opportunity Causes

Rapid Transcriptional Changes in the Social Behaviour Network of the Brain in an

African Cichlid Fish. *Journal of Neuroendocrinology*, *25*(2), 145–157.

https://doi.org/10.1111/j.1365-2826.2012.02382.x

McClelland, S., Korosi, A., Cope, J., Ivy, A., & Baram, T. Z. (2011). Emerging roles of

epigenetic mechanisms in the enduring effects of early-life stress and experience on

learning and memory. *Neurobiology of Learning and Memory*, *96*(1), 79–88.

https://doi.org/10.1016/j.nlm.2011.02.008

McDonald, D. B. (2007). Predicting fate from early connectivity in a social network.

Proceedings of the National Academy of Sciences, *104*(26), 10910–10914.

https://doi.org/10.1073/pnas.0701159104

Meyer-Lindenberg, A., & Tost, H. (2012). Neural mechanisms of social risk for psychiatric

disorders. *Nature Neuroscience*, *15*(5), 663–668. https://doi.org/10.1038/nn.3083

Monclús, R., Cook, T., & Blumstein, D. T. (2012). Masculinized female yellow-bellied marmots

initiate more social interactions. *Biology Letters*, *8*(2), 208–210.

https://doi.org/10.1098/rsbl.2011.0754

Moretz, J. A., Martins, E. P., & Robison, B. D. (2007). The effects of early and adult social

environment on zebrafish (Danio rerio) behaviour. *Environmental Biology of Fishes*,

80(1), 91–101. https://doi.org/10.1007/s10641-006-9122-4

Morgan, T., Pluske, J., Miller, D., Collins, T., Barnes, A. L., Wemelsfelder, F., & Fleming, P. A.

(2014). Socialising piglets in lactation positively affects their post-weaning behaviour.

Applied Animal Behaviour Science, *158*, 23–33.

https://doi.org/10.1016/j.applanim.2014.06.001

Moy, S. S., Nadler, J. J., Perez, A., Barbaro, R. P., Johns, J. M., Magnuson, T. R., Piven, J., &

Crawley, J. N. (2004). Sociability and preference for social novelty in five inbred strains:

An approach to assess autistic-like behaviour in mice. *Genes, Brain and Behaviour*, *3*(5),

287–302. https://doi.org/10.1111/j.1601-1848.2004.00076.x

Näslund, J., & Johnsson, J. I. (2016). State-dependent behaviour and alternative behavioural

strategies in brown trout (Salmo trutta L.) fry. *Behavioural Ecology and Sociobiology*,

70(12), 2111–2125. https://doi.org/10.1007/s00265-016-2215-y

Nikonov, A. A., Butler, J. M., Field, K. E., Caprio, J., & Maruska, K. P. (2017). Reproductive

and metabolic state differences in olfactory responses to amino acids in a mouth brooding

African cichlid fish. *Journal of Experimental Biology*, jeb.157925.

https://doi.org/10.1242/jeb.157925

Nyman, C., Fischer, S., Aubin-Horth, N., & Taborsky, B. (2018). Evolutionary conserved neural

signature of early life stress affects animal social competence. *Proceedings of the Royal*

Society B: Biological Sciences, *285*(1871), 20172344.

https://doi.org/10.1098/rspb.2017.2344

Oliveira, R. F., McGregor, P. K., & Latru, C. (1998). *Know thine enemy: Fighting fish gather*

information from observing conspecific interactions.

Øverli, Ø., Sørensen, C., Pulman, K. G. T., Pottinger, T. G., Korzan, W., Summers, C. H., &

Nilsson, G. E. (2007). Evolutionary background for stress-coping styles: Relationships

between physiological, behavioural, and cognitive traits in non-mammalian vertebrates.

Neuroscience & Biobehavioural Reviews, *31*(3), 396–412.

https://doi.org/10.1016/j.neubiorev.2006.10.006

Parikh, V. N., Clement, T., & Fernald, R. D. (2006). Physiological consequences of social

descent: Studies in Astatotilapia burtoni. *Journal of Endocrinology*, *190*(1), 183–190.

https://doi.org/10.1677/joe.1.06755

Perkeybile, A. M., & Bales, K. L. (2017). Intergenerational transmission of sociality: The role of

parents in shaping social behaviour in monogamous and non-monogamous species.

Journal of Experimental Biology, *220*(1), 114–123. https://doi.org/10.1242/jeb.142182

Perkeybile, A. M., Griffin, L. L., & Bales, K. L. (2013). Natural variation in early parental care

correlates with social behaviours in adolescent prairie voles (Microtus ochrogaster).

Frontiers in Behavioural Neuroscience, *7*. https://doi.org/10.3389/fnbeh.2013.00021

Pike, T. W., Samanta, M., Lindström, J., & Royle, N. J. (2008). Behavioural phenotype affects

social interactions in an animal network. *Proceedings of the Royal Society B: Biological*

Sciences, *275*(1650), 2515–2520. https://doi.org/10.1098/rspb.2008.0744

Prut, L., & Belzung, C. (2003). The open field as a paradigm to measure the effects of drugs on

anxiety-like behaviours: A review. *European Journal of Pharmacology*, *463*(1–3), 3–33.

https://doi.org/10.1016/S0014-2999(03)01272-X

Pryce, C. R., Aubert, Y., Maier, C., Pearce, P. C., & Fuchs, E. (2011). The developmental impact

of prenatal stress, prenatal dexamethasone and postnatal social stress on physiology,

behaviour and neuroanatomy of primate offspring: Studies in rhesus macaque and

common marmoset. *Psychopharmacology*, *214*(1), 33–53.

https://doi.org/10.1007/s00213-010-1989-2

Réale, D., Reader, S. M., Sol, D., McDougall, P. T., & Dingemanse, N. J. (2007). Integrating

animal temperament within ecology and evolution. *Biological Reviews*, *82*(2), 291–318.

https://doi.org/10.1111/j.1469-185X.2007.00010.x

Renn, S. C. P., Carleton, J. B., Magee, H., Nguyen, M. L. T., & Tanner, A. C. W. (2009).

Maternal care and altered social phenotype in a recently collected stock of Astatotilapia

1038    burtoni cichlid fish. *Integrative and Comparative Biology*, *49*(6), 660–673.

1039    https://doi.org/10.1093/icb/icp085

1040 Reyes-Contreras, M., Glauser, G., Rennison, D. J., & Taborsky, B. (2019). Early-life

1041    manipulation of cortisol and its receptor alters stress axis programming and social

1042    competence. *Philosophical Transactions of the Royal Society B: Biological Sciences*,

1043    *374*(1770), 20180119. https://doi.org/10.1098/rstb.2018.0119

1044 Rodriguez-Santiago, M., Nührenberg, P., Derry, J., Deussen, O., Francisco, F. A., Garrison, L.

1045    K., Garza, S. F., Hofmann, H. A., & Jordan, A. (2020). Behavioural traits that define

1046    social dominance are the same that reduce social influence in a consensus task.

1047    *Proceedings of the National Academy of Sciences*, *117*(31), 18566–18573.

1048    https://doi.org/10.1073/pnas.2000158117

1049 Salazar, L. C., Ko, H.-L., Yang, C.-H., Llonch, L., Manteca, X., Camerlink, I., & Llonch, P.

1050    (2018). Early socialisation as a strategy to increase piglets' social skills in intensive

1051    farming conditions. *Applied Animal Behaviour Science*, *206*, 25–31.

1052    https://doi.org/10.1016/j.applanim.2018.05.033

1053 Saltz, J. B., Geiger, A. P., Anderson, R., Johnson, B., & Marren, R. (2016). What, if anything, is

1054    a social niche? *Evolutionary Ecology*, *30*(2), 349–364. https://doi.org/10.1007/s10682-

1055    015-9792-5

1056 Schneider, C. A., Rasband, W. S., & Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years of

1057    image analysis. *Nature Methods*, *9*(7), 671–675. https://doi.org/10.1038/nmeth.2089

1058 Sih, A. (2011). Effects of early stress on behavioural syndromes: An integrated adaptive

1059    perspective. *Neuroscience & Biobehavioural Reviews*, *35*(7), 1452–1465.

1060    https://doi.org/10.1016/j.neubiorev.2011.03.015

Sih, A., Bell, A., & Johnson, J. C. (2004). Behavioural syndromes: An ecological and

evolutionary overview. *Trends in Ecology & Evolution*, *19*(7), 372–378.

https://doi.org/10.1016/j.tree.2004.04.009

Sih, A., Bell, A. M., Johnson, J. C., & Ziemba, R. E. (2004). Behavioural Syndromes: An

Integrative Overview. *The Quarterly Review of Biology*, *79*(3), 241–277.

https://doi.org/10.1086/422893

Sih, A., Mathot, K. J., Moirón, M., Montiglio, P.-O., Wolf, M., & Dingemanse, N. J. (2015).

Animal personality and state–behaviour feedbacks: A review and guide for empiricists.

Trends in Ecology & Evolution, *30*(1), 50–60. https://doi.org/10.1016/j.tree.2014.11.004

Silk, J. B. (2007). Social Components of Fitness in Primate Groups. *Science*, *317*(5843), 1347–

1351. https://doi.org/10.1126/science.1140734

Solomon-Lane, T. K., Butler, R. M., & Hofmann, H. A. (2022). Vasopressin mediates

nonapeptide and glucocorticoid signaling and social dynamics in juvenile dominance

hierarchies of a highly social cichlid fish. *Hormones and Behaviour*, *145*, 105238.

https://doi.org/10.1016/j.yhbeh.2022.105238

Solomon-Lane, T. K., & Hofmann, H. A. (2018). *A tagging method for very small fish* [Preprint].

https://doi.org/10.1101/437772

Solomon-Lane, T. K., & Hofmann, H. A. (2019). Early-life social environment alters juvenile

behaviour and neuroendocrine function in a highly social cichlid fish. *Hormones and*

Behaviour, *115*, 104552. https://doi.org/10.1016/j.yhbeh.2019.06.016

Solomon-Lane, T. K., Pradhan, D. S., Willis, M. C., & Grober, M. S. (2015). Agonistic

reciprocity is associated with reduced male reproductive success within haremic social

1083    networks. *Proceedings of the Royal Society B: Biological Sciences*, *282*(1811),

1084    20150914. https://doi.org/10.1098/rspb.2015.0914

1085 Spencer, K. A. (2017). Developmental stress and social phenotypes: Integrating neuroendocrine,

1086    behavioural and evolutionary perspectives. *Philosophical Transactions of the Royal*

1087    *Society B: Biological Sciences*, *372*(1727), 20160242.

1088    https://doi.org/10.1098/rstb.2016.0242

1089 Stevens, H. E., Leckman, J. F., Coplan, J. D., & Suomi, S. J. (2009). Risk and Resilience: Early

1090    Manipulation of Macaque Social Experience and Persistent Behavioural and

1091    Neurophysiological Outcomes. *Journal of the American Academy of Child & Adolescent*

1092    *Psychiatry*, *48*(2), 114–127. https://doi.org/10.1097/CHI.0b013e318193064c

1093 Stevenson, R. D., & Woods, W. A. (2006). Condition indices for conservation: New uses for

1094    evolving tools. *Integrative and Comparative Biology*, *46*(6), 1169–1190.

1095    https://doi.org/10.1093/icb/icl052

1096 Stevenson, T. J., Alward, B. A., Ebling, F. J. P., Fernald, R. D., Kelly, A., & Ophir, A. G.

1097    (2018). The Value of Comparative Animal Research: Krogh's Principle Facilitates

1098    Scientific Discoveries. *Policy Insights from the Behavioural and Brain Sciences*, *5*(1),

1099    118–125. https://doi.org/10.1177/2372732217745097

1100 Taborsky, B. (2016). Opening the Black Box of Developmental Experiments: Behavioural

1101    Mechanisms Underlying Long-Term Effects of Early Social Experience. *Ethology*,

1102    *122*(4), 267–283. https://doi.org/10.1111/eth.12473

1103 Taborsky, B. (2017). Developmental Plasticity: Preparing for Life in a Complex World. In

1104    *Advances in the Study of Behaviour* (Vol. 49, pp. 49–99). Elsevier.

1105    https://doi.org/10.1016/bs.asb.2016.12.002

Taborsky, B., Arnold, C., Junker, J., & Tschopp, A. (2012). The early social environment affects

social competence in a cooperative breeder. *Animal Behaviour*, *83*(4), 1067–1074.

https://doi.org/10.1016/j.anbehav.2012.01.037

Taborsky, B., Tschirren, L., Meunier, C., & Aubin-Horth, N. (2013). Stable reprogramming of

brain transcription profiles by the early social environment in a cooperatively breeding

fish. *Proceedings of the Royal Society B: Biological Sciences*, *280*(1753), 20122605.

https://doi.org/10.1098/rspb.2012.2605

Thomson, J. S., Watts, P. C., Pottinger, T. G., & Sneddon, L. U. (2011). Physiological and

genetic correlates of boldness: Characterising the mechanisms of behavioural variation in

rainbow trout, Oncorhynchus mykiss. *Hormones and Behaviour*, *59*(1), 67–74.

https://doi.org/10.1016/j.yhbeh.2010.10.010

Thresher, R. E., & Gronell, A. M. (1978). Subcutaneous Tagging of Small Reef Fishes. *Copeia*,

1978(2), 352. https://doi.org/10.2307/1443582

Turecki, G., & Meaney, M. J. (2016). Effects of the Social Environment and Stress on

Glucocorticoid Receptor Gene Methylation: A Systematic Review. *Biological Psychiatry*,

79(2), 87–96. https://doi.org/10.1016/j.biopsych.2014.11.022

Wallace, K. J., & Hofmann, H. A. (2021). Equal performance but distinct behaviours: Sex

differences in a novel object recognition task and spatial maze in a highly social cichlid

fish. *Animal Cognition*, *24*(5), 1057–1073. https://doi.org/10.1007/s10071-021-01498-0

Weaver, I. C. G. (2009). Shaping adult phenotypes through early life environments. *Birth*

Defects Research Part C: Embryo Today: Reviews, *87*(4), 314–326.

https://doi.org/10.1002/bdrc.20164

Weinstein, T. A. R., Bales, K. L., Maninger, N., Hostetler, C. M., & Capitanio, J. P. (2014).

Early involvement in friendships predicts later plasma concentrations of oxytocin and

vasopressin in juvenile rhesus macaques (Macaca mulatta). *Frontiers in Behavioural*

Neuroscience, *8*. https://doi.org/10.3389/fnbeh.2014.00295

Weinstein, T. A. R., & Capitanio, J. P. (2008). Individual differences in infant temperament

predict social relationships of yearling rhesus monkeys, Macaca mulatta. *Animal*

Behaviour, *76*(2), 455–465. https://doi.org/10.1016/j.anbehav.2008.01.024

Weitekamp, C. A., & Hofmann, H. A. (2017). Neuromolecular correlates of cooperation and

conflict during territory defense in a cichlid fish. *Hormones and Behaviour*, *89*, 145–156.

https://doi.org/10.1016/j.yhbeh.2017.01.001

Weitekamp, C. A., Nguyen, J., & Hofmann, H. A. (2017). Social context affects behaviour,

preoptic area gene expression, and response to D2 receptor manipulation during territorial

defense in a cichlid fish. *Genes, Brain and Behaviour*, *16*(6), 601–611.

https://doi.org/10.1111/gbb.12389

Wendelaar Bonga, S. E. (1997). The stress response in fish. *Physiological Reviews*, *77*(3), 591–

625. https://doi.org/10.1152/physrev.1997.77.3.591

White, D. J., Gersick, A. S., Freed-Brown, G., & Snyder-Mackler, N. (2010). The ontogeny of

social skills: Experimental increases in social complexity enhance reproductive success in

adult cowbirds. *Animal Behaviour*, *79*(2), 385–390.

https://doi.org/10.1016/j.anbehav.2009.11.014

Williamson, C. M., Lee, W., & Curley, J. P. (2016). Temporal dynamics of social hierarchy

formation and maintenance in male mice. *Animal Behaviour*, *115*, 259–272.

https://doi.org/10.1016/j.anbehav.2016.03.004

Wilson, E. O. (1980). *Sociobiology: The Abridged Edition*. The Belknap Press of Harvard

University Press.

1153

1154

1155   **TABLES**

1156   **Table 1:** Treatment differences in social behaviour and distances between dyads in the stable

1157   group, socialized pair, and stable pair home tanks.

Behaviour / measure	DF	Test Statistic	p-value	Effect size	*Post hoc* / direction of effect
Approaches	2, 38	F=11.46	0.00013	0.38	**Stable group > social. pair: p<0.0001** **Stable group > stable pair: p=0.0061** Social. pair vs. stable pair: p=0.41
Displacements	2, 37	F=11.37	0.00014	0.38	**Stable group > social. pair: p<0.0001** **Stable group > stable pair: p=0.02** Social. pair vs. stable pair: p=0.21
Efficiency	2	χ^2=1.06	0.59		
Into territory	2	χ^2=10.81	0.0045	0.23	**Stable group > stable pair: p=0.0037** Stable group > social. pair: p=0.071 Social. pair vs. stable pair: p=0.75
Mean dyad distance	2, 46	F=13.62	<0.0001	0.37	**Stable group > stable pair: p<0.0001** **Stable group > social. pair: p<0.0001** Social. pair vs. stable pair: p=0.92
Minimum dyad distance	2	χ^2=7.49	0.0025	0.12	**Stable group > social. pair: p=0.021** Stable group > stable pair: p=0.085 Social. pair vs. stable pair: p=1.0

Results of one-way ANOVAs (F) or nonparametric Kruskal-Wallis tests (χ^2). Eta-squared is

reported for effect sizes. Significant results in bold. Tukey HSD tests were used for *post hoc*

analysis of significant ANOVA results. Dunn's tests were used for *post hoc* analysis of

significant Kruskal-Wallis results.

Table 2: Treatment differences in the frequency entering and time spent in tank zones during the

open field exploration and social cue investigation.

Tank Zones / measure	DF	Test Statistic	p-value	Effect size	*Post hoc* / direction of effect
Open Field Exploration					
Territory (Frequency)	**2,30**	**F=3.70**	**0.037**	**0.2**	**Social. pair > stable pair: p=0.037** Social. pair vs. stable group: p=0.1 Stable group vs. stable pair: p=0.77
Close (Frequency)	2,30	F=3.30	0.051		
Far (Frequency)	2	χ²=4.97	0.083		
Investigate (Frequency)	2	χ²=3.51	0.17		
Territory (Time)	2	χ²=4.34	0.11		
Close (Time)	2	χ²=0.34	0.84		
Far (Time)	**2**	**χ²=7.47**	**0.024**	**0.18**	**Stable group > stable pair: p=0.032** Stable group vs. social. pair: 0.16 Social. pair vs. stable pair: p=1.0
Investigate (Time)	2	χ²=2.13	0.35		
Social Cue Investigation					
Territory (Frequency)	2	χ²=0.76	0.69		
Close (Frequency)	2	χ²=1.51	0.47		
Far (Frequency)	**2**	**χ²=6.60**	**0.037**	**0.15**	**Stable group > stable pair: p=0.036** Stable group vs. social. pair: 0.40 Social. pair vs. stable pair: p=1.0
Investigate (Frequency)	2	χ²=4.49	0.11		
Territory (Time)	**2**	**χ²=6.36**	**0.042**	**0.15**	**Stable pair > stable group: p=0.042** Stable group vs. social. pair: 0.39 Social. pair vs. stable pair: p=1.0
Close (Time)	2	χ²=2.19	0.33		
Far (Time)	**2**	**χ²=7.93**	**0.019**	**0.20**	**Stable group > stable pair: p=0.019** Stable group vs. social. pair: 0.25 Social. pair vs. stable pair: p=1.0
Investigate (Time)	2	χ²=5.00	0.082		

Results of one-way ANOVAs (F) or nonparametric Kruskal-Wallis tests (χ^2). Eta-squared is

reported for effect sizes. Significant results in bold. Tukey HSD tests were used for *post hoc*

analysis of significant ANOVA results. Dunn's tests were used for *post hoc* analysis of

significant Kruskal-Wallis results.

Table 3: Treatment differences in social behaviour and status in the dominant and subordinate

behaviour assays.

Assay / Behaviour	DF	χ^2	p-value	Effect size	*Post hoc* / direction of effect
Dominant Behaviour Assay					
Approaches	**2**	**8.71**	**0.013**	**0.22**	**Social. pair > stable pair: p=0.01** Social. pair vs. stable group: p=0.16 Stable group vs. stable pair: p=0.57
Displacements	**2**	**7.62**	**0.022**	**0.19**	**Social. pair > stable pair: p=0.02** Social. pair vs. stable group: p=0.16 Stable group vs. stable pair: p=0.85
Approaches Received	2	5.44	0.066		
Submissions	2	4.63	0.099		
David's Score	**2**	**8.48**	**0.014**	**0.22**	**Social. pair > stable pair: p=0.043** **Social. pair > stable group: p=0.022** Stable group vs. stable pair: p=1.0
Directional consistency (App.)	2	3.89	0.14		
Subordinate Behaviour Assay					
Approaches	2	5.23	0.073		
Displacements	2	2.69	0.26		
Approaches Received	**2**	**6.13**	**0.047**	**0.14**	**Social. pair > stable pair: p=0.042** Social. pair vs. stable group: p=0.73 Stable group vs. stable pair: p=0.38
Submissions	2	4.05	0.13		
Focal efficiency	2	1.38	0.50		
Directional consistency (App.)	**2**	**13.22**	**0.0014**	**0.37**	**Social. pair < stable pair: p=0.0038** **Social. pair < stable group: p=0.004** Stable group vs. stable pair: p=1.0

1170   Results of Kruskal-Wallis tests (χ^2). Eta-squared is reported for effect sizes. Significant results in

1171   bold. Dunn's tests were used for *post hoc* analysis of significant results.

1172   **Table 4:** Treatment differences for principal components (PC) 1-5.

Principal component (% variation)	DF	Test Statistic	p-value	Effect size	*Post hoc* / direction of effect
PC1 (25.1%)	2,28	**F=4.19**	**0.026**	**0.23**	**Stable pair > stable group: p=0.032** Stable pair > social. pair: p=0.063 Stable group vs. social. pair: p=1.0
PC2 (18.2%)	2,28	F=1.82	0.18		
PC3 (14.3%)	2,28	F=0.97	0.39		
PC4 (10.5%)	2	χ^2=2.69	0.26		
PC5 (8.5%)	2,28	**F=6.62**	**0.0044**	**0.32**	**Social. pair > stable group: p=0.0036** **Social. pair > stable pair: p=0.036** Stable group vs. stable pair: p=0.75

1173   Results of one-way ANOVAs (F) or nonparametric Kruskal-Wallis tests (χ^2) for PC1-PC5.

1174   Percentages indicate the variance explained. Eta-squared is reported for effect sizes. Significant

1175   results in bold. Tukey HSD tests were used for *post hoc* analysis of significant ANOVA results.

1176   Dunn's tests were used for *post hoc* analysis of significant Kruskal-Wallis results.

 FIGURES

Figure 1: A) Juvenile fish were reared in stable groups (n=6, 6 fish each), stable pairs (n=6, 2

fish each), or socialized pairs (n=9, 2 fish each). Every 5 days in the socialized pairs, one fish of

the pair was removed, and a novel juvenile was introduced. The novel juvenile came from a

different socialized pair, and the removed juvenile became the novel partner for a socialized pair.

This was repeated 5 times so that socialized fish had a total of 5 social partners – equal to the

stable group fish. One fish per stable group and pair was also removed from their tank with a

hand net as a control. This fish was immediately returned to its home tank. B) After 26 days in

these rearing environments, individual juvenile behaviour was quantified in a novel experimental

tank. The tank contained a terracotta pot shard, and black lines (drawn in permanent marker)

divided the tank into four zones: territory, close, far, and investigate. In the open field

exploration, the focal fish was alone in the tank. In the social cue investigation, a small juvenile

cue fish inside of a scintillation vial was placed in the circle in the investigate zone. In the

dominance assay, a freely-swimming novel cue fish (smaller than the focal) was added to the

tank. In the subordinate assay, a freely-swimming novel cue fish (larger than the focal) was

added to the tank. C) In the dominance and subordinate assay, we quantified the number of

approaches, where one fish swims within three body lengths directly towards any part of another

fish. If the approached fish swam away in any direction, it was counted as a displacement.

1215

Figure 2: Social behaviour and distances in the stable groups, socialized pairs, and stable pairs.

A) Frequency of social approaches, displacements, and times entering the territory / terracotta

pot in the rearing environments. B) Total agonistic efficiency (total displacements / total

approaches). C) Distance between all dyads of fish was measured twice for each tank (20 time

points each, 40 total time points). Mean distance between dyads was then calculated per tank

(each data point is the mean for one tank). The maximum distance lines are the lengths of the

diagonal along the bottom of the tanks. D) The minimum distance recorded between a dyad

across all time points. $*p < 0.05$, $**p < 0.01$, $***p<0.001$.

Figure 3: Social measures from the dominant (A-C) and subordinate (D-F) behaviour assays. A)

Social approaches (App.), displacements (Displ.), approaches received (App. Rec.), and

submissions (Submit) in the dominance behaviour assay. B) Focal fish David's score in the

dominance behaviour assay. C) Directional consistency (calculated based on approaches) in the

dominance behaviour assay. Filled triangles indicate directionality was significantly greater than

zero. Empty triangles indicate pairs were not significantly directional. D) Social behaviour in the

subordinate behaviour assay. B) Focal fish David's score in the subordinate behaviour assay. C)

Directional consistency (calculated based on approaches) in the subordinate behaviour assay.

*$p<0.05$.

1277 **Figure 4:** Focal fish water-borne cortisol (pg/mL/hr), corrected for body mass (g). Log

1278 transformed data are shown.

1300

Figure 5: Principal components analysis (PCA) of cortisol and behaviour—including open field (OF) time in the far and investigate zones; social cue (SC) time in and frequency entering the territory, close, far, and investigate zones; and dominant and subordinate behaviour assay approaches (App.), displacements (Displ.), approaches received (App. Rec.), submissions, and David's Score. A) A heatmap of eigenvalues showing the PCA variables that load on PC1 (25.1%), PC2 (18.2%), PC3 (14.3%), PC4 (10.5%), and PC5 (8.5%). Numerical values are shown for variables stronger than ±0.25. Rows are hierarchically clustered. B) Treatment differences in PC1. C) Treatment differences in PC5. Percentages refer to the amount of variance explained by that PC. *p<0.05. **p<0.01.