The cation channel mechanisms of subthreshold inward depolarizing currents in the VTA dopaminergic neurons and their roles in the chronic-stress-induced depression-like behavior

Jing Wang[1,3], Min Su[1,4], Dongmei Zhang[1,5], Ludi Zhang[1], Chenxu Niu[1], Chaoyi Li[1], Shuangzhu You[1], Yuqi Sang[1], Yongxue Zhang[1,6], Xiaona Du[1], Hailin Zhang[1,2]*

[1]Department of Pharmacology; The Key Laboratory of Neural and Vascular Biology, Ministry of Education; The Key Laboratory of New Drug Pharmacology and Toxicology, Hebei Medical University, Shijiazhuang, Hebei, China
[2]Department of Psychiatry, The First Hospital of Hebei Medical University, Mental Health Institute of Hebei Medical University, China
[3]Department of Pharmacochemistry, Hebei University of Chinese Medicine, Shijiazhuang, Hebei, China
[4]Yiling Pharmaceutical Company, Shijiazhuang, Hebei, China
[5]Department of Clinical Pharmacy, Xintai Ninth Hoapital, Xintai, Hebei, China
[6]Department of Pharmacy, Handan First Hospital, Handan, Hebei, China

* Correspondence:
Hailin Zhang
zhanghl@hebmu.edu.cn

Abstract:

The slow-intrinsic-pacemaker dopaminergic (DA) neurons originating in the ventral tegmental area (VTA) is implicated in various mood- and emotion-related disorders, such as anxiety, fear, stress and depression. Abnormal activity of projection-specific VTA DA neurons is the key factor in the development of these disorders. Here, we describe the crucial role for the NALCN and TRPC6, non-selective cation channels in mediating the subthreshold inward depolarizing current and driving the firing of action potentials of VTA DA neurons in physiological condition. Furthermore, we demonstrate that down-regulation of TRPC6 protein expression in the VTA DA neurons likely contributes to the reduced activity of projection-specific VTA DA neuron in chronic mild unpredictable stress (CMUS) depressive mice. In consistent with this, selective knockdown of TRPC6 channels in the VTA DA neurons conferred mice with depression-like behavior. This current study suggests down-regulation of

TRPC6 expression/function is involved in reduced VTA DA neuron firing and chronic stress-induced depression-like behavior in the mouse models of chronic stress-induced depression.

Keywords: Ventral tegmental area, Dopaminergic neurons, Cation channels, NALCN, TRPC6 channel

Introduction

Dopaminergic (DA) neurons in the ventral tegmental area (VTA) play a key role in mood, reward and emotion-related behaviors[1]. These DA neurons project, through mesocorticolimbic dopaminergic system, to brain regions including the nucleus accumbens (NAc), medial prefrontal cortex (mPFC) and basolateral amygdala (BLA), and modulate these target neuronal circuits through their regulated activity (firing of action potentials)[2, 3]. As such, understandably, the altered functional activity of the VTA DA neurons is found to be the key determinants in abnormal behaviors related to the diseased conditions such as depression- or anxiety-related states[4]. Thus, the study on the mechanism underlying the firing activity of the VTA DA neurons is crucial for understanding the function of the DA circuitry and the pathogenesis of related mental diseases[4-8].

The VTA DA neurons are slow intrinsic pacemakers, which are further modulated by synaptic inputs from multiple excitatory and inhibitory projections [9, 10]. Firing activity controls the release of DA, thus function of the VTA DA neurons. The altered activity of the VTA DA neurons in a pathological state includes altered firing frequency and switch of the firing patterns[7, 11].

The research on the mechanism for spontaneous firing of action potential (AP) is the key to understand functional regulation of the VTA DA neurons. A variety of ion channels (e.g. voltage-gated Na^+ channels, high-threshold-activated Ca^{2+} channels, Kv2, large conductance calcium-activated potassium channels, A- and M-type potassium channels) are reported to be involved in regulation of the spontaneous firing activity of midbrain DA neurons (including the substantial nigra pars compacta, SNc and VTA)[12-22]. However, it is not completely clear how this spontaneous firing is initiated. AP fires when the resting membrane potential (RMP) is depolarized

to the activation threshold for the voltage-dependent Na^+ channels, which results in
opening of Na^+ channels and the upstroke of AP[12]. Therefore, a controlled
depolarization of the RMP to the activation threshold of AP is the first key step in a
cascade leading to firing of AP. In the VTA DA neurons, the identity of ion channels
contributing to this subthreshold depolarization has not been unambiguously
established.

In some cells with spontaneous firing of AP, such as cardiomyocytes and certain
neurons (including a subset of midbrain SNc DA neurons[23]), the ionic mechanism
for the auto-rhythmicity is the HCN, a non-selective cation channel that would
depolarize the membrane potential after being activated at a hyperpolarized
membrane potential[23]. However, in the VTA DA neurons, results concerning the
HCN channels to the spontaneous firing are controversial[15, 23-25].

Another non-selective cation channel, NALCN, unlike HCN, is a non-voltage
dependent channel[26], and is reported to be among the candidates for molecular
correlates of the persistent background Na^+ leak currents in the VTA DA neurons [27,
28]. The NALCN-mediated background Na^+ leak currents are found to be involved in
modulation of firing activity in a variety of neurons including DA neurons in the
SNc[26, 28-31]. However, it has not been reported if NALCN plays a role in
spontaneous firing of the VTA DA neurons, although the VTA DA neurons have larger
Na^+ leak currents than SNc DA neurons do[15].

TRP channels are a large class of non-selective cation channels, and are widely
distributed in the peripheral and central nervous system. TRPC3, which has a high
homology with TRPC6[32], is found to be involved in the regulation of cardiac and
SNc DA neuronal excitability, generating depolarizing currents, triggering action
potentials and participating in cellular rhythmic firing[31, 33, 34].The role of TRP
channels, except TRPC4[35], in the regulation of VTA DA neuronal excitability has
not been described.

In this study, we drove to find channel conductance which contribute to the
subthreshold depolarization and thus the spontaneous firing of the VTA DA neurons.
And furthermore, we investigated if these channels are the molecular mechanism for
the altered function activity of the VTA DA neurons related to depression-like

behaviors in mouse model of depression. We started by profiling the expression of non-selective cation channels (NSCCs) of the VTA DA neurons using method of Patch-Seq from midbrain DA neurons projecting to different brain regions, and then focused on HCN, NALCN, TRPC6 and TRPV2 channels which are dominantly expressed in these DA neurons.

Results

Inflow of extracellular Na^+ contribute to the subthreshold depolarization of the VTA DA neurons

One key feature of the midbrain DA neurons is that these neurons are relatively depolarized in the resting condition; the resting membrane potential (RMP) is far from the K^+ equilibrium potential (~-55 mV *vs* ~-90 mV)[15], which enable the spontaneous firing. Thus, there must be persistent inward depolarizing conductance counterbalancing the hyperpolarizing K^+ conductance which otherwise would maintain the RMP at a hyperpolarized level. The main aim of this study is to identify the subthreshold depolarizing conductances which contribute to the spontaneous firing of the VTA DA neurons. While we initially performed our experiments using male mice, most of the experiments were also repeated using female mice, results of which were reported separately in the following results. We first observed firing frequency and the RMP of the VTA DA neurons using patch clamp recordings. The midbrain DA neurons, marked by dopamine transporter (DAT), mainly located in the VTA and SNc (**Fig. 1Ai**). We recorded neurons from the VTA brain slices of mice, and performed single-cell PCR post electrophysiologic recordings to identify the DA neurons with established markers for DA neurons (TH, DAT, D_2R (type2 dopamine receptor), and GIRK2 (G protein-gated inward rectifier K inwardly rectifying K^+ channels)). Although the majority of cells in the VTA area are DA neurons (~60%), there are still about 30% GABA neurons and a small number of glutamatergic neurons [36, 37]. Glutamatergic- and GABAergic neurons were also single-cell PCR typed by using markers of vGluT2 (vesicular glutamate transporter 2) and GAD1 (glutamic acid decarboxylase 1), respectively (**Fig. 1Aii**).

To focus on the intrinsic channel conductance for spontaneous firing, potential input modulation from fast-type transmitter transmission to the VTA DA neurons, namely

activation of AMPA/kainite/NMDA receptor and $GABA_A$ receptor were blocked by

receptor antagonists CNQX, DL-AP5 and picrotoxin [15], respectively, in the

following experiments.

We first studied whether inflow of extracellular Ca^{2+} contributed to initiation of

spontaneous firing in the VTA DA neurons. It has been shown that, unlike the SNc

DA neurons, the VTA DA neurons do not have subthreshold Ca^{2+} oscillatory

waves[15], suggesting that subthreshold Ca^{2+} should not be an important component

of subthreshold depolarizing conductance in the VTA DA neurons. In line with this

finding, we found in our study, replacing Ca^{2+} with Mg^{2+} from the extracellular

recording solution (ACSF, artificial cerebral spinal solution) did not reduce, but rather

instead, increased the firing frequency of the VTA DA neurons in male mice (**Fig. 1B**,

from 1.71 ± 0.19 Hz to 2.36 ± 0.28 Hz, n = 16, N = 8, $P < 0.001$). Consistent with this，

the RMP of the VTA DA neurons was not hyperpolarized by replacing Ca^{2+} with Mg^{2+}

from ACSF (**Fig. 1C**, from -50.77 ± 0.85 mV to -49.47 ± 1.09 mV, n = 14, N = 8, $P <$

0.05).

We then, in comparison with Ca^{2+}, observed the role of extracellular Na^+ in the RMP

of the VTA DA neurons. For this, we followed the method of a previous study[15], by

replacing Na^+ in ACSF with NMDG (N-methyl-d-glucamine); TTX (Tetrodotoxin)

was also added to block the TTX-sensitive Na^+ currents, which is known to be the

main contributor to the suprathreshold depolarization component of the action

potential. Replacement of the extracellular Na^+ with NMDG resulted in a

hyperpolarization of the RMP in mice (from -46.32 ± 1.97 mV to -64.53 ± 1.74 mV, n

= 8, N = 5, $P < 0.001$, **Fig. 1D**), indicating a persistent inflow of Na^+ through a

TTX-insensitive conductance which caused significant depolarization of the neurons,

contributing significantly to the subthreshold depolarization of the VTA DA neurons.

Similar results were obtained in the VTA DA neurons of female mice, with increased

firing frequency and hyperpolarized RMP when the extracellular $Ca^{2+}$and Na^+ were

replaced, respectively (**sFig. 1A** and **B**).

Taken together, above results indicate persistent Na^+ influx contribute to the

subthreshold depolarization of the VTA DA neurons.

Potential candidates of channel conductance contributing to subthreshold depolarization of the VTA DA neurons

The channel conductance mediating the above-described persistent Na^+ influx should come from a cation channel(s) permeating to Na^+. We thus investigated the presence and expression level of non-selective cation channels (NSCCs) in the VTA DA neurons using a combination of patch clamp and single-cell RNA-Seq (Patch-seq) technology[38]. In this part of the study, we considered the heterogeneity and projection-specificity of the VTA DA neurons, i.e. VTA DA neurons projecting to different brain regions have different electrophysiological characteristics[39]. Using retrograde labelling techniques using retrobeads, sixty male VTA neurons projecting to five different brain regions (medial prefrontal cortex, mPFC; basal lateral amygdala, BLA; nucleus accumbens core, NAc c; nucleus accumbens lateral shell, NAc ls; nucleus accumbens medial shell, NAc ms) (**sFig. 2**) were collected for RNA-Seq, using patch-clamp electrodes. Consistent with previous study[39], the VTA DA neurons projecting to above different brain regions were anatomically congregated into subregions of the VTA (**sFig. 3**). The high expression of biomarkers (Tyrosine hydroxylase, TH; Dopamine decarboxylase, Ddc and Dopamine transporter, DAT) in the 45 cells of 60 cells (**Fig. 2A**) indicates that these are DA neurons. In addition, less than half of sequenced DA neurons (predominantly those projecting to NAc ms, BLA, and mPFC) have biomarkers (Vesicular glutamate transporter, VGluT1-3) of glutamatergic neurons, predicting a possible coexistence of neurotransmitters (DA and glutamate)[40]. In contrast, fewer GABAergic neuronal markers (Glutamic acid decarboxylase, GAD1/2 and vesicular GABA transporter, VGAT) co-expressed with the DA neurons, which is consistent with previous studies that VTA DA neurons co-expressing GABAergic neuronal markers mainly project to the lateral habenula[41]. It needs to be noted that some neurons in the VTA only expressed markers for glutamatergic or GABAergic neurons, these neurons were excluded from further analysis as we focused our study on the DA neurons.

Expression levels of NSCCs in the VTA DA neurons classified by projection specificity were analyzed and shown in **Fig. 2B**. Different colored strips on the top of the figure corresponded to the projection specificity coded by the same colors in **Fig.**

2A. Multiple transient receptor potential channels (TRP) were present in the VTA DA neurons, with prominent expression level of TRPC6 and TRPV2 (**Fig. 2Bi**). Other prominently expressed NSCCs included HCN2, HCN3, NALCN and pannexin 1 (Panx1) (**Fig. 2Bii**). Summarized averaged relative expression levels for eight most richly expressed NSCCs were shown in **Fig. 2Biii**. Furthermore, projection-specific expression of these dominantly expressed NSCCs were analyzed and shown in **Fig. 2C**; interestingly only TRPC6 seemed expressed in a projection-specific manner, namely, the VTA DA neurons projecting to the NAc have higher TRPC6 expression than the VTA DA neurons projecting to the BLA and the mPFC **(Fig. 2C)**. In our subsequent study, we focused our study on HCN, NALCN, TRPC6 and TRPV2, one for they are the prominently expression channels, and two for some of these channels have been indicated in modulation of excitability of different neuron types[23, 24, 26, 28-31, 42]. Panx1 was investigated in a separate study since it normally composes the semi channels which are different from these more conventional ion channels.

HCN does not contribute to the spontaneous firing of the VTA DA neurons

We mainly used two pharmacological tools as HCN channel blockers, CsCl and ZD7288[15, 24] to explore the role of HCN channels in the spontaneous firing of the male VTA DA neurons. Efficient blocking effect of these blockers on HCN channel was first verified. For this, the sag potential produced by a hyperpolarizing current (-100 pA) injection was used to assess HCN activity[24]. Both CsCl (3 mM) and ZD7288 (60 μM) effectively reduced the sag potential (from 16.43 ± 2.96 mV to 3.84 ± 1.92 mV, n = 4, N = 4, $P < 0.05$; from 19.38 ± 3.14 mV to 4.18 ± 2.60 mV, n = 4, N = 4 respectively, $P < 0.05$) (**sFig. 4A and B**).

However, these same HCN blockers did not inhibit the spontaneous firing of the VTA DA neurons; CsCl slightly increased (1.96 ± 0.19 Hz to 2.14 ± 0.34 Hz, n = 7, N = 4) whereas ZD7288 slightly decreased the firing frequency (1.65 ± 0.25 Hz to 1.50 ± 0.24 Hz, n = 13, N = 5), but none of these effects were statistically significant ($P > 0.05$) (**sFig. 4C and D**). It has been reported that VTA DA neurons projecting to the NAc lateral shell have more pronounced HCN/Ih currents than other VTA DA neurons[39]. However, even the VTA DA neurons projecting to the NAc lateral shell

(retrogradely labelled by retrobeads), no effect of CsCl or ZD7288 on the spontaneous

firing of the male VTA DA neurons were observed (from 2.04 ± 0.24 Hz to $1.93 \pm$

0.22 Hz, n = 5, N = 4, by CsCl, and from 1.71 ± 0.29 Hz to 1.65 ± 0.33 Hz, n = 5, N =

5, by ZD7288, $P > 0.05$) (**sFig. 5A and B**).

HCN are activated by membrane hyperpolarization, with an activation threshold

around -70 ~ -90 mV (HCN2, HCN3)[43, 44]. At a depolarized RMP like these in the

VTA DA neurons (-51.28 ± 1.85 mV, n = 8, N = 8; e.g. -54.50 mV, **sFig. 4Ai**, and

-53.50 mV, **sFig. 4Bi**), HCN are most likely not activated thus would not participate

in generation of spontaneous firing. To further prove this, we tested if a more

hyperpolarized RMP would involve HCN in the spontaneous firing of the VTA DA

neurons. For this, we first lowered the K^+ concentration in the extracellular ACSF

from 3 mM to 1.5 mM to increase the gradient between inside and outside cellular K^+;

this maneuver indeed hyperpolarized the cell membrane from -53.33 ± 2.79 mV to

-65.00 ± 3.02 mV (n = 6, N = 6, $P < 0.01$) (**sFig. 5C**). Interestingly, under 1.5 mM

extracellular K^+, the spontaneous firing of the male VTA DA neurons was almost

totally blocked by ZD7288 (**sFig. 5D**), although an initial enhancement was seen in

some neurons (**sFig. 5Di**).

We tested another possible mechanism for HCN's failure to participate in generation

of VTA DA firing. It has been reported that in the midbrain SNc DA neurons, HCN is

involved in the spontaneous firing in young but not in adult mice, due to a more

hyperpolarization-shifted activation threshold of HCN in SNc DA neurons in adult

mice. We tested if this could also be the case in the VTA DA neurons. Indeed, as

shown in **sFig. 5E**, in contrast to what we saw in adult male mice, the spontaneous

firing frequency of the VTA DA neurons in young male mice (less than 15 days

postnatal) was significantly reduced by ZD7288 (from 1.86 ± 0.29 Hz to 0.98 ± 0.24

Hz, n = 9, N = 5, $P < 0.05$).

Taken together, above results suggest HCN is not involved in the spontaneous firing

of the VTA DA neurons in adult mice, due to a depolarized RMP and a

hyperpolarization-shifted activation property.

NALCN contributes to subthreshold depolarization and spontaneous firing of the VTA DA neurons

The Na^+ currents produced by NALCN have been suggested to be an important component of background Na^+ currents in multiple central neurons including DA neurons in the substantia nigra[28].We next investigated if NALCN also play a role in setting the RMP and in the spontaneous firing of the VTA DA neurons. Consistent with above RNA-Seq results, both immunofluorescence (**Fig. 3A**) and single-cell PCR (**Fig. 3B)** results confirmed broad expression of NALCN in the VTA DA neurons.

$GdCl_3$ (100 μM), a nonspecific NALCN blocker [29, 30], hyperpolarized the RMP of the VTA DA neurons from -52.17 ± 4.15 mV to -67.33 ± 2.68 mV (n = 6, N = 6, $P <$ 0.05) (**Fig. 3C**), reduced the spontaneous firing frequency of the VTA DA neurons from 1.89 ± 0.29 Hz to 1.32 ± 0.37 Hz (n = 7, N = 3, $P < 0.01$) (**Fig. 3D**). L703,606, a specific NALCN blocker[31], also hyperpolarized the RMP of the VTA DA neurons from -49.16 ± 1.48 mV to -54.30 ± 1.67 mV (n = 6, N = 4, $P < 0.001$) (**Fig. 3E)**, reduced the spontaneous firing frequency of the VTA DA neurons from 2.36 ± 0.38 Hz to 0.64 ± 0.32 Hz (n = 8, N = 4, $P < 0.001$) (**Fig. 3F**).

To observe a more specific effect on NALCN, a shRNA against NALCN was used to knockdown NALCN. For this, AAV9 viral construct (AAV9-U6-shRNA (NALCN)-CMV-GFP) was injected into the VTA of a mouse; similar AAV viral construct containing scramble-shRNA of nonsense sequences was utilized as controls. The qPCR results show sufficient knockdown of NALCN mRNA (NALCN-shRNA: 0.32 ± 0.07, N = 6; control scramble-shRNA: 1.07 ± 0.19, N = 6, $P < 0.01$, **Fig. 3G)** in the VTA tissue. Immunofluorescence results (**Fig. 3H**) show efficient infection of the VTA DA neurons (GFP expression in the DAT-positive neurons) by the virus. Knockdown of NALCN in the VTA DA neurons by shRNA almost completely silenced the firing of the male VTA DA neurons (0.07 ± 0.04 Hz *vs* 1.84 ± 0.25 Hz in shRNA (n = 12, N = 3) and scramble-shRNA (n = 10, N = 3) infected VTA DA neurons, respectively, $P < 0.001$, **Fig. 3I**). Furthermore, the RMP of the male VTA DA neurons was significantly hyperpolarized (-60.67 ± 2.33 mV *vs* -48.9 ± 1.45 mV in shRNA and scramble-shRNA infected VTA DA neurons, respectively, n = 18 and 15,

N = 5 and 5, $P < 0.001$, **Fig. 3J**). In female mice, L703,606 also reduced the
spontaneous firing frequency of the VTA DA neurons and knockdown of NALCN
also silenced the firing of the VTA DA neurons (**sFig. 6A** and **B**).

Taken together, above results suggest that NALCN is a major contributor to the
subthreshold depolarization and contribute significantly to the generation of
spontaneous firing in the VTA DA neurons.

TRPC6 contributes to subthreshold depolarization and spontaneous firing of the
288
VTA DA neurons
289

Above RNA-Seq results suggest a broad and strong expression of TRPC6 and TRPV2
channels in the VTA DA neurons. This part of the experiments was focused on the
role of these two TRP channels in subthreshold depolarization and spontaneous firing
of the VTA DA neurons.

Consistent with the RNA-Seq results, single-cell PCR experiments confirmed high
proportion expression of TRPC6 and TRPV2 in the VTA DA neurons (**Fig. 4A**);
among the 28 VTA DA neurons (DAT$^+$), 20 neurons expressed TRPV2 (71.4%), and
18 neurons (64.3%) expressed TRPC6. TRPC3, which is highly homologous with
TRPC6[45] and has been reported to be involved in the firing activity of the SNc DA
neurons[31], was not detected in RNA-Seq study (**Fig. 2B**), nor in the single-cell PCR
experiment (**Fig. 4A**). These results reciprocally confirmed the reliability of both
RNA-Seq and single cell PCR results.

General roles of TRP channels in subthreshold depolarization and spontaneous firing
of the VTA DA neurons were first assessed using a non-specific broad-spectrum TRP
channel blocker, 2-aminoethoxydiphenylborane (2-APB; 100 μM) and a specific
TRPC6 channel blocker, larixyl acetate[46] (LA; 10 μM). Both blockers significantly
reduced the firing frequency of the male VTA DA neurons (2-APB: from 1.32 ± 0.27
Hz to 0.06 ± 0.02 Hz, n = 12, N = 5, $P < 0.001$; LA: from 1.16 ± 0.21 Hz to $0.39 \pm$
0.09 Hz, n = 7, N = 5, $P < 0.01$; **Fig. 4B, 4C**). Accordingly, both blockers also
significantly hyperpolarized the RMP of the male VTA DA neurons (2-APB: from
-51.01 ± 2.20 mV to -61.64 ± 1.28 mV, n = 6, N = 6, $P < 0.001$; LA: from $-53.93 \pm$
1.05 mV to -57.48 ± 1.09 mV, n = 5, N = 4, $P < 0.01$; **Fig. 4D, 4E**).

We then tested a more TRPV channel-selective blocker ruthenium red (RR), to study

possible role of TRPV2 (and other TRPV) in the spontaneous firing of the VTA DA

neurons. RR (60 μM), on average statistically, did not affect the spontaneous firing of

the male VTA DA neurons (from 2.30 ± 0.31 Hz to 2.18 ± 0.32 Hz, n = 9, N = 6, $P >$

0.05) (**sFig. 7**). However, it needs to be noted that among the nine VTA DA neurons

we tested, both increase (**sFig. 7B**) and decrease (**sFig. 7C**) of firing frequency by RR

were seen, thus with opposite changes of firing frequency in different populations of

neurons (**sFig. 7A**), no overall effect of RR was obtained from the current analysis.

We then focused our next study on TRPC6. Consistent with both RNA-Seq (**Fig. 2B**)

and single-cell PCR (**Fig. 4A**) results, the mRNA level of TRPC6 in mPFC-projecting

VTA single DA neuron is significantly lower than that in NAc c-projecting VTA

single DA neuron (0.56 ± 0.12, n = 25, N = 8 for mPFC-projecting TH-positive cells

vs 1.09 ± 0.10, n = 28, N = 7 for NAc c-projecting TH-positive cells,** $P < 0.01$, **Fig.**

5A) The immunofluorescence results also demonstrated strong expression of TRPC6

protein in the VTA DA neurons (**Fig. 5B**, TH-positive). A AAV viral construct with

shRNA against TRPC6 (AAV9-U6-shRNA (TRPC6)-CMV-GFP) was injected into

the VTA of mice, to knockdown TRPC6; efficiency of this knockdown was assessed

by using qPCR measuring the mRNA level of TRPC6 in the VTA tissue (**Fig. 5C**),

and efficiency of viral infection into the VTA DA neurons (marked by DAT) was

observed through visualization of GFP (**Fig. 5D**). Both qPCR (**Fig. 5C**) and

immunofluorescence (**Fig. 5D**) results indicated a sufficient repression of TRPC6 in

the VTA DA neurons. Indicative of a significant role in the spontaneous firing and

subthreshold depolarization, knockdown of TRPC6 (TRPC6-KD) resulted in a

substantial reduction in the spontaneous firing frequency (0.80 ± 0.25 Hz, n = 9, N=3

for TRPC6-KD *vs* 1.90 ± 0.27 Hz, n = 9, N = 3 for scramble-shRNA, $P < 0.01$, **Fig.**

5E), and hyperpolarization of the RMP (-58.21 $\pm$ 1.79 mV, n = 10, N = 6 for

TRPC6-KD *vs* -50.02$\pm$ 0.67 mV, n = 10, N = 6 scramble-shRNA, $P < 0.001$, **Fig. 5G**)

of the male VTA DA neurons. In addition, the inhibitory effect of 2-APB and LA on

the firing of VTA DA neurons infected with TRPC6-shRNA largely diminished, with

no statistical difference in firing frequency before and after dosing (2-APB: from 0.58

$\pm$ 0.15 Hz to 0.40 $\pm$ 0.15 Hz, n = 7, N = 5, $P > 0.05$, **Fig. 5F**; LA: from 0.43 ± 0.05 Hz

to 0.43 ± 0.06 Hz, n = 8, N = 5, $P > 0.05$, **Fig. 5H**). While the above results were observed in the male mice, similar results were observed in the female mice (**sFig.8 A** and **B**).

Consequence of knocking down TRPC6 described above should not be a result of a secondary effect on other TRPC channels with which TRPC6 is known to interact to form heteromers, such as TRPC4 and TRPC7[47, 48], since in experiments using single cell PCR (**sFig. 9A**), it was found only a very small proportion of TRPC6-positive DA cells (DAT$^+$) expressed TRPC4 (**sFig. 9Bi**) or TRPC7 (**sFig. 9Bii**), in consistent with the results of single-cell RNA-seq results (**Fig. 2**).

Taken together, above results indicate TRPC6 is an important contributor in subthreshold depolarization and spontaneous firing of the VTA DA neurons.

Down-regulation of TRPC6 contributes to the altered firing activity of the VTA DA neurons in depression model

In multiple depression models, the depression-like behavior was directly linked to the altered firing activity of the VTA DA neurons[4, 7, 49-51]. In consideration of the evidence we described above that NALCN and TRPC6 play key roles in firing activity of the VTA DA neurons, we went further to study if these channels also contributed to the altered firing activity of the VTA DA neurons and to the development of the depression-like behavior in depression models of mice.

We first established a mice depression model of chronic mild unpredictable stress (CMUS) [52], which, after 5 weeks subjecting to two different stressors every day (**Fig. 6A**), manifested chronic-stress-induced depression-like behaviors in the sucrose preference test (SPT) and the tail suspension test (TST), with reduced sucrose preference and lengthened immobility time, respectively (**Fig. 6B**). Multiple other behaviors tests were also performed on these CMUS mice (**sFig. 10**), all indicating depression/anxiety-like behaviors.

It has been shown that a reduced firing activity in the VTA DA neurons is responsible for the chronic-stress-induced depression-like behavior in the CMUS model[51, 53]. Consistent with this finding, we also found the firing frequency of the VTA DA neurons from the CMUS male mice was significantly reduced as compared with that

from the control male mice (2.83 ± 0.42 Hz, n = 18, N = 5 for control mice *vs* 1.30 ± 0.14 Hz, n = 28, N = 5 for CMUS mice, $P < 0.01$, **Fig. 6C**). In agreement with a role for TRPC6 in this reduced firing activity, TRPC6 protein in the male VTA tissue was found to be down-regulated in western blot experiments (**Fig. 6D**). On the other hand, NALCN protein in the VTA was not altered in the CMUS male mice model of depression (**sFig. 11**).

We also tested if this down-regulation of TRPC6 protein could also be found in a similar but different model of depression. For this, a chronic restraint stress (CRS) model was used. After 3 weeks of restraint stress stimulation, the CRS male mice developed similar depression/anxiety-like behavior in the behavior tests like these similarly performed in the CMUS model (**sFig. 12A-I**). Importantly, male mice with CRS-induced depression also showed a significant downregulation of TRPC6 protein in the VTA (0.43 ± 0.09 fold, N = 6 for control and N = 9 for CRS, $P < 0.001$, **sFig. 12J**). In female CMUS mice (**sFig13. A-I**), TRPC6 protein in the VTA tissue was also found to be down-regulated in western blot experiments (**sFig. 13J**).

It is known that CMUS mainly decreases the activity of VTA dopamine neurons that project to mPFC[11, 51]. Assuming that the decreased firing activity of the mPFC-projecting VTA DA neurons in the CMUS mice was associated with the down-regulation of TRPC6 expression, we reasoned that the firing activity in these CMUS mice would respond less to the TRP channel inhibitors. Indeed, TRP channel inhibitor 2-APB and TRPC6 inhibitor LA decreased their inhibitory effect on the firing activity of the mPFC-projecting VTA DA neurons from the CMUS male mice (2-APB: 1.61 ± 0.23 Hz to 1.19 ± 0.22 Hz, n = 15, N=5, $P > 0.05$; LA: 0.97 ± 0.15 Hz to 0.66 ± 0.09 Hz, n = 19, N = 8, $P > 0.05$) compared with these from the control male mice (2-APB: 3.74 ± 0.42 Hz to 0.17 ± 0.09 Hz, n = 12, N = 5, $P < 0.001$; LA: 2.83 ± 0.17 Hz to 0.10 ± 0.07 Hz, n = 17, N = 8, $P < 0.001$) (**Fig. 6E** and **F**).

Taken together, above results suggest down-regulation of TRPC6 is a key determinant for the altered firing activity of the VTA DA neuron in mice models of depression

Down-regulation of TRPC6 in the VTA DA neurons confer the mice with depression-like behavior

If down-regulation of TRPC6 is crucial for the depression-like behaviors, like that indicated above in the CMUS and the CRS depression models, then down regulation of TRPC6 in non-stressfully treated mice should also develop depression-like behavior. We tested this possibility by selectively knocking down TRPC6 in the VTA DA neurons, using DAT-Cre male mice injected with a AAV viral construct (AAV9-hSyn-DIO-shRNA (TRPC6) -RFP) (**Fig. 7A**), driving the expression of shRNA against TRPC6 selectively in the VTA DA neurons. Three weeks after the AAV injection, the qPCR (**Fig. 7B**) and the immunofluorescence results (**Fig. 7C**) indicated an efficient down-regulation of TRPC6 in the VTA DA neurons.

We next compared the depression- and anxiety-like behaviors in these conditionally TRPC6-knockout male mice (TRPC-cKD) with that in the control mice (infected with virus carrying the scrambled shRNA). As shown in **Fig. 7D**, in the TRPC6-cKD mice, the sucrose preference was significantly reduced (7Di, $72.60 \pm 3.52\%$, N = 13 for TRPC6-cKD, *vs* 92.74 ± 0.86 %, N = 9 for control, $P < 0.001$), the immobility time in the TST was significantly lengthened (7Dii, 192.10 ± 3.13 s *vs* 133.20 ± 8.89 s, $P < 0.001$). In the elevated plus maze test, the percentage of time spent in the open arm was significantly reduced (7Diii, 2.43 ± 0.48 % *vs* 14.78 ± 1.26 %, $P < 0.001$) and the percentage of time spent in the closed arm was also significantly increased (7Div). And the Cre-dependent TRPC6-overexpression in male VTA DAT-Cre mice (TRPC6-cOE) rescued the selective-TRPC6-knockdown-induced depression-like behaviors (**Fig. 7E-H**). Furthermore, selectively knockdown of TRPC6 in the mPFC-projecting VTA DA neurons (virus-retro carrying DIO-TRPC6-shRNA (TRPC6-rcKD) was injected into the mPFC of DAT-Cre male mice) (**Fig. 8A-C**) also conferred the mice with the depression- and anxiety-like behaviors (**Fig. 8D**).

Taken together, these results suggest downregulation of TRPC6 in the VTA DA neurons confers the mice with phenotypes of depression/anxiety.

Discussion

In this study, we made a systematic study on the molecular mechanism for the subthreshold depolarization that drive spontaneous firing of the VTA DA neurons. We identified TRPC6 channels, alongside NALCN, as major contributors to this

subthreshold depolarization and related spontaneous firing, and further importantly, we also demonstrated that TRPC6 contributed to the altered firing activity of the VTA DA neurons under states of chronic-stress-induced depression-like behaviors.

The nature of a relatively depolarized resting membrane potential in the VTA DA neurons imposes a need for a full understanding of the channel conductance underlying this depolarization. It is first clear from these current and previous studies that this conductance is mediated by a persistent Na^+ influx. Unlike the DA neurons in the SNc, where Ca^{2+} influx is needed for the spontaneous firing [54], the results in our and others' studies indicate Ca^{2+} influx are not necessary for the spontaneous firing of the VTA DA neurons, rather, replacement of Ca^{2+} ions in the extracellular fluid with Mg^{2+} accelerates the frequency of spontaneous firing (Fig. 1B) [15], possibly due to a reduced activity of the Ca^{2+}-activated K^+ currents and the increased activity of NALCN[28, 55]. Consistent with these finding and more precisely for a role of Ca^{2+} influx on the subthreshold depolarization, removing extracellular Ca^{2+} did not hyperpolarize the RMP. On the other hand, consistent the previous study [15], a TTX-insensitive Na^+ influx clearly contributed to the subthreshold depolarization (Fig. 1D).

A TTX-insensitive nature of this Na^+ conductance suggests that cation conductance permeable to Na^+ might be the molecular candidates. We first focused on the HCN channels because, (1) it has been suggested HCN channels are important regulators for the spontaneous firing of VTA DA neurons under states of depression-like behaviors[7, 11]; (2) HCN are one of the more highly expressed NSCCs in the VTA DA neurons in our single cell RNA-Seq study; (3) HCN are permeable to Na^+[56]. However, it is clear from our study that HCN does not contribute to the spontaneous firing of the VTA DA neurons in the adult mice; none of the known HCN blockers we used had any significant effects on the firing activity. Presence of HCN has long been used as a signature of DA neurons[57], also as a differentiating factor for the projection-specificity of the VTA DA neurons[39]. Nonetheless, contribution of HCN to the spontaneous firing of the VTA DA neurons has been controversial; the HCN blockers ZD7288 and CsCl have been reported as both effective[23, 24, 58] and

non-effective [15] on the spontaneous firing of the VTA DA neurons. A closer

inspection into our and others' results present a possible explanation for these

different results: in the younger rodents (less than one month old) (sFig.5) [59], the

HCN in the DA neurons play a more important role whereas in the adult mice (~ 8

weeks)[59], like the ones we used in this study, the HCN do not contribute to the

spontaneous firing of the VTA DA neurons. This is likely due to a shift of activation

gating of HCN channels in a hyperpolarization direction with age, thus the

modulation of HCN on the firing activity diminishes with age, as it is described in the

SNc DA neurons[59]. Consistent with this, we found hyperpolarizing the RMP

rendered the VTA DA neurons of the adult mice with sensitivity to the HCN blockers.

Thus, the VTA DA neurons in adult mice do not use HCN to depolarize the membrane

to generate action potentials.

NALCN is widely expressed in central neurons[60, 61], and is a popular candidate

considered for the ion mechanism of the subthreshold depolarization currents and Na^+

leaky currents. NALCN has been shown to be an important component of background

Na^+ currents and to be important for neuronal excitability in the hippocampal neurons,

posterior rhomboid nucleus (Retrotrapezoid nucleus/RTN)) chemo-sensitive neurons

(CO_2/H^+-sensitive neurons), GABA neurons and DA neurons in the substantia nigra,

and spinal projection neurons [26, 28-31, 42]. Our results with single cell RNA-Seq

and immunofluorescence methods validated high level expression of NALCN in the

VTA DA neurons; both the pharmacological and the NALCN-gene-knockout

experiments demonstrated convincingly involvement of NALCN in subthreshold

depolarization and spontaneous firing of the VTA DA neurons.

We did not detect alteration of NALCN protein expression in the depression model of

CMUS in the VTA tissue. However, this does not completely exclude a possible

contribution of NALCN to the altered functional activity of the VTA DA neurons

which underlies the chronic-stress-induced depression-like behaviors. Future study

focusing on the functional property of NALCN in the cellular level of the VTA DA

neurons under a state of depression is needed to clarify this issue. NALCN is after all

a potential drug target in consideration of its important contribution to the resting

membrane potential and functional activity of neurons.

The most interesting and important finding of this study is the role of TRPC6 in

regulation of the firing activity of the VTA DA neurons in both physiological and

depression-state conditions, and its involvement in depression-like behaviors; the

experimental evidence from pharmacological, gene silencing, electrophysiological

and behavior results unambiguously support this conclusion. It is interesting to note,

although with an unclear neuronal circuit mechanism of action, the TRPC6 channel

opener hyperforin was previously reported to have antidepressant effects in

corticosterone-depressed mice, which were reversed by prior administration of the

TRPC6 blocker larixyl acetate[62]. Also, in a CMUS rat model, TRPC6 expression

was found to be downregulated in the hippocampus, and administration of the TRPC6

opener hyperforin was effective in alleviating depression-like behavior[63]. Our study

provides strong and convincing evidence that TRPC6 is a key regulator of the VTA

DA neurons which are known to be a key player in depression states [7, 21, 49, 50,

64].

The above-mentioned role for TRPC6 should come from its contribution to the

subthreshold depolarization, through a persistent permeation to the influx of Na^+.

Although TRPC channels has been better recognized for their permeability to Ca^{2+}

and their regulation on cellular Ca^{2+} homeostasis, the importance of TRPC channels

for the permeability of monovalent cations, especially Na^+, is now gradually being

appreciated [65]. In the substantial nigra GABA and DA neurons, TRPC3 has been

reported to play an important role in maintaining depolarization membrane potential,

pacemaking, and firing regularity[31, 33]. Otherwise, it has been reported that slow

tonic firing of SNc DA neurons, depends on the basal activity of both the NALCN and

TRPC3 channels, but that burst firing does not require TRPC3 channels but relies

only on NALCN channels[66]. TRPC6 and TRPC3 belong to the same TRPC family

subclass and TRPC6 is more than 75% homologous to TRPC3[45]. Interestingly, our

results with single-cell RNA-Seq and single-cell PCR demonstrated the rich

expression of TRPC6 in the VTA DA neurons among all TRP channels but no

expression of TRPC3 was found in these neurons. These results present an

opportunity for selectively targeting a single TRPC channel to exert selective pharmacological results. Besides, TRPC4 expression was detected in a uniformly distributed subset of VTA DA neurons[35], which is consistent with our results with single-cell RNA-Seq, and TRPC4 KO rats showed decreased VTA DA neuron tonic firing and deficits in cocaine reward and social behaviors[35].

It has also been demonstrated in cell types other than neurons, such as rat portal vein smooth muscle cells, the activation of α-adrenergic receptor can open TRPC6, which leads to cell depolarization, activation of voltage-dependent Ca^{2+} channels, and ultimately the contraction of smooth muscle cell[67]; it is also shown that angiotensin II and endothelin I can activate TRPC3/C6 heterodimers via vascular G protein-coupled receptors (GPCR), thereby depolarizing cells [68, 69]. Following this line, it would be very interesting to know if TRPC6 in the VTA DA neurons, where multiple GPCR reside receiving input and auto modulation by neuronal transmitters [70, 71], could also be a target of modulation with similar mechanism. In consideration of the important role, we described here in this study, this type of modulation will present possible explanations for some unsolved mechanism for physiological and pathophysiological functions of the VTA DA neurons.

The facts that down-regulation of TRPC6 proteins was correlated with reduced firing activity of the VTA DA neurons, the chronic-stress-induced depression-like behaviors, and that knocking down of TRPC6 in the VTA DA neurons confers the mice with depression-like behaviors strongly suggest a crucial role for TRPC6 in the development of chronic-stress-induced depression-like behaviors under stressful conditions like CMUS. To reinforce this, down-regulation of TRPC6 was also found in another chronic-stress-induced depression model of CRS, a model with similar neuronal alteration of reduced firing activity in the VTA DA neurons to that described in the CMUS model[72]. These facts with the fact that TRPC6 activator hyperforin is an effective antidepressant in multiple depression models[62, 63], and that hyperforin is the principal component of St. John's wort, a well-known antidepressant herb[73], present TRPC6 as a very attractive drug target for new lines of antidepressants.

Materials and methods

Animal Preparation

Male or female 6-8-week-old C57BL/6 (Vital River, China) and DAT-Cre mice with a C57BL/6 background (Stock No: 006660, the Jackson Laboratories, USA) were used for the studies. All experiments were conducted in accordance with the guidelines of Animal Care and Use Committee of Hebei Medical University and approved by the Animal Ethics Committee of Hebei Medical University.

Brain slice preparation

The details for the preparation of coronal brain slice containing VTA were the same as described in our previously published work[21, 70]. Briefly, mice were anesthetized with chloral hydrate (200 mg/kg, i.p.). After intracardial perfusion with ice-cold sucrose solution, the brains of the mice were removed quickly and placed into the slicing solution. The ice-cold sucrose-cutting solution contained (in mM): sucrose 260, $NaHCO_3$ 25, KCl 2.5, NaH_2PO_4 1.25, $CaCl_2$ 2, $MgCl_2$ 2, and D-glucose 10; osmolarity, 295-305 mOsm). Using the vibratome (VT1200S; Leica, Germany), the coronal midbrain slices (250 μm-thick) containing VTA were sectioned. The slices were incubated for 30 min at 36°C in oxygenated artificial cerebrospinal fluid (ACSF) (in mM: NaCl 130, $MgCl_2$ 2, KCl 3, NaH_2PO_4-$2H_2O$ 1.25, $CaCl_2$ 2, D-Glucose 10, $NaHCO_3$ 26; osmolarity, 280-300 mOsm), and were then left for recovery for 60 min at room temperature (23-25°C) until use.

The brain slices were transferred to the recording chamber and were continuously perfused with fully oxygenated ASCF during recording.

Identification of DA neurons and electrophysiological recordings

Recordings in the slices were performed in whole-cell current-clamp and voltage-clamp configurations on the Axoclamp 700B preamplifier (Molecular Devices, USA) coupled with a Digidata 1550B AD converter (Molecular Devices, USA). Neurons in the VTA were visualized with a 40X water-immersion objective equipped by an optiMOS microscope camera (Qimaging, Canada) on an Olympus-BX51 microscope (Olympus, Japan). Projection-specific or GFP-positiveVTA neurons were identified by infrared-differential interference contrast (IR-DIC) video microscopy

and epifluorescence (Olympus, Japan) for detection of retrobeads (red) positive or GFP-positive neurons.

For whole-cell recording, glass electrodes (3-5 MΩ) were filled with internal solution (in mM): K-methylsulfate 115, KCl 20, $MgCl_2$ 1, HEPES 10, EGTA 0.1, MgATP 2 and Na_2GTP 0.3, pH adjusted to 7.4 with KOH. And the extracellular solution was the ACSF.

In whole-cell current-clamp mode, HCN function was judged by the inwardly rectifying characteristic sag potential generated by giving a hyperpolarizing current (-100 pA).

When recording the effect of removing extracellular Ca^{2+} on spontaneous cell discharge, $CaCl_2$ in the ACSF was replaced with $MgCl_2$, and the final $MgCl_2$ concentration was 4 mM.

The resting membrane potential (RMP) was measured in current clamp mode (I = 0); the composition of the recording solution (mM) contained: NaCl 151, KCl 3.5, $CaCl_2$ 2, $MgCl_2$ 1, glucose 10, and HEPES 10, and the pH was adjusted to about 7.35 with NaOH. 1 μM tetrodotoxin was added to the extracellular solution to abolish action potential during measurement of the RMP.

To observe the effect of extracellular Na^+ on the RMP, NaCl in the original recording solution was replaced with equimolar NMDG, and the NMDG-recording solution (mM) contained: NMDG 151, KCl 3.5, $CaCl_2$ 2, $MgCl_2$ 1, glucose 10, and HEPES 10, and the pH was adjusted to 7.35 with KOH.

For recording spontaneous firing of the neurons, cell-attached "loose-patch" (100-300 MΩ) recordings were used[74]. In this case, patch-pipettes (2-4 MΩ) were filled with ACSF, and the spontaneous activity was recorded in the current-clamp mode (I = 0). The synaptic blockers (CNQX, 10 μM; APV, 50 μM and gabazine, 10 μM) were added to isolate the intrinsic firing properties.

At the end of electrophysiological recordings, the recorded VTA neurons were collected for single-cell PCR. VTA DA neurons were identified by single-cell PCR for the presence of TH and DAT.

Retrograde labelling

Briefly, under general chloral hydrate anaesthesia (200 mg/kg, i.p.) and stereotactic control (RWD Instruments, Guangzhou, China), the skull surface was exposed. All coordinates are relative to bregma in mm using landmarks and neuroanatomical nomenclature that was described in the Franklin and Paxinos mouse brain atlas[75] . Red retrobeads (100 nl for single injection; Lumafluor Inc., Naples, FL, USA) were injected into the following sites using KD scientific syringe pump (KD scientific, Holliston MA, USA): bilaterally into NAc core (NAc c), NAc lateral shell (NAc ls), NAc medial shell (NAc ms) and basolateral amygdala (BLA), 4 separate sites (2 per hemisphere) into medial prefrontal cortex (mPFC). Coordinates for infusions were as follows: NAc c (AP +1.50, LM 0.84, DV -4.0; 100 nl beads); NAc ls (AP +0.86, LM 1.72, DV -4.0; 100 nl beads); NAc ms (AP +1.70, LM 0.53, DV -4.0; 100 nl beads); mPFC (AP +2.05 and 2.15, LM 0.27, DV -2.1 + 1.7; 200 nl beads); BLA (AP -1.46, LM 2.85, DV -4.3; 100 nl beads). Retrobeads were delivered through a pulled glass pipette using a PAP107 Multipipette Puller (MicroData Instrument, Inc., USA) and at a rate of 100 nl/min; the injection needle was left in place for at least 5 minutes after each infusion. Following surgery, mice were returned to single housing. For sufficient labelling, survival periods for retrograde tracer transport depended on respective injection areas: NAc c, NAc ls, NAc ms, 14 days; mPFC, 21 days; BLA 14 days[39]. Coronal sections of injection sites were stained with 4, 6-diamidino-2-phenylindole (DAPI, Sigma, USA) to confirm representative target location. Then, serial analyses of the injection-sites were carried out routinely.

AAV for gene knockdown or overexpression and viral construct and injection

For knockdown of NALCN and TRPC6, AAV9-U6-shRNA (NALCN)-CMV-GFP (300 nl) or AAV9-U6-shRNA (TRPC6)-CMV-GFP or its control AAV9-scramble-shRNA was delivered into the VTA (AP, −3.08 mm; ML, ±0.50 mm; DV, −4. 50 mm) of the mice. AAV9-hSyn-DIO-shRNA (TRPC6)-RFP (300 nl) or its control AAV9-scramble-shRNA was delivered into the VTA of the DAT-Cre C57BL/6 mice. AAV-retro-hSyn-DIO-mCherry-shRNA (TRPC6) (300 nl) or its control AAV-retro-scramble-shRNA was delivered into the mPFC of the DAT-Cre C57BL/6 mice.

The shRNA hairpin sequences used in this study:

NALCN shRNA: AAGATCGCACAGCCTCTTCAT[26];

TRPC6 shRNA: 5'-CCAGGATCAATGCATACAA-3'.

For Cre-dependent overexpression of TRPC6, AAV9-DIO-TRPC6-GFP (300 nl) or its

control AAV9-scramble-RNA was delivered into the VTA of the DAT-Cre C57BL/6

mice, 7 days after injection of AAV9-hSyn-DIO-shRNA (TRPC6)-RFP.

Mice were singly housed with enough food and water to recover for 4-5 weeks after

injection of virus, before behavior tests and electrophysiological recordings.

Single cell PCR

mRNA was reversely transcribed to cDNA by PrimeScript^{TM}II1st Strand cDNA

Synthesis Kit (Takara-Clontech, Kyoto, Japan). At the end of electrophysiological

recordings, the recorded neuron was aspirated into a pipette and then expelled into a

PCR sterile tube containing 1 μl oligo-dT Primer and 1 μl dNTP mixture. The mixture

was heated to 65°C for 5 min to denature the nucleic acids and then cooled on ice for

2 min. Reverse transcription from mRNA into cDNA was performed at 50°C for 50

min and then 85°C for 5 sec. cDNA was stored at -40°C. Then two rounds of

conventional PCR with pairs of gene-specific outside (first round) and inner primers

(second round) for GAPDH (positive control), TH, DAT, D2, GIRK2, Vgult2, GAD1,

NALCN, TRPC6, TRPV2 and TRPC3 using GoTaq Green Master Mix (Promega,

Madison, USA) were performed. After adding the specific outside primer pairs into

each PCR tube, first-round synthesis conditions were as follows: 95 °C (5 min); 30

cycles of 95 °C (50 s), 58-63 °C (50 s), 72 °C (50 s); 72 °C(5 min). Then, the product

of the first PCR was added in the second amplification round by using specific inner

primer (final volume 25 μl). The second amplification round consisted of the

following: 95 °C (5 min); 35 cycles of 95 °C (50 s), 58 °C-62 °C (45 s), 72 °C (50 s)

and 5 min elongation at 72 °C. The final PCR products were separated by

electrophoresis on 2% agarose gels. The negative control reactions with no added

template were also performed in each experiment.

The "outer" primers (from 5′ to 3′) as follows:

GAPDH:

AAATGGTGAAGGTCGGTGTGAACG（sense）

AGTGATGGCATGGACTGTGGTCAT （antisense）

TH:

GCCGTCTCAGAGCAGGATAC

GGGTAGCATAGAGGCCCTTC

DAT:

CTGCCCTGTCCTGAAAGGTGT

GCCCAGTGATCACAGACTCC

D2-R:

AGCATCGACAGGTACACAGC

CCATTCTCCGCCTGTTCACT

GIRK2:

TGGACCAGGATGTGGAAAGC

AAACCCGTTGAGGTTGGTGA

Vgult2:

CTGCTTCTGGTTGTTGGCTACTCT

ATCTCGGTCCTTATAGGTGTACGC

GAD1:

ACAACCTTTGGCTGCATGTGGATG

AATCCCACGGTGCCCTTTGCTTTC

NALCN:

TCCATCTGTGGGAAGCATGT

CAAAAGCTGGTCCTCTTCAGTG

TRPC6:

AGCCTGTCTATTGAGGAAGAAC

AGCGAGAATGATTGGGGTCA

TRPV2:

CTGCACATCGCCATAGAGAA

AGGCTGGTGGTAGGCAACTA

TRPC3:

GCTGGCCAACATAGAGAAGG

CCTGCACGTGACTATCCACA

TRPC4:

GTCTATGTAGGCGATGCGCT

AGGAGGTCCTTGGCAAATTGT

TRPC7:

GCCATCAGCAAGGGCTATGT

CACGCCCACCACAAAATCC

The "inner" primers (from 5′ to 3′) as follows:

GAPDH:

GCAAATTCAACGGCACAGTCAAGG

TCTCGTGGTTCACACCCATCACAA

TH:

AGGAGAGGGATGGAAATGCT

ACCAGGGAACCTTGTCCTCT

DAT:

ATTTTGAGCGTGGTGTGCTG

TGCCTCACAGAGACGGTAGA

D2-R:

CCATTGTCTGGGTCCTGTCC

GTGGGTACAGTTGCCCTTGA

GIRK2:

AGCCGAGACAGGACCAAAAG

ATGTACGCAATCAGCCACCA

Vgult2:

CATCTCCTTCTTGGTGCTTGCAGT

ACAGCGTGCCAACGCCATTTGAAA

GAD1:

AGTCACCTGGAACCCTC

GCTTGTCTGGCTGGAA

NALCN:

GCCTTTGCTGGAGTTGTTCTG

CCCTGCATAATTGCCACAGTC

TRPC6:

TGCTAGAAGAGTGTCATTCCCT

CCTCCACAATCCGTACATAACC

TRPV2:

GTGGGATGTGGTGACCTACC

GCTGGTACAGCCCTGAGAAC

TRPC3:

CCTTGGGTCTTCCATTCCTC

CACAACTGCACGATGTACTCC

TRPC4:

CCACGAGGTCCGCTGTAAC

CTCCCAACTTAACTGAAAGGCA

TRPC7:

CGCTTCTCCCACGACATCAC

ACTGGATAGGGACAGGTAGGC

RT-qPCR

Total RNA was prepared using Trizol. RNA was reversely transcribed into cDNA by

TAKARA PrimeScripttm RT reagent Kit with gDNA Eraser for Reverse transcription.

Subsequently, the SYBR Green Master Mix (TaKaRa) was used to do the qRT-PCR

assays.

The specific primers were:

GAPDH：

GCAAATTCAACGGCACAGTCAAGG

TCTCGTGGTTCACACCCATCACAA

TRPC6：

GCAGAAACACAGAGGAAGTGG

GCTCTTTCCAGCTTGGCATATC

NALCN：

TAATGAGATAGGCACGAGTA

TGATGAAGTAGAAGTAGGAG

Single cell RT-qPCR

Following the retrograde labelling and VTA brain slice preparation, projection-specific VTA neurons were identified by infrared-differential interference contrast (IR-DIC) video microscopy and epifluorescence (Olympus, Japan) for detection of retrobeads (red) positive neurons. mRNA was reversely transcribed to cDNA by PrimeScript^{TM}II1st Strand cDNA Synthesis Kit (Takara-Clontech, Kyoto, Japan). The projection-specific neurons was aspirated into a pipette and then expelled into a PCR sterile tube containing 1 µl oligo-dT Primer and 1 µl dNTP mixture. The mixture was heated to 65°C for 5 min to denature the nucleic acids and then cooled on ice for 2 min. Reverse transcription from mRNA into cDNA was performed at 50°C for 50 min and then 85°C for 5 sec. cDNA was stored at -40°C. The targeted pre-amplification was done, with pairs of targeted primers for GAPDH, TH and TRPC6, to quantify multiple targets per cell. After adding the targeted primer pairs into each PCR tube, the synthesis conditions were as follows: 95 °C (5 min); 10 cycles of 95 °C (50 s), 58-62 °C (50 s), 72 °C (50 s); 72 °C (5 min). After targeted pre-amplification, Quantitative PCR (qPCR) is the final laboratory step of the scRT-qPCR workflow with the SYBR Green Master Mix (TaKaRa).

The targeted primers for pre-amplification (from 5′to 3′) as follows:

GAPDH:

AAATGGTGAAGGTCGGTGTGAACG（sense）

AGTGATGGCATGGACTGTGGTCAT （antisense）

TH:

AAGGTTCATTGGACGGCGG

ACATCGTCAGACACCCGAC

TRPC6:

AGCCTGTCTATTGAGGAAGAAC

AGCGAGAATGATTGGGGTCA

The primers for qPCR (from 5′ to 3′) as follows:

GAPDH:

GCAAATTCAACGGCACAGTCAAGG

TCTCGTGGTTCACACCCATCACAA

TH:

TCTCCTTGAGGGGTACAAAACC

ACCTCGAAGCGCACAAAGT

TRPC6:

TGCTAGAAGAGTGTCATTCCCT

CCTCCACAATCCGTACATAACC

Immunofluorescence

After intracardial perfusion with 4% paraformaldehyde (PFA) in 0.01 M PBS (pH 7.4).

The brains were post-fixed in 4% paraformaldehyde. 48 h later, the brain tissue was

placed in 30% sucrose solution (PBS preparation) to dehydrate. The brain tissue was

sectioned coronally, including the nucleus accumbens (NAc), medial prefrontal cortex

(mPFC), basolateral amygdala (BLA), and VTA, using a vibrating microtome. The

section thickness of NAc, mPFC and BLA was 80μm and the section thickness of

VTA was 40 μm. The sections were placed in PBS solution and stored in a

refrigerator at 4°C for storage.

The brain section was incubated in 0.3% triton/3% BSA for 1 h at room temperature

and then was blocked with 10% donkey serum at 37°C for 1 h. After that, the brain

section was incubated in the corresponding antibodies in PBS at 4°C for 24 h. The

section was washed three times (10 min) with PBS. Finally, the brain section was

incubated in the secondary antibodies for 2 h at 37°C. Images were obtained on a

Leica TCS SP5 confocal laser microscope (Leica, Germany) equipped with laser lines

for 405 mm, 488 mm, 561 mm and 647 mm illumination. Images were analyzed with

LAS-AF-Lite software (Leica, Germany).

Single-cell whole-transcriptome gene sequencing

After retrobeads injection, following brain slice preparation and 1 h recording, the

recorded and retrobeads-labelled neurons were aspirated into the patch pipette and

were then broken into the PCR tube containing 1 μl lysis buffer. For mRNA in

individual cells, mRNA was amplified and cDNA by SMARTer Ultra Low Input RNA

for Illumina Kit, which was qualified and reversely transcribed to cDNA by Qubit and

Agilent Bioanalyzer 2100 electrophoresis. After fragmentation of cDNA (300 bp) by ultrasound, sequencing libraries (end repair, addition of poly(A), and ligation of sequencing connectors) were built using the Ovation Ultralow Library System V2. After that, the constructed libraries were sequenced using Illumina HiseqXten (Sinotech Genomics Co., Ltd.).

Depression models

Chronic Mild Unpredictable Stress (CMUS) procedure

The CMUS procedures consisted of food and water deprivation (24 h), day/night inversion, damp bedding (12 h), cage tilt (12 h), no bedding (12 h), rat bedding (12 h), 4°C cold bath (5 min), restraint (2 h)and tail pinching (30 min). Mice were subjected to consecutive 35 days of CMUS with two stressors per day. Non-stressed controls were handled only for cage changes and behavioral tests.

Chronic Restraint Stress (CRS) procedure

The mice were immobilized in a special restraint device for 6 h per day from 9:00 to 15:00 for 21 days.

Behavior tests

The behavior tests were performed in the following order.

Sucrose preference test

Mice were single-housed and trained to drink from two drinking bottles with 1 bottle of tap water and 1 bottle of 1% sucrose water, for 48 hours, and the positions of the drinking bottles were exchanged every 12 h to exclude the interference of position preference. After the training, the mice were deprived of food and water for 12 h, and then the sucrose preference test was performed for 24 h. One bottle of tap water and one bottle of 1.0 % sucrose water were given again, and the positions of the drinking bottles were exchanged at 12 h. Finally, the consumption of tap water and sugar water at 24 h were recorded to calculate the sucrose solution preference rate. Sucrose preference rate = sucrose solution consumption/(tap water consumption + sucrose solution consumption).

Open field test, OFT

The mice were placed in a topless chamber (40 cm × 40 cm × 30 cm) with a camera

mounted on top of the chamber and connected to ANY-maze video tracking system on the computer to automatically track and record the activity of mice during the experiment and to obtain behavioral data. The bottom of the chamber was divided into 16 small square grids (10 cm × 10 cm) on the computer, and four square grids in the central area were defined as the central zone. During the 10 min test session, Total distance traveled and time spent in the center were recorded.

Elevated plus maze, EPM

The EPM apparatus consists of two closed arms (25 x 5 cm) across from each other and perpendicular to two open arms (25 x 5 cm) that are connected by a center platform (5 x 5 cm), with a camera in the center ceiling to automatically track and record the activity of mice. Mice were placed in the center platform facing a closed arm and allowed to freely explore the maze for 7 min, of which the first 2 min was the adaptation time. The time spent in open arms was analyzed for last 5 min after the adaptation.

Accelerating rotarod test

Locomotor activity was estimated by accelerating the rotarod test. Each group of animals was placed on an accelerating rotarod walking for 300 s from 4 to 40 rpm. After three days of training (three times per day), latency was automatically recorded from the beginning of the trial until the mouse falls off.

Forced swimming test, FST

Forced swimming test (FST) was performed in a glass cylinders, filled with the water depth about 15 cm at room temperature. The experiment was conducted for 5 min, of which the first 1 min was the adaptation time and the mice were allowed to move freely, and the immobility time was recorded for last 4 min after the adaptation.

Tail suspension test, TST

Mouse was suspended by taping the tail (1 cm from the tip of the tail) to the fixation device for 6 min. The cumulative immobility time within last 4 min was recorded, and the time when all limbs were immobile except for respiration was considered as immobility time.

Western Blot

The VTA protein was isolated by100 µl RIPA and 1 µl PMSF. The total protein for

each sample was transferred onto the polyvinylidene difluoride (PVDF) membranes

after SDS-polyacrylamide gel electrophoresis (SDS-PAGE), and blocked 2 h at room

temperature with 5% bovine serum albumin (KeyGen Biotechnology). Subsequently,

all these membranes were incubated overnight at 4 °C with the primary antibodies as

below: TRPC6, NALCN, and GAPDH. Following 2 h secondary antibodies

incubation, all the bands were detected. The relative expression was calculated based

on the internal control GAPDH.

Drugs and reagents

Drugs were bath applied at the following concentrations:

6-cyano-7-nitroquinoxaline-2,3-dione (CNQX; 10 µM; Sigma),

DL-2-amino-5-phosphonopentanoic acid (APV; 50 µM; Sigma) and gabazine (10 µM;

Sigma), CsCl (3 mM; sigma), ZD7288 (60 µm; Abcam), TTX (1 µM; MCE), $GdCl_3$

(100 µM; Sigma), L703,606 (10 µM; Sigma), 2-Aminoethyldiphenylborinate (2-APB,

100 µM; Sigma), Larixyl acetate (10 µM; Sigma), Ruthenium Red (60 µM; TCI).

Commercial antibodies used were: Anti-Tyrosine Hydroxylase Antibody (1:400,

Millipore, MAB318, RRID: AB_2201528, for Immunofluorescence), Anti-Dopamine

Transporter (N-terminal) antibody (1:400, sigma, D6944, RRID: AB_1840807, for

Immunofluorescence), Anti-NALCN (1:100, Thermo Fisher Scientific, MA5-27593,

RRID: AB_2735285, for Immunofluorescence), Anti-TRPC6 (1:100, Alomone,

ACC-017, RRID: AB_2040243, for Immunofluorescence), Anti-GAPDH (1:10000,

Santa Cruz, sc-137179,RRID: AB_2232048, for Western Blot), Anti-NALCN (1:100,

GeneTex, GTX54808, for Western Blot), Anti-TRPC6 (1:100, Cell Signaling

Technology, 16716, RRID:AB_2798768, for Western Blot).

Secondary antibodies: Donkey anti-Mouse IgG (H+L) Highly Cross-Adsorbed

Secondary Antibody (Alexa Fluor 488, Thermo Fisher Scientific, A-21202, 1:1000),

Donkey anti-Mouse IgG (H+L) Highly Cross-Adsorbed Secondary Antibody (Alexa

Fluor 546, Thermo Fisher Scientific, A10037, 1:1000), Donkey anti-Rabbit IgG (H+L)

Highly Cross-Adsorbed Secondary Antibody (Alexa Fluor 488, Thermo Fisher

Scientific, A21206, 1:1000), Donkey anti- Rabbit IgG (H+L) Highly Cross-Adsorbed

Secondary Antibody (Alexa Fluor 546, Thermo Fisher Scientific, A10040, 1:1000),

Donkey anti- Rabbit IgG (H+L) Highly Cross-Adsorbed Secondary Antibody (Alexa

Fluor 647, Thermo Fisher Scientific, A31573, 1:1000)

Quantification and statistical analysis

Software such as GraphPad Prism6, OriginPro 8.0 (Origin Lab) and Adobe Illustrator

CS6 were used for data analysis and image processing. All experimental data were

expressed as mean ± standard error (mean ± S.E.M.). When the data were normally

distributed, the difference between two groups was statistically analyzed by

two-sample T-test or paired-sample T-test; when the data were not normally

distributed, the difference between two groups was statistically analyzed by

Mann-Whitney U test or Wilcoxon matched-pairs signed rank test. $P < 0.05$ was

considered as a statistically significant difference between two groups. For the data

between multiple groups, when the data were normally distributed and there was no

significant variance in homogeneity, one-way ANOVA with Dunnett's multiple

comparisons test or two-way ANOVA with Sidak's multiple comparisons test was

used; when the data were not normally distributed, Kruslal-Wallis-H test with

Dunnett's multiple comparisons test and Student-Newman-Keuls test with Dunnett's

multiple comparisons test were used.

Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial

or financial relationships that could be construed as a potential conflict of interest.

Author Contributions

Hailin Zhang conceived, designed and supervised the experiments; Jing Wang

performed the experiments, acquired and analyzed the data and prepared the figures;

Dongmei Zhang and Min Su performed immunofluorescence of the brain slices and

performed some preliminary electrophysiological experiments; Yuqi Sang and Chaoyi

Li performed part of single-cell PCR; Yongxue Zhang performed part of statistical

analysis; Ludi Zhang and Chenxu Niu took care of mice; Hailin Zhang and Jing Wang

wrote the manuscript; Hailin Zhang and Xiaona Du prepared the final version of the

manuscript.

Funding

This work was supported by the National Natural Science Foundation of China (81871075, 82071533) grants to HZ; National Natural Science Foundation of China (81870872) grants to XD; Science Fund for Creative Research Groups of Natural Science Foundation of Hebei Province (no. H2020206474);Basic Research Fund for Provincial Universities (JCYJ2021010), Natural Science Foundation of Hebei Province (H2023423065) and Scientific research project of Hebei administration of traditional Chinese Medicine (2024091) grants to WJ.

Acknowledgments

We would like to thank colleagues at the Core Facilities and Centers of Hebei Medical University for the technical assistance.

Data Availability Statement

The data that support the findings of this study are available from the corresponding author upon reasonable request.

References

1. Bromberg-Martin, E.S., M. Matsumoto, and O. Hikosaka, *Dopamine in motivational control: rewarding, aversive, and alerting.* Neuron, 2010. **68**(5): p. 815-34.

2. Lammel, S., B.K. Lim, and R.C. Malenka, *Reward and aversion in a heterogeneous midbrain dopamine system.* Neuropharmacology, 2014. **76 Pt B**(0 0): p. 351-9.

3. Van den Heuvel, D.M. and R.J. Pasterkamp, *Getting connected in the dopamine system.* Prog Neurobiol, 2008. **85**(1): p. 75-93.

4. Chaudhury, D., et al., *Rapid regulation of depression-related behaviours by control of midbrain dopamine neurons.* Nature, 2013. **493**(7433): p. 532-6.

5. Kabanova, A., et al., *Function and developmental origin of a mesocortical inhibitory circuit.* Nat Neurosci, 2015. **18**(6): p. 872-82.

6. Badrinarayan, A., et al., *Aversive stimuli differentially modulate real-time dopamine transmission dynamics within the nucleus accumbens core and shell.* J Neurosci, 2012. **32**(45): p. 15779-90.

7. Friedman, A.K., et al., *Enhancing depression mechanisms in midbrain dopamine neurons achieves homeostatic resilience.* Science, 2014. **344**(6181): p. 313-9.

8. Krishnan, V., et al., *Molecular adaptations underlying susceptibility and resistance to social defeat in brain reward regions.* Cell, 2007. **131**(2): p. 391-404.

9. Grace, A.A., et al., *Regulation of firing of dopaminergic neurons and control of goal-directed behaviors.* Trends Neurosci, 2007. **30**(5): p. 220-7.

10. Johnson, S.W. and R.A. North, *Two types of neurone in the rat ventral tegmental area and their synaptic inputs.* J Physiol, 1992. **450**: p. 455-68.

11. Zhong, P., et al., *HCN2 channels in the ventral tegmental area regulate behavioral responses to chronic stress.* Elife, 2018. **7**.

12.   Bean, B.P., *The action potential in mammalian central neurons.* Nat Rev Neurosci, 2007. **8**(6):
p. 451-65.

13.   Guzman, J.N., et al., *Robust pacemaking in substantia nigra dopaminergic neurons.* J Neurosci,
2009. **29**(35): p. 11011-9.

14.   Puopolo, M., E. Raviola, and B.P. Bean, *Roles of subthreshold calcium current and sodium*
current in spontaneous firing of mouse midbrain dopamine neurons. J Neurosci, 2007. **27**(3):
p. 645-56.

15.   Khaliq, Z.M. and B.P. Bean, *Pacemaking in dopaminergic ventral tegmental area neurons:*
depolarizing drive from background and voltage-dependent sodium conductances. J Neurosci,
2010. **30**(21): p. 7401-13.

16.   Beckstead, M.J., et al., *Vesicular dopamine release elicits an inhibitory postsynaptic current in*
midbrain dopamine neurons. Neuron, 2004. **42**(6): p. 939-46.

17.   Ford, C.P., M.J. Beckstead, and J.T. Williams, *Kappa opioid inhibition of somatodendritic*
dopamine inhibitory postsynaptic currents. J Neurophysiol, 2007. **97**(1): p. 883-91.

18.   Philippart, F., et al., *Differential Somatic Ca2+ Channel Profile in Midbrain Dopaminergic*
Neurons. J Neurosci, 2016. **36**(27): p. 7234-45.

19.   Dufour, M.A., A. Woodhouse, and J.M. Goaillard, *Somatodendritic ion channel expression in*
substantia nigra pars compacta dopaminergic neurons across postnatal development. J
Neurosci Res, 2014. **92**(8): p. 981-99.

20.   Kimm, T., Z.M. Khaliq, and B.P. Bean, *Differential Regulation of Action Potential Shape and*
Burst-Frequency Firing by BK and Kv2 Channels in Substantia Nigra Dopaminergic Neurons. J
Neurosci, 2015. **35**(50): p. 16404-17.

21.   Li, L., et al., *Selective targeting of M-type potassium K(v) 7.4 channels demonstrates their key*
role in the regulation of dopaminergic neuronal excitability and depression-like behaviour. Br J
Pharmacol, 2017. **174**(23): p. 4277-4294.

22.   Tarfa, R.A., R.C. Evans, and Z.M. Khaliq, *Enhanced Sensitivity to Hyperpolarizing Inhibition in*
Mesoaccumbal Relative to Nigrostriatal Dopamine Neuron Subpopulations. J Neurosci, 2017.
37(12): p. 3311-3330.

23.   Seutin, V., et al., *Evidence for a modulatory role of Ih on the firing of a subgroup of midbrain*
dopamine neurons. Neuroreport, 2001. **12**(2): p. 255-8.

24.   Neuhoff, H., et al., *I(h) channels contribute to the different functional properties of identified*
dopaminergic subpopulations in the midbrain. J Neurosci, 2002. **22**(4): p. 1290-302.

25.   Krashia, P., et al., *On the properties of identified dopaminergic neurons in the mouse*
substantia nigra and ventral tegmental area. Eur J Neurosci, 2017. **45**(1): p. 92-105.

26.   Shi, Y., et al., *Nalcn Is a "Leak" Sodium Channel That Regulates Excitability of Brainstem*
Chemosensory Neurons and Breathing. J Neurosci, 2016. **36**(31): p. 8174-87.

27.   Lu, B., et al., *Peptide neurotransmitters activate a cation channel complex of NALCN and*
UNC-80. Nature, 2009. **457**(7230): p. 741-4.

28.   Philippart, F. and Z.M. Khaliq, *G(i/o) protein-coupled receptors in dopamine neurons inhibit*
the sodium leak channel NALCN. Elife, 2018. **7**.

29.   Lutas, A., et al., *The leak channel NALCN controls tonic firing and glycolytic sensitivity of*
substantia nigra pars reticulata neurons. Elife, 2016. **5**.

30.   Ford, N.C., D. Ren, and M.L. Baccei, *NALCN channels enhance the intrinsic excitability of spinal*
projection neurons. Pain, 2018. **159**(9): p. 1719-1730.

31.   Um, K.B., et al., *TRPC3 and NALCN channels drive pacemaking in substantia nigra*
1041   *dopaminergic neurons.* Elife, 2021. **10**.

1042   32.   Kim, S.H., et al., *Nonselective cation channels are essential for maintaining intracellular Ca2+*
1043   *levels and spontaneous firing activity in the midbrain dopamine neurons.* Pflugers Arch, 2007.
1044   **455**(2): p. 309-21.

1045   33.   Zhou, F.W., S.G. Matta, and F.M. Zhou, *Constitutively active TRPC3 channels regulate basal*
1046   *ganglia output neurons.* J Neurosci, 2008. **28**(2): p. 473-82.

1047   34.   Hof, T., et al., *Transient receptor potential channels in cardiac health and disease.* Nat Rev
1048   Cardiol, 2019. **16**(6): p. 344-360.

1049   35.   Klipec, W.D., et al., *Loss of the trpc4 gene is associated with a reduction in cocaine*
1050   *self-administration and reduced spontaneous ventral tegmental area dopamine neuronal*
1051   *activity, without deficits in learning for natural rewards.* Behav Brain Res, 2016. **306**: p.
1052   117-27.

1053   36.   German, D.C. and K.F. Manaye, *Midbrain dopaminergic neurons (nuclei A8, A9, and A10):*
1054   *three-dimensional reconstruction in the rat.* J Comp Neurol, 1993. **331**(3): p. 297-309.

1055   37.   Nair-Roberts, R.G., et al., *Stereological estimates of dopaminergic, GABAergic and*
1056   *glutamatergic neurons in the ventral tegmental area, substantia nigra and retrorubral field in*
1057   *the rat.* Neuroscience, 2008. **152**(4): p. 1024-31.

1058   38.   Cadwell, C.R., et al., *Electrophysiological, transcriptomic and morphologic profiling of single*
1059   *neurons using Patch-seq.* Nat Biotechnol, 2016. **34**(2): p. 199-203.

1060   39.   Lammel, S., et al., *Unique properties of mesoprefrontal neurons within a dual*
1061   *mesocorticolimbic dopamine system.* Neuron, 2008. **57**(5): p. 760-73.

1062   40.   Kawano, M., et al., *Particular subpopulations of midbrain and hypothalamic dopamine*
1063   *neurons express vesicular glutamate transporter 2 in the rat brain.* J Comp Neurol, 2006.
1064   **498**(5): p. 581-92.

1065   41.   Stamatakis, A.M., et al., *A unique population of ventral tegmental area neurons inhibits the*
1066   *lateral habenula to promote reward.* Neuron, 2013. **80**(4): p. 1039-53.

1067   42.   Lu, B., et al., *The neuronal channel NALCN contributes resting sodium permeability and is*
1068   *required for normal respiratory rhythm.* Cell, 2007. **129**(2): p. 371-83.

1069   43.   Kusch, J., et al., *Interdependence of receptor activation and ligand binding in HCN2*
1070   *pacemaker channels.* Neuron, 2010. **67**(1): p. 75-85.

1071   44.   Mistrík, P., et al., *The murine HCN3 gene encodes a hyperpolarization-activated cation*
1072   *channel with slow kinetics and unique response to cyclic nucleotides.* J Biol Chem, 2005.
1073   **280**(29): p. 27056-61.

1074   45.   Clapham, D.E., L.W. Runnels, and C. Strübing, *The TRP ion channel family.* Nat Rev Neurosci,
1075   2001. **2**(6): p. 387-96.

1076   46.   Urban, N., et al., *Identification and Validation of Larixyl Acetate as a Potent TRPC6 Inhibitor.*
1077   Mol Pharmacol, 2016. **89**(1): p. 197-213.

1078   47.   Strübing, C., et al., *Formation of novel TRPC channels by complex subunit interactions in*
1079   *embryonic brain.* J Biol Chem, 2003. **278**(40): p. 39014-9.

1080   48.   Goel, M., W.G. Sinkins, and W.P. Schilling, *Selective association of TRPC channel subunits in rat*
1081   *brain synaptosomes.* J Biol Chem, 2002. **277**(50): p. 48303-10.

1082   49.   Tye, K.M., et al., *Dopamine neurons modulate neural encoding and expression of*
1083   *depression-related behaviour.* Nature, 2013. **493**(7433): p. 537-541.

50. Chang, C.H. and A.A. Grace, *Amygdala-ventral pallidum pathway decreases dopamine activity*
after chronic mild stress in rats. Biol Psychiatry, 2014. **76**(3): p. 223-30.

51. Moreines, J.L., Z.L. Owrutsky, and A.A. Grace, *Involvement of Infralimbic Prefrontal Cortex but*
not Lateral Habenula in Dopamine Attenuation After Chronic Mild Stress.
Neuropsychopharmacology, 2017. **42**(4): p. 904-913.

52. Willner, P., *The chronic mild stress (CMS) model of depression: History, evaluation and usage.*
Neurobiol Stress, 2017. **6**: p. 78-93.

53. Liu, D., et al., *Brain-derived neurotrophic factor-mediated projection-specific regulation of*
depressive-like and nociceptive behaviors in the mesolimbic reward circuitry. Pain, 2018.
159(1): p. 175.

54. Kang, Y. and S.T. Kitai, *A whole cell patch-clamp study on the pacemaker potential in*
dopaminergic neurons of rat substantia nigra compacta. Neurosci Res, 1993. **18**(3): p. 209-21.

55. Lu, B., et al., *Extracellular calcium controls background current and neuronal excitability via*
an UNC79-UNC80-NALCN cation channel complex. Neuron, 2010. **68**(3): p. 488-99.

56. Benarroch, E.E., *HCN channels: function and clinical implications.* Neurology, 2013. **80**(3): p.
304-10.

57. Mercuri, N.B., et al., *Properties of the hyperpolarization-activated cation current Ih in rat*
midbrain dopaminergic neurons. Eur J Neurosci, 1995. **7**(3): p. 462-9.

58. Okamoto, T., M.T. Harnett, and H. Morikawa, *Hyperpolarization-activated cation current (Ih) is*
an ethanol target in midbrain dopamine neurons of mice. J Neurophysiol, 2006. **95**(2): p.
619-26.

59. Chan, C.S., et al., *'Rejuvenation' protects neurons in mouse models of Parkinson's disease.*
Nature, 2007. **447**(7148): p. 1081-6.

60. Lee, J.H., L.L. Cribbs, and E. Perez-Reyes, *Cloning of a novel four repeat protein related to*
voltage-gated sodium and calcium channels. FEBS Lett, 1999. **445**(2-3): p. 231-6.

61. Lu, T.Z. and Z.P. Feng, *NALCN: a regulator of pacemaker activity.* Mol Neurobiol, 2012. **45**(3): p.
415-23.

62. Pochwat, B., et al., *Hyperforin Potentiates Antidepressant-Like Activity of Lanicemine in Mice.*
Front Mol Neurosci, 2018. **11**: p. 456.

63. Liu, Y., et al., *The change of spatial cognition ability in depression rat model and the possible*
association with down-regulated protein expression of TRPC6. Behav Brain Res, 2015. **294**: p.
186-93.

64. Cao, J.L., et al., *Mesolimbic dopamine neurons in the brain reward circuit mediate*
susceptibility to social defeat and antidepressant action. J Neurosci, 2010. **30**(49): p. 16453-8.

65. Eder, P., et al., *Na(+) entry and modulation of Na(+)/Ca(2+) exchange as a key mechanism of*
TRPC signaling. Pflugers Arch, 2005. **451**(1): p. 99-104.

66. Hahn, S., et al., *Proximal dendritic localization of NALCN channels underlies tonic and burst*
firing in nigral dopaminergic neurons. J Physiol, 2023. **601**(1): p. 171-193.

67. Inoue, R. and Y. Mori, *Molecular candidates for capacitative and non-capacitative Ca2+ entry*
in smooth muscle. Novartis Found Symp, 2002. **246**: p. 81-90; discussion 221-7.

68. Nishida, M., et al., *Phosphorylation of TRPC6 channels at Thr69 is required for*
anti-hypertrophic effects of phosphodiesterase 5 inhibition. J Biol Chem, 2010. **285**(17): p.
13244-53.

69. Nishioka, K., et al., *Cilostazol suppresses angiotensin II-induced vasoconstriction via protein*

kinase A-mediated phosphorylation of the transient receptor potential canonical 6 channel.
Arterioscler Thromb Vasc Biol, 2011. **31**(10): p. 2278-86.

70.   Su, M., et al., *Kv7.4 Channel Contribute to Projection-Specific Auto-Inhibition of Dopamine*
Neurons in the Ventral Tegmental Area. Front Cell Neurosci, 2019. **13**: p. 557.

71.   McCall, N.M., E. Marron Fernandez de Velasco, and K. Wickman, *GIRK Channel Activity in*
Dopamine Neurons of the Ventral Tegmental Area Bidirectionally Regulates Behavioral
Sensitivity to Cocaine. J Neurosci, 2019. **39**(19): p. 3600-3610.

72.   Qu, N., et al., *A POMC-originated circuit regulates stress-induced hypophagia, depression,*
and anhedonia. Mol Psychiatry, 2020. **25**(5): p. 1006-1021.

73.   Butterweck, V., *Mechanism of action of St John's wort in depression : what is known?* CNS
Drugs, 2003. **17**(8): p. 539-62.

74.   Burlet, S., C.J. Tyler, and C.S. Leonard, *Direct and indirect excitation of laterodorsal tegmental*
neurons by Hypocretin/Orexin peptides: implications for wakefulness and narcolepsy. J
Neurosci, 2002. **22**(7): p. 2862-72.

75.   Jan-Pieter and Konsman, *The mouse brain in stereotaxic coordinates: Second Edition (Deluxe)*
By Paxinos G. and Franklin, K.B.J., Academic Press, New York, 2001, ISBN 0-12-547637-X.
Psychoneuroendocrinology, 2003.

1145

1146

Figure legends

Fig.1 The effects of extracellular Ca^{2+} and Na^+ on firing activity of the male VTA DA neurons.

A. Identification of the VTA DA neurons. i: confocal images showing the anatomical distribution of DA neurons in the VTA and SNc; the DA neurons were identified to be DAT-immunofluorescence positive (scale bar, 100 μm). ii: left panel: single-cell PCR results; four cells showed the presence of genes indicated. right panel: Percentage of neuronal markers (DA neuron: TH, DAT, D_2R, Girk2; Glu neuron: Vglut2 and GABA neuron: GAD1) positive neurons in the VTA neurons (GAPDH positive). n = 63 cells, N = 10. **B**. The effect of replacing extracellular Ca^{2+} with Mg^{2+} on firing frequency of the VTA DA neurons. Example time-course (i) and traces (ii) of the spontaneous firing before and after replacement of extracellular Ca^{2+} (2 mM) by an equimolar amount of Mg^{2+}.The inset on the top of ii shows a map of a coronal midbrain slice indicating the location of neurons that were recorded and subsequently identified as DA neurons which were DAT positive with single cell-PCR (red dots). iii: Summarized data for ii. Firing of the VTA DA neurons were recorded using loose cell-attached patch from a brain slice of the VTA (n = 16, N = 8). Paired-sample T test, t = 4.698, P = 0.0003. **C**. The effect of replacing extracellular Ca^{2+} with Mg^{2+} on the resting membrane potential (RMP) of the VTA DA neurons, in the presence of 1 μM TTX. Example time-course (i) and summarized data (ii) for the resting membrane potential before and after replacement of extracellular Ca^{2+} (2 mM) by an equimolar amount of Mg^{2+} (n = 14, N = 8). The inset on the top of ii shows a map of a coronal midbrain slice indicating the location of neurons that were recorded and subsequently identified as DA neurons which were DAT positive with single cell-PCR (red dots). Wilcoxon matched-pairs signed rank test, W = 66.00, P = 0.037. **D.** i: Replacement of external Na^+ by equimolar NMDG resulted in hyperpolarization of the resting membrane potential of the VTA DA neurons, in the presence of 1 μM TTX. ii: Summarized data for i (n = 8, N = 5). The inset on the top of ii shows a map of a coronal midbrain slice indicating the location of neurons that were recorded and subsequently identified as DA neurons which were DAT positive with single cell-PCR (red dots). Paired-sample T test, t = 7.803, P = 0.0001. * P < 0.05, *** P < 0.001. n is number of neurons recorded and N is number of mice used.

Fig. 2 Gene expression profile of non-selective cation channels (NSCCs) in the male VTA DA neurons.

Single-cell RNA-Seq was performed on the VTA neurons projecting to five different brain regions of mice (NAc c, NAc ms, NAc ls, BLA, mPFC). **A**. Gene expression profile of markers for neuron subtypes from 45 VTA neurons; the neuron subtypes included: dopaminergic (Th, Ddc, DAT), GABAergic (Gad1, Gad2, VGAT) and glutamatergic (VGluT2, VGluT1, VGluT3), which was arranged in different rows indicated in the right labels and in the left-colored vertical lines; projection targets of these neurons were indicated at the top by the colored lines and labels. Relative expression levels of these genes were indicated by the dark-blue color intensity which was transformed from the log2 values of the number of transcripts per million (FPKM) plus 1. **B**. Relative gene expression levels of TRP channels (i) and other NSCCs (ii) in 45 individual VTA DA neurons and the population average (mean, right columns). Each column of the individual neurons in (i) and (ii) corresponded to the columns in A. (iii) Bar graph of the mean log2 (FPKM +1) for the top 8 NSCCs in descending order. Error bars indicate SEM. **C.** Gene expression profile of aforementioned top 8 NSCCs from 45 VTA neurons.

Fig. 3 NALCN contributes to subthreshold depolarization and spontaneous firing of the male VTA DA neurons.

A. Confocal images showing co-expression of NALCN (green) and DAT (red) representing DA neurons (scale bar, 10 μm). **B**. i: Single-cell PCR from the VTA DA neurons with the expression of NALCN. ii: Percentage of NALCN positive neurons from 15 DA neurons (with expression of DAT). **C**. Example time course (i) and summarized data (ii) showed that NALCN channel blockers (GdCl$_3$) significantly hyperpolarized the resting membrane potential (RMP) (n = 6, N = 6). The inset on the top of Cii shows a map of a coronal midbrain slice indicating the location of neurons that were recorded and subsequently identified as DA neurons which were DAT positive with single cell-PCR (red dots). Wilcoxon matched-pairs signed rank test, W = -21.00, *P* = 0.0313. **D**. Example time-course and traces (i) and summarized data (ii) of the effect of GdCl$_3$ on spontaneous firing frequency in DA neurons (n = 7, N = 3). The inset on the top of Dii shows a map of a coronal midbrain slice indicating the location of neurons that were recorded and subsequently identified as DA neurons

which were DAT positive with single cell-PCR (red dots). Paired-sample T test, t = 5.742, P = 0.0012. **E**. Example time course (i) and summarized data (ii) showed that NALCN channel blocker (L703,606) significantly hyperpolarized the resting membrane potential (RMP) (n = 6, N = 4). The inset on the top of Eii shows a map of a coronal midbrain slice indicating the location of neurons that were recorded and subsequently identified as DA neurons which were DAT positive with single cell-PCR (red dots). Paired-sample T test, t = 14.95, P = 0.0001. **F**. Example time-course and traces (i) and summarized data (ii) of the effect of L703,606 on spontaneous firing frequency in DA neurons (n = 8, N = 4). The inset on the top of Fii shows a map of a coronal midbrain slice indicating the location of neurons that were recorded and subsequently identified as DA neurons which were DAT positive with single cell-PCR (red dots). Paired-sample T test, t = 6.236, P = 0.0004. **G**. shRNA against NALCN carried by AAV virus (i) (NALCN-shRNA) was injected into the VTA of mice, the mRNA level in the shRNA-NALCN transfected VTA (NALCN-KD) and the scramble shRNA transfected VTA (Control) was analyzed using qPCR (ii) (N = 6). Two-sample T test, t = 3.742, P = 0.0038. **H**. Confocal images showing the expression of AAV9-NALCN-shRNA-GFP (green) in the VTA DA neurons (DAT, red) (scale bar, 10 μm). **I**. Loose cell-attached current clamp recordings of the spontaneous firing of the VTA DA neurons transfected with either NALCN-shRNA (ii, n = 12, N = 3) or scramble-shRNA (Con, i, n = 10, N = 3) (both GFP and DAT positive). Examples of firing traces in the middle of i and ii and summarized data in the bottom of i were shown. The inset on the top of i and ii shows a map of a coronal midbrain slice indicating the location of GFP$^+$ neurons that were recorded and subsequently identified as DA neurons which were DAT positive with single cell-PCR (i: Con, grey dots; ii: NALCN-KD, red dots). Mann-Whitney U test, U = 0, P = 0.0001. **J**. Whole-cell current clamp recordings of the resting membrane potential (RMP) of the VTA DA neurons transfected with either NALCN-shRNA (NALCN-KD, ii, n = 18, N = 5) or scramble-shRNA (Con, i, n = 15, N = 5) (both GFP and DAT positive). Examples of RMP in the middle of i and ii and summarized data in the bottom of i were shown. The inset on the top of i and ii shows a map of a coronal midbrain slice indicating the location of GFP$^+$ neurons that were recorded and subsequently identified as DA neurons which were DAT positive with single cell-PCR (i: Con, grey dots; ii: NALCN-KD, red dots). Mann-Whitney U test, U = 0, P = 0.0001. * P < 0.05,

1246   ** $P <0.01$, *** $P < 0.001$. n is number of neurons recorded and N is number of mice

1247   used.

1248

Fig. 4 TRP channels contribute to subthreshold depolarization and spontaneous firing

of the male VTA DA neurons.

A. The expression of TRP channels in VTA DA neurons. i: Single-cell PCR from 5

VTA cells (C1-C5). ii: Percentage of TRP channels (C3, C6 and V2) positive neurons

in the VTA DA neurons (DAT positive). **B** and **C**. Example time-course (i), traces (ii)

and the summarized data (iii) for the effect of a nonselective cation channel blocker

2-APB (100 μM, n = 12, N = 5) (B) and a potent TRPC6 inhibitor LA (10 μM, n = 7,

N = 5) (C) on the spontaneous firing frequency in the VTA DA neurons. The inset on

the top of Bii and Cii shows a map of a coronal midbrain slice indicating the location

of neurons that were recorded and subsequently identified as DA neurons which were

DAT positive with single cell-PCR (red dots). B: Wilcoxon matched-pairs signed rank

test, W = -78.00, $P = 0.0005$. C: Paired-sample T test, t = 4.089, $P = 0.0064$. **D** and **E**.

Example time-course (i) and the summarized data (ii) for the effect of a nonselective

cation channel blocker 2-APB (n = 6, N = 6) (D) and a potent TRPC6 inhibitor LA (n

= 5, N = 4) (E) on the resting membrane potential (RMP) of the VTA DA neurons.

Paired-sample T test, D: t = 7.976, $P = 0.0005$, E: t = 5.610, $P = 0.005$.   ** $P < 0.01$,

*** $P < 0.001$. n is number of neurons recorded and N is number of mice used.

1266

Fig. 5 TRPC6 contributes to subthreshold depolarization and spontaneous firing of the

male VTA DA neurons.

A. The normalized expression profile of TRPC6 in NAc c- and mPFC-projecting VTA

DA neurons was verified using single cell-qPCR (NAc c: n = 28, N = 7; mPFC: n =

25, N = 8, both TH positive), Mann-Whitney U test, U = 174, $P = 0.0014$. **B**.

Confocal images showing co-expression of TRPC6 (green) and TH (red) representing

DA neurons (scale bar, 10 μm). **C**. The efficiency of shRNA knockdown of TRPC6 in

the VTA verified by qPCR. shRNA against TRPC6 carried by AAV virus (i)

(TRPC6-shRNA) was injected into the VTA of mice, the mRNA level in the

shRNA-TRPC6 transfected VTA (TRPC6-KD, N = 5) and the scramble shRNA

transfected VTA (Con, N = 5) was analyzed using qPCR (ii). Two-sample T test,

t=4.184, P=0.0031. **D**. Immunofluorescence labelling showing the expression of

AAV9-shRNA (TRPC6)-GFP (green) and DAT (red) in the VTA (scale bar, 10 μm). **E**.

Loose cell-attached current clamp recordings of the spontaneous firing of the VTA

DA neurons from the mice transfected with either TRPC6-shRNA (TRPC6-KD, ii, n =

9, N = 3) or scramble-shRNA (Con, i, n = 9, N = 3) (both GFP and DAT positive).

Examples of firing traces in the middle of i and ii and summarized data in the bottom

of ii were shown. The inset on the top of i and ii shows a map of a coronal midbrain

slice indicating the location of GFP$^+$ neurons that were recorded and subsequently

identified as DA neurons which were DAT positive with single cell-PCR (i: Con, grey

dots; ii: TRPC6-KD, red dots). Two-sample T test, t = 3.005, P = 0.0084. **F**. Example

time-course (i) and summarized data (ii) showed that shRNA knockdown of TRPC6

in the VTA decreased the 2-APB-inhibited firing responses of the VTA DA neurons.

The inset on the top of ii shows a map of a coronal midbrain slice indicating the

location of GFP$^+$ neurons that were recorded and subsequently identified as DA

neurons which were DAT positive with single cell-PCR (TRPC6-KD, red dots). (n = 7,

N = 5, both GFP and DAT positive), Wilcoxon matched-pairs signed rank test, W =

-18.00, P = 0.1563. **G**. Whole-cell current clamp recordings of the resting membrane

potential (RMP) of the VTA DA neurons transfected with either TRPC6-shRNA

(TRPC6-KD, ii, n = 10, N = 6) or scramble-shRNA (Con, i, n = 10, N = 6) (both GFP

and DAT positive). Examples of RMP in the middle of i and ii and summarized data

in the bottom of i were shown. The inset on the top of i and ii shows a map of a

coronal midbrain slice indicating the location of GFP+ neurons that were recorded

and subsequently identified as DA neurons which were DAT positive with single

cell-PCR (i: Con, grey dots; ii: TRPC6-KD, red dots). Mann-Whitney U test, U = 8, P

= 0.0007. **H**. Example time-course (i) and summarized data (ii) showed that shRNA

knockdown of TRPC6 in the VTA decreased the LA-inhibited firing responses of the

VTA DA neurons. The inset on the top of ii shows a map of a coronal midbrain slice

indicating the location of GFP$^+$ neurons that were recorded and subsequently

identified as DA neurons which were DAT positive with single cell-PCR (TRPC6-KD,

red dots). (n = 8, N = 5, both GFP and DAT positive), Paired-sample T test, t = 0.000,

P = 0.9999. n.s. P > 0.05, ** P <0.01, *** P < 0.001. n is number of neurons recorded

and N is number of mice used.

1310

1311 **Fig. 6** CMUS depression male mice have a decreased firing activity and

down-regulated TRPC6 expression in the VTA DA neurons.

A. Experimental procedure timeline. **B**. i: Sucrose preference test for CMUS mice. Two-sample T test, t = 5.810, P = 0.0001, compared with control mice. ii: Tail suspension test for CMUS mice, Two-sample T test, t = 4.689, P = 0.0005, compared with control mice. **C**. Loose cell-attached current clamp recordings of the spontaneous firing of the VTA DA neurons from the CMUS (n = 28, N = 5, DAT positive) and the control (n = 18, N = 5, DAT positive) mice. The inset on the top of i and ii shows a map of a coronal midbrain slice indicating the location of neurons that were recorded and subsequently identified as DA neurons which were DAT positive with single cell-PCR (i: Con, grey dots; ii: CMUS, red dots). Examples of firing traces in the bottom of i as well as in the bottom left of ii and summarized data in the bottom right of ii were shown. Mann-Whitney U test, U = 118, P = 0.0020. **D**. Representative Western blot assay (i) and summarized data (ii) showing the expression of TRPC6 and GAPDH in the VTA of the control (Con, N = 6) and the CMUS (N = 6) mice. Two-sample T test, t = 4.848, P = 0.0007. **E**. i and ii: Example traces (up) and time-course (bottom) of firing frequency of mPFC-projecting VTA DA neurons, effect of 2-APB. (i) Con, (ii) CMUS. iii. Summarized effect of 2-APB in the control (n = 12 cells, N = 5, DAT positive) and CMUS (n = 15 cells, N=5, DAT positive) mice. Two-way ANOVA with Sidak's multiple comparisons test, Con vs. CMUS: $F_{(1, 25)}$ = 3.411, Con-ACSF vs. CMUS-ACSF, P = 0.0001; Con-2-APB vs. CMUS-2-APB, P = 0.0164. ACSF vs. 2-APB: $F_{(1, 25)}$ = 92.43, Con-ACSF vs. Con-2-APB, P = 0.0001; CMUS-ACSF vs. CMUS-2-APB, P = 0.2259. iv. The inhibition rate (%) on firing rate by 2-APB in the control group (n = 12, N=5) and the CMUS group (n = 15, N = 5). Mann-Whitney U test, U = 1, P = 0.0001. The inset on the top of iii and iv shows a map of a coronal midbrain slice indicating the location of mPFC-projecting neurons that were recorded and subsequently identified as DA neurons which were DAT positive with single cell-PCR (iii: Con, grey dots; iv: CMUS, red dots). **F**. i and ii: Example traces (up) and time-course (bottom) of firing frequency of mPFC-projecting VTA DA neurons, effects of LA.(i) Con, (ii) CMUS. iii. Summarized effect of LA on the control (n = 17 cells, N = 8, DAT positive) and CMUS (n = 19 cells, N = 8, DAT positive) mice. Kruslal-Wallis-H test with Dunnett's multiple comparisons test, H = 53.08, Con-ACSF vs. Con-LA: P = 0.0001, Con-ACSF vs. CMUS-ACSF: P = 0.0023, CON-LA vs. CMUS-LA: P = 0.0187, CMUS-ACSF vs. CMUS-LA: P > 0.9999. iv.

The inhibition rate (%) on firing rate by LA in the control group (n = 17, N = 8) and the CMUS group (n = 19, N = 8). Mann-Whitney U test, U = 0, P = 0.0001. The inset on the top of iii and iv shows a map of a coronal midbrain slice indicating the location of mPFC-projecting neurons that were recorded and subsequently identified as DA neurons which were DAT positive with single cell-PCR (iii: Con, grey dots; iv: CMUS, red dots). n.s. $P > 0.05$, * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$. n is number of neurons recorded and N is the number of mice used.

Fig. 7 Selective knockdown of TRPC6 in the VTA DA neurons confers the male mice with depression-like and anxiety-like behaviors which are reversed by overexpression of TRPC6.

A. shRNA against TRPC6 carried by AAV-loxp virus (TRPC6-shRNA) was injected into the VTA of the DAT-Cre mice. **B**. The mRNA level in the shRNA-TRPC6 transfected VTA (TRPC6-cKD) and the scramble shRNA transfected VTA (Con) was analyzed using qPCR (Con: N = 4; TRPC6-cKD: N = 6). Mann-Whitney U test, U = 0, P = 0.0159. **C**. Immunofluorescence labelling showing the expression of AAV9-hSyn-DIO-shRNA (TRPC6)-RFP (red) and DAT (green) in the VTA of the DAT-Cre mice (scale bar, 10 µm). **D**. The effects of Cre-induced conditional knockdown of TRPC6 in the VTA DA neurons on the behaviors of mice in the sucrose preference test (i), the tail suspension test (ii), and the elevated plus-maze test (iii, iv) (Con: N = 9; TRPC6-cKD: N = 13). Two-sample T test, Sucrose preference test: t = 4.663, P = 0.0001; Tail suspension test: t = 7.143, P = 0.0001; Elevated plus-maze test: open arm t = 10.36, P = 0.0001, closed arm t = 11.18, P = 0.0001.**E**. The effects of TRPC6 over-expression. AAV9-DIO-TRPC6-GFP (TRPC6-cOE) was injected into the VTA of the DAT-Cre mice, 7 days after TRPC6 knocking down (AAV9-hSyn-DIO-shRNA (TRPC6)-RFP injection). **F**. The protein level TRPC6 of VTA (Con: the scramble shRNA for TRPC6, N = 7; C6-cKO: TRPC6-cKD, N = 8; c6-cKO+cOE: TRPC6-cKD+ TRPC6-cOE, N = 7). Kruskal-Wallis H test with Dunnett's multiple comparisons test, H = 14.75, Con *vs.* C6-cKO: P = 0.0016, C6-cKO *vs.* C6-cKO+cOE: P = 0.0062. **G**. Immunofluorescence labelling showing the expression of AAV9-hSyn-DIO-shRNA (TRPC6)-RFP (red), AAV9-DIO-TRPC6-GFP (green) and DAT (blue) in the VTA of the DAT-Cre mice (scale bar, 10 µm). **H**. The effects of TRPC6 over-expression on depression-like

behaviors of TRPC6-cKD mice: the sucrose preference test (i), the tail suspension test (ii), and the elevated plus-maze test (iii) (Con: N = 11; TRPC6-cKD: N = 11; TRPC6-cKD+TRPC6-cOE: N = 11). One-way ANOVA with Dunnett's multiple comparisons test, sucrose preference test: $F_{(2, 30)}$ = 37.22, Con *vs*. C6-cKO: P = 0.0001, C6-cKO *vs*. C6-cKO+cOE: P = 0.0001; tail suspension test: $F_{(2, 30)}$ = 20.10, Con *vs*. C6-cKO: P = 0.0146, C6-cKO *vs*. C6-cKO+cOE: P = 0.0001; elevated plus-maze test: $F_{(2, 30)}$ = 17.10, Con *vs*. C6-cKO: P = 0.0001, C6-cKO *vs*. C6-cKO+cOE: P = 0.0095. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.n is number of neurons recorded and N is the number of mice used.

Fig. 8 Selective knockdown of TRPC6 in the mPFC-projecting VTA DA neurons produces depression-like and anxiety-like behaviors.

A. shRNA against TRPC6 carried by retro-AAV-loxp virus (TRPC6-shRNA) was injected into the mPFC of the DAT-Cre male mice. **B**. The mRNA level in the shRNA-TRPC6 transfected VTA (TRPC6-rcKD) DA neurons and the scramble shRNA transfected VTA DA neurons (Con) was analyzed using single-cell quantitative PCR (scqPCR) (Con: n = 8, N = 4; TRPC6-rcKD: n = 10, N = 4). Two-sample T test, t = 4.506, P = 0.0004. **C**. Immunofluorescence labelling showing the expression of AAV-retro-hSyn-DIO-mCherry-shRNA (TRPC6) (red) and DAT (green) in the VTA of the DAT-Cre mice (scale bar, 10 μm). **D**. The effects of Cre-induced conditional knockdown of TRPC6 in mPFC-projecting VTA DA neurons on the behaviors of mice: (i) the sucrose preference test, (ii) the tail suspension test, and (iii) the elevated plus-maze test (Con: N = 8; TRPC6-rcKD: N = 10). Two-sample T test, sucrose preference test: t = 5.215, P = 0.0001; tail suspension test: t = 2.993, P = 0.0097; elevated plus-maze test: open arm t = 5.802, P = 0.0001. ** $P < 0.01$, *** $P < 0.001$. n is number of neurons recorded and N is the number of mice used.

Fig. 2

1416

1417

1418

1420

1421

1422

1423

1424

Fig. 5

A

B

C i

ii

D

E i

ii

F i

ii

G i

ii

H i

ii

1431

1432

1433

1434

Fig. 7

1442

52