1   **Netrin-1 directs vascular patterning and maturity in the developing kidney**

3   Samuel Emery Honeycutt[1], Pierre-Emmanuel Yoann N'Guetta[1], Deanna Marie Hardesty[1], Yubin Xiong[1],

4   Shamus Luke Cooper[1], Lori Lynn O'Brien[1]*

6   [1]Department of Cell Biology and Physiology

7   University of North Carolina at Chapel Hill, Chapel Hill, NC 27599, USA

9   *Corresponding author: lori_obrien@med.unc.edu

Abstract

Blood filtering by the kidney requires the establishment of an intricate vascular system that

works to support body fluid and organ homeostasis. Despite these critical roles, little is known about

how vascular architecture is established during kidney development. More specifically, how signals from

the kidney influence vessel maturity and patterning remains poorly understood. Netrin-1 (*Ntn1*) is a

secreted ligand critical for vessel and neuronal guidance. Here, we demonstrate that *Ntn1* is expressed

by stromal progenitors in the developing kidney, and conditional deletion of *Ntn1* from Foxd1+ stromal

progenitors (*Foxd1$^{GC/+}$;Ntn1$^{fl/fl}$*) results in hypoplastic kidneys that display extended nephrogenesis.

Despite expression of the netrin-1 receptor *Unc5c* in the adjacent nephron progenitor niche, *Unc5c*

knockout kidneys develop normally. The netrin-1 receptor *Unc5b* is expressed by embryonic kidney

endothelium and therefore we interrogated the vascular networks of *Foxd1$^{GC/+}$;Ntn1$^{fl/fl}$* kidneys.

Wholemount, 3D analyses revealed the loss of a predictable vascular pattern in mutant kidneys. As

vascular patterning has been linked to vessel maturity, we investigated arterialization in these mutants.

Quantification of the CD31+ endothelium at E15.5 revealed no differences in metrics such as the

number of branches or branch points, whereas the arterial vascular smooth muscle metrics were

significantly reduced at both E15.5 and P0. In support of these results, whole kidney RNA-seq showed

upregulation of angiogenic programs and downregulation of muscle-related programs which included

smooth muscle-associated genes. Together, our findings highlight the significance of netrin-1 to proper

vascularization and kidney development.

Keywords

Vasculature, stroma, patterning, netrin-1, kidney development

Introduction

Introduction

The organization of blood vessels into a network capable of maintaining organ and body fluid

homeostasis requires both proper development of the vasculature and specialization unique to the

tissue. Inappropriate vascular formation and patterning can result in pathologies in many organs(Blei

and Bittman, 2016; Heimann and Damato, 2010; Queisser et al., 2021; Wälchli et al., 2023). Vascular

endothelial cells organize into branched, hierarchical networks that form a continuous system for blood

flow. Larger arteries branch into smaller arterioles, which further divide into capillaries. Venules then

return the blood to larger veins and continue its circulation(Potente and Mäkinen, 2017). The vessels

within these networks are distinct in their morphology and function, which is often tailored to the

precise needs of the organ. For example, the endothelium of the kidney glomerular capillaries and

ascending vasa recta are fenestrated to enable their unique functions in filtration and diffusion,

respectively(Molema and Aird, 2012; Pallone et al., 2003). In association with their functions, the

endothelium of these different vascular beds also have distinct molecular signatures(Barry et al., 2019;

Daniel et al., 2018). Additionally, endothelial association with mural cells delineates the various types of

vessels within an organ. Larger vessels such as arteries are coated by vascular smooth muscle cells

whereas smaller capillaries are covered by pericytes, with each type of mural cell supporting distinct

vascular functions(Potente and Mäkinen, 2017). Altogether, these vascular properties highlight the need

for precise developmental programs that support the proper formation of these complex networks.

Recent studies have begun to shed light on the progressive formation of kidney vascular

networks during development. The temporal progression and predictable rudimentary patterning of the

major renal arteries through mid-gestation were elucidated through comprehensive imaging

studies(Daniel et al., 2018; Munro et al., 2017). Arteries which connect to the aorta undergo angiogenic

growth towards the developing kidney forming a vascular ring around the base of the ureteric bud by

E11.25(Munro et al., 2017). Continued growth, remodeling, and maturation of the endothelial network

leads to the establishment of an arterial tree by E13.5 that forms a predictable pattern(Daniel et al.,

2018). Renin expressing cells are localized along the developing arterial vasculature and mark points of

branching, and renin itself is important for proper vascular development (Gomez et al., 1989; Reddi et

al., 1998; Sequeira-Lopez et al., 2015b; Takahashi et al., 2005). At the molecular level, vascular

patterning was shown to be regulated by the association of mural cells with the nascent endothelium

and modulated in part by Pbx1(Hurtado et al., 2015). Deletion of the transcription factor *Pbx1* from the

Foxd1+ stromal progenitors induces precocious differentiation and premature association of vascular

mural cells with the endothelium, leading to stochastic arterial patterning and impaired renal

function(Hurtado et al., 2015). Another study showed that Dicer activity in the kidney stroma is

important for patterning smaller vessels, such as the peritubular and glomerular capillaries(Nakagawa et

al., 2015). While these studies have garnered some of the first insights into how the kidney vascular

network is established and organized, we still lack a clear understanding of how this process is

modulated at the molecular level.

Signaling molecules play important roles during development, often providing instructional cues

which help pattern the developing tissues. In respect to vasculature, a number of signaling pathways,

which also regulate axon guidance, provide angiogenic guidance. This includes Slit/Robo, Eph/Ephrin,

Semaphorins/Plexin-Neuropilins, and Netrins/UNC5(Adams and Eichmann, 2010). These guidance cues

direct vascular development during embryogenesis, and unsurprisingly, the removal of many of these

cues or their receptors results in vascular defects and mortality in the forming embryo (Bin et al., 2015;

Gerety et al., 1999; Gitler et al., 2004; Gu et al., 2005, 2003; Jones et al., 2008; Lu et al., 2004). Netrin-1

(*Ntn1*) is a classical neuronal guidance cue well-known for its role in commissural axon

guidance(Kennedy et al., 1994; Serafini et al., 1994). However, roles for netrin-1 have expanded to

include mediating pancreas development, inner ear morphogenesis, mammary gland development, lung

branching, and angiogenesis (Larrivée et al., 2007; Liu et al., 2004; Lu et al., 2004; Salminen et al., 2000;

Srinivasan et al., 2003; Sun et al., 2011; Wilson et al., 2006; Yebra et al., 2003). Endothelial guidance by

netrin-1 is accomplished through chemotactic interactions with its cognate receptors in the vasculature,

Unc5b and CD146 (*Mcam1*), which regulate blood vessel growth, branching, and patterning by

promoting either a repulsive or attractive response, respectively (Larrivée et al., 2007; Lu et al., 2004; Tu

et al., 2015). Whether netrin-1 plays a similar role in guiding vascular development and patterning in the

kidney remains unexplored.

Here we report that netrin-1 (*Ntn1*) is expressed and secreted by stromal progenitors in the

developing kidney. Deletion of *Ntn1* from this population results in hypoplastic kidneys and delayed

cessation of nephrogenesis. In correlation with its known role in vascular development, we found that

netrin-1 supports proper patterning and maturation of the vasculature during kidney organogenesis.

Our findings correlate with those in the accompanying manuscript by **Luo, Gu et al.**, and together we

establish a critical role for netrin-1 in mediating proper kidney development through regulation of

vascular maturity.

Results

Ntn1 is expressed and secreted by Foxd1+ stromal progenitors in the developing kidney

While netrin family members including netrin-1 play important roles in other branching

organs during development, such as the lung, pancreas, and mammary gland, whether they

have a role in kidney development has not been investigated to date(Liu et al., 2004; Srinivasan

et al., 2003; Yebra et al., 2003). To begin, we examined the expression pattern of *Ntn1* mRNA in

the developing kidney and found that *Ntn1* is expressed by stromal progenitors around the

nephron progenitor caps at embryonic day (E)15.5 and, to a lesser extent, in the proximal

tubule which is more specifically detected at later postnatal (P2) stages of development (Fig.

1A; Fig. S1A). This cellular expression pattern is confirmed by publicly available single cell RNA-

seq data of the kidney at E18.5(Combes et al., 2019; England et al., 2020). As early as E11.5,

secreted netrin-1 protein was observed surrounding the Six2+ nephron progenitors where the

stromal progenitors coalesce (Fig. 1B). At E15.5, netrin-1 was similarly secreted into the area

surrounding the nephron progenitor niche (Fig. 1C, arrow). Some protein was diffused into the

area below the nephron progenitor niche where ureteric tips lie, potentially acting as a ligand

sink to prevent further diffusion (Fig. 1C). The stromal progenitors expressing *Ntn1* give rise to

the renal interstitium, mesangial cells, and vascular mural cells, among others, and function in

advancing nephron progenitor differentiation(Hatini et al., 1996; Kobayashi et al., 2014;

Sequeira-Lopez et al., 2015a), suggesting that netrin-1 signaling may support proper kidney

development.

Deletion of Ntn1 from stromal progenitors disrupts normal kidney development

126   Germline deletion of *Ntn1* was previously shown to result in embryonic lethality by

E14.5(Bin et al., 2015). To circumvent this early lethality and address the role of netrin-1 in

kidney development, we conditionally deleted *Ntn1* from the Foxd1+ stromal progenitors by

using the *Foxd1^{GC}* allele in combination with a *Ntn1* floxed allele(Bin et al., 2015; Humphreys et

al., 2010; Kobayashi et al., 2014). The resulting *Foxd1$^{GC/+}$; Ntn1$^{fl/fl}$* mice have very little or no

netrin-1 protein present in the interstitial regions surrounding the nephron progenitors,

confirming efficient deletion (Fig. 1C). Our initial observations found that *Foxd1$^{GC/+}$; Ntn1$^{fl/fl}$*

mutant animals have hypoplastic kidneys at E15.5 and this phenotype persists postnatally (Fig.

1D). Immunostaining of the ureteric tree at E14.5 also revealed qualitative differences in kidney

size (Fig. S1B). Measurements of kidney length showed these reductions became most

significant after E13.5 (Fig. 1E).

As a means of assessing whether *Ntn1* deletion affects either the onset of kidney

development or cessation of nephrogenesis, we examined the presence of Six2+ nephron

progenitors at the initial branching event (E11.5) and after the cessation of nephrogenesis (P4),

which normally occurs around P2-P3 in the mouse(Hartman et al., 2007; Self et al., 2006). At

E11.5, we observed no empirical differences in the condensation of Six2+ nephron progenitors

around the ureteric buds suggesting kidney development initiated at the same time in *Ntn1*

conditional mutants (Fig. 2A). Analysis of nephron progenitor niches at E15.5 revealed no

obvious differences in their organization, suggesting normal progression of nephrogenesis (Fig.

2B). Strikingly though, at P4, $Foxd1^{GC/+}$; $Ntn1^{fl/fl}$ kidneys maintained small clusters of Six2+ cells

whereas controls had exhausted their Six2+ population as expected (Fig. 2C). Our previous bulk

nephron progenitor RNA-seq data as well as recent scRNA-seq data indicate the netrin receptor

Unc5c is expressed by nephron progenitors(Combes et al., 2019; O'Brien et al., 2018, 2016). *In*

situ hybridization of *Unc5c* confirmed expression by the nephron progenitors, as well as

podocytes and smooth muscle cells of the ureter (Fig. 2D). Due to the nephron progenitor

expression, we wondered whether disrupted signaling via netrin-1 to the *Unc5c+* nephron

progenitors might account for the smaller kidney size and delay in cessation of nephrogenesis.

We assessed the kidney phenotype of whole animal *Unc5c* knockouts(Ackerman et al., 1997)

and found no difference in kidney size compared to controls (Fig. 2E). Additionally, the Six2+

nephron progenitor niches appear normal, and no other overt phenotypes were observed at

any stages examined (Fig 2F). Therefore, the absence of netrin-1 signaling through Unc5c is

unlikely to cause the observed hypoplastic phenotype.

In developing $Foxd1^{GC/+}$; $Ntn1^{fl/fl}$ kidneys, the loss of $Ntn1$ could affect the number of

progenitors in nephron progenitor or ureteric tip niches, leading to hypoplasia. To assess cell

numbers, we analyzed kidney sections at multiple timepoints and counted Six2+ nephron

progenitors and Ecad+ ureteric tip cells within the niche (Fig. 2G). Apart from a transient

difference in tip cell numbers at E13.5, neither population showed any significant differences

between $Foxd1^{GC/+}$; $Ntn1^{fl/fl}$ kidneys and controls when assessed at E13.5, E15.5, and P0 (Fig.

2H). Using the same method, we investigated whether the proliferative capacity of niche

progenitors was impacted by measuring EdU incorporation into these populations as a measure

of cell cycling (Fig. 2G). We found no significant differences in the number of EdU+ cells in

either Six2+ or Ecad+ progenitor populations suggesting cell proliferation, at least at the stages

analyzed, was not causing the hypoplastic phenotype (Fig. 3C).

Netrin-1 mediates vascular patterning in the developing kidney

Both axons and endothelium respond to guidance cues such as netrin-1 using specific

receptors. Unc5b has been shown to mediate angiogenesis (Larrivée et al., 2007; Lu et al., 2004)

and *in situ* hybridization at E15.5 shows that *Unc5b* is expressed by the kidney endothelium as

well as the ureteric epithelium, although is excluded from the most peripheral ureteric

branches(Lindström et al., 2018) (Fig 3A). During kidney development, axonal projections tightly

associate and track with the vasculature, forming tight neurovascular bundles (Fig 3B).

Therefore, we included both nerves and vessels in our analyses since both have the potential to

respond to netrin-1 guidance. To assess overall organization and patterning of these networks,

we utilized wholemount immunostaining in combination with light-sheet microscopy to

generate 3D visualizations. Control kidneys show the formation of an extensive CD31+

endothelial network by E15.5 (Fig. 3C). Tuj1+ axons track closely with the main arterial

branches at this stage which form a pattern consistent with those observed by Daniel et

al.(Daniel et al., 2018) (Fig 3C, Suppl. Video 1). In contrast, we observed aberrant neuronal and

vascular patterning in the $Foxd1^{GC/+}$; $Ntn1^{fl/fl}$ mutant kidneys (Fig. 3C, Suppl. Video 2). The

example shown features looping of the arterial tree within the kidney cortex and ectopic

vessels on the kidney surface (arrowheads, Fig. 3C). Tracing of the main CD31+ arterial branches

shows that controls form a predictable pattern, as expected, which can be overlayed into 5

main branches from a lateral view(Daniel et al., 2018) (Fig. 3D). However, the $Foxd1^{GC/+}$; $Ntn1^{fl/fl}$

kidneys have a breakdown of predictable patterning and overall stochastic patterning (Fig. 3D).

Though not a focus of this study, the cortical vasculature around the developing nephron

progenitor niches of $Foxd1^{GC/+}$; $Ntn1^{fl/fl}$ kidneys is also disrupted, featuring disorganized and

ectopic vasculature (Fig. S2A). Notably, the presence of Ter119+ erythroid cells indicates that

the blood vessels are perfused in both control and $Foxd1^{GC/+}$; $Ntn1^{fl/fl}$ kidneys at E15.5, although

contribution from hemo-vasculogenesis is also possible (Sequeira Lopez et al., 2003)(Fig. S2B).

$Ntn3$ (netrin-3), a functionally related guidance cue that can engage netrin-1 receptors(Wang et

al., 1999), is also expressed, albeit at lower levels, by the stromal progenitors(Combes et al.,

2019; England et al., 2020) (Fig S3A). However, we did not observe any neurovascular

patterning defects or differences in kidney size in $Ntn3$ knockouts or any exacerbation of the

$Foxd1^{GC/+}$;$Ntn1^{fl/fl}$ phenotype when double knockouts ($Foxd1^{GC/+}$;$Ntn1^{fl/fl}$;$Ntn3^{-/-}$) were

generated (Fig. S3B,C). Together, these data suggest netrin-1 is the primary netrin guidance cue

responsible for the neurovascular patterning defects.

As both axons and endothelium can respond to netrin-1 guidance cues, we sought to

differentiate the primary responder to stromal-derived netrin-1 signals. Deletion of the *TrkA*

receptor for Nerve Growth Factor (NGF) induces apoptosis of peripheral neurons such as those

that innervate the kidney(Smeyne et al., 1994). We investigated vascular patterns in the setting

of reduced kidney innervation utilizing a *TrkA* knockout allele(Liebl et al., 2000). In *TrkA*

knockout kidneys, the vascular pattern was qualitatively normal at E16.5 (Fig. 3E). However,

when combined with the *Foxd1$^{GC/+}$;Ntn1$^{fl/fl}$* alleles to generate *Foxd1$^{GC/+}$;Ntn1$^{fl/fl}$;TrkA$^{-/-}$* animals,

the kidney endothelium still displayed abnormal patterning including looping and ectopic

vessels (Fig. 3E). The ectopic and looping vessels were not associated with Tuj1+ axons (Fig. 3E,

arrowheads), suggesting the endothelium is the primary responder to netrin-1 guidance cues

with the nerves following the vascular pattern, and in particular the smooth muscle coated

(alpha-smooth muscle actin, SMA+) endothelium (Fig. 3E). We also wanted to investigate

whether *Unc5c* knockout kidneys displayed altered patterning, even though they were not

hypoplastic. The neurovascualture appeared largely normal with no ectopic vessels, aberrant

looping, or qualitative reductions suggesting Unc5c is not a major receptor for netrin-1 in the

kidney or is functionally redundant with another receptor (Fig. S3D).

Vascular smooth muscle cell coverage is reduced in Foxd1$^{GC/+}$; Ntn1$^{fl/fl}$ kidneys

As the endothelium matures it is covered by a layer of vascular smooth muscle cells

(vSMCs). These cells provide support, maintain arterial wall integrity, and can direct innervation

through the release of guidance molecules(Brunet et al., 2014; Potente and Mäkinen, 2017). To

determine if vSMCs were impacted in *Foxd1*^{GC/+}; *Ntn1*^{fl/fl} animals, we investigated SMA coverage

of the endothelium in wholemount immunostained kidneys. Despite the lack of predictable

patterning, quantification of the CD31+ endothelium in *Foxd1*^{GC/+}; *Ntn1*^{fl/fl} kidneys revealed that

branching metrics and total vessel length were not significantly affected at E15.5 (Fig S4A, Fig

4A). However, the CD31+ endothelium showed qualitative reductions in SMA coverage in

Foxd1^{GC/+}; *Ntn1*^{fl/fl} kidneys compared to controls (Fig. 4B, arrowheads, Suppl. Videos 3,4).

Furthermore, *Tuj1*+ axons tracked exclusively with SMA+ vSMCs, representing arteries, and

were not found anywhere where endothelium singularly expressed CD31. These data suggest

that *Ntn1* loss impacts smooth muscle coverage of the vasculature, which is associated with

vessel stabilization and maturity.

While vascular patterning defects in *Foxd1*^{GC/+}; *Ntn1*^{fl/fl} kidneys are apparent, they are

stochastic and highly variable. Since some kidneys appear to have a more significant degree of

mis-patterning, they may also display differences in vSMC coverage. However, despite the

stochastic patterning differences, quantification of the SMA+ arterial vasculature showed

reductions at E15.5 and P0 in *Foxd1*^{GC/+}; *Ntn1*^{fl/fl} kidneys (Fig. 4C,D). On average, E15.5 kidneys

from *Foxd1*^{GC/+}; *Ntn1*^{fl/fl} animals have greater than a 20% reduction in SMA+ arterial metrics

including branch number, branch points, and the total length of the vSMC-lined vasculature

(Fig. 4D). At P0, reductions in these three vascular parameters were still significant highlighting

the persistence of this phenotype throughout development. To determine whether the defects

observed in development extended into adulthood, we labelled the vasculature of 2-3 month

old *Foxd1*^{GC/+}; *Ntn1*^{fl/fl} animals with Evans blue (Fig 4C). Evans blue is a bis-azo dye that

fluoresces only while bound to albumin and labels the perfused renal vasculature(Honeycutt

determine if vSMCs were impacted in $Foxd1^{GC/+}$; $Ntn1^{fl/fl}$ animals, we investigated SMA coverage

of the endothelium in wholemount immunostained kidneys. Despite the lack of predictable

patterning, quantification of the CD31+ endothelium in $Foxd1^{GC/+}$; $Ntn1^{fl/fl}$ kidneys revealed that

branching metrics and total vessel length were not significantly affected at E15.5 (Fig S4A, Fig

4A). However, the CD31+ endothelium showed qualitative reductions in SMA coverage in

$Foxd1^{GC/+}$; $Ntn1^{fl/fl}$ kidneys compared to controls (Fig. 4B, arrowheads, Suppl. Videos 3,4).

Furthermore, *Tuj1*+ axons tracked exclusively with SMA+ vSMCs, representing arteries, and

were not found anywhere where endothelium singularly expressed CD31. These data suggest

that *Ntn1* loss impacts smooth muscle coverage of the vasculature, which is associated with

vessel stabilization and maturity.

While vascular patterning defects in $Foxd1^{GC/+}$; $Ntn1^{fl/fl}$ kidneys are apparent, they are

stochastic and highly variable. Since some kidneys appear to have a more significant degree of

mis-patterning, they may also display differences in vSMC coverage. However, despite the

stochastic patterning differences, quantification of the SMA+ arterial vasculature showed

reductions at E15.5 and P0 in $Foxd1^{GC/+}$; $Ntn1^{fl/fl}$ kidneys (Fig. 4C,D). On average, E15.5 kidneys

from $Foxd1^{GC/+}$; $Ntn1^{fl/fl}$ animals have greater than a 20% reduction in SMA+ arterial metrics

including branch number, branch points, and the total length of the vSMC-lined vasculature

(Fig. 4D). At P0, reductions in these three vascular parameters were still significant highlighting

the persistence of this phenotype throughout development. To determine whether the defects

observed in development extended into adulthood, we labelled the vasculature of 2-3 month

old $Foxd1^{GC/+}$; $Ntn1^{fl/fl}$ animals with Evans blue (Fig 4C). Evans blue is a bis-azo dye that

fluoresces only while bound to albumin and labels the perfused renal vasculature(Honeycutt

and O'Brien, 2021). At two months, quantification of $Foxd1^{GC/+}$; $Ntn1^{fl/fl}$ kidneys labeled with

Evans blue revealed no significant differences in vascular metrics, although all three metrics

trended towards a reduction (Fig. 5D). At both 2 and 3 months of age, ectopic vessels were no

longer observed and the kidney vasculature has fewer visible defects, although this is difficult

to ascertain due to the highly complex vascular network that has formed. However, differences

were noted such as meandering, elongated vessels near the cortex (Fig 4C and Fig S4B, red

dotted lines) and areas of reduced branching (Fig S4B, yellow arrows). Additionally, we

occasionally observed clouds of Evans blue signal outside the vascular network (Fig S4B, white

arrowhead) indicating possible leakage and compromise to vessel integrity.

RNA-seq reveals differences in vascular and metabolic programs of Foxd1$^{GC/+}$; Ntn1$^{fl/fl}$ kidneys

To better understand alterations to transcriptional programs relevant to the vascular

immaturity and the hypoplastic phenotype observed in $Foxd1^{GC/+}$; $Ntn1^{fl/fl}$ animals, we

performed bulk RNA-seq on whole kidneys (Table S1). To account for any exacerbated

differences due to the loss of one *Foxd1* allele from the *GFP-Cre* insertion, we compared

$Foxd1^{GC/+}$; $Ntn1^{fl/fl}$ mutant kidneys to $Foxd1^{GC/+}$; $Ntn1^{fl/+}$ heterozygous kidneys. We utilized a

cutoff of 1.2-fold as more significant changes may be drowned out due to the bulk analysis.

Gene ontology analyses were then performed on up- and downregulated genes to identify

biological processes enriched in either category (Tables S2 and S3). In support of our imaging

analyses, processes related to the circulatory system and blood vessel development were

upregulated in $Foxd1^{GC/+}$; $Ntn1^{fl/fl}$ kidneys and included genes such as *Notch1*, *Flt1*, *Nrp2*, and

Unc5b (Fig. 5A, Table S2). Programs related to cell adhesion and cell migration were also

upregulated and included genes in the cadherin superfamily and planar cell polarity related

genes, respectively (Fig 5A, Table S2). Such pathways may affect the vSMC association with the

endothelium in addition to affecting overall kidney development, thus leading to the

hypoplastic phenotype. In support of the reduction in vSMC coverage of the endothelium in

mutant kidneys, muscle related processes were downregulated and genes in these categories

included *Acta2* which encodes for α-SMA and *Myh11* which is a smooth muscle myosin (Fig 5B,

Table S3). Additional biological processes downregulated in *Foxd1$^{GC/+}$; Ntn1$^{fl/fl}$* kidneys were

related to more general cellular processes such as translation, RNA processing, and glycolysis

(Fig 5B). Altogether, these data suggest that in the absence of netrin-1 signaling, the kidney

vasculature is more immature (increased angiogenic processes vs arterialization) which may

affect metabolic and cellular processes, leading to the hypoplastic phenotype (Fig. 5C).

Discussion

In this study, we demonstrate that stromal progenitor *Ntn1* is necessary for proper

patterning and maturation of the kidney arterial network. *Foxd1$^{GC/+}$; Ntn1$^{fl/fl}$* kidneys have

visible and quantifiable stochastic vascular patterning defects with reductions in most arterial

metrics during embryogenesis and postnatally. These differences in vascular metrics largely

resolve in adulthood, although patterning defects are still observed. Furthermore, deletion of

Ntn1 resulted in a delay in the cessation of nephrogenesis and kidney hypoplasia. Altogether,

our studies and those in the accompanying manuscript by **Luo, Gu et. al.,** which uncover similar

findings, highlight the significance of netrin-1 to proper kidney vascularization and associated

development.

While we find differences in vascularization in the absence of netrin-1, our analyses

relate specifically to the larger arteries and arterioles of the vascular tree. SMA will only label

vasculature with sufficient smooth muscle coverage, typically larger arteries and arterioles,

missing the capillaries which are primarily covered by NG2+ pericytes. However, **Luo, Gu et. al.**

demonstrate a lack of proper NG2+ pericyte localization in early development, suggesting

netrin-1 is an important signaling molecule for all vascular mural cells. Additionally, we did not

investigate venous formation and patterning during development, primarily owing to a lack of

appropriate and agreed-upon markers of the venous vasculature(Daniel et al., 2018). Veins are

also covered with significantly fewer SMCs due to a decreased need to regulate pressure and

tone[38]. This makes SMA an unreliable venous marker at lower magnifications, such as those

used by light-sheet microscopy to image the vasculature. Therefore, additional studies are

necessary to determine to what extent the entire vascular plexus is disrupted in *Foxd1$^{GC/+}$;*

Ntn1$^{fl/fl}$ kidneys.

Vascular patterning and maturity are both disrupted in the *Foxd1$^{GC/+}$; Ntn1$^{fl/fl}$* kidneys in

our study. **Luo, Gu et. al.** further define the maturity phenotype as being associated with the

loss of Klf4 and accumulation of mural cells on the surface of the kidneys. We did not observe

any obvious accumulation of mural cells at E15.5 or later stages, and whether Klf4 is associated

with the phenotype in our studies is unknown. We did not find a difference in *Klf4* expression in

our E15.5 RNA-seq, although this is 2 days later than the RNA-seq performed by **Luo, Gu et. al.**

and Klf4 levels may have normalized by this time. Therefore, the observed maturation

phenotypes may initiate at the earliest stages of arterialization, around E13.5, and prior to our

analysis at E15.5(Daniel et al., 2018). The timing of the phenotype could explain the differences

between our study and that of **Luo, Gu et. al.,** highlighting the complementary nature of our

analyses. VSMCs *in vitro* express netrin receptors such as Neogenin (*Neo1*) and Unc5b and

signaling through these receptors supports endothelial vSMC coverage(Lejmi et al., 2014, 2008).

As such, deletion of *Ntn1* from the stroma may ultimately impact SMC maturation or

recruitment in the kidney and, in turn, affect vascular patterning. This is similar to the findings

of Hurtado et al. who showed that Pbx1-mediates mural cell maturation ultimately influencing

proper vascular patterning within the developing kidney(Hurtado et al., 2015). Alternatively,

patterning of the endothelium may influence vascular maturation. Deletion of *Unc5b* from the

kidney endothelium and the assessment of vascular patterning and maturity would potentially

help sort out if the patterning defects we observed are independent of vSMC coverage. The lack

of significant phenotypes in the *Unc5c* knockouts suggest it is either not a major receptor for

netrin-1 in the kidney, or it is functionally redundant. Considering netrin-1 can signal through a

large number of receptors, including integrins, and display receptor independent cross-

repressive activities with LRIG proteins, uncovering the receptor(s) and downstream signaling

mechanisms of netrin-1 in the kidney may prove challenging (Abdullah et al., 2021; Abraira et

al., 2008; Stanco et al., 2009; Sun et al., 2011; Yebra et al., 2003).

The kidney hypoplasia observed in our *Foxd1$^{GC/+}$; Ntn1$^{fl/fl}$* mutants may be linked to the

relative vascular immaturity we observed. This could perturb metabolic processes within the

kidney leading to downregulation of the glycolysis pathways and associated genes in our RNA-

seq data, and ultimately affect proper kidney growth and prolong the nephrogenic period(Liu et

al., 2017). Alternatively, a mechanism independent of the vasculature may perturb proper

kidney development. Altered interactions between netrin-1 and Bmp4 could lead to the

hypoplastic phenotype. Bmp4 is a negative regulator of ureteric branching through its

modulation of Gdnf signaling(Miyazaki et al., 2000). Dysregulation of Gdnf signaling can result

in hypoplasia and agenesis at the extreme(Moore et al., 1996; Pichel et al., 1996; Sánchez et al.,

1996). Netrin-1 suppresses Bmp4 signaling *in vitro,* and netrin-1 mediated modulation of Bmp

signaling within the kidney could help balance Gdnf signaling(Abdullah et al., 2021). It is

possible that loss of *Ntn1* results in increases in Bmp4 and a subsequent reduction in Gdnf

activity, leading to renal hypoplasia. Our RNA-seq analyses did not reveal any significant

perturbations to *Bmp4* or *Gdnf* levels, although assessing an earlier timepoint when the

hypoplasia is initiated may be more informative and help sort out the cause of the hypoplastic

phenotype.

Whether the development of a kidney with abnormal vascular patterns affects

physiological function remains unknown and will be the focus of future studies. $Foxd1^{GC/+}$;

$Ntn1^{fl/fl}$ animals do live into adulthood, with no significant kidney or health phenotypes.

However, if the smaller vascular beds in important functional regions of the nephron such as

the glomerulus and tubules are organized properly, then physiological function may be largely

unperturbed. However, injury, disease, or aging may exacerbate or induce changes in

physiological function in these animals regardless. Understanding whether precise vascular

patterning is critical to function, both in health and under physiological stress, will help inform

efforts to vascularize tissues for functional transplant.

As the kidney lacks regenerative mechanisms, *de novo* kidney engineering and *in situ*

repair using *in vitro* generated nephrons or iPSC-derived renal organoids are considered

attractive therapeutic possibilities(Morizane et al., 2015; Taguchi et al., 2014; Takasato et al.,

2015; Wu et al., 2018). Unfortunately, neither treatment is currently approaching clinical

relevance(Bantounas et al., 2018). With current protocols, renal organoids form basic nephron

structures but eventually become necrotic from a deficit of oxygen and nutrients since

functional endothelium is not present(Grebenyuk and Ranga, 2019). Interestingly, cells

expressing the pan-endothelial marker CD31 occur *de novo* in organoids, but without perfusion

they do not form connected, functional networks, resulting in a lack of proper lumenization,

flow, and patterning and their eventual regression(Homan et al., 2019; Ryan et al., 2021;

Taguchi et al., 2014; Wu et al., 2018). Additionally, the vessels would need to obtain

appropriate mural cell coverage to support function. Therefore, studies such as ours and those

of **Luo, Gu et. al.** provide critical insights into how the renal vasculature is patterned, matures,

and contributes to kidney development and will help guide kidney engineering strategies.

Material and methods

Animals

All animal studies were approved by the Office of Animal Care and Use at the University

of North Carolina at Chapel Hill and the UNC Institutional Animal Care and Use Committee

(IACUC). Procedures were performed under IACUC-approved protocols for mice (16-276.0, 19-

183.0, 22-136). Animal husbandry is performed by UNC's Division of Comparative Medicine

(DCM). All animals in this study were kept in environmental conditions consisting of

temperatures between 68-74°F, with humidity ranges of 30-70%. No more than five adult

animals were housed per cage, and all cages were provided a 12-hour light/dark cycle. Animals

are fed LabDiet PicoLab Select Rodent 50 IF/6F (Catalog #, 5V5R). Timed matings were set up

and plugs were ascertained by visual and probed inspection. The presence of a plug was

considered 0.5 days gestation. The day of birth (E19.5) was considered postnatal day 0 (P0),

pups born before or after E19.5 were not utilized in these studies.

All mouse lines are maintained on the C57Bl6/J background (Jackson Labs, Strain

#000664). Details of genetically edited mouse lines utilized are as follows: $Foxd1^{GC/+}$ (Strain

#012463)(Humphreys et al., 2010; Kobayashi et al., 2014) and $Ntn1^{fl/fl}$ (Strain #028038)(Bin et

al., 2015) mice were purchased from Jackson Labs, $TrkA$ knockout mice(Liebl et al., 2000) were

a gift from Lino Tessarollo at the NIH National Cancer Institute Center for Cancer Research, and

$Unc5c$ knockout mice(Ackerman et al., 1997) were a gift from Susan Ackerman at the University

of California, San Diego. $Foxd1^{GC/+}$; $Ntn1^{fl/fl}$ mice were generated by crossing Cre-expressing

$Foxd1^{GC/+}$; $Ntn1^{f/+}$ to $Ntn1^{fl/fl}$ mice unless otherwise noted. Controls were either $Ntn1^{c/c}$ or

$Foxd1^{GC/+}$; $Ntn1^{fl/+}$ as we did not find any qualitative or quantitative differences between

genotypes for any of the analyses.

$Ntn3$ knockout mice were generated by the UNC Animal Models Core using

CRISPR/Cas9-mediated genome editing. Guide RNAs $Ntn3$-gRNA1 (CGAGGAATCGCAGACGCGAT,

Exon 1, chr17:24208777-24208796) and $Ntn3$-gRNA2 (GAAAGCGTGTCCACGAGCCG, Intron 5,

chr17:24206891-24206910) were designed to delete the majority of the $Ntn3$ coding region

spanning Exons 1-6. Guide RNAs were generated within our lab and provided to the animal

models core at estimated concentrations of 500ng/µl and 450ng/µl, respectively. Guide RNAs

were prepared as in O'Brien et al.(O'Brien et al., 2018). Recombinant Cas9 protein was

expressed and purified by the UNC Protein Expression Core using a plasmid provided by the

Animal Models Core. Two hundred and ten fertilized embryos (zygotes) were harvested from 9

super-ovulated $Ntn1^{fl/fl}$ females crossed to $Foxd1^{GC/+}$; $Ntn1^{fl/fl}$ males. Zygotes were

electroporated with two different reagent concentrations diluted in OptiMEM medium (Gibco)

and then implanted in pseudo-pregnant recipient females. Mix 1: 1.2 µM Cas9 protein, 47 ng/µl

each guide RNA. 102 embryos electroporated; 85 embryos implanted in 4 B6D2F1 recipients.

Mix2: 400 nM Cas9 protein, 25 ng/µl each guide RNA. 108 embryos electroporated; 95 embryos

implanted in 4 B6D2F1 recipients. Of the resulting pups, one male containing the correct

deletion as confirmed by Sanger sequencing over the deletion site was used as the founder.

Brain and kidney tissue was harvested, and qPCR performed to confirm the loss of *Ntn3*

transcripts in knockout animals.

Genotyping primers utilized in this study: *Ntn1* F: GGCAGTGAATGTGTTTCCGTT, *Ntn1* R:

ATCGCGGGATAGTGGGTTTC; *Foxd1* WT F: CTCCTCCGTGTCCTCGTC, *Foxd1* Mut F:

GGGAGGATTGGGAAGACAAT, *Foxd1* Common R: TCTGGTCCAAGAATCCGAAG; *Ntn3* Internal

(WT) F: GGCCTGTGGTCTGGTTACAG, *Ntn3* Internal (WT) R: TAGCCTGGGGACTTCTGACC, *Ntn3*

External (deletion) F: CACCTCCCAGCAGCAAGTAAC, *Ntn3* External (deletion) R:

ACGCAGAGTAGCGGACTAGG; *TrkA* 1: TGTACGGCCATAGATAAGCAT, *TrkA* 2:

TGCATAACTGTGTATTTCAC, *TrkA* 3: CGCCTTCTTGACGAGTTCTTCTG; *Unc5c* WT F:

CACTCTATGGAAATGGCTGAAT, *Unc5c* WT R: GTCCTCCAATCCAAGAACTG, *Unc5c* Mut F:

CAGGAGAAGATACATTTAACCAC, *Unc5c* Mut R: GACAGAAGAGCATAGCATTCAC

Kidney size measurements

Kidney length measurements were performed using FIJI software to analyze images

taken at 2x magnification on a Leica dissecting scope. Standardization and calibration of FIJI

measurements were performed using a 1µm-1mm micrometer taken at the same

magnification.

In situ hybridization

In situ hybridizations were performed on wildtype cryosectioned E15.5 and P2 Swiss

Webster (SW, Taconic) kidneys using the "Section In Situ Hybridization (SISH) on kidney

sections" protocol (Krautzberger and Guo; McMahon Lab;

https://www.gudmap.org/chaise/record/#2/Protocol:Protocol/RID=N-H9AC) on GUDMAP

(www.gudmap.org). Digoxygenin (DIG) labeled probes were generated following the

"Digoxigenin-labelled Riboprobe Synthesis from a PCR-generated DNA templates" protocol

found on GUDMAP (Little Group, GUDMAP Consortium). Primers utilized for probes are as

follows: *Ntn1* F:CTTCCTCACCGACCTCAATAAC, *Ntn1* (SP6) R:

GCGATTTAGGTGACACTATAGTTGTGCCTACAGTCACACACC; *Ntn3* F:GGCACGCTCTCCGCTGCA,

Ntn3 (T7) R:TAATACGACTCACTATAGGGAGTGGCTGAGTAGGAGTA; *Unc5c*

F:ATGAGGAAAGGTCTGAGGGCGACAG, *Unc5c* (T7)

R:TAATACGACTCACTATAGGGAGAGAGTTGAAGGTACCAA; *Unc5b* F:TCAAGTGTAATGGCGAGTG,

Unc5b (T7) R:TAATACGACTCACTATAGGGGCCTCCATTCACATAGACGA

Section immunostaining

For all samples, kidneys were harvested at the necessary time point. Tissue was then

fixed in 4% paraformaldehyde (PFA) in 1 x PBS for 15-30 minutes (embryonic/postnatal) to 24

hours (adult). Kidneys were washed 2x in PBS and stored for up to two weeks before

processing. Samples were incubated in 30% sucrose in PBS overnight at 4 ^{o}C. Tissue was

embedded in OCT media, frozen with 100% EtOH and dry ice, and stored at -80 ^{o}C for up to two

years. Frozen tissue was sectioned at 10-12μm. Sections were air-dried at room temperature

(RT) for a minimum of 10 minutes. Samples were then incubated in 1x PBS for 5 minutes to

remove OCT media. Next, sections were placed in blocking buffer (PBS with 3% BSA, 1% donkey

serum, and 0.25% Triton x-100) for 20 minutes at RT. The blocking buffer was removed, then

sections were incubated with primary antibodies in the blocking buffer for 1 hour at RT.

Sections were then rinsed 3x in wash buffer (PBS with 0.1% Triton x-100). Next, sections were

incubated with appropriate Alexa Fluor conjugated secondary antibodies at 1:1000 (Invitrogen)

in blocking buffer for 20 minutes. Sections were then rinsed 3x in wash buffer and mounted

with Prolong Gold with DAPI (Invitrogen). Antibodies utilized and their dilution are listed in

Table S4.

Quantification of proliferation

Mice were injected intraperitoneally with 1.25 mg of EdU (10mg/ml) 1.5 hours before

euthanasia and tissue collection. For analysis of proliferation, the following procedure was

used. The tissue was fixed, then sectioned and stained as described previously in this report.

The Click-It EdU kit was utilized to detect EdU positive cells. Images were then collected on a

Zeiss 880 confocal using a 40x oil objective. The whole kidney images were created through

automated tiling of the tissue section using Zen Black software. Reconstruction of the images

into a single image was performed using Imaris stitcher (Bitplane). For proliferation counts, a

minimum of 10 nephrogenic caps were counted per section for each genotype and age.

Quantitation of proliferation for nephron progenitors was determined as the number of cells

colocalized for both EdU and Six2 divided by the total number of Six2+ cells. Ureteric bud (UB)

tip niches were defined as the Ecad+ cells within the UB but above the ventral end of the

nephrogenic cap. Quantitation for UB proliferation was determined as the number of cells

within the UB niche colocalized with Ecad and EdU divided by the total number of Ecad+ cells

within the UB niche.

Wholemount immunostaining

The wholemount immunostaining protocol was adapted from Jafree et al.(Jafree et al.,

2019) and Renier et al.(Renier et al., 2014). All following steps are performed with rocking

unless otherwise noted. Samples are collected and fixed in 4% PFA for 15-30 minutes

(embryonic/postnatal). Tissue is washed with rocking 2x for 20 minutes at RT. Samples are then

dehydrated in an increasing MeOH/H_2O series of 25%, 50%, 75%, and 100% for 30 minutes or 1

hour, depending on tissue size. Next, tissue is bleached overnight in 5% H_2O_2 (from 30% stock)

in 100% MeOH. Samples are then rehydrated following a reverse MeOH/H_2O series (100%, 75%,

50%, 25%, PBS) for 30 minutes or 1 hour, depending on tissue size. Tissue is then permeabilized

overnight at 4 ^{o}C in PBS with 0.2% Triton x-100, 2.3% glycine, and 20% DMSO. The

permeabilization solution is removed, and blocking buffer (PBS with 0.2% triton x-100, 6%

donkey serum, and 10% DMSO) is added; the tissue is then incubated for one day at RT. The

blocking buffer is then removed, and primary antibody in staining solution (PBS with 0.2%

Tween-20, 0.1% heparin [from 10mg/ml stock], 3% donkey serum, and 5% DMSO) is added to

samples, which are then incubated for three days at 4^{o}C, followed by washing 4-6x for 1 hour at

RT in wash buffer (PBS with 0.2% Tween-20 and 0.1% heparin [from 10mg/ml stock]). The

appropriate Alexa fluor labeled secondary antibody in staining solution is then added (1:250)

and incubated overnight at 4^{o}C. Samples are washed 4-6x for 1 hour at RT in wash buffer.

Samples are then immediately moved to clearing or stored at 4^{o}C in 1x PBS, protected from

light until ready to proceed to clearing. Antibodies utilized and their dilution are listed in Table

S4.

Evans blue

The full protocol for Evans blue dye is from Honeycutt and O'Brien(Honeycutt and

O'Brien, 2021) and is described here briefly. Animals are anesthetized to a surgical plane under

isoflurane. Next, the chest cavity and peritoneum are cut to expose the diaphragm. 2% Evans

blue dye in 0.9% NaCl/H_2O at 3 μl/g of body weight is injected into the left ventricle with either

a 25g needle or 31g insulin syringe. The dye is allowed to circulate until the exposed areas

(paws, tail, chin) turn blue, roughly 2-5 minutes. The animal is subsequently euthanized, and

kidney is collected, rinsed briefly in PBS several times to remove excess dye, then fixed in 4%

PFA for 24 hours at 4^{o}C. Samples are washed in PBS and stored at 4^{o}C until clearing.

Tissue clearing and imaging

This protocol was adapted from the iDISCO method of tissue clearing(Renier et al.,

2014). First, embryonic and small samples are placed into agarose blocks (1% agarose in tris-

acetate-EDTA [TAE]). Adult kidneys are not embedded. All following steps are performed with

rocking unless otherwise noted. These blocks or adult kidneys are dehydrated in an increasing

MeOH/H_2O series of 25%, 50%, 75%, and 100% for 30 minutes or 1 hour at RT, depending on

tissue size. Samples are then incubated in 66% dichloromethane (DCM) / 33% MeOH for 3

hours at RT. Next, two rounds of incubation with 100% DCM at RT are performed. DCM is

removed, and 100% dibenzyl ether (DBE) is added. Samples are incubated, with no rocking,

until clear (1 – 24 hours) and stored indefinitely in 100% DBE at either RT or 4 ^{o}C, protected

from light, until ready to proceed to imaging. Light-sheet imaging of whole kidneys was

performed using a LaVision UltraMicroscope II (LaVision BioTec). 3D reconstruction of images

and vascular network quantification was conducted using Imaris imaging software (Bitplane).

Quantification of wholemount 3D vasculature

To conduct quantification of renal vasculature within Imaris, a surface of the 3D image

was first created to digitally render the vasculature. Next, a binary mask of the rendered

surface was created using the "Mask channel" function with "Constant inside/outside" settings

for voxel intensity set to 0 outside and 100 inside. This was necessary to render the vasculature

effectively solid, so that the filament tracer module within Imaris could be implemented to

develop wireframes for statistical analysis. Kidneys were batched with similar parameters per

age group. Then supervised automatic tracing was performed to ensure that automatically

generated paths were not duplicated, or artifacts inaccurately rendered as vasculature. As both

kidneys from the same animal have highly similar statistical values, with little deviation when

averaged, for most studies, only one kidney per animal was needed for quantification. This

method was used to quantify renal arterial trees for Evans blue in adults and SMA and CD31

staining in the embryonic and neonatal kidneys. Cortical arterioles and vascular beds were

excluded from analysis.

For graphical display of vascular patterning and imaging that were not quantified, a

surface was created for the desired channel. This surface was then masked using the same

procedure, but "Constant inside/outside" settings for voxel intensity were set to 0 outside with

the inside setting left unchecked.

RNA-seq

Whole kidneys were collected at E15.5 from 3 separate $Ntn1^{fl/fl}$ x $Foxd1^{GC/+}$;$Ntn1^{fl/+}$

crosses. Three kidneys were collected of each genotype: $Foxd1^{GC/+}$;$Ntn1^{fl/+}$ (heterozygous),

$Foxd1^{GC/+}$;$Ntn1^{fl/fl}$ (homozygous), and $Ntn1^{fl/fl}$ (wildtype). Kidneys were placed in RNAlater

(Thermo Fisher) until all samples were collected. RNA was extracted with TRIzol (Invitrogen).

RNA was sent to Cancer Genetics Inc (Morrisville, NC) for sequencing and data processing. RNA

samples were converted to sequencing libraries using the TruSeq Stranded mRNA kit (Illumina

Cat. #20020595). Resulting libraries were assessed for quality via analysis on a 2100 Bioanalyzer

(Agilent Technologies) and quantitated by qPCR (KAPA Library Quantification; Roche: KK4824,

Cat# 07960140001). Normalized and pooled libraries were sequenced on the Illumina Nextseq

550 system. Alignment and analysis were performed following a modified Tuxedo workflow.

Briefly, alignment was done with STAR aligner followed by cufflinks/cuffmerge for assembled

transcripts. FPKM was calculated with CuffDif and analyzed with the R package. Heterozygous

samples were used as a control and comparison to the homozygous samples to account for any

effects of the hemizygous $Foxd1$ allele due to the knock-in of GFP-Cre into the $Foxd1$ locus. A

count of 5 FPKM was considered minimum criteria for expression in the whole kidney RNA-seq,

genes with less than this FPKM (unless downregulated compared to the reference) were

excluded from gene expression analyses. Up- and downregulated gene lists were pasted into

the Database for Annotation, Visualization and Integrated Discovery (DAVID) to obtain Gene

Ontology information and enrichment.

<u>Acknowledgments</u>

We thank members of the O'Brien lab for technical help and critical discussions, Dr.

Pablo Ariel, director of the UNC Microscopy Services Laboratory, for assistance with light-sheet

imaging and Imaris rendering/analysis, Dr. Robert Currin (retired) of the UNC Hooker Imaging

Core for confocal microscopy imaging assistance, and Dr. Dale Cowley, director of the UNC

Animal Models Core, for help with *Ntn3* knockout mouse generation. We thank Drs. Lino

Tessarollo (NCI) and Susan Ackerman (UCSD) for their kind gifts of *TrkA* and *Unc5c* knockout

mice, respectively.

<u>Author Contributions</u>

LLO conceived and designed the project, performed experiments, analyzed data, and helped

write the manuscript. SEH helped design the experiments, performed experiments, analyzed

data, and wrote the manuscript. P-EYN, DMH, YX, and SLC performed experiments and analyzed

data.

<u>Competing Interests</u>

The authors have nothing to declare and no competing interests, relevant affiliations, financial

involvement, or financial conflicts with any entities or organizations that may have financial

interests in the research, subject matter, or materials in this manuscript apart from those

disclosed.

<u>Funding</u>

This work was supported by NIH R01DK121014 (LLO), 5T32HL069768 (SEH), and the

Ross and Charlotte Johnson Family Dissertation Fellowship (SEH). The UNC Animal Models Core,

the Microscopy Services Laboratory, and UNC Hooker Imaging Core Facility are supported in

part by the P30 CA016086 Cancer Center Core Support Grant to the UNC Lineberger

Comprehensive Cancer Center.

References

572
Abdullah, A., Herdenberg, C., Hedman, H., 2021. Netrin-1 functions as a suppressor of bone
morphogenetic protein (BMP) signaling. Sci Rep 11, 8585. https://doi.org/10.1038/s41598-021-
87949-7
Abraira, V.E., del Rio, T., Tucker, A.F., Slonimsky, J., Keirnes, H.L., Goodrich, L.V., 2008. Cross-repressive
interactions between Lrig3 and netrin 1 shape the architecture of the inner ear. Development
135, 4091–4099. https://doi.org/10.1242/dev.029330
Ackerman, S.L., Kozak, L.P., Przyborski, S.A., Rund, L.A., Boyer, B.B., Knowles, B.B., 1997. The mouse
rostral cerebellar malformation gene encodes an UNC-5-like protein. Nature 386, 838–842.
https://doi.org/10.1038/386838a0
Adams, R.H., Eichmann, A., 2010. Axon Guidance Molecules in Vascular Patterning. Cold Spring Harb
Perspect Biol 2, a001875. https://doi.org/10.1101/cshperspect.a001875
Bantounas, I., Ranjzad, P., Tengku, F., Silajdžić, E., Forster, D., Asselin, M.-C., Lewis, P., Lennon, R.,
Plagge, A., Wang, Q., Woolf, A.S., Kimber, S.J., 2018. Generation of Functioning Nephrons by
Implanting Human Pluripotent Stem Cell-Derived Kidney Progenitors. Stem Cell Reports 10, 766–
779. https://doi.org/10.1016/j.stemcr.2018.01.008
Barry, D.M., McMillan, E.A., Kunar, B., Lis, R., Zhang, T., Lu, T., Daniel, E., Yokoyama, M., Gomez-Salinero,
J.M., Sureshbabu, A., Cleaver, O., Di Lorenzo, A., Choi, M.E., Xiang, J., Redmond, D., Rabbany,
S.Y., Muthukumar, T., Rafii, S., 2019. Molecular determinants of nephron vascular specialization
in the kidney. Nat Commun 10, 5705. https://doi.org/10.1038/s41467-019-12872-5
Bin, J.M., Han, D., Lai Wing Sun, K., Croteau, L.-P., Dumontier, E., Cloutier, J.-F., Kania, A., Kennedy, T.E.,
2015. Complete Loss of Netrin-1 Results in Embryonic Lethality and Severe Axon Guidance
Defects without Increased Neural Cell Death. Cell Rep 12, 1099–1106.
https://doi.org/10.1016/j.celrep.2015.07.028
Blei, F., Bittman, M.E., 2016. Congenital vascular anomalies: current perspectives on diagnosis,
classification, and management. JVD 4, 23–37. https://doi.org/10.2147/JVD.S63244
Brunet, I., Gordon, E., Han, J., Cristofaro, B., Broqueres-You, D., Liu, C., Bouvrée, K., Zhang, J., del Toro,
R., Mathivet, T., Larrivée, B., Jagu, J., Pibouin-Fragner, L., Pardanaud, L., Machado, M.J.C.,
Kennedy, T.E., Zhuang, Z., Simons, M., Levy, B.I., Tessier-Lavigne, M., Grenz, A., Eltzschig, H.,
Eichmann, A., 2014. Netrin-1 controls sympathetic arterial innervation. J Clin Invest 124, 3230–
3240. https://doi.org/10.1172/JCI75181
Combes, A.N., Phipson, B., Lawlor, K.T., Dorison, A., Patrick, R., Zappia, L., Harvey, R.P., Oshlack, A.,
Little, M.H., 2019. Single cell analysis of the developing mouse kidney provides deeper insight

into marker gene expression and ligand-receptor crosstalk. Development 146, dev178673.
https://doi.org/10.1242/dev.178673
Daniel, E., Azizoglu, D.B., Ryan, A.R., Walji, T.A., Chaney, C.P., Sutton, G.I., Carroll, T.J., Marciano, D.K.,
Cleaver, O., 2018. Spatiotemporal heterogeneity and patterning of developing renal blood
vessels. Angiogenesis 21, 617–634. https://doi.org/10.1007/s10456-018-9612-y
England, A.R., Chaney, C.P., Das, A., Patel, M., Malewska, A., Armendariz, D., Hon, G.C., Strand, D.W.,
Drake, K.A., Carroll, T.J., 2020. Identification and characterization of cellular heterogeneity
within the developing renal interstitium. Development 147, dev190108.
https://doi.org/10.1242/dev.190108
Gerety, S.S., Wang, H.U., Chen, Z.F., Anderson, D.J., 1999. Symmetrical mutant phenotypes of the
receptor EphB4 and its specific transmembrane ligand ephrin-B2 in cardiovascular development.
Mol Cell 4, 403–414. https://doi.org/10.1016/s1097-2765(00)80342-1
Gitler, A.D., Lu, M.M., Epstein, J.A., 2004. PlexinD1 and semaphorin signaling are required in endothelial
cells for cardiovascular development. Dev Cell 7, 107–116.
https://doi.org/10.1016/j.devcel.2004.06.002
Gomez, R.A., Lynch, K.R., Sturgill, B.C., Elwood, J.P., Chevalier, R.L., Carey, R.M., Peach, M.J., 1989.
Distribution of renin mRNA and its protein in the developing kidney. Am J Physiol 257, F850-858.
https://doi.org/10.1152/ajprenal.1989.257.5.F850
Grebenyuk, S., Ranga, A., 2019. Engineering Organoid Vascularization. Front Bioeng Biotechnol 7, 39.
https://doi.org/10.3389/fbioe.2019.00039
Gu, C., Rodriguez, E.R., Reimert, D.V., Shu, T., Fritzsch, B., Richards, L.J., Kolodkin, A.L., Ginty, D.D., 2003.
Neuropilin-1 conveys semaphorin and VEGF signaling during neural and cardiovascular
development. Dev Cell 5, 45–57. https://doi.org/10.1016/s1534-5807(03)00169-2
Gu, C., Yoshida, Y., Livet, J., Reimert, D.V., Mann, F., Merte, J., Henderson, C.E., Jessell, T.M., Kolodkin,
A.L., Ginty, D.D., 2005. Semaphorin 3E and plexin-D1 control vascular pattern independently of
neuropilins. Science 307, 265–268. https://doi.org/10.1126/science.1105416
Hartman, H.A., Lai, H.L., Patterson, L.T., 2007. Cessation of Renal Morphogenesis in Mice. Dev Biol 310,
379–387. https://doi.org/10.1016/j.ydbio.2007.08.021
Hatini, V., Huh, S.O., Herzlinger, D., Soares, V.C., Lai, E., 1996. Essential role of stromal mesenchyme in
kidney morphogenesis revealed by targeted disruption of Winged Helix transcription factor BF-
2. Genes Dev 10, 1467–1478. https://doi.org/10.1101/gad.10.12.1467
Heimann, H., Damato, B., 2010. Congenital vascular malformations of the retina and choroid. Eye 24,
459–467. https://doi.org/10.1038/eye.2009.310
Homan, K.A., Gupta, N., Kroll, K.T., Kolesky, D.B., Skylar-Scott, M., Miyoshi, T., Mau, D., Valerius, M.T.,
Ferrante, T., Bonventre, J.V., Lewis, J.A., Morizane, R., 2019. Flow-enhanced vascularization and
maturation of kidney organoids in vitro. Nat Methods 16, 255–262.
https://doi.org/10.1038/s41592-019-0325-y
Honeycutt, S.E., O'Brien, L.L., 2021. Injection of Evans blue dye to fluorescently label and image intact
vasculature. Biotechniques 70, 181–185. https://doi.org/10.2144/btn-2020-0152
Humphreys, B.D., Lin, S.-L., Kobayashi, A., Hudson, T.E., Nowlin, B.T., Bonventre, J.V., Valerius, M.T.,
McMahon, A.P., Duffield, J.S., 2010. Fate tracing reveals the pericyte and not epithelial origin of
myofibroblasts in kidney fibrosis. Am J Pathol 176, 85–97.
https://doi.org/10.2353/ajpath.2010.090517
Hurtado, R., Zewdu, R., Mtui, J., Liang, C., Aho, R., Kurylo, C., Selleri, L., Herzlinger, D., 2015. Pbx1-
dependent control of VMC differentiation kinetics underlies gross renal vascular patterning.
Development 142, 2653–2664. https://doi.org/10.1242/dev.124776
Jafree, D.J., Moulding, D., Kolatsi-Joannou, M., Perretta Tejedor, N., Price, K.L., Milmoe, N.J., Walsh, C.L.,
Correra, R.M., Winyard, P.J., Harris, P.C., Ruhrberg, C., Walker-Samuel, S., Riley, P.R., Woolf, A.S.,

Scambler, P.J., Long, D.A., 2019. Spatiotemporal dynamics and heterogeneity of renal lymphatics
in mammalian development and cystic kidney disease. Elife 8, e48183.
https://doi.org/10.7554/eLife.48183
Jones, C.A., London, N.R., Chen, H., Park, K.W., Sauvaget, D., Stockton, R.A., Wythe, J.D., Suh, W.,
Larrieu-Lahargue, F., Mukouyama, Y.-S., Lindblom, P., Seth, P., Frias, A., Nishiya, N., Ginsberg,
M.H., Gerhardt, H., Zhang, K., Li, D.Y., 2008. Robo4 stabilizes the vascular network by inhibiting
pathologic angiogenesis and endothelial hyperpermeability. Nat Med 14, 448–453.
https://doi.org/10.1038/nm1742
Kennedy, T.E., Serafini, T., de la Torre, JoséR., Tessier-Lavigne, M., 1994. Netrins are diffusible
chemotropic factors for commissural axons in the embryonic spinal cord. Cell 78, 425–435.
https://doi.org/10.1016/0092-8674(94)90421-9
Kobayashi, A., Mugford, J.W., Krautzberger, A.M., Naiman, N., Liao, J., McMahon, A.P., 2014.
Identification of a multipotent self-renewing stromal progenitor population during mammalian
kidney organogenesis. Stem Cell Reports 3, 650–662.
https://doi.org/10.1016/j.stemcr.2014.08.008
Larrivée, B., Freitas, C., Trombe, M., Lv, X., DeLafarge, B., Yuan, L., Bouvrée, K., Bréant, C., Toro, R.D.,
Bréchot, N., Germain, S., Bono, F., Dol, F., Claes, F., Fischer, C., Autiero, M., Thomas, J.-L.,
Carmeliet, P., Tessier-Lavigne, M., Eichmann, A., 2007. Activation of the UNC5B receptor by
Netrin-1 inhibits sprouting angiogenesis. Genes Dev. 21, 2433–2447.
https://doi.org/10.1101/gad.437807
Lejmi, E., Bouras, I., Camelo, S., Roumieux, M., Minet, N., Leré-Déan, C., Merkulova-Rainon, T., Autret,
G., Vayssettes, C., Clement, O., Plouët, J., Leconte, L., 2014. Netrin-4 promotes mural cell
adhesion and recruitment to endothelial cells. Vasc Cell 6, 1. https://doi.org/10.1186/2045-
824X-6-1
Lejmi, E., Leconte, L., Pédron-Mazoyer, S., Ropert, S., Raoul, W., Lavalette, S., Bouras, I., Feron, J.-G.,
Maitre-Boube, M., Assayag, F., Feumi, C., Alemany, M., Jie, T.X., Merkulova, T., Poupon, M.-F.,
Ruchoux, M.-M., Tobelem, G., Sennlaub, F., Plouët, J., 2008. Netrin-4 inhibits angiogenesis via
binding to neogenin and recruitment of Unc5B. Proceedings of the National Academy of
Sciences 105, 12491–12496. https://doi.org/10.1073/pnas.0804008105
Liebl, D.J., Klesse, L.J., Tessarollo, L., Wohlman, T., Parada, L.F., 2000. Loss of brain-derived neurotrophic
factor-dependent neural crest-derived sensory neurons in neurotrophin-4 mutant mice.
Proceedings of the National Academy of Sciences 97, 2297–2302.
https://doi.org/10.1073/pnas.040562597
Lindström, N.O., Guo, J., Kim, A.D., Tran, T., Guo, Q., De Sena Brandine, G., Ransick, A., Parvez, R.K.,
Thornton, M.E., Basking, L., Grubbs, B., McMahon, J.A., Smith, A.D., McMahon, A.P., 2018.
Conserved and Divergent Features of Mesenchymal Progenitor Cell Types within the Cortical
Nephrogenic Niche of the Human and Mouse Kidney. J Am Soc Nephrol 29, 806–824.
https://doi.org/10.1681/ASN.2017080890
Liu, J., Edgington-Giordano, F., Dugas, C., Abrams, A., Katakam, P., Satou, R., Saifudeen, Z., 2017.
Regulation of Nephron Progenitor Cell Self-Renewal by Intermediary Metabolism. J Am Soc
Nephrol 28, 3323–3335. https://doi.org/10.1681/ASN.2016111246
Liu, Y., Stein, E., Oliver, T., Li, Y., Brunken, W.J., Koch, M., Tessier-Lavigne, M., Hogan, B.L.M., 2004. Novel
Role for Netrins in Regulating Epithelial Behavior during Lung Branching Morphogenesis. Curr
Biol 14, 897–905. https://doi.org/10.1016/j.cub.2004.05.020
Lu, X., Le Noble, F., Yuan, L., Jiang, Q., De Lafarge, B., Sugiyama, D., Bréant, C., Claes, F., De Smet, F.,
Thomas, J.-L., Autiero, M., Carmeliet, P., Tessier-Lavigne, M., Eichmann, A., 2004. The netrin
receptor UNC5B mediates guidance events controlling morphogenesis of the vascular system.
Nature 432, 179–186. https://doi.org/10.1038/nature03080

Miyazaki, Y., Oshima, K., Fogo, A., Hogan, B.L., Ichikawa, I., 2000. Bone morphogenetic protein 4
regulates the budding site and elongation of the mouse ureter. J Clin Invest 105, 863–873.
https://doi.org/10.1172/JCI8256
Molema, G., Aird, W.C., 2012. Vascular Heterogeneity in the Kidney. Seminars in Nephrology,
Endothelium and the Kidney 32, 145–155. https://doi.org/10.1016/j.semnephrol.2012.02.001
Moore, M.W., Klein, R.D., Fariñas, I., Sauer, H., Armanini, M., Phillips, H., Reichardt, L.F., Ryan, A.M.,
Carver-Moore, K., Rosenthal, A., 1996. Renal and neuronal abnormalities in mice lacking GDNF.
Nature 382, 76–79. https://doi.org/10.1038/382076a0
Morizane, R., Lam, A.Q., Freedman, B.S., Kishi, S., Valerius, M.T., Bonventre, J.V., 2015. Nephron
organoids derived from human pluripotent stem cells model kidney development and injury.
Nat Biotechnol 33, 1193–1200. https://doi.org/10.1038/nbt.3392
Munro, D.A.D., Hohenstein, P., Davies, J.A., 2017. Cycles of vascular plexus formation within the
nephrogenic zone of the developing mouse kidney. Sci Rep 7, 3273.
https://doi.org/10.1038/s41598-017-03808-4
Nakagawa, N., Xin, C., Roach, A.M., Naiman, N., Shankland, S.J., Ligresti, G., Ren, S., Szak, S., Gomez, I.G.,
Duffield, J.S., 2015. Dicer1 activity in the stromal compartment regulates nephron
differentiation and vascular patterning during mammalian kidney organogenesis. Kidney Int 87,
1125–1140. https://doi.org/10.1038/ki.2014.406
O'Brien, L.L., Guo, Q., Bahrami-Samani, E., Park, J.-S., Hasso, S.M., Lee, Y.-J., Fang, A., Kim, A.D., Guo, J.,
Hong, T.M., Peterson, K.A., Lozanoff, S., Raviram, R., Ren, B., Fogelgren, B., Smith, A.D., Valouev,
A., McMahon, A.P., 2018. Transcriptional regulatory control of mammalian nephron progenitors
revealed by multi-factor cistromic analysis and genetic studies. PLoS Genet 14, e1007181.
https://doi.org/10.1371/journal.pgen.1007181
O'Brien, L.L., Guo, Q., Lee, Y., Tran, T., Benazet, J.-D., Whitney, P.H., Valouev, A., McMahon, A.P., 2016.
Differential regulation of mouse and human nephron progenitors by the Six family of
transcriptional regulators. Development 143, 595–608. https://doi.org/10.1242/dev.127175
Pallone, T.L., Zhang, Z., Rhinehart, K., 2003. Physiology of the renal medullary microcirculation. American
Journal of Physiology-Renal Physiology 284, F253–F266.
https://doi.org/10.1152/ajprenal.00304.2002
Pichel, J.G., Shen, L., Sheng, H.Z., Granholm, A.C., Drago, J., Grinberg, A., Lee, E.J., Huang, S.P., Saarma,
M., Hoffer, B.J., Sariola, H., Westphal, H., 1996. Defects in enteric innervation and kidney
development in mice lacking GDNF. Nature 382, 73–76. https://doi.org/10.1038/382073a0
Potente, M., Mäkinen, T., 2017. Vascular heterogeneity and specialization in development and disease.
Nat Rev Mol Cell Biol 18, 477–494. https://doi.org/10.1038/nrm.2017.36
Queisser, A., Seront, E., Boon, L.M., Vikkula, M., 2021. Genetic Basis and Therapies for Vascular
Anomalies. Circulation Research 129, 155–173.
https://doi.org/10.1161/CIRCRESAHA.121.318145
Reddi, V., Zaglul, A., Pentz, E.S., Gomez, R.A., 1998. Renin-expressing cells are associated with branching
of the developing kidney vasculature. J Am Soc Nephrol 9, 63–71.
https://doi.org/10.1681/ASN.V9163
Renier, N., Wu, Z., Simon, D.J., Yang, J., Ariel, P., Tessier-Lavigne, M., 2014. iDISCO: a simple, rapid
method to immunolabel large tissue samples for volume imaging. Cell 159, 896–910.
https://doi.org/10.1016/j.cell.2014.10.010
Ryan, A.R., England, A.R., Chaney, C.P., Cowdin, M.A., Hiltabidle, M., Daniel, E., Gupta, A.K., Oxburgh, L.,
Carroll, T.J., Cleaver, O., 2021. Vascular deficiencies in renal organoids and ex vivo kidney
organogenesis. Dev Biol 477, 98–116. https://doi.org/10.1016/j.ydbio.2021.04.009
Salminen, M., Meyer, B.I., Bober, E., Gruss, P., 2000. Netrin 1 is required for semicircular canal formation
in the mouse inner ear. Development 127, 13–22. https://doi.org/10.1242/dev.127.1.13

Sánchez, M.P., Silos-Santiago, I., Frisén, J., He, B., Lira, S.A., Barbacid, M., 1996. Renal agenesis and the
absence of enteric neurons in mice lacking GDNF. Nature 382, 70–73.
https://doi.org/10.1038/382070a0
Self, M., Lagutin, O.V., Bowling, B., Hendrix, J., Cai, Y., Dressler, G.R., Oliver, G., 2006. Six2 is required for
suppression of nephrogenesis and progenitor renewal in the developing kidney. EMBO J 25,
5214–5228. https://doi.org/10.1038/sj.emboj.7601381
Sequeira Lopez, M.L.S., Cherñavvsky, D.R., Nomasa, T., Wall, L., Yanagisawa, M., Gomez, R.A., 2003. The
embryo makes red blood cell progenitors in every tissue simultaneously with blood vessel
morphogenesis. American Journal of Physiology-Regulatory, Integrative and Comparative
Physiology 284, R1126–R1137. https://doi.org/10.1152/ajpregu.00543.2002
Sequeira-Lopez, M.L.S., Lin, E.E., Li, M., Hu, Y., Sigmund, C.D., Gomez, R.A., 2015a. The earliest
metanephric arteriolar progenitors and their role in kidney vascular development. Am J Physiol
Regul Integr Comp Physiol 308, R138–R149. https://doi.org/10.1152/ajpregu.00428.2014
Sequeira-Lopez, M.L.S., Nagalakshmi, V.K., Li, M., Sigmund, C.D., Gomez, R.A., 2015b. Vascular versus
tubular renin: role in kidney development. Am J Physiol Regul Integr Comp Physiol 309, R650–
R657. https://doi.org/10.1152/ajpregu.00313.2015
Serafini, T., Kennedy, T.E., Gaiko, M.J., Mirzayan, C., Jessell, T.M., Tessier-Lavigne, M., 1994. The netrins
define a family of axon outgrowth-promoting proteins homologous to C. elegans UNC-6. Cell 78,
409–424. https://doi.org/10.1016/0092-8674(94)90420-0
Smeyne, R.J., Klein, R., Schnapp, A., Long, L.K., Bryant, S., Lewin, A., Lira, S.A., Barbacid, M., 1994. Severe
sensory and sympathetic neuropathies in mice carrying a disrupted Trk/NGF receptor gene.
Nature 368, 246–249. https://doi.org/10.1038/368246a0
Srinivasan, K., Strickland, P., Valdes, A., Shin, G.C., Hinck, L., 2003. Netrin-1/neogenin interaction
stabilizes multipotent progenitor cap cells during mammary gland morphogenesis. Dev Cell 4,
371–382. https://doi.org/10.1016/s1534-5807(03)00054-6
Stanco, A., Szekeres, C., Patel, N., Rao, S., Campbell, K., Kreidberg, J.A., Polleux, F., Anton, E.S., 2009.
Netrin-1–α3β1 integrin interactions regulate the migration of interneurons through the cortical
marginal zone. Proceedings of the National Academy of Sciences 106, 7595–7600.
https://doi.org/10.1073/pnas.0811343106
Sun, K.L.W., Correia, J.P., Kennedy, T.E., 2011. Netrins: versatile extracellular cues with diverse functions.
Development 138, 2153–2169. https://doi.org/10.1242/dev.044529
Taguchi, A., Kaku, Y., Ohmori, T., Sharmin, S., Ogawa, M., Sasaki, H., Nishinakamura, R., 2014. Redefining
the In Vivo Origin of Metanephric Nephron Progenitors Enables Generation of Complex Kidney
Structures from Pluripotent Stem Cells. Cell Stem Cell 14, 53–67.
https://doi.org/10.1016/j.stem.2013.11.010
Takahashi, N., Lopez, M.L.S.S., Cowhig, J.E., Taylor, M.A., Hatada, T., Riggs, E., Lee, G., Gomez, R.A., Kim,
H.-S., Smithies, O., 2005. Ren1c homozygous null mice are hypotensive and polyuric, but
heterozygotes are indistinguishable from wild-type. J Am Soc Nephrol 16, 125–132.
https://doi.org/10.1681/ASN.2004060490
Takasato, M., Er, P.X., Chiu, H.S., Maier, B., Baillie, G.J., Ferguson, C., Parton, R.G., Wolvetang, E.J., Roost,
M.S., Chuva de Sousa Lopes, S.M., Little, M.H., 2015. Kidney organoids from human iPS cells
contain multiple lineages and model human nephrogenesis. Nature 526, 564–568.
https://doi.org/10.1038/nature15695
Tu, T., Zhang, C., Yan, H., Luo, Y., Kong, R., Wen, P., Ye, Z., Chen, J., Feng, J., Liu, F., Wu, J.Y., Yan, X.,
2015. CD146 acts as a novel receptor for netrin-1 in promoting angiogenesis and vascular
development. Cell Res 25, 275–287. https://doi.org/10.1038/cr.2015.15

Wälchli, T., Bisschop, J., Carmeliet, P., Zadeh, G., Monnier, P.P., De Bock, K., Radovanovic, I., 2023.
Shaping the brain vasculature in development and disease in the single-cell era. Nat Rev
Neurosci 1–28. https://doi.org/10.1038/s41583-023-00684-y
Wang, H., Copeland, N.G., Gilbert, D.J., Jenkins, N.A., Tessier-Lavigne, M., 1999. Netrin-3, a mouse
homolog of human NTN2L, is highly expressed in sensory ganglia and shows differential binding
to netrin receptors. J Neurosci 19, 4938–4947. https://doi.org/10.1523/JNEUROSCI.19-12-
04938.1999
Wilson, B.D., Ii, M., Park, K.W., Suli, A., Sorensen, L.K., Larrieu-Lahargue, F., Urness, L.D., Suh, W., Asai, J.,
Kock, G.A.H., Thorne, T., Silver, M., Thomas, K.R., Chien, C.-B., Losordo, D.W., Li, D.Y., 2006.
Netrins promote developmental and therapeutic angiogenesis. Science 313, 640–644.
https://doi.org/10.1126/science.1124704
Wu, H., Uchimura, K., Donnelly, E.L., Kirita, Y., Morris, S.A., Humphreys, B.D., 2018. Comparative Analysis
and Refinement of Human PSC-Derived Kidney Organoid Differentiation with Single-Cell
Transcriptomics. Cell Stem Cell 23, 869-881.e8. https://doi.org/10.1016/j.stem.2018.10.010
Yebra, M., Montgomery, A.M.P., Diaferia, G.R., Kaido, T., Silletti, S., Perez, B., Just, M.L., Hildbrand, S.,
Hurford, R., Florkiewicz, E., Tessier-Lavigne, M., Cirulli, V., 2003. Recognition of the Neural
Chemoattractant Netrin-1 by Integrins α6β4 and α3β1 Regulates Epithelial Cell Adhesion and
Migration. Developmental Cell 5, 695–707. https://doi.org/10.1016/S1534-5807(03)00330-7
813

<u>Figure Legends</u>

Figure 1. *Ntn1* is expressed and secreted by stromal progenitors in the kidney and deletion

results in kidney hypoplasia

(A) Light microscopy image of an E15.5 section showing the *in situ* hybridization expression

pattern of *Ntn1*. *Ntn1* expression is observed in the stromal progenitors (SP) surrounding the

nephron progenitors (NP). Dotted lines demarcate NP niches. Scale bar=50 μm. (B)

Immunostaining of E11.5 kidney sections shows that netrin-1 (magenta) is secreted by the

stromal progenitors surrounding the Six2+ (green) nephron progenitors. Inset shows a higher

magnification of the netrin-1 signal adjacent to the nephron progenitors. Scale bar=50 μm. (C)

Immunofluorescence images showing netrin-1 (magenta) signal in the stromal progenitors (SP)

and adjacent to the Six2+ nephron progenitors (NP, green) (left panel). Secreted netrin-1 signal

is also observed in the ureteric tip region. Genetic deletion of *Ntn1* in the *Foxd1*+ stromal

population results in a significant reduction in netrin-1 signal (right panel). Scale bar=25 µm. (D)

Kidney hypoplasia is observed at various stages of development. Shown are micrographs of

whole urogenital systems at E15.5 and P0. Insets for P0 show the same kidneys micro dissected

and side-by-side for size comparison. (E) Kidney length was measured in Fiji imaging software at

E13.5 and E15.5, with the size reduction becoming statistically significant by E15.5. Columns

and error represent mean and standard deviation (SD), respectively, with significance derived

from Students' T-test. Respective p-values displayed on graphs.

Figure 2. Loss of *Ntn1* delays the cessation of nephrogenesis but is not due to signaling

through nephron progenitor expressed *Unc5c* or differences in progenitor dynamics

A) Immunostaining of whole metanephric kidneys at E11.5 shows that metanephric kidney

development is initiated normally in *Ntn1* conditional knockouts, with the expected T-shaped

ureteric bud (cytokeratin, magenta) surrounded by condensed nephron progenitors (Six2,

green). Scale bar=200 µm. B) Nephron progenitor niches (Six2, green) appear normally

organized at E15.5 in surface views of wholemount immunostained kidneys. Scale bar=30 µm.

C) Wholemount immunostains of P4 kidneys show the persistence of Six2+ nephron progenitors

(green) on the surface of the kidney in *Ntn1* conditional knockouts. Scale bar=30 µm. D) *In situ*

hybridization showing the expression of the netrin-1 receptor *Unc5c* in the nephron progenitors

(black arrowhead), podocytes (yellow arrowhead), and smooth muscle layer of the ureter

(green arrowhead) at E15.5. Scale bar=50 µm. E) Light microscopy images of whole kidneys at

P0 showing no difference in kidney size of *Unc5c* knockout kidneys compared to controls. F)

Wholemount immunostaining and surface views show the normal organization of nephron

progenitor niches in *Unc5c* knockouts at P0 (Six2-green, cytokeratin-magenta). Scale bar=40

µm. G) Example of a sectioned kidney labeled with EdU (white) and immunostained with Six2

(green) and E-cadherin (Ecad, magenta) to label the nephron progenitors and collecting duct

system, respectively. Inset shows a high magnification image of a niche with dotted lines

highlighting the extent of what was considered either a nephron progenitor (NP) niche or

ureteric bud (UB) tip niche for quantification purposes. Scale bar=200 µm. H) Quantification of

Six2+ nephron progenitors or Ecad+ ureteric tip cells at E13.5, E15.5, and P0 show no significant

differences in total cell numbers except for Ecad+ cells at E13.5. Bars represent mean;

significance was determined using Students t-test. I) Quantification of the % of EdU+ Six2+

nephron progenitors or EdU+ Ecad+ ureteric tip cells reveals no significant difference at E13.5,

E15.5, or P0 between $Foxd1^{GC/+}$; $Ntn1^{fl/fl}$ kidneys or controls. Bars represent mean; significance

was determined using Students t-test.

Figure 3. Stromal deletion of *Ntn1* results in aberrant neurovascular patterning

A) *In situ* hybridization on E15.5 kidney sections shows the expression of the netrin-1 receptor

Unc5b in the endothelium of the kidney (black arrowhead and inset of glomerular endothelium,

left panel), the secondary branches/generations of the collecting duct (yellow arrowhead, left

panel), and ureter (right panel). Scale bars=50 µm B) Wholemount immunostain of an E15.5

wildtype kidney show the tracking of Tuj1+ (cyan) axonal projections with the renal vasculature

(CD31, red). Far right panel shows an independent close-up image of a neurovascular bundle.

Merge scale bar=300 µm, close-up scale bar=50 µm. C) Wholemount immunostains show the

organization of the nerves and vasculature at E15.5 in *Nnt1* conditional knockouts and controls.

The nerves (Tuj1, cyan) track with the endothelium (CD31, red) in all genotypes. White

arrowheads highlight the abnormal patterning observed in *Ntn1* conditional mutants such as

looping and ectopic neurovascular bundles. Scale bar=300 μm. D) Depiction of tracing (white

lines) of the neurovascular bundles (left panel). Tracing of five control kidneys (top panel,

colored lines) show predictable patterns of 5 main lateral arterial branches which can be

overlayed. *Ntn1* conditional knockout kidneys (bottom panel, colored lines) show stochastic

patterning with the absence of any predictable patterning. Scale bar=300 μm. E) Wholemount

immunostains of control, *TrkA$^{-/-}$*, and *Foxd1$^{GC/+}$*; *Ntn1$^{fl/fl}$*; *TrkA$^{-/-}$* kidneys at E16.5 show the

organization of nerves (Tuj1, cyan), endothelium (CD31, red), and smooth muscle cells (SMA,

green). Innervation of the kidney is reduced in *TrkA* knockouts, but abnormal patterning

persists with vasculature not associated with axons (arrowheads) still showing abnormal

patterning, suggesting the endothelium is the primary responder to netrin-1 guidance cues.

Scale bar=300 μm.

Figure 4. *Ntn1* conditional knockouts have normal endothelial metrics but reduced smooth

muscle coverage throughout development

A) Quantitation of CD31+ endothelial metrics including the number of branches, branch points,

and total endothelium (total additive length, μM) at E15.5 shows no statistical differences

between controls and *Ntn1* conditional knockouts. Significance was calculated with Students t-

test, p<0.05 was considered significant, columns represent mean. Error bars = SD. B)

Wholemount immunostaining of E15.5 kidneys with smooth muscle actin (SMA, green) in

addition to the axonal (Tuj1, cyan) and endothelial (CD31, red) markers show a qualitative delay

in smooth muscle cell association with the endothelium (white arrowheads, zoom) suggesting a

potential delay is vascular maturation. Scale bars=300 μm. C) Wholemount immunostains of

E15.5 and P0 kidneys with SMA (white) show a qualitative reduction on SMA vascular coverage

in *Ntn1* conditional mutants. The injection of Evans Blue dye at 2 months to label the adult

kidney vasculature shows a well-developed arterial tree with some remnants of abnormal

patterning such as elongated branches in *Ntn1* conditional knockouts. Red dotted lines overlay

normal (top) and abnormal (bottom) branching. E15.5 scale bar=300 μm, P0 scale bar=500 μm,

2-month scale bar=1000 μm. D) Quantitation of SMA+ arterial metrics including the number of

branches, branch points, and total vasculature length (μM) at E15.5 and P0 show a deficit in

smooth muscle metrics suggesting a delay in arterial differentiation. Quantitation of Evans blue

labeled arterial vascular metrics at 2 months reveals that *Ntn1* conditional knockouts have

attained levels similar to controls. Significance was calculated with Students t-test, $p<0.05$ was

considered significant, individual p-values are displayed on the respective graphs, while

columns represent mean. Error bars = SD. Demarcation between P0 and two months denotes

the differences in labeling, with early stages labeled by SMA and 2 months perfused with Evans

blue.

Figure 5. Gene expression analyses support a delay in vascular maturation in *Ntn1* conditional

knockouts

A) Gene ontology (GO) analysis of genes upregulated (top) or B) downregulated (bottom) in

Ntn1 conditional knockouts. Shown are the GO Biological Processes enriched in both categories.

Genes were considered up- or downregulated with a >1.2 fold-change. In the insets are

examples of genes that are upregulated (top) or downregulated (bottom). C) Schematic of the

role of netrin-1 in guiding vascular patterning and maturation during development.

Figure 1

A E15.5 — *Ntn1* — SP, NP, NP

B Wildtype — E11.5 — Netrin-1, Six2

C E15.5 — Control / *Foxd1^{GC/+}; Ntn1^{fl/fl}* — Netrin-1, Six2 — SP, NP

D Control / *Foxd1^{GC/+}; Ntn1^{fl/fl}* — E15.5, P0

E Kidney length — p=0.003 — Millimeters — 13.5, E15.5 — Control, *Foxd1^{GC/+}; Ntn1^{fl/fl}*

Figure 2

Figure 3

Figure 4

Figure 5

A

Upregulated in *Foxd1GC/+; Ntn1fl/fl* (>1.2-fold *Mutant vs Het*)

GO Biological Process
GO:0007155:cell adhesion
GO:0007156:homophilic cell adhesion via plasma membrane adhesion molecules
GO:0009653:anatomical structure morphogenesis
GO:0098609:cell-cell adhesion
GO:0032502:developmental process
GO:0048731:system development
GO:0007275:multicellular organism development
GO:0030154:cell differentiation
GO:0007399:nervous system development
GO:0009888:tissue development
GO:0009887:animal organ morphogenesis
GO:0016477:cell migration
GO:0072359:circulatory system development
GO:0050793:regulation of developmental process
GO:0048468:cell development

Genes
Nrp2
Flt1
Notch1
Unc5b
Bmpr2
Tgfb1

-log10(p-value)

B

Downregulated in *Foxd1GC/+; Ntn1fl/fl* (>1.2-fold *Mutant vs Het*)

GO Biological Process
GO:0006412:translation
GO:0006364:rRNA processing
GO:0006397:mRNA processing
GO:0008380:RNA splicing
GO:0090131:mesenchyme migration
GO:0006937:regulation of muscle contraction
GO:0033700:phospholipid efflux
GO:0034384:high-density lipoprotein particle clearance
GO:0003009:skeletal muscle contraction
GO:0061621:canonical glycolysis
GO:0043484:regulation of RNA splicing
GO:0034380:high-density lipoprotein particle assembly
GO:0033344:cholesterol efflux
GO:0002181:cytoplasmic translation
GO:0006782:protoporphyrinogen IX biosynthetic process

Genes
Acta2
Myh11
Gpx1
Eno1
Aldoa

-log10(p-value)

C