Longitudinal Survey of Astrovirus infection in different bat species in Zimbabwe: Evidence of high genetic *Astrovirus* diversity

Chidoti Vimbiso[1], De Nys Hélène[2,3,4], Abdi Malika[5], Mashura Getrude[1], Pinarello Valérie [1,2,3,4], Chiweshe Ngoni[4], Matope Gift[1], Guerrini Laure[2,3], Pfukenyi Davies[1,6], Cappelle Julien [2,3], Mwandiringana Ellen[1], Missé Dorothée[2], Gori Elizabeth[1,7], Bourgarel Mathieu[2,3,4] and Liégeois Florian [1,5,*].

[1]Faculty of Veterinary Science, University of Zimbabwe, P.O. Box MP167 Harare, Zimbabwe

[2]ASTRE, Univ Montpellier, CIRAD, INRAE, Montpellier, France.

[3]CIRAD, UMR ASTRE, Montpellier, France.

[4] CIRAD, UMR ASTRE, Harare, Zimbabwe

[5]MIVEGEC, University of Montpellier, IRD, CNRS, 34394 Montpellier, France

[6]Faculty of Animal and Veterinary Sciences, Botswana University of Agriculture and Natural Resources, Private Bag 0027, Gaborone, Botswana

[7]Department of Biochemistry, Molecular Biology and Genetics, University of Rwanda, Rwanda

*Corresponding Author

Florian Liegeois

University of Zimbabwe, Faculty of Veterinary Science, Veterynay Biosciences Departement

Room 2F7, P.O. Box MP167 Harare, Zimbabwe.

Florian.liegeois@ird.fr

Abstract

Astroviruses (AstVs) have been discovered in over 80 animal species including diverse bat species and avian species. A study on *Astrovirus* circulation and diversity in different insectivorous and frugivorous chiropteran species roosting in trees, caves and building basements was carried out at 11 different sites across Zimbabwe. Pooled and individual faecal sampling methods were used for this study, collection date ranged from June 2016 to July 2021. In two sites, Magweto and Chirundu, sampling was carried out at monthly intervals from August 2020 to July 2021. Astroviruses and bat mitochondrial genes were amplified using pan-AstVs and CytB /12S RNA PCR systems respectively. Phylogenetic analysis of the *RdRp Astrovirus* sequences revealed a high genetic diversity of astroviruses. All the bat astroviruses tested in this study clustered with the *Mamastrovirus* genus. Two distinct groups of the amplified sequences were identified. One group comprised of sequences isolated from *Hipposideros, Rhinolophus* and *E. helvum spp* clustered with Human Astrovirus strains: *HuAstV* types1-6, *HuAstV*-MLB1-3 and *HuAstV*-VA-1. A second group comprised of the majority of the sequences clustered with different strains of Bat AstVs. Results from the longitudinal study at Magweto and Chirundu showed an overall AstV prevalence of 13.7% and 10.4% respectively. Peaks of AstV infection at Chirundu coincided with the period when juveniles are 4-6 months old. Coinfection of bats with CoVs and AstVs at Magweto and Chirundu sites was 2.6% and 3.5% respectively.

Introduction

Astrovirus are single-stranded positive sense RNA (+ssRNA) viruses. They are non-enveloped with an icosahedral morphology and a genome length of approximately 6.2 to 7.7 Kb [1]. They infect a wide variety of both domestic and wild marine and terrestrial mammals, including humans (*Mamastrovirus*) as well as avian hosts (*Avastrovirus*) [2]. Recently *Astrovirus* infections in fish and insects have been reported [3,4]. Bats and wild birds are considered natural reservoirs of astroviruses [5]. Astroviruses have been found to occur in over 80 non-human species including a diverse number of bat species from Europe, Africa, America and Asia. Human astroviruses (*HuAstVs*) have been identified as causal agents of acute viral gastrointestinal illness worldwide particularly in children, immunocompromised people and the elderly [3–5]. Beyond this well-known clinical manifestation of *Astrovirus* infections, neurovirulent *Astrovirus* infections have also been reported in both humans and domestic animals [6,7]. To note, in humans, the majority of *Astrovirus*-associated encephalitis or meningitis were reported in immunocompromised people [8–10]. According to the International Committee on Taxonomy of Viruses (https://talk.ictvonline.org), *Mamastrovirus* are classified within 19 recognized species, *MAstV*-1 to -19, and two genogroups GI and GII. All classic and novel Human Astrovirus belong to four different species, *MAstV*-1, *MAstV*-6, *MAstV*-8 and *MastV*-9 [2,11,12]. Astroviruses do not seem to have a common reservoir [12]. However, numerous cross-species transmissions of *Astrovirus* have been documented [4]. More particularly, certain *HuAstV* strains are closely related to rodent, feline, mink, ovine and porcine *Astrovirus* [2,4,12].

To date, no human *AstV* strains have been associated with Bat *AstV*. Bat *Astrovirus* belong to *MAstV*-12, and *MastV*-14 to -19 species. Bat-borne viruses represent an extensive research field owing to the plethora of viruses carried out by the Chiropterans. Bats are known to be persistently infected by astroviruses [13]. This is inclusive of many species of insectivorous bats which harbor these viruses [14]. To date, a high diversity of Bat *Astroviruses* from different bat families has been reported worldwide [9,15]. In Europe a variety of bat species belonging to the Yangochiroptera suborder have been discovered to harbor several astroviruses [16]. In Africa, bat astroviruses were reported in Egypt, Gabon, Madagascar, Kenya and Mozambique [5,15,17–19]. Astroviruses are known to occur in bat species with high prevalence and exceedingly high genetic diversity whereby infections in these hosts are minimally pathogenic [20,21]. Despite all these studies, large gaps still exist to map the extent of existence of these viruses in bats' populations [22].

Astrovirus prevalence and detection in bat colonies seemed rather correlated to abiotic factor factors such as seasons and year of sampling than biotic factors such as sex and reproductive status or yet viral co-infection [23]. Viral shedding in bats occurs in spatial and temporal pulses that can drive spillover to other animals or humans [24] Shedding pulses in bats determine varying degrees in the

prevalence of viruses where if they do not occur, viruses are rarely detected or at low prevalence and

when they occur, higher prevalence is detected [24].

In this study we enlarge the spectrum of bat *Astroviruses* knowledge in Africa by identifying and

analyzing Bat *Astroviruses* from different bat species colonies according to season in Zimbabwe.

Material and Methods

Sampling approaches and sites

Two different approaches were followed in this study: bat community sampling and individual bat

sampling.

Bat community pooled sampling

Between February 2016 and December 2020, faecal samples from both insectivorous and frugivorous

bat species were respectively collected in different sites including caves, an ancient mine and trees in

Zimbabwe (Figure 1, Table 1).

Figure 1: The study sites highlighted on the map show the sampling areas for insectivorous and frugivorous

bats colonies across Zimbabwe. Red circles represent community approach study sites; yellow circles represent

longitudinal survey sites (individual approach).

Table 1: Number of Astrovirus positives per each site/sampling year in pooled faecal samples from frugivorous and insectivorous bats at different 14 sites.

Sampling Sites	type of roost	Date of collect	Bat Species	Number of collected tubes	Number of Tested pools	Number of positive pools
Magweto Cave	Cave	June 2016	*Hipposideros spp.*	24	8	5
Bashongwe Cave	Cave	March 2018	*Rhinolophus and Hipposideros spp.*	59	7	0
		July 2018		57	20	0
		December 2018		135	45	0
Mabura Cave	Cave	June 2016	*Hipposideros spp.*	36	12	6
		February 2017	*Hipposideros spp.*	22	7	0
Mugunza Cave	Cave	April 2017	*Roussetus aegyptus*	50	13	0
Honde Valley						
Machiri		August 2017	*Eidolon helvum*	25	9	0
Muwomba		November 2017	*Eidolon helvum*	66	22	0
Nyamajo		November 2017	*Eidolon helvum*	58	19	0
Nyamhosva		November 2017	*Eidolon helvum*	35	11	0
Nyamajo	Trees	January 2018	*Eidolon helvum*	17	6	1
Machiri		January 2018	*Eidolon helvum*	7	2	2
Muwomba		January 2018	*Eidolon helvum*	8	2	0
Nyamhosva		January 2018	*Eidolon helvum*	28	10	1
Nyamajo		February 2020	*Eidolon helvum*	92	92	0
Chirundu	Baobab	December 202	*Rhinolophus spp.*	21	21	1
Total				**740**	**306**	**16**

All these sites were chosen according to the presence of bat colonies and the existence of anthropic activities. For instance, the selected caves and ancient mine selected are regularly visited by local people to collect bat guano which is used as fertilizer, and to hunt bats for consumption. Frugivorous bats roosting sites in trees were close to maize crops or/and fruit tree cultures.

All sites, except for three were visited at different times during the sampling period (Table 1). The same sampling method was used at all sites every session as previously described [25]. Briefly, two square meters of plastic sheets were laid down at each site/per sampling session, underneath the bat colonies and left overnight. Faeces were collected from each plastic sheet at a rate of ≈ 6 grams of pooled faeces in a 15 ml tube with 6 ml of homemade RNA stabilization solution (*https://protocol-online.org/*). Back in the laboratory, samples were stored at -80°C until analysed.

Bat individual sampling

Individual bat samples which had already been collected from two study sites for a study on coronaviruses by Chidoti *et al*. were used in this study (Figure 1) [26]. These two sites, one cave (Magweto) and one building basement (Chirundu Farm) had been visited from August 2021 to July 2022 on a monthly basis (Figure 1, Table 2) [26]. Unfortunately, this study was conducted during the COVID 19 crisis and we were not able to access the study sites every month as planned owing to the

imposed lock downs. Faecal samples had been collected by placing two square meters plastic sheets underneath the bat colonies. Only one faecal dropping (clean and not contaminated by other faeces or urine) per 20 cm^2 was collected, assuming it represented one individual. Faeces were conserved individually in a 1.5 ml tube filled up with 0.5 ml of home-made RNA stabilization solution (*https://protocol-online.org/*) and stored at -80°C before further laboratory analyses.

Table 2: Prevalence of Astroviruses and Coinfection with Coronaviruses and confidence intervals (CIs) per month at both Magweto and Chirundu sites in faecal samples from individual sampling of insectivorous bat communities

Site	Reproduction cycle	Month sampled	No of samples tested	No of Astrovirus positives	Prevalence (%) + CI (95%)	No of coinfection CoV and AstV	Prevalence (%) + CI (95%)
Chirundu	**Non-gestation**	August, 2020	153	0	0.0 (0.0-0.2)	0	0.0 (0.0-0.2)
Chirundu	**Pregnancy**	October, 2020	297	3	1.0 (0.3-2.9)	0	0.0 (0.0-1.3)
Chirundu	**Parturition**	November, 2020	241	3	1.2 (0.4-3.6)	1	0.4 (0.05-2.3)
Chirundu	**Lactation**	December, 2020	159	16	10.1 (6.3-15.7)	7	4.4 (2.1-8.8)
Chirundu	**Weaning**	February, 2021	170	29	17.1 (12.1-23.4)	18	10.6 (6.8-16.1)
Chirundu	**4-6 Month juveniles**	March, 2021	240	51	21.3 (16.5-26.8)	19	7.9 (5.1-12.0)
Chirundu	**Non-gestation**	May, 2021	243	47	19.3 (14.8-24.7)	9	3.7 (1.9-6.8)
Chirundu	**Non-gestation**	July,2021	225	30	13.3 (9.5-18.4)	9	4.0 (2.1-7.4)
	Overall prevalence		**1728**	**179**	**10.4 (9.0-11.8)**	**63**	**3.6 (3.8-5.8)**
Magweto	**Non-gestation**	September, 2020	348	18	5.2 (3.3-8.0)	1	0.3 (0.05-1.6)
Magweto	**Pregnancy**	October, 2020	257	46	17.9 (13.7-23.0)	7	2.7 (1.3-5.5)
Magweto	**Parturition**	November, 2020	228	31	13.6 (9.7-18.6)	3	1.3 (0.4-3.8)
Magweto	**4-6 Month juveniles**	March, 2021	242	46	19.0 (14.5-24.4)	20	8.3 (5.4-12.4)
Magweto	**Non-gestation**	April, 2021	242	26	10.7 (7.4-15.2)	6	2.5 (1.1-5.3)
Magweto	**Non-gestation**	June, 2021	242	47	19.4 (14.9-24.9)	3	1.2 (0.4-3.6)
	Overall prevalence		**1559**	**214**	**13.7 (12.1-15.5)**	**40**	**2.6 (1.8-3.5)**

Periods of gestation (pregnancy), parturition, lactation, weaning and presence of 4-6 months old juveniles had been determined based on observations (from captures and observation of the roosting bats) combined with literature, as already reported in Chidoti et *al.* [26]. The reproductive season at Chirundu and Magweto site had been observed to begin in September to February for the predominant bat species. The different insectivorous bat families observed at both sites had been found to be synchronous regarding their reproductive cycles, and consensus reproduction periods had been determined based on the literature and observations [26, 27].

Nucleic acid extraction and RT-PCR.

Community samples.

Nucleic acids extraction was done from all faecal samples as previously described [26]. Briefly, biological material (Faeces) of three or four sample tubes collected from the same plastic sheet were pooled and transferred in a 50 ml tube with 20 ml of PBS 1X then vigorously mixed. Tubes were centrifuged at 4500 rpm for 10 min. The supernatant was first filtered using gauze in order to eliminate faecal matter and transferred in fresh tubes then re-centrifuged at 4500 rpm for 10 min. The

supernatant was filtered through a 0.45 µm filter to remove eukaryotic and bacterial sized particles.

Seven millilitres of filtered samples were centrifuged at 250,000 g for 2.5 h at 4°C. The pellets were

re-suspended in 600 µl H$_2$0 molecular grade and 150 µl were used to extract RNA and DNA using

NucleoSpin® RNA Kit (Macherey-Nagel, Hoerdt, France) according to the manufacturer's protocol.

Individual samples

Nucleic acids were extracted from 200 µl of faecal samples preserved in 0.5 ml RNA stabilization

solution using 5X MagMax Pathogen RNA/DNA Kit (ThermoFisher Scientific, Illkirch-

Graffenstaden, France), as already described in [26]. The faeces were vortexed vigorously (30Hz)

using Retsch MM400 Tissue lysser for 5 min to fully homogenise and mix the faecal particles,

followed by centrifugation at 16000 g for 3 min to fully separate the supernatant from the faecal

debris. A volume of 130µl of the supernatant was used for the isolation and purification stage of the

nucleic acids using Mag Max extraction kit with the automatic KingFisher Duo Prime Purification

System extractor (ThermoFisher Scientific, Illkirch-Graffenstaden, France). A final volume of 80µl

of eluted RNA/DNA was stored at -80◦C.

Bat species identification

All positive *Astrovirus* bat species were identified by sequencing *cytochrome b* [28]. Sequences were

then compared to available bat sequences in the GenBank database using *Basic Local Alignment*

Search Tool (BLAST) program.

Astrovirus detection

Reverse Transcription (RT) was done using random hexamers were done on 5µl of RNA sample

template using 1µl random hexamers, 0.5µl Oligo dT primer, 0.4µl of dNTPs (10mM) (ThermoFisher

Scientific, Illkirch-Graffenstaden, France) and 5.5µl molecular grade water incubated at 65°C for 5

min. This was followed by addition of 4µl of Buffer 5X, 2µl of 0.1M DTT (M-MLV Reverse

Transcriptase, Invitrogen, ThermoFisher Scientific) and 1µl of RNAse OUT, incubated at 37°C for 2

min. A volume of 1µl of M-MLV (M-MLV Reverse Transcriptase, Invitrogen, ThermoFisher

Scientific, Illkirch-Graffenstaden, France) reverse transcriptase was added to the mixture followed

by incubation at 25ºC for 10 min, 37ºC for 50 min and 70ºC for 15 min. The cDNA obtained was then

used to partially amplify the *Astrovirus RNA-dependent-RNA polymerase* gene (*RdRp*) by using a

semi-nested Pan-Astrovirus PCR system developed by Chu et *al* [13]. Positive PCR products (422

bp) were gel-agarose purified (Geneclean Turbo Kit, MP Biomedicals, Illkirch-Graffenstaden, France)

and then sequenced in both 5' and 3' directions (LGC, Berlin, Germany) by using Sanger method.

For the community approach, purified PCR products were cloned by using Topo PCR Cloning kit

according to the manufacturer's protocol (ThermoFisher Scientific, Illkirch-Graffenstaden, France).

Ten clones per PCR product were sequenced in both 5' and 3' direction using the Sanger method

(Eurofins, Germany).

Phylogenetic analyses

Overlapping sequences were assembled into contiguous sequences using Geneious software package

V. 2021.2.2 (Biomatters Ltd, Auckland, New Zealand). Partial non-concatenated nucleic acid

sequences of the new *Astrovirus* were aligned using MEGA 7 [29], with minor manual adjustments.

Sites that could not be unambiguously aligned and divergent regions were excluded from subsequent

analyses. Phylogenies were inferred using Maximum Likelihood (ML) method implemented in

PhyML [30]. The suited evolution model was selected by Akaike's Information criterion (AIC) using

Topali software [31]. The reliability of branching orders was tested using the bootstrap approach

(1000 replicates) [32] and the GTR + F+ I substitution model was determined as the best suited

evolution model.

Temporal variations of Astrovirus prevalence and bat reproductive phenology

In the same way as in Chidoti et *al*, the prevalence of astrovirus infection was calculated at the

community level for small insectivorous bat species from the two longitudinal study sites, Magweto

and Chirundu farm [26]. The proportion of RNA AstV-positive samples were estimated per month

and site with 95% confidence intervals (CI) using Wilson score test [33]. A descriptive analysis of

AstVs prevalence per month and corresponding reproductive season at each site was done.

The influence of the different phases of the bat reproductive cycle (periods of pregnancy, parturi-

tion/lactation, weaning, weaned juveniles of 4 to 6 months old) and of *Coronavirus* infection/shed-

ding on the prevalence of astroviruses was tested by running a generalized linear mixed model

(GLMM) for each site (Magweto and Chirundu) as described in Chidoti et *al* [26]. Parturition and

lactation were analysed as one season because the observations did not allow the separation of the

two periods as they were interlinked. The response variable with a binomial distribution was the

Astrovirus PCR result of the samples, and the explanatory variables with fixed effects were the dif-

ferent phases of the reproduction cycle. Given *Coronavirus* PCR results on the same samples were

available through our previous study (see below), *Coronavirus* status was also integrated into the

model as fixed variable to investigate any effect of co-infection on *Astrovirus* infection/shedding. The

reproductive phases were coded as 1 if it was during the corresponding reproduction phase and 0 if it

was not. A session identification code was included as a random effect to control for repeated

measures from the same sampling session to account for clustered samples collected.

Astrovirus and Coronavirus co-infection

Coronavirus data, produced by Chidoti et *al* [26], on the same sample sets were used in this study to assess the *Astrovirus* and *Coronavirus* co-infection in bat communities from Magweto and Chirundu sites and a descriptive analysis of prevalences was carried out.

Results

Astrovirus detection in pooled sampling sites

The sampling at these sites varied from June 2016 to December 2020 (Table 1). At Magweto 24 sampled were collected. In Bashongwe collection was carried out in March, July and December 2018, and 251 samples were collected. In Mabura cave site, sampling was done once in 2016 and 2017 and a total of 58 samples were collected. In Mugunza cave site, sampling was done once in 2017 and 50 samples were collected. In the Honde Valley site, 336 samples were collected under 9 *Eidolon helvum* roost trees (Table 1). Five (62.5%) pooled samples from Magweto site were positive for AstVs out of 8 pooled samples analysed and all 72 pooled samples from Bashongwe cave were negative for AstV (Table 1). Six (31.6%) out of 19 pooled samples from Mabura cave tested for AstVs were positive and all of the 13 samples from Mugunza tested for AstVs were negative (Table 1). In Honde Valley, 173 pooled samples were tested for AstVs and only four (2.3%) were positive (Table 1). Only one (4.8%) out of the 21 samples collected in Chirundu Baobab tree site was positive for AstV.

Prevalence and seasonality of Astroviruses at Chirundu and Magweto site

We analysed 1559 samples from insectivorous bats from the Magweto cave site and a total of 1728 faecal samples from bats at the Chirundu farm site (Table 2).The overall prevalence of *Astrovirus* at bat community level of Chirundu was 10.4% [95% CI: 9.0-11.8] (179 positives out of the 1728 samples). The highest prevalence of 21.3% [95% CI: 16.5-26.8] was observed in March 2021 and corresponded to the 4-6 months old juvenile period (Table 2). The prevalence of *AstV* increased from December 2020 to March 2021, which corresponded to the lactation, weaning and 4-6 months juvenile periods. The lowest prevalence was observed in August (0%); October (1% [95% CI: 0.3-2.9]) and November (1.2% [95% CI: 0.4-3.6]), which correspond to the transition between non-gestation and gestation period, as well as the beginning of the parturition period (Table 1, Figure 2a).

Figure 2: Bat community *Astrovirus* prevalence per month at Chirundu site. **Fig 2a & 2b** show estimation of the astrovirus prevalence (with CI 95%) at Chirundu and Magweto sites respectively

246   for the months sampled. The graphs were plotted with y-axes as the number of AstVs positive samples,

247   and the prevalence of AstVs and x-axis as the sampling date and corresponding reproductive season.

Figure 2a

	14/08/20	14/09/20	15/10/20	15/11/20	15/12/20	15/01/21	15/02/21	14/03/21	15/04/21	15/05/21	15/06/21	14/07/21
	Non-gestation		Pregnancy	Parturition	Parturition & Lactation	Lactation	Weaning	4-6_months Juveniles		Non-gestation		
● Chirundu AstV Prevalence (%)	0		1	1.2	10.1		17.1	21.2		19.3		13.3
Number of AstVs positives	0		3	3	16		29	51		47		30
Total number of samples analysed	153		297	241	159		170	240		243		225

Date/Reproductive

In Magweto site, the overall prevalence of Astroviruses for the insectivorous bat community was 13.7%

[95% CI:12.1-15.5] (214 positives out of 1559 samples). There was no clear pattern for the prevalence

of AstVs according to reproductive seasons in this site (Table 2, Figure 2b).The highest prevalence

of 19.4% [95% CI: 14.9-24.9] was observed in June 2021 during the non-gestation period , and very

close prevalences observed during the pregnancy (17.9% [95% CI: 13.7-23]) and 4-6 months old

juvenile (19% [95% CI: 14.5-24.4]) periods (Table 2). Lower prevalence was observed in September

at the end of the non-gestation period (5,2% [95% CI: 3.3-8]).

Figure 2b

	14/08/20	14/09/20	15/10/20	15/11/20	15/12/20	15/01/21	15/02/21	14/03/21	15/04/21	15/05/21	15/06/21	15/07/21
	Non-gestation		Pregnancy	Parturition	Parturition & Lactation	Lactation	Weaning	4-6_months Juveniles		Non-gestation		
Magweto AstV Prevalence (%)		5.2	17.9	13.6				19	10.7		19.4	
Number of AstVs positives		18	46	31				46	26		47	
Total number of samples analysed		348	257	228				242	242		242	

Date/Reproductive

Results from the GLMM didn't show an effect bats seasonal reproduction periods on detection of RNA-AstVs positive samples at Magweto site (Table 3a). For Chirundu site, the GLMM showed significantly higher probability of being positive to astrovirus during the period of weaned 4-6 months old juveniles (odds ratio (OR) = 5.088, 95% CI = 1.33- 36.32, p= 0.016), and significantly lower probability of astrovirus positivity during the pregnancy period (odds ratio (OR) = 0.136, 95% CI = 0.03- 0.81, p=0.006) (Table 3b).

Table 3: Results of the GLMM with the following explanatory variables (a/ Magweto, b/Chirundu): PCR_Pan-*Coronavirus*, Pregnancy, Weaning, Weaned juveniles 4 to 6 months. With references OR odds ratio, CI95 95% confidence interval.

a/

Variables	Odds ratios	p-value	OR CI95
Intercept	0.235	0.0005	0.088-0.627
PCR_Pan-*Coronavirus*	1.238	0.290	0.825-1.825
Pregnancy	0.473	0.145	0.138-1.548
Weaned_juveniles 4 to 6 months	0.678	0.448	0.202-2.256

272 *b/*

Variables	Odds ratios	p-value	OR CI95
Intercept	0.05	6.68e-09	0.010- 0.121
PCR_ Pan-*Coronavirus*	1.116	0.536	0.785 -1.574
Pregnancy	0.4	0.006	0.027-0.810
Weaning	3.91	0.094	0.721-39.593
Weaned_juveniles 4 to 6 months	5.09	0.016	1.333-36.322

Prevalence and seasonality of coinfection with Astroviruses and Coronavirus at Chirundu and
Magweto site.

The prevalence of co-infection of bat communities from Chirundu and Magweto sites was fairly low
however it showed peaks that correlated with peaks in coronavirus and astrovirus infections as
observed in the graphs. The overall prevalence of co-infection of insectivorous bats from Chirundu
site by Astroviruses and Coronaviruses was 3.6% [95% CI: 3.8-5.8] (Table 2). The highest co-
infection prevalence observed in this site was 10.6% [95% CI: 6.8-16.1] in February 2021 during the
weaning period while the lowest prevalence of 0% was observed in August [95% CI: 0.0-0.2] and
October [95% CI: 0.0-1.3] during the end of the non-gestation and the pregnancy period (Table 2,
Figure 3a).

The overall prevalence for co-infection of insectivorous bats in Magweto site by coronaviruses and
astroviruses was 2.6% [95% CI: 1.8-3.5] (Table 2, Figure 3b). Co-infection in this site was recorded
in all the months, from September 2020 to June 2021, except from December to February during
which no sampling took place (Table 2, Figure 3b). The highest prevalence of co-infection was 8.3%
[95% CI: 5.4-12.4] in March during the 4-6 months old juvenile period; while the lowest 0.3% [95%
CI: 0.05-1.6] was observed in September 2020 during the non-gestation period (Table 2, Figure 3b).
290

Figure 3a

	14/08/20	14/09/20	15/10/20	15/11/20	15/12/20	15/01/21	15/02/21	14/03/21	15/04/21	15/05/21	15/06/21	14/07/21
	Non-gestation		Pregnancy	Parturition	Parturition & Lactation	Lactation	Weaning	4-6_months Juveniles		Non-gestation		
◆AstV+CoV co-infection Prevalence (%)	0		0	0.4	4.4		10.6	7.9		3.7		3.5
■Chirundu CoV Prevalence (%)	5.8		1.35	18.6	37.5		44.2	35		25.2		31.4
▲Chirundu AstV Prevalence (%)	0		1	1.2	10.1		17.1	21.2		19.3		13.3

2
292

Figure 3b

	14/08/20	14/09/20	15/10/20	15/11/20	15/12/20	15/01/21	15/02/21	14/03/21	15/04/21	15/05/21	15/06/21
	Non-gestation		Pregnancy	Parturition	Parturition & Lactation	Lactation	Weaning	4-6_months Juveniles		Non-gestation	
◆AstV+CoV co-infection Prevalence (%)		0.3	2.7	1.3				8.3	2.5		1.2
■Magweto CoV Prevalence (%)		2	10.5	35.9				34.7	16.6		7.02
▲Magweto AstV Prevalence (%)		5.2	17.9	13.6				19	10.7	0	19.4

Figure 3*:* Prevalences of astroviruses and coronavirus co-infection (in green), coronaviruses (in blue) and astroviruses (in red) per month in bat communities for a/ Chirundu site and b/ Magweto site.

Genetic diversity of RNA Astroviruses

In this study we sequenced all *Astrovirus* positive samples from 5 sites: (Chirundu farm N=179; Magweto cave N=229; Mabura cave N=7; Honde Valley N=7 and Baobab tree N=1). Of the 423 samples, 214 (50.6%) were amplified from *Hipposideros* spp, 63 (14.9%) from *Rhinolophus* spp, 53 (12.5%) from *Miniopterus*, 10 (2.4%) from *Nycteris* spp, 7 (1.7%) from *Eidolon* spp and for the remaining 76 samples, the bat genus / species could not be determined mainly owing to bad sequence qualities.

Phylogenetic analysis of the 155 *RdRp AstV* nucleotide sequences generated for this study showed that they all clustered with the genus *Mamastrovirus* of the *Astroviridae* family (Figure 4).

Two sequences isolated from Magweto cave, 1 from *Rhinolophus spp* (MAG-573) and 1 from *Hipposideros spp* (MAG-292) clustered with a group of Human astroviruses (*HuAstVs*) (Figure 4). This cluster was well sustained (Bootstrap >90%) and the nucleotidic acid identities between Bat Astroviruses MAG-573, MAG-292 and HuAstV-2 were 96 and 93% respectively (Data not shown). The other specific cluster of bat astrovirus showing a close relation to human astroviruses, is a sequence isolated from *Hipposideros spp* (MAG-1236) in Magweto, it clustered with *HuAstV-MLB-1; -2* and *3* (Figure 4). Six sequences isolated from the frugivorous bats *E.helvum* found in Honde Valley and one from *Rhinolophus spp* collected in a Baobab tree formed a sister clade with another group of Human astroviruses *HuAstV-VA-1;-2;-3;-4* and *HuAstV-HMO-A* (Figure 4). The majority of the sequences, comprises sequences closely related to insectivorous Bat Astroviruses (Figure 4). The majority of these insectivorous bat related astroviruses sequences were amplified from *Hipposideros* (HC-1 to -9) and *Rhinolophus* (Rhi-1 to -4) bat species. One cluster comprised of sequences isolated from *Rhinolophus spp*. Most of these sequences were isolated from Chirundu and a few of them from Magweto, and clustered together with *AstV* strains from *Nyctalus* and *Verpitillio* bat species isolated in Czech Republic (references Y & Z in the phylogenetic tree) (Figure 4). Another cluster comprising of majority of sequences from Chirundu and some from Magweto site, isolated from *Miniopterus* and *Rhinolopus* spp showed phylogenetic clustering with strains of bat astroviruses isolated from *Miniopterus* spp from Madagascar and China (references N, O, R, U, T), *Rhinolophus* spp from Korea (references P, Q) as well as *Rousettus* and *Paratrianops BAstV* from Madagascar (references V, W) (Figure 4)

**Figure 4*: Phylogenetic tree of *Astroviruses* partial *RdRp* gene.* The sequences detected at Chirundu site are represented by red branches, at Magweto site by blue branches, at Honde Valley in by green branches, at Mabura cave by brown branches and at Baobab site by an orange branch. The red rectangle highlighted the phylogenetic relationship between MAG-292, MAG-573 and *HuAstV* strains. The tree was built using the maximum likelihood method based on the GTR + G4 + I model. The robustness of nodes was assessed with 1000 bootstrap replicates. Bootstrap values >70 are in asterisk and those <70 are not shown. HC=*Hipposideros caffer*; RS=*Rhinolophus simulator*; NT=*Nycteris thebaica*; EH= *Eidolon helvum*; Min=*Miniopterus spp*; Rhi=*Rhinolophus species*. MAG=Magweto site; CHI=Chirundu site; BAOB= Baobab; MAB= Mabura site; HV=Honde valley site.

Human *Astrovirus* are highlighted in red and bold letters represented the different bat AstV references used to build this tree:

Bat Astrovirus references:
A - MH013972.1: Hipposideros caffer, Mozambique; **B** - MH013971.1: Hipposideros caffer, Mozambique; **C** - MH013970.1: Hipposideros caffer, Mozambique; **D** - KX858349.1: Rhinolophus spp., Laos; **E** – MC841039.1: Rhinolophus Ferrumquinum, Korea; **F** - MH013989.1: Trianop afer, Mozambique ; **G** - MH013986.1: Trianop afer, Mozambique ; **H** - MH013971.1: Hipposideros caffer, Mozambique ; **I** - KX858371.1: Hipposidoros larvatus, Laos **J** - KY575656.1: Myotis goudoti, Madagascar ;
K - KY575655.1: Myotis goudoti, Madagascar; **L** - MH013988.1: Trianops afer, Mozambique
M - MH013076.1: Nycteris thebaica, Mozambique ; **N**- KY575649.1: Miniopterus griveaudi, Madagascar ; **O** - KY575647.1: Miniopterus griveaudi, Madagascar ; **P** - MC840967.1: Rhinolophus ferrumquinum, Korea ; **Q** - MC840969.1: Rhinolophus ferrumquinum, Korea ; **R** - KY575648.1: Miniopterus griveaudi, Madagascar ; **S** - EU847151.1 : Miniopterus magnater, Chine ; **T** - KY575674.1: Miniopterus griveaudi, Madagascar ; **U**- KY575666.1: Rousettus madagascariensis, Madagascar ; **V** - KY575661.1 : Paratrianops furculus, Madagascar ; **W** - KY575644.1 : Miniopterus griveaudi, Madagascar ; **X** - EU847155.1: Miniopterus pussilus, Chine ; **Y**- KP843561.1: Vespertillio Murinus, Czech ; **Z**- KP843558.1: Nyctalus noctule, Czech ; **AA** - KX858513.1: Megaderma lyra, Cambodge ; **AB** - HM368175.1: Myotis myotis, Germany ; **AC** - KX858367.1: Roussetus spp., Laos ; **AD** - FJ571067.1: Taphozous melanopogon, Chine ; **AE** - KX858377.1 : Roussetus spp, Cambodge ; **AF** - KX858378.1 : Roussetus spp, Laos
Human Astrovirus references:
 GQ502195.1: Human Astrovirus VA2; JQ673585.1: Human Astrovirus HMO-A; JX857869.1: Human Astrovirus VA4; KJ920196.1: Human Astrovirus VA1; GQ502196.1: Human Astrovirus VA3; JX857870.1: Human Astrovirus MLB3; JQ673581.1: Human Astrovirus MLB1; GQ502188.1: Human Astrovirus MLB2; L23509.1: Human Astrovirus Type 2; AY20891.1: Human Astrovirus Type 4; DQ028633.1: Human Astrovirus Type 5; GQ495608.1: Human Astrovirus Type 6; JF491430.1: Human Astrovirus Type 3
Other Astrovirus references:
GQ914773.1: Porcine Astrovirus (Po AstV); FJ890355.1: Bottlenose Dolphin Astrovirus (BDo AstV); JN 420351.1: California Sea Lion Astrovirus (CSL AstV); æEU669001.1: Chicken Astrovirus (Ch AstV)

Two major clades from Chirundu and one from Magweto, isolated from *Miniopterus, Rhinolophus* and *Hipposideros spp* of bats also phylogenetically clustered with a strain from Mozambique isolated from *Nycteris thebaica* (reference M). One sequence from Magweto isolated from *Rhinolophus spp* and two from Mabura cave isolated from *Hipposideros* spp, clustered with a strain of bat astrovirus isolated from *Myotis* spp in Madagascar (references K, L) (Figure 4). A cluster, comprising two clades from Magweto and one sequence from Chirundu and three sequences from Mabura cave isolated from *Rhinolophus* and *Hipposideros,* were closely related to bat astrovirus strains from *Hipposideros* spp of Mozambique and Laos (references H, I) (Figure 4). The last small group comprising one clade from Chirundu, four sequences from Magweto, all isolated from *Rhinolophus* spp, and one sequence from Mabura isolated from *Hipposideros* spp clustered together with bat astrovirus strains from *Rhinolophus* spp of LAOS and Korea (references D, E, F) (Figure 4).

One of the large clusters shows the phylogenetic relationship amongst sequences derived from, *Hipposideros* spp (Clades HC-5 to -8), these making up the majority of the sequences in this clade

and one isolated from *Miniopterus* spp, one from *E. helvum* and one from a *Nycteris* spp. The bat

astroviruses in this clade, all showed a close relation to a bat astrovirus strain isolated from

Hipposideros spp of Mozambique (reference C). The largest cluster constituted mainly by Magweto

site sequences and some Chirundu sequences, from *Hipposideros* spp (clades HC-1 to 4) and a few

Rhinolophus and *Nycteris* spp, clustered with a strain from *Hipposideros* spp of Mozambique

(references A, B) (**Figure 4**).

Discussion

The overall prevalence of *AstVs* in bats was 10% and 13.7% in Chirundu and Magweto sites,

indicating a fairly high circulation of this family of viruses in bats in Zimbabwe. However, compared

to studies in other African countries, the observed prevalence was low: Hoarau et *al* and

Lebarbenchon et *al* detected *AstV* in 20-22% of individual bats in Mozambique and Madagascar,

respectively [17,19]. This difference in prevalence of *AstV* compared to other countries can be

attributed to the lack of longitudinal survey in the Mozambican and Madagascar studies as sampling

effort was done for periods of two separate months in both studies [17,19].

In Chirundu site increasing RNA-*AstV* prevalences were observed from the lactation to the 4-6

months old juvenile seasons with 21.2% of RNA -*AstV* detection during the latter. Astroviruses have

been described to display seasonal variations in prevalence [17]. Drexler et al observed different

peaks of astrovirus detection associated with different stages of the reproductive season with peaks

correlating to maternal aggregations during breeding season and after parturition season due to

establishment of a susceptible subpopulation of weaned new-born bats who did not yet have their

own adaptive immunity [34]. In our study the prevalence was high during some of the months

compared to overall prevalence, which might be related to the effect of reproductive seasons on the

shedding of viruses, as demonstrated for other viruses. During the reproductive season with juveniles

and immature individuals, the prevalence of CoV was very high as compared to prevalence in the

absence of juveniles and presence of sexually mature bats [35,36]. In this study we observed a similar

trend for AstVs for one of the study sites. Furthermore, Mendenhall et *al*, also identified the bat

juvenile stage as exhibiting a greater *Astrovirus* viral burden than any other stage of the reproductive

season [37]. The range of prevalence observed during the highest peaks coincides with the ≥20%

overall AstV detection reported by Hoarau et *al* [19] during the time of their sampling in February

and May. Here we observe a peak during the 4-6 months old juvenile season in Chirundu site which

corresponds to a high influx of immunologically immature individuals in the bat population, similar

to the what is observed in the Chidoti et *al*, (2022) for CoV infection [26].

During this period, pups are known to contribute to shedding pulses of viruses as they develop
productive infections during the acute phase [24], which coincides with the waning protection of
maternal antibodies [26], thus increasing the susceptibility and rate of infection in the young.
We could not observe a clear trend in Magweto site, which might be related to the absence of samples
from December 2020 to March 2021, which correspond to the lactation and weaning periods.

In our study we also investigated co-infection of bats by coronaviruses and astroviruses. We compared
the co-infection prevalence with the prevalence of astroviruses and coronaviruses described at each
site during the same reproductive cycles. The co-infection prevalence showed a similar trend to that
observed in the individual viruses, whereby during observed peaks of high AstVs and CoVs
detections, the co-infection also peaked at similar phases of the reproductive cycle. Thus the co-
infection of bats by both CoVs and AstVs is high when there is high infection of bats by either virus.
The overall prevalence of co-infection was 3.5% and 2.6 % in Chirundu and Magweto respectively
while the bats from Madagascar and Mozambique respectively showed a *AstV-CoV* co-infection of
5% (±2.7%) and of 2.7% (±1,1%) [17,19]. Co-infection and recombination of viruses in bats have
been reported on several occasions including co-infection of bats by coronaviruses and astroviruses
[17,23]. Bats infected with either *CoV* or *AstV* were shown to be more likely co-infected with the
respective virus [23]. In our study, no effect of coronavirus infection/shedding on *Astrovirus*
infection/shedding was observed.

In the current study we detected bat Astroviruses from *Hipposideridae, Rhinolophidae, Pteropodidae,*
*Nycterida*e and *Miniopterida*e families. Astroviruses are known to show no host restriction and are
widespread within the Chiroptera order [13,20]. The majority of the astroviruses amplified were from
Hipposideros and *Rhinolophus* species and these clustered with *AstV* sequences derived from
Mozambique, Madagascar and China of the same families [17,19,20]. The observed trend is due to
the two genera being the most dominant species of bats at both sites where sampling was done.
Therefore, the probability of higher detection of *AstV* in these species is expected and evidently
observed to be higher than other rare or less dominant species. Active transmission events of
astroviruses amongst *Rhinolophidae* and *Hipposideridae* bat species are known to occur [20]. In this
study, a high degree of species-specific tropism, especially in bat astrovirus related clusters was
observed with astrovirus strains isolated from *Hipposideros spp* clustering with each other and similar
trends for *Rhinolophus* and *Miniopterus spp*. However, the clusters were not site-specific as
sequences from Magweto, Mabura and Chirundu sites formed clusters with each other. There is a
significant phylogenetic clustering of isolated bat astrovirus according to specific species. Previous
studies on astroviruses in bats have also described this family of viruses to be involved solely in
tropism transmission which is species specific [38–40]. *Astrovirus* sequences isolated from different
bat species of the same site clustered together. This indicates potential cross species transmission of

astroviruses within each site, as reported by Xiao et *al* in bats sharing the same habitat [40]. We also

observed phylogenetic clustering of astroviruses from Chirundu and Magweto into same clades

showing a great diversity of astroviruses in circulation within the studied bat communities. Novel

astroviruses species have been reported in bats hosts [21], and in this study it is evident due to the

diverse astroviruses described. The *AstV* isolated from *E. helvum*, frugivorous bats, clustered together

and showed close phylogenetic relation to *BAstV* isolated from *Rousettus spp.* bats in China and

Taphozous spp. bats in Laos. One strain isolated from *E. helvum* clustered with *Hipposideros* bat

astroviruses sequences isolated in Chirundu site. This could mean that frugivorous bats and

insectivorous bats can be infected by the same type of AstVs strain thus indicative of the multi-host

spectrum for the Astroviridae family. Detection of *Astrovirus* in the frugivorous bats was low

compared to insectivorous bats, and this can be attributed to the insectivorous bats ability to adapt

and harbour more astroviruses than frugivorous bats [40].

Two strains, Mag-292 *H. Caffer*, Mag-573 *Rhinolophus spp* clustered within the *HuAstv* type 1 to 6,

that are known to cause infection and diarrhoea in children and infants [40]. Six of our strains isolated

from *E. helvum* and one from *Rhinolophus spp* formed sister clades to Human *AstV* HMO, associated

with severe extra-intestinal illnesses in humans [12] and VA-1-4 strains. Another *AstV* strain detected

in *Hipposideros* bat species was close and formed a sister clade with *HuAstv* MLB-1, -2 and -3 all

known to cause gastro-intestinal and neurological diseases with mild-severe symptoms in humans

[41]. This phylogenetic clustering could suggest that *HuAstV* and *BAstV* may have a shared common

ancestor. However, with limited sequence data the evolutionary history is merely a speculation that

needs to be further investigated through ORFs sequencing and MRCA reading. Another hypothesis

for the observed relation is that, since bats are known to be important reservoir hosts of astroviruses

with great diversity, they could be ancestors of the *HuAstV* due to the high prevalence of *AstV* detected

in bats. *HuAstV* transmission occurs via the faecal-oral route and contaminated food or water [2],

therefore it is unlikely that anthroponosis may have occurred. Third, this observed relation could be

due to a rare event or mutation, or a new strain of bat astrovirus that has not been described yet.

Further full genomic data as well as epidemiological data on these bat colonies are needed to reach a

full and true conclusion.

As novel strains and host species continue to be discovered, understanding *AstV* transmission and

their zoonotic potential is essential. Disseminating *HuAstV* and *AstV* from zoonotic reservoir hosts

remain significant threats to public health [4]. Astroviruses have a high genetic diversity, multiple

mechanism of generating additional diversity, and infect a wide range of host species therefore

understanding their prevalence in their wildlife hosts can help to predict or prevent the emergence of

novel astrovirus strains into domestic animals and human populations.

Acknowledgments: We thank Billy Butete for his field assistance. The authors gratefully acknowledge and thank the International Atomic Energy Agency for making available the sequencing services. We thank the Research Council of Zimbabwe for approving this study (research registration certificate N° 03006) and the Hurungwe Rural District council for their assistance and facilitation. We thank the Animal Research Ethics Committee of Zimbabwe for their approval (ref number 002/2017). This work was conducted within the framework of the Research Platform "Production and Conservation in Partnership" (RP–PCP).

Funding: This work was supported by grants of the French Ministry of Europe and Foreign Affairs (Fond de Solidarité pour les Projets Innovants, les sociétés civiles, la francophonie et le développement humain—CAZCOM Project, FSPI N°2019/88) and the AFD (French Agency for development-Projet PACMAN, AFD CWZ 1019 01 V).

Conflicts of Interest: We declare that we have no conflicts of interest

Author's contribution: Conceptualization: VC, MA, HdN, MB, LG and FL.; **Methodology**: VC, MA, LG, HdN, DP, EG, GM, EM and FL.; **Software:** VC, MA, HdN and FL; **Validation**: VC, MA HdN, JC, and FL; **Formal Analysis**: VC, HdN, JC, and FL.; **Investigation**: VC, VP, NC, GM, and FL.; **Resources**: DP, GM, MP, MB, HdN, LG, and FL.; **Data Curation**: VC, HdN, and FL.; **Writing – Original Draft Preparation**: VC and FL; **Writing – Review & Editing**: HdN, MA, JC, MB, LG, VP, EG, DM, DP, EM, GM and FL.; **Supervision**: HdN, DP, GM and FL.; **Project Administration**: LG, MB, HdN and FL; **Funding Acquisition**: MB, HdN, LG and FL.

Data, scripts, code, and supplementary information availability

GenBank Accession Numbers

The *Astrovirus* sequences have been deposited to the GenBank under the following numbers: OQ271049 - OQ271203

The Cytochrome B sequences have been deposited to the GenBank under the following numbers: OM487705-OM488020

Statistical analysis

The script used for the GLMM analysis is available at: https://doi.org/10.5281/zenodo.7847934

Reference

[1] Bosch A, Pintó RM, Guix S. Human Astroviruses. Clin Microbiol Rev . 2014;27:1048–1074. doi/10.1128/CMR.00013-14.

[2] Donato C, Vijaykrishna D. The Broad Host Range and Genetic Diversity of Mammalian and Avian Astroviruses. Viruses . 2017;9:102. doi/10.3390/v9050102.

[3] Wu H, Pang R, Cheng T, et al. Abundant and Diverse RNA Viruses in Insects Revealed by RNA-Seq Analysis: Ecological and Evolutionary Implications. Cristea IM, editor. mSystems. 2020;5. doi/10.1128/mSystems.00039-20.

[4] Roach SN, Langlois RA. Intra- and Cross-Species Transmission of Astroviruses. Viruses. 2021;13:1127. doi/10.3390/v13061127.

[5] El Taweel A, Kandeil A, Barakat A, et al. Diversity of Astroviruses Circulating in Humans, Bats, and Wild Birds in Egypt. Viruses . 2020;12:485. doi/10.3390/v12050485.

[6] Vu D-L, Cordey S, Brito F, et al. Novel human astroviruses: Novel human diseases? J Clin Virol . 2016;82:56–63. doi/10.1016/j.jcv.2016.07.004.

[7] Johnson C, Hargest V, Cortez V, et al. Astrovirus Pathogenesis. Viruses . 2017;9:22. doi/10.3390/v9010022.

[8] Frémond M-L, Pérot P, Muth E, et al. Next-Generation Sequencing for Diagnosis and Tailored Therapy: A Case Report of Astrovirus-Associated Progressive Encephalitis. J Pediatric Infect Dis Soc . 2015;4:e53–e57. doi/10.1093/jpids/piv040.

[9] Cortez V, Meliopoulos VA, Karlsson EA, et al. Astrovirus Biology and Pathogenesis. Annu Rev Virol . 2017;4:327–348. doi/10.1146/annurev-virology-101416-041742.

[10] Janowski AB, Klein RS, Wang D. Differential In Vitro Infection of Neural Cells by Astroviruses. Griffin DE, editor. MBio . 2019;10. doi/10.1128/mBio.01455-19.

[11] Boujon CL, Koch MC, Seuberlich T. The Expanding Field of Mammalian Astroviruses: Opportunities and Challenges in Clinical Virology. Adv Virus Res. 2017. p. 109–137.10.1016/bs.aivir.2017.07.002

[12] Wohlgemuth N, Honce R, Schultz-Cherry S. Astrovirus evolution and emergence. Infect Genet Evol . 2019;69:30–37. doi.org/10.1016/j.meegid.2019.01.009.

[13] Chu DKW, Poon LLM, Guan Y, et al. Novel Astroviruses in Insectivorous Bats. J Virol . 2008;82:9107–9114. doi/10.1128/JVI.00857-08.

[14] Dufkova L, Straková P, Širmarová J, et al. Detection of Diverse Novel Bat Astrovirus Sequences in the Czech Republic. Vector-Borne Zoonotic Dis . 2015;15:518–521.doi/10.1089/vbz.2015.1813.

[15] Rougeron V, SuquetT E, Maganga GD, et al. Characterization and phylogenetic analysis of new bat astroviruses detected in Gabon, Central Africa. Acta Virol . 2016;60:386–392. doi/10.4149/av_2016_04_386.

[16] Lacroix A, Duong V, Hul V, et al. Diversity of bat astroviruses in Lao PDR and Cambodia.
Infect Genet Evol . 2017;47:41–50. doi/10.1016/j.meegid.2016.11.013.

[17] Lebarbenchon C, Ramasindrazana B, Joffrin L, et al. Astroviruses in bats, Madagascar. Emerg
Microbes Infect . 2017;6:1–3. doi/10.1038/emi.2017.47.

[18] Waruhiu C, Ommeh S, Obanda V, et al. Molecular detection of viruses in Kenyan bats and
discovery of novel astroviruses, caliciviruses and rotaviruses. Virol Sin . 2017;32:101–114.
doi/10.1007/s12250-016-3930-2.

[19] Hoarau F, Le Minter G, Joffrin L, et al. Bat Astrovirus in Mozambique. Virol J . 2018;15:104.
doi/10.1186/s12985-018-1011-x.

[20] Lee S-Y, Son K-D, Yong-Sik K, et al. Genetic diversity and phylogenetic analysis of newly
discovered bat astroviruses in Korea. Arch Virol . 2018;163:3065–3072. doi/10.1007/s00705-
018-3992-6.

[21] Bergner LM, Mollentze N, Orton RJ, et al. Characterizing and Evaluating the Zoonotic Poten-
tial of Novel Viruses Discovered in Vampire Bats. Viruses. 2021;13:252.
doi/10.3390/v13020252.

[22] Fischer K, Zeus V, Kwasnitschka L, et al. Insectivorous bats carry host specific astroviruses
and coronaviruses across different regions in Germany. Infect Genet Evol . 2016;37:108–116.
doi/10.1016/j.meegid.2015.11.010.

[23] Seltmann A, Corman VM, Rasche A, et al. Seasonal Fluctuations of Astrovirus, But Not Coro-
navirus Shedding in Bats Inhabiting Human-Modified Tropical Forests. Ecohealth.
2017;14:272–284. doi/10.1007/s10393-017-1245-x.

[24] Plowright RK, Peel AJ, Streicker DG, et al. Transmission or Within-Host Dynamics Driving
Pulses of Zoonotic Viruses in Reservoir–Host Populations. Remais J V., editor. PLoS Negl
Trop Dis . 2016;10:e0004796. doi/10.1371/journal.pntd.0004796.

[25] Bourgarel M, Pfukenyi DM, Boué V, et al. Circulation of Alphacoronavirus, Betacoronavirus
and Paramyxovirus in Hipposideros bat species in Zimbabwe. Infect Genet Evol.
2018;58:253–257. doi/10.1016/j.meegid.2018.01.007.

[26] Chidoti V, De Nys H, Pinarello V, et al. Longitudinal Survey of Coronavirus Circulation and
Diversity in Insectivorous Bat Colonies in Zimbabwe. Viruses. 2022;14:781.
doi/10.3390/v14040781.

[27] Monadjem A, Taylor PJ, Schoeman FPD, et al. Bats of Southern and Central Africa: A Bioge-
ographic and Taxonomic Synthesis. African Zool. 2011;46:437–437.
doi/10.1080/15627020.2011.11407657.

[28] Kocher TD, Thomas WK, Meyer A, et al. Dynamics of mitochondrial DNA evolution in ani-
mals: amplification and sequencing with conserved primers. Proc Natl Acad Sci.
1989;86:6196–6200. doi/10.1073/pnas.86.16.6196.

[29] Kumar S, Stecher G, Tamura K. MEGA7: Molecular Evolutionary Genetics Analysis Version
7.0 for Bigger Datasets. Mol Biol Evol . 2016;33:1870–1874. doi/10.1093/molbev/msw054.

[30] Guindon S, Dufayard J-F, Lefort V, et al. New Algorithms and Methods to Estimate Maxi-
mum-Likelihood Phylogenies: Assessing the Performance of PhyML 3.0. Syst Biol.
2010;59:307–321. doi/10.1093/sysbio/syq010.

[31] Milne I, Lindner D, Bayer M, et al. TOPALi v2: a rich graphical interface for evolutionary
analyses of multiple alignments on HPC clusters and multi-core desktops. Bioinformatics.
2009;25:126–127. doi/25/1/126/302670.

[32] Lemoine F, Wilkinson E, Correia D, et al. Renewing Felsenstein's Phylogenetic Bootstrap in
the Era of Big Data. Nature. 2018. doi/10.1038/s41586-018-0043-0

[33] Wilson, E. B. Probable inference, the law of succession, and statistical inference. Journal of
the American Statistical Association. 1927; **22** (158): 209–212. doi/10.2307/2276774.

[34] Drexler JF, Corman VM, Wegner T, Tateno AF, Zerbinati RM, Gloza-Rausch F, et al.
Amplification of emerging viruses in a bat colony. Emerg Infect Dis. 2011;17:449–56.
doi/10.3201/eid1703.100526.

[35] Wacharapluesadee, S.; Duengkae, P.; Chaiyes, A.; Kaewpom, T.; Rodpan, A.;
Yingsakmongkon, S.; Petcharat, S.; Phengsakul, P.; Maneeorn, P.; Hemachudha, T.
Longitudinal study of age-specific pattern of coronavirus infection in Lyle's flying fox
(Pteropus lylei) in Thailand. Virol. J. 2018, 15, 1–10. doi/10.1186/s12985-018-0950-6.

[36] Cappelle J, Furey N, Hoem T, Ou TP, Lim T, Hul V, Heng O, Chevalier V, Dussart P, Duong V.
Longitudinal monitoring in Cambodia suggests higher circulation of alpha and
betacoronaviruses in juvenile and immature bats of three species. Sci Rep. 2021;11(1):24145.
doi/10.1038/s41598-021-03169-z.

[37] Mendenhall IH, Skiles MM, Neves ES, Borthwick SA, Low DHW, Liang B, Lee BPYH, Su
YCF, Smith GJD. Influence of age and body condition on astrovirus infection of bats in
Singapore: An evolutionary and epidemiological analysis. One Health .2017; 4: 27-33.
doi.org/10.1016/j.onehlt.2017.10.001.

[38] George DB, Webb CT, Farnsworth ML, et al. Host and viral ecology determine bat rabies
seasonality and maintenance. Proc Natl Acad Sci. 2011;108:10208–10213.
doi/10.1073/pnas.1010875108.

[39] Baxendale W, Mebatsion T. The isolation and characterisation of astroviruses from chickens.
Avian Pathol . 2004;33:364–370. doi/10.1080/0307945042000220426.

[40] Xiao J, Li J, Hu G, et al. Isolation and phylogenetic characterization of bat astroviruses in
southern China. Arch Virol . 2011;156:1415–1423. doi/10.1007/s00705-011-1011-2.

[41] Kapoor A, Li L, Victoria J, et al. Multiple novel astrovirus species in human stool. J Gen Virol.
2009;90:2965–2972. doi/10.1099/vir.0.014449-0.