Longitudinal Survey of Astrovirus infection in different bat species in Zimbabwe: **Evidence of**

high genetic Astrovirus diversity

Chidoti Vimbiso[1], De Nys Hélène[2,3,4], Abdi Malika[5], Mashura Getrude[1], Pinarello Valérie [1,2,3,4],
Chiweshe Ngoni[4], Matope Gift[1], Guerrini Laure[2,3], Pfukenyi Davies[1,6], Cappelle Julien [2,3],
Mwandiringana Ellen[1], Missé Dorothée[2], Gori Elizabeth[1,7], Bourgarel Mathieu[2,3,4] and Liégeois
Florian [1,5,*].

[1]Faculty of Veterinary Science-Biotechnology Center, University of Zimbabwe, P.O. Box MP167

Harare, Zimbabwe

[2]ASTRE, Univ Montpellier, CIRAD, INRAE, Montpellier, France.

[3]CIRAD, UMR ASTRE, Montpellier, France.

[4] CIRAD, UMR ASTRE, Harare, Zimbabwe

[5]MIVEGEC, University of Montpellier, IRD, CNRS, 34394 Montpellier, France

[6]Faculty of Animal and Veterinary Sciences, Botswana University of Agriculture and Natural

Resources, Private Bag 0027, Gaborone, Botswana

[7]Department of Biochemistry, Molecular Biology and Genetics, University of Rwanda, Rwanda

*Corresponding Author

Florian Liegeois

University of Zimbabwe, Faculty of Veterinary Science, Veterinary Biosciences Departement

Room 2F7, P.O. Box MP167 Harare, Zimbabwe.

Florian.liegeois@ird.fr

Keywords: Astrovirus, Bats, Zimbabwe, Phylogeny,

Abstract

Astroviruses (AstVs) have been discovered in over 80 animal species including diverse bat species and avian species. A study on Astrovirus circulation and diversity in different insectivorous and frugivorous chiropteran species roosting in trees, caves and building basements was carried out at 11 different sites across Zimbabwe. Pooled and individual faecal sampling methods were used for this study, with collection dates ranging from June 2016 to July 2021. In two sites, Magweto and Chirundu, sampling was carried out at monthly intervals from August 2020 to July 2021. Astroviruses and bat mitochondrial genes were amplified using pan-AstVs and CytB /12S RNA PCR systems respectively. Phylogenetic analysis of the *RdRp* Astrovirus sequences revealed a high genetic diversity of astroviruses. All the bat astroviruses tested in this study clustered with the *Mamastrovirus* genus. Two distinct groups of amplified sequences were identified. One group was comprised of sequences isolated from *Hipposideros, Rhinolophus* and *Eidolon helvum* spp. clustered with Human Astrovirus strains: *HuAstV* types1-6, *HuAstV*-MLB1-3 and *HuAstV*-VA-1. The second group comprising the majority of the sequences clustered with different strains of Bat AstVs. Results from the longitudinal study at Magweto and Chirundu showed an overall AstV prevalence of 13.7% and 10.4% respectively. Peaks of AstV infection at Chirundu coincided with the period when juveniles are 4-6 months old. By combining data from our previous work on Coronaviruses, we also anaylzed co-infection of bats with Coronaviruses and Astroviruses at Magweto and Chirundu sites for which the prevalence of co-infection was 2.6% and 3.5% respectively.

Introduction

Astroviruses (AstVs) are non-enveloped single-stranded positive sense RNA viruses with a genome length of approximately 6.2 to 7.7 Kb [1]. They have been described in over 80 non-human species and are classified into two genera: *Mamastrovirus* and *Avastrovirus,* respectively infecting mammals and avian species [2]. According to the International Committee on Taxonomy of Viruses (https://talk.ictvonline.org), *Mamastrovirus* are classified into 19 recognized species, *MAstV*-1 to -19, and two genogroups GI and GII [3]. All classic and novel Human Astrovirus (*HuAstVs*) belong to four different species, *MAstV*-1, *MAstV*-6, *MAstV*-8 and *MAstV*-9 [2,4,5]. Recently, studies have reported AstVs infections in fish and insects [6,7], however bats and wild birds are considered to be natural reservoirs of the *Astroviridae* family [8,9]. Generally *HuAstVs* have been identified as causal agents of acute viral gastrointestinal illness worldwide particularly in children, immunocompromised people and the elderly [6–8]. Of note, in humans, the majority of AstV-associated encephalitis or meningitis illnesses were reported in immunocompromised people [10–12]. Furthermore, beyond this well-known clinical manifestation of AstVs infections, neurovirulent AstVs infections have also been reported in both humans and domestic animals [13,14]. Moreover, studies have reported numerous cross-species transmissions of AstVs [7], where some AstVs detected in humans closely clustered with strains identified in rodents, minks, feline species and ovine [2,5,7]. Bat-borne viruses represent an extensive research field owing to the plethora of viruses hosted by Chiropterans [15]. Furthermore, this mammalian order is known to be persistently infected by astroviruses [16] and this is inclusive of many species of insectivorous bats [17]. To date, a high diversity of bat AstVs from different bat families has been reported worldwide [11,18-19]. Bat AstVs belong to *MAstV*-12, and *MAstV*-14 to -19 species [1]. In Europe a variety of bat species belonging to the Yangochiroptera sub-order have been discovered to harbor several astroviruses [20]. In Africa, bat astroviruses were reported in Egypt, Gabon, Madagascar, Kenya and Mozambique [18, 21–23].

AstVs are known to occur in bat species with high prevalence and exceedingly high genetic diversity whereby infections in these hosts are usually non-pathogenic [24,25]. Despite all these studies, large gaps exist on astroviruses prevalence and infection dynamics in bats [9]. Astroviruses prevalence and detection in bat colonies seemed rather correlated to abiotic factors such as seasons and year of sampling and biotic factors such as sex and reproductive status and also viral co-infection [26]. Viral shedding in bats occurs in spatial and temporal pulses that can drive spillover to other animals or humans [27-28]. The varying prevalence of viruses in bats is determined by shedding pulses, where rare or very low prevalences are detected in the absence of shedding pulses and higher prevalence is detected when shedding pulses occur [27,29].

Until now, no data was available on genetic diversity and prevalence of circulating AstVs in insectivorous bats in Zimbabwe. In this study we enlarge the spectrum of bat AstVs knowledge in

88   Africa by identifying and analyzing bat-AstVs, their diversity and prevalence from different bat
89   species colonies according to reproductive season phenology in Zimbabwe. Furthermore, we will also
90   show and highlight the presence of co-infection of bats by astroviruses and coronaviruses (CoVs) and
91   how it varies according to reproductive phenology.
92

Material and Methods
Sampling approaches and sites
Two different approaches were followed in this study: bat community sampling and individual bat
sampling.

Bat community pooled sampling
Between February 2016 and December 2020, faecal samples from both insectivorous and frugivorous
bat species were respectively collected in different sites including caves, an ancient mine and trees in
Zimbabwe (Figure 1, Table 1).

101
102   **Figure 1:** The study sites highlighted on the map show the sampling areas for insectivorous and frugivorous
103   bats colonies across Zimbabwe. In Honde valley, the *Eidolon helvum* colony was established in at least four
104   different sites represented by the rectangle at the right of the figure. Red circles represent community approach
105   study sites; yellow circles represent longitudinal survey sites (individual approach).

Table 1: Number of Astrovirus positives per site/sampling year in pooled faecal samples from frugivorous and insectivorous bats for the selected 14 sites.

Sampling Sites	Type of roost	Collection Date	Bat Species	Number of collected tubes	Number of Tested pools	Number of positive pools
Magweto Cave	Cave	June 2016	*Hipposideros spp.*	24	8	5
Bashongwe Cave	Cave	March 2018	*Rhinolophus and Hipposideros spp.*	59	7	0
		July 2018		57	20	0
		December 2018		135	45	0
Mabura Cave	Cave	June 2016	*Hipposideros spp.*	36	12	6
		February 2017	*Hipposideros spp.*	22	7	0
Mugunza Cave	Cave	April 2017	*Roussetus aegyptus*	50	13	0
Honde Valley *Machiri*		August 2017	*Eidolon helvum*	25	9	0
Muwomba		November 2017	*Eidolon helvum*	66	22	0
Nyamajo		November 2017	*Eidolon helvum*	58	19	0
Nyamhosva		November 2017	*Eidolon helvum*	35	11	0
Nyamajo	Trees	January 2018	*Eidolon helvum*	17	6	1
Machiri		January 2018	*Eidolon helvum*	7	2	2
Muwomba		January 2018	*Eidolon helvum*	8	2	0
Nyamhosva		January 2018	*Eidolon helvum*	28	10	1
Nyamajo		February 2020	*Eidolon helvum*	92	92	0
Chirundu	Tree	December 2020	*Rhinolophus spp.*	21	21	1
Total				**740**	**306**	**16**

All these sites were chosen according to the presence of bat colonies and the existence of anthropic activities. For instance, the selected caves and ancient mine are regularly visited by local people to collect bat guano, which is used as fertilizer, and/or to hunt bats for consumption. In Honde valley, frugivorous bats roosting sites in trees were close to maize crops or/and orchards.

All sites, except for three, were visited at different times during the sampling period (Table 1). The same sampling method was used at all sites and in every session as previously described by Bourgarel et al, [30]. Briefly, two square meters of plastic sheets were laid down at each site/per sampling session, underneath the bat colonies and left overnight. Six grams of fresh faeces from different bat individuals were collected from each plastic sheet in a 15 ml tube with 6 ml of RNA stabilization solution (*https://protocol-online.org/*). Back in the laboratory, samples were stored at -80°C for subsequent analysis. Of note, for all sampling sessions, personnel protection equipment (double pair of gloves, FFP3 mask, Tyvek overall, goggles, boots) were used in order to respect the biosafety procedures and to protect the sampling teams. Good biosafety practices training are regularly dispensed for our sampling teams.

Bat individual sampling

Individual bat samples which had already been collected from two study sites for a study on coronaviruses by Chidoti et *al*. were used in this study (Figure 1) [31]. These two sites, one cave (Magweto) and one building basement (Chirundu Farm) had been visited from August 2021 to July 2022 on a monthly basis (Figure 1, Table 2) [31]. Unfortunately, this study was conducted during the COVID 19 crisis and we were not able to access the study sites every month as planned owing to the imposed lock downs. Faecal samples had been collected by placing two square meters plastic sheets underneath the bat colonies. Only one faecal dropping (not contaminated by other faeces or urine) per 20 cm^2 was collected, assuming it represented one individual. Faeces were conserved individually in a 1.5 ml tube filled up with 0.5 ml of home-made RNA stabilization solution (*https://protocol-online.org/*) and stored at -80°C before further laboratory analyses.

Table 2: Prevalence of Astroviruses and co-infection with Coronaviruses and confidence intervals (CIs) per month at both Magweto and Chirundu sites in faecal samples from individual sampling of insectivorous bat communities and per reproduction cycle stage.

Site	Reproduction cycle	Month sampled	No of samples tested	No of Astrovirus positives	Prevalence (%) + CI (95%)	No of coinfection CoV and AstV	Prevalence (%) + CI (95%)
Chirundu	Non-gestation	August, 2020	153	0	0.0 (0.0-0.2)	0	0.0 (0.0-0.2)
Chirundu	Pregnancy	October, 2020	297	3	1.0 (0.3-2.9)	0	0.0 (0.0-1.3)
Chirundu	Parturition	November, 2020	241	3	1.2 (0.4-3.6)	1	0.4 (0.05-2.3)
Chirundu	Lactation	December, 2020	159	16	10.1 (6.3-15.7)	7	4.4 (2.1-8.8)
Chirundu	Weaning	February, 2021	170	29	17.1 (12.1-23.4)	18	10.6 (6.8-16.1)
Chirundu	4-6 Month juveniles	March, 2021	240	51	21.3 (16.5-26.8)	19	7.9 (5.1-12.0)
Chirundu	Non-gestation	May, 2021	243	47	19.3 (14.8-24.7)	9	3.7 (1.9-6.8)
Chirundu	Non-gestation	July, 2021	225	30	13.3 (9.5-18.4)	9	4.0 (2.1-7.4)
Overall prevalence			**1728**	**179**	**10.4 (9.0-11.8)**	**63**	**3.6 (3.8-5.8)**
Magweto	Non-gestation	September, 2020	348	18	5.2 (3.3-8.0)	1	0.3 (0.05-1.6)
Magweto	Pregnancy	October, 2020	257	46	17.9 (13.7-23.0)	7	2.7 (1.3-5.5)
Magweto	Parturition	November, 2020	228	31	13.6 (9.7-18.6)	3	1.3 (0.4-3.8)
Magweto	4-6 Month juveniles	March, 2021	242	46	19.0 (14.5-24.4)	20	8.3 (5.4-12.4)
Magweto	Non-gestation	April, 2021	242	26	10.7 (7.4-15.2)	6	2.5 (1.1-5.3)
Magweto	Non-gestation	June, 2021	242	47	19.4 (14.9-24.9)	3	1.2 (0.4-3.6)
Overall prevalence			**1559**	**214**	**13.7 (12.1-15.5)**	**40**	**2.6 (1.8-3.5)**

Periods of gestation (pregnancy), parturition, lactation, weaning and presence of 4-6 months old juveniles had been determined based on observations (from captures and observation of the roosting bats) combined with literature, as already reported in Chidoti et *al*. [31]. The reproductive season at Chirundu and Magweto site had been observed to begin in September to February for the predominant bat species. The different insectivorous bat families observed at both sites had been found to be synchronous regarding their reproductive cycles, and consensus reproduction periods had been determined based on the literature and observations [31,32].

Nucleic acid extraction and RT-PCR

Community samples.

Nucleic acids extraction was done from all faecal samples as previously described [30]. Briefly,

biological material (faeces) of three or four sample tubes collected from the same plastic sheet were

pooled and transferred in a 50 ml tube with 20 ml of PBS 1X then vigorously mixed. Tubes were

centrifuged at 4500 rpm for 10 min. The supernatant was first filtered using gauze in order to eliminate

faecal matter and transferred in fresh tubes then re-centrifuged at 4500 rpm for 10 min. The

supernatant was filtered through a 0.45 µm filter to remove eukaryotic and bacterial sized particles.

Seven millilitres of filtered samples were centrifuged at 250,000 g for 2.5 h at 4°C. The pellets were

re-suspended in 600 µl H_2O molecular grade and 150 µl were used to extract RNA and DNA using

NucleoSpin® RNA Kit (Macherey-Nagel, Hoerdt, France) according to the manufacturer's protocol.

Individual samples

Nucleic acids were extracted from 200 µl of faecal samples preserved in 0.5 ml RNA stabilization

solution using 5X MagMax Pathogen RNA/DNA Kit (ThermoFisher Scientific, Illkirch-

Graffenstaden, France), as already described in [30]. The faeces were vortexed vigorously (30Hz)

using Retsch MM400 Tissuelyser for 5 min to fully homogenise and mix the faecal particles, followed

by centrifugation at 16000 g for 3 min to fully separate the supernatant from the faecal debris. A

volume of 130µl of the supernatant was used for the isolation and purification stage of the nucleic

acids using Mag Max extraction kit with the automatic KingFisher Duo Prime Purification System

extractor (ThermoFisher Scientific, Illkirch-Graffenstaden, France). A final volume of 80µl of eluted

RNA/DNA was stored at -80◦C.

Bat species identification

For all bat samples that tested positive for AstVs, bat species identification was carried out by the

amplification and sequencing of mitochondrial C*ytochrome b* gene as previously described by Kocher

et *al.* [33]. Sequences were then compared to available bat sequences in the GenBank database using

Basic Local Alignment Search Tool (BLAST).

Astrovirus detection

Reverse Transcription (RT) using random hexamers were done on 5µl of RNA sample template using

1µl random hexamers, 0.5µl Oligo dT primer, 0.4µl of dNTPs (10mM) (ThermoFisher Scientific,

Illkirch-Graffenstaden, France) and 5.5µl molecular grade water incubated at 65°C for 5 min. This

was followed by addition of 4µl of Buffer 5X, 2µl of 0.1M DTT (M-MLV Reverse Transcriptase,

Invitrogen, ThermoFisher Scientific) and 1µl of RNAse OUT, incubated at 37°C for 2 min. A volume

of 1µl of M-MLV (M-MLV Reverse Transcriptase, Invitrogen, ThermoFisher Scientific, Illkirch-

Graffenstaden, France) reverse transcriptase was added to the mixture followed by incubation at 25ºC

for 10 min, 37ºC for 50 min and 70ºC for 15 min. The cDNA obtained was then used to partially

amplify the *Astrovirus RNA-dependent-RNA polymerase* gene (*RdRp*) by using a semi-nested Pan-

Astrovirus PCR system developed by Chu et *al* [16]. Visualization of positive PCR products was performed with agarose gel electrophoresis using SYBR green on a 1% gel. Positive PCR products (422 bp) were gel-agarose or directly purified (Geneclean Turbo Kit, MP Biomedicals, Illkirch-Graffenstaden, France) and then sequenced in both 5' and 3' directions (LGC, Berlin, Germany) by using Sanger method.

For the community approach, purified PCR products were cloned by using Topo PCR Cloning kit according to the manufacturer's protocol (ThermoFisher Scientific, Illkirch-Graffenstaden, France). Ten clones per PCR product were sequenced in both 5' and 3' direction using the Sanger method (Eurofins, Germany).

Phylogenetic analyses

Bat AstVs overlapping sequences were assembled into contiguous sequences using Geneious software package V. 2021.2.2 (Biomatters Ltd, Auckland, New Zealand). Partial non-concatenated nucleic acid sequences of the new AstVs were aligned using Clustal W [34] implemented in MEGA 7 [35], with minor manual adjustments. Ambiguously aligned and divergent regions were excluded from subsequent analyses. Phylogenies were inferred using Maximum Likelihood (ML) method implemented in PhyML V. 3.0 [36]. The suited evolution model was selected by Akaike's Information criterion (AIC) using Topali software [37]. The reliability of branching orders was tested using the bootstrap approach (1000 replicates) [38] and the GTR + F+ I substitution model was determined as the best suited evolution model.

Temporal variations of Astrovirus prevalence and bat reproductive phenology

In the same way as in Chidoti et *al*, the prevalence of astrovirus infection was calculated at the community level for small insectivorous bat species from the two longitudinal study sites, Magweto and Chirundu farm [31]. The proportion of RNA AstV-positive samples were estimated per month and site with 95% confidence intervals (CI) using Wilson score test *(https://epitools.ausvet.com.au/ciproportion)* [39].

The influence of the different phases of the bat reproductive cycle stages (periods of pregnancy, parturition/lactation, weaning, weaned juveniles of 4 to 6 months old) and of *Coronavirus* infection/shedding on the prevalence of astroviruses was tested by running a generalized linear mixed model (GLMM) for each site (Magweto and Chirundu) as described in Chidoti et *al* [31]. Parturition and lactation were analysed as one season because the observations did not allow the separation of the two periods as they were interlinked. The response variable with a binomial distribution was the AstVs PCR result of the samples, and the explanatory variables with fixed effects were the different phases of the reproduction cycle. Given Coronavirus PCR results on the same samples were available

through our previous study (see below), Coronavirus status was also integrated into the model as fixed variable to investigate any effect of co-infection on AstVs infection/shedding. The reproductive phases were coded as 1 if it was during the corresponding reproduction phase and 0 if it was not. A session identification code was included as a random effect to control for repeated measures from the same sampling session to account for clustered samples collected.

Astrovirus and Coronavirus co-infection

Coronavirus data, produced by Chidoti et *al* [31], on the same sample sets were used in this study to assess the AstVs and CoVs co-infection in bat communities from Magweto and Chirundu sites and a descriptive analysis of prevalences was carried out (**Table 2**).

Results

Astrovirus detection in pooled sampling sites

The sampling was carried out from June 2016 to December 2020 (Table 1) at the selected sites. At Magweto cave site, 24 samples were collected. In Bashongwe collection was carried out in March, July and December 2018, and 251 samples were collected. For Mabura cave site, a total of 58 samples were collected, in 2016 and 2017, with sampling only carried out once in both years. In Mugunza cave site, a once off sampling session was done in 2017 and a total of 50 samples were collected. In the Honde Valley site, 336 samples were collected under nine *Eidolon helvum* roost trees (Table 1). In Magweto, five of the eight pooled samples (62.5%) tested positive for AstV. In Bashongwe cave, all the 72 samples collected tested negative for AstV (Table 1). In Mabura, Six (31.6%) out of 19 pooled samples tested positive for AstVs and all of the 13 samples from Mugunza tested for AstVs were negative (Table 1). In Honde Valley, 173 pooled samples were tested for AstVs and only four (2.3%) were positive (Table 1). In Chirundu Baobab tree only one (4.8%) out of the 21 samples collected was positive for AstV.

Prevalence and seasonality of Astroviruses at Chirundu and Magweto site

In total, we analyzed 1559 samples from Magweto cave site, and 1728 samples from Chirundu farm. (Table 2). The overall prevalence of AstVs at bat community level at Chirundu was 10.4% [95% CI: 9.0-11.8] (179 positives out of the 1728 samples). The highest prevalence of 21.3% [95% CI: 16.5-26.8] was observed in March 2021 and corresponded to the 4-6 months old juvenile period (Table 2). The prevalence of AstV increased from December 2020 to March 2021, which corresponded to the lactation period, followed by the weaning of four to six months old juveniles. The lowest prevalence was observed in August (0%), October (1% [95% CI: 0.3-2.9]) and November (1.2% [95% CI: 0.4-3.6]), which correspond to the transition between non-gestation and gestation period, as well as the beginning of the parturition period (Table 1, Figure 2a).

Figure 2:

Bat community AstVs prevalence per month at Magweto site. **Fig 2a & 2b** show estimation of the astrovirus prevalence (with CI 95%) at Chirundu (a) and Magweto (b) sites respectively for the months and year sampled. The graphs were plotted with y-axes as the percentage of AstVs prevalence and x-axis as the sampling collection date and the corresponding reproductive cycle stage, each colour code representing a different season.

In Magweto site, the overall prevalence of Astroviruses for the insectivorous bat community was 13.7% [95% CI:12.1-15.5] (214 positives out of 1559 samples). There was no clear pattern for the prevalence of AstVs according to reproductive cycle stages in this site (Table 2, Figure 2b). The highest prevalence of 19.4% [95% CI: 14.9-24.9] was observed in June 2021 during the non-gestation period , and very close prevalences observed during the pregnancy (17.9% [95% CI: 13.7-23]) and 4-6 months old juvenile (19% [95% CI: 14.5-24.4]) periods (Table 2). Lower prevalence was observed in September at the end of the non-gestation period (5.2% [95% CI: 3.3-8]).

Results from the GLMM didn't show any effect of bats seasonal reproduction periods on detection of AstV RNA in samples collected at Magweto site (Table 3). For Chirundu site, the GLMM showed significantly higher probability of being positive to astrovirus during the period of weaned 4-6 months old juveniles (odds ratio (OR) = 5.088, 95% CI = 1.33- 36.32, p= 0.016), and significantly lower probability of AstV positivity during the pregnancy period (odds ratio (OR) = 0.136, 95% CI = 0.03-0.81, p=0.006) (Table 3).

Table 3: Results of the GLMM with the following explanatory variables for both site (Chirundu and Magweto): PCR_Pan-*Coronavirus*, Pregnancy, Weaning, Weaned juveniles 4 to 6 months. With references OR odds ratio, CI95 95% confidence interval.

Variables	Odds ratios	p-value	OR CI95
Chirundu Site			
intercept	0.05	6.68e-09	0.010- 0.121
PCR_Pancoronavirus	1.116	0.536	0.785 -1.574
Pregnancy	0.14	0.006	0.027-0.810
Weaning	3.91	0.094	0.721-39.593
Weaned_juveniles 4 to 6 months	5.09	0.016	1.333-36.322
Magweto Site			
intercept	0.235	0.0005	0.088-0.627
PCR_Pancoronavirus	1.238	0.290	0.825-1.825
Pregnancy	0.473	0.145	0.138-1.548
Weaned_juveniles 4 to 6 months	0.678	0.448	0.202-2.256

Prevalence and seasonality of coinfection with Astroviruses and Coronavirus at Chirundu and Magweto site

The co-infection prevalence in bat communities from Chirundu and Magweto sites was relatively low.
The co-infection prevalence was observed to follow a similar trend as that of CoVs and AstVs
prevalences, where it showed peaks that correlated with peaks of the individual viral families (Figure
3a & b). The overall prevalence of co-infection at Chirundu site was 3.6% [95% CI: 3.8-5.8] (Table
2) with the highest co-infection prevalence of 10.6% [95% CI: 6.8-16.1] observed in February 2021
during the weaning period. The lowest prevalences of 0% were observed in August [95% CI: 0.0-0.2]
and October [95% CI: 0.0-1.3] during the end of the non-gestation and the pregnancy period,
respectively (Table 2, Figure 3a).
The overall prevalence of co-infection at Magweto site was 2.6% [95% CI: 1.8-3.5] (Table 2, Figure
3b). Co-infection in this site was recorded from September 2020 to December 2020 and from March
2021 to June 2021 (Table 2, Figure 3b). The highest prevalence of co-infection was 8.3% [95% CI:
5.4-12.4] in March during the 4-6 months old juvenile period; while the lowest 0.3% [95% CI: 0.05-
1.6] was observed in September 2020 during the non-gestation period (Table 2, Figure 3b).
296
297 **Figure 3***:* Prevalence of *Astrovirus*, *Coronavirus* and co-infections involving both viruses per month: co-
298 infections are represented in green, coronaviruses in blue and astroviruses in red for a/ Chirundu site and b/
299 Magweto site according to reproduction cycle stages.
300
301

Astroviruses genetic diversity

All AstV positive samples (N=423) were sequenced, representing 179 samples from Chirundu, 229 from Magweto, seven from Mabura, seven from Honde valley, and one from Baobab tree. Of the 423 samples, 214 (50.6%) were amplified from *Hipposideros* spp., 63 (14.9%) from *Rhinolophus* spp., 53 (12.5%) from *Miniopterus* spp., 10 (2.4%) from *Nycteris* spp., seven (1.7%) from *Eidolon* spp. and for the remaining 76 samples known to originate from bats for which we couldn't determine the genus/species.

A phylogenetic analysis was conducted with 155 partial sequences generated during this study and representative of the genetic diversity observed. All clustered within the *Mamastrovirus* genus (Figure 4).

Two sequences isolated from Magweto cave, one from *Rhinolophus* spp. (MAG-573) and 1 from *Hipposideros* spp. (MAG-292) clustered with a group of Human astroviruses (*HuAstVs*) (Figure 4). This cluster was well supported (Bootstrap >90%) and the nucleotide acid identities between Bat Astroviruses MAG-573, MAG-292 and *HuAstV*-2 were 96 and 93% respectively (Data not shown). The other specific cluster of bat astrovirus showing a close relation to *HuAstVs* is a sequence isolated from *Hipposideros* spp. (MAG-1236) in Magweto, it clustered with *HuAstV-MLB-1, -2* and *3* (Figure 4). Six sequences isolated from the frugivorous bats *E. helvum* found in Honde Valley and one from *Rhinolophus* spp. collected in a Baobab tree formed a sister clade with another group of Human astroviruses *HuAstV-VA-1, -2, -3, -4* and *HuAstV-HMO-A* (Figure 4). In this study, the majority of the sequences were closely related to astroviruses identified in insectivorous bats (Figure 4). These were mainly amplified from *Hipposideros* spp. (HC-1 to -9) and *Rhinolophus* spp. (Rhi-1 to -4) bat species. One cluster comprised specifically of sequences isolated from *Rhinolophus* spp. Most of these

sequences were isolated from Chirundu and a few of them from Magweto, and clustered together with AstV strains from *Nyctalus* spp. and *Verpitillio* spp. bat species isolated in Czech Republic (references Y & Z in the phylogenetic tree) (Figure 4). Another cluster comprising of the majority of AstV sequences from Chirundu and some from Magweto site, were characterized from *Miniopterus* spp. and *Rhinolopus* spp. All together, they formed a phylogenetic cluster with strains of bat astroviruses isolated from *Miniopterus* spp. from Madagascar and China (references N, O, R, U, T), *Rhinolophus* spp. from Korea (references P, Q) as well as *Rousettus* spp. and *Paratrianops* spp. Bat AstVs *(BAstV)* from Madagascar (references V, W) (Figure 4)

Two major clades from Chirundu and one from Magweto, isolated from *Miniopterus, Rhinolophus* and *Hipposideros* spp. of bats also phylogenetically clustered with a strain from Mozambique isolated from *Nycteris thebaica* (reference M). One sequence from Magweto isolated from *Rhinolophus* spp. and two from Mabura cave isolated from *Hipposideros* spp*.,* clustered with a strain of bat astrovirus isolated from *Myotis* spp. in Madagascar (references K, L) (Figure 4). A cluster, comprising two clades from Magweto and one sequence from Chirundu and three sequences from Mabura cave isolated from *Rhinolophus* and *Hipposideros,* were closely related to bat astrovirus strains from *Hipposideros* spp. of Mozambique and Laos (references H, I) (Figure 4). The last small group comprising one clade from Chirundu, four sequences from Magweto, all isolated from *Rhinolophus* spp*.,* and one sequence from Mabura isolated from *Hipposideros* spp. clustered together with bat astrovirus strains from *Rhinolophus* spp. of Laos and Korea (references D, E, F) (Figure 4).

One of the large clusters shows the phylogenetic relationship amongst sequences derived from, *Hipposideros* spp. (Clades HC-5 to -8), these making up the majority of the sequences in this clade and one isolated from *Miniopterus* spp., one from *E. helvum* and one from a *Nycteris* spp.. The bat astroviruses in this clade, all showed a close relation to a bat astrovirus strain isolated from *Hipposideros spp.* of Mozambique (reference C). The largest cluster constituted mainly by Magweto site sequences and some Chirundu sequences, from *Hipposideros* spp. (clades HC-1 to 4) and a few *Rhinolophus* and *Nycteris* spp., clustered with a strain from *Hipposideros* spp. of Mozambique (references A, B) (Figure 4).

HC-1

MAG-1022 HC
MAG859 RS
CHI-1161 HC
CHI-1224 NT
MAG-938 HC
MAG-862 Rhi spp.

HC-2

MAG-883
MAG-845 HC

HC-3

CHI-1213 HC
MAG-556 RS
CHI-1122 HC
MAG-617

HC-4

MAG-661
MAG-1271 HC
MAG-1119 HC

HC-5

CHI-1207 NT

HC-6

CHI-1123
CHI-1235 HC
HV-031 EH

HC-7

MAG-1183 HC
CHI-1320 Min spp.

HC-8

MAB-038a HC

Rhi-1

D
E
F
G

HC-9

H

Rhi-2

MAB-023c HC

I
J

MAB-021a HC
MAG-09 RS
MAB-038c HC

K
L

CHI-1105
CHI-1045
CHI-1048 HC
CHI-1068 HC

M

MAG-036a HC
MAG-573 Rhispp.
MAG-197 Min spp.
MAG-591 Rhispp.
CHI-1275 HC

Rhi-3

CHI-1717 Min spp.
MAG-595
CHI-1476
CHI-1477
CHI-1471
CHI-1726 Min spp.
CHI-1139 Rhi spp.

N
O
P
Q
R
S
T
U
V
W

CHI-1303 Min spp.
CHI-1449 Min spp.
MAG-1143 Rhi spp.
CHI-1551

Rhi-4

X

MAG-580 Min spp.
MAG-425
CHI-450 Rhi spp.
CHI-425 Rhi spp.
MAG-568 Rhi spp.
CHI-1704
CHI-1565
CHI-1569

Y
Z
AA

VA2
HMO-A
VA4
VA1
VA3

HuAstV

BAOB-016 Rhi pp.

AB

EH

AC

AD

MBL3
MBL1
MBL2
MAG-1236 HC

HuAstV

MAB-033 HC
MAG-1292 HC

AE
AF

MAG-573 Rhispp.
MAG-292 Rhispp.

2
4
6
3

HuAstV

4.0

PoAstV
BDoAstV
CSLAstV
Ch AstV

356
357

Figure 4: **Phylogenetic tree of** *Astroviruses* **partial** *RdRp* **gene**. The sequences detected at Chirundu site are represented
by red branches, at Magweto site by blue branches, at Honde Valley by green branches, at Mabura cave by brown branches
and at Baobab tree site by an orange branch. Human *Astrovirus* are highlighted in red rectangles and bold letters represent
the different bat AstV references used to build this tree (See Supplementary Data: Figure 4). The tree was built using the
maximum likelihood method based on the GTR + G4 + I model. The robustness of nodes was assessed with 1000 bootstrap
replicates. Bootstrap values >70 are represented by asterisks and those <70 are not shown. HC=*Hipposideros caffer*;
RS=*Rhinolophus simulator*; NT=*Nycteris thebaica*; EH= *Eidolon helvum*; Min=*Miniopterus* spp.; Rhi=*Rhinolophus* spp..
MAG=Magweto site; CHI=Chirundu site; BAOB= Baobab; MAB= Mabura site; HV=Honde valley site. The scale bar
represented the number of substitutions per site. The AstV sequences used as references are detailed in supplementary
data: DOI **10.5281/zenodo.8399361**
368

Discussion

The overall AstVs prevalence was respectively 10.0% and 13.7% for Chirundu and Magweto sites,
confirming their circulation in bat populations from Zimbabwe. However, compared to studies in
other African countries, the observed prevalence was low: Hoarau et *al* and Lebarbenchon et *al*
detected AstV in 20-22% of individual bats in Mozambique and Madagascar, respectively [21,23].
This difference in prevalence of AstV compared to other countries can be attributed to different study
designs, longitudinal sampling in our case versus transversal studies in Mozambique and Madagascar
[21,23]. Furthermore, these differences could also result from the studied bat species having varying
immunological responses and susceptibilities to different viruses [40], and the different localities
resulting in subtle changes that may impact the infection dynamics.

We also observed difference between the two main sites studied (Chirundu and Magweto). Further-
more, the differences in prevalence between the two main individual populations can be attributed
but not limited to differences between habitats, bat population structures and bat species composition
of bat communities and also anthropogenic factors. Therefore there is need for a more comprehensive
study, covering several years of follow up, to fully establish the factors contributing to the AstV viral
dynamics among bat populations at these two sites.

At Chirundu site increasing RNA-AstV prevalences were observed from the lactation to the 4-6
months old juvenile seasons with 21.2% of RNA-AstV detection during the latter. Astroviruses have
been described to display seasonal variations in prevalence [23]. Drexler et *al* observed different
peaks of astrovirus detection associated with different stages of the reproductive season with peaks
correlating to maternal aggregations during breeding season and after parturition season due to
establishment of a susceptible sub-population of weaned new-born bats who did not yet have their
own adaptive immunity [24]. In our study the prevalence was high during some of the months
compared to overall prevalence, which might be related to the effect of reproductive cycle stages on
the shedding of viruses, as demonstrated for other viruses. In another study the reproductive season
with juveniles and immature individuals showed a very high prevalence of CoV as compared to
prevalences in the absence of juveniles and presence of sexually mature bats thus suggesting different
stages of the reproduction having effect on viral shedding [31, 41–43]. In this study we observed a

similar trend for AstVs for one of the study sites. Furthermore, Mendenhall et *al*, also identified the
bat juvenile stage as exhibiting a greater AstVs viral burden than any other stage of the reproductive
season [43]. The range of prevalence observed during the highest peaks coincides with the ≥20%
overall AstV detection reported by Hoarau et *al* [21] during the time of their sampling in February
and May. Here we observe a peak during the 4-6 months old juvenile season in Chirundu site which
corresponds to a high influx of immunologically immature individuals in the bat population, similar
to what was observed in the Chidoti et *al*, (2022) for CoV infection [31].

Juveniles and immature individuals are known to contribute to shedding pulses of viruses as they
develop productive infections during the acute phase [27], which coincides with the waning
protection of maternal antibodies [31], thus increasing the susceptibility and rate of infection in the
young.

We could not observe a clear trend at Magweto site, which might be related to the absence of samples
from December 2020 to March 2021, which correspond to the lactation and weaning periods.

In our study we also investigated co-infection of bats by coronaviruses and astroviruses. We compared
the co-infection prevalence with the prevalence of astroviruses and coronaviruses described at each
site during the same reproduction cycle stages. The co-infection prevalence showed a similar trend to
that observed in the individual viruses, whereby during observed peaks of high AstVs and CoVs
detections, the co-infection also peaked at similar phases of the reproduction cycle stages. Thus the
co-infection of bats by both CoVs and AstVs is high when there is high infection of bats by either
virus.

The overall prevalence of co-infection was 3.5% and 2.6 % in Chirundu and Magweto respectively
while the bats from Madagascar and Mozambique respectively showed a AstV-CoV co-infection of
5% (±2.7%) and of 1.7% (±1,1%) [9,40]. Co-infection and recombination of viruses in bats have been
reported on several occasions including co-infection of bats by coronaviruses and astroviruses
[9,40,44]. Bats infected with either CoV or AstV were shown to be more likely co-infected with the
respective virus [26]. In our study, no effect of coronavirus infection/shedding on astrovirus
infection/shedding was observed.

In the current study we detected bat Astroviruses from *Hipposideridae, Rhinolophidae, Pteropodidae,*
*Nycterida*e and *Miniopterida*e families. Astroviruses are known to show no host restriction and are
widespread within the Chiroptera order [16,24]. The majority of the astroviruses amplified were from
Hipposideros and *Rhinolophus* species and these clustered with AstV sequences derived from
Mozambique, Madagascar and China of the same families [21,23,24]. The observed trend is due to
the two genera being the most dominant species of bats at both sites where sampling was done.
Therefore, the probability of higher detection of AstV in these species is expected and evidently
observed to be higher than other rare or less dominant species. Active transmission events of

astroviruses amongst *Rhinolophidae* and *Hipposideridae* bat species are known to occur [24]. In this study, a high degree of species-specific tropism, especially in bat astrovirus related clusters was observed with astrovirus strains isolated from *Hipposideros* spp. clustering with each other and similar trends for *Rhinolophus* and *Miniopterus* spp. However, the clusters were not site-specific as sequences from Magweto, Mabura and Chirundu sites formed clusters with each other. We also observed a large phylogenetic cluster which comprised of specific *Hipposideros and Rhinolophus* AstV sub-clades (HC-1 to -8 and Rhi-1 to -4) as well as a few AstV sequences from other bat species (Figure 4). Previous studies on astroviruses in bats have also described this family of viruses to be involved in species tropism. However recent studies challenged this assumption and highlighted the fact that AstV cross-species transmissions are more frequent than previously thought [4,45]. In our study, AstVs sequences isolated from different bat species of the same site clustered together. This indicates potential cross species transmission of astroviruses within each site, as reported by Xiao et *al* in bats sharing the same habitat [45]. We also observed phylogenetic clustering of astroviruses from Chirundu and Magweto sites into same clades showing a great diversity of astroviruses in circulation within the studied bat communities. Novel astroviruses species have been reported in bats hosts [25], and in this study it is evident due to the diverse astroviruses described. The AstV isolated from *E. helvum*, frugivorous bats, clustered together and showed close phylogenetic relation to *BAstV* isolated from *Rousettus* spp. bats in China and *Taphozous* spp. bats in Laos. One strain isolated from *E. helvum* clustered with *Hipposideros* bat astroviruses sequences isolated in Chirundu site. This could mean that frugivorous bats and insectivorous bats can be infected by the same type of AstVs strain thus indicative of the multi-host spectrum for the Astroviridae family. Detection of AstVs in the frugivorous bats was low compared to insectivorous bats, and this can be attributed to the insectivorous bats ability to adapt and harbour more astroviruses than frugivorous bats [45].

Two strains, Mag-292 *H. Caffer*, Mag-573 *Rhinolophus* spp. clustered within the *HuAstv* type 1 to 6, that are known to cause infection and diarrhoea in children and infants [1]. Six of our strains isolated from *E. helvum* and one from *Rhinolophus* spp. formed sister clades to *HuAstV* HMO, associated with severe extra-intestinal illnesses in humans [5] and VA-1-4 strains. Another AstV strain detected in *Hipposideros* bat species clustered with *HuAstv* MLB-1, -2 and -3 which are all known to cause gastro-intestinal and neurological diseases with mild-severe symptoms in humans [46]. This phylogenetic clustering could suggest that *HuAstV* and *BAstV* may have a shared common ancestor. However, with limited sequence data the evolutionary history is merely a speculation that needs to be further investigated through full genome analysis. Another hypothesis for the observed relation is that, since bats are known to be important reservoir hosts of astroviruses with great diversity, they could be ancestors of the *HuAstV* due to the high prevalence of AstV detected in bats. *HuAstV* transmission occurs via the faecal-oral route and contaminated food or water [1], therefore it is unlikely that

anthroponosis may have occurred. However, this observed relation could be due to a rare event or
mutation, or a new strain of bat astrovirus that has not been described yet. Further full genomic data
as well as epidemiological data on these bat colonies are needed to reach a full conclusion.As novel
strains and host species continue to be discovered, understanding AstV transmission and their
zoonotic potential is essential. Dissemination of *HuAstV* and AstV from zoonotic reservoir hosts
remain a significant threat to public health [7].

Conclusion

This study offered important contributions to the understanding of virus diversity in an under-sampled
region of the world, and the sequences contribute to a better picture of co-evolution and transmission
among bat species. Multiple sampling points as in the study further give a more comprehensive pic-
ture of how viruses vary over space and time and host species. Astroviruses have a high genetic
diversity, multiple mechanism of generating additional diversity, and infect a wide range of host spe-
cies therefore understanding their prevalence and infection dynamics in their wildlife hosts can help
to predict or prevent the emergence of novel astrovirus strains into domestic animals and human
populations.

Acknowledgments: We thank Billy Butete for his field assistance. The authors gratefully
acknowledge and thank the International Atomic Energy Agency for making available the sequencing
services. We thank the Research Council of Zimbabwe for approving this study (research registration
certificate N° 03006) and the Hurungwe Rural District council for their assistance and facilitation.
We thank the Animal Research Ethics Committee of Zimbabwe for their approval (ref number
002/2017). This work was conducted within the framework of the Research Platform "Production
and Conservation in Partnership" (RP–PCP).

Funding: This work was supported by grants of the French Ministry of Europe and Foreign Affairs
(Fond de Solidarité pour les Projets Innovants, les sociétés civiles, la francophonie et le développe-
ment humain—CAZCOM Project, FSPI N°2019/88) and the AFD (French Agency for development-
Projet PACMAN, AFD CWZ 1019 01 V).

Conflicts of Interest: We declare that we have no conflicts of interest

Author's contribution: Conceptualization: VC, MA, HdN, MB, LG and FL.; **Methodology**: VC,
MA, LG, HdN, DP, EG, GM, EM and FL.; **Software:** VC, MA, HdN and FL; **Validation**: VC, MA
HdN, JC, and FL; **Formal Analysis**: VC, HdN, JC, and FL.; **Investigation**: VC, VP, NC, GM, and
FL.; **Resources**: DP, GM, MP, MB, HdN, LG, and FL.; **Data Curation**: VC, HdN, and FL.; **Writing**
– Original Draft Preparation: VC and FL; **Writing – Review & Editing**: HdN, MA, JC, MB, LG,
VP, EG, DM, DP, EM, GM and FL.; **Supervision**: HdN, DP, GM and FL.; **Project Administration**:
LG, MB, HdN and FL; **Funding Acquisition**: MB, HdN, LG and FL.

Data, scripts, code, and supplementary information availability

Sampling Data, Astrovirus sequences, Graphs and Figures

https://doi.org/10.5281/zenodo.8399450

https://doi.org/10.5281/zenodo.7849008
506
GenBank Accession Numbers
The AstVs sequences have been deposited to the GenBank under the following numbers: OQ271049
- OQ271203
The Cytochrome B sequences have been deposited to the GenBank under the following numbers:
OM487705-OM488020

Statistical analysis

The script used for the GLMM analysis is available at: https://doi.org/10.5281/zenodo.7847934

Supplementary data: https://doi.org/10.5281/zenodo.8399361
515

Reference

[1] Bosch A, Pintó RM, Guix S. Human Astroviruses. Clin Microbiol Rev 2014;27:1048–1074. doi/10.1128/CMR.00013-14.

[2] Donato C, Vijaykrishna D. The Broad Host Range and Genetic Diversity of Mammalian and Avian Astroviruses. Viruses 2017;9:102. doi/mdpi.com/1999-4915/9/5/102.

[3] ICTV. ICTV Positive Sense RNA Viruses: *Astroviridae*. VIRUS Taxon. 2020; https://talk.ictvonline.org.

[4] Boujon CL, Koch MC, Seuberlich T. The Expanding Field of Mammalian Astroviruses: Opportunities and Challenges in Clinical Virology. Adv Virus Res 2017. p. 109–137. doi/S0065352717300143.

[5] Wohlgemuth N, Honce R, Schultz-Cherry S. Astrovirus evolution and emergence. Infect Genet Evol 2019;69:30–37. doi/S1567134818307226.

[6] Wu H, Pang R, Cheng T, et al. Abundant and Diverse RNA Viruses in Insects Revealed by RNA-Seq Analysis: Ecological and Evolutionary Implications. Cristea IM, editor. mSystems 2020;5. doi/10.1128/mSystems.00039-20.

[7] Roach SN, Langlois RA. Intra- and Cross-Species Transmission of Astroviruses. Viruses 2021;13:1127. doi/mdpi.com/1999-4915/13/6/1127.

[8] El Taweel A, Kandeil A, Barakat A, et al. Diversity of Astroviruses Circulating in Humans, Bats, and Wild Birds in Egypt. Viruses 2020;12:485. doi/mdpi.com/1999-4915/12/5/485.

[9] Fischer K, Zeus V, Kwasnitschka L, et al. Insectivorous bats carry host specific astroviruses and coronaviruses across different regions in Germany. Infect Genet Evol 2016;37:108–116. doi/S156713481530040X.

[10] Frémond M-L, Pérot P, Muth E, et al. Next-Generation Sequencing for Diagnosis and Tailored Therapy: A Case Report of Astrovirus-Associated Progressive Encephalitis. J Pediatric Infect Dis Soc 2015;4:e53–e57. doi/10.1093/jpids/piv040.

[11] Cortez V, Meliopoulos VA, Karlsson EA, et al. Astrovirus Biology and Pathogenesis. Annu Rev Virol 2017;4:327–348. https://www.annualreviews.org/doi/10.1146/annurev-virology-101416-041742.

[12] Janowski AB, Klein RS, Wang D. Differential In Vitro Infection of Neural Cells by Astroviruses. Griffin DE, editor. MBio 2019;10. doi/10.1128/mBio.01455-19.

[13] Vu D-L, Cordey S, Brito F, et al. Novel human astroviruses: Novel human diseases? J Clin Virol2016;82:56–63. doi.org/10.1016/j.jcv.2016.07.004.

[14] Johnson CK, Hitchens PL, Pandit PS, et al. Global shifts in mammalian population trends reveal key predictors of virus spillover risk. Proc R Soc B Biol Sci. 2020;287.

[15] Baker KS, Todd S, Marsh GA, et al. Novel, Potentially Zoonotic Paramyxoviruses from the African Straw-Colored Fruit Bat Eidolon helvum. J Virol. 2013;87:1348–1358.

[16] Chu DKW, Poon LLM, Guan Y, et al. Novel Astroviruses in Insectivorous Bats. J Virol 2008;82:9107–9114. doi/10.1128/JVI.00857-08.

[17] Dufkova L, Straková P, Širmarová J, et al. Detection of Diverse Novel Bat Astrovirus Sequences in the Czech Republic. Vector-Borne Zoonotic Dis 2015;15:518–521. doi/10.1089/vbz.2015.1813.

[18] Rougeron V, Suquet E, Maganga GD, et al. Characterization and phylogenetic analysis of new bat astroviruses detected in Gabon, Central Africa. Acta Virol 2016;60:386–392.doi/10.4149/av_2016_04_386.

[19] Fischer K, Pinho dos Reis V, Balkema-Buschmann A. Bat Astroviruses: Towards
Understanding the Transmission Dynamics of a Neglected Virus Family. Viruses
2017;9(2):34.doi.org/10.3390/v9020034.

[20] Lacroix A, Duong V, Hul V *et al.* Diversity of bat astroviruses in Lao PDR and Cambodia,
Infect Genet Evol 2017;47:41-50.doi:10.1016/j.meegid.2016.11.013.

[21] Hoarau F, Le Minter G, Joffrin L, et al. Bat Astrovirus in Mozambique. Virol J 2018;15:104.
doi/10.1186/s12985-018-1011-x.

[22] Waruhiu C, Ommeh S, Obanda V, et al. Molecular detection of viruses in Kenyan bats and
discovery of novel astroviruses, caliciviruses and rotaviruses. Virol Sin 2017;32:101–114.
doi/10.1007/s12250-016-3930-2.

[23] Lebarbenchon C, Ramasindrazana B, Joffrin L, et al. Astroviruses in bats, Madagascar. Emerg
Microbes Infect 2017;6:1–3. doi/full/10.1038/emi.2017.47.

[24] Lee S-Y, Son K-D, Yong-Sik K, et al. Genetic diversity and phylogenetic analysis of newly
discovered   bat   astroviruses   in   Korea.   Arch   Virol   2018;163:3065–3072.
doi.org/10.1007/s00705-018-3992-6.

[25] Bergner LM, Mollentze N, Orton RJ, et al. Characterizing and Evaluating the Zoonotic Poten-
tial of Novel Viruses Discovered in Vampire Bats. Viruses 2021;13:252. doi/1999-
4915/13/2/252.

[26] Seltmann A, Corman VM, Rasche A, et al. Seasonal Fluctuations of Astrovirus, But Not Coro-
navirus Shedding in Bats Inhabiting Human-Modified Tropical Forests. Ecohealth
2017;14:272–284. doi:10.1007/s10393-017-1245-x.

[27] Plowright RK, Peel AJ, Streicker DG, et al. Transmission or Within-Host Dynamics Driving
Pulses of Zoonotic Viruses in Reservoir–Host Populations. Remais J V., editor. PLoS Negl
Trop Dis 2016;10:e0004796. doi:10.1371/journal.pntd.0004796.

[28] Drexler JF, Corman VM, Wegner T, et al. Amplification of emerging viruses in a bat colony.
Emerg Infect Dis 2011;17(3):449-56.doi/10.3201/eid1703.100526

[29] Amman BR, Nyakarahuka L, McElroy AK, et al. Marburgvirus resurgence in Kitaka Mine
bat population after extermination attempts, Uganda. Emerg Infect Dis. 2014;20(10):1761-4.
doi:10.3201/eid2010.140696.

[30] Bourgarel M, Pfukenyi DM, Boué V, et al. Circulation of Alphacoronavirus, Betacoronavirus
and Paramyxovirus in Hipposideros bat species in Zimbabwe. Infect Genet Evol 2018;58:253–
257. doi.org/10.1016/j.meegid.2018.01.007.

[31] Chidoti V, De Nys H, Pinarello V, et al. Longitudinal Survey of Coronavirus Circulation and
Diversity in Insectivorous Bat Colonies in Zimbabwe. Viruses 2022;14:781. doi/1999-
4915/14/4/781.

[32] Monadjem A, Taylor PJ, Cotterill FPD, et al. Bats of Southern and Central Africa: A Biogeo-
graphic and Taxonomic Synthesis. WITS Press. 2011.

[33] Kocher TD, Thomas WK, Meyer A, et al. Dynamics of mitochondrial DNA evolution in ani-
mals: amplification and sequencing with conserved primers. Proc Natl Acad Sci
1989;86:6196–6200. doi/full/10.1073/pnas.86.16.6196.

[34] Higgins DG, Sharp PM.CLUSTAL: a package for performing multiple sequence alignment on
a microcomputer. Gene,1988;73(1):237-244.doi/10.1016/0378-1119(88)90330-7.

[35] Kumar S, Stecher G, Tamura K. MEGA7: Molecular Evolutionary Genetics Analysis Version
7.0 for Bigger Datasets. Mol. Biol. Evol. 2016;33:1870–1874. doi/33/7/1870/2579089.

[36] Guindon S, Dufayard J-F, Lefort V, et al. New Algorithms and Methods to Estimate Maximum-Likelihood Phylogenies: Assessing the Performance of PhyML 3.0. Syst. Biol. 2010;59:307–321.doi/10.1093/sysbio/syq010.

[37] Milne I, Lindner D, Bayer M, et al. TOPALi v2: a rich graphical interface for evolutionary analyses of multiple alignments on HPC clusters and multi-core desktops. Bioinformatics 2009;25:126–127. doi/25/1/126/302670.

[38] Lemoine F, Wilkinson E, Correia D, et al. Renewing Felsenstein's Phylogenetic Bootstrap in the Era of Big Data. Nature. 2018.

[39] Wilson E. Probable Inference , the Law of Succession , and Statistical Inference. J Am Stat Assoc. 1927;22:209–212.

[40] Hoarau A, Köster M, Dietrich M, et al. Synchronicity of viral shedding in molossid bat maternity colonies. Epidemiol Infect. 2023;151:1–7.

[41] Wacharapluesadee S, Duengkae P, Chaiyes A, et al. Longitudinal study of age-specific pattern of coronavirus infection in Lyle's flying fox (Pteropus lylei) in Thailand. Virol J. 2018;15:1–10.

[42] Cappelle J, Furey N, Hoem T, et al. Longitudinal monitoring in Cambodia suggests higher circulation of alpha and betacoronaviruses in juvenile and immature bats of three species. Sci Rep 2021;11:1–11. doi.org/10.1038/s41598-021-03169-z.

[43] Mendenhall IH, Skiles MM, Sena E, et al. In fl uence of age and body condition on astrovirus infection of bats in Singapore : An evolutionary and epidemiological analysis. One Heal. 2017;4:27–33.

[44] Hoarau AOG, Goodman SM, Halabi D Al, et al. Investigation of astrovirus , coronavirus and paramyxovirus co-infections in bats in the western Indian Ocean. Virol J. 2021;18:1–8. doi.org/10.1186/s12985-021-01673-2.

[45] Xiao J, Li J, Hu G, et al. Isolation and phylogenetic characterization of bat astroviruses in southern China. Arch. Virol. 2011;156:1415–1423. doi/10.1007/s00705-011-1011-2.

[46] Kapoor A, Li L, Victoria J, et al. Multiple novel astrovirus species in human stool. J Gen Virol 2009;90:2965–2972. doi/10.1099/vir.0.014449-0.