Polymerization force-regulated actin filament-Arp2/3 complex interaction dominates self-adaptive cell migrations

Xindong Chen[1,2*], Yuhui Li[3], Ming Guo[4], Bowen Xu[1], Yanhui Ma[2], Hanxing Zhu[2*], Xi-Qiao Feng[1*]

[1] Institute of Biomechanics and Medical Engineering, AML, Department of Engineering Mechanics, Tsinghua University, Beijing 100084, China

[2] School of Engineering, Cardiff University, Cardiff CF24 3AA, UK

[3] Univ. Grenoble-Alpes, CEA, CNRS, INRA, Interdisciplinary Research Institute of Grenoble, Laboratoire de Phyiologie Cellulaire & Végétale, CytoMorpho Lab, 38054 Grenoble, France.

[4] Department of Mechanical Engineering, Massachusetts Institute of Technology, Cambridge, MA 02139, USA

[*] Corresponding authors: chenxin7@tsinghua.edu.cn; zhuh3@cardiff.ac.uk; fengxq@tsinghua.edu.cn;

Abstract

Cells migrate by adapting their leading-edge behaviours to heterogeneous extracellular microenvironments (ECMs) during cancer invasions and immune responses. Yet it remains poorly understood how such complicated dynamic behaviours emerge from millisecond-scale assembling activities of protein molecules, which are hard to probe experimentally. To address this gap, we established a spatiotemporal "resistance-adaptive propulsion" theory based on the protein interactions between Arp2/3 complexes and polymerizing actin filaments, and a multiscale dynamic modelling system spanning from molecular proteins to the cell. Combining spatiotemporal simulations with experiments, we quantitatively find that cells can accurately self-adapt propulsive forces to overcome heterogeneous ECMs via a resistance-triggered positive feedback mechanism, dominated by polymerization-induced actin filament bending and the bending-regulated actin-Arp2/3 binding. However, for high resistance regions, resistance triggered a negative feedback, hindering branched filament assembly, which adapts cellular morphologies to circumnavigate the obstacles. Strikingly, the synergy of the two opposite feedbacks not only empowers cells with both powerful and flexible migratory capabilities to deal with complex ECMs, but also endows cells to use their intracellular proteins efficiently. In addition, we identify that the nature of cell migration velocity depending on ECM history stems from the inherent temporal hysteresis of cytoskeleton remodelling. We also quantitatively show that directional cell migration is dictated by the competition between the local stiffness of ECMs and the local polymerizing rate of actin network caused by chemotactic cues. Our results reveal that it is the polymerization force-regulated actin filament-Arp2/3 complex binding interaction that dominates self-adaptive cell migrations in complex ECMs, and we provide a predictive theory and a spatiotemporal multiscale modelling system at the protein level.

Introduction

Cells migrate through coupling their propulsive forces generated by assembling cytoskeletons to extracellular microenvironments (ECMs) (1-4). Actin-based lamellipodia protrusion is a powerful force-generating system that drives cell migrations during the cancer invasion, immune surveillance, and embryonic development (1, 5-9). Arp2/3 complex binds on an existing actin filament and then nucleates a daughter filament, assembling into branched actin networks in lamellipodia and invadopodia (10, 11). The polymerization of the network generates a pushing force to open a sufficient wide channel in ECM to drive individual or collective cell migrations (12-14).

Clinical studies show that Arp2/3 complex-medicated migration is tightly associated with cancer invasion (15-20), and patients over expressing Arp2/3 complex have poor survivals in lung (16), breast(17), pancreatic(20), and colorectal cancers(15). In addition, the migrations of immune cells, such as dendritic cells and T cells, in three-dimensional ECMs are also extensively driven by the Arp2/3 complex formed lamellipodial protrusions (6, 7). However, ECMs *in vivo* are highly mechanically heterogeneous(21, 22). Both invasive cancer cells and immune cells need to migrate long distances to establish new tumors(21, 23) and to find killing targets(7, 24), respectively. Experimental studies show that the magnitude of ECM resistance can affect the density of lamellipodial branched actin filaments(1, 5, 11, 25), and the lamellipodial leading-edge velocity exhibits resistance-history dependent(1, 26). In addition, cells predominantly migrate along the path of least resistance in heterogeneous extracellular microenvironments (HTECMs)(6). All these studies suggested that the leading edge of migratory cells actively mechano-senses variations of ECMs and adapts its migrating behaviour (Fig. 1a)(1, 5, 6, 27) in response to complex ECMs. However, the migratory leading edge involves highly dynamic interplays of various proteins, including Arp2/3 complexes, actin monomers, actin filaments, Wiskott-Aldrich syndrome proteins (WASPs), ATP, capping

proteins, leading-edge membrane, integrin-based adhesions and ECMs(5, 10, 11, 28). Due to

both the temporal and spatial cross-scale complexities(29) (Fig. 1a), live imaging protein

behaviours at millisecond and nanometre scales is intrinsically difficult(30). Thus,

quantitatively interpreting how these complicated cell-scale self-adaptive migration behaviours

(Fig. 1a) emerge from the dynamic activities of molecular proteins has been a grand and long-

lasting challenge (31, 32). It not only seriously hampers our in-depth mechanistic

understandings of *in vivo* cell migrations, but also hinders us from discovering target proteins

for designing new medicines and gene editing therapies to prevent cancer cells from invasions

or enhance immune cell infiltrations for solid cancer immunotherapy.

Constructing a predictive spatiotemporal multiscale modelling system that describes cellular

dynamic behaviours from the molecular to cellular scales at the intersection of biology, physics,

chemistry and computer science will greatly accelerate biomedical advancements(33, 34).

From the level of molecular protein-protein interactions, we here derive a spatiotemporal

'resistance-adaptive propulsion' (RAP) theory based on the geometric nonlinear deformation

mechanisms of polymerizing actin filaments and the mechano-chemical assembling behaviours

of Arp2/3 complexes. The theory describes both spatial and temporal mechanical interactions

between the polymerizing branched actin filaments and the bent leading-edge membrane

constrained by ECM. On the basis of this RAP theory, we develop a multiscale spatiotemporal

modelling system, which encompasses dynamic actin polymerization, capping protein

inhabiting actin filament polymerization, nonlinear deformation of actin filaments, actin

monomer diffusion, ATP binding on actin monomers and Arp2/3 complex, mechano-chemical

assembly of Arp2/3 complex, detachments of molecular linkers, bent leading-edge membrane

and HTECMs. It can not only simulate dynamic cell migrations in complex environments but

also quantitatively shed light on synchronous interacting and assembling behaviours of

multiple proteins. Combining spatiotemporal simulations with experiments tracking actin

dynamics during cell migration, we quantitatively discovered a resistance-triggered positive

feedback mechanism at the protein level for adapting propulsive forces to overcome the

heterogeneity in ECM, and a resistance-triggered negative feedback mechanism for adapting

cell morphology to circumnavigate regions with high ECM resistance. Strikingly, the

combination of the two opposite feedback mechanisms along the broad lamellipodial leading

edge shows formidable synergistic effects, empowering cells with both powerful and flexible

migratory capabilities to deal with complex ECMs, and meanwhile endowing intracellular ATP

resource with an optimal efficiency in fuelling cell migration. These insights explain why it is

hard to prevent metastasis by cancer cells once they have acquired invasive ability. By

monitoring the assembling behaviours of proteins, we further find that the nature of migration

velocity depending on the resistance history of ECM is derived from the temporal hysteresis

of adaptive actin cytoskeleton remodelling. In addition, we reveal that directional cell

migration is dictated by the competition between the local stiffness of ECMs and the local

polymerizing rates of the leading-edge actin cytoskeleton in response to gradients of

chemotactic cues. Overall, we establish the first, to our knowledge, spatiotemporal biophysical

theory and multiscale modelling system, which can accurately predict self-adaptive cell

migrations in complex ECMs from protein behaviours that happen at the milliseconds *in vivo*.

Establishing such predictive biophysical theory and spatiotemporal modelling system will

allow computer simulations to replace some laboratory experiments to quantitatively test how

new drugs and gene editing technologies targeting on proteins affect cell dynamics in the future.

Spatiotemporal self-adaptive propulsion theory and multiscale modelling system

115   Branched actin network protrusion is an important way that drives cell migrations in ECMs.

116   Through adding actin monomers to the barbed ends, polymerizing branched actin filaments

117   grow and thus generate pushing force on the leading-edge membrane (5). Before being capped

118   by capping proteins, the growing length of filaments with the polymerizing time t can be

119   expressed as

120
$$l(t) = \delta \int_{t^{nuc}}^{t^{cap}} \left[\gamma(\Phi) \cdot \psi(D_a) \cdot \mathbf{C}_a \cdot k_P(\mathbf{p}, \theta) \cdot k_{on} - k_{off} \right] dt \qquad (1)$$

where δ is the radius of an actin monomer; t^{nuc} and t^{cap} are the nucleation time and the

capping time, respectively; $\mathbf{C}_a$ is the local concentration of actin monomers in the cell; $\gamma(\Phi)$

is the consuming factor of actin monomers, introducing the relation that polymerizing rate is

proportional to the ratio of the concentration of actin monomers to the density of polymerizing

filaments Φ (35); ψ is a scaled diffusion coefficient D_a of actin monomers and is to introduce

the effect of actin filament density on the actin diffusion flux towards the polymerizing barbed

ends based on the Fick's first law of diffusion; $k_P(\mathbf{p}, \theta)$ is an interacting force-induced

probability density distribution in the rate of filament polymerization due to size-dependent

insertion of monomers $k_P(\mathbf{p}, \theta_b) = \int_{\Delta}^{\infty} \lambda \exp(-\lambda \vartheta) d\vartheta \bigg/ \int_0^{\infty} \lambda \exp(-\lambda \vartheta) d\vartheta$ (13, 36); $\mathbf{p}$ is the

interaction force that the polymerizing actin filament acts on the leading-edge membrane; θ_b

is the angle of the filament barbed end relative to the normal vector of the local membrane; ϑ

is a function of $\mathbf{p}$ and θ_b, and Δ is the size of sufficient gap to permit intercalation of

monomers(13, 36). k_{on} and k_{off} are the polymerization and depolymerization rate constants,

respectively.

The leading-edge membrane under the polymerizing force of filaments is in a bending state.

We simplify it into several continuous inclined planes based on the theory of differential

geometry (Supplementary Fig. 2b). Though the mechanical interactions between all

polymerizing actin filaments and the leading-edge membrane are in three-dimensional space,

the interaction between a single polymerizing actin filament and the local membrane can be

described in a two-dimensional deformation plane (Fig. 1b and Supplementary Fig. 2c). Then,

based on the geometric nonlinear deformation theory of continuum mechanics, the spatial and

temporal mechanical interactions between the growing (polymerizing) actin filament and the

leading-edge membrane is derived (Fig. 1b, Supplementary Methods):

144
$$\beta(s,t) = \frac{\mathbf{p}}{EI} \int_{s(t)}^{l(t)} [(1-\varepsilon(s,t)) \int_0^{s(t)} (1-\varepsilon(r,t)) \sin\beta(r,t)\,\mathrm{d}\,r]\,\mathrm{d}\,s + \frac{2\kappa(1+\nu)\mathbf{p}\sin\beta(s,t)}{EA} + \theta \quad (2)$$

145   where $\beta(s,t)$ is the deformed angle along the actin filament due to the combined effects of

146   bending, axial compression and transverse shear under the polymerizing growth; E, ν, A, I,

147   κ are the Young's modulus, Poisson's ratio, cross-sectional area, the second moment of the

148   cross-sectional area and shape factor of actin filaments, respectively. Using the deformation

149   compatibility condition,

150
$$\int_0^{l(t)} [1-\varepsilon(s,t)]\cos\beta(s,t)\,\mathrm{d}\,s = h(t), \tag{3}$$

the nonlinear deformation function in Eq. (2) can be solved through iteration. h is the distance

from the pointed end of the polymerizing actin filament to the local leading-edge membrane.

Then, the interacting force $\mathbf{p}(t)$, the propulsive force $\mathbf{f}_p(\theta,t)$ in cell migration direction, the

total deformation energy $U(t)$ and the mean bending curvature $\xi(t) = \dfrac{\beta(0) - \beta(l(t))}{l(t)}$ of the

actin filament can all be solved (Supplementary Fig. 4a-f). The total deformation energy $U(t)$

is expressed as

157
$$U(t) = \frac{\mathbf{p}^2}{2EI} \int_0^{l(t)} \left[\int_0^{s(t)} (1 - \frac{\mathbf{p}\cos\beta}{EA})\sin\beta ds \right]^2 \mathrm{d}\,s + \frac{1}{2} \int_0^{l(t)} \frac{(\mathbf{p}\cos\beta)^2}{EA}(1 - \frac{\mathbf{p}\cos\beta}{EA})\,\mathrm{d}\,s$$
$$+ \int_0^{l(t)} \frac{\kappa(1+\nu)(\mathbf{p}\sin\beta)^2}{EA}(1 - \frac{\mathbf{p}\cos\beta}{EA})\,\mathrm{d}\,s \tag{4}$$

Our mechanical analysis also shows that convex side surface of the bending actin filament

is stretched while the concave side surface is compressed (Fig. 1c). Their relative strain

$\varepsilon_r = 2r_0\xi$ where r_0 is the radius of actin filament. Experiments(37) show that *in vitro* long actin

filaments (~10μm) exhibit large bending deformations under thermal fluctuations, and Arp2/3

complexes prefer to bind onto the convex sides of bending actin filaments. We fit the

experimental data of relative branched density P with an inverted sigmoid function

$P = 2/(1+e^{0.73\xi})$ (Fig. 1d) and analyse it. Strikingly, we have , for the first time, obtained the

relative   curvature-dependent   Arp2/3-actin   filament   dissociation   constant

$K_d(\mathbf{p})/K_d^0 = 0.5(1+e^{0.73\xi})$ where $K_d(\mathbf{p})$ and K_d^0 are the dissociation constants in the bending

and straight states, respectively (Fig. 1e and Supplementary information). The relative

$K_d(\mathbf{p})/K_d^0$ shows that the affinity of Arp2/3 complex binding on the convex surface (negative

curvature side) of actin filament is higher than the straight surface, which is higher than the

concave surface (positive curvature side). This could be explained by the combination of our

mechanical analysis and the recent cryo-electron structure of Arp2/3 complex-actin filament

junction(38-40) (Fig. 1c and Supplementary Fig. 1). There are five actin subunits in mother

actin filament that contact with Arp2/3 complex, with many of the contact surfaces in the

grooves between these subunits (supplementary Fig. 1a)(38-40). ArpC1of Arp2/3 even has a

protrusion helix that inserts into actin subdomains for binding (supplementary Fig. 1b)(38, 40).

During the polymerization of actin filaments, they generate bending deformations under the

constraints of cell membrane and ECM, leading to compressions on their concave sides (Fig.

1b and c). As a result, some of the binding surfaces in the grooves on the concave side is buried

(Fig. 1c), inducing the decrease of the binding affinity of Arp2/3. On the contrary, the groove

sites on the convex side of the bending actin filaments undergoes stretch (Fig. 1c), which

facilitates Arp2/3 complex binding and improves Arp2/3 binding affinity. To better show this

force-dependent binding affinity, we also calculate the relative dissociation constant

$K_d^1(\mathbf{p})/K_d^2(\mathbf{p})$ where $K_d^1(\mathbf{p})$ and $K_d^2(\mathbf{p})$ are the dissociation constants on the convex and

concave surfaces, respectively, demonstrating that the binding affinity on the convex surface

is much higher than the concave surface (Fig. 1f). Our theoretical analysis has shown that

although *in vivo* actin filaments are relatively short (~250nm), their polymerization force still

can induce them to generate significant backward bending deformations, which could be

verified by the actin retrograde flow phenomenon (41) and the recent measurement that the

bending curvature can reach up to 10 μm^{-1} (42). This indicates that the force-dependent Arp2/3

complex- actin filament binding affinity occurs *in vivo*. To incorporate that Arp2/3 complex

has a higher binding affinity with the convex surface of actin filaments than with straight

surface, we introduce a bending curvature-dependent binding factor $d^{arp}(\xi^{max})$, which is

defined as the space between two adjacent Arp2/3 complex branches along an actin filament,

where ξ^{max} is the biggest bending curvature in the deformation history of the actin filament (Eq.

S14). The number of Arp2/3 complexes binding on the *i*th actin filament with a polymerization

length $l_i(t)$ of can be determined as $n_i^{arp} = l_i / d^{arp}(\xi_i^{max})$. The probabilities that an Arp2/3

complex binding on the convex and concave side of an actin filament are

$P(|\xi|)/[P(|\xi|)+P(-|\xi|)]$ and $P(-|\xi|)/[P(|\xi|)+P(-|\xi|)]$, respectively. Thus, the total number of

actin filaments $N(t)$ pushing against the leading-edge membrane at time t is

$$N(t) = \sum_{i=1}^{m(t)} n_i^{arp}(\xi_i^{max}) + m(t), \tag{5}$$

where $m(t)$ is the number of mother filaments. During cell migrations, some filaments are

linked to the leading-edge membrane through molecular linkers, such as Ezrin and N-

WASPs(12, 43-45), and generate a resultant attachment force $\sum_{k=1}^{\Omega(t)} \mathbf{f}_{a,k}(t)$ to pull back the

membrane where $\Omega(t)$ and $\mathbf{f}_a$ are the total number of attaching molecular linkers and

attaching force of each molecular linker, respectively. The total number of tethered filaments

$\Omega(t) = \alpha N(t)$, where the parameter α is the percentage of the total number of actin filaments

contacting with the leading-edge membrane. In the static state, the leading edge of migrating

cells follows force balance (Fig. 1g):

$$\sum_{j=1}^{N(t)} \mathbf{f}_{p,j}(\theta_j, t) = \sum_{k=1}^{\Omega(t)} \mathbf{f}_{a,k}(t) + \mathbf{f}_m(t) + \mathbf{f}_{ECM}(t) , \tag{6}$$

where $\mathbf{f}_m(t)$ and $\mathbf{f}_{ECM}(t)$ are a backward tension force from the leading-edge membrane and a

resistance force from ECM, respectively. The total elastic deformation energy stored in all

branched filaments pushing the leading-edge membrane is $\prod(t) = \sum_{j=1}^{N(t)} U_j(\theta_j, t)$. Experiments

showed that the motion of branched actin filaments is saltatory with a step size $\Delta\mathbf{S}$ of about 1-

10nm with time (46, 47) due to the detachment of molecular linkers(13, 43, 48). Actually, in

vivo, cell migration involves not only the detachments of molecular linkers from leading-edge

membrane(43), but also the local ruptures of some nascent integrin adhesions (49) and

extracellular crosslinking matrix networks owing to the propulsive force (21, 50). Since these

processes are very complex and also involves different energy barriers, in order to capture the

key characteristics of cell migration, we assume that when the resultant propulsive force

$\sum_{j=1}^{N(t)} \mathbf{f}_{p,j}(\theta_j, t)$ is larger than the maximum resultant stall force $\sum_{k=1}^{\Omega(t)} \mathbf{f}_{a,k}(t) + \mathbf{f}_m(t) + \mathbf{f}_{ECM}(t)$, some

molecular linkers will detach from the leading-edge membrane (43) and thus cell will migrate

forward with a leaping step size $\Delta\mathbf{S}$. Then, the position of the leading edge of the cell at time

$t+1$ can be expressed by

$$\mathbf{S}(t+1) = \begin{cases} \mathbf{S}(t) & \text{if } \sum_{j=1}^{N(t+1)} \mathbf{f}_{p,j}(\theta_j, t+1) \leq \sum_{k=1}^{\Omega(t+1)} \mathbf{f}_{a,k}(t+1) + \mathbf{f}_m(t+1) + \mathbf{f}_{ECM}(t+1) \\ \mathbf{S}(t) + \Delta\mathbf{S} & \text{if } \sum_{j=1}^{N(t+1)} \mathbf{f}_{p,j}(\theta_j, t+1) > \sum_{k=1}^{\Omega(t+1)} \mathbf{f}_{a,k}(t+1) + \mathbf{f}_m(t+1) + \mathbf{f}_{ECM}(t+1) \end{cases} \tag{7}$$

and the transient average migration velocity $\mathbf{V}$ is $\Delta\mathbf{S}/\Delta t$. The dynamic leading-edge

migration should meet the force balance defined by

$$\sum_{j=1}^{N(t+1,\ \mathbf{S}+\Delta\mathbf{S})} \mathbf{f}_{p,j}(\theta_j, t+1) = \sum_{k=1}^{\Omega(t+1,\ \mathbf{S}+\Delta\mathbf{S})} \mathbf{f}_{a,k}(t+1) + \mathbf{f}_m(t+1) + \mathbf{f}_{ECM}(t+1) \qquad (8)$$

Because the present theory is derived from the spatiotemporal mechanical interactions between

the polymerizing branched actin filaments and the leading-edge membrane constrained by

extracellular resistance, and incorporates the force-dependent assembling behaviours of

proteins, we name it spatiotemporal 'resistance-adaptive propulsion' (RAP) model.

To quantitatively shed light on how the assembling behaviours of multiple proteins impact

on the dynamics of cell migration, we further develop a multiscale spatiotemporal modelling

system by applying the RAP theory and integrating the stochastic behaviours of proteins

(Supplementary Methods). This modelling framework systematically encompasses the *in vivo*

actin monomer nucleation, actin filament polymerization, capping protein terminating filament

polymerization, mechano-chemical nucleation of Arp2/3 complex, ATP binding, bent leading-

edge membrane, detachments of molecular linkers, protein gradients caused by chemotactic

cues, integrin-based adhesion and heterogeneous ECMs. Using this bottom-up approach, we

can span multiple scales in both space and time to investigate cell migration behaviours in

complex ECMs, and thus shed light on their dominating biophysical principles from the level

of specific proteins.

ECM resistance-triggered positive feedback adapts propulsive force

High extracellular resistance results in denser lamellipodial branched actin filaments during

cell migrations(1, 5). To explore quantitatively whether our spatiotemporal RAP theory can

reproduce this significant behaviour and reveal its underlying biophysical mechanism, we

perform spatiotemporal simulations of cell migrations in both mechanically homogeneous

extracellular microenvironment (HMECM) and HTECM (Fig. 2a). we observe that when the

resistance $\mathbf{f}_{ECM}$ is 0.5 nN/μm (the normal range is 0.1–2.0 nN/μm of ECM(51)) in the HMECM

condition, the density of polymerizing branched actin filaments Φ stably fluctuates in a very

narrow range 230–270 /μm (Fig. 2a), agreeing well with the experimental data 150–350

/μm(51). However, in the HTECM, when the extracellular resistance $\mathbf{f}_{ECM}$ increases from the

low resistance (LR) 0.5 nN/μm to a higher resistance (HR) 1.0 nN/μm, the branched actin

density Φ increases by ~35% (320–350 /μm, Fig. 2a). After $\mathbf{f}_{ECM}$ returning to the low

resistance (RLR) 0.5 nN/μm, the density also decreases to its previous level (Fig. 2a). The

density of Arp2/3 complex assembling actin cytoskeleton cofluctuates with the resistance (Fig.

2b). Strikingly, our model well predicts the experimentally measured correlations between the

extracellular resistance and the lamellipodial branched actin filament density during cell

migration in refs.(1, 5, 11) (Fig. 2a). The architecture of the lamellipodial branched actin

network generated in our spatiotemporal simulations is also consistent with the experimental

measurements (Fig. 2c) (52). In addition, we show that when cells encounter higher resistances,

the consumption rate of ATP (Fig. 2d) increases to fuel cell migrations. This prediction is also

validated by the experimental data (Fig. 2d) that mitochondria and ATP levels are higher at the

invasive cell leading edge in a stiffer ECM confinement(11) and that cancer cells overproduce

ATP to boost their lamellipodia formations and invasions(53).

Next, we analyse the spatiotemporal propulsive force (Fig. 2e) and the elastic deformation

energy Π stored in the branched filaments (Fig. 2f). It is found that both synchronously

fluctuate with the density of branched actin filaments. We further quantitatively identify that

the adaptation of filament density is to meet the propulsive force and energy demands for

overcoming the varying extracellular resistance. To shed light on the more fundamental cross-

scale biophysical mechanism of such significant adaptive behaviours in response to ECMs, we

then examine the assembly of protein molecules that happens at milliseconds, finding an ECM

resistance-triggered positive feedback (Fig. 2g). When the resistance $\mathbf{f}_{ECM}$ increases,

polymerizing (growing) actin filaments under the ECM confinement will automatically have

larger nonlinear bending deformations (Supplementary Movie), which increase the probability

that Arp2/3 complex will bind and nucleate daughter actin filaments, and vice versa. Through

this mechanism, cells can adapt the density of branched actin network and thus the propulsive

force and energy for migrations. More importantly, through the extent of bending deformations,

cells can sense the ECM resistance, so a larger resistance induces a larger bending deformation.

Thus, from the protein level, we reveal that migrating cells can sensitively sense the immediate

variations of ECM resistance through the polymerizing growth of actin filaments, and then

accordingly make accurate adaptive responses in filament density, propulsive force and energy

through the mechano-triggered Arp2/3 complex-actin filament assembling behaviours (Fig. 2g).

This mechanism also endows migrating cells with an optimization ability to accurately employ

their intracellular proteins and ATP (Fig. 2a, b and d) according to their demands in complex

ECMs.

Leading-edge velocity depends on the temporal hysteresis of filament density

adaptation to varying ECM

We next explore how the leading-edge migration velocity responds to the varying stiffness

of the ECMs and how the underlying biophysical mechanisms operate at the level of protein

molecules. The spatiotemporal simulations in HTECMs show that when migrating cell

encounters an increased extracellular resistance $\mathbf{f}_{ECM}$, its leading-edge migrating velocity $\mathbf{V}$

suddenly decreases from 3.3 μm/min to 0.7 μm/min (from stage I to II in the black ellipse in

Fig. 2h), which is lower than the velocity 1.7–5.2 μm/min in the HMECMs with $\mathbf{f}_{ECM} = 0.5$

nN/μm. However, with the continuing polymerization of branched filaments, the decreased

velocity partially recovers. This is owing to the gradual increase in the filament density based

on the ECM resistance-triggered positive feedback. Afterwards, we reduce the extracellular

resistance from 1 nN/μm to its previous value 0.5 nN/μm for the HTECMs. Strikingly, the

leading-edge velocity abruptly increases to a very high value 10.7 μm/min, and then gradually

decreases to the previous range 1.7-5.2 μm/min (Fig. 2h). The spatiotemporal predictions from

stage I to V are validated by the experimental data (Fig. 2h) (1, 26). To gain insight into the

protein behaviours, we check the evolution of the spatiotemporal remodelling of the growing

lamellipodial actin network. Previous studies(26) described this phenomenon as velocity

dependence on loading history. Here, we quantitatively show that the nature of the leading-

edge velocity variations in the HTEMCs stems from that the adaptation of branched actin

filament density is always temporally hysteretic to its triggering reason, i.e., varying ECM

resistance (Fig. 2i). This is because the generation of daughter filaments and their growths to

the leading-edge membrane always cost some time due to actin monomer nucleation, filament

polymerization, Arp2/3 activation and assembly (Fig. 2i). Although the extracellular resistance

$\mathbf{f}_{ECM}$ has increased or decreased, the density, propulsive force and deformation energy of

polymerizing actin filaments keep unchanged in this process. Thus, the velocity suddenly

decreases because of incapable of overcoming the increased resistance, and increases because

of easily overcoming the decreased resistance to release the excess deformation energy,

respectively (Fig. 2e and f). This also indicates that, in HTECMs, there exists no one-to-one

correspondence between $\mathbf{V}\text{-}\mathbf{f}_{ECM}$ that can describe the leading-edge migrating behaviours.

However, in HMECMs, the one-to-one $\mathbf{V}\text{-}\mathbf{f}_{ECM}$ relationship exists.

High ECM resistance-triggered negative feedback adapts cell morphology for pathfinding

The above study focuses on the low ranges of ECM resistance. However, the leading edge usually faces some local dense collagen regions with high resistances in ECM. Thus, we here investigate migratory pathfinding in this kind of complex ECMs and shed light on its underlying mechanistic operating basis at the protein level. We design an ECM as demonstrated by Fig. 3a. The ECM is divided into two stages. In the first stage (from 0 to 110 nm), it is mechanically homogeneous and has a resistance $\mathbf{f}_{ECM}$ of 0.5 nN/μm. However, in the second stage (from 110 to 300 nm), its resistance becomes 1.0 nN/μm, and there are two very dense collagen regions with a very high resistance $\mathbf{f}_{ECM}^{right} = \mathbf{f}_{ECM}^{left} = 5$ nN/μm.

Spatiotemporal simulations show that, in the first homogeneous mechanical environment, the left, central and right parts of the leading edge migrate forward synchronously with similar velocities (Fig. 3b). The density of branched actin filaments is approximately homogeneous (Fig. 3c). However, when the cell encounters the two dense collagen ECM regions, it stops moving forward from the left and right sides, and turns to squeeze out from the central region where the resistance is weak (Fig. 3b). Unexpectedly, even though the leading edges on the left and right sides have not overcome their local resistances, the increase of the density of actin filaments stops (red ellipse area in Fig. 3c). Interestingly, this indicates that when the ECM resistance is very high, the mechanism of resistance-triggered positive feedback no longer work. We perform experiments to validate this spatiotemporal model prediction. We record Human retinal pigment epithelial-1 (RPE1) cells expressing green fluorescent protein linked to a small peptide (Lifeact-GFP) with affinity for actin microfilaments. We image the dynamic cell migration processes in the microchannels with constrictions (Fig. 3d), and track the temporal actin intensity when the cell encounters the constriction regions. The experimental

results of the temporal variations of leading-edge actin intensity are strongly consistent with

our modelling predictions (Fig. 3e). We then check how our spatiotemporal simulations operate

at the protein level (Fig. 3f and g). Migrating cells apply the resistance-triggered positive

feedback to adapt filament density and propulsive force to try to overcome the obstacles until

high forces between the leading-edge membrane and actin filaments (Fig. 3h) compromise the

intercalations of actin monomers with the barbed ends (Fig. 3f), which stops the bending

deformation of filaments (Fig. 3i) and the assembly of Arp2/3 complex (Fig. 3g). These results

of our simulation are strongly validated by *in vitro* experiments (25). We also find that the

porosity of the leading-edge actin network is reduced due to the increased density of actin

filaments (Fig. 3j). This lowers the diffusion of actin monomers to the free barbed ends, and

reduces the polymerization rate at the leading edge (Fig. 3k).

Unexpectedly, we find that an ECM resistance-triggered negative adaptation feedback (Fig.

3l) coexists with the resistance-triggered positive adaptation feedback (Fig. 2g) in migrating

cells. While the positive feedback adapts cell propulsive force to overcome ECMs, the negative

feedback can adapt cell morphology by stopping actin polymerization to circumnavigate the

high resistance regions in ECMs. This behaviour can avoid unnecessary consumptions of

intracellular proteins and ATP resources (Fig. 3f and g), and thus improve cell migration

efficiency. The synergy of the two opposite feedbacks allow the leading edge to discriminate

whether the local ECM confinement is weak to overcome or strong enough to require

circumnavigation with a formidable efficiency. It endows cells with both powerful and flexible

migration capabilities to widely adapt morphologies to the complex ECMs (Fig. 3m). This may

explain why cancer metastasis is extremely hard to prevent once they acquire invasive ability.

In addition, our results reveal that the initiation of cellular morphology adaptations, a prominent

characteristic of invasive cancer cells, derives from the leading edge sensing strong barriers

and escaping from them.

Directional cell migration is steered by a balanced and competing relation

Cells typically follow the gradients of chemotactic cues to migrate(6). The nucleus of migrating cells acts as a mechanical gauge to make a temporary adaptations while choosing a passable path (6, 54, 55). Given the global migration direction predefined by chemotactic cues, leading edge is much more important in persistently directing cell migration towards the final destination. Our simulations demonstrate that different local stiffness of ECM results in different local densities of branched filaments along the leading edge. Here, we investigate another challenging problem of how the complex interplays between the multiple simultaneous factors of extracellular resistance, density heterogeneity of branched actin filaments, and the external diffusible chemotactic stimuli synergistically steer directional cell migration in HTECM. We introduce a gradient diffusion of a localised chemotactic cue sensed by transmembrane receptors (Fig. 4a), rendering a gradient distribution of intracellular actin monomers. We design and simulate three cases A–C (Fig. 4b), in which local leading edges at the positions of $\mathbf{x}_1$ and $\mathbf{x}_2$ simultaneously drive cell migrations. Based on the experimental observations (6), it is reasonable to hypothesize that the fastest migration direction of local leading edges is the main migration direction of a cell. Our simulations in case A show that the migration distance of the leading edge at the position $\mathbf{x}_2$ are larger than those at the position $\mathbf{x}_1$ (Fig. 4c), concluding that when the extracellular mechanical microenvironments are homogeneous, cells more actively migrate toward the site where the local concentration of actin monomers is higher. Since homogeneous ECM induces homogeneous density of branch filaments in the whole leading edge (Fig. 4d), a higher local concentration of actin monomers means that sufficient actin monomers can be supplied to the polymerizing barbed ends of the local branched actin filaments. The chemotactic cue determines the local protein concentration to steer directional cell migration. Then, we test case B, in which the left and right sides are designed to have the same local concentrations of actin monomers due to homogeneous

distribution of the chemotactic cue, but different extracellular resistances. Cells migrate toward

the low resistance $\mathbf{x}_1$ side (Fig. 4e), because higher resistance on the right-side results in denser

branched actin filaments there (Fig. 4f), which deplete action monomers locally and thus slows

polymerization rate at barbed ends. This case highlights that extracellular resistance plays a

determining role in directing cell migration. However, the results of case C (Supplementary

Fig. 5), in contrary to case B, shows that cells migrate toward the high resistance $\mathbf{x}_2$ side (Fig.

4g). Even though the denser polymerizing filaments consume more action monomers on the $\mathbf{x}_2$

side (Fig. 4h), the strong chemotactic cue can sustain a high local concentration of actin

monomers, which enable branched actin filaments to polymerize at a higher rate than that on

the $\mathbf{x}_1$ side. Although here we take actin monomers as an example, local variations in the

concentrations of other intracellular proteins, such as Arp2/3 complex, WASPs, FMNL, aprin

and profilin, have the same effect on cell migration by enhancing or inhibiting nucleation,

branch formation and filament polymerization.

From the distinct results of cases A–C, we find that directional cell migration is not solely

steered by either the gradients of intracellular proteins induced by chemotactic cues or the local

stiffness of ECMs. It is a balanced competing consequence between the two (Fig. 4i and j).

While strong chemotactic cues improve the local nucleation, branch formation and

polymerization rate of actin filaments, stiffer local ECMs result in denser local branched actin

filaments, which in turn reduce these rates and slow down cell migration. In addition, through

the collaboration of the resistance-triggered positive feedback mechanism and the high

resistance-triggered negative feedback mechanism, here our results further indicate that as long

as the chemotactic cues are strong enough in sustaining the high gradients of proteins for

keeping a high branching and polymerizing rates, the leading edge will drive cells to migrate

globally and persistently toward the prescribed final destination by overcoming weak

confinements and circumnavigating strong barriers encountered.

Discussions

Although cell migrations have been studied many decades, how cells mechano-sense and make self-adaptive responses to complex ECMs at the protein level still remains elusive. In this study, by analysing the spatiotemporal nonlinear deformation of polymerizing actin filaments and the polymerization force-regulated Arp2/3 complex-actin filament binding interactions, we derive a spatiotemporal 'resistance-adaptive propulsion' (RAP) theory for cell migrations. Then, with the RAP theory, we develop a spatiotemporal multiscale modelling system, which can not only simulate dynamic cell migrations in ECMs, but also shed light on the assembling behaviours of single proteins. Our simulations predict many important spatial and temporal adaptive cell migration behaviours observed experimentally and clinically (Table 1), and reveal their underlying operating mechanisms emerged from proteins behaviours.

We find that it is the polymerization force-regulated actin filament-Arp2/3 complex binding interaction that dominates self-adaptive cell migrations in complex ECMs, through the synergistic effects of positive and negative feedbacks. The polymerization-induced bending state of actin filaments serves as the mechano-sensor of cells. Our finding is highly consistent with two recent studies that show bending force (42) and polymerization force (56) evoke nucleotide-state conformational changes of actin filaments and thus affect the binding behaviours of actin-binding proteins. In fact, mechanical force-induced conformational change of proteins is a key regulator of protein-protein interactions (57), thereby regulating physiological and pathophysiological cellular behaviours.

Previous studies indicated that increased extracellular resistance induced denser branched actin filaments in the migration leading edge(1, 5, 25). Here, we identify that the enhanced leading-edge actin filament density is to improve the propulsive force and energy to overcome the resistance. Our results physically interpret why Arp2/3 complex overexpression is tightly

associated with cancer cell invasions(15-19), poor patient survival in cancers(58) and high

migratory force(59). We indicate that Arp2/3 is a key target protein for developing anti-cancer

drugs, especially for patients in the advanced stages of cancer. It should be mentioned that the

current simulations are for the conditions that integrin-based adhesions are sufficient for fixing

the branched actin network. In the event that the levels of integrin or vinculin are not abundant,

it is conceivable that high ECM resistance could trigger a rapid actin retrograde flow (41, 60)

based on the positive feedback, resulting in an increased backward movement of the branched

actin network and subsequently reducing the velocity of cell protrusion. This represents an

additional type of negative feedback mechanism that can arise from excessively high ECM

resistance.

It should be emphasized that polymerization of Arp2/3 complex-branched actin filament

network is the most important way for cells to generate propulsive forces to interact with their

surrounding microenvironments and perform their functions. It participates in phagocytosis of

immune responses(61), endocytosis(62), dendritic spine formation of cortical neurons(63), T-

cell and cancer cell interactions(64), and promoting of DNA fork repair (65) and chromatin

organization(66, 67). All these processes require cells to mechano-sense the surrounding

microenvironments and then make adaptive force generation responses. Thus, besides cell

migrations, our cross-scale findings and the spatiotemporal multiscale modelling system can

also be applied to investigate these dynamic physiological cell activities.

Materials and Methods

Cell culture

Human telomerase-immortalized, retinal-pigmented epithelial cells expressing GFP-LifeAct (CellLight Actin-GFP, BacMam 2.0 from Thermo Fischer Scientific) were grown in a humidified incubator at 37 °C and 5% CO_2 in DMEM/F12 supplemented with 10% fetal bovine serum and 1% penicillin-streptomycin. All cell culture products were purchased from GIBCO/Life Technologies. Cell lines were regularly checked for mycoplasma contamination (MycoAlert, Lonza).

Cell fixation and immunostaining

Cells were pre-permeabilized in 0.5% Triton X-100 in cytoskeleton buffer for 15 s for tubulin and then fixed in 0.5% glutaraldehyde (no.00216-30; Polysciences) in cytoskeleton buffer with 0.5% Triton X-100 and 10% sucrose for 15 min at room temperature. Cells were then washed three times with PBS-tween 0.1% and incubated in a quenching agent of 1mg ml-1 sodium borohydride for 10 min at room temperature. After fixation, the cells were washed with PBS-Tween 0.1% and then blocked with 3% bovine serum albumin (BSA) overnight. The cells were incubated with appropriate dilutions of primary antibodies in PBS containing 3% BSA and 0.1% Tween overnight at 4 °C in a humid chamber. After washing three times with PBS-tween 0.1%, the coverslips were then incubated with appropriate dilutions of secondary antibodies diluted in PBS containing 3% BSA and 0.1% for 1 h at room temperature in a humid chamber. After washing three times with PBS-Tween 0.1%, coverslips were then mounted onto slides using Prolong Gold antifade reagent (no. P36935; Invitrogen). Here, we used rat monoclonal antibodies against α-tubulin (no. ab6160, Abcam) and Alexa Fluor 555 goat anti-rat (1:500, no.A21434; Invitrogen) as secondary antibody. Antibodies were diluted as followed for immunofluorescence: α-tubulin (primary: 1:500) and secondary (1:1000). For actin

483 immunofluorescence, Phalloidin-Atto 488 (49409, Sigma-Aldrich) was used to and diluted as

484 1:1000 to stabilize actin filaments in cells.

Cell migration under constriction

Micro-channels were prepared as previously described (doi:10.1007/978-1-61779-207-

6_28). Briefly, polydimethylsiloxane (PDMS, 10/1 w/w PDMS A / crosslinker B) (GE

Silicones) was used to prepare 12 μm wide and 5μm high micro-channels with a constriction

of 2 μm. For confined migration in Fig. 3d, coverslip and micro-channels were treated by

plasma for 1mins and then were stuck together at 70 ℃ for 10mins. Before cell seeding, the

microchannels were treated with fibronectin at 10 μg ml-1 for 30mins and incubate with culture

medium for 3 hrs at room temperature. GFP-LifeAct RPE1 cells were then seeded in the

microchannels with a concentration at 2x105 cells ml-1. Imaging was performed after the

overnight incubation of the microchannels.

Imaging

Images of the immunostainings were acquired on a Zeiss LSM900 confocal microscopes

(Axio Observer) using a 63x magnification objective (Plan- Apochromat 63X/1.4 oil). Image

acquisition for time-lapse of GFP life-act RPE1 cells in constrictions was performed on a

confocal spinning-disc system (EclipseTi-E Nikon inverted microscope equipped with a

CSUX1-A1 Yokogawa confocal head, an Evolve EMCCD camera from Roper Scientific,

Princeton Instruments) through an 63x magnification objective (Nikon CFI Plan Fluor 60X/0.7

oil) objective every 4 s during 2 hrs for each time-lapse. The set-up was equipped with a live

cell chamber, and the temperature was constantly kept at 37 ℃. Labelled-actin was excited

with a 491 nm laser line, and emission was observed with a standard GFP filter. The microscope

was monitored with MetaMorph software (Universal Imaging).

Spatiotemporal theoretical model and dynamic multiscale modelling system

The derivation of the spatiotemporal biophysical theory and the developing process of the spatiotemporal multiscale dynamic modelling system are provided in the Supplementary Text.

Data availability

The data that support the findings of this study are available from the corresponding author on reasonable request.

Author contributions

X.D.C., X.Q.F., H.X.Z. and M.G. designed the research. X.D.C., H.X.Z. and X.Q.F. developed the theory and the spatiotemporal simulation framework. X.D.C performed the simulations. X.D.C., B.W.X. and Y.H.M analysed the data. X.D.C. and Y.H.L. did the experiments and analysed experimental data. X.D.C., X.Q.F., H.X.Z., Y.H.L. and M.G. wrote the manuscript.

Acknowledgements

We gratefully thank Professor Thomas D. Pollard for his invaluable insights and great help for manuscript preparation, and Professor Laurent Blanchoin for help with the experiments. We also thank Professors Xin Liang, Congying Wu, Bo Li and Yue Shao for valuable discussions. X.D.C., X.Q.F, and B.W.X. acknowledge the support from the National Natural Science Foundation of China (Grant no. 11921002 and 12032014)

Competing interests

The authors declare no conflict of interest.

Code availability

All computer codes are available from the corresponding authors on reasonable request.

References

1. Mueller J, *et al.* (2017) Load adaptation of lamellipodial actin networks. *Cell* 171(1):188-200. e116.

2. Bera K, *et al.* (2022) Extracellular fluid viscosity enhances cell migration and cancer dissemination. *Nature*:1-9.

3. Chen X, *et al.* (2020) Predictive assembling model reveals the self-adaptive elastic properties of lamellipodial actin networks for cell migration. *Communications biology* 3(1):1-17.

4. Broedersz CP & MacKintosh FC (2014) Modeling semiflexible polymer networks. *Reviews of Modern Physics* 86(3):995.

5. Bieling P, *et al.* (2016) Force Feedback Controls Motor Activity and Mechanical Properties of Self-Assembling Branched Actin Networks. *Cell* 164(1-2):115-127.

6. Renkawitz J, *et al.* (2019) Nuclear positioning facilitates amoeboid migration along the path of least resistance. *Nature* 568(7753):546.

7. Liu T-l, *et al.* (2018) Observing the cell in its native state: Imaging subcellular dynamics in multicellular organisms. *Science* 360(6386):eaaq1392.

8. Chen B-C, *et al.* (2014) Lattice light-sheet microscopy: imaging molecules to embryos at high spatiotemporal resolution. *Science* 346(6208).

9. Svitkina TM & Borisy GG (1999) Arp2/3 complex and actin depolymerizing factor/cofilin in dendritic organization and treadmilling of actin filament array in lamellipodia. *The Journal of cell biology* 145(5):1009-1026.

10. Blanchoin L, *et al.* (2000) Direct observation of dendritic actin filament networks nucleated by Arp2/3 complex and WASP/Scar proteins. *Nature* 404(6781):1007.

11. Kelley LC, *et al.* (2019) Adaptive F-actin polymerization and localized ATP production drive basement membrane invasion in the absence of MMPs. *Developmental cell* 48(3):313-328. e318.

12. Mehidi A, *et al.* (2021) Forces generated by lamellipodial actin filament elongation regulate the WAVE complex during cell migration. *Nature Cell Biology*:1-15.

13.   Mogilner A & Oster G (2003) Force generation by actin polymerization II: the elastic
ratchet and tethered filaments. *Biophysical journal* 84(3):1591-1605.

14.   Lin Y (2009) Mechanics model for actin-based motility. *Physical Review E*
79(2):021916.

15.   Frentzas S, *et al.* (2016) Vessel co-option mediates resistance to anti-angiogenic
therapy in liver metastases. *Nature medicine* 22(11):1294-1302.

16.   Semba S, *et al.* (2006) Coexpression of actin-related protein 2 and Wiskott-Aldrich
syndrome family verproline-homologous protein 2 in adenocarcinoma of the lung.
Clinical cancer research 12(8):2449-2454.

17.   Kazazian K, *et al.* (2017) Plk4 promotes cancer invasion and metastasis through Arp2/3
complex regulation of the actin cytoskeleton. *Cancer research* 77(2):434-447.

18.   Molinie N & Gautreau A (2017) The Arp2/3 regulatory system and its deregulation in
cancer. *Physiological reviews* 98(1):215-238.

19.   Zhao K, *et al.* (2020) WDR63 inhibits Arp2/3‑dependent actin polymerization and
mediates the function of p53 in suppressing metastasis. *EMBO reports* 21(4):e49269.

20.   Juin A, *et al.* (2019) N-WASP Control of LPAR1 Trafficking Establishes Response to
Self-Generated LPA Gradients to Promote Pancreatic Cancer Cell Metastasis. *Dev Cell*
51(4):431-445 e437.

21.   Wisdom KM, *et al.* (2018) Matrix mechanical plasticity regulates cancer cell migration
through confining microenvironments. *Nature communications* 9(1):4144.

22.   Park D, *et al.* (2019) Extracellular matrix anisotropy is determined by TFAP2C-
dependent regulation of cell collisions. *Nature Materials*:1-12.

23.   Di Martino JS, *et al.* (2021) A tumor-derived type III collagen-rich ECM niche
regulates tumor cell dormancy. *Nature Cancer*:1-18.

24.   Rafiq S, Hackett CS, & Brentjens RJ (2020) Engineering strategies to overcome the
current roadblocks in CAR T cell therapy. *Nature reviews Clinical oncology* 17(3):147-
167.

25.   Bieling P, Weichsel J, Mullins RD, & Fletcher DA (2022) The molecular mechanism
of load adaptation by branched actin networks. *Elife* 11:e73145.

26.   Parekh SH, Chaudhuri O, Theriot JA, & Fletcher DA (2005) Loading history
determines the velocity of actin-network growth. *Nature cell biology* 7(12):1219.

27.   Boujemaa-Paterski R, *et al.* (2017) Network heterogeneity regulates steering in actin-
based motility. *Nature communications* 8(1):655.

28.   Simon C, *et al.* (2019) Actin dynamics drive cell-like membrane deformation. *Nature*
Physics:1.

29.   Qin Y, *et al.* (2021) A multi-scale map of cell structure fusing protein images and
interactions. *Nature*:1-7.

30.   Li D, *et al.* (2015) Extended-resolution structured illumination imaging of endocytic
and cytoskeletal dynamics. *Science* 349(6251).

31.   Stuelten CH, Parent CA, & Montell DJ (2018) Cell motility in cancer invasion and
metastasis: insights from simple model organisms. *Nature reviews Cancer* 18(5):296.

32.   Mogilner A (2006) On the edge: modeling protrusion. *Current opinion in cell biology*
18(1):32-39.

33.   Singla J, *et al.* (2018) Opportunities and Challenges in Building a Spatiotemporal
Multi-scale Model of the Human Pancreatic β Cell. *Cell* 173(1):11-19.

34.   Sun S-Y, *et al.* (2022) Bio-chemo-mechanical coupling models of soft biological
materials: A review. *Advances in Applied Mechanics* 55:309.

35.   Lee CW, *et al.* (2013) Dynamic localization of G-actin during membrane protrusion in
neuronal motility. *Current Biology* 23(12):1046-1056.

36.   Mogilner A & Oster G (1996) Cell motility driven by actin polymerization. *Biophysical*
journal 71(6):3030-3045.

37.   Risca VI, *et al.* (2012) Actin filament curvature biases branching direction. *Proceedings*
of the National Academy of Sciences 109(8):2913-2918.

38.   Fäßler F, Dimchev G, Hodirnau V-V, Wan W, & Schur FKM (2020) Cryo-electron
tomography structure of Arp2/3 complex in cells reveals new insights into the branch
junction. *Nature Communications* 11(1):6437.

39.   Chou SZ, Chatterjee M, & Pollard TD (2022) Mechanism of actin filament branch
formation by Arp2/3 complex revealed by a high-resolution cryo-EM structureof the
branch   junction.   *Proceedings   of   the   National   Academy   of   Sciences*
119(49):e2206722119.

40.   Ding B, *et al.* (2022) Structure of Arp2/3 complex at a branched actin filament junction
resolved by single-particle cryo-electron microscopy. *Proceedings of the National*
Academy of Sciences 119(22):e2202723119.

41.   Bangasser BL, *et al.* (2017) Shifting the optimal stiffness for cell migration. *Nature*
communications 8(1):15313.

42.   Reynolds MJ, Hachicho C, Carl AG, Gong R, & Alushin GM (2022) Bending forces
and nucleotide state jointly regulate F-actin structure. *Nature* 611(7935):380-386.

43. Welf ES*, et al.* (2020) Actin-membrane release initiates cell protrusions. *Developmental Cell*.

44. Senju Y*, et al.* (2017) Mechanistic principles underlying regulation of the actin cytoskeleton by phosphoinositides. *Proceedings of the National Academy of Sciences* 114(43):E8977-E8986.

45. Co C, Wong DT, Gierke S, Chang V, & Taunton J (2007) Mechanism of actin network attachment to moving membranes: barbed end capture by N-WASP WH2 domains. *Cell* 128(5):901-913.

46. Shaevitz JW & Fletcher DA (2007) Load fluctuations drive actin network growth. *Proceedings of the National Academy of Sciences* 104(40):15688-15692.

47. Kuo SC & McGrath JL (2000) Steps and fluctuations of Listeria monocytogenes during actin-based motility. *Nature* 407(6807):1026.

48. Alberts JB & Odell GM (2004) In silico reconstitution of Listeria propulsion exhibits nano-saltation. *PLoS biology* 2(12):e412.

49. Thoumine O, Kocian P, Kottelat A, & Meister J-J (2000) Short-term binding of fibroblasts to fibronectin: optical tweezers experiments and probabilistic analysis. *European Biophysics Journal* 29(6):398-408.

50. Adebowale K*, et al.* (2021) Enhanced substrate stress relaxation promotes filopodia-mediated cell migration. *Nature Materials*:1-10.

51. Abraham VC, Krishnamurthi V, Taylor DL, & Lanni F (1999) The actin-based nanomachine at the leading edge of migrating cells. *Biophysical journal* 77(3):1721-1732.

52. Maly IV & Borisy GG (2001) Self-organization of a propulsive actin network as an evolutionary process. *Proceedings of the National Academy of Sciences* 98(20):11324-11329.

53. Delaunay S*, et al.* (2022) Mitochondrial RNA modifications shape metabolic plasticity in metastasis. *Nature*:1-11.

54. Lomakin A*, et al.* (2020) The nucleus acts as a ruler tailoring cell responses to spatial constraints. *Science* 370(6514).

55. Venturini V*, et al.* (2020) The nucleus measures shape changes for cellular proprioception to control dynamic cell behavior. *Science* 370(6514).

56. Oosterheert W, Klink BU, Belyy A, Pospich S, & Raunser S (2022) Structural basis of actin filament assembly and aging. *Nature*:1-6.

57.   Wang Y, *et al.* (2021) Force-dependent interactions between talin and full-length
660   vinculin. *Journal of the American Chemical Society* 143(36):14726-14737.

661   58.   Broutier L, *et al.* (2017) Human primary liver cancer–derived organoid cultures for
662   disease modeling and drug screening. *Nature medicine* 23(12):1424-1435.

663   59.   Krndija D, *et al.* (2019) Active cell migration is critical for steady-state epithelial
664   turnover in the gut. *Science* 365(6454):705-710.

665   60.   Isomursu A, *et al.* (2022) Directed cell migration towards softer environments. *Nature*
666   *Materials* 21(9):1081-1090.

667   61.   Jaumouillé V, Cartagena-Rivera AX, & Waterman CM (2019) Coupling of β2 integrins
668   to actin by a mechanosensitive molecular clutch drives complement receptor-mediated
669   phagocytosis. *Nature cell biology* 21(11):1357-1369.

670   62.   Yang C, *et al.* (2022) Actin polymerization promotes invagination of flat clathrin-
671   coated lattices in mammalian cells by pushing at lattice edges. *Nature communications*
672   13(1):1-20.

673   63.   Kim IH, *et al.* (2015) Spine pruning drives antipsychotic-sensitive locomotion via
674   circuit control of striatal dopamine. *Nature neuroscience* 18(6):883.

675   64.   Basu R, *et al.* (2016) Cytotoxic T cells use mechanical force to potentiate target cell
676   killing. *Cell* 165(1):100-110.

677   65.   Lamm N, *et al.* (2020) Nuclear F-actin counteracts nuclear deformation and promotes
678   fork repair during replication stress. *Nature Cell Biology* 22(12):1460-1470.

679   66.   Schrank BR, *et al.* (2018) Nuclear ARP2/3 drives DNA break clustering for homology-
680   directed repair. *Nature* 559(7712):61-66.

681   67.   Debaugnies M, *et al.* (2023) RHOJ controls EMT-associated resistance to
682   chemotherapy. *Nature*:1-8.

688 **Figures**

a Time/Space Scales: min/µm ⟵ ms/nm

ECM triggers

Proteins determine

Adaptive cell migration behaviours

Assembling behaviours of proteins

b $\vec{\mathbf{n}}$

membrane

$\mathbf{p}$ point/line contact

polymerizing

$\beta(s)$ s mechano-triggered binding

pointed end

h

θ

c $\varepsilon_1 = r_0\xi$

ds

$\varepsilon_2 = -r_0\xi$ r_0 r_0

$\varepsilon_r = \varepsilon_1 - \varepsilon_2$

R

$d\beta$

O

d

$$P = \frac{2}{1+e^{0.73\xi}}$$

— Fitting curve
○ Experimental data (37)

Relative branch density P

Curvature ξ (µm⁻¹)

e

$0.5(1+e^{0.73\xi})$

$(0,1)$

$K_d(\mathbf{p}) / K_d^0$

Curvature ξ (µm⁻¹)

f

— $K_d^1(\mathbf{p}) / K_d^2(\mathbf{p})$

$$\frac{1+e^{-0.73|\xi|}}{1+e^{0.73|\xi|}}$$

ε_r

$K_d^1(\mathbf{p}) / K_d^2(\mathbf{p})$

Relative strain ε_r (%)

Curvature $|\xi|$ (µm⁻¹)

g Mechanically heterogeneous ECM

$\mathbf{f}_{ECM}$

$\mathbf{f}_m$ $\mathbf{f}_p$ $\mathbf{f}_a$

Arp2/3 • Actin monomer • WASP • Capping protein • Molecular linker

ATP • Activated Arp2/3 • ATP-actin monomer • ECM

Fig. 1. Polymerizing branched actin filaments at lamellipodial leading edge drive cells to migrate in ECMs. (**a**) Cell migrations in ECMs are spatial and temporal cross-scale biophysical behaviours performed by proteins. (**b**) Demonstration of the two-dimensional mechanical interaction for theoretical analysis. (**c**) Polymerization-induced deformation analysis of a small segment ds of actin filament. The convex side surface is stretched with a strain $\varepsilon_1 = r_0 \xi$ while the concave side surface is compressed with a strain $\varepsilon_2 = -r_0 \xi$ where r_0 is the radius of actin filaments. The colour bar regions denote the binding surfaces of Arp2/3 complex on the convex and concave sides of the actin filaments. (**d**) Relationship between relative branched density and actin filament curvature from experimental data (37). It is fitted by an inverted sigmoid function, which is more reasonable than a linear fitting because the relative branched density should be neither minus nor excessively high. The relative branched density P is the ratio of the number of branch points with different curvatures to the corresponding number of mother actin filaments. (**e**) Relationship between the relative dissociation constants $K_d(\mathbf{p})/K_d^0$ and actin filament curvature where $K_d(\mathbf{p})$ and K_d^0 are the dissociation constants in the bending and straight states, respectively (Supplementary information). (**f**) The relative dissociation constants $K_d^1(\mathbf{p})/K_d^2(\mathbf{p})$ shows that the binding affinity of Arp2/3 complex on the convex side ($K_d^1(\mathbf{p})$) of a bending actin filament is much higher than the concave surface ($K_d^2(\mathbf{p})$). (**g**) Forces acting on the leading-edge membrane when a cell is migrating in ECMs. $\mathbf{f}_p$ is the propulsive force generated by the polymerization of a branched actin filament. $\mathbf{f}_{ECM}$ is the extracellular resistance from the ECM. $\mathbf{f}_m$ is the tension force of the top and bottom lamellipodial membrane. $\mathbf{f}_a$ is the attachment force of a molecular linker, which links actin filament and the leading-edge membrane.

Fig. 2. Spatiotemporal simulations from protein behaviours accurately predict self-

adaptive cell migrations in complex ECMs and reveal their underlying biophysical

mechanisms. (**a**) Resistance force-induced branched actin network density adaption.

Model predictions: temporal fluctuations of polymerizing actin cytoskeletal density at

the leading edge in HMECM (blue line) and HTECM (red line). The extracellular

resistance $\mathbf{f}_{ECM}$ of the HMECM is 0.5 nN/μm. For the HTECM condition (red line), the

$\mathbf{f}_{ECM}$ is 1.0nN/μm in the time frame 2000–5000 ms (shadow region), and is 0.5 nN/μm

in the time frames of 0–2000 ms and 5000–7000 ms. The induced actin density along the

leading edge in the two conditions is 200–350 /μm, agreeing well with the experimental

data 150–350 /μm(51). The first experimental data is the temporal fluctuations of actin

reporter lifeact: GFP intensity at the leading edge of migrating fish keratocytes in

HTECMs from experiments in ref. (1). The second experimental data is the fluctuations

of polymerizing branched actin density in responding to varying external loads from

experiments in ref.(5). Actin density is normalized by that of the unloaded condition(5).

Green region is the standard deviation. (**b**) Resistance force-induced Arp2/3 complex

density adaption. Model predictions: temporal fluctuations of the leading-edge Arp2/3

density in responding to HTECMs (n=30 each group). The Arp2/3 density is normalized

by that of the homogeneous ECM condition. The first experimental data: adaptive

fluorescent intensity of ARX-2::GFP for Arp2/3 complex at the leading edge when cells

invade ECMs with low resistance and high resistance (n=10 each group) from

experiments in ref. (11). The second experimental data: Arp2/3 complex density under

increasing resistance (n=15 each group) from experiments in ref. (25). The HR condition

is normalized by the LR condition. (**c**) Model predictions and published experimental

measurements(52) of the orientation frequency of lamellipodial actin filaments relative

to the migration direction, which is defined as $0°$ direction. (**d**) Adaptive ATP

consumption rate for fuelling cells migrating through HTECMs. Model predictions: the ATP consumption rate is the average number of ATP used for assembling actin cytoskeleton per micro-second, and is normalized by that of the HMECM condition (n=20 each group). Experiments: adaptive fluorescent intensity of mitochondria and ATP at the leading edge when cells invade ECMs with low resistance and high resistance (n=10 each group) from ref. (11). Scale bar is 5 μm. Temporal fluctuations of propulsive force (**e**) and deformation energy (**f**) generated by the leading-edge actin cytoskeleton in HMECM and HTECM. (**g**) Molecular protein-level biophysical mechanism of the positive feedback that migrating cells mechano-sense the variation of ECM and then accurately adapt their propulsive forces to overcome it. (**h**) Velocity adaptation. Model predictions: temporal fluctuations of leading-edge migration velocities in HMECM and HTECM. Experiments: temporal fluctuations of leading-edge migration velocity in responding to HTECM from the experiments in ref.(1). (**i**) Molecular protein-level biophysical mechanism of leading-edge migration velocity depending on extracellular resistance history is the temporal hysteresis of the leading-edge actin cytoskeleton remodelling and adaptation. The red and green time lines correspond to the velocity adaptations caused by the increased and decreased ECM resistances, respectively. Velocity adaptation stages I–V in (**i**) correspond to the stages I–V in (**h**), respectively.

a

f_{ECM}

170nm | 1.0nN/μm

$f_{ECM}^{right} = 5nN/μm$

20nm

110nm | 0.5nN/μm

migration

$f_{ECM}^{left} = 5nN/μm$

Left   Central   Right

b Model Predictions

Left
Central
Right

Migration distance (nm)
Time (10^3ms)

c Model Predictions

Left
Central
Right

Actin filament density Φ (/μm)
Time (10^3ms)

d Microfludic device

2μm   12μm

t=0s
t=4s
t=8s
t=20s

e Experiments

Actin flourecent intensity (a.u.)
Time (10^3ms)

f Model Predictions

Left   Central
Right

Actin monomer consumption rate (/ms)
Time (10^3ms)

g

Left   Central
Right

Arp2/3 complex consumption rate (/ms)
Time (10^3ms)

h Model Predictions

Propulsive force (pN)
Left   Central   Right

i Model Predictions

Left   Central
Right

Bending curvature (/μm)
Time (10^3ms)

j Model Predictions

Left   Central
Right

Actin cytoskeleton porosity
Time (10^3ms)

k Model Predictions

Left   Central
Right

Actin diffusion coefficient D_a (μm²/s)
Time (10^3ms)

l

① very high ECM resistance

② a. pushing force f_p increases;
b. actin diffusion coefficient decreases

f_p

⑤ Arp2/3 assembling stops

③ actin monomer insertion stops (polymerization stops)

④ filament bending curvature stop increasing

⑥ filament density and propulsive force stop increasing

stop overcoming

m

resistance-triggered positive feedback

high resistance-triggered negative feedback

high resistance -triggered negative feedback

Fig. 3. High extracellular resistance triggers negative feedback to adapt cell

morphology to circumnavigate obstacles for compromisingly pathfinding. (**a**) The

designed HTECM in simulations. It is divided into two stages, where extracellular

resistances in the first (in the range of 0–110 nm in front of the migrating cell leading

edge) and second stages (in the range of 110–300 nm in front of the migrating cell leading

edge) are 0.5 nN/μm and 1.0 nN/μm, respectively. However, the second stage has two

dense ECM regions with a very high resistance $\mathbf{f}_{ECM}^{left} = \mathbf{f}_{ECM}^{right} = 5$ nN/μm. The red arrow

denotes cell migration direction. Spatiotemporal migration distances (**b**) and leading-

edge actin filament densities (**c**) at the left, central and right parts of the leading edge. (**d**

and **e**) Microfluidic device and temporal fluctuation of actin reporter GFP-lifeact: the

actin intensity as a function of the time at the leading edge when migrating cells

encounter an obstacle (n=10 each group). Scale bar in (**d**) is 5 μm. Temporal actin

monomer (**f**) and Arp2/3 complex (**g**) consumption rates at the left, central and right local

leading edges. Actin monomer consumption rate is the ratio of the number of actin

monomers adding to the barbed ends of actin filaments to the number of uncapped actin

filaments per micro-second. Arp2/3 complex consumption rate is the number of Arp2/3

complex assembling for the actin cytoskeleton in a time span of 500 ms. (**h**) Propulsive

forces produced by each polymerizing actin filaments at the left, central and right local

leading edges. Temporal fluctuations of the average bending curvature of polymerizing

actin filaments(**i**), the leading-edge actin cytoskeleton porosity (**j**) and the actin diffusion

coefficient (**k**). (**l**) Biophysical mechanism of the high ECM resistance-triggered negative

adaptation feedback. (**m**) Demonstration of simulation results that the leading edge

circumnavigates high ECM resistance regions based on the negative feedback and opens

a channel in the weak region based on the positive feedback. The positive and negative

feedbacks work cooperatively to adapt cell morphology and drive cell migration.

Fig. 4. Directional cell migration is steered by a balanced relation between

intracellular proteins, HTECMs and chemotactic cues. (**a**) Demonstration of initial

simulation conditions that local leading edges drive cell migration at the two locations of

x_1 and x_2. Yellow and violet arrows denote the migration directions at x_1 and x_2,

respectively. The dots represent chemoattractant of actin. (**b**) Local extracellular

resistance of ECMs and local concentration of actin monomers caused by a gradient

chemoattractant at the positions of x_1 and x_2 in cases A-C. Migration distances (**c**, **e** and

g) and leading-edge actin cytoskeleton densities (**d**, **f** and **h**) in cases A-C (n=4). (**i**)

Leading-edge migration velocity for varying ECM resistance and actin concentration. (**j**)

Biophysical mechanism of directional cell migration is that it is steered by a balanced

relation between intracellular proteins for assembling leading-edge actin cytoskeleton,

HTECMs and chemotactic cues.

Table 1. RAP model systematically predicts all the key adaptive migratory behaviours
discovered in experiments.

Key lamellipodial leading edge behaviours during cell migrations

---

- ➕ Adaptive propulsive force generation based on ECM resistance (1)

- ➕ The orientation of actin filaments in lamellipodial is about $\pm 35\,^\circ$ relative to the migration direction (1, 52).

- ➕ Increased ECM resistance induces the increase of branched actin filament density (1, 5, 11).

- ➕ Increased resistance induces the decrease in the rates of actin nucleation and capping of barbed ends (25).

- ➕ Increased resistance induces the decrease in the rates of Arp2/3 branching (25).

- ➕ Overexpression of Arp2/3 complex is positively related to cell migratory ability and velocity (15, 17, 59).

- ➕ Increased ECM resistance leads to denser mitochondria at the Arp2/3 complex-assembled invasive cell leading edge and ATP can boost cancer leading edge invasion (11, 53).

- ➕ Migratory leading edge can circumnavigate obstacles and select the path of least resistance (6).

- ➕ Leading edge migration velocity depends on extracellular resistance history (1, 26).

Arp2/3 complex side
facing mother actin filament

Mother actin filament

| Arp2 | ArpC1 | ArpC3 | ArpC5 |
| Arp3 | ArpC2 | ArpC4 | Ⓜ Mother filament subunit # |

Supplementary Fig. 1. Actin-Arp2/3 complex binding interaction surfaces within the branch junction(38)**. (a)** Binding interaction surface between mother actin filament and Arp2/3 complex. Most of the binding surface locates in the groove between actin subunits of the mother actin filaments. (**b**) The protrusion helix of ArpC1 of Arp2/3 complex insert into the mother actin filament subunit M4.

Supplementary Fig. 2. Polymerizing branched actin network drives cell migration.

(**a**) Confocal image of a labelled migrating cell (actin: violet color; microtubule: cyan

color); scale bar: 10 μm. (**b**) Polymerizing branched actin filaments under the leading-

edge membrane. (**c**) The interaction between a single polymerizing branched actin

filament and the local leading-edge membrane confined by ECM is a two-dimensional

behaviour.

 Supplementary Fig. 3. The nonlinear theory is strongly validated by the finite

 element method (FEM). θ and l are inclined angle and length of an actin filament.

 (**a-d**) Nonlinear deformation behaviours of actin filaments with different lengths and

 inclined angles.

Supplementary Fig. 4. The dynamic interactions between a polymerizing actin filament and the leading-edge membrane during cell migration. (**a-c**) The dynamic evolutions of the propulsive force (**a**), deformation energy (**b**) and mean bending curvature (**c**) when an actin filament is polymerizing toward the leading-edge membrane.

(**d-f**) The dynamic evolutions of the propulsive force (**d**), deformation energy (**e**) and

mean bending curvature (**f**) when the leading-edge membrane is migrating forward. The

pink circle in (**a-c**) is the moment that the barbed end of the polymerizing actin filament

begins to attach the leading-edge membrane and the contact between them is a point. The

purple circle in (**a-c**) is the moment that the contact between the polymerizing actin

filament and the leading-edge membrane changes from a point to a line. The yellow circle

in (**a-c**) is the moment that a capping protein caps the barbed end of the polymerizing

actin filament and thus the polymerization stops. The pink circle in (**d-f**) is the moment

that the contact between the actin filament and the migrating leading-edge membrane

changes from a line to a point. The black circle in (**d-f**) is the moment that the actin

filament falls behind of the migrating leading-edge membrane and thus they lose contact

with each other.

Supplementary Fig. 5. Demonstrations of the initial condition and the simulation results of case C in the directional cell migration section. (**a**) Initial condition of case C. The distribution of actin monomers induced by a chemoattractant has a gradient from the left to the right side. The right-side ECM has a larger resistance. (**b**) Simulation result. It shows that directional cell migration is not solely steered by either the gradients of intracellular proteins induced by chemoattractant or the local stiffness of ECMs. It is a balanced competing consequence between the two.

Polymerization force-regulated actin filament-Arp2/3 complex interaction dominates self-adaptive cell migrations

Xindong Chen[1,2*], Yuhui Li[3], Ming Guo[4], Bowen Xu[1], Yanhui Ma[2], Hanxing Zhu[2*], Xi-Qiao Feng[1*]

[1] Institute of Biomechanics and Medical Engineering, AML, Department of Engineering Mechanics, Tsinghua University, Beijing 100084, China

[2] School of Engineering, Cardiff University, Cardiff CF24 3AA, UK

[3] Univ. Grenoble-Alpes, CEA, CNRS, INRA, Interdisciplinary Research Institute of Grenoble, Laboratoire de Phyiologie Cellulaire & Végétale, CytoMorpho Lab, 38054 Grenoble, France.

[4] Department of Mechanical Engineering, Massachusetts Institute of Technology, Cambridge, MA 02139, USA

This supplementary material provides the derivations of the spatiotemporal biophysical theory, and the development of the spatiotemporal multiscale dynamic modelling system.

Supplementary method

Derivations of the spatiotemporal "resistance-adaptive propulsion" (RAP) theoretical model

Lamellipodium is one of important ways driving cell migration (Supplementary Fig. 2)(1, 2). Firstly, we investigate the mechanical interactions between a polymerizing actin filament and the curved leading-edge membrane. Near the leading-edge membrane of the lamellipodium, the pointed ends of actin filaments connect with the branched actin network via Arp2/3 complex, which is a relatively rigid protein-protein binding connection(3). The branched actin network has a high stiffness(4, 5) and is fixed by adhesions during cell migration(6). Therefore, the polymerizing actin filaments are assumed to be fixed at their pointed ends. The leading-edge membrane is bent due to the propelling force of the growing branched actin filaments(7). According to the theory of differential geometry, we divide the curved leading-edge membrane into several continuous inclined flat plane segments (Supplementary Fig. 2b). The leading-edge membrane is first assumed to be fixed due to the confinement of the extracellular matrix. When a polymerizing actin filament gets in contact with the membrane, it becomes bent and the interaction force between the bent actin filament and the local plat segment of the membrane is in the direction normal to the flat segment. Though the force interaction between all the bent polymerizing actin filaments and the leading-edge membrane is in three-dimensional space, the interaction between a single polymerizing actin filament and the local flat plane segment of the membrane (i.e., inclined plane) can be described in a two-dimensional deformation plane, which contains the whole bent actin filament (Supplementary Fig. 2c and Fig. 1b in the Main Text).

In our model, the barbed and pointed ends of the polymerizing actin filament are denoted as $s=0$ and $s=l(t)$, respectively. Before the polymerizing barbed end reaches the inclined local leading-edge membrane, i.e., $l(t)\cos\beta < h$, they have no interaction force ($p=0$) and thus the actin filament is straight.

As the actin filament grows in length with the polymerizing time, its barbed end gets in contact with the membrane, i.e., $l(t)\cos\theta \geq h$, the growing filament is compressed and bent due to the constraint from the leading-edge membrane. When the angle $\beta(0)$ of the barbed end is less than $\pi/2$, the growing barbed end slides along the membrane in the deformation plane and they have a point contact. The deformed configuration of the actin filament and the interaction force p acting on the leading-edge membrane is shown in Fig. 1b in the Main Text. It is noted that for simplicity, the possible friction between the bent actin filament and the membrane is ignored, thus the force p is always in the normal direction of the local leading-edge membrane. The relation between the bending curvature and the bending moment is expressed as

$$\frac{d\beta_b(s,t)}{ds} = -\frac{p}{EI}\int_0^{s(t)}[1-\varepsilon(r,t)]\sin\beta(r,t)dr \tag{S1}$$

$$\varepsilon(r,t) = \frac{p\cos\beta(r,t)}{EA} \tag{S2}$$

where $0 \leq r \leq s$; E is Young's modulus of actin filaments and is taken as 2GPa(8); A and I are the cross-sectional area and the second moment of the cross-sectional area of the actin filaments, and they can be calculated from the diameter (7 nm) of actin filaments. At the pointed end of the actin filament, i.e., $s=l(t)$, $\beta_b[l(t),t]=\theta$. It is noted that β_b equals θ plus the local rotation of the filament resulted from bending, and β equals θ plus the local rotation of the filament resulted

from both bending and transverse shear. Thus, the angle $\beta_b(s)$ along the deformed actin filament as a result of the combined effects of bending, axial compression and polymerization growth can be determined as

$$\beta_b(s,t) = \theta + \frac{\mathrm{p}}{EI}\int_{s(t)}^{l(t)}[(1-\varepsilon(s,t))\int_0^{s(t)}(1-\varepsilon(r,t))\sin\beta(r,t)dr]ds \qquad \text{(S3)}$$

The angle β_s induced by the transverse shear deformation of the polymerizing actin filament is expressed as

$$\beta_s(s,t) = \frac{\kappa\mathrm{p}\sin\beta(s,t)}{GA} = \frac{2\kappa(1+\nu)\mathrm{p}\sin\beta(s,t)}{EA} \qquad \text{(S4)}$$

where ν is the Poisson's ratio of the actin filament material and is taken to be 0.3(8); and κ is the shape factor and is 10/9 for a circular cross-section. Thus, the total angle of the deformed actin filament is

$$\beta(s,t) = \beta_b(s,t) + \beta_s(s,t) \qquad \text{(S5)}$$

For a polymerization length of the actin filament at time t, the nonlinear function $\beta(s,t)$ can be determined using the iterative method and the following deformation compatibility condition:

$$\int_0^{l(t)}[1-\varepsilon(s,t)]\cos\beta(s,t)ds = h(t) \qquad \text{(S6)}$$

To solve the function $\beta(s,t)$, we reverse the interacting process by using the local leading-edge membrane (i.e., a plat plane) to compress the actin filament to meet the condition given by equation (S6). More specifically, the actin filament is first assumed to be straight. Then, we divide it into 100000 equal linear segments and apply the load p in the normal direction of the local leading-edge membrane to compress the actin filament at its barbed end, as shown in Fig. 1b in the Main

Text, by very small increments of Δp (0.01pN) using an iterative method. The initial axial

compressive strain ε is chosen as $\dfrac{\Delta p \cos\beta(s,t)}{EA}$ with the initial angle $\beta(s,t) = \theta$. Using the

deformation compatibility condition $h(t)$ in equation (S6), the convergent solution of $\beta(s,t)$ and

the corresponding force p acting on the leading-edge membrane can be obtained by the iterative

method. Our nonlinear theory is validated by the finite element method (Supplementary Fig. 5).

The mean curvature ξ of the geometrically nonlinear deformed actin filament is given as

$$\xi(t) = \frac{\beta(0) - \beta(l(t))}{l(t)} \tag{S7}$$

The total elastic strain energy in the actin filament, $U(t)$, is the summation of the bending energy

$U_b(t)$, axial compression energy $U_c(t)$ and transverse shearing energy $U_s(t)$, that is,

$$U(t) = U_b(t) + U_c(t) + U_s(t) \tag{S8}$$

where

$$U_b(t) = \frac{1}{2} EI \int_0^{l(t)} \left(\frac{\beta_b(t)}{ds}\right)^2 ds = \frac{p^2}{2EI} \int_0^{l(t)} \left[\int_0^{s(t)} (1 - \frac{p\cos\beta}{EA}) \sin\beta ds\right]^2 ds \tag{S9}$$

$$U_c(t) = \frac{1}{2} \int_0^{l(t)} \frac{(p\cos\beta)^2}{EA} (1 - \frac{p\cos\beta}{EA}) ds \tag{S10}$$

$$U_s(t) = \int_0^{l(t)} \frac{\kappa(1+\nu)(p\sin\beta)^2}{EA} (1 - \frac{p\cos\beta}{EA}) ds \tag{S11}$$

When the angle $\beta(0)$ at the barbed end increases to $\pi/2$, the barbed end of the actin filament will

have a line contact with the local leading-edge membrane (Fig. 1b in the Main Text). Thus, with

the continuing polymerization, the actin filament has no further deformation and the interaction

force p and the deformation energy U keep constant. It is worth mentioning that the interaction

force p is in the direction normal to the local leading-edge membrane, rather than in the direction of cell migration. The propulsive force f_p (in the direction of cell migration) acting on the local leading-edge membrane by the actin filament can be expressed by

$$f_p(t) = p(t)\cos\phi \qquad (S12)$$

where ϕ denotes the angle between the normal direction of the local leading-edge membrane and the direction of cell migration. The dynamic interactions between a polymerizing actin filament and the leading-edge membrane, which is confined by the ECM, are shown in Supplementary Fig. 4a-f. It shows that the polymerizing actin filaments experience large backward nonlinear deformations due to the constraint of the leading-edge membrane and extracellular matrix, which is validated by experimental observations (4, 5, 9), and is also the nature of the actin retrograde flow formation.

Arp2/3 complex has a heterodimer formed by the two subunits, Arp2 and Arp3. The heterodimer bonded with ATP mimics a filamentous actin dimer and thereby creates a template for branching out a daughter actin filament from a mother filament (10, 11). We analyze the conformational deformations of the curved surfaces, both convex and concave sides, of a bending mother filament that potentially interacts with Arp2/3 complex, showing that the convex surface is stretched while the concave surface is compressed (Fig. 1c). Their relative strain ε_r relating to the polymerization force-induced bending states can be quantitatively described by $\varepsilon_r = 2r_0\xi$ where r_0 is the radius of actin filaments (Fig. 1c). Experiments(12) show that *in vitro* long actin filaments (~10μm) exhibit large bending deformations under thermal fluctuations, and the relative branched density P, which is the ratio of the number of branch points with different curvatures to the corresponding number of mother actin filaments, is larger in the convex side than in the concave side. We fit the experimental data of P with an inverted

sigmoid function $P = 2/(1 + e^{0.73\xi})$ (Fig. 1d), which is more reasonable than a linear fitting because the relative branched density should be neither minus nor excessively high. The dissociation constant K_d between Arp2/3 complex and actin filaments is expressed by $K_d = [\text{Arp}][\text{AF}]/[\text{ArpAF}]$ where $[\text{Arp}]$, $[\text{AF}]$, $[\text{ArpAF}]$ are the equilibrium concentrations of Arp2/3 complex, actin filaments, and the Arp2/3-actin filament branches, respectively. Since $[\text{ArpAF}]/[\text{AF}] = P$, the dissociation constant becomes $K_d = [\text{Arp}]/P$. Thus, compared to the straight state of actin filaments, the relative curvature-dependent Arp2/3-actin filament disassociation constant $K_d(\text{p})/K_d^0$ is

$$K_d(\text{p})/K_d^0 = \frac{[\text{Arp}]/P(\xi)}{[\text{Arp}]/P(0)} = \frac{P(0)}{P(\xi)} = 0.5(1 + e^{0.73\xi}) \tag{S13}$$

where $K_d(\text{p})$ and K_d^0 are the dissociation constants in the bending and straight states, respectively (Fig. 1e). The relative $K_d(\text{p})/K_d^0$ shows that the affinity of Arp2/3 complex binding on the convex surface (negative curvature side) of actin filament is higher than the straight surface, which is higher than the concave surface (positive curvature side). This could be explained by the combination of our mechanical analysis and the recent cryo-electron structure of Arp2/3-actin filament junction(11, 13, 14) that the stretched convex surface of a bending actin filament exposes the groove binding sites for Arp2/3 complex, which facilitate Arp2/3 complex binding, while the compressed concave surface buries some binding sites (Fig. 1c and Supplementary Fig. 1). To better show this force-dependent binding affinity, we also calculate the relative disassociation constant $K_d^1(\text{p})/K_d^2(\text{p}) = P(-|\xi|)/P(|\xi|)$ where $K_d^1(\text{p})$ and $K_d^2(\text{p})$ are the dissociation constants on the convex and concave surfaces, respectively, demonstrating that the binding affinity on the convex surface is much higher than the concave surface (Fig. 1f). Our theoretical analysis has shown that although *in vivo* actin filaments are relatively short (~250nm), their polymerization force still can induce them to generate significant backward bending deformations, which could be verified by the actin retrograde flow phenomenon and the recent measurement that the bending curvature can reach up to 10 μm⁻

[1] (9). This indicates that the force-dependent actin-Arp2/3 binding affinity occurs *in vivo*. To incorporate that Arp2/3 complex has a higher binding affinity with the convex surface of actin filaments than with straight surface, we introduce a bending curvature-dependent binding factor $d^{arp}(\xi^{\max})$ to the model (Eq. S14), which is a segmented function and is defined as the space between two adjacent Arp2/3 complex branches along an actin filament, where $\xi^{\max}$ is the biggest bending curvature in the deformation history of the actin filament. Since experiments have shown that most of the space d^{arp} between two adjacent Arp2/3 branches is 100nm(1, 15), it is appropriate to hypothesize that the relation between d^{arp} and the maximum curvature $\xi^{\max}$ in deformation history follows

$$d^{arp}(\xi^{\max}) = \begin{cases} 300 & if \ 0 \le \xi^{\max} < 0.3 \\ 150 & if \ 0.3 \le \xi^{\max} < 1 \\ 100 & if \ 1 \le \xi^{\max} \end{cases} \tag{S14}$$

The units of the space d^{arp} is nm. In the sufficient amount of Arp2/3 complex condition, the number of Arp2/3 complexes binding on an actin filament with a length of l_i can be determined as

$$n_i^{arp} = l_i / d^{arp}(\xi^{\max}) \tag{S15}$$

The probabilities that an Arp2/3 complex binding on the convex and concave side of an actin filament are $P(|\xi|)/[P(|\xi|)+P(-|\xi|)]$ and $P(-|\xi|)/[P(|\xi|)+P(-|\xi|)]$, respectively. Then, Arp2/3 complexes branch out daughter actin filaments from mother actin filaments and thus form the lamellipodial branched actin filament network. During cell migration, the extracellular resistance force is normally in the range of 0.4 nN/μm to 2 nN/μm(16). The leading-edge membrane tension force is about 360 pN/μm(17). The molecular linking attachment force between the barbed ends of actin filaments and the leading-edge membrane, which drag the membrane backward(18-20), is estimated at 20 pN per molecular linker(21-24). The polymerization force of the branched actin

network pushes leading-edge membrane to overcome these resistance forces and thus drive the cell to migrate in the ECM.

Spatiotemporal multi-scale modelling system

Next, we develop a modelling system to spatiotemporally simulate that the three-dimensional lamellipodial self-assembling branched actin filaments drive cells migration forward. In this framework, the interactions between the growing branched actin filaments and the curved leading-edge membrane, which is confined by the varying immediate ECM, are described by the RAP model. The complicated stochastic assembling process and mechano-chemical reactions, such as the highly dynamic actin polymerization, capping protein inhabiting filament growth, Arp2/3 complex nucleation and molecular linker detachment between the filament barbed end and curved leading-edge membrane, are systematically incorporated.

The lamellipodium usually has a thickness of about 200 nanometers and a width of about 20-50 micrometres (25). Thus, we generate a lamellipodial domain of $10,000 \text{ nm} \times 1,000 \text{ nm} \times 200$ nm and use periodic boundaries in the width direction. During cell migration, the propulsive force of the actin filaments bends leading-edge membrane(7). By referring to the differential geometry theory, we divide the smoothly curved leading-edge membrane into N inclined flat plane segments $\mathbf{P} = [p_1, p_2, \dots, p_N]$ as shown in Supplementary Fig. 2. The kth membrane plane has a point with coordinates of $\mathbf{x}_k^p$ and a normal vector $\mathbf{x}_k^{p_n}$. Let $\mathbf{x}^p$ and $\mathbf{x}^{p_n}$ be the concatenations $[\mathbf{x}_1^{pT}, \mathbf{x}_2^{pT}, \dots, \mathbf{x}_N^{pT}]^T$ and $[\mathbf{x}_1^{p_nT}, \mathbf{x}_2^{p_nT}, \dots, \mathbf{x}_N^{p_nT}]^T$, respectively. During lamellipodium driving cell migration, the leading-edge protrude forward in the extracellular matrix. Thus, the equations $\mathbf{M}(\mathbf{P}, t)$ of these inclined flat plane segments $\mathbf{P}$ in three-dimensional space can be written as

$$\mathbf{M}(P,t) = (\mathbf{x} - \mathbf{x}^P(t)) \cdot \mathbf{x}^{P_n}(t)^T = 0 \tag{S16}$$

In our simulations, the leading-edge membrane is divided into six inclined planes. Then, we generate a certain concentrations of actin monomers C_a, Arp2/3 complex C_{arp}, capping proteins and ATP in the lamellipodial space. The distribution of actin monomers is defined by a gradient extracellular chemoattractant (Supplementary Fig. 5). These actin monomers stochastically nucleate into a certain number n of short actin filaments, and the pointed end coordinates $\mathbf{x}_i^{fp}$ of the ith actin filament are randomly and independently generated. For each an actin filament growing in the lamellipodial sheet-like space, a local spherical coordinate system is created by regarding its pointed end as the origin. This local spherical coordinate system is used to determine the orientation and the barbed end coordinates $(r_i^{fb}, \varphi_i^{fb}, \theta_i^{fb})$ of each actin filament confined in the lamellipodium. Published experimental results showed that the orientation of actin filaments relative to the cell migration direction is generally around $\pm 35°$(26). Thus, the orientation angles of the initial actin filaments are randomly produced by a normal distribution and a uniform distribution defined as

$$\varphi_i \sim U(70°,110°); \; \theta_i \sim N(\pm 35°,15°) \tag{S17}$$

The behaviour of actin is coupled to the nucleotide bound to its active site, and ATP hydrolysis is activated during polymerization(27). Thus, the polymerization process of actin filaments is simulated by adding individual actin monomers bonded by an ATP to the barbed ends. The growth rate V_{fil} of the polymerizing free barbed end of an actin filament is expressed as

$$V_{fil}(t) = \delta[\gamma(\Phi,t) \cdot \psi(D_a,t) \cdot C_a \cdot k_P(p,\theta) \cdot k_{on} - k_{off}] \tag{S18}$$

where δ is the radius of an actin monomer; C_a is the local concentration of actin monomers in the cell; $\gamma(\Phi)$ is the consuming factor of actin monomers, introducing the relation that polymerizing rate is proportional to the ratio of the concentration of actin monomers to the density of polymerizing filaments Φ (28); t^{cap} and t^{cap} are the nucleation and capping time of the actin filament, respectively. $k_P(p,\theta)$ is an interacting force-induced probability density distribution in the rate of filament polymerization due to size-dependent insertion of monomers $k_P(p,\theta_b) = \int_\Delta^\infty \lambda \exp(-\lambda\vartheta)d\vartheta / \int_0^\infty \lambda \exp(-\lambda\vartheta)d\vartheta$ (29, 30). p is the interaction force that the polymerizing actin filament acts on the leading-edge membrane; θ_b is the angle of the filament barbed end relative to the normal vector of the local membrane; ϑ is a function of p and θ_b and Δ is the size of sufficient gap to permit intercalation of monomers (29, 30). k_{on} and k_{off} are the polymerization and depolymerization rate constants, respectively. p and θ_b are calculated based on the RAP theory. ψ is a scaled diffusion coefficient of actin monomers and is to introduce the effect of actin filament density on the actin diffusion flux to the polymerizing barbed ends based on the Fick's first law of diffusion. D_a is the diffusion coefficient of actin monomers for supplying the polymerization of the leading-edge actin filaments, and is described as

$$D_a = \frac{k_B T}{6\pi\eta\delta}\exp^{(-k_C\delta\varpi^{0.75})} \tag{S19}$$

in which k_B is Boltzmann's constant, T is temperature, η is the viscosity of water at T, ϖ is the volume fraction of branched actin filament near the leading edge, and k_C is a regressed constant(31). Let $\mathbf{x}^{fp}$ be the concatenation $[\mathbf{x}_1^{fpT}, \mathbf{x}_2^{fpT}, \ldots, \mathbf{x}_n^{fpT}]^T$. Thus, before the barbed ends attach the leading-edge membrane, their coordinates $\mathbf{x}^{fb}(t)$ at time t are specified by

$$\mathbf{x}^{fb}(t) = \mathbf{r}^{fb}(t) \cdot \mathbf{A} + \mathbf{x}^{fp} \tag{S20}$$

$$\mathbf{r}^{fb}(t) = V_{fil}(t\mathbf{I} - \mathbf{t}^s) \tag{S21}$$

$$\mathbf{A} = \begin{bmatrix} \sin\varphi_1^{fb}\cos\theta_1^{fb} & \sin\varphi_1^{fb}\cos\theta_1^{fb} & \cos\varphi_i^{fb} \\ \sin\varphi_2^{fb}\cos\theta_2^{fb} & \sin\varphi_2^{fb}\cos\theta_2^{fb} & \cos\varphi_2^{fb} \\ . & . & . \\ . & . & . \\ \sin\varphi_n^{fb}\cos\theta_n^{fb} & \sin\varphi_n^{fb}\cos\theta_n^{fb} & \cos\varphi_n^{fb} \end{bmatrix} \tag{S22}$$

where $\mathbf{I}$ denotes the identity matrix and $\mathbf{t}^{nuc} = diag(t_1^{nuc}, t_2^{nuc}, \ldots, t_n^{nuc})$. t_i^{nuc} is the time when the ith actin filament nucleates and begins to polymerize. In lamellipodia, actin filaments can normally grow to a length of about 200 nm(1, 32). Thus, we assume that when an actin filament grows to its maximal length, which is generated by the normal distributions $N(200,50)$, capping proteins will cap its barbed end and the polymerization will stop. Thus, the final lengths of actin filaments can also be expressed by

$$\mathbf{r}^{fb}(t) = \int_{t^{nuc}}^{t^{cap}} V_{fil} dt = \delta \int_{t^{nuc}}^{t^{cap}} \left[\gamma(\Phi) \cdot \psi(D_a) \cdot C_a \cdot k_P(p,\theta) \cdot k_{on} - k_{off} \right] dt \quad \text{when } (t > t^{cap}) \tag{S23}$$

where $\mathbf{t}^{cap} = diag(t_1^{cap}, t_2^{cap}, \ldots, t_n^{cap})$ with t_i^{cap} being the time that a capping protein binds on the barbed end of the ith actin filament. The final coordinate z_i^{fb} of its barbed end in the lamellipodial sheet-like space should be within the domain given by

$$\left\{ z_i^{fb} \mid 0 \le z_i^{fb} \le 200 \right\} \tag{S24}$$

Once their barbed ends reach the leading-edge membrane, the spatiotemporal mechanical interaction between the polymerizing actin filaments and the bent protruding leading-edge membrane can be described by our RAP theoretical model. The whole process of lamellipodia protrusion is in a highly dynamic state. N-WASP proteins form molecular linkers between the

barbed ends of actin filaments and the leading-edge membrane. Meanwhile, Arp2/3 complexes are activated by Wiskott-Aldrich syndrome proteins (WASPs). Then, they bind on actin filaments and branch out new actin filaments. The probabilities of the Arp2/3 complex binding on the convex and concave sides of an actin filament are $P(|\xi|)/[P(|\xi|)+P(-|\xi|)]$ and $P(-|\xi|)/[P(|\xi|)+P(-|\xi|)]$, respectively. To identify the binding coordinates of the Arp2/3 complex along actin filaments during this dynamic process, we create a series of local Cartesian coordinate systems (x',y',z') by regarding the pointed end of each actin filament as the origin O'. Plane $.O'x'y'.$ is the deformation plane of the bent actin filament and axis $O'x'$ is the normal vector of the inclined leading-edge membrane flat plane segment. For the jth Arp2/3 complex on the ith actin filament, which pushes against the inclined leading-edge membrane plane segment k, the deflection vector $\Delta\mathbf{x}'_{ikj}(t)$ of the binding point of the Arp2/3 complex in the local coordinate system $O'_{ik}x'_{ik}y'_{ik}z'_{ik}$ at time t can be calculated with the RAP theoretical model. If there are m Arp2/3 complexes binding on the ith actin filament, let $\Delta\mathbf{x}'_{ik}(t)=[\Delta\mathbf{x}'_{ik1}(t)^T,\Delta\mathbf{x}'_{ik2}(t)^T,......,\Delta\mathbf{x}'_{ikm}(t)^T]^T$. Then, with Jacobian determinant $\mathbf{J}_{ik}$, the dynamic coordinates of the start point of Arp2/3 complex $\mathbf{x}^{as}_{ik}(t)=[\mathbf{x}^{as}_{ik1}(t)^T,\mathbf{x}^{as}_{ik2}(t)^T,......,\mathbf{x}^{as}_{ikm}(t)^T]^T$ in the global Cartesian coordinate system at time t can be obtained as

$$\mathbf{x}^{as}_{ik}(t)=\Delta\mathbf{x}'_{ik}(t)\cdot\mathbf{J}^T_{ik}+V_{fil}\boldsymbol{\lambda}_{ik}\cdot(\mathbf{t}^{arp}_{ik}-t^{nuc}_i\mathbf{I}_i)\cdot\mathbf{B}_i+\mathbf{x}^{fp}_i \tag{S25}$$

$$\mathbf{J}_{ik}=\begin{vmatrix} \dfrac{\partial x}{\partial x'_{ik}} & \dfrac{\partial x}{\partial y'_{ik}} & \dfrac{\partial x}{\partial z'_{ik}} \\[2mm] \dfrac{\partial y}{\partial x'_{ik}} & \dfrac{\partial y}{\partial y'_{ik}} & \dfrac{\partial y}{\partial z'_{ik}} \\[2mm] \dfrac{\partial z}{\partial x'_{ik}} & \dfrac{\partial z}{\partial y'_{ik}} & \dfrac{\partial z}{\partial z'_{ik}} \end{vmatrix} \tag{S26}$$

$$\mathbf{B}_i = \begin{bmatrix} \sin\varphi_i^{fb}\cos\theta_i^{fb} & \sin\varphi_i^{fb}\sin\theta_i^{fb} & \cos\varphi_i^{fb} \\ \sin\varphi_i^{fb}\cos\theta_i^{fb} & \sin\varphi_i^{fb}\sin\theta_i^{fb} & \cos\varphi_i^{fb} \\ . & . & . \\ . & . & . \\ \sin\varphi_i^{fb}\cos\theta_i^{fb} & \sin\varphi_i^{fb}\sin\theta_i^{fb} & \cos\varphi_i^{fb} \end{bmatrix} \tag{S27}$$

$$\mathbf{x}_i^{fp} = \begin{bmatrix} x_i & y_i & z_i \\ x_i & y_i & z_i \\ . & . & . \\ . & . & . \\ x_i & y_i & z_i \end{bmatrix} \tag{S28}$$

where $\boldsymbol{\lambda}_{ik} = diag(\lambda_{ik1}, \lambda_{ik2}, ..., \lambda_{ikm})$ and $\mathbf{t}_{ik}^{arp} = diag(t_{ik1}^{arp}, t_{ik2}^{arp}, ..., t_{ikm}^{arp})$. t_{ikj}^{arp} denotes the time that the jth Arp2/3 complex attaches the ith actin filament due to the mechanical interaction with the local leading-edge membrane plane segment k. λ_{ikj} is randomly generated in the range of (0, 1). Matrices $\mathbf{B}_i$, $\mathbf{x}_i^{fp}$ and $\mathbf{I}_i$ all have m rows. The Jacobian determinant is calculated from the transformation relationship between the local ($x'_{ik} y'_{ik} z'_{ik}$) and global (xyz) Cartesian coordinate systems.

The length of Arp2/3 complex r^{arp} is normally about 10 nm (33). The subunits Arp2 and Arp3 bind ATP via nucleotide-binding domains, and then assemble branched actin network through hydrolysis of ATP(34). During this process, Arp2/3 complex binds on the convex side of a bent actin filament (12) with an angle of ~70 °(3). Thus, the possible branching position of Arp2/3 complex is constrained on a half conical surface around the actin filament. The azimuthal angle φ_{ikj}^{arp} of Arp2/3 complex in its local spherical system is also generated by a uniform distribution $U(70°,110°)$. Because the forthcoming daughter actin filament will grow out from the Arp2/3 complex, angle φ_{ikj}^{arp} also should allow the daughter filament to polymerize to a length r_{ikj}^{df} specified by a normal distribution $N(200,50)$. Therefore, the azimuthal angle φ_{ikj}^{arp} and the end point

coordinates $\mathbf{x}_{ikj}^{ae}(t)$, i.e., the vertical vector $[x_{ikj}^{ae}(t), y_{ikj}^{ae}(t), z_{ikj}^{ae}(t)]$, of the jth Arp2/3 complex on the ith actin filament interacting with the local leading-edge membrane segment k at time t should satisfy

$$\left\{ \varphi_{ikj}^{arp} \mid 0 \leq z_{ikj}^{ae}(t) + r^{df} \cos \varphi_{ikj}^{arp} \leq 150 \right\} \tag{S29}$$

$$r^{arp2} - (\mathbf{x}_{ikj}^{ae}(t) - \mathbf{x}_{ikj}^{as}(t))^{T} \cdot (\mathbf{x}_{ikj}^{ae}(t) - \mathbf{x}_{ikj}^{as}(t)) = 0 \tag{S30}$$

$$\left[\Delta\mathbf{x}_{ikj}^{local}(t)^{T} \cdot (\mathbf{x}_{ikj}^{ae}(t) - \mathbf{x}_{ikj}^{as}(t)) \right]^{2} = (\cos\alpha)^{2} \Delta\mathbf{x}_{ikj}^{local}(t)^{T} \cdot \Delta\mathbf{x}_{ikj}^{local}(t)(\mathbf{x}_{ikj}^{ae}(t) - \mathbf{x}_{ikj}^{as}(t))^{T} \cdot (\mathbf{x}_{ikj}^{ae}(t) - \mathbf{x}_{ikj}^{as}(t))$$

(S31)

where $\Delta\mathbf{x}_{ikj}^{local}(t)$ is the local tangent vector of the bent actin filament segment where Arp2/3 complex binds. This vector updates with time because of actin filament polymerization, leading-edge membrane protrusion and their dynamic interaction, and can be calculated with the RAP model. α denotes the angle between the mother filament and the Arp2/3 complex, and is generated by a Gaussian distribution of $N(70°, 2°)$. Thus, the end point coordinates $\mathbf{x}_{ikj}^{ae}(t)$ of Arp2/3 complex can be specified and updated by using the equations (S29-31).

After the binding of Arp2/3 complex, daughter actin filaments will nucleate there. The concatenation matrix of the pointed ends of daughter actin filaments $\mathbf{x}^{dfp}(t) = [\mathbf{x}_1^{ae}(t)^{T}, \mathbf{x}_2^{ae}(t)^{T}, \ldots, \mathbf{x}_q^{ae}(t)^{T}]^{T}$ and q is the total number of Arp2/3 complex. Then, the daughter filaments polymerize by adding actin monomers onto their barbed ends in the branching direction of the Arp2/3 complex. The positions of the daughter filaments are also in a dynamic state with Arp2/3 complex, and can be determined by the deflection equations of their mother filaments calculated from the RAP model. Like the mother actin filaments, the polymerizing daughter filaments will also mechanically interact with the leading-edge membrane and be bent. Their spatiotemporal positions are updated by the RAP model. New activated Arp2/3 complex by

WASPs will also bind on these daughter filaments and branch out next generation daughter actin filaments. With the increase in the number of actin filaments pushing against the leading-edge membrane, the leading-edge membrane will protrude forward. Consequently, these bent actin filaments will straighten, either partially or totally. Meanwhile, the assembling and mechano-chemical reactions between these different proteins are constantly taking place, remodeling and organizing new lamellipodial cytoskeleton to drive cell migrations in ECMs.

Supplementary Table 1. Parameter values

Symbol	Meaning	Value	Source
C_a	Concentration of actin monomer	10-50 μM	(29, 35, 36)
δ	Radius of actin monomer	2.5 nm	(37, 38)
l	Length of actin filament	10-300 nm	(32, 39)
k_{on}	Polymerization rate	$10 \text{ s}^{-1} \text{ μM}^{-1}$	(29, 37)
k_{off}	Depolymerization rate	1 s^{-1}	(29, 37)
ξ	Mean curvature of a bent actin filament	$0\text{-}1.5 \text{ μm}^{-1}$	(12, 40)
E	Young's modulus of actin filament	1-2GPa	(8, 41)
v	Poisson's ratio of actin filament	0.3	(42)
A	Cross-sectional area	38.5 nm^2	(8, 43)
I	The second moment of the cross-sectional area	118 nm^4	(8, 43)
k	Shape factor of the circular cross section	10/9	(44)
f_a	Attachment force per actin filament	20 pN	(21-24)
f_m	Tension force of membrane	320 pN/μm	(17)
f_{ECM}	Resistance force of ECM	400-2600 pN/μm	(16, 45, 46)
k_B	Boltzmann constant	$1.380649 \times 10^{-23} \text{ J/K}^{-1}$	(30, 31)
η	Viscosity of water	$1.0 \times 10^{-3} \text{ Pa s}$	(31)

Supplementary Table 2. Other notations

Symbol	Meaning
p	Interacting force between a polymerizing actin filament and the leading-edge membrane (pN)

ε	Compressive strain
θ	Angle between the normal direction of the local leading-edge membrane and the direction of cell migration (°)
f_p	Propulsive force generated by a polymerizing actin filament in cell migration direction (pN) $f_p = p\cos\phi$
β	Local angle of a bent actin filament relative to the normal direction of the local leading-edge membrane it interact with (°)
U	Total elastic strain energy of an actin filament (10^{-19}N·m)
U_b	Bending deformation energy of an actin filament (10^{-19}N·m)
U_c	Compression deformation energy of an actin filament (10^{-19}N·m)
U_s	Transverse shearing deformation energy of an actin filament (10^{-19}N·m)
Φ	Density of free barbed ends of actin filaments (when C_a =40 μM, Φ_0 = 240/μm)
γ	Scaled density of free barbed ends. $\gamma = \Phi / \Phi_0$
d^{arp}	Distance between two adjacent Arp2/3 complex branches along an actin filament (nm)
n^{arp}	Number of Arp2/3 complexes binding on an actin filament
t	Time (ms)
t_{nuc}	Starting time of actin filament nucleation (ms)
N	Number of actin filaments pushing the leading-edge membrane
α	Percentage of actin filaments attaching the leading-edge membrane ($\alpha = 10\%$)
Ω	Number of attaching molecular linker between the barbed ends of actin filaments and the membrane. $\Omega = \alpha N$
S	Migration distance of the leading edge (nm)
V	Migration velocity of the leading edge (μm/min)
D_a	Diffusion coefficient of actin monomers (μm²/s). When Φ_0 =240/μm, diffusion coefficient is D_a^0
ϖ	Volume fraction of branched actin filament at the leading edge
T	Absolute temperature (K)
ψ	Scaled diffusion coefficient of actin monomers ($\psi = D_a / D_a^0$, where D_a^0 is the diffusion coefficient of actin monomers when Φ_0 = 240/μm)
m	Number of mother filaments
Δ	size of sufficient gap to permit intercalation of monomers(29, 30)

References

1. Pollard TD & Borisy GG (2003) Cellular motility driven by assembly and disassembly of actin filaments. *Cell* 112(4):453-465.

2. Pollard TD (2003) The cytoskeleton, cellular motility and the reductionist agenda. *Nature* 422(6933):741.

3. Blanchoin L*, et al.* (2000) Direct observation of dendritic actin filament networks nucleated by Arp2/3 complex and WASP/Scar proteins. *Nature* 404(6781):1007.

4. Chaudhuri O, Parekh SH, & Fletcher DA (2007) Reversible stress softening of actin networks. *Nature* 445(7125):295.

5. Bieling P*, et al.* (2016) Force Feedback Controls Motor Activity and Mechanical Properties of Self-Assembling Branched Actin Networks. *Cell* 164(1-2):115-127.

6. Case LB & Waterman CM (2015) Integration of actin dynamics and cell adhesion by a three-dimensional, mechanosensitive molecular clutch. *Nature cell biology* 17(8):955.

7. Simon C*, et al.* (2019) Actin dynamics drive cell-like membrane deformation. *Nature Physics*:1.

8. Broedersz CP & MacKintosh FC (2014) Modeling semiflexible polymer networks. *Reviews of Modern Physics* 86(3):995.

9. Reynolds MJ, Hachicho C, Carl AG, Gong R, & Alushin GM (2022) Bending forces and nucleotide state jointly regulate F-actin structure. *Nature* 611(7935):380-386.

10. Shaaban M, Chowdhury S, & Nolen BJ (2020) Cryo-EM reveals the transition of Arp2/3 complex from inactive to nucleation-competent state. *Nature Structural & Molecular Biology* 27(11):1009-1016.

11. Chou SZ, Chatterjee M, & Pollard TD (2022) Mechanism of actin filament branch formation by Arp2/3 complex revealed by a high-resolution cryo-EM structureof the branch junction. *Proceedings of the National Academy of Sciences* 119(49):e2206722119.

12. Risca VI*, et al.* (2012) Actin filament curvature biases branching direction. *Proceedings of the National Academy of Sciences* 109(8):2913-2918.

13. Fäßler F, Dimchev G, Hodirnau V-V, Wan W, & Schur FKM (2020) Cryo-electron tomography structure of Arp2/3 complex in cells reveals new insights into the branch junction. *Nature Communications* 11(1):6437.

14. Ding B*, et al.* (2022) Structure of Arp2/3 complex at a branched actin filament junction resolved by single-particle cryo-electron microscopy. *Proceedings of the National Academy of Sciences* 119(22):e2202723119.

15. Svitkina TM & Borisy GG (1999) Arp2/3 complex and actin depolymerizing factor/cofilin in dendritic organization and treadmilling of actin filament array in lamellipodia. *The Journal of cell biology* 145(5):1009-1026.

16. Abraham VC, Krishnamurthi V, Taylor DL, & Lanni F (1999) The actin-based nanomachine at the leading edge of migrating cells. *Biophysical journal* 77(3):1721-1732.

17. Lieber AD, Schweitzer Y, Kozlov MM, & Keren K (2015) Front-to-rear membrane tension gradient in rapidly moving cells. *Biophysical journal* 108(7):1599-1603.

18. Welf ES*, et al.* (2020) Actin-membrane release initiates cell protrusions. *Developmental Cell*.

19. Mehidi A*, et al.* (2021) Forces generated by lamellipodial actin filament elongation regulate the WAVE complex during cell migration. *Nature Cell Biology*:1-15.

20. Co C, Wong DT, Gierke S, Chang V, & Taunton J (2007) Mechanism of actin network attachment to moving membranes: barbed end capture by N-WASP WH2 domains. *Cell* 128(5):901-913.

21. Dudko OK, Hummer G, & Szabo A (2008) Theory, analysis, and interpretation of single-molecule force spectroscopy experiments. *Proceedings of the National Academy of Sciences* 105(41):15755-15760.

22. Müller DJ, Helenius J, Alsteens D, & Dufrêne YF (2009) Force probing surfaces of living cells to molecular resolution. *Nature chemical biology* 5(6):383-390.

23. Evans EA & Calderwood DA (2007) Forces and bond dynamics in cell adhesion. *Science* 316(5828):1148-1153.

24. Benoit M, Gabriel D, Gerisch G, & Gaub HE (2000) Discrete interactions in cell adhesion measured by single-molecule force spectroscopy. *Nature cell biology* 2(6):313-317.

25. Giri A*, et al.* (2013) The Arp2/3 complex mediates multigeneration dendritic protrusions for efficient 3-dimensional cancer cell migration. *The FASEB Journal* 27(10):4089-4099.

26. Weichsel J & Schwarz US (2010) Two competing orientation patterns explain experimentally observed anomalies in growing actin networks. *Proceedings of the National Academy of Sciences* 107(14):6304-6309.

27. Merino F*, et al.* (2018) Structural transitions of F-actin upon ATP hydrolysis at near-atomic resolution revealed by cryo-EM. *Nature structural & molecular biology* 25(6):528-537.

28. Lee CW*, et al.* (2013) Dynamic localization of G-actin during membrane protrusion in neuronal motility. *Current Biology* 23(12):1046-1056.

29. Mogilner A & Oster G (1996) Cell motility driven by actin polymerization. *Biophysical journal* 71(6):3030-3045.

30. Mogilner A & Oster G (2003) Force generation by actin polymerization II: the elastic ratchet and tethered filaments. *Biophysical journal* 84(3):1591-1605.

31. Amsden B (1998) Solute diffusion within hydrogels. Mechanisms and models. *Macromolecules* 31(23):8382-8395.

32. Bear JE*, et al.* (2002) Antagonism between Ena/VASP proteins and actin filament capping regulates fibroblast motility. *Cell* 109(4):509-521.

33. Robinson RC*, et al.* (2001) Crystal structure of Arp2/3 complex. *Science* 294(5547):1679-1684.

34. Dayel MJ, Holleran EA, & Mullins RD (2001) Arp2/3 complex requires hydrolyzable ATP for nucleation of new actin filaments. *Proceedings of the National Academy of Sciences* 98(26):14871-14876.

35. Cooper JA (1991) The role of actin polymerization in cell motility. *Annual review of physiology* 53(1):585-605.

36. Marchand J-B*, et al.* (1995) Actin-based movement of Listeria monocytogenes: actin assembly results from the local maintenance of uncapped filament barbed ends at the bacterium surface. *The Journal of Cell Biology* 130(2):331-343.

37. Boujemaa-Paterski R*, et al.* (2017) Network heterogeneity regulates steering in actin-based motility. *Nature communications* 8(1):655.

38. Bieling P, Weichsel J, Mullins RD, & Fletcher DA (2022) The molecular mechanism of load adaptation by branched actin networks. *Elife* 11:e73145.

39. Bailly M*, et al.* (1999) Relationship between Arp2/3 complex and the barbed ends of actin filaments at the leading edge of carcinoma cells after epidermal growth factor stimulation. *The Journal of cell biology* 145(2):331-345.

40. Small JV, Winkler C, Vinzenz M, & Schmeiser C (2011) Reply: Visualizing branched actin filaments in lamellipodia by electron tomography. *Nature Cell Biology* 13(9):1013.

41. de Pablo PJ, Schaap IA, MacKintosh FC, & Schmidt CF (2003) Deformation and collapse of microtubules on the nanometer scale. *Physical review letters* 91(9):098101.

42. Lu L, Oswald SJ, Ngu H, & Yin FC-P (2008) Mechanical properties of actin stress fibers in living cells. *Biophysical journal* 95(12):6060-6071.

43. Blanchoin L, Boujemaa-Paterski R, Sykes C, & Plastino J (2014) Actin dynamics, architecture, and mechanics in cell motility. *Physiological reviews* 94(1):235-263.

44. Gere & JamesM (*Mechanics of materials. 7th ed* (Mechanics of materials. 7th ed).

45. Chang J, Pang EM, Adebowale K, Wisdom KM, & Chaudhuri O (2020) Increased stiffness inhibits invadopodia formation and cell migration in 3D. *Biophysical Journal* 119(4):726-736.

46. Marcy Y, Prost J, Carlier M-F, & Sykes C (2004) Forces generated during actin-based propulsion: a direct measurement by micromanipulation. *Proceedings of the National academy of Sciences of the United States of America* 101(16):5992-5997.