Title: Learning induces unique transcriptional landscapes in the auditory cortex

Authors: Graham, G.[1,2], Chimenti, M.S.[3], Knudtson, K.L.[3], Grenard, D.N.[2], Co, L.[2], Sumner, M.[2], Tchou, T.[2], Bieszczad, K.M.[1,2,4,5]

[1]Neuroscience Graduate Program, Rutgers Univ., Piscataway, NJ
[2]Behavioral and Systems Neuroscience, Dept. of Psychology, Rutgers Univ., Piscataway, NJ
[3]Iowa Institute of Human Genetics, Univ. of Iowa Carver College of Medicine, Iowa City, IA
[4]Rutgers Center for Cognitive Science, Rutgers Univ., Piscataway, NJ
[5]Dept. of Otolaryngology-Head and Neck Surgery, Rutgers Robert Wood Johnson Medical School, New Brunswick, NJ

Abstract: Learning can induce neurophysiological plasticity in the auditory cortex at multiple timescales. Lasting changes to auditory cortical function that persist over days, weeks, or even a lifetime, require learning-induced *gene expression.* Indeed, *de novo* transcription is the molecular determinant for whether transient experiences transform into long-term memories with a lasting impact on behavior. However, auditory cortical genes that support auditory learning, memory, and acquired sound-specific behavior are largely unknown. This report is the first to identify genome-wide changes in learning-induced gene expression within the auditory cortex thought to underlie the formation of auditory memory. Bioinformatic analyses on gene enrichment profiles from RNA sequencing identified biological pathways that include *cholinergic synapses* and *neuroactive receptor interactions*. The findings characterize key candidate effectors underlying changes in cortical function that support the formation of long-term auditory memory in the adult brain. The molecules and mechanisms identified are potential therapeutic targets to facilitate long-term and sound-specific changes to auditory function in adulthood and are now prime for future gene-targeted investigations.

Manuscript:

A well-accepted concept in the field of learning and memory is that memories are stored where they are processed (Nadel & Hardt, 2011). Biological events known as memory consolidation can stabilize transient neural representations evoked by a sensory experience (Lechner, Squire, & Byrne, 1999; McGaugh, 2000; Dudai, 2012). Fundamental and evolutionarily conserved mechanisms that initiate memory consolidation are transcription and translation, defined as the active expression of genes and their subsequent protein products, respectively (Alberini & Kandel, 2014; Costa-Mattioli et al., 2009). We hypothesized that sound discrimination learning induces *de novo* gene expression within the auditory cortex. Highly specific representations of sound cues can outlast the transience of experience (seconds and minutes) by consolidating into long-term memory (hours to days) that later guides sound-cued behavior. While distinctive regional transcriptomic profiles are thought to be responsible for memory consolidation (Katzman, et al., 2021), learning-induced transcription events that support memory formation in the central auditory system are severely under-described. In contrast, the auditory cortex has been exceptionally well-described in studies of auditory learning and memory at the level of *neurophysiological* changes particularly in receptive fields and tonotopic maps (Schreiner & Polley, 2014; Weinberger N. M., 2015; Pienkowski & Eggermont, 2011), cortico-cortical and cortico-fugal connectivity (Souffi et al., 2021; Lesicko & Geffen, 2022; Schreiner & Polley, 2014;

Liu et al., 2011; Xiong, Znamenskiy, & Zador, 2015), including at multiple time scales (Froemke & Martins, 2011; Froemke & Schreiner, 2015; Fritz, Elhilali, David, & Shamma, 2007; Tchernichovski & Margoliash, 2013). Moreover, neurophysiological changes are linked to behavior, e.g., for cue-directed action (Letzkus, Wolff, & Lüthi, 2015), attention (Fritz et al., 2007; Elhilali et al., 2007) and memory for sound signals (Bieszczad & Weinberger 2010; Grosso et al., 2015; Aschauer & Rumpel, 2016; Concina, Renna, Grosso, & Sacchetti, 2019; Letzkus, Wolff, & Lüthi, 2015; Ghosh & Zador, 2021). Decades of evidence for learning-induced neurophysiological plasticity in the auditory cortex has shown shared behavioral characteristics of auditory memory (Weinberger 2007a; 2007b), making it a top candidate region for auditory memory consolidation. By investigating auditory cortex bioinformatically, we also leverage an unbiased opportunity to uncover brain-wide common or distinct biological gatekeepers of neuroplasticity underlying adaptive auditory function. From a broader perspective, regional transcriptomic profiling can lead to a more complete understanding of how the sensory system is modified by experience in the service of memory. Learning-induced transcriptomic profiles identified within the auditory cortex can validate, extend, or lead to new biological models of processes that support or impair adaptive auditory processing. The consolidation of auditory memory is likely a product of experience-dependent changes in gene expression that affect cellular functions within the auditory cortex, which in turn promote lasting changes to sound-evoked neurophysiological responsivity that alter sound-cued behaviors.

To identify learning-induced transcripts, expression profiling analysis, using RNA sequencing (RNAseq) was performed on samples of anatomically defined auditory cortex (Bregma -3.10 mm, Interaural 6.90 mm; Paxinos & Watson, 2007) from water-restricted adult rats trained to bar press to pure tones for water rewards. Responses to a target pure tone frequency (5.0 kHz; 65 dB SPL) resulted in reward, while responses to the non-target (11.5 kHz; 65 dB SPL) were unrewarded and initiated a time-out period that extended the time to the next trial. This two-tone discrimination task (2TD) is not difficult perceptually; the acoustic frequencies are over an octave apart and easily distinguished by rodents (Talwar & Gerstein, 1998). Rather, the behavioral challenge was associative in nature: 2TD performance demands memory for which tone is associated with reward (vs. no reward). Trained rats (N=8) were sacrificed after three consecutive daily 45-minute 2TD training sessions and compared to a group of sound-naïve rats (N=4). This is a time point early in training when animals are still acquiring the 2TD task. Early task acquisition was targeted to capture initial learning-induced transcriptional events that set the stage for later increases in 2TD performance observed in behavior over weeks of training. For example, the average performance was above chance, but only 66±7.81% after 3 days, compared to ≥90% after extended training (Shang, Bylipudi, & Bieszczad, 2019). To capitalize on an opportunity to identify the expression of genes relevant for successful sound-specific associative memory, we leveraged an HDAC-inhibitor that targets epigenetic regulation of activity-dependent gene expression in learning and memory formation (McQuown, et al., 2011; Kwapis, et al., 2017; Malvaez, et al., 2012). Importantly, a decade of work has shown that HDAC-inhibition can facilitate learning-induced neurophysiological plasticity in sound-evoked auditory cortical responses (relative to vehicle) and enhances auditory discriminative behaviors (Bieszczad et al., 2015; Phan et al., 2017; Shang, Bylipudi, & Bieszczad, 2019; Rotondo & Bieszczad, 2020; Rotondo & Bieszczad 2021a; 2021b) including in humans (Gervain, et al., 2013). Half of the trained animals were treated with the HDAC-inhibitor,

RGFP966 (N=4; 10 mg/kg, ApexBio, cat#A8803; *sub.cu.* injection), while the other half were identically trained but administered vehicle solution (N=4; matched volume, *sub.cu.* injection; **Fig. 1a**). There were no differences in 2TD performance between groups at the time of brain collection (RGFP966: 61±5.0% vs. Vehicle: 71±7.0%; $t_{(5.9166)}$ = -1.21, p=0.272; Welch's t-test). Brains were promptly collected and flash-frozen at a timepoint consistent with the peak concentration of the inhibitor in auditory cortex, one hour after the post-session injection of either the HDAC-inhibitor or vehicle (Bieszczad et al., 2015).

The findings herein are the first to identify a transcriptomic profile of associative learning within the auditory cortex. Learning in the two-tone discrimination task induced changes in the transcription levels of hundreds of genes (compared to sound-naïve) (**Fig. 1b**). A hierarchical clustering algorithm showed that genes are uniquely upregulated or downregulated, revealing a complex network of cortical gene expression events induced by auditory learning (**Fig. 2a**). Enrichment analyses (iPathwayGuide™; Impact Analysis method) identified the *cholinergic synapse* (differentially expressed genes (DEG) / all genes (ALL): 22/101; p = 0.004, Bonferroni correction) as the top biological pathway (**Table 1**). This result is consistent with research since the 1990s highlighting the sufficiency of cholinergic signaling in the auditory cortex for neurophysiological plasticity and related auditory behavior (Froemke & Martins, 2011; Weinberger, 2003; Bakin & Weinberger, 1996; Kilgard & Merzenich, 1998). Transcript levels induced by learning under HDAC-inhibition were either amplified in the same direction (by further increases or further decreases in unique gene transcript levels) (**Fig. 2b**), or blunted compared to learning alone (**Fig. 2c**). One top biological pathway under these conditions was the *Neuroactive ligand-receptor interaction* (DEG/ALL: 30/194; p = 0.016; **Table 1**) which involved effectors crucial to neural activation. Another top pathway was the *extracellular matrix-receptor (ECM-receptor) interaction* (DEG/ALL: 14/69; p = 0.04; **Table 1**). Components of the ECM are critical for cortical plasticity and memory consolidation (Happel, et al., 2014; Banerjee, et al., 2017; El-Tabbal, et al., 2021; Sonntag, et al., 2015). These pathways offer potential alternatives, perhaps complementary, to cholinergic signaling that can facilitate lasting experience-dependent changes in auditory function (Ji, Gao, & Suga, 2001; Luo & Yan, 2013; Metherate, 2011). Other genes whose expression changed with auditory learning but no further with HDAC-inhibition were likely related to procedural conditions of the task, rather than to sound-specific associative memory. For example, the *apelin signaling pathway* was identified (DEG/ALL: 20/116; p = 0.0005), which is important in the brain for the homeostatic regulation of water intake (Hu, et al., 2021). A direct comparison of transcript levels between the two groups of trained rats (learning with or without HDAC-inhibition) showed very few uniquely differentially expressed genes (DEGs). A threshold appreciably used to identify the most likely DEGs (p = 0.05) found only *Adamts13*, *U6*, *Rexo4*, and *Cabin1* were differentially expressed (**Fig. 2d**). Thus, the major effect of HDAC-inhibition appears to modulate the expression of genes induced by auditory learning under normal conditions, rather than to newly recruit unique genes to subserve sound-specific memory. Together, these findings show large-scale transcriptomic changes occur in the auditory cortex early in training as adult animals learn to discriminate associative relationships between sound cues. Together with established neurophysiological and behavioral reports of HDAC-inhibition to promote auditory function, these findings support the tactic of using an HDAC-inhibitor to distinguish genes whose expression determines the success of auditory memory formation.

Several genes of interest (GOIs) were selected from the enrichment analyses in biological pathways to validate genome-wide sequencing with a gene-targeted approach. Samples were collected from separate cohorts that replicated the two groups of trained and treated animals at the same early-in-training timepoint (i.e., one hour after the third 2TD training session) for use in gene expression analysis using quantitative real-time polymerase chain reaction (qRT-PCR). There were no 2TD performance differences in replicate cohorts (RGFP966$_1$: 61±5.0% vs. RGFP966$_2$: 67±6.0%; $t_{(8.441)}$ = -0.834, p = 0.4274; and Vehicle$_1$: 71±6.0% vs. Vehicle$_2$: 74±3.0%; $t_{(4.0809)}$ = -0.444, p = 0.6796; Welch's t-test). The first GOI was *Egr1,* a well-studied, activity-dependent "immediate-early gene" known to peak within the first hours of learning and critical for memory formation (Duclot & Kabbaj, 2017). It was a top learning-induced gene with confirmed increased expression that was likewise amplified with HDAC-inhibition in the separate cohort of animals in gene-targeted study (**Fig. 3**). The same was true for *Per2*, a gene involved in the regulation of circadian rhythms and serotoninergic pathways in the brain (Bae, et al., 2001; Albrecht, et al., 2001; Cuesta, et al., 2009; Reh, et al., 2020). *Per2* is part of the "clock gene" family that has been linked to HDAC modulation in other brain regions (Kwapis, et al., 2018) and may have a role in auditory function as well (Reh et al., 2020). In contrast, some GOIs identified genome-wide were only partially confirmed. *Chrna7* encodes a subunit of the dynamic nicotinic acetylcholine receptor (nAChR) that was found to be downregulated with auditory learning, consistent with neurophysiological evidence (Takesian, et al., 2018; Kuchibhotla, et al., 2017). Thus, we expected the gene-targeted study to confirm *Chrna7* down-regulation in all 2TD trained animals, but down-regulation was apparent only in vehicle-treated animals that learned 2TD without HDAC-inhibition. Additional selected GOIs were from the *Nr4a* orphan nuclear receptor family, *Nr4a1* and *Nr4a2*, which are immediate early genes reported to be necessary for propagating the downstream effects of the selective HDAC target of RGFP966 in other brain regions (McQuown et al., 2011; Kwapis, et al., 2019). Consistent with prior reports of task-dependent regional differences in *Nr4a* family gene expression (McNulty, et al., 2012), *Nr4a1* but not *Nr4a2* was up-regulated in auditory cortex after auditory learning. Other DEGs, *Htr1a* or *Adamts13*, were not confirmed by gene-targeted study likely due to differences in known transcript variants not detected by our custom-designed gene-targeted probes, or due to type I error, or to subtle behavioral differences between trained cohorts of animals that went undetected in 2TD performance measures. We also investigated *Lynx1*, a gene with a long history of function for re-opening "critical period"-like plasticity in sensory cortex (Morishita et al., 2010), likely by its action on cholinergic and serotonergic modulation (Takesian, et al., 2018). However consistent with prior genome-wide reports that failed its detection (Kalish, et al., 2020), it was neither detected in our RNAseq dataset, nor in gene-targeted study. Though it remains a challenge to determine whether discrepancies can be explained by real biological or by individual variability between animals, or due to technical variability with low abundance or cell-type specific expressing transcripts, we report exceptionally consistent results for some GOIs. As the learning-induced expression of *Chrna7*, *Egr1*, and *Per2* is consistent between the gene-targeted and genome-wide approaches, and between different cohorts of trained animals, it stands to reason that these genes may be of the most fundamental players for lasting changes to auditory function and memory and are now prime for future investigation.

Considered together, the findings reveal a dynamic transcriptional landscape in the adult auditory cortex that could support emergent auditory function in neurophysiology and behavior. Given the growing evidence of epigenetic controls on sensory system function (c.f., Shang & Bieszczad, 2022), it is exciting to consider how epigenetic mechanisms play a role in balancing stability with the experience-dependent plasticity of auditory cortical circuits. For example, the current genome-wide dataset also identified a learning-induced reduction in the expression of a repressive histone deacetylase, *Hdac9*, that was absent with HDAC-inhibition. It is tempting to liken the effect of reduced HDAC9 expression to the effect of pharmacologically inhibiting HDACs to promote learning-induced transcription. Another important class of epigenetic regulators alter DNA rather than histones. Examples are *Tet1* and *Gadd45b* that both impact DNA methylation (Bayraktar & Kreutz, 2018) and were found to be down- and up-regulated with learning, respectively, under HDAC-inhibition. Further, a class of microRNAs appeared as the top biological pathway induced by learning with HDAC-inhibition (see **Table 1**), which have gained traction in the field as key epigenetic regulators of neural transcriptional control (Saab & Mansuy, 2014). Higher-order molecular interactions between epigenetic players and the protein products of expressed genes are also likely. For example, the identified up-regulated *Egr1* can recruit *Tet1* to remove repressive methylation marks to activate downstream genes (Sun, et al., 2019). Further molecular studies will be needed to parse out the significance of interactions between epigenetic regulators in the auditory system during learning. Moreover, this report presents an immediate opportunity to establish links between select auditory cortical genes and their downstream effectors. Bridging the gap from these genes and molecules to their influences on sound-evoked neurophysiological events is essential to understand how the adult auditory system adapts with experience to alter behavior. Indeed, the translation of *de novo* transcripts are required for long-term changes to behavioral function because they produce lasting changes to cellular function. For example, key target effectors of learning-induced transcriptional processes may alter the availability of channels or receptors (Metherate, Intskirveli, & Kawai, 2012; Brown & Kaczmarek, 2011; Henton, Zhao, & Tzounopoulos, 2023) within circuits that determine sound-evoked threshold, responsivity, and receptive field architecture in the auditory system. It is reasonable to assume that different auditory tasks would recruit unique gene networks for biological pathways relevant to the particular cellular functions that would support learning for the task-relevant sound feature or task structure.

Overall, this report serves to act as a starting point to make RNAseq datasets from the learning, adult, auditory cortex available (see **Data Repository**). An effort to narrow the gap in knowledge exacerbated by the severe lack of studies that focus on molecular genetic processes in adult auditory cortex is now welcome. We encourage study beyond the spatio-temporal limitations of bulk RNA-sequencing whose sensitivity is limited technically also by read depth (Li & Wang, 2021), especially since modern *Omics* technologies are quickly evolving and improving. While research in the auditory cortex has made some initial strides in molecular genetics that have identified essential molecular cascades (Schicknick, et al., 2008), permissive IEG profiles (Mello, Velho, & Pinaud, 2006; de Hoz, et al., 2018; Peter, et al., 2011) and even chromatin dynamics (Peter, et al., 2021), there is precedent in the auditory periphery where transcription dynamics are beautifully studied (Kwan, 2016; Barta, et al., 2018; Li, et al., 2020; Ebeid, et al., 2017). Existing molecular tools like for single cell RNA sequencing (scRNAseq) and small-molecule fluorescent in situ hybridization (smFISH) will be useful to provide insights into

experience-dependent cell-to-cell variation and molecular interactions within auditory cortical cells and circuits. Unlike bulk RNAseq, these methods honor the profound cellular diversity and higher order organization in auditory cortex, such as its layer-specific microcircuitry and lemniscal topography. Clever approaches to tag and sequence transcripts from only recently active auditory cortical cells (Cho , Huang, & Gray, 2016) may increase sensitivity enough to robustly obtain the most functionally relevant sets of genes from the most functionally relevant cell types and populations. Similar approaches could be used also subcortically to capture and contrast activity-dependent regional transcriptomic profiles in the cortex that honors the incredible integration of cortical with subcortical function of the auditory system under different listening conditions or behavioral demands. Fully characterizing cellular and regional distinctions in genetic and molecular mechanisms underlying auditory learning in the adult brain is paramount for developing site-selective and molecule-targeted precision therapeutics that enable robust and persistent functional changes to support specific hearing and listening abilities across the lifespan.

Methods

Subjects: A total of 24 adult male Sprague-Dawley rats (250 – 300 g on arrival; Charles River Laboratories, Wilmington, MA) were used in behavioral and molecular experiments. All animals were individually housed in a temperature-controlled (24 ˚C) colony room on a 12-hour light/dark cycle. Subjects had ad libitum access to food and water prior to behavioral training. All procedures were approved and conducted according to guidelines by the Institutional Animal Care and Use Committee (IACUC) at Rutgers, the State University of New Jersey (Protocol No.: 999900026 (K.M.B)).

Behavioral apparatus and sound stimuli: All behavioral sessions were conducted in two identical instrumental conditioning chambers (H10-112TC-NSF; Coulbourn Instruments, Holliston, MA) within a sound-attenuated box. Daily training sessions were counterbalanced to ensure equal exposure to both chambers. Each chamber (12" W x 10" D x 12" H; wire mesh floor was fitted with a response lever (H21-03R), house light (H11-01R), a speaker (H12-01R), and a water delivery system (H14-05R). During training phases, animals could depress the response lever ("barpress"), which triggered the presentation of a water cup (~0.02cc) in the reward port (1.25" W x 1.625" H). A hand switch (H21-01) was used during early session to shape the animal's barpress response to trigger presentations of the water cup that allowed access to the water reward. Behavioral responses were recorded using Graphic State 4 software (Coulbourn Instruments, Holliston MA) for offline analysis.

All auditory stimuli were generated using Tucker-Davis Technologies (TDT, Alachua, FL) and RPvdsEx software, and presented via the operant chamber's wall-mounted speaker. White noise (during procedural training; **Fig. 1a**) was presented for 7 or 9 s in duration (75 dB SPL). Pure tones (during two-tone discrimination training; **Fig. 1a**) were always presented for 8 s (70 dB SPL). Sound levels were calibrated daily using a digital sound meter (Larson Davis SoundTrack LxT1).

Behavioral training and pharmacological inhibition of HDAC3: After one day of acclimating to the vivarium, rats were handled daily for a minimum of 3 days. Prior to the start of behavioral

training, rats were placed on a schedule of restricted water until they reached 85% of the non-restricted weight of age-matched control animals. Water-restricted rats were then shaped inside sound-attenuated chambers to barpress for water rewards (**Fig. 1a**). Barpress shaping and subsequent sound training were as previously described (Shang, Bylipudi, & Bieszczad, 2019). Briefly, all animals were shaped to barpress over 5 days, and then trained in a procedural task to learn to bar-press to sound, which in this phase was a broad-band Gaussian noise stimulus (1-12.5 kHz band-pass filtered white noise; 75 dB SPL), to obtain a water reward. All animals learned successfully learned to associate this sound with reward prior to continuing to the next phase of training. Procedural training showed individual variability of how quickly the animals could learn the task to high levels of performance. Nonetheless, all animals were able to achieve a performance level of ≥90% for two consecutive days (mean = 92.85%, s.e.m. = 0.01%) after an average of 12.67 days (s.d. = 2.85 days). Performance was calculated using $Performance = \frac{\#BPs\ to\ white\ noise}{Total\ \#BPs} \times 100\%$.

The next phase of training was the two-tone discrimination (2TD) task (**Fig. 1a**), in which rats were trained to discriminate between two spectrally distinct sound frequencies. Barpresses to the S+ tone (5.0 kHz; 70 dB SPL) would result in the presentation of a water reward, while barpresses to the S- tone (11.5 kHz; 70 dB SPL) resulted in an error signal (flashing house light) and a "time-out" (an additional 6 s wait for the start of the next trial). S+ and S- trials were randomized and lasted for 8 s. Inter-trial intervals (ITIs) were on average 15 s (range: 5-25 s, randomized). Barpresses during a silent ITI were inconsequential (no time-out, no error signal, nor water reward). Daily sessions were 45 minutes in length. All animals performed the 2TD task for a total of three consecutive days. Rats were paired by a trained observer so that animals with similar rates of acquiring the procedural task were assigned different treatment conditions in the 2TD phase. Performance-matched pairs were received systemic injections of either the pharmacological class I HDAC3 inhibitor RGFP966 (Abcam Inc., ab144819; 10 mg/kg; s.c.; N = 12) or vehicle solution (N = 9) immediately after each 2TD session. Performance of the 2TD task was calculated as previously described: $Performance = \frac{BP^{S+}}{BP^{S+} + BP^{S-}} \times 100\%$ (Shang, Bylipudi, & Bieszczad, 2019).

Tissue collection and isolation of RNA: One hour after animals received their third and final injection of RGFP966 after the third session of 2TD, brains were quickly dissected and flash-frozen in a beaker of 2-methylbutane placed upon dry ice. Flash-frozen brains were then stored at -80° C until future processing. To prepare brains for cryosectioning, brains were encased in optimal cutting temperature compound (OCT) and stored at -20° C for 12-24 hours to ensure brain tissue would reach -20° C for cutting. Brains encased in OCT would then be sliced horizontally in a cryostat (Leica CM 3050S) at a thickness of 250 µm. Using a 1 mm round tissue micropunch, 2 mm³ of auditory cortical tissue from each hemisphere (combined into one sample for a total of 4 mm³ per brain region) was sampled for RNA extraction. Auditory cortex location was identified using Paxinos and Watson's *The Rat Brain in Stereotaxic Coordinates* (6th edition) as a reference (approximately A/P = -3.6 to -5.8 mm, M/L = ±6.4 mm D/V = -4.2 to -5.8 mm; and using hippocampus as landmark).

Total RNA from each sample was isolated using the PureLink™ RNA Mini Kit (ThermoFisher) using the manufacturer's protocol. RNA samples were then purified with the RNA Clean and Concentrator™ -25 Kit (Zymo Research).

Gene expression analysis, Gene Ontology (GO) biological process analysis, and gene set enrichment analysis (GSEA): RNA-seq analysis was conducted by the Iowa Institute of Human Genetics (IIHG; Iowa City, IA, USA). Briefly, using 500 ng total RNA (all RIN values >8), sequencing libraries were generated using the Illumina TruSeq® Stranded mRNA Library Prep kit according to the manufacturer's recommended protocols. Libraries were pooled and sequencing was performed on an Illumina NovaSeq 6000 running 100 bp paired-end SBS chemistry. Reads were processed with the 'bcbio-nextgen.py' open-source informatics pipeline developed primarily at Harvard Chan Bioinformatics (v.1.2.4) running on the Argon HPC resource at the University of Iowa. This pipeline includes 'best practices' approaches for read quality control, read alignment and quantitation. The 'bcbio-nextgen.py' pipeline was run in "RNA-seq" mode with the 'rn6' key as the selected genome build (internally referencing Ensembl assembly and genebuild 'Rnor_6.0'). The pipeline aligned reads to the Rnor_6.0 genome using the splice-aware, ultra-rapid hisat2 aligner (2.2.1) and concurrently quantified reads to the transcriptome using the 'salmon' (1.4.0) aligner. Qualimap (2.2.2), a computational tool that examines hisat2 BAM alignment files, was used to examine the read data for quality control. Sequence quality scores passed basic checks and sequence duplication rates were within acceptable parameters. All samples passed QC for read alignment to exonic regions. Salmon-derived transcript quantifications (TPM or "transcripts per million") were imported and summarized to estimated counts at the gene level using tximport (1.12.3) in Rstudio, as described in the best-practices DESeq2 vignette (https://bioconductor.org/packages/release/bioc/vignettes/DESeq2/inst/doc/DESeq2.html).
Genes with fewer than 5 estimated counts across all samples were pre-filtered from downstream analysis, as per recommended procedure. Differential gene expression analysis was conducted with DESeq2 (v.1.24.0) on estimated gene-level counts. An FDR of 5% and X < abs(logFC) < 10 was set as a cutoff for differential expression (DEGs). Heatmaps, line graphs, and volcano plots were generated using clusterProfiler packages in R/Bioconductor. Gene level, pathway, and DEG analyses were generated using iPathwayGuide (Advaita Bioinformatics, https://www.advaitabio.com/ipathwayguide; last accessed November 15, 2022). iPathwayGuide scores pathways using the Impact Analysis method (Draghici et al., 2007); Tarca et al., 2009, Khatri et al., 2007). The underlying pathway topologies, comprised of genes and their directional interactions, are obtained from the KEGG database (Kanehisa et al., 2000; Kanehisa et al., 2010; Kanehisa et al., 2012; Kanehisa et al., 2014).

RNA-seq data validation with qRT-PCR: RNAseq data were validated by qRT-PCR on mRNA extracted from distinct cohorts of rats. Five transcripts were chosen from a list of novel DEGs found enriched in each region at baseline. Three of the genes were hypothesis-driven based on auditory cortical physiological and behavioral literature, while another three genes were novel genes identified from gene level and DEG analyses in a comparison between Drug and Vehicle groups. All primer sequences were selected either from previous literature on brain tissue in rats (sequences provided in table below) or designed using NCBI Primer Blast and validated for target specificity by assessing melt-curves and PCR amplicon product sequencing. qRT-PCR

analyses were performed in a QuantStudio 3 (Applied Biosystems) with SsoAdvanced[TM] Universal SYBR Green Supermix (Bio-Rad). For each sample, 500 ng of cDNA was amplified using an iScript[TM] cDNA Synthesis Kit (Bio-Rad).

Gene	Forward primer	Reverse primer	Reference
Gapdh	GTGGACCTCATGGCCTACAT	TGTGAGGGAGATGCTCAGTG	Malvaez et al., 2018
18S	CGGACAGGATTGACAGATTG	CAAATCGCTCCACCAACTAA	Chang et al., 2016
Egr1	AACCCTACGAGCACCTGAC	CGGGTAGTTTGGCTGGGATA	Malvaez et al., 2018
Nr4a1	CCGGTGACGTGCAGCAATTTTATGAC	GGCTAGAATGTTGTCTATCCAGTC	Malvaez et al., 2018
Nr4a2	ATTGCTGCCCTGGCTATGGT	GACCATCCCATTATTGAAAGTCACATGGTC	Malvaez et al., 2018
Chrna7	TGCACGTGTCCCTGCAAGGC	GTACACGGTGAGCGGCTGCG	Bychkov et al., 2018
Lynx1	ACCACTCGAACTTACTTCACC	ATCGTACACGGTCTCAAAGC	Bychkov et al., 2018
Htr1a	GACCACGGCTACACCATCTAC	CTGTCCGTTCAGGCTCTTCTT	Qiu et al., 2016
Per2	TCCGAGTATATCGTGAAGAACG	CAGGATCTTCCCAGAAACCA	Kwapis et al., 2018
Adamts13	AAACACTGCCTGATGTCCCG	GCACGGCAACCATAAGCAAA	custom-design

Acknowledgements

The authors would like to thank Dr. Andrea Shang, Sean Tsaur, and Sooraz Bylipudi for their assistance in behavioral procedures and analysis; Dr. Mimi L. Phan for her assistance with RNA extraction and sample collection protocols; Dr. Troy A. Roepke, Ali Yasrebi, and Christopher O'Brien for their assistance with RNA sample purification analysis; and Alisa Ray for technical assistance. We would like to thank all current and former CLEF Lab personnel for their assistance and support.

This work was supported by the National Institutes of Health, National Institute of Deafness and Communication Disorders [R01-DC014753 to K.M.B.]; the School of Arts and Sciences at Rutgers University; the Aresty Foundation at Rutgers University with small grant funding for undergraduate research; and the Project SUPER program at Rutgers University with small grant funding for undergraduate research.

Data Repository:

Raw sequencing files are publicly available here: rutgers.box.com/s/x1oi6d3vh7q46niumyyvh3qs5c1zbqbe

References:

Alberini, C. M., & Kandel, E. R. (2014). The Regulation of Transcription in Memory Consolidation. *Cold Spring Harbor Perspectives in Biology, 7*(1), a021741. doi:https://doi.org/10.1101/cshperspect.a021741

Albrecht, U., Zheng, B., Larkin, D., Sun, Z., & Lee, C. (2001). MPer1 and mper2 are essential for normal resetting of the circadian clock. *J Biol Rhythms, 16*, 100-104. doi:10.1177/074873001129001791.

Aschauer, D., & Rumpel, S. (2016). The Sensory Neocortex and Associative Memory. *Behavioral Neuroscience of Learning and Memory, 37*, 177-211. doi:https://doi.org/10.1007/7854_2016_453

Bae, K., Jin, X., Maywood, E. S., Hastings, M. H., Reppert, S. M., & Weaver, D. R. (2001). Differential functions of mPer1, mPer2, and mPer3 in the SCN circadian clock. *Neuron, 30*, 525-536. doi:10.1016/S0896-6273(01)00302-6.

Bakin, J. S., & Weinberger, N. M. (1996). Induction of a physiological memory in the cerebral cortex by stimulation of the nucleus basalis. *Proceedings of the National Acadmey of Sciences of the United States of America, 93*(20), 11219-11224. doi:https://doi.org/10.1073/pnas.93.20.11219

Banerjee, S. B., Gutzeit, V. A., Baman, J., Aoued, H. S., Doshi, N. K., Liu, R. C., & Ressler, K. J. (2017). Perineuronal nets in the adult sensory cortex are necessary for fear learning. *Neuron, 95*(1), 169-179. doi:10.1016/j.neuron.2017.06.007

Barta, C. L., Liu, H., Chen, L., Giffen, K. P., Li, Y., Kramer, K. L., . . . He, D. Z. (2018). RNA-seq transcriptomic analysis of adult zebrafish inner ear hair cells. *Scientific Data, 5*, 180005. doi:https://doi.org/10.1038/sdata.2018.5

Bayraktar, G., & Kreutz, M. R. (2018). The Role of Activity-Dependent DNA Demethylation in the Adult Brain and in Neurological Disorders. *Frontiers in Molecular Neuroscience, 11*, 169. doi:10.3389/fnmol.2018.00169

Bieszczad, K. M., Bechay, K., Rusche, J. R., Jacques, V., Kudugunti, S., Miao, W., . . . Wood, M. A. (2015). Histone Deacetylase Inhibition via RGFP966 Releases the Brakes on Sensory Cortical Plasticity and the Specificity of Memory Formation. *J Neurosci, 35*(38), 13124-13132. doi:https://doi.org/10.1523/JNEUROSCI.0914-15.2015

Brown, M. R., & Kaczmarek, L. K. (2011). Potassium channel modulation and auditory processing. *Hearing Research, 279*(1-2), 32-42. doi:https://doi.org/10.1016/j.heares.2011.03.004

Bychkov, M. L., Vasilyeva, N. A., Shulepko, M. A., Balaban, P. M., Kirpichnikov, M. P., & Lyukmanova, E. N. (2018). Lynx1 Prevents Long-Term Potentiation Blockade and Reduction of Neuromodulator Expression Caused by Aβ1-42 and JNK Activation. *Acta Naturae, 10*(3), 57-61.

Chang , R., Chou, M., Hung, L., Wang, M., Hsu, M., & Chiu, C. (2016). Study of Valproic Acid-Enhanced Hepatocyte Steatosis. *BioMed Research International, 2016*, 9576503. doi:https://doi.org/10.1155/2016/9576503

Cho , J., Huang, B. S., & Gray, J. M. (2016). RNA sequencing from neural ensembles activated during fear conditioning in the mouse temporal association cortex. *Scientific Reports, 6*, 31753. doi:https://doi.org/10.1038/srep31753

Concina, G., Renna, A., Grosso, A., & Sacchetti, B. (2019). The auditory cortex and the emotional valence of sounds. *Neuroscience & Biobehavioral Reviews, 98*, 256-264. doi:https://doi.org/10.1016/j.neubiorev.2019.01.018

Costa-Mattioli, M., Sossin, W. S., Klann, E., & Sonenberg, N. (2009). Translational Control of Long-Lasting Synaptic Plasticity and Memory. *Neuron, 16*(1), 10-26. doi:https://doi.org/10.1016/j.neuron.2008.10.055

Cuesta, M., Clesse, D., Pevet, P., & Challet, E. (2009). New light on the serotonergic paradox in the rat circadian system. *J Neurochem, 110*, 231-243. doi:10.1111/j.1471-4159.2009.06128.x.

de Hoz, L., Gierej, D., Lioudyno, V., Jaworski, J., Blazejczyk, M., Cruces-Solis, H., . . . Kaczmarek, L. (2018). Blocking c-Fos Expression Reveals the Role of Auditory Cortex Plasticity in Sound Frequency Discrimination Learning. *Cerebral Cortex, 28*(5), 1645-1655. doi:https://doi.org/10.1093/cercor/bhx060

Draghici, S., Khatri, P., Tarca, A. L., Amin, K., Done, A., Voichita, C., . . . Romero, R. (2007). A systems biology approach for pathway levelanalysis. *Genome Research, 17*(10), 1537-1545.

Duclot, F., & Kabbaj, M. (2017). The Role of Early Growth Response 1 (EGR1) in Brain Plasticity and Neuropsychiatric Disorders. *Frontiers in Behavioral Neuroscience, 11*, 1-20. doi:https://doi.org/10.3389/fnbeh.2017.00035

Dudai, Y. (2012). The Restless Engram: Consolidations Never End. *Annual Reveiw of Neuroscience, 35*, 227-247. doi:https://doi.org/10.1146/annurev-neuro-062111-150500

Ebeid, M., Sripal, P., Pecka, J., Beisel, K. W., Kwan, K. Y., & Soukup, G. A. (2017). Transcriptome-wide comparison of the impact of Atoh1 and miR-183 family on pluripotent stem cells and multipotent otic progenitor cells. *Plos One, 12*(7), e0180855. doi:https://doi.org/10.1371/journal.pone.0180855

Elhilali, M., Fritz, J. B., Chi, T.-S., & Shamma, S. A. (2007). Auditory Cortical Receptive Fields: Stable Entities with Plastic Abilities. *Journal of Neuroscience, 27*(39), 10372-10382. doi:https://doi.org/10.1523/JNEUROSCI.1462-07.2007

El-Tabbal, M., Niekisch, H., Henschke, J. U., Budinger, E., Frischknecht, R., Deliano, M., & Happel, M. F. (2021). The extracellular matrix regulates cortical layer dynamics and cross-columnar frequency integration in the auditory cortex. *Communications Biology, 4*, 322. doi:https://doi.org/10.1038/s42003-021-01837-4

Fritz, J. B., Elhilali, M., David, S. V., & Shamma, S. A. (2007). Does attention play a role in dynamic receptive field adaptation to changing acoustic salience in A1? *Hearing Research, 229*(1-2), 186-203. doi:https://doi.org/10.1016/j.heares.2007.01.009

Froemke, R. C., & Martins, A. R. (2011). Spectrotemporal dynamics of auditory cortical synaptic receptive field plasticity. *Hearing Research, 279*(1-2), 149-161. doi:https://doi.org/10.1016/j.heares.2011.03.005

Froemke, R. C., & Schreiner, C. E. (2015). Synaptic plasticity as a cortical coding scheme. *Current opinion in neurobiology, 35*, 185-199. doi:https://doi.org/10.1016/j.conb.2015.10.003

Gervain, J., Vines, B. W., Chen, L. M., Seo, R. J., Hensch, T. K., Werker, J. F., & Young, A. H. (2013). Valproate reopens critical-period learning of absolute pitch. *Frontiers in system neuroscience, 7*, 102. doi:https://doi.org/10.3389/fnsys.2013.00102

Ghosh, S., & Zador, A. M. (2021). Corticostriatal Plasticity Established by Initial Learning Persists after Behavioral Reversal. *eNeuro, 8*(2), ENEURO.0209-20.2021. doi:https://doi.org/10.1523/ENEURO.0209-20.2021

Grosso, A., Cambiaghi, M., Concina, G., Sacco, T., & Sacchetti, B. (2015). Auditory cortex involvement in emotional learning and memory. *Neuroscience, 299*, 45-55. doi:https://doi.org/10.1016/j.neuroscience.2015.04.068

Happel, M. F., Niekisch, H., Rivera, L. L., Ohl, F. W., Deliano, M., & Frischknecht, R. (2014). Enhanced cognitive flexibility in reversal learning induced by removal of the extracellular matrix in auditory cortex. *PNAS, 111*(7), 2800-2805. doi:https://doi.org/10.1073/pnas.1310272111

Henton, A., Zhao, Y., & Tzounopoulos, T. (2023). A role for KCNQ channels on cell-type-specific plasticity in mouse auditory cortex after peripheral damage. *J Neurosci*, JN-RM-1070-22. doi:https://doi.org/10.1523/JNEUROSCI.1070-22.2023

Hu, G., Wang, Z., Zhang, R., Sun, W., & Chen, X. (2021). The Role of Apelin/Apelin Receptor in Energy Metabolism and Water Homeostasis: A Comprehensive Narrative Review. *Frontiers in Physiology, 12*, 632886. doi:10.3389/fphys.2021.632886

Ji, W., Gao, E., & Suga, N. (2001). Effects of acetylcholine and atropine on plasticity of central auditory neurons caused by conditioning in bats. *J. Neurophysiol., 86*, 211-225. doi:https://doi.org/10.1152/jn.2001.86.1.211

Kalish, B. T., Barkat, T. R., Diel, E. E., Zhang, E. J., Greenberg, M. E., & Hensch, T. K. (2020). Single-nuecleus RNA sequencing of mouse auditory cortex reveals critical periods triggers and brakes. *Proc Natl Acad Sci U S A, 117*(21), 11744-11752. doi:10.1073/pnas.1920433117

Kanehisa, M., & Goto, S. (2000). KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic Acids Research, 28*, 27-30.

Kanehisa, M., Goto, S., Furumichi, M., Tanabe, M., & Hirakawa, M. (2010). KEGG for representation and analysis of molecular networks involving diseasesand drugs. *Nucleic Acids Research, 38*, D355-D360.

Kanehisa, M., Goto, S., Sato, Y., Furumichi, M., & Tanabe, M. (2012). KEGG for integration and interpretation of large-scale molecular datasets. *Nucleic Acids Research, 40*, D109-D114.

Kanehisa, M., Goto, S., Sato, Y., Kawahima, M., Furumichi, M., & Tanabe, M. (2014). Data, information, knowledge and principle: back to metabolism inKEGG. *Nucleic Acids Research, 42*, D199-D205.

Katzman, A., Khodadadi-Jamayran, A., Kapeller-Libermann, D., Ye, X., Tsirigos, A., Heguy, A., & Alberini, C. M. (2021). Distinct Transcriptomic Profiles in the Dorsal Hippocampus and Prelimbic Cortex Are Transiently Regulated following Episodic Learning. *J Neurosci*, 41(12):2601-2614.

Khatri, P., Draghici, S., Tarca, A. D., Hassan, S. S., & Romero, R. (2007). A system biology approach for the steady-state analysis of gene signaling networks. *Lecture Notes in Computer Science (LNCS), 4756*, 32-41.

Kilgard, M. P., & Merzenich, M. M. (1998). Cortical map reorganization enabled by nucleus basalis activity. . *Science, 279*(5357), 1714-1718. doi:https://doi.org/10.1126/science.279.5357.1714

Kuchibhotla, K. V., Gill, J. V., Lindsay, G. W., Papadoyannis, E. S., Field, R. E., Hindmarsh Sten, T. A., . . . Froemke, R. C. (2017). Parallel processing by cortical inhibition enables context-dependent behavior. *Nat Neurosci, 20*(1), 62-71. doi:10.1038/nn.4436

Kwan, K. Y. (2016). Single-Cell Transcriptome Analysis of Developing and Regenerating Spiral Ganglion Neurons. *Current Pharmacological Report, 2*, 211-220. doi:10.1007/s40495-016-0064-z

Kwapis, J. L., Alaghband, Y., Kramár, E. A., López, A. J., Ciernia, A. V., White, A. O., . . . Wood, M. A. (2018). Epigenetic regulation of the circadian gene Per1 contributes to age-related changes in hippocampal memory. *Nature Communications, 9*, 3323. doi:https://doi.org/10.1038/s41467-018-05868-0

Kwapis, J. L., Alaghband, Y., López, A. J., Long, J. M., Li, X., Shu, G., . . . Wood, M. A. (2019). HDAC3-Mediated Repression of the Nr4a Family Contributes to Age-Related Impairments in Long-Term Memory. *J Neurosci, 39*(25), 4999-5009. doi:10.1523/JNEUROSCI.2799-18.2019

Kwapis, J. L., Alaghband, Y., Lopez, A. J., White, A. O., Campbell, R. R., Dang, R. T., . . . Wood, M. A. (2017). Context and Auditory Fear are Differentially Regulated by HDAC3 Activity in the Lateral and Basal Subnuclei of the Amygdala. *Neuropsychopharmacology, 42*(6), 1284-1294. doi:https://doi.org/10.1038/npp.2016.274

Lechner, H. A., Squire, L. R., & Byrne, J. H. (1999). 100 years of consolidation--remembering Müller and Pilzecker. *Learning & memory (Cold Spring Harbor, N.Y.), 6*(2), 77-87.

Lesicko, A. M., & Geffen, M. N. (2022). Diverse functions of the auditory cortico-collicular pathway. *Hearing Research, 425*, 108488. doi:https://doi.org/10.1016/j.heares.2022.108488

Letzkus, J. J., Wolff, S. B., & Lüthi, A. (2015). Disinhibition, a Circuit Mechanism for Associative Learning and Memory. *Neuron, 88*(2), 264-276. doi:https://doi.org/10.1016/j.neuron.2015.09.024

Li, C., Li, X., Bi, Z., Sugino, K., Wang, G., Zhu, T., & Liu, Z. (2020). Comprehensive transcriptome analysis of cochlear spiral ganglion neurons at multiple ages. *eLife, 9*, e50491. doi:https://doi.org/10.7554/eLife.50491

Li, X., & Wang, C. Y. (2021). From bulk, single-cell to spatial RNA sequencing. *Int J Oral Sci, 13*, 36. doi:https://doi.org/10.1038/s41368-021-00146-0

Liu, X., Basvaraj, S., Krishnan, R., & Yan, J. (2011). Contributions of the thalamocortical system towards sound-specific auditory plasticity. *Neuroscience & Biobehavioral Reviews, 35*(10), 2155-2161. doi:https://doi.org/10.1016/j.neubiorev.2011.02.010

Luo, F., & Yan, J. (2013). Sound-specific plasticity in the primary auditory cortex as induced by the cholinergic pedunculopontine tegmental nucleus. *Eur J Neurosci, 37*(3), 393-399. doi:doi.10.1111/ejn.12046

Malvaez, M., Greenfield, V. Y., Matheso, D. P., Angelillis, N. A., Murphy, M. D., Kennedy, P. J., . . . Wassum, K. M. (2018). Habits Are Negatively Regulated by Histone Deacetylase 3 in the Dorsal Striatum. *Biol Psychiatry, 84*(5), 383-392. doi:https://doi.org/10.1016/j.biopsych.2018.01.025

Malvaez, M., McQuown, S. C., Rogge, G. A., Astarabadi, M., Jacques, V., Carreiro, S., . . . Wood, M. A. (2012). HDAC3-selective inhibitor enhances extinction of cocaine-seeking behavior in a persistent manner. *PNAS, 110*(7), 2647-2652. doi:https://doi.org/10.1073/pnas.1213364110

McGaugh, J. L. (2000). Memory -- a century of consolidation. *Science, 287*(5451), 248-251. doi:10.1126/science.287.5451.248

McNulty, S. E., Barrett, R. A., Vogel-Ciernia, A., Malvaez, M., Hernandez, N., Davatolhagh, M. F., . . . Wood, M. A. (2012). Differential roles for Nr4a1 and Nr4a2 in object location vs. object recognition long-term memory. *Learn Mem, 19*(12), 588-592. doi:10.1101/lm.026385.112.

McQuown, S. C., & Wood, M. A. (2011). HDAC3 and the molecular brake pad hypothesis. *Neurobiol Learn Mem.*, 96(1):27-34.

McQuown, S. C., Barrett, R. M., Matheos, D. P., Post, R. J., Rogge, G. A., Alenghat, T., . . . Wood, M. A. (2011). HDAC3 Is a Critical Negative Regulator of Long-Term Memory Formation. *The Journal of Neuroscience, 31*(2), 764-774. doi:https://doi.org/10.1523/JNEUROSCI.5052-10.2011

Mello, C. V., Velho, T. A., & Pinaud, R. (2006). Song-Induced Gene Expression: A Window on Song Auditory Processing and Perception. *Annals of the New York Academy of Sciences, 1016*(1), 263-281. doi:https://doi.org/10.1196/annals.1298.021

Metherate, R. (2011). Functional connectivity and cholinergic modulation in audiory cortex. *Neurosci Biobehav Rev., 35*(10), 2058-2063. doi:10.1016/j.neubiorev.2010.11.010

Metherate, R., Intskirveli, I., & Kawai, H. D. (2012). Nicotinic filtering of sensory processing in auditory cortex. *Frontiers in Behavioral Neuroscience*, 44. doi:https://doi.org/10.3389/fnbeh.2012.00044

Morishita, H., Miwa, J. M., Heintz, N., & Hensch, T. K. (2010). Lynx1, a Cholinergic Brake, Limits Plasticity in Adult Visual Cortex. *Science (New York, N.Y.), 330*(6008), 1238-1240. doi:10.1126/science.1195320

Nadel, L., & Hardt, O. (2011). Update on Memory Systems and Processes. *Neuropsychopharmacol, 36*, 251-273. doi:https://doi.org/10.1038/npp.2010.169

Paxinos, G., & Watson, C. (2007). *The Rat Brain in Stereotaxic Coordinates.* London: Academic Press.

Peter, M., Aschauser, D. F., Rose, R., Sinning, A., Grössl, F., Kargl, D., . . . Rumpel, S. (2021). Rapid nucleus-scale reorganization of chromatin in neurons enables transcriptional adaptation for memory consolidation. *Plos One, 16*(5), e0244038. doi:https://doi.org/10.1371/journal.pone.0244038

Peter, M., Scheuch, H., Burkard, T. R., Tinter, J., Wernle, T., & Rumpel, S. (2011). Induction of immediate early genes in the mouse auditory cortex after auditory cued fear conditioning to complex sounds. *Genes, Brain and Behavior, 11*(3), 314-324. doi: https://doi.org/10.1111/j.1601-183X.2011.00761.x

Phan, M. L., Gergues, M. M., Mahidadia, S., Jimenez-Castillo, J., Vicario, D. S., & Bieszczad, K. M. (2017). HDAC3 Inhibitor RGFP966 Modulates Neuronal Memory for Vocal Communication Signals in a Songbird Model. *Frontiers in Systems Neuroscience, 11*, 65. doi:https://doi.org/10.3389/fnsys.2017.00065

Pienkowski, M., & Eggermont, J. J. (2011). Cortical tonotopic map plasticity and behavior. *Neuroscience & Biobehavioral Reviews, 35*(10), 2117-2128. doi:https://doi.org/10.1016/j.neubiorev.2011.02.002

Qiu , Y., Chen, D., Huang, X., Huang, L., Tang, L., Jiang, J., . . . Li, S. (2016). Neuroprotective effects of HTR1A antagonist WAY-100635 on scopolamine-induced delirium in rats and underlying molecular mechanisms. *BMC Neuroscience, 17*, 66. doi:10.1186/s12868-016-0300-9

Reh, R. K., Dias, B. G., Nelson , C. A., Werker, J. F., Kolb, B., Levine, J. D., & Hensch, T. K. (2020). Critical period regulation across multiple timescales. *PNAS, 117*(38), 23242-23251. doi:https://doi.org/10.1073/pnas.1820836117

Reh, R. K., Dias, B. G., Nelson III, C. A., Kaufer, D., Werker, J. F., Kolb, B., . . . Hensch, T. K. (2019). Critical period regulation across multiple timescales. *PNAS, 117*(38), 23242-23251. doi:https://doi.org/10.1073/pnas.1820836117

Rotondo, E. K., & Bieszczad, K. M. (2020). Precise memory for pure tones is predicted by measures of learning-induced sensory system neurophysiological plasticity at cortical and subcortical levels. *Learning & memory (Cold Spring Harbor, N.Y.), 27*(8), 328-339. doi:https://doi.org/10.1101/lm.051318.119

Rotondo, E. K., & Bieszczad, K. M. (2021a). Sensory cortical and subcortical auditory neurophysiological changes predict cue-specific extinction behavior enabled by the pharmacological inhibition of an epigenetic regulator during memory formation. *Brain Research Bulletin, 169*, 167-183. doi:10.1016/j.brainresbull.2021.01.016

Rotondo, E. K., & Bieszczad, K. M. (2021b). Memory specific to temporal features of sound is formed by cue-selective enhancements in temporal coding enabled by inhibition of an epigenetic regulator. *The Journal of Neuroscience, 41*(44), 9192-9209. doi:https://doi.org/10.1523/JNEUROSCI.0691-21.2021

Saab, B. J., & Mansuy, I. M. (2014). Neuroepigenetics of memory formation and impairment: the role of microRNAs. *Neuropharmacology, 80*, 61-69. doi:https://doi.org/10.1016/j.neuropharm.2014.01.026

Schicknick, H., Schott, B. H., Budinger, E., Smalla, K. H., Riedel, A., Seidenbecher, C. I., . . . Tischmeyer, W. (2008). Dopaminergic modulation of auditory cortex-dependent memory consolidation through mTOR. *Cerebral Cortex, 18*(11), 2646-2658. doi:https://doi.org/10.1093/cercor/bhn026

Schreiner, C. E., & Polley, D. B. (2014). Auditory map plasticity: diversity in causes and consequences. *Current Opinion in Neurobiology, 24*, 143-156. doi:https://doi.org/10.1016/j.conb.2013.11.009

Schreiner, C. E., & Winer, J. A. (2007). Auditory Cortex Mapmaking: Principles, Projections, and Plasticity. *NEuron, 56*(2), 356-365. doi:https://doi.org/10.1016/j.neuron.2007.10.013

Shang, A., Bylipudi, S., & Bieszczad, K. M. (2019). Inhibition of histone deacetylase 3 via RGFP966 facilitates cortical plasticityunderlying unusually accurate auditory associative cue memory forexcitatory and inhibitory cue-reward associations. *Behavioural Brain Research*, 453-469.

Sonntag, M., Blosa, M., Schmidt, S., Rübsamen, R., & Morawski, M. (2015). Perineuronal nets in the auditory system. *Hearing Research, 329*, 21-32. doi:https://doi.org/10.1016/j.heares.2014.12.012

Souffi, S., Nodal, F. R., Bajo, V. M., & Edeline, J. M. (2021). When and How Does the Auditory Cortex Influence Subcortical Auditory Structures? New Insights About the Roles of Descending Cortical Projections. *Frontiers in Neuroscience, 15*, 690223. doi:https://doi.org/10.3389/fnins.2021.690223

Sun, Z., Xu , X., He, J., Murray, A., Sun, M.-a., Wei, X., . . . Xie, H. (2019). EGR1 recruits TET1 to shape the brain methylome during development and upon neuronal activity. *Nature Communications*, 10(1), 3892.

Takesian, A. E., Bogart, L. J., Lichtman, J. W., & Hensch, T. K. (2018). Inhibitory circuit gating of auditory critical period plasticity. *Nat Neurosci, 21*(2), 218-227. doi:10.1038/s41593-017-0064-2

Talwar, S. K., & Gerstein, G. L. (1998). Auditory frequency discrimination in the white rat. *Hearing Research, 126*(1-2), 135-150. doi:https://doi.org/10.1016/s0378-5955(98)00162-2

Tchernichovski, O., & Margoliash, D. (2013). Time Scales of Vocal Learning in Songbirds. In *Animal Models of Speech and Language Disorders* (pp. 43-60). Springer, New York, NY. doi:https://doi.org/10.1007/978-1-4614-8400-4_3

Weinberger, N. M. (2003). The nucleus basalis and memory codes: auditory cortical plasticity and the induction of specific, associative behavioral memory. *Neurobiology of learning and memory, 80*(3), 268-284. doi:https://doi.org/10.1016/s1074-7427(03)00072-8

Weinberger, N. M. (2007a). Auditory associative memory and representational platsicity in the primary auditory cortex. *Hearing Research, 229*(1-2), 54-68. doi:https://doi.org/10.1016/j.heares.2007.01.004

Weinberger, N. M. (2007b). Associative representational plasticiy in the auditory cortex: a synthesis of two disciplines. *Learning & Memory (Cold Spring Harbor, N.Y.), 14*(1-2), 1-16. doi:https://doi.org/10.1101/lm.421807

Weinberger, N. M. (2015). New perspectives on the auditory cortex: learning and memory. . *Handbook of clinical neurology, 129*, 117-147. doi:https://doi.org/10.1016/B978-0-444-62630-1.00007-X

Xiong, Q., Znamenskiy, P., & Zador, A. M. (2015). Selective corticostriatal plasticity during acquisition of an auditory discrimination task. *Nature, 521*, 348-351. doi:https://doi.org/10.1038/nature14225

Name	Learning-induced			Learning-induced with RGFP966		
	Rank	Genes (DE/ALL)	PV	Rank	Genes (DE/ALL)	PV
Cholinergic synapse	1	**22/101**	**0.004**	1	**26/143**	**0.011**
Oxytocin signaling pathway	2	**24/133**	**0.014**	2	**41/275**	**0.012**
Morphine addiction	3	**18/82**	**0.025**	3	**30/194**	**0.016**
Parathyroid hormone synthesis, secretion, and action	4	**18/86**	**0.032**	4	**14/69**	**0.040**
PI3K-Akt signaling pathway	--	37/275	0.127	--	22/133	0.085
MicroRNAs in cancer	--	23/143	0.133	--	17/86	0.182
ECM-receptor interaction	--	13/69	0.465	--	16/101	1.000
Neuroactive ligand-receptor interaction	--	25/194	0.776	--	13/82	1.000

Figure Legends

Figure 1. *Behavioral timeline for tissue collection and use in RNA-sequencing.* **(a) Shaping:** Rats were water restricted and trained to bar press for a water reward in single daily sessions over 5 days. **Procedural Training:** To put bar-pressing behavior under the control of sound, all rats were trained in daily 45-minute sessions to bar-press to broad-band noise for water rewards to a performance criterion of 80% correct, which all subjects achieved over 1-2 weeks with individual differences in the rate of acquisition. Thus, pairs of rats that similarly acquired the procedural aspects of the task were assigned to different treatment groups in the next phase of training (with the selective HDAC inhibitor, RGFP966, vs. Vehicle) so that both groups were matched in prior auditory experience and basic learning ability (as in Shang et al., 2019). **Two-tone discrimination (2TD):** Rats were trained in single daily 45-minute 2TD sessions, each immediately followed by injections of the HDAC-inhibitor, RGFP966, or Vehicle for 3 consecutive days. 2TD required rats to bar-press to a rewarded cue (5.0 kHz; 65 dB SPL) for water reward, and to avoid responding to the unrewarded cue (11.5 kHz; 65 dB SPL) (see inset). **Brain collection:** One hour after the last injection of RGFP966/Vehicle on the third day of 2TD training, rats were sacrificed for brain collection. Whole brains were rapidly removed and immediately flash frozen for storage at -80° C. To prepare for cryosectioning, brains were encased in optimal cutting temperature compound (OCT) and then stored at -20° C for 12-24 hours. Brains were sliced horizontally in a cryostat (Leica CM 3050S) at a thickness of 250 μm in four adjacent slices targeting auditory cortex (see inset) to collect 2 mm^3 of cortical tissue from each hemisphere using a 1 mm round tissue micropunch (Harris). Thus, a total of 4 mm^3 per brain region per subject was used to produce a single sample for subsequent RNA extraction and sequencing. **(b)** Volcano plot showing auditory cortical learning-induced gene levels. Learning-induced genes were identified in transcript level comparisons between 2TD-trained vehicle-treated rats and sound-Naïve rats. Each dot represents a gene; some are labeled with gene names or identifiers. Significantly expressed genes (red) were determined by a set threshold of -log10(p-value) of 5.301 (*y-axis*; actual adjusted p-value = 0.000005). Genes included in enrichment analyses were only those that were significantly expressed and passed a log2FoldChange threshold of 0.65 (*x-axis*; actual fold-change value greater than 1.5692).

Figure 2. *RNA-sequencing reveals abundant transcriptional changes in auditory cortex.* RNA-sequencing of individual subject samples of auditory cortical tissue from naïve (N=4), or 2TD-trained vehicle treated ("Vehicle": N=4), and 2TD-trained HDAC-inhibitor treated ("RGFP966": N=4) rats. **(a)** Heatmap of the top 250 genes with the most different levels of expression between trained rats treated with RGFP966 vs. Naïve. Transcript levels for these 250 genes are shown for all three groups. A hierarchical clustering algorithm applied to categorize relative transcript levels across the 3 groups revealed two major clusters of effects: genes that are *upregulated* with 2TD learning and further upregulated by adding HDAC-inhibition (RGFP966) to 2TD learning, as opposed to genes that are *downregulated* with learning and further down by adding HDAC-inhibition to 2TD learning (see cluster brackets on the left of the heatmap plot). **(b)** Box and whisker plots show the same analysis extended to the top 500 genes. Each dot represents relative gene abundance of the same gene across the 3 conditions (Naïve, Vehicle or RGFP966). The left panel shows that 283/500 of the genes identified are upregulated with 2TD learning and further upregulated by adding HDAC-inhibition, while the right panel shows

217/500 DEGs are downregulated with learning and further downregulated by adding HDAC-inhibition. **(c)** Box and whisker plots of the top 500 genes with the most different levels of expression between trained rats treated with Vehicle vs. Naïve. As in (b), each dot represents relative gene abundance of the same gene across the 3 conditions (Naïve, Vehicle or RGFP966). The left panel shows that 250/500 of genes identified are *downregulated* with 2TD learning (Vehicle), which is blunted by adding HDAC-inhibition (RGFP966). The right panel shows that the other 250/500 of genes identified are *upregulated* with 2TD learning (Vehicle), which is blunted by adding HDAC-inhibition (RGFP966). **(d)** A direct comparison of differentially expressed genes (DEGs) between 2TD-trained groups, Vehicle vs. RGFP966, identified only four DEGs determined with an adjusted threshold of p<0.05.

Table 1. *Enrichment analysis of learning-dependent transcriptional changes identifies potential downstream pathways important for auditory function.* A meta-analysis shows the rank list order of unique biological pathways involved in learning-induced transcriptional changes identified in iPathwayGuide© ("Learning-induced": Vehicle vs. Naïve; "Learning-induced with RGFP966": RGFP966 vs. Naïve). Ranks are assigned by p-value (with Bonferroni correction) to identify top pathways. The number of genes reported as DEGs (DE) out of all genes involved in a pathway according to the iPathwayGuide© databases (ALL) are listed as "DE/ALL". *PV: p-value.*

Figure 3. *A gene-targeted approach using qRT-PCR validates selected genes of interest identified from genome-wide RNAseq.* A second cohort of animals was used for qRT-PCR analysis. They were trained identically to the animals that generated samples for RNAseq (Naïve: N=7; Vehicle: N=5; RGFP966: N=8). As for RNAseq, brains were collected 1 hour post injection of RGFP966 after the third daily session of 2TD training to generate auditory cortical RNA samples. All values were normalized to the housekeeping genes *Gapdh* or *18S* (values plotted show normalization to *Gapdh*). qRT-PCR results demonstrate robust learning-induced increases in expression of *Egr1, Nr4a1*, and *Per2*. All data are expressed as means (±standard errors). *p<0.05 (Kolmogorov-Smirnov test).