SWI/SNF complex-mediated chromatin remodelling is vital for cell surface adhesin repression and immune evasion

Kundan Kumar[1], Aditi Pareek[1,2] and Rupinder Kaur[1,*]

[1]Laboratory of Fungal Pathogenesis, Centre for DNA Fingerprinting and Diagnostics, Hyderabad-500039, India

[2]Graduate studies, Manipal Academy of Higher Education, Manipal-576104, Karnataka, India

[*]For correspondence: Rupinder Kaur, Centre for DNA Fingerprinting and Diagnostics (CDFD), Hyderabad, Telangana, India. Tel.: 91-40-27216137; Fax: 91-40-27216006; E-mail: rkaur@cdfd.org.in.

Key words: SWI/SNF chromatin remodelling complex, nucleosome mapping, pro-inflammatory cytokines, Epa1 adhesin, Yapsins, *Candida glabrata*, Macrophages, NLRP3 inflammasome, NF-κB

Running title: Role of chromatin remodelling in immune escape

Abstract

Immune evasion is critical for fungal virulence. However, how the human opportunistic pathogen *Candida glabrata* (*Cg*) accomplishes this is unknown. Here, using micrococcal nuclease-sequencing, RNA-sequencing, macrophage-signalling and genetic analyses, we demonstrate that chromatin remodelling in macrophage-internalized *Cg*, via CgSnf2 [SWI/SNF chromatin remodelling complex-ATPase subunit], leads to transcriptional upregulation and downregulation of immunosuppressive seven-gene mannosyltransferase-cluster (*CgMT-C*) and immunostimulatory cell surface adhesin-encoding *EPA1* gene, respectively. Consistently, *EPA1* overexpression and *CgMT-C* deletion led to increased proinflammatory cytokine IL-1β production and reduced *Cg* proliferation in macrophages. Further, *CgSNF2* deletion evoked increased IL-1β secretion, and the consequent killing of macrophage-internalized *Cg*, with elevated IL-1β levels reversed partially in Akt-, p38-, NF-κB- or NLRP3 inflammasome-inhibited macrophages. Importantly, macrophages respond to multiple *Candida* pathogens via NF-κB-dependent IL-1β production, underscoring NF-κB signalling's role in fungal diseases. Altogether, our findings directly link nucleosome positioning-based chromatin remodelling to fungal immunomodulatory molecule expression, which dictates *Cg* fate in host immune cells.

Introduction

Fungal infections cause >1.6 million deaths annually[1]. *Candida* species are the most common cause of invasive fungal infections, with *Candida glabrata* (*Cg*) being the second-to-fourth most prevalent infectious agent, based upon the geographical location[1–4]. *Cg* invasive

infections are associated with mortality rate exceeding 35%[5–7]. *Cg* is recently classified as a fungal pathogen of high-priority by the World Health Organization[8].

Cg belongs to the *Nakaseomyces* genus, also refers to as *Nakaseomyces glabrata*, and is evolutionarily closer to the budding yeast *Saccharomyces cerevisiae*[4,9]. *Cg* lacks secreted proteolytic activity and hyphal formation, and predominantly banks upon the stealth-strategy to persist in the host[4,10]. *Cg* circumvents the host immune system by inhibiting phagolysosome acidification and suppressing proinflammatory cytokine IL-1β production in macrophages[4]. Further, the type I interferon signalling contributes to *Cg* immune evasion by dysregulating iron and zinc homeostasis in macrophages[11,12].

Cg infection to macrophages induces transcriptomic changes in both macrophages and macrophage-internalized *Cg*, with *Cg* relying on multiple mechanisms including chromatin heterochromatinization, carbon metabolism reprogramming, autophagy induction and cell surface reconfiguration to thwart macrophage anti-*Cg* response, and proliferate in macrophages[4,10]. Macrophages activate NLRP3 inflammasome and produce IL-1β in spleen tyrosine kinase (Syk) signalling-dependent manner, upon *Cg* infection[13]. *Cg* utilizes a family of eleven putative glycosylphosphatidylinositol (GPI)-linked cell surface-associated proteases (CgYps1-11; CgYapsins) to restrict IL-1β secretion in infected-macrophages[13]. Consistently, *CgYPS1-11* loss led to *Cg* killing in macrophages and clearance from mice organs[13,14].

Another important intracellular survival strategy of *Cg* involves chromatin reconfiguration, with CgRtt109 (Histone H3 lysine-56 acetyltransferase) and two subunits of RSC chromatin remodelling complex (CRC), CgRsc3A and CgRsc3B, being required for *Cg* proliferation in macrophages[15]. Macrophage-internalized *Cg* displayed elevated lysine deacetylase activity, and increase and decrease in closed and open-chromatin marks, respectively, with remodelled-chromatin linking metabolic adaptation and energy homeostasis to *Cg* replication[15]. Further, 6 h and 10 h of macrophage internalization rendered *Cg* chromatin resistant to micrococcal nuclease (MNase) digestion, indicating a widespread chromatin reorganization[15]. A nucleosome, consisting of 147 bp DNA wrapped around the histone octamer, is the fundamental repeating subunit of chromatin, and nucleosome dynamics governs chromatin structure and functions[16]. Nucleosomes occupy defined positions in the genome, with DNA sequence, CRCs, transcriptional regulators and RNA polymerase II (RNAPII) transcription machinery acting in conjunction to maintain nucleosome locations[16]. Nucleosome positioning control DNA accessibility to various enzymes and regulatory proteins[16,17]. The nucleosome-depleted regions (NDRs), and/or chromatin regions opened up locally by CRCs, facilitate binding of transcription factors and RNAPII machinery to DNA, thereby promoting gene expression[16,17].

ATP-dependent CRCs mobilize nucleosomes via DNA translocation, histone exchange and/or DNA-histone disengagement[17]. These belong to four subfamilies: imitation switch (ISWI), chromodomain helicase DNA-binding (CHD), SWI/SNF (SWItch/Sucrose Non-Fermentable) and INO80, with their catalytic ATPase subunit aiding in DNA translocation[17]. The SWI/SNF complex in *Cg* and *S. cerevisiae* consists of 12 subunits[18]. Transcription factors recruit SWI/SNF complex at specific DNA sites in *S. cerevisiae*, with ScSnf2 involved in DNA binding[17]. Notably, ScSnf2 localizes to NDRs, and controls transcription initiation by governing +1 nucleosome (transcription start site-associated) position[19].

Despite chromatin architecture being pivotal to intracellular survival of *Cg*, the nucleosome landscape of macrophage-internalized *Cg* is undefined. Here, we report the first genome-wide nucleosome map of macrophage-internalized *Cg* consisting of ~70,000 nucleosomes, including 12,000 nucleosomes that exhibited changes in position, occupancy or fuzziness. Additionally, we present roles of seven ATP-dependent chromatin remodelling complexes in *Cg* pathobiology, underscoring the essentiality of CgSnf2 for immune evasion and virulence. Altogether, our findings unify two central virulence mechanisms, chromatin restructuring and immune response evasion, in *Cg*.

Results

MNase-sequencing analysis reveals nucleosome dynamics in macrophage-internalized *Cg*

Nucleosome dynamics encompasses nucleosome position shifts, occupancy changes and fuzziness changes[16,17,20]. To investigate nucleosome dynamics, we performed MNase-seq analysis in 2 h and 10 h macrophage-internalized and RPMI-grown *Cg* cells (Fig. 1a). We identified 56,637-71,823 nucleosomes under four-studied conditions (Fig. 1b and Supplementary Table 1), with 12-24% exhibiting changes in position, occupancy or fuzziness (Fig.1 c and d and Supplementary Table 2). Most nucleosomes displayed a fuzziness score of 50-60 (Extended Data Fig. 1a). Most chromosomes contained a high number of dynamic nucleosomes at their ends (Fig.1e). Further, gene bodies displayed the largest number of nucleosomes (Extended Data Fig. 1b and c), with macrophage-internalized *Cg* showing 5-6% more nucleosomes mapped to intergenic regions (Extended Data Fig. 1d). Although overall genomic nucleosome distribution remained unaltered (Supplementary Fig. 1), 5% of dynamic nucleosomes were present in promoter regions across four compared datasets (Extended Data Fig. 1e). Notably, nucleosome fuzziness was the least observed change in all comparisons (Fig. 1d and Extended Data Fig. 2a-c; and Supplementary Table 2).

Consistent with macrophage-induced largescale chromatin changes[15], nucleosomes in 10 h macrophage-internalized *Cg* were more dynamic, compared to 2 h macrophage-internalized *Cg* (Fig. 1c), with position shift being the predominant change (Extended Data Fig. 2b). 10 h macrophage-internalization also led to highest gain in nucleosome occupancy (Extended Data Fig. 1f) and lowest dynamic nucleosome numbers at gene promoters (Extended Data Fig. 1e and Supplementary Table 3), compared to growth in RPMI medium.

Functional analysis revealed that promoters of genes, belonging to cell surface, translation, copper ion import and tricarboxylic acid (TCA) cycle GO terms, displayed altered nucleosome configurations in 2 h macrophage-internalized *Cg*, with translation gene promoters showing high nucleosome occupancy (Fig. 1f and Supplementary Tables 3 and 4). Notably, TCA cycle and translation genes in *Cg* are upregulated and downregulated, respectively, in response the macrophage milieu[14,15]. Further, compared to 2 h, 10 h macrophage internalization led to dynamic nucleosomes in promoters of genes belonging to fungal-type cell wall, translation and adhesion of symbiont to host (Extended Data Fig. 2d and Supplementary Table 4). Notably, fungal-type cell wall gene promoters contained dynamic nucleosomes in both 10 h macrophage-internalized and 10 h RPMI-grown *Cg*, compared to 10 h and 2 h RPMI-grown *Cg*, respectively (Extended Data Fig. 2e and f, and Supplementary Table 4), probably reflecting cell wall gene expression plasticity.

We draw six major conclusions from our MNase-Seq data which report nucleosome dynamics for the first time in a host cell-internalized eukaryotic pathogen. First, promoter regions in *Cg* are nucleosome-depleted, consistent with the previous study[21]. Second, 10-20% nucleosomes displayed changes in position, occupancy or fuzziness upon macrophage internalization. Third, macrophage-induced transcriptional downregulation and upregulation, respectively, of translation and TCA cycle genes may be governed by increased and decreased nucleosome occupancy at their promoters. Fourth, nucleosome positioning may govern expression of cell wall genes in response to the environmental cues. Fifth, compared to gene bodies, dynamic nucleosomes were double the number in promoter and intergenic regions after 2 h macrophage internalization. Finally, position shift was the most common nucleosome dynamic event in *Cg*. Altogether, besides revealing macrophage milieu-induced gene-specific nucleosome-pattern in *Cg*, our findings raise the possibility that CRCs, which are essential for nucleosome dynamics, may drive *Cg* survival in macrophages.

CRCs modulate intracellular survival of *Cg*

Cg possesses seven ATP-dependent CRCs. To elucidate their functions, we generated six deletion strains, *Cgsnf2Δ*, *Cgisw1Δ*, *Cgisw2Δ*, *Cgchd1Δ*, *Cgino80Δ* and *Cgswr1Δ*, for non-essential genes that encode putative ATPase subunits of CRCs (Extended Data Fig. 3a), and examined their growth profiles and virulence-associated traits. Since *Cgrsc3aΔbΔ* previously displayed reduced survival in macrophages and mice[15], we included this mutant in our analysis as well. Phenotypic profiling revealed pleiotropic stress, and low pH and high iron sensitivity of *Cgsnf2Δ* and *Cgswr1Δ*, respectively, (Fig. 2a). *Cgsnf2Δ* grew extremely slowly with a doubling time of 5 h in YPD medium (Extended Data Fig. 3b and c), and exhibited elongated cell morphology (Extended Data Fig. 3d). *Cgrsc3aΔbΔ* and *Cgino80Δ* could not utilize alternate carbon sources and displayed reduced growth in iron-surplus medium, with *Cgino80Δ* also exhibiting elevated thermal stress susceptibility (Fig. 2a), and 40% longer doubling time than *wild-type* (*wt*) (Extended Data Fig. 3c).

These distinct stress sensitivity phenotypes notwithstanding, three mutants, *Cgsnf2Δ*, *Cgino80Δ* and *Cgswr1Δ,* were attenuated in their ability to form biofilms on polystyrene surface (Fig. 2b). Further, human THP-1 macrophage-infection analysis revealed differential survival of chromatin remodeller-mutants. While *Cgisw1Δ*, *Cgisw2Δ*, *Cgchd1Δ* and *Cgswr1Δ* replicated like *wt* in THP-1 cells, *Cgrsc3aΔbΔ* and *Cgino80Δ* were defective in intracellular proliferation (Fig. 2c). Contrarily, *Cgsnf2Δ* was killed in macrophages (Fig. 2c), underscoring CgSnf2 essentiality for intracellular survival of *Cg*. Importantly, all mutants were engulfed similarly to *wt* (Extended Data Fig. 3e), and *Cgsnf2Δ* and *Cgino80Δ* replicated in RPMI medium, albeit slower than *wt* (Extended Data Fig. 3f). Further, CgSnf2 was crucial for *Cg* survival in all three target organs, kidneys, liver and spleen, in the mouse model of disseminated candidiasis, as 5- to 100-fold lower organ fungal load was observed in *Cgsnf2Δ*-infected mice, compared to *wt*-infected mice (Fig. 2d). Notably, *Cgisw2Δ*- and *Cgino80Δ*-infected mice displayed 20-fold lower and 7-fold higher renal fungal burden, respectively (Fig. 2d), indicating positive and antagonistic role for CgIsw2 and CgIno80, respectively, in *Cg* survival in kidneys. Altogether, besides implicating ATP-dependent CRCs in diverse stress survival *in vitro*, these data underscore that while CgChd1 and CgIsw1 are dispensable for the

examined pathogenesis-associated traits, CgSnf2 is pivotal to *Cg* virulence. Notably, CgSnf2 loss is known to adversely affect growth and biofilm formation[22].

CgSnf2 suppresses macrophage activation

Among CRC mutants, *Cgsnf2Δ* was most severely compromised for pathogenicity, we therefore decided to characterize *Cgsnf2Δ* further. First, we showed through complementation analysis that the diverse stress susceptibility (Extended Data Fig. 4a), and elongated cell morphology of *Cgsnf2Δ* (Extended Data Fig. 4b) is due to *CgSNF2* absence. Similarly, *CgSNF2*-expressing *Cgsnf2Δ* displayed *wt*-like survival in macrophages and mice (Extended Data Fig. 4c and d).

Since *Cg* survival in THP-1 macrophages was associated with NLRP3 inflammasome- and Syk-dependent IL-1β secretion inhibition[13], we next determined if intracellular killing of *Cgsnf2Δ* is owing to elevated IL-1β secretion. We found 15-fold increased IL-1β secretion by *Cgsnf2Δ*-infected THP-1 macrophages, compared to *wt*-infected macrophages (Fig. 3a), which was brought down by 4-fold by the NLRP3 inhibitor MCC950 (Fig. 3b), with MCC950 also partially rescuing *Cgsnf2Δ*'s intracellular survival defect (Fig. 3c). *Cgsnf2Δ* also exhibited 2-fold higher survival in R406 (Syk inhibitor)-treated macrophages, compared to untreated macrophages (Fig. 3d). Further, *Cgsnf2Δ* was killed in primary murine macrophages (Fig. 3e) which also exhibited 10-fold increased IL-1β levels, with MCC950 treatment diminishing IL-1β secretion (Fig. 3f), underscoring the conserved response of cultured and primary, as well as, human and mouse macrophages to *Cgsnf2Δ* infection. Altogether, these results implicate CgSnf2 in *Cg*-mediated suppression of NLRP3 inflammasome activation, and reinforce that impeding IL-1β production is crucial for *Cg* survival in macrophages.

CgSnf2 governs transcriptional response to the macrophage internal milieu

Cg persists in macrophages in three-distinct phases, Early, Mid and Late stage, with genome heterochromatinization being a trademark of the Mid-stage (6-12 h of macrophage-ingestion)[15]. Notably, CRCs facilitate heterochromatin formation as well as orchestrate transcriptional regulatory proteins-target DNA interactions, that govern gene expression[17,23]. Therefore, to determine the molecular basis underlying CgSnf2-mediated macrophage activation abolishment, we profiled transcriptomes of macrophage-internalized *wt* and *Cgsnf2Δ* at 2 h and 10 h post-infection via RNA-sequencing, and identified differentially expressed genes (DEGs; genes showing ≥ 2-fold change in expression). A total of 1410 (834 upregulated and 576 downregulated) and 192 (162 upregulated and 30 downregulated) genes displayed differential expression in 2 h macrophage-ingested *wt* and *Cgsnf2Δ*, compared to corresponding RPMI-grown cells (Fig 4a and b, and Supplementary Table 5). Similarly, 10 h macrophage-internalization led to differential expression of 622 (482 upregulated and 140 downregulated) and 212 (155 upregulated and 57 downregulated) genes in *wt* and *Cgsnf2Δ*, respectively (Extended Data Fig. 5a-c, and Supplementary Table 5), indicating that the magnitude of *Cg* transcriptional response to the macrophage milieu is dependent upon CgSnf2. Consistently, of all DEGs, only 42 genes were found to be CgSnf2-independent (Fig. 4c).

DAVID functional analysis revealed fatty acid β-oxidation (FAO) and tricarboxylic acid (TCA) cycle enrichment, in addition to other GO terms, in upregulated gene sets in 2 h and 10 h macrophage-internalized *wt* (Extended Data Fig. 5d and Supplementary Fig. 2 and Table 6). Contrarily, while rRNA processing, translational frameshifting and ribosome biogenesis terms were enriched in downregulated genes in 2 h macrophage-internalized *wt*, 10 h macrophage internalization led to downregulation of de novo' IMP biosynthetic process, DNA replication and iron ion homeostasis (Extended Data Fig. 5d and Supplementary Fig. 2 and Table 6). These data are consistent with earlier microarray-based studies reporting similar gene expression patterns for macrophage-internalized *Cg*[14,15], and underscore carbon metabolism reprogramming and translational machinery shutdown as characteristic components of *Cg* transcriptional response to the macrophage environment.

Intriguingly, comparison of 2 h RPMI-grown *wt* and *Cgsnf2Δ* transcriptomes revealed that *CgSNF2* deletion led to upregulation and downregulation of FAO, TCA cycle, glyoxylate cycle and adhesion of symbiont to host, and rRNA processing, translational frameshifting and ribosome biogenesis, respectively (Fig. 4d and Supplementary Table 6). Since this transcriptional signature is hallmark of 2 h macrophage-internalized *wt*, it could partly account for *Cgsnf2Δ*'s muted-transcriptional response to the macrophage ingestion. Further, the 'fungal-type cell wall organization' term was enriched in upregulated genes in 10 h RPMI-grown *Cgsnf2Δ*, compared to corresponding *wt* (Extended Data Fig. 5d and Supplementary Table 6). We verified the physiological relevance of this observation by performing cell wall component analysis, and found elevated β-glucan and chitin content in *Cgsnf2Δ* (Extended Data Fig. 5e), which suggests that CgSnf2 maintains cell wall composition probably by regulating the expression of cell wall organization genes.

Cgsnf2Δ responded to 2 h of macrophage-ingestion by upregulating iron-sulfur cluster assembly, TCA cycle and inositol biosynthetic process, and downregulating axial cellular bud site selection (Fig. 4e and Supplementary Table 6). Notably, TCA cycle and translational frameshifting were upregulated and downregulated, respectively, in 10 h macrophage-internalized *Cgsnf2Δ* (Extended Data Fig. 5d and Supplementary Table 6). Besides showing the *wt*-like carbon metabolism reconfiguration, these data collectively underscore an impaired transcriptional response of *Cgsnf2Δ* to closedown translational machinery, upon macrophage ingestion, probably because of already-downregulated translation genes in the mutant.

Notably, contrary to *wt*, *Cgsnf2Δ* showed no activation of the seven mannosyltransferase gene-containing cluster (*MT-C*), which included five β-1,2-mannosyltransferase-encoding genes, *CgBMT2-6*, and two α-1,3-mannosyltransferase-encoding genes, *CAGL0B02981g* and *CAGL0B03014g*, after 2 h of macrophage internalization (Extended Data Fig. 6a-c). Further, comparative transcriptome analysis of 2 h macrophage-ingested *Cgsnf2Δ* and *wt* revealed TCA cycle, gluconeogenesis and FAO downregulation, and NAD transport, adhesion of symbiont to host and fungal-type cell wall organization upregulation (Extended Data Fig. 5d and Supplementary Fig. 2 and Table 6). Since *Cg* adherence to macrophages is the first step in its phagocytosis by macrophages, we examined closely how *CgSNF2* loss impacts expression of cell wall adhesin-encoding genes. Of known 81 adhesins[24], 21 adhesin genes, including 9 subtelomeric adhesin genes, were differentially expressed in *Cgsnf2Δ* (Fig. 5a). Notably, some *EPA* (Epithelial adhesin) genes are subjected to

subtelomeric silencing owing to their location at chromosome ends, with CgSnf2 negatively affecting subtelomeric silencing[22,25]. CgSnf2 has also been reported to regulate *EPA1* and *EPA6* expression negatively and positively, respectively[22].

Based on these data, we hypothesized that impaired α- and β-mannan, and adhesin-dependent cell surface remodelling, due to perturbed transcriptional regulatory networks, may contribute to increased IL-1β production and *Cgsnf2Δ* death in macrophages.

CgSnf2-dependent *EPA1* regulation is pivotal to *Cg*-mediated suppression of IL-1β secretion in macrophages

To test our hypothesis, we first confirmed *CgBMT2* upregulation in macrophage-internalized *wt* by qRT-PCR (Extended Data Fig. 6d). Next, we generated a mutant lacking all seven genes of the mannosyltransferase cluster, and found that *MT-C* deletion led to reduced intracellular proliferation and increased IL-1β secretion (Extended Data Fig. 6e and f). These results suggest that β-1,2-, and α-1,3-oligomannosides in *Cg* cell wall aid in immunosuppression. Of note, CgBmt2-6 were required for gut colonization in the murine colitis model[26].

Next, to examine the role of adhesins in immunosuppression, we selected *EPA1*, from all differentially-expressed adhesin genes, for further analysis primarily for 3 reasons: (1) Epa1 is the major adhesin for *Cg* adherence to host cells[27,28], (2) *EPA1*-expressing *S. cerevisiae* adhered well to macrophages and stimulated inflammatory cytokine production[28], and (3) the eleven aspartyl protease-deficient *Cgyps1-11Δ* mutant, with elevated surface-exposed Epa1, was killed in macrophages, due to increased IL-1β secretion[13,14]. We first verified RNA-seq results by qRT-PCR and showed that *EPA1* transcription was activated and repressed upon *CgSNF2* deletion and macrophage-internalization of *wt* cells, respectively (Fig. 5b and c), raising two possibilities: (1) *EPA1* downregulation aids in suppressing macrophage pro-inflammatory response, and (2) Increased *EPA1* expression is deleterious to *Cg* survival.

To address these, we performed four experiments. First, we overexpressed *EPA1* from strong *PDC1* promoter in *wt* and profiled growth in THP-1 cells. We found *wt/EPA1* to display reduced proliferation (Fig. 5d), and 3-fold increased IL-1β secretion in macrophages (Fig. 5e). Since Epa1 is a calcium-dependent lectin, lactose treatment inhibits its binding to host asialo-lactosyl-containing carbohydrates[27]. Consistently, THP-1 infection with lactose-treated *wt/EPA1* reduced IL-1β secretion significantly (Fig. 5e), reinforcing the role of Epa1 in modulating IL-1β production. Second, we deleted *EPA1* gene in *Cgsnf2Δ* background, and found that the *Cgsnf2Δepa1Δ* infection invoked 1.5-fold less IL-1β production in THP-1 macrophages, compared to infection with the single *Cgsnf2Δ* mutant (Fig. 5f). IL-1β secretion was similar in response to *wt* and *epa1Δ* infection, probably due to functional redundancy among Epa adhesins (Fig. 5f). Third, we performed the same analysis with *Cgyps1-11Δ* after deleting *EPA1*, and found that *Cgyps1-11Δepa1Δ*-infected macrophages secreted 1.3-fold less IL-1β than *Cgyps1-11Δ*-infected cells (Fig. 5g). Notably, CgYapsins are required for Epa1 processing from the cell wall, and *Cgyps1-11Δ* contained increased Epa1 levels in its cell wall[14]. Fourth, we checked the intracellular survival of *epa1Δ* in THP1-macrophages, and found *wt*-like intracellular proliferation, while *Cgsnf2Δepa1Δ* showed 27% better survival than

Cgsnf2Δ (Fig. 5h)*,* highlighting Epa1's adverse contribution to *Cgsnf2Δ* survival in macrophages.

Altogether, these data suggest that Epa1 is immunostimulatory, acts as a fungal activator of IL-1β induction, and that *EPA1* levels are probably regulated transcriptionally by CgSnf2 via nucleosome repositioning, and posttranslationally by CgYapsins through cleavage at the cell wall.

CgSnf2 binds to *EPA1* promoter

To demonstrate that CgSnf2 directly regulates *EPA1* expression, we performed two experiments. First, through chromatin immunoprecipitation, we showed 2-fold enrichment of CgSnf2 on *EPA1* promoter (Fig. 5i), suggesting that CgSnf2 keeps *EPA1* expression in check under normal growth conditions. Second, we mutated the conserved serine (Ser-861) residue in the ATPase domain of CgSnf2 (Extended Data Fig. 7a and b) to aspartate, and found CgSnf2^{S861D} to express well (Extended Data Fig. 7c). However, CgSnf2^{S861D} could neither rescue stress susceptibility of *Cgsnf2Δ* and nor abolish elevated IL-1β secretion in *Cgsnf2Δ*-infected macrophages (Extended Data Fig. 7d and e), underscoring CgSnf2 catalytic functions, and by extension SWI/SNF complex-mediated chromatin remodelling, in *Cg*-mediated suppression of the host immune response.

CgSnf2-dependent IL-1β suppression involves NF-κB activation

NLRP3 inhibition in macrophages neither completely reversed IL-1β secretion nor *Cgsnf2Δ* viability (Fig. 3), which could be due to other activated immune signalling pathways. Therefore, to define CgSnf2-repressed IL-1β-producing host signalling pathways, we focussed on three pathways, nuclear factor-kappa B (NF-κB), phosphoinositide-3-kinase(PI3K)-protein kinase B/Akt (Akt) and p38 mitogen-activated protein kinase (MAPK) signalling pathways. The NF-κB, a heterodimer of p50 and p65 proteins, is a transcriptional factor which regulates NLRP3 expression and controls macrophage inflammatory gene expression[29]. Akt and p38 pathways are also implicated in antifungal immunity[29]. We found 1.8-, 2- and 10-fold higher phosphorylation of p65 (regulatory protein of NF-κB), Akt serine/threonine kinase (downstream effector of PI3K signalling), and p38 MAPK in *Cgsnf2Δ*-infected macrophages, compared to uninfected macrophages, respectively (Fig. 6a and Extended Data Fig. 8a and b), indicating a role for CgSnf2 in dampening activation of three other host pathways. Consistently, IL-1β levels were lower in BAY 11-7082 (blocks the inhibitory kappa B protein of NF-κB activation), SH-6 (Akt-specific inhibitor), and SB203580 (p38 inhibitor)-treated *Cgsnf2Δ*-infected macrophages, compared to untreated mutant-infected macrophages (Fig. 6b and Extended Data Fig. 8c and d). Further, *Cgsnf2Δ* infection also resulted in elevated secretion of two other pro-inflammatory cytokines, TNF-α and IL-8, in macrophages, which were partially reversed by specific-immune signalling inhibitors (Fig. 6c and d). These data suggest that *C. glabrata* suppresses macrophage activation by controlling inflammatory response via multiple signalling pathway modulation.

Since fungal PAMP (pathogen-associated molecular pattern)-induced signalling pathways may converge on NF-κB activation[29], which is associated with p65 nuclear translocation, we examined p65 cellular localization in J774A.1 macrophages, as these murine

macrophage-like cells have larger cytosol volume. We found 1.5-fold higher translocation of p65 to the nucleus (Fig. 6e) in *Cgsnf2Δ*-infected, compared to *wt*-infected macrophages. Further, the role of Epa1 in *Cg*-mediated suppression of NF-κB pathway was demonstrated by enhanced and diminished nuclear translocation of p65 in J774A.1 macrophages infected with *EPA1*-overexpressing *Cg* (Fig. 6e) and *Cgsnf2Δepa1Δ* cells (Fig. 6e), respectively. Altogether, these data reveal a hitherto unknown facet of *Cg*-mediated host response modulation, via Epa1 adhesin, that involves the master regulator, NF-κB.

Finally, to investigate if NF-κB-dependent inflammatory response represents a pan anti-*Candida* response of macrophages, we infected THP-1 cells with four *Candida* species, *Cg*, *C. albicans*, *C. auris* and *C. tropicalis*. We found increased IL-1β production in all infections, which was reduced upon treatment with BAY 11-7082 (Fig. 6f), indicating that NF-κB governs macrophage inflammatory response against *Candida* pathogens. Notably, *S. cerevisiae* infection of THP-1 cells invoked no IL-1β secretion (Extended Data Fig. 8e). Altogether, these data indicate that macrophage respond to pathogenic *Candida* yeasts via IL-1β production, and controlled augmentation of IL-1β has potential for host-directed anti-*Candida* therapy.

Discussion

Cg resides on mucosal surfaces in healthy humans but causes superficial mucosal and life-threatening invasive infections in immunocompromised patients[30]. Macrophages constitute key host immune defence cells against candidemia, with fungal cell wall structural components, β-glucan and mannan, involved in immune recognition[31]. Here, we present the first genome-wide nucleosome landscape of macrophage-internalized *Cg*, and establish CgSnf2 as a key determinant of fungal immune evasion. We show that *Cg* modulates NF-κB signalling, by regulating the probable host-recognizable immunostimulatory Epa1 adhesin and immunosuppressive β-1,2- oligomannoside expression, via CgSnf2, to subvert macrophage proinflammatory cytokine production.

The active immune suppression mechanisms in *Cg* are poorly-understood, and involve impeding phagolysosome acidification and suppressing Syk-dependent IL-1β production[13,15]. *Cg* cell wall, the first point-of-contact with the host, is a multilayered organelle consisting of an inner structural network of chitin, immunostimulatory β-1,3-glucan and β-1,6-glucan, with the outer mannan layer, comprised of N-linked or O-linked mannosylated proteins, shielding the inner core from immune cells[4,31]. Dectin-1, Dectin-2, mannose, TLR4, galectin 3 and TLR9 receptors are involved in β-1,3-glucan, α-mannan, mannan and mannoprotein, O-linked mannan, β-mannan, and chitin and DNA recognition, respectively[31]. Epa1, founding member of the Epa-adhesin family, is a GPI-anchored, glucan cross-linked, calcium-dependent adhesin, possesses a N-terminal PA14 ligand-binding domain, and resides in the outer layer of the cell wall[27,32,33]. Epa1 is shaved off the cell wall in CgYapsins-dependent manner[14]. Epa1 binds to host glycan ligands containing terminal galactose residue, and mediates adhesion to host epithelial, endothelial and macrophage cells[27,28,33,34]. Our data establish Epa1 as a fungal PAMP, whose finely-tuned expression is pivotal to suppress the pro-inflammatory innate immune response. Notably, *EPA1* overexpression, due to gain-of-function mutation in CgPdr1 transcriptional activator, was associated with decreased *Cg* phagocytosis by macrophages in a

Cg strain-dependent manner, and increased adherence to epithelial cells in all *Cg* strain backgrounds[35], underscoring two distinct roles for Epa1 in host-*Cg* interaction. The *wt*-like phagocytosis rate of *Cgsnf2Δ* (Extended Data Fig. 3e) is consistent with CgPdr1 mutation-dependent *EPA1* overexpression not affecting internalization of its parent strain, BG2[35]. Epa1 binding to NKp46 receptor led to *Cg* killing in natural killer cells[36], therefore, the decrease in surface-exposed Epa1 is likely to constitute a principal *Cg* defence strategy against multiple immune cell-types.

CgSnf2 regulates both basal and macrophage milieu-responsive expression of *EPA1* and other adhesin genes. *Cg* possesses 81 adhesins, with many adhesin genes encoded at subtelomeric regions[24,33]. Nicotinic acid limitation relieves subtelomeric adhesin gene silencing due to restricted NAD^+ availability lowering activity of the NAD^+-dependent histone deacetylase Sir2[33,37]. Expression of *EPA1*, which is localized 25 kb away from the right telomere of chromosome E, is negatively regulated by the subtelomeric silencing-protein complex, protosilencer Sil2126 and a negative-sequence element[33]. We add another regulatory layer of CgSnf2-dependent chromatin remodelling to *EPA1* transcription control which may involve +1 nucleosome position shift, as evidenced in a 60 bp shift towards the *EPA1* gene body in 2 h macrophage-internalized *Cg*, compared to RPMI-grown *Cg* (Extended Data Fig. 9a). Notably, CgSnf2 regulates +1 nucleosome position in multidrug transporter gene promoter in *S. cerevisiae*[19]. Additionally, we observed nucleosome compaction at *EPA1* promoter and 3' UTR, with average internucleosomal distance being 206 bp and 177 bp in RPMI-grown and macrophage-internalized *Cg*, respectively (Extended Data Fig. 9b). This complex Epa1 regulation could be pivotal to induce *EPA1* expression for initial infection-stages involving *Cg* adherence to host epithelial and endothelial cells, and repress *EPA1* transcription later to suppress immune recognition and activation. Consistently, early Syk inhibition, following *Cg* ingestion, is pivotal to control *Cg* proliferation effectively in macrophages[38], and timely fungal immunogenic molecule-masking and unmasking aids in manipulating immune responses[39].

We speculate/propose that in the absence of the hyphal form, *Cg* primarily relies on CgSnf2-dependent dynamic cell surface organization to avoid immune surveillance/sentinel cells (Extended Data Fig. 10). Accordingly, RNAPII ChIP-seq analysis revealed that *Cg* adhesin genes were upregulated and downregulated, marking immediate (30 min) and late (2 h onwards) response, respectively, to macrophage infection, with CgSnf2 being transcriptionally downregulated at 30 min post-infection[40]. This study also reported 1589 genes to be either constitutively-expressed or temporally-induced in macrophage-internalized *Cg*[40], which is similar to 1410 DEGs, we identified in 2 h macrophage-ingested *Cg*. Our data suggest that macrophage activation is likely attained by multiple immune signalling pathways. Consistently, loss of Dectin-1 or Dectin-2 (α-mannan receptor)-mediated Syk signalling individually had no effect on *Cg* replication in macrophages[38]. Altogether, although our data unequivocally show epigenetic regulation of *Cg* immunomodulatory factors, the nature and distribution of these PAMPs (cell wall-associated or shedded Epa1; α- or β-mannosides), and their immune receptors remain to be identified. Similarly, whether and how Epa1 and mannosides co-operate to antagonize immune-signalling pathways, thereby generating *Cg*-specific immune evasion mechanisms, warrant further studies.

Microbial pathogens often employ heterochromatin structure to control virulence gene expression[23], and nucleosome position changes regulate stress-response genes, with growth and stress-gene promoters exhibiting distinct chromatin features in hemiascomycetous yeasts, including Cg[21]. Cg dynamically switches its chromatin between transcriptionally-active euchromatin and transcriptionally-silent heterochromatin[15]. We mapped 86-102 nucleosomes to the 18.64 kb mannosyl transferase gene-cluster region under R2, M2, R10 and M10 conditions, with 4-22% nucleosomes exhibiting altered occupancy or position (Supplementary Tables 1 and 2). Additionally, an increased occupancy at +1 and -2 nucleosomes for $CgBMT5$, and a 90 bp shift towards TSS was observed for $CAGL0B02981$g gene, in 2 h macrophage-internalized Cg (Extended Data Fig. 9c). Notably, the dynamic nucleosome number at MT-C loci was 3- to 6-fold lower in 10 h macrophage-internalized/10 h RPMI-grown Cg, compared to other three datasets (Supplementary Table 2). Altogether, these results implicate nucleosome dynamics in controlling MT-C gene expression, though the significance of observed nucleosome position and occupancy changes remains to be determined. Overall, our data also raise the possibility that the nucleosome pattern on immunomodulatory genes in pathogenic fungi may differ from the gene-averaged nucleosome pattern, which can potentially be useful for therapeutic interventions.

CRCs remodel chromatin by changing composition of, mobilizing or evicting nucleosomes[17]. Cg has seven ATP-dependent CRCs, of which RSC complex ATPase CgSth1 may be essential for Cg growth. Since 59% of DEGs in macrophage-internalized Cg contained CgRsc3-binding sites[15], multiple chromatin remodelers may direct nucleosome position. Identifying genomic loci with repositioned/shifted nucleosomes in macrophage-internalized $Cgrsc3a\Delta b\Delta$ and $Cgsnf2\Delta$ mutants, and genome-wide mapping of CgSnf2 and CgSth1, will unveil individualistic CRC contribution. However, given CgSnf2's essentiality for intracellular survival, its ability to create facile-chromatin regions and recruit specific transcription factors and/or RNAPII machinery may contribute majorly to Cg immune evasion.

In conclusion, chromatin reorganization in Cg impacts macrophages' inflammatory response and Cg survival. Due to CgSnf2 indispensability for immune evasion, exploring fungal-specific subunits of the SWI/SNF complex[18] as new therapeutic targets holds promise.

Methods

Ethics statement

Mice infection experiments were conducted at the Animal House Facility of Centre for DNA Fingerprinting and Diagnostics (CDFD; www.cdfd.org.in) or the CDFD animal facility, VIMTA Labs Limited (http://www.vimta.com), Hyderabad, India in accordance with guidelines of the Committee for the Purpose of Control and Supervision of Experiments on Animals, Government of India. The protocols were approved by the Institutional Animal Ethics Committees of CDFD (EAF/RK/CDFD/22) and VIMTA Labs Ltd (PCD/CDFD/05). Procedures were designed to minimize animal suffering.

Strains, plasmids and growth conditions

C. glabrata (*Cg*) *wild-type* (*wt*) and mutant strains, derivatives of the *Cg*-BG2 strain, were maintained on yeast extract-peptone-dextrose (YPD) medium or minimal yeast nitrogen base medium containing Casamino Acid (CAA) at 30°C. The *Escherichia coli* DH5α strain was

used for gene cloning and plasmid propagation, and grown in LB medium containing ampicillin at 37°C. To collect logarithmic (log)-phase *Cg* cells, overnight-grown *Cg* strains were re-inoculated at 0.1 OD_{600} in fresh YPD/CAA medium and cultured for 4-5 h at 30°C.

      Cg deletion strains were generated, using the homologous recombination-based approach, wherein the *Cg* gene was replaced with the *nat1* gene, that imparts nourseothricin resistance as a selection marker, as described previously[41]. Despite multiple attempts, *CgSTH1* could not be deleted, which may reflect essentiality of CgSth1 for *Cg* growth. For generation of double mutants, *Cgsnf2Δepa1Δ* and *Cgyps1-11Δepa1Δ*, *CgSNF2* (*CAGL0M04807g*) and *EPA1* (*CAGL0E06644g*) genes were deleted in *Cgsnf2Δ* and *Cgyps1-11Δ* strain backgrounds, respectively. For complementation analysis, *CgSNF2* (5.193 kb) gene along with 5' (671 bp)- and 3' (720 bp)-UTR sequences, was cloned at *BamHI* and *XmaI* sites in the pRK1016 (pGRB2.1) vector. For ChIP analysis, *CgSNF2* gene was tagged with the SFB [S-protein(S)-FLAG(F)-Sterptavidin-binding-peptide(B)] epitope at the C-terminus via cloning at Spe*I* and Xma*I* restriction sites in the pRK1349 plasmid, which carries *PDC1* promoter. *CgSNF2^{S861D}* was generated by rolling circle method using pRK2404 (*PDC1-CgSNF2-SFB*) plasmid DNA as template. For overexpression analysis, *EPA1* (3.105 kb) gene was cloned in *XmaI* and *XhoI* sites, downstream of the strong constitutive *CgPDC1* promoter in the pRK999 vector. The *Candida* strains, plasmids, primers, and antibodies and inhibitors used in this study are listed in and Supplementary Tables 7, 8, 9 and 10, respectively.

Growth analysis

Growth profiles of *Cg* strains in the presence of different stressors were analysed by serial dilution spot or time course assays in solid and liquid medium, respectively. For spot assay, OD_{600} of overnight-grown *Cg* cultures was normalized to 1.0, and cultures were 10-fold serially diluted in phosphate-buffered saline (PBS), followed by spotting 3 μl of each dilution on appropriate medium, and recording growth after 1 to 2 days of incubation at 30°C. For time-course analysis, overnight-YPD medium-grown *Cg* strains were inoculated in fresh YPD medium at an initial OD_{600} of 0.1 and incubated at 30ºC with shaking. OD_{600} of cultures was monitored at regular intervals till 36 h, followed by plotting of the absorbance readings as a function of time. Generation time for each strain was calculated during the log-phase (2 h - 8 h) of growth. *Cg* replication in RPMI medium was determined by colony-forming unit-based assay, wherein overnight-YPD medium-grown *Cg* strains were inoculated in RPMI medium containing 10% FBS (fetal bovine serum) at an initial OD_{600} of 0.1 and incubated at 37ºC with 5% CO_2. OD_{600} of cultures was recorded periodically, and appropriate dilution of cultures were plated on YPD medium. After 1-2 days of incubation at 30ºC, *Cg* colonies were counted, and CFUs were plotted as a function of time. Mutants showing ≥ 20% longer doubling time than *wt* cells were considered to have retarded growth rate.

Biofilm formation assay

Log-phase, YPD medium-grown *Cg* cells seeded in a 24-well polystyrene plate at an OD_{600} of 0.5. After 90 min incubation at 37°C, non-adherent *Cg* cells were washed off with PBS, and adherent cells were grown in RPMI 1640 medium containing 10% FBS for 48 h at 37°C, with a medium change at 24 h. After removing unbound cells with PBS washes, adherent *Cg* cells were stained with 0.4% (w/v) crystal violet for 45 min, followed by destaining in 95% ethanol. Absorbance of the destaining solution was recorded at 595 nm, *Cg*-lacking blank well

absorbance values were subtracted from *Cg*-containing well absorbance values, and biofilm-forming ability of *Cg* mutants was plotted as the biofilm ratio which represents the mutant/*wt* absorbance values. Mutants showing ≥ 1.25-fold reduction in biofilm ratio were considered to be attenuated for their biofilm-forming capacity.

Mice infection analysis

For animal infection studies, overnight-grown *Cg* strains in YPD medium at 30°C were collected, washed twice with sterile PBS and suspended in PBS to a final cell density of 20 OD_{600}. 100 µl of cell suspension was injected into the tail vein of 6-8 week-old female BALB/c mice. At 7th day post-infection, mice were euthanized using CO_2 and three organs (kidneys, liver and spleen) were excised out. The harvested organs were homogenized in PBS, and homogenates were appropriately diluted and plated on YPD medium supplemented with penicillin and streptomycin antibiotics. After 1-2 days of incubation at 30°C, *Cg* colonies were counted. The fungal burden in each organ was calculated by multiplying the colony number with the appropriate dilution factor.

Cg-**macrophage interaction analysis**

Human THP-1 monocytic cells (1 x 10^6 cells in each well of a 24-well tissue culture plate) were treated with phorbol 12-myristate 13-acetate (16 nM) for 12 h for differentiation into macrophages at 37°C with 5% CO_2, followed by 12 h cell recovery in RPMI-10% FBS medium. THP-1 macrophages were infected with overnight, YPD medium-grown *Cg* cells at a multiplicity of infection (MoI) of 0.1. Non-phagocytosed *Cg* cells were removed after 2 h, and the infection was continued for another 22 h. At 2 h and 24 h post-infection, extracellular *Cg* cells were removed with PBS, THP-1 cells were lysed in water and lysates were plated on YPD medium. After 2 days of growth at 30°C, *Cg* colonies were counted and the fold replication of *Cg* in THP-1 cells was calculated by dividing 24 h CFUs by 2 h CFU counts. % phagocytosis was determined by dividing 2 h CFUs by 0 h CFUs (Number of *Cg* cells infected to THP-1 macrophages). Mutants showing ≥ 2-fold change in fold replication, as compared to *wt* cells, were considered to have altered survival in macrophages.

Secreted cytokines were measured by enzyme-linked immunosorbent assay (ELISA). For this, THP-1 macrophages were infected with *Cg* at a MoI of 1.0 for 24 h, as described above, and culture supernatants were collected and centrifuged at 1000 rpm to remove any cell debris. IL-1β, IL-8 and TNF-α in cleared supernatants were measured using the ELISA kits following manufacturer's protocol. For inhibitor treatment, PMA-treated THP-1 macrophages were pre-treated with inhibitors or DMSO solvent for 2 h, prior to *Cg* infection, and *Cg*-THP1 co-culture was carried out in the presence of inhibitor or DMSO for 24 h.

For isolation of primary macrophages, 6-8-week-old BALB/c mice were intraperitoneally injected with 1.5 ml of 4% (w/v) thioglycollate. After 5 days, mice were euthanized, and macrophages were collected in DMEM from the mouse peritoneal cavity. Macrophages were centrifuged, suspended in DMEM medium containing 10% heat-inactivated FBS and were seeded in a 24-well plate. *Cg* survival and secreted IL-1β in primary mouse peritoneal macrophages were measured in a similar way, as that for THP-1 cells.

Nucleosome dynamics analysis by MNase-Seq

For MNase-seq, PMA-activated THP-1 (2.2 x 10^7) cells were infected with overnight YPD medium-grown *wt* cells at a MOI of 1:1, and macrophage-internalized *Cg* cells were collected

post 2 h and 10 h infection, after lysing infected macrophages in ice-cold water. Macrophage lysates were gently vortexed to separate *Cg* cells form macrophage debris, followed by centrifugation at 6000 rpm for 5 min at 4°C. The compact *Cg* cell pellet was collected and suspended in ice-cold water, followed by centrifugation at 10,000 rpm for 5 min at 4°C. This step was repeated 3-4 times to obtain *Cg* cell pellet that was devoid of macrophage debris. As a control, overnight YPD medium-grown *wt* cells were grown in RPMI-10% FBS medium for 2 h and 10 h, and cell pellets were collected. After PBS washes, 2 h and 10 h RPMI-grown and macrophage-internalized *wt* cell pellets were suspended in MNase-digestion buffer [10 mM Tris-Cl (pH 8.0) and 1 mM $CaCl_2$], and lysed using glass beads. After lysate centrifugation at 13000 rpm at 4°C for 10 min, lysates (1 mg) were incubated with MNase (10 units/1 mg lysate; NEB #M0247S) at 37°C for 60 min. The MNase digestion was stopped by adding Stop Buffer (8.6% SDS and 0.007 M EDTA), followed by digestion with proteinase K (20 mg/ml) first at room temperature for 30 min, and later at 65°C for overnight. DNA was isolated with PCI (Phenol:Chloroform:Isoamyl alcohol) extraction, precipitated with sodium acetate and ethanol, and was suspended in nuclease-free water, followed by RNase A digestion for 30 min at 37°C. DNA was run on 2% agarose gel, and bands of ~ 150 bp, corresponding to mononucleosomal DNA fragments, were excised, purified using QIAquick extraction kit, and were sent to Scigenom, Kochi, India (http://www.scigenom.com/) for library preparation and sequencing. The concentration and quality of purified mononucleosomal DNA was determined by Qubit, Nanodrop and Agilent Tapestation analysis, followed by library generation using the NEBNext Ultra II DNA Library Prep Kit (Illumina), following the manufacturer's instructions. The library quality was assessed using the Agilent Bioanalyzer High Sensitivity DNA chip.

The prepared libraries were sequenced (2x100 bp paired-end sequencing) on the HiSeq 2500 (Illumina) platform, and 40-60 million, high-quality 100 bp reads, with 93 % of the total reads passing ≥30 phread score, were generated for each sample. Data were analysed by DeepSeeq Bioinformatics, Bengaluru, India (https://www.deepseeq.com/). The adapter sequences and low-quality reads were removed using Trimmomatic (version 0.39). Trimmed sequence reads were aligned to the *C. glabrata* genome version s04-m01-r06 (www.candidagenome.org) using bowtie2, version 2.5.0. The percentage alignment varied between RPMI-grown (89-97%) and macrophage-internalized (6-26%) samples. The SAM files generated by bowtie2 were converted to BAM format, sorted and indexed using Samtools, version 1.7. The DANPOS3 software[20] was used to identify the occupancy, position and fuzziness of nucleosomes. For this, clonal reads from BAM files were first removed, followed by quantile normalization. Next, individual dpos, dpeak, and dregion functions from DANPOS3 were used to identify positions, peaks, and regions occupied by nucleosomes in individual samples as well as for compared datasets. Annotation of the called nucleosome position was done using gene start and stop coordinates for 5604 genes, obtained from the C_glabrata_CBS138_version_current_features.gff file version S04-m01-r06 downloaded from http://www.candidagenome.org/download/gff/C_glabrata_CBS138/, by employing the closest function from the Bedtools software version v2.28.0. The –flank_up and –flank_down parameters were set to 500 and 1000, respectively, for TSS (Transcription Start Site) -based analysis, and to 500 and 500, respectively, for TTS (Transcription Termination Site)-based analysis. Nucleosomes were identified in about 5300 genes in all four-studied conditions. Lastly, three filters, Position shift: ≥ 50 bp shift, Occupancy change: ≥ 2-fold change (FDR ≤

0.05) or Fuzziness change: ≥ 1.5-fold change (FDR ≤ 0.05), were applied to identify dynamic nucleosomes.

Immunoblotting analysis

For signalling pathway analysis, PMA-activated macrophages were either left uninfected or infected with *wt* and *Cgsnf2Δ* cells at a MOI of 1:1. After 4 h infection, macrophages were washed with ice-cold PBS, scrapped and centrifuged at 2000 rpm for 5 min. Cell pellets were suspended in freshly-prepared NETN lysis buffer [250 mM NaCl, 5 mM EDTA (pH 8.0), 50 mM Tris-Cl (pH 8.0), and 0.5% Nonidet P-40] containing protease (cOmplete Mini, Roche) and phosphatase (PhosSTOP, Roche) inhibitors, and incubated for 30 min at 4°C. Cell lysates were sonicated for 15 cycles (30 sec ON/30 sec OFF), with the Diagenode bioruptor sonicator and centrifuged at 13000 rpm for 10 min. 120 µg proteins were resolved on 10% SDS-PAGE, and probed with different antibodies. For quantification, intensities of the bands of interest were quantified from independent immunoblots using the ImageJ densitometry software, and normalized with respective GAPDH signal intensities. Values were plotted as a bar graph, considering the signal in control samples as 1.0. For *Cg* protein analysis, log-phase *Cg* cells were lysed using the glass beads, and lysates (80 µg) were resolved on 10% SDS-PAGE, followed by probing with anti-FLAG antibody.

Cg cell wall analysis

For quantification of major cell wall components, overnight YPD medium-grown *wt* and *Cgsnf2Δ* mutant strains were inoculated in fresh YPD medium at 0.1 OD_{600} and incubated at 30°C for 4 h. Log-phase cells (2.0 OD_{600}) were collected, washed and suspended in PBS. Cells were either left untreated or stained with calcofluor white (2.5 µg/ml), fluorescein isothiocyanate-labeled concanavalin A (1 µg/ml) or aniline blue (12.5 µg/ml) for 15 min at room temperature for measurement of chitin, mannan and β-glucan, respectively. After PBS washes, fluorescence intensity of ~50,000 cells was recorded on BD FACS ARIA III flow cytometer at an excitation wavelength of 355 nm and an emission wavelength of 433 nm. All data were analysed with the FlowJo software, and the background fluorescence was corrected by subtracting the mean intensity fluorescence value of unstained samples from that of respective stained samples. Data were presented as the mean intensity ratio which indicates *Cgsnf2Δ*/*wt* fluorescence intensity values.

RNA-sequencing analysis

2 h and 10 h RPMI-medium and macrophage-internalized *wt* and *Cgsnf2Δ* cells were harvested and total RNA was extracted using RNeasy kit (Qiagen), followed by removal of DNA contamination, if any, using DNase I. The purified total RNA was sent to National Genomics Core (NGC) at CDFD, Hyderabad (http://ngc.cdfd.org.in/) for sequencing, which involved mRNA enrichment using Poly(A) mRNA Magnetic Isolation Module, library preparation using NEB Ultra II Directional RNA Library Prep Kit and 150 bp paired-end sequencing on the Illumina NextSeq 2000 platform. Sequenced reads were processed and mapped on to the *C. glabrata* CBS138 reference genome using HISAT 2.1.0 aligner.

For gene expression quantification, the counts of mapped reads for each gene were considered using the Feature counts tool, followed by sequencing depth normalization of raw read counts using the DESeq2 tool. Differentially expressed genes were classified based on two criteria: ≥ 2-fold change in expression and adjusted p-value of ≤ 0.05.

Imaging analysis

For morphology analysis, log-phase *Cg* cells were visualized using the LSM700 confocal microscope equipped with 63X/1.44 NA objective. For cell wall chitin staining, log-phase *Cg* cells were stained with calcofluor white (1.25 μg/ml) for 30 min at room temperature. After PBS washes, cells were imaged using LSM700 confocal microscope equipped with 63X/1.44 NA objective. All images were processed using the ZEN blue software.

For immunofluorescence analysis, J774.1 murine macrophage-like cells (1 x 10^6) were seeded on a coverslip, and infected with *wt* and *Cgsnf2Δ* cells at a MoI of 1:1. At 2 h post-infection, un-phagocytosed *Cg* cells were removed, followed by 10% FBS-containing DMEM medium addition and incubation for 2 h at 37°C, 5% CO_2. As a control, uninfected J774A.1 cells were grown under similar conditions. J774. 1 cells were washed with ice-cold PBS and fixed with 3.7% (v/v) formaldehyde for 15 min. After fixation, cells were permeabilized with 0.5% (v/v) Triton X-100 in PBS for 5 min, washed twice with PBS and blocked with 10% (v/v) FBS in PBS for 1 h at room temperature. Cells were probed overnight with anti-NF-κB p65 antibody at 4°C, followed by incubation with AlexaFluor 568-conjugated secondary antibody for 1 h. Coverslips were washed, air-dried and mounted in the VECTASHIELD anti-fade mounting medium containing DAPI. Cells were visualized, and Z-stack images were acquired throughout the nucleus at 1 μm interval using the Leica confocal microscope (63X/1.44 NA objective), and processed using the LAS X software. Each maximum-intensity projection (MIP) image was constructed using 8-12 confocal image subsets.

Quantitative RT-PCR (qRT-PCR) analysis

RNA was isolated from 2 h and 10 h RPMI-grown and macrophage-internalized *Cg* cells using the RNeasy kit (Qiagen), followed by DNase I digestion. 1 μg DNase I-treated total RNA was used for cDNA synthesis using the Superscript III reverse transcriptase, DyNAmo ColorFlash SYBR Green qPCR kit was used to perform qRT-PCR, and gene expression was determined by the 2$^{-ΔΔCT}$ method. *GAPDH/ACT1* expression was used as an internal reference control to normalize the gene expression data.

Chromatin immunoprecipitation (ChIP) analysis

ChIP was performed, as described previously[42]. Briefly, *CgSNF2-SFB*-expressing *Cgsnf2Δ* was grown in CAA medium for 4 h, followed by formaldehyde crosslinking. Next, cells were lysed in buffer containing 1 mM EDTA [pH 8.0], 50 mM HEPES [pH 7.5], 0.1% sodium deoxycholate [w/v], 140 mM NaCl, 1× protease inhibitor cocktail, and 1% Triton X-100 by bead-beating, and sonicated for 40 min [30 s pulses of on and off] to obtain 250-500 bp small chromatin fragments. A fraction of the clear supernatant was used as Input fraction, while the remainder was subjected to immunoprecipitation with anti-IgG or anti-FLAG antibody. After de-crosslinking, DNA was purified using phenol:chloroform:isoamyl alcohol, and used as template for qRT-PCR.

Functional enrichment and statistical analysis

The David tool (https://david.ncifcrf.gov/) with default settings was used for enrichment of gene ontology (GO) terms for biological process, cellular component and molecular function, in genes with dynamic nucleosomes at their promoters and in differentially-expressed genes. Terms with the q-value of ≤ 0.05 were considered as enriched in gene datasets. Statistical significance was determined using the two-tailed Student t-test or the nonparametric Mann-Whitney test in the GraphPad Prism software. Error bars indicate standard error of the mean

(s.e.m). The multiple amino acid sequence alignment was done using the Clustal W tool. Asterisks were used to represent p-values: * $p \leq 0.05$, ** $p \leq 0.01$; *** $p \leq 0.005$; **** $p \leq 0.001$. The p value of ≤ 0.05 was considered significant.

Data availability

The raw MNase-seq and RNA-seq data will be deposited in the NCBI's Gene Expression Omnibus (GEO).

Acknowledgements

We would like to thank Dr. Pranjali Pore and staff of the Experimental Animal Facility of CDFD for their help with mice experiments. We are indebted to Vandana Sharma and Romila Moirangthem for her help with MNase-Seq experiments and *C. glabrata* deletion strain generation, respectively.

Funding

This work was supported by the DBT/Wellcome Trust India Alliance Senior Fellowship to RK [IA/S/15/1/501831; www.indiaalliance.org/], and grants from the Department of Biotechnology [BT/PR40336/BRB/10/1921/2020 and BT/PR42015/MED/29/1561/2021; www.dbtindia.gov.in/], and Science and Engineering Research Board, Department of Science and Technology [CRG/2021/000530; www.serb.gov.in/], Government of India, to RK. KK was a recipient of the Shyama Prasad Mukherjee Fellowship of the Council of Scientific and Industrial Research, New Delhi, India [www.csirhrdg.res.in/]. AP is a recipient of the INSPIRE fellowship of the Department of Science and Technology, New Delhi, India (https://online-inspire.gov.in/). The funders had no role in study design, data collection and analysis, or decision to publish or preparation of the manuscript.

Author contributions

KK conceived the study. KK and RK designed the study. KK and AP performed experiments and acquired data. KK, AP and RK analysed data. KK prepared tables and figures. KK and RK wrote the manuscript with input from AP.

Competing interests

The authors declare no competing interests.

Additional information

This manuscript contains extended data (Figures 1 to 10).

Supplementary information

This manuscript contains supplementary data (Supplementary Figures 1 and 2, and Tables 1 to 10).

References

1.   Bongomin, F., Gago, S., Oladele, R. & Denning, D. Global and Multi-National Prevalence of Fungal Diseases—Estimate Precision. *J. Fungi* **3**, 57 (2017).

2.	Chakrabarti, A. Candida glabrata candidemia. *Indian J. Crit. Care Med.* **19**, 138–139 (2015).

3.	Pfaller, M. A., Diekema, D. J., Turnidge, J. D., Castanheira, M. & Jones, R. N. Twenty Years of the SENTRY Antifungal Surveillance Program: Results for Candida Species From 1997-2016. *Open forum Infect. Dis.* **6**, S79–S94 (2019).

4.	Rasheed, M., Battu, A. & Kaur, R. Host–pathogen interaction in Candida glabrata infection: current knowledge and implications for antifungal therapy. *Expert Rev. Anti. Infect. Ther.* **18**, 1093–1103 (2020).

5.	Wisplinghoff, H. *et al.* Nosocomial Bloodstream Infections in US Hospitals : Analysis of 24 , 179 Cases from a Prospective Nationwide Surveillance Study. **0019**, 309–317 (2004).

6.	Gupta, A., Varma, A. & Gupta, A. Candida glabrata candidemia : An emerging threat in critically ill patients. *Indian J. Crit. Care Med.* **19**, 151–154 (2015).

7.	Meyahnwi, D., Siraw, B. B. & Reingold, A. Epidemiologic features , clinical characteristics , and predictors of mortality in patients with candidemia in Alameda County , California ; a 2017 – 2020 retrospective analysis. 1–10 (2022).

8.	*WHO fungal priority pathogens list to guide research, development and public health action*.

9.	Gabaldón, T. *et al.* Comparative genomics of emerging pathogens in the Candida glabrata clade. (2013).

10.	Kumar, K., Askari, F., Sahu, M. S. & Kaur, R. Candida glabrata: A lot more than meets the eye. *Microorganisms* **7**, 1–22 (2019).

11.	Riedelberger, M. *et al.* Type I Interferon Response Dysregulates Host Iron Homeostasis and Enhances Candida glabrata Infection. *Cell Host Microbe* **27**, 454-466.e8 (2020).

12.	Riedelberger, M. *et al.* Type I Interferons Ameliorate Zinc Intoxication of Candida glabrata by Macrophages and Promote Fungal Immune Evasion. *iScience* **23**, 101121 (2020).

13.	Rasheed, M., Battu, A. & Kaur, R. Aspartyl proteases in Candida glabrata are required for suppression of the host innate immune response. *J. Biol. Chem.* **293**, 6410–6433 (2018).

14.	Kaur, R., Ma, B. & Cormack, B. P. A family of glycosylphosphatidylinositol-linked aspartyl proteases is required for virulence of Candida glabrata. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 7628–33 (2007).

15.	Rai, M. N., Balusu, S., Gorityala, N., Dandu, L. & Kaur, R. Functional genomic analysis of Candida glabrata-macrophage interaction: role of chromatin remodeling in virulence. *PLoS Pathog.* **8**, e1002863 (2012).

16.	Struhl, K. & Segal, E. Determinants of nucleosome positioning. *Nat. Struct. Mol. Biol.* **20**, 267–273 (2013).

17.	Clapier, C. R., Iwasa, J., Cairns, B. R. & Peterson, C. L. Mechanisms of action and regulation of ATP-dependent chromatin-remodelling complexes. *Nat. Rev. Mol. Cell Biol.* **18**, 407–422 (2017).

18.	Tebbji, F., Chen, Y., Sellam, A. & Whiteway, M. The Genomic Landscape of the Fungus-Specific SWI/SNF Complex Subunit, Snf6, in Candida albicans. *mSphere* **2**, (2017).

19.	Nikolov, V. N., Malavia, D. & Kubota, T. SWI/SNF and the histone chaperone Rtt106 drive expression of the Pleiotropic Drug Resistance network genes. *Nat. Commun.* **13**, 1968 (2022).

20.	Chen, K. *et al.* DANPOS: Dynamic analysis of nucleosome position and occupancy by sequencing. *Genome Res.* **23**, 341–351 (2013).

21.	Tsankov, A. M., Thompson, D. A., Socha, A., Regev, A. & Rando, O. J. The Role of

Nucleosome Positioning in the Evolution of Gene Regulation. **8**, (2010).

22. Riera, M., Mogensen, E., d'Enfert, C. & Janbon, G. New regulators of biofilm development in Candida glabrata. *Res. Microbiol.* **163**, 297–307 (2012).

23. Juárez-Reyes, A. & Castaño, I. Chromatin architecture and virulence-related gene expression in eukaryotic microbial pathogens. *Curr. Genet.* **65**, 435–443 (2019).

24. Xu, Z. *et al.* De novo genome assembly of Candida glabrata reveals cell wall protein complement and structure of dispersed tandem repeat arrays. 1209–1224 (2020). doi:10.1111/mmi.14488

25. López-Fuentes, E. *et al.* Candida glabrata's genome plasticity confers a unique pattern of expressed cell wall proteins. *J. Fungi* **4**, (2018).

26. Jawhara, S. *et al.* Murine Model of Dextran Sulfate Sodium-induced Colitis Reveals Candida glabrata Virulence and Contribution of β-Mannosyltransferases. *J. Biol. Chem.* **287**, 11313–11324 (2012).

27. Cormack, B. P., Ghori, N. & Falkow, S. An adhesin of the yeast pathogen Candida glabrata mediating adherence to human epithelial cells. *Science* **285**, 578–82 (1999).

28. Kuhn, D. M. & Vyas, V. K. The Candida glabrata adhesin Epa1p causes adhesion, phagocytosis, and cytokine secretion by innate immune cells. *FEMS Yeast Res.* **12**, 398–414 (2012).

29. Patin, E. C., Thompson, A. & Orr, S. J. Pattern recognition receptors in fungal immunity. *Semin. Cell Dev. Biol.* **89**, 24–33 (2019).

30. Fidel, P. L., Vazquez, J. A. & Sobel, J. D. Candida glabrata: Review of epidemiology, pathogenesis, and clinical disease with comparison to C. albicans. *Clin. Microbiol. Rev.* **12**, 80–96 (1999).

31. Erwig, L. P. & Gow, N. A. R. Interactions of fungal pathogens with phagocytes. *Nat. Rev. Microbiol.* **14**, 163–176 (2016).

32. Frieman, M. B., McCaffery, J. M. & Cormack, B. P. Modular domain structure in the Candida glabrata adhesin Epa1p, a β1,6 glucan-cross-linked cell wall protein. *Mol. Microbiol.* **46**, 479–492 (2002).

33. Timmermans, B., Peñas, A. D. Las & Castaño, I. Adhesins in Candida glabrata. *Fungi J.* 1–16 (2018). doi:10.3390/jof4020060

34. Zupancic, M. L. *et al.* Glycan microarray analysis of Candida glabrata adhesin ligand specificity. *Mol. Microbiol.* **68**, 547–559 (2008).

35. Vale-Silva, L. A. *et al.* Upregulation of the Adhesin Gene EPA1 Mediated by PDR1 in Candida glabrata Leads to Enhanced Host Colonization. *mSphere* **1**, 1–16 (2016).

36. Vitenshtein, A. *et al.* NK Cell Recognition of Candida glabrata through Binding of NKp46 and NCR1 to Fungal Ligands Epa1, Epa6, and Epa7. *Cell Host Microbe* **20**, 527–534 (2016).

37. Domergue, R. *et al.* Nicotinic acid limitation regulates silencing of Candida adhesins during UTI. *Science* **308**, 866–70 (2005).

38. Dagher, Z. *et al.* Fluorescent Tracking of Yeast Division Clarifies the Essential Role of Spleen Tyrosine Kinase in the Intracellular Control of Candida glabrata in Macrophages. **9**, (2018).

39. Cottier, F. *et al.* Remasking of Candida albicans β-Glucan in Response to Environmental pH Is Regulated by Quorum Sensing. *MBio* **10**, (2019).

40. Rai, M. N. *et al.* Temporal transcriptional response of *Candida glabrata* during macrophage infection reveals a multifaceted transcriptional regulator CgXbp1 important for macrophage response and drug resistance. *bioRxiv* (2021). doi:10.1101/2021.09.28.462173

41. Kumar, K., Moirangthem, R. & Kaur, R. Histone H4 dosage modulates DNA damage response in the pathogenic yeast Candida glabrata via homologous recombination

pathway. *PLoS Genet.* **16**, e1008620 (2020).

42.   Bhakt, P., Raney, M. & Kaur, R. The SET-domain protein CgSet4 negatively regulates antifungal drug resistance via the ergosterol biosynthesis transcriptional regulator CgUpc2a. *J. Biol. Chem.* **298**, 1–22 (2022).

Figure legends

Figure 1: Genome-wide nucleosome dynamics in response to the macrophage milieu. a, Experimental design diagram for nucleosome mapping in *Cg wild-type* cells that were grown either in RPMI medium or infected to THP-1 macrophages for 2 h and 10 h. **b,** Total nucleosomes identified using the dpos function of DANPOS3 tool with default parameters. R2: 2 h RPMI-grown, R10: 10 h RPMI-grown, M2: 2 h macrophage-internalized and M10: 10 h macrophage-internalized *Cg* cells. **c,** Dynamic nucleosomes identified in indicated compared datasets. **d,** Venn diagram showing the overlap in the number of genes that exhibited position shift, occupancy and fuzziness changes at their promoter regions in M2 versus R2 comparison. **e,** Chromosome-wise distribution of dynamic nucleosomes in M2 versus R2 comparison. The colour scale indicates nucleosome density. **f,** Gene Ontology (GO) terms for biological process (BP), cellular component (CC) and molecular function (MF) enriched in genes with dynamic nucleosomes at their promoters in M2 versus R2 comparison.

Figure 2: CgSnf2 is essential for intracellular survival and virulence. a, Heatmap showing growth profiles of *Cg* strains. The colour code depicts differences between mutant and *wt* growth under the same condition. **b,** Biofilm formation on polystyrene. Data represent mean ± SEM (n = 3-4). Paired two-tailed Student's t-test. **c,** CFU-based measurement of *Cg* survival in human THP-1 macrophages. Data (mean ± SEM; n =4) represent fold replication at 24 h. Unpaired two-tailed Student's t-test. **d,** *Cg* survival in the murine model of systemic candidiasis. Diamonds and horizontal line represent fungal CFUs recovered from each mouse, and CFU geometric mean (n = 6-9) for each organ, respectively. Mann-Whitney test.

Figure 3: CgSnf2 suppresses host immune cell activation. a, Secreted IL-1β levels by uninfected (UI) and *Cg*-infected THP-1 macrophages. Data represent mean ± SEM (n = 6). Unpaired two-tailed Student's t-test. Black and red asterisks indicate statistically significant differences in IL-1β, compared to UI and *wt*-infected THP-1 cells, respectively. **b,** IL-1β secretion in uninfected or *Cg*-infected THP-1 macrophages that were treated with DMSO (solvent control) or NLRP3 inflammasome inhibitor (MCC950; 15 μM). Data represent mean ± SEM (n = 4). Unpaired two-tailed Student's t-test. Black and green asterisks represent statistically significant differences in IL-1β, compared to DMSO-treated uninfected and inhibitor-treated uninfected macrophages, respectively. **c and d,** *Cg* survival measurement in DMSO-, 15 μM MCC950 (**c**) or 5 μM R406 (Syk inhibitor) (**d**)-treated THP-1 macrophages. Data (mean ± SEM; n = 3-4) represent fold replication at 24 h. Unpaired two-tailed Student's t-test. **e,** *Cg* survival measurement in murine peritoneal macrophages. Data (mean ± SEM; n =3) represent fold replication at 24 h. Unpaired two-tailed Student's t-test. **f,** IL-1β secretion in uninfected or *Cg*-infected murine peritoneal macrophages that were treated with DMSO or

15 µM MCC950. Data represent mean ± SEM (n = 3). Unpaired two-tailed Student's t-test. Black asterisks represent statistically significant differences in IL-1β, compared to DMSO-treated uninfected macrophages.

Figure 4: CgSnf2 is required for *Cg* transcriptional response to the macrophage environment. a, Volcano plots showing mean fold change of differentially-expressed genes (DEGs) and their adjusted p (false discovery rate-corrected) values. Green, red and black dots indicate upregulated, downregulated and non-DEGs, respectively. **b,** Venn diagrams showing overlap in DEGs among four compared datasets. **c,** Venn diagram illustrating overlap in CgSnf2-regulated genes in 2 h RPMI-grown or 2 h macrophage-internalized *Cg*. Of 1410 DEGs in macrophage-internalized *wt*, expression of 1269 (90%) and 61 (4.3%) genes are dependent and independent of CgSnf2, respectively, with CgSnf2 regulating basal expression of 680 (48%) genes. Additionally, induction of 80 *Cg* genes (5.7%) in macrophages require CgSnf2. **d and e,** Bubble plots showing enriched GO-BP terms for upregulated and downregulated genes in 2 h RPMI-grown *Cgsnf2Δ* (**d**) and 2 h macrophage-internalized *Cgsnf2Δ* (**e**), compared to 2 h RPMI-grown *wild-type* and 2 h RPMI-grown *Cgsnf2Δ*, respectively.

Figure 5: *EPA1* expression is deleterious for *Cg* survival in macrophages. a, Heatmap showing differential expression of 21 adhesin genes. Adhesin genes encoded at subtelomeric regions (25 kb from the chromosome ends) are marked in blue colour. R2 and M2 refer to *Cg* grown in RPMI medium and infected to THP-1 macrophages for 2 h, respectively. **b,** Integrative genome viewer (IGV) snapshot of RNA-seq signal at *EPA1* locus (ChrE: 682420 to 685524 bp). All IGV tracks have the same scaling factor [0-750] for the Y-axis. **c,** qPCR-based analysis of *EPA1* expression after 2 h growth in RPMI medium or macrophage internalization. Data mean ± SEM (n = 3) were normalized with *ACT1* mRNA control, and plotted as fold change in gene expression, compared to RPMI-grown *wt* (considered as 1.0). Paired two-tailed Student's t-test. **d,** *Cg* replication in THP-1 macrophages. Data represent mean ± SEM (n = 5-7). Unpaired two-tailed Student's t-test. **e,** IL-1β secretion in uninfected (UI) and *Cg*-infected THP-1 macrophages. *Cg* strains were either untreated- or treated with 10 mM lactose, 1 h prior to THP-1 infection, and the infection was continued for 24 h in the presence of lactose. Data represent mean ± SEM (n = 4). Unpaired two-tailed Student's t-test. **f and g,** IL-1β secretion in uninfected (UI) and *Cg*-infected THP-1 macrophages. Data represent mean ± SEM (n = 4-6). Unpaired two-tailed Student's t-test. **h,** *Cg* survival in THP-1 macrophages. Data represent mean ± SEM (n = 3-4). Unpaired two-tailed Student's t-test. **i,** ChIP-qPCR quantification of the level of bound, ectopically expressed SFB-tagged CgSnf2 to *EPA1* promoter. Y-axis label is ChIP % input, with immunoglobulin G (IgG)-control and anti-FLAG (CgSnf2) antibodies. Data represent mean ± SD (n = 2). Unpaired two-tailed Student's t-test.

Figure 6: *EPA1* expression governs nuclear localization of NF-κB. a, Representative western blots of phosphorylated p65 expression. Bar graphs show fold change in phosphorylation, normalized to GAPDH signal. Data represent mean ± SEM (n = 4). Paired

two-tailed Student's t-test. **b,** IL-1β in uninfected (UI) and *Cg*-infected THP-1 macrophages which were treated with DMSO or NF-κB inhibitor (BAY 11-7082; 10 μM). Unpaired two-tailed Student's t-test. Black and green asterisks represent statistically significant differences in IL-1β levels, compared to DMSO-treated uninfected and BAY 11-7082-treated uninfected macrophages, respectively. **c and d,** TNF-α **(c)** and IL-8 **(d)** secretion in uninfected (UI) and *Cg*-infected THP-1 macrophages which were treated with DMSO, BAY 11-7082 (10 μM), NLRP3 inhibitor (MCC950; 15 μM), Akt inhibitor (SH-6; 5 μM) or p38 MAPK inhibitor (SB203580; 10 μM). Data represent mean ± SEM (n = 3-4). Unpaired two-tailed Student's t-test. Black and red asterisks represent statistically significant differences in TNF-α and IL-8 levels, compared to DMSO-treated uninfected and the corresponding inhibitor-treated uninfected macrophages, respectively. **e,** Confocal images showing cellular localization of p65 in uninfected or *Cg*-infected J774A.1 cells, as detected with anti-p65 antibody. DAPI was used to stain macrophage nuclei. Bar = 10 μm. The p65 signal was quantified using ImageJ, and data represent (mean ± SEM; n = 30-50 cells) ratio of nuclear to cytosolic p65. **f,** IL-1β secretion in DMSO or BAY 11-7082 (10 μM)-treated uninfected (UI) THP-1 and THP-1 cells infected with four *Candida* species, *C. albicans*, *C. auris*, *C. glabrata* and *C. tropicalis* at a MoI of 5:1 for 4 h. Data represent mean ± SEM (n = 3-4). Unpaired two-tailed Student's t-test. Black and red asterisks represent statistically significant differences in IL-1β levels, compared to DMSO-treated uninfected and respective DMSO-treated *Candida spp*-infected THP-1 macrophages, respectively.

Extended data: Extended figures 1-10.

Extended Data Figure Legends
Extended Data Figure 1: Nucleosome dynamics in macrophage-internalized *Cg*. a, Percentage of fuzzy nucleosomes. R2: 2 h RPMI-grown, R10: 10 h RPMI-grown, M2: 2 h macrophage-internalized and M10: 10 h macrophage-internalized *Cg*. **b,** A schematic of a genomic locus divided into three regions for nucleosome mapping: Promoter [0-500 bp upstream of the Transcription Start Site (TSS)], intragenic (gene-encoding sequence) and intergenic (region between the stop codon of one gene and the promoter or the stop codon of the next gene). The arrow indicates the direction of transcription. **c,** Nucleosome distribution across three genomic regions. **d,** Dynamic nucleosomes across three genomic regions. **e,** Dynamic nucleosomes in gene promoters. The exact number is indicated in the upper edge of the bar. **f,** Percentage of four types of nucleosome occupancy dynamics across three genomic regions.

Extended Data Figure 2: Genes involved in host adhesion contain dynamic nucleosomes at their promoters. a-c, Venn diagrams illustrating overlap in the number of genes which showed nucleosome position shift, occupancy and fuzziness changes at their promoters in M10 versus R10 **(a)**, M10 versus M2 **(b)** and R10 versus R2 **(c)** comparison. **d-f,** GO terms for biological process (BP), cellular component (CC) and molecular function (MF) enriched in genes with dynamic nucleosomes at their promoters in M10 versus M2 **(d)**, M10 versus R10 **(e)** and R10 versus R2 **(f)** comparison.

Extended Data Figure 3: *CgSNF2* deletion leads to growth retardation. a, ATPase subunits of seven ATP-dependent chromatin remodelling complexes in *C. glabrata*, as identified by BLAST analysis of their *S. cerevisiae* orthologs. **b,** Time course analysis of *Cg* growth in YPD medium. **c,** Generation time, calculated during the logarithmic-phase (between 2 and 8 h) of the growth curve shown in *b* panel. Data represent mean ± SEM (n = 3). Unpaired two-tailed Student's *t* test. **d,** Representative confocal images of YPD-grown, log-phase *Cg* cells. Scale bar = 5 μm. **e,** % phagocytosis of *Cg* strains by THP-1 macrophages. Data represent mean ± SEM (n = 4). **f,** CFU-based growth analysis of *Cg* strains in RPMI medium containing 10% FBS at 37ºC, 5% CO_2. At indicated time points, appropriate dilutions of RPMI-grown cultures were plated on YPD medium. After incubation at 30ºC for 2-3 days, colonies were counted and plotted. Data represent mean ± SEM (n = 3-4).

Extended Data Figure 4: *CgSNF2* complements virulence defect of *Cgsnf2Δ*. a, Serial dilution spotting assay showing *Cg* growth. Bleomycin (Bleo), methyl methanesulfonate (MMS), hydroxyurea (HU), β -mercaptoethanol (β-ME), menadione (MD), hydrogen peroxide (H_2O_2), calcium chloride ($CaCl_2$), zinc chloride ($ZnCl_2$), manganese chloride ($MnCl_2$), copper sulphate ($CuSO_4$), congo red (CR), caffeine and sodium dodecyl sulfate (SDS) were used at 5 μM, 0.03%, 25 mM, 20 mM, 100 μM, 25 mM, 500 mM, 8 mM, 3 mM, 8 mM, 2 mg/ml, 10 mM and 0.05% concentrations, respectively. **b,** Representative confocal images of log-phase CAA medium-grown, calcofluor white-stained *Cg* cells. Scale bar = 2 μm. **c,** CFU-based intracellular survival analysis of *Cg* strains in THP-1 macrophages. Data represent mean ± SEM (n = 3). Unpaired two-tailed Student's t-test. **d,** CFU-based analysis of organ fungal burden in the murine model of systemic candidiasis. Diamonds and horizontal line represent CFUs recovered from each mouse and geometric CFU mean (n = 8-10) for each organ, respectively. Mann-Whitney test. CFU values plotted for *wt* and *Cgsnf2Δ* strains are the same, as shown in Figure 2d.

Extended Data Figure 5: CgSnf2 regulates gene expression. a and b, Volcano plots showing mean fold change of differentially-expressed genes (DEGs) and their adjusted p (false discovery rate-corrected) values in 10 h **(a)** and 10 h vs 2 h comparison **(b)**. Green, red and black dots indicate upregulated, downregulated and non- DEGs, respectively. **c,** Venn diagrams showing overlap in DEGs among four compared datasets. **d,** Heatmap showing enriched GO-BP terms in four compared datasets. **e,** Bar graph showing mannan, glucan and chitin amounts in *Cgsnf2Δ*, relative to *wt*. Data represent mean fluorescence intensity ratios ± SEM (n = 4). Paired two-tailed Student t test.

Extended Data Figure 6: CgSnf2-dependent activation of a 7-mannosyltransferase gene cluster is pivotal to supress host activation. a, Heatmap showing normalized read counts (presented as z-score) of 35 mannosyltransferase (MT)-encoding genes in 2 h RPMI-grown (R2) and macrophage-internalized (M2) *wt* and *Cgsnf2Δ*. **b,** JBrowse snapshot (www.candidagenome.org) of the seven (five-β and two-α) mannosyltransferase genes present in tandem on the Chromosome B. **c,** Integrative genome viewer (IGV) snapshot of RNA-seq signal at *MT* loci (ChrB: 282186 to 300029 bp) in *wild-type* and *Cgsnf2Δ* grown either in

RPMI-medium (R2) or incubated with THP-1 macrophages (M2) for 2 h. All IGV tracks have the same scaling factor [0-500] for the Y-axis. **d,** qPCR-based analysis of *CgBMT2* expression in 2 h macrophage-internalized *wt* and *Cgsnf2Δ*. Data (mean ± SEM; n = 3) were normalized with *ACT1* mRNA control, and represent fold change in gene expression, compared to RPMI-grown *wt* (considered as 1). Paired two-tailed Student's t-test. **e,** *Cg* intracellular replication analysis. The *CgMT-CΔ* lacks ~19 kb genomic region containing seven mannosyltransferase genes. Data represent mean ± SEM (n = 4). Unpaired two-tailed Student's t-test. **f,** IL-1β secretion in THP-1 macrophages. Data represent mean ± SEM (n = 4). Unpaired two-tailed Student's t-test. Black and red asterisks represent statistically significant differences in IL-1β compared to uninfected (UI) and *wild-type*-infected (*wt*) macrophages, respectively.

Extended Data Figure 7: The conserved serine-861 residue in the ATPase domain of CgSnf2 is required for the host immune response suppression. a, Schematic representation of different domains in CgSnf2 (CAGL0M04807p), as predicted by the SMART tool (http://smart.embl-heidelberg.de/). QLQ: Gln, Leu, Gln motif; DEXDc: DEAD-like helicases superfamily; HELICc: helicase superfamily c-terminal domain; SnAC: Snf2-ATP coupling, chromatin remodelling complex; AT_hook: DNA binding domain with preference for A/T rich regions and BROMO: bromo domain. Green and pink-colored regions represent coiled-coil and low-complexity regions, respectively. The SnAC domain is involved in binding to histones and required for mobilising nucleosomes. **b,** Multiple amino acid sequence alignment, generated using CLUSTAL-W program, illustrating the conserved serine-861 residue in CgSnf2. **c,** Representative western blot showing similar expression of *wild-type* and serine-to-aspartate-mutated (CgSnf2^{S861D}) CgSnf2 tagged with the triple epitope SFB [S protein-Flag-Streptavidin-binding peptide] at the C-terminus, as detected using anti-FLAG antibody. **d,** Serial dilution spotting analysis illustrating non-complementation of *Cgsnf2Δ* mutant phenotypes by CgSnf2^{S861D}. Methyl methanesulfonate (MMS), hydroxyurea (HU), menadione (MD), zinc chloride (ZnCl$_2$), manganese chloride (MnCl$_2$), β -mercaptoethanol (β-ME), caffeine and sodium dodecyl sulfate (SDS) were used at 0.03%, 25 mM, 100 μM, 8 mM, 3 mM, 20 mM, 10 mM and 0.05% concentrations, respectively. **e,** IL-1β secretion in THP-1 macrophages. Data represent mean ± SEM (n = 4). Unpaired two-tailed Student's t-test.

Extended Data Figure 8: CgSnf2 supresses Akt and p38 activation. a and b, Representative western blots of phosphorylated Akt (**a**) and p38 (**b**) expression. Bar graphs show fold change in phosphorylation, normalized to GAPDH signal. Data represent mean ± SEM (n = 5-6). Paired two-tailed Student's t-test. **c and d,** IL-1β secretion in uninfected and *Cg*-infected THP-1 macrophages which were treated with DMSO or Akt inhibitor (SH-6; 5 μM) (**c**; n=4) and p38 inhibitor (SB203580; 10 μM) (**d**; n=4). Unpaired two-tailed Student's t-test. Black and green asterisks represent statistically significant differences in IL-1β levels, compared to DMSO-treated uninfected and inhibitor-treated uninfected macrophages, respectively. **e,** IL-1β secretion in uninfected and *S. cerevisiae*-infected THP-1 macrophages. Data represent mean ±SEM (n = 4).

Extended Data Figure 9: CgSnf2 binds to *EPA1* promoter. a and c, Integrative genome viewer (IGV) snapshot of MNase-seq signal at *EPA1* **(a)** and *MT* **(c)** loci in *wild-type* cells grown either in RPMI-medium (R2) or incubated with THP-1 macrophages (M2) for 2 h. Green and red peaks in M2-R2 indicate an increase and a decrease in nucleosome occupancy, respectively, upon macrophage internalization. The scaling factor for the Y-axis is indicated at the top of each IGV track. The arrow indicates the direction of transcription. **b,** Nucleosomes at *EPA1* locus (ChrE: 681312 – 686692 bp) are more compactly arranged in 2 h macrophage-internalized (M2), compared to 2 h RPMI-grown *Cg* (R2). Dynamic nucleosomes are shown underneath both *EPA1* and *MT-C* IGV tracks.

Extended Data Figure 10: *Cg* uses chromatin remodelling to escape host immune response. A schematic illustrating how CgSnf2 ATPase-mediated chromatin remodelling, during *Cg*-macrophage interaction, results in cell surface reconfiguration, and aids *Cg* suppress signalling pathways in host macrophages.

Supplementary Information: Supplementary figures 1 and 2, and Supplementary tables 1-10

Supplementary Figure Legends

Supplementary Figure 1: Nucleosome organization in *Cg*. a and b, Overlays of average nucleosome occupancy profiles of 5300 *Cg* genes, relative to the transcription start site (TSS) **(a)** and the transcription termination site (TTS) **(b)**. R2: 2 h RPMI-grown, R10: 10 h RPMI-grown, M2: 2 h macrophage-internalized and M10: 10 h macrophage-internalized *Cg* cells. **c,** Heatmaps depicting global nucleosome occupancy profiles around TSS which are clustered into 10 distinct gene clusters. **d,** Overlays of gene-cluster-specific nucleosome occupancy profiles around TSS. The box color corresponds to the color code of each cluster depicted in panel *c*.

Supplementary Figure 2: Heatmap showing enriched Gene Ontology-Biological Process (GO-BP) terms in four compared datasets.

Supplementary Tables
Table 1: Nucleosome mapping in *C. glabrata* (*Cg*).
Table 1a: Total nucleosomes identified in 2 h RPMI-grown *wild-type* (w*t*) cells.
Table 1b: Total nucleosomes identified in 2 h macrophage-internalized *wt* cells.
Table 1c: Total nucleosomes identified in 10 h RPMI-grown *wt* cells.
Table 1d: Total nucleosomes identified in 10 h macrophage-internalized *wt* cells.

Table 2: Nucleosome dynamics in *Cg*.
Table 2a: Nucleosomes showing position shifts in 2 h macrophage-internalized *Cg*, as compared to 2 h RPMI-grown *Cg*.
Table 2b: Nucleosomes showing occupancy changes in 2 h macrophage-internalized *Cg*, as compared to 2 h RPMI-grown *Cg*.
Table 2c: Nucleosomes showing fuzziness changes in 2 h macrophage-internalized *Cg*, as compared to 2 h RPMI-grown *Cg*.

Table 2d: Nucleosomes showing position shifts in 10 h macrophage-internalized *Cg*, as compared to 10 h RPMI-grown *Cg*.
Table 2e: Nucleosomes showing occupancy changes in 10 h macrophage-internalized *Cg*, as compared to 10 h RPMI-grown *Cg*.
Table 2f: Nucleosomes showing fuzziness changes in 10 h macrophage-internalized *Cg*, as compared to 10 h RPMI-grown *Cg*.
Table 2g: Nucleosomes showing position shifts in 10 h macrophage-internalized *Cg*, as compared to 2 h macrophage-internalized *Cg*.
Table 2h: Nucleosomes showing occupancy changes in 10 h macrophage-internalized *Cg*, as compared to 2 h macrophage-internalized *Cg*.
Table 2i: Nucleosomes showing fuzziness changes in 10 h macrophage-internalized *Cg*, as compared to 2 h macrophage-internalized *Cg*.
Table 2j: Nucleosomes showing position shifts in 10 h RPMI-grown *Cg*, as compared to 2 h RPMI-grown *Cg*.
Table 2k: Nucleosomes showing occupancy changes in 10 h RPMI-grown *Cg*, as compared to 2 h RPMI-grown *Cg*.
Table 2l: Nucleosomes showing fuzziness changes in 10 h RPMI-grown *Cg*, as compared to 2 h RPMI-grown *Cg*.

Table 3: Nucleosome dynamics at gene promoters in *Cg*.
Table 3a: Genes with dynamic nucleosomes at their promoter regions in 2 h macrophage-internalized *Cg*, as compared to 2 h RPMI-grown *Cg*.
Table 3b: Genes with dynamic nucleosomes at their promoter regions in 10 h macrophage-internalized *Cg*, as compared to 10 h RPMI-grown *Cg*.
Table 3c: Genes with dynamic nucleosomes at their promoter regions in 10 h macrophage-internalized *Cg*, as compared to 2 h macrophage-internalized *Cg*.
Table 3d: Genes with dynamic nucleosomes at their promoter regions in 10 h RPMI-grown *Cg*, as compared to 2 h RPMI-grown *Cg*.

Table 4: Gene Ontology (GO) enrichment analysis for *Cg* genes that contain dynamic nucleosomes at their promoters.
Table 4a: Enriched GO terms in 2 h macrophage-internalized *Cg*, as compared to 2 h RPMI-grown *Cg*.
Table 4b: Enriched GO terms in 10 h macrophage-internalized *Cg*, as compared to 10 h RPMI-grown *Cg*.
Table 4c: Enriched GO terms in 10 h macrophage-internalized *Cg*, as compared to 2 h macrophage-internalized *Cg*.
Table 4d: Enriched GO terms in 10 h RPMI-grown *Cg*, as compared to 2 h RPMI-grown *Cg*.

Table 5: Differential gene expression in response to macrophage internalization and *CgSNF2* deletion.
Table 5a: Differentially-expressed genes (DEGs) in 2 h macrophage-internalized *wild-type* (*wt*), as compared to 2 h RPMI-grown *wt*.
Table 5b: DEGs in 2 h RPMI-grown *Cgsnf2Δ*, as compared to 2 h RPMI-grown *wt*.
Table 5c: DEGs in 2 h macrophage-internalized *Cgsnf2Δ*, as compared to 2 h RPMI-grown *Cgsnf2Δ*.
Table 5d: DEGs in 2 h macrophage-internalized *Cgsnf2Δ*, as compared to 2 h macrophage-internalized *wt*.
Table 5e: DEGs in 10 h macrophage-internalized *wt*, as compared to 10 h RPMI-grown *wt*.
Table 5f: DEGs in 10 h RPMI-grown *Cgsnf2Δ*, as compared to 10 h RPMI-grown *wt*.

Table 5g: DEGs in 10 h macrophage-internalized *Cgsnf2Δ*, as compared to 10 h RPMI-grown *Cgsnf2Δ*.

Table 5h: DEGs in 10 h macrophage-internalized *Cgsnf2Δ*, as compared to 10 h macrophage-internalized *wt*.

Table 5i: DEGs in the 10 h RPMI-grown *wt*, as compared to 2 h RPMI-grown *wt*.

Table 5j: DEGs in 10 h macrophage-internalized *wt*, as compared to 2 h macrophage-internalized *wt*.

Table 5k: DEGs in 10 h RPMI-grown *Cgsnf2Δ*, as compared to 2 h RPMI-grown *Cgsnf2Δ*.

Table 5l: DEGs in 10 h macrophage-internalized *Cgsnf2Δ*, as compared to 2 h macrophage-internalized *Cgsnf2Δ*.

Table 6: Gene Ontology (GO) enrichment analysis for differentially-expressed genes.

Table 6a: Enriched GO terms in 2 h macrophage-internalized *wt*, as compared to 2 h RPMI-grown *wt*.

Table 6b: Enriched GO terms in 2 h RPMI-grown *Cgsnf2Δ*, as compared to 2 h RPMI-grown *wt*.

Table 6c: Enriched GO terms in 2 h macrophage-internalized *Cgsnf2Δ*, as compared to 2 h RPMI-grown *Cgsnf2Δ*.

Table 6d: Enriched GO terms in 2 h macrophage-internalized *Cgsnf2Δ*, as compared to 2 h macrophage-internalized *wt*.

Table 6e: Enriched GO terms in 10 h macrophage-internalized *wt*, as compared to 10 h RPMI-grown *wt*.

Table 6f: Enriched GO terms in 10 h RPMI-grown *Cgsnf2Δ*, as compared to 10 h RPMI-grown *wt* cells.

Table 6g: Enriched GO terms in 10 h macrophage-internalized *Cgsnf2Δ*, as compared to 10 h RPMI-grown *Cgsnf2Δ*.

Table 6h: Enriched GO terms in 10 h macrophage-internalized *Cgsnf2Δ*, as compared to 10 h macrophage-internalized *wt*.

Table 6i: Enriched GO terms in 10 h RPMI-grown *wt*, as compared to 2 h RPMI-grown *wt*.

Table 6j: Enriched GO terms in 10 h macrophage-internalized *wt*, as compared to 2 h macrophage-internalized *wt*.

Table 6k: Enriched GO terms DEGs in 10 h RPMI-grown *Cgsnf2Δ*, as compared to 2 h RPMI-grown *Cgsnf2Δ*.

Table 6l: Enriched GO terms in 10 h macrophage-internalized *Cgsnf2Δ*, as compared to 2 h macrophage-internalized *Cgsnf2Δ*.

Table 7: List of *C. glabrata* strains used in the study.

Table 8: List of plasmids used in the study.

Table 9: List of primers used in the study.

Table 10: List of antibodies and inhibitors used in the study.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Extended Data Figure 1

Extended Data Figure 2

a M10 Vs R10

Position — Occupancy
129 | 19 | 123
1 | 2 | 2
0
Fuzziness

b M10 Vs M2

Position — Occupancy
458 | 33 | 80

c R10 Vs R2

Position — Occupancy
270 | 27 | 86
3 | 2 | 1
1
Fuzziness

d M10 Vs M2

positive regulation of ascospore-type prospore
membrane assembly
NADPH regeneration
(1->3)-beta-D-glucan biosynthetic process
1,3-beta-D-glucan synthase activity
'de novo' cotranslational protein folding
adhesion of symbiont to host
acetate biosynthetic process
phosphatidic acid biosynthetic process
translational frameshifting
cellular calcium ion homeostasis
glycogen biosynthetic process
drug transmembrane export
cell surface
cellular response to drug
glycolytic process
ascospore formation
mitochondrial nucleoid
transmembrane transporter activity
transcription factor activity, sequence-specific
DNA binding
extracellular region
oxidoreductase activity
vesicle-mediated transport
fungal-type cell wall
ribosome
translation
plasma membrane
structural constituent of ribosome
integral component of membrane
ATP binding

Fold enrichment

count: 30, 60, 90
- log10(pvalue): 2, 3, 4, 5, 6
class: BP, CC, MF

e M10 Vs R10

'de novo' cotranslational protein folding
translational frameshifting
regulation of translational fidelity
ribosomal subunit export from nucleus
positive regulation of autophagy
cytoplasmic translation
cell surface
fungal-type cell wall
transmembrane transporter activity
integral component of membrane

Fold enrichment

- log10(pvalue): 1.6, 2.0, 2.4
class: BP, CC, MF
count: 10, 20, 30, 40, 50

f R10 Vs R2

gluconeogenesis
translational termination
nucleosome
cell surface
cellular response to drug
GTPase activator activity
protein heterodimerization activity
fungal-type cell wall
extracellular region
ribosome
plasma membrane

Fold enrichment

count: 4, 8, 12, 16
- log10(pvalue): 2, 3, 4, 5, 6
class: BP, CC, MF

Extended Data Figure 3

Extended Data Figure 4

Extended Data Figure 5

Extended Data Figure 6

a

b

H. sapiens	VSLYNNNLNGILADEMGLGKTIQTIALITYLMEHKRINGPFLIIVPL⟦S⟧TLSNWAYEFDKW	825
C. albicans	VSLFNNHLNGILADEMGLGKTIQTISLLTYLVEVKKIPGPFLVIVPL⟦S⟧TVTNWNLEFEKW	852
S. cerevisiae	VSLFNNHLNGILADEMGLGKTIQTISLLTYLYEMKKIRGPYLVIVPL⟦S⟧TLSNWSSEFAKW	838
C. glabrata	VSLFNNHLNGILADEMGLGKTIQTISLLTYLYEMKNIKGPFLIIVPL⟦S⟧TLPNWSSEFAKW	873

c

Cgsnf2Δ

d

	wt	*Cgsnf2Δ*	*CgSNF2-SFB*	*CgSNF2^{S861D}-SFB*
YPD				
MMS (0.03%)				
HU (25 mM)				
MD (100 µM)				
ZnCl$_2$ (8 mM)				
MnCl$_2$ (3 mM)				
β-ME (20 mM)				
Caffeine (10 mM)				
SDS (0.05%)				

Cgsnf2Δ

e

Extended Data Figure 8

Extended Data Figure 9

Supplementary Figure 1

Supplementary Figure 2