Strong gene flow across an urbanised coastal landscape in a dune specialist digger wasp

Authors:

Femke Batsleer[1]*(ORCID 0000-0001-5893-5966)

Fabien Duez[1,2] (ORCID 0000-0003-2730-6694)

Dirk Maes[3,4] (ORCID 0000-0002-7947-3788)

Dries Bonte[1] (ORCID 0000-0002-3320-7505)

[1] Ghent University, Department of Biology, Terrestrial Ecology Unit, Karel Lodewijk Ledeganckstraat 35, 9000 Gent, Belgium

[2] Institut des Sciences de l'Evolution, Université de Montpellier, CNRS, IRD, Montpellier, France

[3] Research Institute for Nature and Forest (INBO), Herman Teirlinckgebouw, Havenlaan 88 bus 73, 1000 Brussel, Belgium

[4] Radboud Institute for Biological and Environmental Sciences (RIBES), Radboud University, PO Box 9010, NL-6500 GL Nijmegen, The Netherlands

*Corresponding author: F. Batsleer, Ghent University, Department of Biology, Terrestrial Ecology Unit. Karel Lodewijk Ledeganckstraat 35, 9000 Ghent, Belgium. E-mail: femke.batsleer@ugent.be.

Telephone: +32 (0)9 264 52 58

Abstract

Context

Genetic connectivity is often disrupted by anthropogenic habitat fragmentation, and therefore often

a focus in landscape-scale conservation. Landscape genetics methods allow for studying functional

connectivity in heterogenous landscapes in detail and have the potential to inform conservation

measures for a species' regional persistence. Yet, for insects, functional connectivity through

landscape genetics remains largely unexplored.

Objectives

We aimed to assess the functional connectivity in the dune-specialist digger wasp *Bembix rostrata*, in

a human-altered 15 km section along the Belgian coast, based on landscape genetics methods.

Methods

We optimised landscape resistance distances according to individual genetic distances with the

package ResistanceGA. We combined this with a multi-model inference approach to deduce relative

conductance or resistance of gene flow to natural and anthropogenic landscape types.

Results

Overall, the populations of this dune-specialist insect are genetically well-connected. We detected—

on top of the prominent background process of isolation-by-distance—a weak but consistent signal

of urban features facilitating gene flow.

Conclusions

Urban areas are not a barrier to genetic connectivity of the dune-specialist insect *B. rostrata* in the

focal human-altered landscape. However, because urbanisation leads to larger scale fragmentation,

its impact on the distribution of populations in the landscape and related effective regional gene flow

remains substantial. As this species depends on early-succession dune vegetations, restoring and

increasing sand dynamics at the local and landscape scale should be the focus of conservation aimed

at the regional species' persistence, rather than trying to increase habitat connectivity in this human-

altered dune ecosystem.

Keywords: landscape genetics, landscape resistance modelling, resistanceGA, resistance

optimisation, dispersal, Hymenoptera

Introduction

Movement of individuals through landscapes is an inherent component of dispersal and can lead to exchange between populations, resulting in gene flow and connectivity. Connectivity can ultimately support a species' regional persistence, as it maintains or supports a wide range of natural processes, such as resource use, demographic rescue, genetic diversity, range expansion, or metapopulation dynamics (Hanski 1998; Crooks and Sanjayan 2006; McRae et al. 2008). The environment in between habitat patches (i.e. the matrix) can impede or facilitate the movement or dispersal of an organism. This matrix is in reality more heterogenous than the practical simplification of a homogeneous matrix, especially in human-altered landscapes (Hein et al. 2003; Manel and Holderegger 2013). The realised connectivity of a species is influenced by distances between populations and the specific landscape context: the structural connectivity. However, it also depends heavily on the behavioural responses of an organism to different components of the landscape (Hodgson et al. 2022). These responses vary inherently between species and individuals (Baguette and Van Dyck 2007; Knowlton and Graham 2010; Logghe et al. 2024). The integration of movement into measures of functional connectivity (Tischendorf & Fahrig, 2003) is, therefore, crucial for understanding the dynamics of spatially structured populations (or metapopulations) and to inform effective conservation measures (McRae 2006).

A general view on functional connectivity is that many organisms will avoid unfamiliar habitat altogether—they exhibit high reluctance to cross gaps and favour instead considerable detours—making unfamiliar habitat a barrier to movement (Develey and Stouffer 2008; Knowlton and Graham 2010; Driscoll et al. 2013). This is the general assumption in for instance least cost mapping, which considers only the most cost-effective path through a landscape (Adriaensen et al. 2003). Unfamiliar habitat can, however, also act as a facilitator to movements when search behaviour of individuals in unfamiliar habitat is more directional—and thus faster—than in favoured habitat (Van Dyck and Baguette 2005; Knowlton and Graham 2010). Or conversely, movements in more suitable habitats

can be impeded because of the availability of resources, leading to less directed—and thus slower—

exploratory, routine movements than in unfamiliar habitat (Van Dyck and Baguette 2005; McRae et

al. 2008; Keller et al. 2012). The behavioural differences between more directional movement paths

(covering greater distances) during displacement and (slower) exploratory, more random, routine

movements are for insects best studied in butterfly species (Schtickzelle et al. 2006; Kuefler et al.

2010). Independent of the exact mechanism behind this potential (relative) increase in movement

through the matrix, the eventual functional connectivity will remain dependent on the distance

between habitats and thus the scale of the fragmentation. The same matrix typology or landscape

category can therefore potentially facilitate or constrain connectivity in different landscapes,

depending on the eventual distance-related costs in terms of survival and energy-use (Segelbacher et

al. 2010; Spear et al. 2010; Bonte et al. 2012; Richardson et al. 2016; Haran et al. 2017).

How a landscape influences the mobility of an organism has implications for the design of regional

conservation measures. If animals show high reluctance towards crossing habitat boundaries,

creating corridors might be an effective structural connectivity measure to increase functional

connectivity between subpopulations. Such regional landscape-scale conservation measures,

including corridors and stepping-stones, which are based on logical and sound scientific principles,

have been widely embraced by conservation practitioners and policy-makers (Crooks and Sanjayan

2006; Watts et al. 2016). However, there is much debate about the effectiveness of landscape-scale

conservation and the emphasis on increasing connectivity for species' regional persistence in

conservation decisions (Hodgson et al. 2009; Humphrey et al. 2015; Watts et al. 2016). The emphasis

of practitioners and policy-makers on landscape-scale conservation is probably biased by a focus

towards larger terrestrial vertebrates, for which structural connectivity—such as corridors—seems

effective (Humphrey et al. 2015). Empirical evidence for many other groups is more limited and

equivocal (Ovaskainen et al. 2008; Watts et al. 2016). In insects, it is generally assumed that the

regional persistence of spatially structured populations is mostly determined by local demography

(Richardson et al. 2016; Drake et al. 2022).

In such cases, site-based conservation focusing on improving habitat area and quality could be more

effective than increasing structural connectivity on the landscape-scale. To effectively direct the

limited resources available for conservation measures towards implementations with the highest

success rate or biodiversity gain, implementations should be founded on empirical evidence

(Sutherland et al. 2004; Ferraro and Pattanayak 2006; Samways et al. 2020). Especially for insects—

whose decline and biological value get widespread attention (Didham et al. 2020; Wagner et al.

2021)—scientific evidence on the effects of landscapes on functional connectivity and gene flow is

still limited (Keyghobadi et al. 1999; Wilcock et al. 2007; Keller et al. 2012; Pérez-Espona et al. 2012;

Watts et al. 2016; Haran et al. 2017; Trense et al. 2021).

An indirect way of measuring functional connectivity is through molecular genetic data. With

landscape genetics methods, one can quantify the effects of landscape composition between habitat

patches on gene flow between populations (Manel et al. 2003; Manel and Holderegger 2013;

Balkenhol et al. 2015). Patterns of gene flow reflect realised or successful dispersal and depend on

post-dispersal survival and reproduction (Spear et al. 2010). Landscape genetics methods allow for

inferences regarding connectivity without the need to collect individual movement data. Collecting

such field data—for example from telemetry or mark-recapture methods—requires intensive effort

or is even unfeasible for many organisms, such as the majority of insects (Spear et al. 2010; Batsleer

et al. 2020). Only some larger insect species have been employed with tracking devices, and even for

these, the biases on the resulting measurements remain insufficiently considered (Batsleer et al.

2020). Both mark-recapture and telemetry rarely detect, and thus underestimate, long-distance

dispersal (Ugelvig et al. 2012; Trense et al. 2021; De Ro et al. 2021). The advancing tools in molecular

and landscape genetics, such as in landscape resistance modelling, can help us understand

connectivity and landscape use in species for which direct estimates of habitat use and movement

are difficult to obtain with more classical field methods (Spear et al. 2010). However, such methods

are rarely applied to insects. Studies on a few specialist insects point towards non-habitat and/or

anthropogenic features forming barriers, such as in in the Sooty Copper butterfly *Lycaena tityrus*

(Trense et al. 2021) and the army ant *Eciton burchellii* (Pérez-Espona et al. 2012). While for the long-

horned beetle *Monochamus galloprovincialis*, which is considered an excellent disperser and even a

vector for diseases in pinewoods (David et al. 2014), open areas do not seem a barrier to gene flow

(Haran et al. 2017).

Coastal dunes in Flanders are geological young formations with calcareous sandy soils forming a

narrow, elongated ecosystem along the coast (Provoost et al. 2004; Decleer 2007). They harbour a

threatened, specific insect biodiversity typical of sandy early-succession and pioneer vegetations

(Maes and Bonte 2006). In northwest Europe, and more specifically Flanders (Belgium), coastal dunes

have gone through extensive landscape changes and fragmentation during the past century

(Provoost et al. 2011). The fragmentation of coastal sandy habitats had two main impacts: loss of

total area and decrease in habitat quality of the remaining patches. Urbanisation from the

interbellum period (1920-1940) onwards decreased the total dune area by half during the previous

century and separated the major dune entities physically (Provoost and Bonte 2004). The resulting

obstruction of wind and sand dynamics, combined with the loss of agricultural practices (such as

grazing of livestock), stimulated succession and scrub development, decreasing the amount of open

dune habitats (Provoost et al. 2011). Large herbivores have been introduced in many coastal dune

reserves to revitalise dune dynamics as a substitute for natural wind dynamics, but have—due to

trampling—a variable effect on local arthropod species (Bonte and Maes 2008; van Klink et al. 2015;

Batsleer et al. 2022b). Altogether, coastal dunes in Flanders are a human-altered landscape with

complex nature management considerations to consider in both landscape scale and site-based

nature management. In such a landscape, understanding functional connectivity—with its many

specialist dune species—is vital to make evidence-based conservation decisions.

Conservation and connectivity studies have historically focused on plants and vertebrates, with

insects receiving less attention. When insects are considered, the focus is typically on butterflies and

pollinators (Clark and May 2002; Potts et al. 2016). Even for pollinators, the importance of nesting

resources is often overlooked in nature management (Kimoto et al. 2012; Buckles and

Harmon-Threatt 2019). As a result, ground-nesting Hymenoptera remain particularly neglected in

both conservation and connectivity studies. The digger wasp *Bembix rostrata* is considered an

emblematic representative of this group, associated with open and early-succession dune

vegetations in Belgium (Batsleer et al. 2022b).

Here, we report on the genetic (functional) connectivity of the dune-habitat specialist digger wasp *B.*

rostrata in a human-altered coastal landscape in Belgium using landscape genetics methods. While it

is known from an earlier population genetic study that the species' regional connectivity is large

(Batsleer et al. 2024), it remains unknown which specific landscape components facilitate or

constrain gene flow. Given the strong urbanisation of the coastal dunes, we expect this land-use to

form a barrier to gene flow in this dune specialist insect. Given the species biology, we expect the

dune areas and beaches to function as facilitators to gene flow. These hypotheses are parallel to

results for other specialist insects in landscape genetics studies (Pérez-Espona et al. 2012; Trense et

al. 2021). We use a resistance raster approach to quantify resistance values of the distinctive

landscape types in our study area. The resistance values represent the degree to which each

landscape type impedes genetic connectivity for the study species (Spear et al. 2010). To determine

the resistance values of each landscape type, these values are optimised, by maximising the

correlation between the landscape resistance distances and observed genetic distances between all

pairwise individuals (Peterman 2018). All possible combinations of landscape categories were

optimised to be able to perform multi-model inference. This allows us to deduce possible barriers or

facilitators in the landscape for this dune-specialist digger wasp. Based on this estimate of functional

connectivity, we can make recommendations on the balance between landscape- and site-based

conservation.

Material & Methods

Study species

Bembix rostrata (Linnaeus, 1785; Hymenoptera, Crabronidae, Bembicinae) is a univoltine dune-specialist, gregariously nesting digger wasp from sandy habitats with sparse vegetation, sensitive to grazing management (Larsson 1986; Klein and Lefeber 2004; Bonte 2005; Batsleer et al. 2022b). Adults are active in summer, showing protandry: females are immediately mated upon emergence by the guarding males, who emerge one to five days earlier (Wiklund and Fagerström 1977; Schöne and Tengö 1981; Evans and O'Neill 2007). Females show brood care: one individual constructs one nest burrow at a time in which it progressively provisions a single larva with flies (Nielsen 1945; Field et al. 2020). It is estimated that up to 5 nests (in northern regions) are produced, each with one offspring (Larsson and Tengö 1989). Nesting locations of *B. rostrata* usually span a width of less than 100 m, with nests arranged in several clusters that have a maximum diameter of 10 m (Batsleer et al. 2022a). Within these clusters, nest densities typically reach up to 10 nests per square meter (Nielsen 1945; Larsson 1986). There are no overlapping generations, as the species overwinters as prepupa and there is only one generation per year. *Bembix rostrata* is often considered to be a philopatric species, as they do not easily colonise vacant habitat and exhibit site fidelity throughout the nesting season (Nielsen 1945; Larsson 1986; Bogusch et al. 2021; Batsleer et al. 2022a). However, a population genetics study across Belgium showed that this does not preclude gene flow between existing populations, including some dispersal to distant, more isolated populations (Batsleer et al. 2024). This population genetics study showed that *B. rostrata* has relatively high genetic connectivity in the coastal dunes in Belgium: genetic distances were low, genetic compositions between populations substantially overlapped, a large number of immigrant links between coastal population were detected and an overall pattern of isolation-by-distance was observed (Batsleer et al. 2024).

Genetic sampling

Sampling took place in the summer of 2018 in the dunes along the Belgian west coast (Fig. 1), from the French-Belgian border (De Panne) to the Yser estuary (Nieuwpoort), stretching 14 km along the coast and reaching maximum 4 km inland. As nests are clustered together at nesting aggregates in grey dunes (EU Habitats Directive habitat 2130), we tried to spatially cover all known nesting locations in the landscape and locally took samples spatially spread within a nesting location, across clusters of nests. Sample locations were based on a spatially exhaustive survey of potential nesting sites the previous summer. The coordinates of each individual sample were saved. Only females were sampled, to solely use diploid individuals in the genetic analyses for this haplodiploid mating system (Zayed 2009; Batsleer et al. 2024). To minimise the impact of sampling, we used a non-lethal sampling procedure with wing clips, a method shown to produce good-quality DNA for microsatellite PCR amplification and having no clear behavioural effects for honey bee queens and workers (Châline et al. 2004). Such wing clips of less than 10% of the wing area mimic natural wear and are shown not to have an effect on metabolic flight costs in bumblebees (Hedenström et al. 2001). Forewing tips from live digger wasps were cut on both sides and stored in absolute ethanol, kept at 4°C after each day of sampling and transferred to a freezer (-18°C) for longer term storage. The individual digger wasps displayed normal behaviour when flying off after sampling.

DNA extraction and PCR amplification were performed as described in Batsleer et al. (2024) and summarised hereafter. DNA is extracted from the wing tips with a protocol based on Chelex (Biorad). 33 polymorphic microsatellites are PCR-amplified for each sampled individual and electropherograms are scored using Geneious Prime 2021.0 (Biomatters). Microsatellites that had a lot of stutter in amplification (five microsatellites) or that were excluded after assumption testing (Hardy-Weinberg, null alleles and linkage disequilibrium; seven microsatellites) in the population genetics study in (Batsleer et al. 2024) were also excluded here, resulting in 21 microsatellites (out of 33) for the subsequent genetic analysis. If an individual had more than 8 missing loci, it was discarded from the

analyses (2.4%, 13 out of 552). A total of 539 sampled individuals were successfully amplified. The

resulting data contained an overall of 0.7% missing data, which were replaced by the mean allele

frequencies for the calculation of the genetic distances (see further).

Land cover categories

We combined two types of land cover datasets available for the study area. For landscape categories

within dune areas, we used detailed vegetation maps (resolution 0.4m) available for the region

derived from remote sensing data with a classification machine learning technique (Bonte et al.

2021). For landscape categories outside dune areas, we used the ground cover map 2018 provided

by the Flemish government ('Bodembedekkingskaart' BBK 1m resolution, source: www.geopunt.be).

We aligned, reclassified and combined the raster datasets (see online resource 1 for detailed

information on reclassificaitons) to have one raster file with resolution of 1m of the study area with 7

simplified landscape categories: urbanised (*urban*), *water*, agricultural (*agric*), *beach*, *scrub*, *trees* and

open dune (*opend*) vegetation (Fig. 1). Roads and urban features were analysed as a single category,

as roads are almost always embedded or aligned within urban area, resulting in a high degree of

spatial dependency. Category *beach* was added by rasterizing a manually drawn shapefile based on

aerial maps and the high water line. For further analyses, we resampled the rasters to work with

resolutions of 20 m and 50 m. We deem these resolutions (spatial grains) adequate to capture the

turnover in heterogeneity of the focal landscape and to be linked with the ecological processes

investigated. When relevant landscape elements are captured by the resolution, landscape genetic

analyses should be relatively robust for changes in cell size (McRae et al. 2008; Cushman and

Landguth 2010).

243
244

Figure 1: overview map of study area: west coast in Belgium from the French-Belgian border (west; dashed red

line) to the Yser estuary (east). Sample locations of each individual (blue dots) are shown and four dune area

clusters (blue dashed lines) are shown. Four dune area clusters (blue dashed lines) are ecological clusters

deemed relevant to nature management; 1) Westhoek 2) Cabour 3) Doornpanne 4) Ter Yde. The landscape

raster categories are shown with a 20m × 20m resolution (cell size). Neighbouring countries of Belgium (BE):

France (FR), Luxembourg (LU), Germany (DE), The Netherlands (NL), United Kingdom (GB). Map made with

QGIS v3.22 (QGIS Development Team 2020) in Cartesian coordinate reference system Belgian Lambert 72,

EPSG:31370. Background aerial photographs (summer 2018) from Agency for Information Flanders (source:

geopunt.be).

Genetic distance measure

We used individual-based genetic distances between pairwise individuals. We used a metric based on multiple axes of a principle components analysis (PCA), which was shown to perform better in landscape genetics analyses than other individual-based metrics, especially when sample size and genetic structure are low (Shirk et al. 2017; Kimmig et al. 2020). First, principal components (PC) were calculated from a multiple contingency table (0, 1, 2 values for each allele for each individual). Second, the Euclidean distance based on the first 16 PCs was calculated to create a genetic distance matrix between all individuals (Shirk et al. 2017). To calculate these genetic distances, we used the R-packages adegenet v2.1.10 (Jombart 2008) and ecodist v2.1.3 (Goslee and Urban 2007).

Landscape resistance optimisation

The unknown values are the resistance values of each landscape type, which quantify at each grid cell the constrain to gene flow (Spear et al. 2010). Based on these resistance values, resistance distances between pairwise individuals are iteratively recalculated during the optimisation (see further for details). Such resistance distances reflect a measure of separation between pairwise individuals and incorporate the effects of different permeabilities of the landscape (Spear et al. 2010). The resistance values are optimised based on maximising the correlation between genetic distances and landscape resistance distances between pairwise individuals. Thus, it is expected that the effect of the landscape on genetic connectivity has a detectable signal on the observed genetic distances. The optimisation was implemented with the RESISTANCEGA R package v4.0-0 (Peterman 2018).

Resistance distance measure. We used the commute resistance distance in RESISTANCEGA (van Etten 2017; Peterman 2018), which is the expected time of a random walk (the stochastic process of random steps) between a pair of points (sampled individuals) along a circuit-based resistance surface (the raster with resistance values). This type of distance accounts for several possible pathways through the landscape (McRae et al. 2008; Peterman 2018). The pairwise resistance distances are

recalculated at each iteration based on the resistance values of the landscape types during the

previous iteration (with initial values of 1).

Optimisation procedure. RESISTANCEGA uses a linear mixed effects model with maximum likelihood

population effects (MLPE) parameterisation to fit pairwise resistance distances (predictor) to

pairwise genetic distances (response). MLPE parameterisation accounts for the non-independence

among pairwise data (Clarke et al. 2002; Bates et al. 2015). The optimisation procedure iteratively

searches many possible resistance value combinations to maximise the MLPE model fit (based on log-

likelihood), resulting in the best fit between resistance distances and genetic distances (Peterman

2018). This optimisation procedure is based on a genetic algorithm (Scrucca 2013). For each raster,

we used the SS_OPTIM function in RESISTANCEGA to optimise the resistance values (see online code).

Parameters were set to default, only the maximum number of iterations (*max.iter*) was changed to

300. There are no a priori expectations for the resistance values of the landscape categories, as no

information is present regarding habitat selection and movement behaviour at the landscape scale

for this species (Knowlton and Graham 2010). Consequently, and by default, the resistance values of

each category have an initial value of 1.

Multicategorical rasters. We optimised multi-categorical rasters, which consider several landscape

categories at once. Although more complex to parameterise, multi-categorical raster optimisation

adds more biological realism to the analysis than considering only one single category at a time,

which would assume an organism's perception of a landscape to be binary (Spear et al. 2010). We

ran optimisations for each possible combination of all 7 categories. A raster then consists of one to

seven focal landscape categories and all else or 'the rest' (for example when categories *urban* and

scrub are considered, all five other categories are lumped into a third rest category). The total

number of all possible combinations of 7 categories (*urban, water, agric, beach, scrub, trees* and

opend vegetation) is 127: $\sum_{k=0}^{7} \binom{7}{k}$. Ascii-files needed as input for RESISTANCEGA were created for

each possible combination with the R-package terra v1.7.78 (Hijmans 2024), see online code. Thus,

optimisations are run for each possible combination for the 7 categories.

Bootstrap analysis. To compare the performance of each of the 127 optimised rasters and the

Isolation-By-Distance (IBD) as null model, we performed a bootstrap analysis. With the RESIST.BOOT

function from RESISTANCEGA, 75% (default) of the individuals are subsampled and the MPLE model

refitted for each of the 127 optimised rasters and IBD, this for 1000 iterations. Relative support for

each model is then compared using several bootstrap-derived metrics for model performance,

importance and stability. The primary metric used is the difference in corrected Akaike information

criterion (AICc, correction for small sample sizes; measure for predictive accuracy of a model). In

general, it is considered that models with ΔAICc < 2 are competing models, and even models with

ΔAICc ranging between 2-7 have some support and should not be dismissed (Burnham et al. 2011).

Other metrics included are: number of times the model was the top model (n.top), the average rank

(avg.rank, measure for overall relative importance), and the average Akaike weight (avg.weight,

measure for relative contribution), average marginal R^2 (explained variance) and log-likelihood

(predictive fit) across bootstrap samples. The extra bootstrap-metrics indicate, next to ΔAICc, if

certain models are estimated to be consistently more important than other models across bootstrap

samples.

Independent optimisations runs. As the optimisation with RESISTANCEGA is a stochastic process,

optimised values can differ from run to run, and it is advised to run all optimisations at least twice to

confirm convergence and relative relationships among resistance values (Peterman 2018). To check

this convergence and account for a possible bias due to the starting cell values of a category in multi-

categorical rasters (i.e. the initial order: urban, water, agricultural, beach, scrub, trees, open dune)

(Peterman 2018; Kimmig et al. 2020), we ran the multi-categorical raster optimisations and bootstrap

analysis a second time with the order of categories inverted (starting with *opend*, ending with *urban*).

To confirm convergence more conclusively, we ran optimisations and bootstrap analyses of the

seven single category rasters (that consider only one category) four times each (online resource 2).

This is computationally less intensive than running all possible combinations, but gives a broad idea

of the robustness of the optimisation.

Multi-model inference. As all possible combinations of categories were optimised to obtain

resistance values, we were able to use a multi-model inference approach to obtain robust estimates

of average weight for each category and average relative resistance value. Multi-model inference is a

procedure to account for the uncertainty of parameter estimates related to model selection and is

especially useful when no single model is overwhelmingly supported by the data (Johnson and

Omland 2004). We used Akaike weights from the bootstrap analysis considering all the models (i.e.

optimised rasters) for each area type. From these, we calculated the summed Akaike weight per

category by taking into account a model if the focal category was included (Burnham and Anderson

2004; Johnson and Omland 2004). These summed Akaike weights give an indication for the relative

importance of each category to explain the pairwise genetic distances. We also calculated relative

resistance values of each category. As optimised resistance values depend on the other categories

included in a multi-categorical model, interpretation for a focal category should be done relative to

these other categories. To be able to do this for each landscape category, we rescaled the resistance

values per optimised raster: we subtract the resistance value of the focal category from the other

resistance values that were simultaneously optimised. As such, the value of the focal category

becomes zero and is considered facilitating to gene flow compared to categories with values larger

than zero and a barrier to gene flow compared to categories with values smaller than zero (for

further information on interpretation, see results section). All models and the relative resistance

values per focal category were then summarised by plotting boxplots for each focal category with

ggplot2 v3.5.1 (Wickham 2016). We consider this as a qualitative type of model averaging in multi-

model inference, where parameters are summarised across all models (Johnson and Omland 2004).

We remain descriptive with these relative resistance values, as values used in these boxplots are not

independent because categories are optimised simultaneously.

To get an idea of the possible small scale variation in landscape effects on gene flow, we repeated

the analyses for each of the four dune area clusters within our study area (Fig. 1). These dune areas

are seen as ecological clusters relevant for nature management as these are clusters of the largest

local nature reserves (Fig. 1). These optimisations for the dune area clusters are run with a resolution

of 20 m (in contrast to 50 m for the complete study area). Details and results can be found in online

resource 4.

The total computational time (runs for all possible combinations of categories) was very high and

could not be run on a single desktop. Therefore, we made use of the high-performance computing

infrastructure of Ghent University (HPC-UGent) to run the optimisations and bootstrapping analyses.

By doing this, we could run all possible combinations in the multi-categorical raster optimisations,

which is the most ideal for robust statistical inference and allows for model averaging to assess the

importance of the different landscape categories. Previous studies reported computational

constraints for optimisations with RESISTANCEGA considering all combinations and used alternative

approaches for model comparison based on model selection (e.g. Lourenço et al. 2019; Kimmig et al.

2020).

The best-supported models (i.e. optimised rasters) for the complete study area were used to

visualise the putative gene flow currents in the landscape using Circuitscape (McRae et al. 2013;

Kimmig et al. 2020). Gene flow currents in Circuitscape represent the putative genetic connectivity in

a landscape, represented by conductance values for a grid cell, where most gene flow passes

between sample locations in the given landscape. Current maps were inferred for each pair of

sample locations (representing multiple possible pathways in the landscape). The conductances of

these current maps were all summed for the final Circuitscape current map, to give a representation

of the overall genetic connectivity of the landscape (McRae et al. 2008; Kimmig et al. 2020).

The main analyses (see online code) were performed in RSTUDIO V1.4 (RStudio Team 2022) using R

v4.1.3 (R Core Team 2020), except for calculations for the current map with the GUI of Circuitscape

v.4.0.5 (McRae et al. 2013) and cartographic visualisations in QGIS v3.22 (QGIS Development Team

2020).

Results

Below, we first describe the results of the optimisation of the resistance values of the multi-categorical landscape maps after bootstrapping. The results show which optimised landscape resistance models relatively perform best to explain the gene flow of *Bembix rostrata* in the study area. The optimisations provide which landscape categories are facilitating or forming a barrier to gene flow for each landscape model. Second, we give results on the multi-model inference to deduce overall patterns of landscape categories across all optimised models. The results indicate which landscape categories are, among all optimised landscape models, the most important in explaining the gene flow and what their relative resistance values are. We further provide a visualisation of Circuitscape connectivity maps of the three best-supported models and describe the results of the analyses on the smaller scale dune area clusters within the study area.

Bootstrap analysis of landscape resistance optimisation

Results of the top five best supported models with optimisations of multi-categorical models (all possible combinations of landscape categories) after bootstrapping can be found in table 1 (full table in online resource 3). In general, there was no overwhelming support for one or a few models based on $\Delta AICc$, as isolation-by-distance (null model) was one of the best supported models (Distance; $\Delta AICc = 1.20$ run 1 and 1.42 run 2; table 1) and was a competing model considering $\Delta AICc < 2$ with *urban* (the best supported model). In general, R^2m's were low (table 1; overall <1%), Akaike weights were small (avg.weight in table 1) and all $\Delta AICc$'s were relatively small (table 1; maximum $\Delta AICc$ was 14.58; online resource 3), and even models with $\Delta AICc$ ranging between 2-7 have some support and should not be dismissed (Burnham et al. 2011).

Multi-model inference

When no single model is overwhelmingly supported by the data, multi-model inference is ideal to deduce average weights and relative resistance values of categories across all models (Johnson and Omland 2004). From such a multi-model inference based on the optimisation of all possible

combinations, we deduced more general patterns from the data based on the summed Akaike

weights per landscape category (table 2) and the relative resistance values (Fig. 2). *Urban* was an

important explanatory category to explain pairwise genetic distances (table 2), which acted mainly as

facilitator to gene flow compared to other landscape categories (table 1, Fig. 2). Natural landscape

categories (*beach, scrub, trees, open dune*; bottom row in Fig. 2) were in general more resistant to

gene flow than anthropogenic categories (*urban, agriculture, water*; top row in Fig. 2) in the focal

landscape. As most categories, apart from urban, were of little (and inconclusive) importance to

explain pairwise genetic distances (table 2), the uncertainty of relative resistance values between

these categories (especially natural categories, Fig. 2 second row) should be considered high. The

second independent optimisation run (included in table 1 and 2; Fig. 2) confirmed these patterns and

showed that category *urban* was consistently present in the top best supported models (table 1) and

the most important (and only conclusive) category to explain pairwise genetic distances (table 2).

This was also confirmed by the four independent runs of the single categories, which consistently

had urban as the first supported model and IBD as the second best competing model (online

resource 2).

Visualisation current maps

422 Circuitscape current (or flow) maps (Fig. 3) were made for the three best supported models (table 1).

423 As all ΔAICc's were relatively small (online resource 3), there were many more models worth

424 considering (as was done in the multi-model inference, Fig. 2 and table 2). However, these

425 Circuitscape current maps give post-hoc visualisations of how the landscape likely influences genetic

426 connectivity within the study area according to the best three models.

Small-scale dune area clusters

The analysis repeated at the level of each of the four dune area clusters (Fig. 1) showed Isolation-By-

Distance (IBD) to be the overall best supported model on this smaller scale (online resources 4 and

5). Best supported models and the multi-model inference showed some smaller variation among the

studied clusters (online resource 4) and generally confirmed the overall pattern from the complete

study area: anthropogenic categories (*urban*, *agric*, *water*) were more facilitating to gene flow than

natural landscape categories. There were two exceptions: for the dune area Doornpanne (dune

cluster 3 in Fig. 1), *trees* was a facilitating landscape property to gene flow, and for Cabour (dune

cluster 2 in Fig. 1), *agriculture* showed to be a strong barrier to gene flow. This is probably related to

the specific habitat composition and landscape context for these dune areas: Doornpanne has

relatively much encroached area (trees and shrubs), while Cabour is the dune area most embedded

in agricultural land (Fig. 1).

Table 1: Top five bootstrap results for the multi-categorical model optimisations for the complete study area for the two independent optimisation runs. Full table with all

440 127 models (and null model) can be found in online resource 3.

| Independent run | Model | k | Bootstrap-derived metrics for model performance | | | | | | | Optimised resistance values | | |
			avg.AICc	ΔAICc	avg.weight	avg.rank	avg.R^{2}m	avg.LL	n.top	Res.value 1st	Res.value 2nd	Res.value other
	urban	3	239704.83	0.00	0.10	4.50	0.00010	-119849.38	326	1	-	2500
	urban.water	4	239705.86	1.03	0.06	7.47	0.00015	-119848.88	94	8	1	2500
Run 1	Distance	2	239706.03	1.20	0.05	6.75	0.00001	-119851.00	180	-	-	-
	beach.scrub	4	239706.67	1.84	0.05	11.92	0.00005	-119849.29	136	1	2500	776
	urban.beach	4	239706.87	2.04	0.04	12.29	0.00010	-119849.38	0	1	2500	2438
	urban	3	239650.74	0	0.10	4.83	0.00011	-119822.34	343	1	-	2500
	water.urban	4	239651.83	1.09	0.06	8.52	0.00016	-119821.87	94	1	2.5	2500
Run 2	Distance	2	239652.16	1.42	0.05	8.43	0.00001	-119824.07	180	-	-	-
	beach.urban	4	239652.78	2.04	0.04	13.16	0.00011	-119822.34	0	2500	1	2488.5
	opend.urban	4	239652.89	2.14	0.03	13.79	0.00011	-119822.39	0	1884	1	2500

Notes: combinations of 7 landscape categories compared in bootstrap analysis after optimisation of multi-categorical models. The two independent optimisation runs (Independent run) have an inverted order of categories, to check convergence and account for a possible bias due to the starting cell values. Urbanised (*urban*), *beach, water, trees*, agriculture (*agric*), *scrub*, open dune (*opend*). Distance is the isolation-by-distance null model: increasing genetic distance with increasing Euclidean geographic

distance. Predictor, landscape category or univariate model; k, number of parameters (number of categories in model + 2); avg.AICc, average AICc across all bootstrap

445   iterations; ΔAICc, difference in avg.AICc compared to the lowest avg.AICc (the best supported model); avg.weight, average Akaike weight across iterations; avg.rank,

average rank across iterations; avg.R^2m, average marginal R^2 across iterations; avg.LL, average log-likelihood across iterations; n.top, number of times the model was the

top model across iterations (does not sum op to 1000, as many more models were considered, see online resource 3); Res.value 1^{st}, the optimised resistance value for the

first mentioned landscape category (e.g. *urban* in the second row); Res.value 2^{nd}, the optimised resistance value for the second mentioned landscape category (e.g. water in

the second row); notice the inverted order of categories in run 2 compared to run 1; Res.value other, optimised resistance value for all else (combined into one landscape

450   variable).

451   Table 2: Summed Akaike weights per category for multi-categorical model optimisations, as part of the multi-

452   model inference. The sum of the weights indicates the importance of a category in all possible multi-categorical

453   optimised rasters to explain the pairwise genetic distances.

Category	Sum weights run 1	Sum weights run 2
urban	0.496	0.501
agric	0.357	0.335
beach	0.356	0.376
trees	0.338	0.307
water	0.297	0.315
scrub	0.286	0.259
opend	0.246	0.258

Notes: the Akaike weight for an optimised model was added to the summation (Sum weights) if the focal

category (Category) was in it.

Figure 2: Relative resistance values from the optimisation of multi-categorical models for the complete study

area, as part of the multi-model inference. Anthropogenic landscape categories (*urban*, *water*,

agric=agriculture) are in the top row, natural landscape categories (*beach*, *scrub*, *trees*, *opend*=open dune) in

the bottom row. The two independent runs are combined as paired boxplots for each category in each panel

(left run 1, right run 2). Resistance values are rescaled for each focal category: as optimised resistance values

depend on the other categories included in a model, all models that include the focal category are selected.

The focal category is then held constant by subtracting its optimised resistance value in each model from the

optimised resistance values of the other models (resistance values - resistance value of focal category). These

rescaled values are then plotted in boxplots for each focal category. Each panel should then be interpreted by

checking what lies above zero (category more resistant than focal category) and below zero (category more

facilitating than focal category). The red dashed line is zero, the boundary of equal resistance values between

the focal category (panel) and other categories. For example, in the *urban* panel (left upper corner), the

medians of *beach*, *scrub*, *trees* and *open dune* are situated above the red line, consequently, those four

categories are in general more resistant than *urban* (*urban* is more facilitating) to gene flow. As most

categories, apart from *urban,* proved to be of little importance to explain pairwise genetic distances (table 2),

the uncertainty on the relative resistance values of these categories should be considered high.

Circuitscape current maps

Summed conductance
-4.6   3.5

· Sampled individuals

A) urban

B) urban + water

C) isolation-by-distance (IBD)

0   1   2   3 km

Figure 3: putative genetic connectivity with Circuitscape current maps for the complete study area for the three best supported models. These maps represent in which grid cells of the landscape the most gene flow passes between the sampled nesting locations (black dots), based on A) best supported model with *urban* as

facilitator, B) 2nd best supported model *urban* and *water* combined, both facilitators and C) 3rd best supported

model Isolation-by-distance (resistance values for all cells was set to 1). The values of the grid cells are log10-

transformed summed conductance (a relative measure with no unit), which is based on the summation of all

current maps between pairwise sampled individuals (black dots). High conductance in dark red, medium

conductance in blue, low conductance in white-transparent. A map with the corresponding land cover

categories can be found in figure 1. Map made with QGIS v3.22 (QGIS Development Team 2020). Background

aerial photographs (summer 2018) from Agency for Information Flanders (source: geopunt.be).

Discussion

We used landscape genetics to unravel patterns of functional connectivity of the digger wasp *Bembix rostrata* in a human-altered coastal region. Despite the species' strong gene flow at regional scales and overall pattern of isolation-by-distance (Batsleer et al. 2024), we were able to discern signals of the landscape on the genetic patterns within the coastal dune metapopulation. Contrary to expectations, we found a consistent signal of urban features facilitating gene flow. In general, anthropogenic landscape categories (urban, agriculture, water) seem more facilitating than natural habitat types in this human-altered dune landscape context.

In line with the previous study at larger spatial scales (Batsleer et al. 2024), isolation-by-distance (IBD) remained a prominent pattern of the species' landscape genetics. In itself, an IBD pattern is not in contradiction with an overall high genetic connectivity or an effect of the landscape (Keyghobadi et al. 1999). An IBD pattern emerges naturally when dispersal is limited at a certain spatial scale, i.e. dispersal is more likely to nearby populations than distant ones, independent of the landscape (Kimura and Weiss 1964). Consequently, IBD is seen as a biological realistic null model (opposed to panmictic gene flow) in landscape genetics when considering the influence of the landscape on gene flow, i.e. isolation-by-resistance (IBR) (Wagner and Fortin 2015). Our results showed that IBD was consistently a competing model and one of the best supported models (table 1, considering ΔAICc < 2 with *urban*). Because overall support for models was low (low values of ΔAICc's), we applied multi-model inference to deduce average weights and relative resistance values of categories (Johnson and Omland 2004). This confirmed the importance of the urban category for discriminating pairwise genetic distances and deduce that generally anthropogenic landscapes seem more facilitating than

natural habitat types. Therefore, considering all possible combinations for landscape resistance

optimisation can be powerful when landscape resistance signals are weak and hard to detect.

The current maps in figure 3 visualise the putative realised conductance of grid cells based on all

sample locations according to the three best supported and competing models (*urban*, *urban-water*,

IBD; table 1). These current maps are derived from the optimised resistance raster and serve as post

hoc visualisations of the inferred relationship between genetic distance and resistance distance. In

this context, the few isolated stretches of road with exceptionally high conductance values (Fig. 3B,

3C) are likely artifacts of this post hoc approach, resulting in overly inflated values. Therefore, the

primary focus should be on the overall patterns. In general, we can see from these current maps that

the best-supported models should be considered as complementary rather than opposing. First,

waterways often run parallel to roads and waterbodies are embedded along or within urban areas

(Fig. 1). Such water—which is also a scarce landscape type—adds only a small extra effect on top of

the urban areas for the resulting current map (Fig. 3B). Consequently, we cannot easily disentangle

the effect of water from that of urban within the given landscape (Angelone et al. 2011; Keller et al.

2012). The Akaike weights from multi-model inference (table 2) indicate that the urban category is

overall the most important to explain pairwise genetic distances. Second, the IBD pattern (Fig. 3C) is

also present within the dune areas in the current map for the *urban* and *urban-water* models (high

conductance around sampled individuals; Fig. 3A, 3B). In the latter models, the natural habitat

categories (*open dune, scrub, trees, beach*) all have the same resistant values (relatively higher than

urban and/or water; table 1), resulting in a homogenous resistance landscape within dunes, resulting

in, on average, an IBD pattern. The importance of IBD on this scale is also confirmed by the analyses

repeated on the dune area clusters within our study area (online resource 4). This interpretation is

also in agreement with the multi-model inference, as natural habitat categories proved to be of little

importance to explain pairwise genetic distances (table 2) and the uncertainty on the relative

resistance values of these categories was high (Fig. 2). Consequently, IBD can be considered as the

main background process, wherein gene flow depends on the pairwise, straight-line distances. The

extra signal from anthropogenic features is then at play between dune areas where especially urban

areas give a weak but consistent extra signal of conductance.

Contrary to our hypothesis, we found that gene flow of a specialised digger wasp nesting in grey

dune habitat was not facilitated by typical beach and dune habitats, but rather by urban features in a

dune landscape. This finding does not align with other landscape genetics studies on specialist

insects, for which non-habitat landscape types were mostly barriers (Pérez-Espona et al. 2012;

Trense et al. 2021). Importantly, urban features, although found to be facilitating to gene flow,

should not be interpreted as being favourable for this specialist dune insect. Patterns of gene flow

only reflect a certain component of movement in a species and interpretation of results in a larger

ecological framework should consider other crucial components of a species survival and resource

requirements (Spear et al. 2010; Cushman and Lewis 2010). As *B. rostrata* is a dune-specialist insect,

urban areas should be considered as inhospitable environments. As the deduced resistance values

per category are relative values (Fig. 2), the pattern could be explained by both slower, confined

movement in dune habitat and/or faster, compensatory movement in inhospitable areas. Indeed,

search behaviour in unfamiliar environments can be more directional and faster than in natural,

familiar habitats, where movements are more exploratory (Van Dyck and Baguette 2005; Schtickzelle

et al. 2006; Knowlton and Graham 2010). Suitable habitat intervening the landscape might hinder

gene flow between more distant populations, as individuals are likely to settle at encountered

habitat during dispersal (Adriaensen et al. 2003; McRae et al. 2008; Keller et al. 2012). On the other

hand, compensatory movement through inhospitable habitats can result from a lack of resources or

can be a strategy to reduce mortality risk (Schtickzelle et al. 2006; Peterman et al. 2014). Both can be

counterbalanced by moving faster in inhospitable environments, resulting in increased genetic

connectivity between fragmented habitats (Schtickzelle et al. 2006).

The deduced resistance values are values per grid cell and relative compared to the other considered

landscape types. Consequently, the deduced resistance values of a category will depend on the

habitat composition and the spatial configuration of how populations are embedded within the

landscape (Richardson et al. 2016; Haran et al. 2017). In our study, urbanisation induces large-scale

fragmentation compared to more natural processes like shrub and woodland encroachment. Hence,

a low resistance of urban infrastructure does not conflict with an increased isolation-by-distance in a

more fragmented dune landscape, resulting in lower amounts of gene flow between isolated dune

fragments. Differences in a species' ecology (and evolutionary history) across regions could further

complicate and diversify possible responses to a certain landscape type in different landscapes,

especially at larger scales and across regions (Segelbacher et al. 2010; Spear et al. 2010). Therefore,

results regarding the influence of anthropogenic landscape types on the gene flow of *B. rostrata*

cannot easily be extrapolated or generalised. Nevertheless, our results for this species will

reasonably remain valid for similar landscapes at comparable scales and with similar compositions,

such as several human-altered coastal landscapes of northwest Europe. But extrapolation to inland

sandy regions, where the species occurs as well, would be invalid. There, the degree of isolation

between habitat patches (Batsleer et al. 2024) might decrease the tendency of the species to cross

inhospitable matrix, increasing the resistance values (Schtickzelle et al. 2006).

We find that the connectivity of *B. rostrata* does not seem limited by the landscape composition

within dune clusters and between the different dune entities. Even on the contrary, on top of the

general process of IBD, inhospitable urban features seem to facilitate gene flow in this landscape for

a given distance between different habitats. This indicates that for *B. rostrata*'s persistence in this

study area, landscape-based conservation measures aimed at increasing connectivity between dune

areas (e.g. with habitat corridors within and/or peripheral to the urbanised regions) would not be an

effective management measure unless they are able to give rise to the establishment of new

populations. It is probably more critical to increase or maintain large population sizes, by increasing

habitat area and habitat quality (Richardson et al. 2016; Watts et al. 2016). The effectiveness of

increasing connectivity in this landscape might be worth more consideration for cursorial species

experiencing high mortality costs when crossing urban infrastructure (e.g., natterjack toad, Cox et al.

2017). However, benefits gained from landscape-scale and site-based conservation should always be

balanced to maximise biodiversity benefits with the limited resources at hand (Watts et al. 2016). For

the conservation of *B. rostrata*, other processes linked with dune habitat fragmentation are anyway

more fundamental to consider than landscape connectivity, such as the decrease in habitat area

itself and shifts in ecological processes (e.g. edge effects and 'ecosystem decay' such as inhibition of

sand dynamics in dunes; Pfeifer et al. 2017; Chase et al. 2020). This fragmentation due to

urbanisation decreases the overall number of potentially suitable locations, increases isolation and

affects regional levels of exchange among populations in a spatially altered population network

(Cheptou et al. 2017). Since urban areas are not a barrier to direct gene flow, these more indirect

effects of fragmentation should be the focus to secure persistence of *B. rostrata* in the human-

altered dune landscape. In the contemporary context of dune nature management, a site-based

conservation approach to maintain and improve open dune habitat is recommended, for which

grazing with large herbivores (cattle, horses) is a crucial management tool. However, a reconciliation

is needed between short-term negative effects of trampling by grazers on nest densities of *B.*

rostrata and long-term positive effects of maintaining open dune vegetations within dune areas.

Therefore, a local adaptive management approach with sheep grazing is recommended (Bonte 2005;

Batsleer et al. 2022b). Additionally, improving habitat quality of open dune vegetations through

landscape-scale conservation aimed at restoring and increasing sand dynamics—rather than

connectivity—would be a larger scale, sustainable measure for the persistence of *B. rostrata* in the

coastal dunes in Belgium.

Statements & Declarations

Acknowledgements

We thank the following persons and instances for permission and access to nature reserves: Johan Lamaire, Guy Vileyn, Koen Maertens, Evy Dewulf and Klaar Meulebrouck from ANB (Agency for Nature and Forests – Flemish government); Rika Driessens from IWVA/Aquaduin. We thank two anonymous reviewers for their constructive peer-review comments.

The computational resources (Stevin Supercomputer Infrastructure) and services used in this work were provided by the VSC (Flemish Supercomputer Center), funded by Ghent University, FWO and the Flemish Government – department EWI.

Funding

F.B. was supported by Research Foundation – Flanders (FWO).

Competing interests

The authors have no relevant financial or non-financial interests to disclose.

Author contributions

All authors contributed to the study conception and design. Data collection and laboratory work were performed by FB. Data analyses were done by FB and FD. The first draft of the manuscript was written by FB and all authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

Data availability

Scripts and data are made available on github (https://github.com/FemkeBatsleer/LandGenBembix) for review and will be published on zenodo upon acceptance.

References

Adriaensen F, Chardon JP, De Blust G, et al (2003) The application of 'least-cost' modelling as a functional landscape model. Landscape and Urban Planning 64:233–247. https://doi.org/10.1016/S0169-2046(02)00242-6

Angelone S, Kienast F, Holderegger R (2011) Where movement happens: scale-dependent landscape effects on genetic differentiation in the European tree frog. Ecography 34:714–722. https://doi.org/10.1111/j.1600-0587.2010.06494.x

Baguette M, Van Dyck H (2007) Landscape connectivity and animal behavior: functional grain as a key determinant for dispersal. Landscape Ecol 22:1117–1129. https://doi.org/10.1007/s10980-007-9108-4

Balkenhol N, Cushman SA, Storfer AT, Waits LP (eds) (2015) Landscape genetics: concepts, methods, applications. John Wiley & Sons Ltd

Bates D, Mächler M, Bolker B, Walker S (2015) Fitting Linear Mixed-Effects Models Using **lme4**. J Stat Soft 67:. https://doi.org/10.18637/jss.v067.i01

Batsleer F, Bonte D, Dekeukeleire D, et al (2020) The neglected impact of tracking devices on terrestrial arthropods. Methods Ecol Evol 11:350–361. https://doi.org/10.1111/2041-210X.13356

Batsleer F, Gallin M, Delafonteyne M, et al (2024) Asymmetrical gene flow between coastal and inland dunes in a threatened digger wasp. Conserv Genet 25:259–275. https://doi.org/10.1007/s10592-023-01566-7

Batsleer F, Maes D, Bonte D (2022a) Behavioral Strategies and the Spatial Pattern Formation of Nesting. Am Nat 199:E15–E27. https://doi.org/10.1086/717226

Batsleer F, Van Uytvanck J, Lamaire J, et al (2022b) Rapid conservation evidence for the impact of sheep grazing on a threatened digger wasp. Insect Conserv Diversity 15:149–156. https://doi.org/10.1111/icad.12532

Bogusch P, Heneberg P, Šilhán K (2021) What are the main factors limiting the distribution of *Bembix rostrata* (Hymenoptera: Crabronidae) at early-succession sites? J Insect Conserv 25:571–583. https://doi.org/10.1007/s10841-021-00324-9

Bonte D (2005) Anthropogenic induced changes in nesting densities of the dune-specialised digger wasp *Bembix rostrata* (Hymenoptera: Sphecidae). Eur J Entomol 102:809–812. https://doi.org/10.14411/eje.2005.114

Bonte D, Batsleer F, Provoost S, et al (2021) Biomorphogenic Feedbacks and the Spatial Organization of a Dominant Grass Steer Dune Development. Front Ecol Evol 9:761336. https://doi.org/10.3389/fevo.2021.761336

Bonte D, Maes D (2008) Trampling affects the distribution of specialised coastal dune arthropods.
Basic Appl Ecol 9:726–734. https://doi.org/10.1016/j.baae.2007.09.008

Bonte D, Van Dyck H, Bullock JM, et al (2012) Costs of dispersal. Biol Rev 87:290–312.
https://doi.org/10.1111/j.1469-185X.2011.00201.x

Buckles BJ, Harmon-Threatt AN (2019) Bee diversity in tallgrass prairies affected by management and
its effects on above- and below-ground resources. J Appl Ecol 56:2443–2453.
https://doi.org/10.1111/1365-2664.13479

Burnham KP, Anderson DR (eds) (2004) Model Selection and Multimodel Inference. Springer New
York, New York, NY

Burnham KP, Anderson DR, Huyvaert KP (2011) AIC model selection and multimodel inference in
behavioral ecology: some background, observations, and comparisons. Behav Ecol Sociobiol
65:23–35. https://doi.org/10.1007/s00265-010-1029-6

Châline N, Ratnieks FLW, Raine NE, et al (2004) Non-lethal sampling of honey bee, *Apis mellifera*,
DNA using wing tips. Apidologie 35:311–318. https://doi.org/10.1051/apido:2004015

Chase JM, Blowes SA, Knight TM, et al (2020) Ecosystem decay exacerbates biodiversity loss with
habitat loss. Nature 584:238–243. https://doi.org/10.1038/s41586-020-2531-2

Cheptou P-O, Hargreaves AL, Bonte D, Jacquemyn H (2017) Adaptation to fragmentation:
evolutionary dynamics driven by human influences. Phil Trans R Soc B 372:20160037.
https://doi.org/10.1098/rstb.2016.0037

Clark JA, May RM (2002) Taxonomic Bias in Conservation Research. Science 297:191–192.
https://doi.org/10.1126/science.297.5579.191b

Clarke RT, Rothery P, Raybould AF (2002) Confidence limits for regression relationships between
distance matrices: Estimating gene flow with distance. JABES 7:361–372.
https://doi.org/10.1198/108571102320

Cox K, Maes J, Van Calster H, Mergeay J (2017) Effect of the landscape matrix on gene flow in a
coastal amphibian metapopulation. Conserv Genet 18:1359–1375.
https://doi.org/10.1007/s10592-017-0985-z

Crooks KR, Sanjayan M (2006) Connectivity conservation. Cambridge University Press

Cushman SA, Landguth EL (2010) Scale dependent inference in landscape genetics. Landscape Ecol
25:967–979. https://doi.org/10.1007/s10980-010-9467-0

Cushman SA, Lewis JS (2010) Movement behavior explains genetic differentiation in American black
bears. Landscape Ecol 25:1613–1625. https://doi.org/10.1007/s10980-010-9534-6

David G, Giffard B, Piou D, Jactel H (2014) Dispersal capacity of *Monochamus galloprovincialis* , the
692   European vector of the pine wood nematode, on flight mills. J Applied Entomology 138:566–
693   576. https://doi.org/10.1111/jen.12110

694   De Ro A, Vanden Broeck A, Verschaeve L, et al (2021) Occasional long-distance dispersal may not
695   prevent inbreeding in a threatened butterfly. BMC Ecol Evo 21:224.
696   https://doi.org/10.1186/s12862-021-01953-z

697   Decleer K (2007) Europees beschermde natuur in Vlaanderen en het Belgisch deel van de Noordzee.
698   Habitattypen | Dier- en plantensoorten. Mededelingen van het Instituut voor Natuur- en
699   Bosonderzoek INBO.M.2007.01. Instituut voor Natuur- en Bosonderzoek, Brussels

700   Develey PF, Stouffer PC (2008) Effects of Roads on Movements by Understory Birds in Mixed-Species
701   Flocks in Central Amazonian Brazil. Conservation Biology 15:1416–1422.
702   https://doi.org/10.1111/j.1523-1739.2001.00170.x

703   Didham RK, Basset Y, Collins CM, et al (2020) Interpreting insect declines: seven challenges and a way
704   forward. Insect Conserv Divers 13:103–114. https://doi.org/10.1111/icad.12408

705   Drake J, Lambin X, Sutherland C (2022) The value of considering demographic contributions to
706   connectivity: a review. Ecography 2022:. https://doi.org/10.1111/ecog.05552

707   Driscoll DA, Banks SC, Barton PS, et al (2013) Conceptual domain of the matrix in fragmented
708   landscapes. Trends in Ecology & Evolution 28:605–613.
709   https://doi.org/10.1016/j.tree.2013.06.010

710   Evans HE, O'Neill KM (2007) The sand wasps: natural history and behavior. Harvard University Press,
711   Cambridge, MA, United States

712   Ferraro PJ, Pattanayak SK (2006) Money for Nothing? A Call for Empirical Evaluation of Biodiversity
713   Conservation Investments. PLoS Biol 4:e105. https://doi.org/10.1371/journal.pbio.0040105

714   Field J, Gonzalez-Voyer A, Boulton RA (2020) The evolution of parental care strategies in subsocial
715   wasps. Behav Ecol Sociobiol 74:78. https://doi.org/10.1007/s00265-020-02853-w

716   Goslee SC, Urban DL (2007) The **ecodist** Package for Dissimilarity-based Analysis of Ecological Data. J
717   Stat Soft 22:. https://doi.org/10.18637/jss.v022.i07

718   Hanski I (1998) Metapopulation dynamics. Nature 396:41–49. https://doi.org/10.1038/23876

719   Haran JM, Rossi J-P, Pajares J, et al (2017) Multi-scale and multi-site resampling of a study area in
720   spatial genetics: implications for flying insect species. PeerJ 5:e4135.
721   https://doi.org/10.7717/peerj.4135

722   Hedenström A, Ellington CP, Wolf TJ (2001) Wing wear, aerodynamics and flight energetics in
723   bumblebees (*Bombus terrestris*): an experimental study. Functional Ecology 15:417–422.
724   https://doi.org/10.1046/j.0269-8463.2001.00531.x

Hein S, Gombert J, Hovestadt T, Poethke H-J (2003) Movement patterns of the bush cricket *Platycleis*
albopunctata in different types of habitat: matrix is not always matrix: Movement patterns of
P. albopunctata. Ecological Entomology 28:432–438. https://doi.org/10.1046/j.1365-
2311.2003.00531.x

Hijmans RJ (2024) terra: Spatial Data Analysis. R package version 1.7-78, https://CRAN.R-
project.org/package=terra

Hodgson JA, Randle Z, Shortall CR, Oliver TH (2022) Where and why are species' range shifts
hampered by unsuitable landscapes? Glob Chang Biol 28:4765–4774.
https://doi.org/10.1111/gcb.16220

Hodgson JA, Thomas CD, Wintle BA, Moilanen A (2009) Climate change, connectivity and
conservation decision making: back to basics. Journal of Applied Ecology 46:964–969.
https://doi.org/10.1111/j.1365-2664.2009.01695.x

Humphrey JW, Watts K, Fuentes-Montemayor E, et al (2015) What can studies of woodland
fragmentation and creation tell us about ecological networks? A literature review and
synthesis. Landscape Ecol 30:21–50. https://doi.org/10.1007/s10980-014-0107-y

Johnson JB, Omland KS (2004) Model selection in ecology and evolution. Trends in Ecology &
Evolution 19:101–108. https://doi.org/10.1016/j.tree.2003.10.013

Jombart T (2008) adegenet: a R package for the multivariate analysis of genetic markers.
Bioinformatics 24:1403–1405. https://doi.org/10.1093/bioinformatics/btn129

Keller D, Van Strien MJ, Holderegger R (2012) Do landscape barriers affect functional connectivity of
populations of an endangered damselfly?: Functional connectivity in a damselfly. Freshwater
Biology 57:1373–1384. https://doi.org/10.1111/j.1365-2427.2012.02797.x

Keyghobadi N, Roland J, Strobeck C (1999) Influence of landscape on the population genetic structure
of the alpine butterfly Parnassius smintheus (Papilionidae). Mol Ecol 8:1481–1495.
https://doi.org/10.1046/j.1365-294x.1999.00726.x

Kimmig SE, Beninde J, Brandt M, et al (2020) Beyond the landscape: Resistance modelling infers
physical and behavioural gene flow barriers to a mobile carnivore across a metropolitan area.
Mol Ecol 29:466–484. https://doi.org/10.1111/mec.15345

Kimoto C, DeBano SJ, Thorp RW, et al (2012) Short-term responses of native bees to livestock and
implications for managing ecosystem services in grasslands. Ecosphere 3:88.
https://doi.org/10.1890/ES12-00118.1

Kimura M, Weiss GH (1964) The stepping stone model of population structure and the decrease of
genetic correlation with distance. Genetics 49:561–576.
https://doi.org/10.1093/genetics/49.4.561

Klein WF, Lefeber V (2004) Crabronidae – graafwespen. In: De wespen en mieren van Nederland (ed.
by T.M.J. Peeters). Naturalis en KNNV, Utrecht, The Netherlands, pp 356–430

Knowlton JL, Graham CH (2010) Using behavioral landscape ecology to predict species' responses to
land-use and climate change. Biological Conservation 143:1342–1354.
https://doi.org/10.1016/j.biocon.2010.03.011

Kuefler D, Hudgens B, Haddad NM, et al (2010) The conflicting role of matrix habitats as conduits and
barriers for dispersal. Ecology 91:944–950. https://doi.org/10.1890/09-0614.1

Larsson FK (1986) Increased nest density of the digger wasp *Bembix rostrata* as a response to
parasites and predators (Hymenoptera: Sphecidae). Entomol Gener 12:71–75

Larsson FK, Tengö J (1989) It is not always good to be large; some female fitness components in a
temperate digger wasp, *Bembix rostrata* (Hymenoptera: Sphecidae). J Kans Entomol Soc
62:490–495

Logghe G, Taelman C, Van Hecke F, et al (2024). Unravelling arthropod movement in natural
landscapes: Small-scale effects of body size and weather conditions. Journal of Animal Ecology 93:
1365–1379. https://doi.org/10.1111/1365-2656.14161

Lourenço A, Gonçalves J, Carvalho F, et al (2019) Comparative landscape genetics reveals the
evolution of viviparity reduces genetic connectivity in fire salamanders. Molecular Ecology
28:4573–4591. https://doi.org/10.1111/mec.15249

Maes D, Bonte D (2006) Using distribution patterns of five threatened invertebrates in a highly
fragmented dune landscape to develop a multispecies conservation approach. Biological
Conservation 133:490–499. https://doi.org/10.1016/j.biocon.2006.08.001

Manel S, Holderegger R (2013) Ten years of landscape genetics. Trends in Ecology & Evolution
28:614–621. https://doi.org/10.1016/j.tree.2013.05.012

Manel S, Schwartz MK, Luikart G, Taberlet P (2003) Landscape genetics: combining landscape ecology
and population genetics. Trends in Ecology & Evolution 18:189–197.
https://doi.org/10.1016/S0169-5347(03)00008-9

McRae BH (2006) Isolation by resistance. Evolution 60:1551–1561. https://doi.org/10.1111/j.0014-
3820.2006.tb00500.x

McRae BH, Dickson BG, Keitt TH, Shah VB (2008) Using circuit theory to model connectivity in
ecology, evolution, and conservation. Ecology 89:2712–2724. https://doi.org/10.1890/07-
1861.1

McRae BH, Shah VB, Mohapatra TK (2013) Circuitscape 4 User Guide

Nielsen ET (1945) Moeurs des Bembex. Spoolia Zoologica Musei Hauniensis VII. Universitetets
Zoologiske Museum, København, Denmark

Ovaskainen O, Luoto M, Ikonen I, et al (2008) An Empirical Test of a Diffusion Model: Predicting
Clouded Apollo Movements in a Novel Environment. The American Naturalist 171:610–619.
https://doi.org/10.1086/587070

Pérez-Espona S, McLeod JE, Franks NR (2012) Landscape genetics of a top neotropical predator. Mol
Ecol 21:5969–5985. https://doi.org/10.1111/mec.12088

Peterman WE (2018) ResistanceGA: An R package for the optimization of resistance surfaces using
genetic algorithms. Methods Ecol Evol 9:1638–1647. https://doi.org/10.1111/2041-
210X.12984

Peterman WE, Connette GM, Semlitsch RD, Eggert LS (2014) Ecological resistance surfaces predict
fine-scale genetic differentiation in a terrestrial woodland salamander. Mol Ecol 23:2402–
2413. https://doi.org/10.1111/mec.12747

Pfeifer M, Lefebvre V, Peres CA, et al (2017) Creation of forest edges has a global impact on forest
vertebrates. Nature 551:187–191. https://doi.org/10.1038/nature24457

Potts SG, Imperatriz-Fonseca V, Ngo HT, et al (2016) Safeguarding pollinators and their values to
human well-being. Nature 540:220–229. https://doi.org/10.1038/nature20588

Provoost S, Ampe C, Bonte D, et al (2004) Ecology, management and monitoring of grey dunes in
Flanders. J Coast Conserv 10:33–42. https://doi.org/10.1652/1400-
0350(2004)010[0033:EMAMOG]2.0.CO;2

Provoost S, Bonte D (2004) Levende duinen: Een overzicht van de biodiversiteit aan de Vlaamse kust.
Instituut voor Natuurbehoud: Brussel

Provoost S, Jones MLM, Edmondson SE (2011) Changes in landscape and vegetation of coastal dunes
in northwest Europe: a review. J Coast Conserv 15:207–226. https://doi.org/10.1007/s11852-
009-0068-5

QGIS Development Team (2020) QGIS Geographic Information System

R Core Team (2020) R: A Language and Environment for Statistical Computing

Richardson JL, Brady SP, Wang IJ, Spear SF (2016) Navigating the pitfalls and promise of landscape
genetics. Mol Ecol 25:849–863. https://doi.org/10.1111/mec.13527

RStudio Team (2022) RStudio: Integrated Development Environment for R

Samways MJ, Barton PS, Birkhofer K, et al (2020) Solutions for humanity on how to conserve insects.
Biological Conservation 242:108427. https://doi.org/10.1016/j.biocon.2020.108427

Schöne H, Tengö J (1981) Competition of males, courtship behaviour and chemical communication in
825   the digger wasp *Bembix rostrata* (Hymenoptera, Sphecidae). Behaviour 77:44–66

826   Schtickzelle N, Mennechez G, Baguette M (2006) Dispersal depression with habitat fragmentation in
827   the Bog fritillary butterfly. Ecology 87:1057–1065. https://doi.org/10.1890/0012-
828   9658(2006)87[1057:DDWHFI]2.0.CO;2

829   Scrucca L (2013) GA: A Package for Genetic Algorithms in R. Journal of Statistical Software 37

830   Segelbacher G, Cushman SA, Epperson BK, et al (2010) Applications of landscape genetics in
831   conservation biology: concepts and challenges. Conserv Genet 11:375–385.
832   https://doi.org/10.1007/s10592-009-0044-5

833   Shirk AJ, Landguth EL, Cushman SA (2017) A comparison of individual-based genetic distance metrics
834   for landscape genetics. Mol Ecol Resour 17:1308–1317. https://doi.org/10.1111/1755-
835   0998.12684

836   Spear SF, Balkenhol N, Fortin M-J, et al (2010) Use of resistance surfaces for landscape genetic
837   studies: considerations for parameterization and analysis. Molecular Ecology 19:3576–3591.
838   https://doi.org/10.1111/j.1365-294X.2010.04657.x

839   Sutherland WJ, Pullin AS, Dolman PM, Knight TM (2004) The need for evidence-based conservation.
840   Trends in Ecology & Evolution 19:305–308. https://doi.org/10.1016/j.tree.2004.03.018

841   Trense D, Schmidt TL, Yang Q, et al (2021) Anthropogenic and natural barriers affect genetic
842   connectivity in an Alpine butterfly. Mol Ecol 30:114–130. https://doi.org/10.1111/mec.15707

843   Ugelvig LV, Andersen A, Boomsma JJ, Nash DR (2012) Dispersal and gene flow in the rare, parasitic
844   Large Blue butterfly *Maculinea arion*: DISPERSAL OF THE LARGE BLUE BUTTERFLY. Molecular
845   Ecology 21:3224–3236. https://doi.org/10.1111/j.1365-294X.2012.05592.x

846   Van Dyck H, Baguette M (2005) Dispersal behaviour in fragmented landscapes: Routine or special
847   movements? Basic Appl Ecol 6:535–545. https://doi.org/10.1016/j.baae.2005.03.005

848   van Etten J (2017) *R* Package **gdistance**: Distances and Routes on Geographical Grids. J Stat Soft 76:.
849   https://doi.org/10.18637/jss.v076.i13

850   van Klink R, van der Plas F, van Noordwijk CGE (Toos), et al (2015) Effects of large herbivores on
851   grassland arthropod diversity. Biol Rev 90:347–366. https://doi.org/10.1111/brv.12113

852   Wagner DL, Grames EM, Forister ML, et al (2021) Insect decline in the Anthropocene: Death by a
853   thousand cuts. PNAS 118:e2023989118. https://doi.org/10.1073/pnas.2023989118

854   Wagner HH, Fortin M-J (2015) Basics of Spatial Data Analysis: Linking Landscape and Genetic Data for
855   Landscape Genetic Studies. In: Balkenhol N, Cushman SA, Storfer AT, Waits LP (eds)
856   Landscape Genetics: concepts, methods and applications. John Wiley & Sons, Ltd, pp 77–98

Watts K, Vanhala T, Connolly T, Cottrell J (2016) Striking the right balance between site and
landscape-scale conservation actions for a woodland insect within a highly fragmented
landscape: A landscape genetics perspective. Biological Conservation 195:146–155.
https://doi.org/10.1016/j.biocon.2015.12.039

Wickham H (2016) ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York

Wiklund C, Fagerström T (1977) Why do males emerge before females? A hypothesis to explain the
incidence of protandry in butterflies. Oecologia 31:153–158.
https://doi.org/10.1007/BF00346917

Wilcock HR, Bruford MW, Nichols RA, Hildrew AG (2007) Landscape, habitat characteristics and the
genetic population structure of two caddisflies. Freshwater Biol 52:1907–1929.
https://doi.org/10.1111/j.1365-2427.2007.01818.x

Zayed A (2009) Bee genetics and conservation. Apidologie 40:237–262.
https://doi.org/10.1051/apido/2009026

Supplementary information

S1 Landscape types_1.pdf: detailed description of the how landscape categories from original

sources were combined and reclassified.

S2 Single category optimisation runs_2.pdf: results for the optimisation and bootstrap analyses of the

7 single category rasters for 4 independent runs.

S3 bootstrapsummary_westcoast_3.xlsx: full results of bootstrapping of all possible multi-categorical

models after optimisation of the complete study area.

S4 dune area clusters_4.pdf: detailed results for the analyses repeated on the 4 dune area clusters

situated within the complete study area.

S5 bootstrapsummary_areasdunes_5.xlsx: full results of the bootstrapping of all possible multi-

categorical models after optimisation dune area clusters situated within the study area.