1 **Bird communities in a selectively logged tropical montane forest are dominated by small, low-**
2 **elevation species**
3

4 Ritobroto Chanda[1*], Shambu Rai[1], Bharat Tamang[1], Binod Munda[1], D.K. Pradhan[1], Mangal Rai[1],
5 Aman Biswakarma[1], Umesh Srinivasan[1]
6
7 [1] Centre for Ecological Sciences, Indian Institute of Science, Bengaluru 560 012, India
8
9 * Corresponding author: Ritobroto Chanda, ritobrotochanda@gmail.com

10
11
12

13 **Abstract**
14

Tropical montane forests are critical centres of terrestrial biodiversity and endemism, and face a
range of threats, including selective logging and climate change. However, few studies have
explored the joint influence of these threats, particularly in tropical mountains, despite how crucial
it is to understand the cumulative impact of forest loss and climate change on biodiversity. We used
mist-netting and bird ringing data from a long-term (10-year) community and population
monitoring program to examine how the composition of the mid-elevation Eastern Himalayan
understorey bird community changed in the mid-elevation primary and logged forest. Because
logged forest is warmer than primary forest, we hypothesised that the bird community would shift
towards lower elevation species, with a faster transition in logged forest. Further, because logged
forest has lower arthropod abundance, we hypothesised that the primary forest community would
have larger species than the logged forest community. Our study shows that the bird community in
the logged forest shifted towards lower-elevation species than in the primary forest (although this
was not statistically significant). Moreover, we found that smaller species were better able to
colonise warmer logged forest, whereas larger species reached higher densities in primary forest.
These trends are likely to be driven by climate change causing upslope range shifts and logged
forest selecting for species that are likely to tolerate the consequently altered abiotic (higher
temperatures) and biotic (lower resources) environment. These findings have significant
implications for understanding the impacts of forest loss and climate change on biodiversity,
particularly in tropical montane forests.

34
35

36 *Keywords*: Bird community, Selective logging, Climate change, Body mass, Elevational range
37 shifts, Montane forests

Introduction

Tropical montane forests are the richest centres of terrestrial biodiversity and endemism globally (Rahbek et al., 2019). These forests face multiple threats (Bradshaw et al. 2009), of which climate change and habitat degradation/conversion are major drivers of biodiversity declines (Asner et al., 2009; Edwards and Laurance, 2013; Constantini, Edwards, & Simons, 2016; Burivalova et al., 2014; Dawson et al., 2011; Corlett, 2012).

While the separate impacts of habitat loss and climate change on biodiversity have been studied extensively (Bellard et al., 2012; Matthews et al., 2014; Fahrig, 2003), few studies have examined the influence of both these threats together (Manytka-Pringle et al., 2012; Guo et al., 2018; Srinivasan and Wilcove, 2021), especially in the tropics, and particularly in tropical mountains. With climate change and forest loss accelerating dramatically (Mitchard, 2018; Arneth et al., 2020), understanding their cumulative impacts on biodiversity is essential. This is especially true because forest degradation (e.g., from selective logging) alters forest structure and creates novel habitats that tend to be hotter and drier than primary forest (Srinivasan and Wilcove, 2021), complicating climate impacts. Further, tropical species are thermal specialists and poor dispersers and tend to have narrow elevational ranges, and how climate- *and* land-use-driven changes in the abiotic environment might impact such species remains poorly known (Janzen, 1967; McCain, 2009; Sheard et al., 2020).

Tropical montane species are shifting their ranges to higher elevations rapidly, with strong evidence that such upslope shifts are a result of rising temperatures globally (Forero-Medina et al., 2011; Campos-Cerqueira et al., 2017; Freeman et al., 2018; Neate-Clegg et al., 2018, Freeman et al. 2021). As tropical species shift their ranges upslope in response to climate change, populations might either encounter primary forest or degraded forest (e.g., selectively logged forest) at higher elevations. With populations of the same species likely shifting their ranges into primary and logged forest at higher elevations, it is crucial to understand how such range shifts might lead to differences in community assembly in undisturbed versus disturbed forest. Although selectively logged forest can retain more biodiversity than other forms of forest loss/degradation such as outright conversion to agriculture (Putz et al., 2012; Gibson et al., 2011), logging does lead to the altered composition of ecological communities (Constantini, Edwards, & Simons, 2016; Burivalova et al., 2014).

Using mist-netting and bird ringing data from a long-term community and population monitoring program, we asked how the composition of the mid-elevation Eastern Himalayan understorey bird community changed over a ten-year period in primary and in logged forest. At any given elevation, the primary forest bird community should become progressively dominated by historically "low-elevation" species as they move upwards to replace "higher-elevation" species that are moving further upslope. Logged forest being warmer than primary forest (Srinivasan and Wilcove, 2021) should be thermally more similar to the lower-elevation primary forest than to the primary forest at the same elevation. We, therefore, expected that upslope range shifts should lead to faster species turnover from "higher-elevation" to "lower-elevation" in logged than in primary forest. Further, we

expected that smaller species, being more thermally tolerant (Sheridan and Bickford, 2011; Speakman et al., 2003) would be better able to colonize warmer logged forest, whereas larger species would reach higher densities in primary forest.

We predicted that:

1. The bird community would gradually shift towards lower elevation species and the transition would occur faster in the logged forest.
2. The bird community in primary forest would have larger species as compared to the logged forest community.

Methods

Field Methods

We carried out this study in mid-elevation (2000m ASL) montane broadleaved forest (Champion and Seth, 1968) in Eaglenest Wildlife Sanctuary (27°06′0″N 92°24′0″E, EWLS henceforth), Arunachal Pradesh, Eastern Himalaya, India. In this habitat, we established three plots each in primary and logged forest (a total of 9ha each in primary and logged forest; Fig. 1). We selected the plots based on interviews with past logging managers. Intensive logging occurred until 2002 after which it was halted (see Srinivasan 2013). The logged patches have visible differences from the intact forest with a very open canopy and the understorey dominated by bamboo. We sampled each of these plots for three consecutive days from 0500 hrs to 1200 hrs in the early breeding season (April-May) from 2011 to 2021 using 24-28 mist nets (12m length, 2.4m height, 16mm mesh, 4-shelf). In 2020, a nationwide COVID-19-related lockdown precluded any bird sampling. We placed the mist nets in a grid-like pattern within the plot with ~40m distance between neighbouring nets (Fig. 1). We then checked each net every 20-30 minutes and any birds captured were brought back to the ringing station at the centre of the sampling plot. We identified and weighed each bird caught with a jeweller's scale, ringed it with a uniquely numbered aluminium ring and then released the bird immediately.

Figure 1: Map of the study area in Eaglenest Wildlife Sanctuary, Arunachal Pradesh, India. Primary forest and logged forest plots are outlined in green and brown, respectively. Arrows show the locations of mist nets.

Data Curation

We excluded high-elevation breeders which are passage migrants to higher-elevation breeding sites and do not breed in our study area (Table S1), aquatic species, carnivores, canopy dwellers, and non-invertivores (which included frugivores, granivores and nectarivores). We finally analysed the understorey insectivorous bird community of 61 species (Table S1). We pooled the raw counts of each species in each habitat in each year pooled from for the first two days of sampling effort in each site (i.e., six days each in logged and primary forest in each year). We did this because in some years in some plots, rain prevented the full three-day sampling protocol and sampling was limited to two consecutive days.

138
139   **Analytical Methods:**
140
141   *Differences in community composition*
142
We first created a year-by-species matrix in primary forest and logged forest separately with the
raw count data (Table S2) transformed using the Hellinger transformation (Bocard et al., 2011)
which reduces the weightage of rare species in contributing to compositional differences between
communities (Legendre and Gallagher, 2001). We then performed a non-metric multidimensional
scaling (NMDS) ordination analysis to visualise the bird community in logged and primary forests
each year using the Bray-Curtis dissimilarity index on the transformed data in the R package *vegan*
(Okansen et al., 2022; R Core Team 2023). To test whether the bird community composition
differed significantly between primary and logged forest, we used permutational multivariate
analysis of variance (PERMANOVA), again using the Bray-Curtis dissimilarity index.
152
To understand the relative contributions of different species to community-level differences
between primary and logged forest, we used similarity percentage routines (SIMPER, Clarke and
Warwick 2001). From the SIMPER routine, we then selected the top species that cumulatively
contributed to 50% of dissimilarity in bird community composition between primary and logged
forest (Table S3). To visualise the importance of these species to the difference in community
composition across habitats, we plotted the location of these species in NMDS space. We used
SIMPER since it gives us a much clearer picture of the possible patterns than using all of the 61
species, especially rarer species that might have had only a few individuals captured over the study
period and might skew the results disproportionately, appearing to contribute much to dissimilarity
between communities, but not indeed doing so.
163
To understand potential differences between the morphology (i.e., body mass) and distribution (i.e.,
elevational ranges) of species that contributed in large part to community-level differences between
primary and logged forest, we used:
(a) A generalized linear model (GLMs) with habitat type (logged/primary) as the categorical
predictor variable and body mass as the response variable.
(b) An ANOVA to compare the elevation range midpoints of species in primary and logged forest.
170
171   The body mass of the species was averaged from the birds caught in logged and primary forests
172   over the ten years. We extracted the breeding elevation range data of each from Spierenburg (2005)
173   and it was supplemented by data from Srinivasan et al. (2018). We used this data to calculate the
174   elevation range midpoint.

180   *Trends over time*
181

To understand the trends of body mass and elevation range midpoints over time, we came up with
two metrics, community mass index (CMI) and community elevation index (CEI). These were
developed using a metric called the community temperature index (Devictor et al., 2008). We took
the community weighted mean of species mass and the breeding elevation range midpoint to
understand the temporal shift. We carried this out using the top species that cumulatively
contributed to 50% of dissimilarity in bird community composition between primary and logged
forest.
189

$$CMI = \sum_{s=1}^{n} Mass_s \times Abundance_s$$

$$CEI = \sum_{s=1}^{n} Breeding\ elevation\ range\ midpoint_s \times Abundance_s$$

190
191
192
193
194   **Results:**
195

196   We recorded 9,036 captures of 6,189 individuals from 130 species in our mist nets across all years
197   and across all sampling plots. Upon curating the data, we finally analysed 6,926 captures of 4,801
198   individuals of 61 understorey insectivorous species. The raw counts of the 61 species are in Table
199   S2.
200

201   *Community composition in primary and logged forest*
202

NMDS axis 1 separated bird communities in primary and logged habitats (stress = 0.20, Fig. 2).
Primary and logged forest communities were compositionally significantly different from each
other (PERMANOVA ($R^2 = 0.14$, $F_{1,18} = 2.98$, $p < 0.01$). The first 20 species (Table S3) explained
50% of the variation in community composition between primary and logged forest and were used
in subsequent analyses. We did not use *Schoeniparus cinereus* and *Tesia castaneocoronata* out of
these since they did not fall in either primary or logged forest clusters.

Figure 2: Non-metric multidimensional scaling (NMDS) showing dissimilarity between primary and logged forest communities over a ten-year period (see Table S3 for details of species codes and contribution to the SIMPER routine).

Body mass trends in primary and logged forest

Habitats were significantly related to body mass (Poisson ANOVA, $\boldsymbol{\beta}_{primary}$ = 0.26 [0.02, 0.50], McFadden's Pseudo R^2 = 0.04. On average, species typical to primary forest were 1.3g heavier than species typical to logged forest. Median body mass across species was 5g higher in primary forest (Fig. 3).

Figure 3: Mass of species that contributed to 50% of dissimilarity in community composition between primary and logged forests. Species typical to primary forest are, on average, larger than those typical to logged forest. The box shows the middle 50% of the data, with the horizontal line inside the box indicating the median value. The whiskers of the boxplot extend from the box to show the minimum and maximum values of the data. Each dot represents individual data points and is plotted with a jitter to avoid overlap.

237 *Trends of Breeding elevational range midpoints in primary and logged forest*

239 Elevation range midpoints of bird species typical to primary forest was, on average 172.5m [-185.3,
240 530.3; $R^2 = 0.06$] higher than species representative of logged forest. While the mean difference in
241 the mid-points of species' elevational ranges was as expected, the effect of habitat was weak (i.e.,
242 95% CIs overlapping zero).

244
Figure 4: Mean breeding elevation range of the species that contributed to 50% dissimilarity
in each habitat. The box shows the middle 50% of the data, with the horizontal line inside the
box indicating the median value. The whiskers of the boxplot extend from the box to show the
minimum and maximum values of the data. Each dot represents individual data points and
is plotted with a jitter to avoid overlap.

The mean low elevation range limits of species in the primary forest were 175m [-188.9, 538.9; R^2
$= 0.06$] higher than in logged forest. Similarly, the mean high elevation range limit of the species
was higher in primary forest as well by 170 m [-270.3, 610.3; $R^2 = 0.04$], supporting the overall
trend (although weak) of species representative of primary forest being "higher elevation" species
than those representative of logged forest in the mid-elevation montane temperate forest (See Supp.
Figs. S1 and S2).

Trends over time

We tried to look at the trends of mass and mid-point of breeding elevation range over time. We
found major overlaps in the confidence intervals of CEI and CMI over time (see Supp. Figs. S3 and
S4).

263
264
265

Discussion:

We investigated how climate change and selective logging might combine to impact tropical
understorey bird communities in primary and selectively logged forest using a 10-year dataset from
the Eastern Himalayas. We find that the bird community in primary forest was, on average,
composed of larger species than in logged forest (Fig. 3). We also find weak support that the
community in logged forest has a lower mean, minimum and maximum summer elevation range
than the primary forest community, which indicates that the logged forest community might
resemble lower elevation communities (Fig. 4). However, this effect was not as strong as the effect
for body mass. We tried to understand trends of mass and mid-point of breeding elevation range
over time but this seemed to be complicated by natural trends in abundance fluctuations in each
year (Supp. Fig S3 and S4). We can likely see stronger trends with more long-term data.

Logging can lead to the loss of large-bodied, old-growth-dependent species and decrease overall
biodiversity (Cleary et al., 2007). Understorey insectivores, which are often highly specialized, are
negatively influenced by logging and show steep abundance declines (Chaves et al. 2017). Further,
the body size of birds is an important factor affecting their response to logging. Birds with larger
body sizes were found to be less common in logged forests, whereas smaller-bodied species were
found to be more resilient to logging (Burivalova et al., 2015). We find a similar pattern, with
species typical of primary forest being significantly larger than species typical of logged forest (Fig.
2). Smaller-bodied bird species were more abundant in logged forest, while larger-bodied species
declined (Velho et al., 2012, Srinivasan 2013). Such patterns might arise from changes in the abiotic
environment with logging leading to changes in arthropod prey composition and abundance, in turn
resulting in reduced resource availability for birds (Aggarwal et al., 2023). Earlier studies from our
field site have shown that logged forest is warmer than old-growth forest because of changes in
forest structure and microclimates (Bharadwaj et al., 2022). On average, primary forest was found
to be 2.3°C cooler and 14.6% more humid than logged forest (Aggarwal et al., 2023), and have
lower densities of foliage-dwelling arthropods (Aggarwal et al., 2023). Because large species have
higher absolute energy requirements, a reduction in resource availability is likely to
disproportionately reduce the abundance of large species (Srinivasan 2013).

The altered abiotic environment in logged forest can not only reduce the abundance of large species
through indirect impacts on prey availability but can also directly influence the composition of bird
communities in terms of body mass and elevational range. Climate warming is believed to have
driven a decrease in the body size of warm-blooded animals and recent work in the Neotropics has
found mean body mass and the mass: wing length ratio of birds decreasing over the past four

decades (Jirinec et al., 2021). Smaller individuals and species might do better in warmer habitats
for several reasons. Smaller individuals tend to have a higher surface area to volume ratio, which
means they lose heat more quickly and can adapt to warmer temperatures more easily (Sheridan
and Bickford, 2011; Speakman et al., 2003). Further, smaller organisms tend to have higher
metabolic rates, which enables them to grow and reproduce faster in warmer environments
(Angilletta, 2009).

Across the world's tropical mountains, bird populations are shifting to higher elevations in response
to rising temperatures (Forero-Medina et al., 2011; Freeman et al., 2018; Neate-Clegg et al., 2018;
Freeman and Class Freeman, 2014), including in the Eastern Himalaya (Srinivasan & Wilcove,
2021; Girish and Srinivasan, 2022). At our study site, as populations shift their elevational ranges
upslope, they might encounter primary or logged forest. Our results show that logged forest bird
communities are dominated by species that have a mean elevation range midpoint that is lower than
that of primary forest species, although this effect was weak (Fig. 4). Logged forest, which is
consistently warmer than primary forest at our site (with a mean midday difference of 2.3°C,
Aggarwal et al., 2023) might be more suitable abiotically for lower-elevation, comparatively warm-
adapted, species compared with primary forest.

These indirect impacts of climate change on bird species highlight the complex interplay between
biotic and abiotic factors in shaping ecological communities. The weak shift in species composition
of understory birds in logged forest towards communities resembling those at lower elevations
suggests that while climate change will cause upslope range shifts, habitat degradation might then
select for species that are adapted to the altered abiotic conditions typical of such habitats. At our
site, bird populations at the cold-edge elevational range limit have increased survival rates over
time, while those at the warm-edge elevational range limit suffer declines (Srinivasan and Wilcove,
2021). This suggests that species are adapting to climate change by tracking favourable conditions
and shifting their ranges upslope. However, individuals of the same species are unable to maintain
demographic vital rates in higher-elevation logged forests in response to climate change in the long
term (Srinivasan and Wilcove, 2021). This could be due to a decrease in resource availability in
logged forest which can severely impact bird demographics (Grames et al., 2023). Therefore,
protecting large tracts of well-preserved primary forests across elevational gradients is essential for
enabling tropical montane species to survive the dual effects of climate change and land-use change.

This study aimed to investigate the effects of climate change and selective logging on the
insectivorous bird communities in tropical understorey in primary and selectively logged forests in
the Eastern Himalayas of India. The results indicate that smaller species, more typical of low
elevations, tended to dominate the bird communities in logged forests. This shift in body size and
range may be due to various factors, such as changes in food resources, habitat, and microclimate
resulting from logging and warming temperatures. This emphasises the importance of considering
long-term changes in bird communities when developing conservation strategies for bird species
and their habitats. In light of our findings, continuous monitoring and management of bird habitats
are crucial to prevent declines in bird diversity and abundance.

Acknowledgements:

We thank the Arunachal Pradesh Forest Department for their continuous support and for providing us with the permits to carry out this work. We are thankful to the Indian Institute of Science, Science and Engineering Research Board, GoI, and the Department of Biotechnology, GoI for funding our study. We thank Indi Glow, Nima Tsering, Joli Borah, Ramesh Kumar Pradhan, Kushal Hazarika, LN Thongchi, Millo Tasser, Nana Khrimey, Pema Mosobi, Ramana Athreya, Puspalal Sharma, Dorjee Pema, Satu Narah, Shreesh Kaulgud and Tarun Menon for all their help during fieldwork. We are grateful towards the people of Ramalingam, West Kameng for their constant support.

Ethics statement:

We thank the Arunachal Pradesh Forest Department for providing us with permission for this study (No: CWL/GEN/2018-19/Pt.IX/NG/5210-13). This study has been approved by the Institutional Animal Ethics Committee of the Indian Institute of Science (No: CAF/Ethics/883/2022).

References:

1. Rahbek, C., Borregaard, M. K., Colwell, R. K., Dalsgaard, B. O., Holt, B. G., Morueta-Holme, N., Nogues-Bravo, D., Whittaker, R. J., & Fjeldså, J. (2019). Humboldt's enigma: What causes global patterns of mountain biodiversity?. *Science*, *365*(6458), 1108-1113.

2. Bradshaw, C. J., Sodhi, N. S., & Brook, B. W. (2009). Tropical turmoil: a biodiversity tragedy in progress. *Frontiers in Ecology and the Environment*, *7*(2), 79-87.

3. Asner, G. P., Rudel, T. K., Aide, T. M., Defries, R., & Emerson, R. (2009). A contemporary assessment of change in humid tropical forests. *Conservation Biology*, *23*(6), 1386-1395.

4. Edwards, D. P., & Laurance, W. F. (2013). Biodiversity despite selective logging. *Science*, *339*(6120), 646-647.

5. Costantini, D., Edwards, D. P., & Simons, M. J. (2016). Life after logging in tropical forests of Borneo: A meta-analysis. *Biological Conservation*, *196*, 182-188.

6. Burivalova, Z., Şekercioğlu, Ç. H., & Koh, L. P. (2014). Thresholds of logging intensity to maintain tropical forest biodiversity. *Current Biology*, *24*(16), 1893-1898.

7. Dawson, T. P., Jackson, S. T., House, J. I., Prentice, I. C., & Mace, G. M. (2011). Beyond predictions: biodiversity conservation in a changing climate. *Science*, *332*(6025), 53-58.

8. Corlett, R. T. (2012). Climate change in the tropics: the end of the world as we know it?. *Biological Conservation*, *151*(1), 22-25.

9. Bellard, C., Bertelsmeier, C., Leadley, P., Thuiller, W., & Courchamp, F. (2012). Impacts of climate change on the future of biodiversity. *Ecology Letters*, *15*(4), 365-377.

10. Matthews, T. J., Cottee-Jones, H. E., & Whittaker, R. J. (2014). Habitat fragmentation and the species-area relationship: a focus on total species richness obscures the impact of habitat loss on habitat specialists. *Diversity and Distributions*, *20*(10), 1136-1146.

11. Fahrig, L. (2003). Effects of habitat fragmentation on biodiversity. *Annual review of ecology, evolution, and systematics*, *34*(1), 487-515.

12. Mantyka-pringle, C. S., Martin, T. G., & Rhodes, J. R. (2012). Interactions between climate and habitat loss effects on biodiversity: a systematic review and meta-analysis. *Global Change Biology*, *18*(4), 1239-1252.

13. Guo, F., Lenoir, J., & Bonebrake, T. C. (2018). Land-use change interacts with climate to determine elevational species redistribution. *Nature communications*, *9*(1), 1315.

14. Srinivasan, U., & Wilcove, D. S. (2021). Interactive impacts of climate change and land-use change on the demography of montane birds. *Ecology*, *102*(1), e03223.

15. Mitchard, E. T. (2018). The tropical forest carbon cycle and climate change. *Nature*, *559*(7715), 527-534.

16. Arneth, A., Shin, Y. J., Leadley, P., Rondinini, C., Bukvareva, E., Kolb, M., ... & Saito, O. (2020). Post-2020 biodiversity targets need to embrace climate change. *Proceedings of the National Academy of Sciences*, *117*(49), 30882-30891.

17. Janzen, D. H. (1967). Why mountain passes are higher in the tropics. *The American Naturalist*, *101*(919), 233-249.

18. McCain, C. M. (2009). Vertebrate range sizes indicate that mountains may be 'higher' in the tropics. *Ecology Letters*, *12*(6), 550-560.

19. Sheard, C., Neate-Clegg, M. H., Alioravainen, N., Jones, S. E., Vincent, C., MacGregor, H. E., ... & Tobias, J. A. (2020). Ecological drivers of global gradients in avian dispersal inferred from wing morphology. *Nature communications*, *11*(1), 1-9.

20. Forero-Medina, G., Terborgh, J., Socolar, S. J., & Pimm, S. L. (2011). Elevational ranges of birds on a tropical montane gradient lag behind warming temperatures. *PloS one*, *6*(12), e28535.

21. Campos-Cerqueira, M., Arendt, W. J., Wunderle Jr, J. M., & Aide, T. M. (2017). Have bird distributions shifted along an elevational gradient on a tropical mountain?. *Ecology and Evolution*, *7*(23), 9914-9924.

22. Freeman, B. G., Scholer, M. N., Ruiz-Gutierrez, V., & Fitzpatrick, J. W. (2018). Climate change causes upslope shifts and mountaintop extirpations in a tropical bird community. *Proceedings of the National Academy of Sciences*, *115*(47), 11982-11987.

23. Neate-Clegg, M. H., Jones, S. E., Burdekin, O., Jocque, M., & Şekercioğlu, Ç. H. (2018). Elevational changes in the avian community of a Mesoamerican cloud forest park. *Biotropica*, *50*(5), 805-815.

24. Freeman, B. G., Song, Y., Feeley, K. J., & Zhu, K. (2021). Montane species track rising temperatures better in the tropics than in the temperate zone. *Ecology Letters*, *24*(8), 1697-1708.

25. Putz, F. E., Zuidema, P. A., Synnott, T., Peña-Claros, M., Pinard, M. A., Sheil, D., ... & Zagt, R. (2012). Sustaining conservation values in selectively logged tropical forests: the attained and the attainable. *Conservation Letters*, *5*(4), 296-303.

26. Gibson, L., Lee, T. M., Koh, L. P., Brook, B. W., Gardner, T. A., Barlow, J., ... & Sodhi, N. S. (2011). Primary forests are irreplaceable for sustaining tropical biodiversity. *Nature*, *478*(7369), 378-381.

27.. R Core Team (2023). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

28. Borcard, D., Gillet, F., & Legendre, P. (2011). *Numerical ecology with R* (Vol. 2, p. 688). New
York: Springer.
29. Legendre, P., & Gallagher, E. D. (2001). Ecologically meaningful transformations for
ordination of species data. *Oecologia*, *129*, 271-280.
30. Oksanen, J., Simpson, G., Blanchet, F., Kindt, R., Legendre, P., Minchin, P., O'Hara, R.,
Solymos, P., Stevens, M., Szoecs, E., …, Weedon, J. (2022). _vegan: Community Ecology
Package_. R package version 2.6-4, https://CRAN.R-project.org/package=vegan.
31. Clarke, K. R., & Warwick, R. M. (2001). Change in marine communities. *An approach to*
statistical analysis and interpretation, *2*, 1-68.
32. Spierenburg, P. (2005). *Birds in Bhutan: status and distribution*. Oriental Bird Club.
33. Srinivasan, U., Elsen, P. R., Tingley, M. W., & Wilcove, D. S. (2018). Temperature and
competition interact to structure Himalayan bird communities. *Proceedings of the Royal Society B:*
Biological Sciences, *285*(1874), 20172593.
34. Devictor, V., Julliard, R., Couvet, D., & Jiguet, F. (2008). Birds are tracking climate warming,
but not fast enough. *Proceedings of the Royal Society B: Biological Sciences*, *275*(1652), 2743-
2748.
35. Cleary, D. F., Boyle, T. J., Setyawati, T., Anggraeni, C. D., Loon, E. E. V., & Menken, S. B.
(2007). Bird species and traits associated with logged and unlogged forest in Borneo. *Ecological*
Applications, *17*(4), 1184-1197.
36. Chaves, W. A., Sieving, K. E., & Fletcher Jr, R. J. (2017). Avian responses to reduced-impact
logging in the southwestern Brazilian Amazon. *Forest ecology and management*, *384*, 147-156.
37. Burivalova, Z., Lee, T. M., Giam, X., Şekercioğlu, Ç. H., Wilcove, D. S., & Koh, L. P. (2015).
Avian responses to selective logging shaped by species traits and logging practices. *Proceedings of*
the Royal Society B: Biological Sciences, *282*(1808), 20150164.
38. Velho, N., Ratnam, J., Srinivasan, U., & Sankaran, M. (2012). Shifts in community structure of
tropical trees and avian frugivores in forests recovering from past logging. *Biological Conservation*,
153, 32-40.
39. Srinivasan, U. (2013). A slippery slope: logging alters mass–abundance scaling in ecological
communities. *Journal of Applied Ecology*, *50*(4), 920-928.
40. Freeman, B. G., & Class Freeman, A. M. (2014). Rapid upslope shifts in New Guinean birds
illustrate strong distributional responses of tropical montane species to global warming.
Proceedings of the National Academy of Sciences, *111*(12), 4490-4494.
41. Bharadwaj, A., Chanda, R., & Srinivasan, U. (2022). Abiotic niche predictors of long-term
trends in body mass and survival of Eastern Himalayan birds. *bioRxiv*, 2022-08.
42. Aggarwal, K., Chanda, R., Rai, S., Rai, M., Pradhan, D. K., Munda, B., ... & Srinivasan, U.
(2023). The effect of selective logging on microclimates, arthropod abundance and the foraging
behaviour of Eastern Himalayan birds. *Forest Ecology and Management*, *541*, 121076.
43. Jirinec, V., Burner, R. C., Amaral, B. R., Bierregaard Jr, R. O., Fernández-Arellano, G.,
Hernández-Palma, A., ... & Stouffer, P. C. (2021). Morphological consequences of climate change
for resident birds in intact Amazonian rainforest. *Science Advances*, *7*(46), eabk1743.
44. Sheridan, J. A., & Bickford, D. (2011). Shrinking body size as an ecological response to climate
change. *Nature Climate Change*, *1*(8), 401-406.

45. Speakman, J. R., Thomas, D. W., Kunz, T. H., & Fenton, M. B. (2003). Physiological ecology
and energetics of bats. *Bat ecology*, 430-490.
46. Angilletta Jr, M. J. (2009). Thermal adaptation: a theoretical and empirical synthesis.Oxford
University Press.
47. Forero-Medina, G., Terborgh, J., Socolar, S. J., & Pimm, S. L. (2011). Elevational ranges of
birds on a tropical montane gradient lag behind warming temperatures. *PloS one*, *6*(12), e28535.
48. Girish, K. S., & Srinivasan, U. (2022). Community science data provide evidence for upward
elevational range shifts by Eastern Himalayan birds. *Biotropica*, *54*(6), 1457-1465.
49. Grames, E. M., Montgomery, G. A., Youngflesh, C., Tingley, M. W., & Elphick, C. S. (2023).
The effect of insect food availability on songbird reproductive success and chick body condition:
Evidence from a systematic review and meta-analysis. *Ecology Letters*.

501 **Supplementary materials:**

503 **Supp. Table S1: Final list of bird species used for analyses after excluding high-elevation**
504 **breeders that do not breed in our study area, aquatic species, carnivores, canopy dwellers,**
505 **and non-invertivores (which included frugivores, granivores and nectarivores).**

506
507

	Scientific Name	Species Code	English Name	Mass (g)	Forest Layer	Diet	Migr ant	Flock ing	Elevatio n - Summer Low	Elevation - Summer High	Elevation - Summer Mean
1	*Abroscopus schisticeps*	ABRSCH	Black-faced Warbler	5.78	Midsto rey	Insec tivor e	No	Yes	1400	2400	1900
2	*Actinodura egertoni*	ACTEGE	Rusty-fronted Barwing	32.77	Midsto rey	Insec tivor e	No	Yes	1000	2000	1500
3	*Actinodura nipalensis*	ACTNIP	Hoary-throated Barwing	46.46	Midsto rey	Insec tivor e	No	Yes	1800	3200	2500
4	*Actinodura waldeni*	ACTWAL	Streak-throated Barwing	47.5	Midsto rey	Insec tivor e	No	Yes	2400	3300	2850
5	*Aegithalos concinnus*	AEGCON	Black-throated Bushtit	6.14	Midsto rey	Insec tivor e	No	Yes	1200	2600	1900
6	*Alcippe castaneceps*	ALCCAS	Rufous-winged Fulvetta	9.7	Midsto rey	Insec tivor e	No	Yes	1800	3200	2500
7	*Alcippe cinerea*	ALCCIN	Yellow-throated Fulvetta	9.34	Unders torey	Insec tivor e	No	Yes	1400	2400	1900
8	*Anthipes monileger*	FICMON	White-gorgeted Flycatcher	11.41	Midsto rey	Insec tivor e	Yes	No	1600	2000	1800
9	*Blythipicus pyrrhotis*	BLYPYR	Bay Woodpecke r	164.4 9	Midsto rey	Insec tivor e	Yes	Yes	800	2200	1500
10	*Brachypteryx leucophris*	BRALEU	Lesser Shortwing	14.73	Terrest rial	Insec tivor e	Yes	No	1400	2800	2100
11	*Brachypteryx montana*	BRAMON	White-browed Shortwing	18.57	Terrest rial	Insec tivor e	Yes	No	2200	3400	2800

12	*Certhia discolor*	CERDIS	Brown-throated Treecreeper	10.86	Midstorey	Insectivore	Yes	Yes	160	2600	1380
13	*Cettia fortipes*	CETFOR	Brown-flanked Bush Warbler	10.23	Understorey	Insectivore	Yes	No	1400	2800	2100
14	*Cissa chinensis*	CISCHI	Common Green Magpie	127	Midstorey	Omnivore	No	No	400	2000	1200
15	*Dendrocitta frontalis*	DENFRO	Collared Treepie	80.99	Midstorey	Omnivore	No	No	600	2000	1300
16	*Ficedula hyperythra*	FICHYP	Snowy-browed Flycatcher	8.65	Understorey	Insectivore	Yes	No	1800	2800	2300
17	*Ficedula westermanni*	FICWES	Little Pied Flycatcher	7.8	Midstorey	Insectivore	Yes	No	1000	2400	1700
18	*Garrulax caerulatus*	GARCAE	Grey-sided Laughingthrush	80.61	Understorey	Omnivore	No	Yes	1600	2400	2000
19	*Harpactes erythrocephalus*	HARERY	Red-headed Trogon	100.81	Midstorey	Omnivore	Yes	Yes	400	2000	1200
20	*Hydrornis nipalensis*	PITNIP	Blue-naped Pitta	124	Terrestrial	Insectivore	Yes	No	1400	1800	1600
21	*Leiothrix argentauris*	LEIARG	Silver-eared Leiothrix	21.89	Understorey	Omnivore	No	No	600	1800	1200
22	*Leiothrix lutea*	LEILUT	Red-billed Leiothrix	22.36	Understorey	Omnivore	No	No	1800	3200	2500
23	*Liocichla bugunorum*	LIOBUG	Bugun Liocichla	39	Midstorey	Omnivore	No	Yes	2000	2600	2300
24	*Lioparus chrysotis*	ALCCHR	Golden-breasted Fulvetta	8.12	Understorey	Insectivore	No	Yes	2000	2800	2400
25	*Machlolophus spilonotus*	PARSPI	Yellow-cheeked Tit	18.39	Midstorey	Insectivore	Yes	Yes	1200	2450	1825
26	*Muscicapa ferruginea*	MUSFER	Ferruginous Flycatcher	10.21	Midstorey	Insectivore	Yes	No	1400	2800	2100

27	*Muscicapella hodgsoni*	MUSHOD	Pygmy Flycatcher	5.76	Midstorey	Insectivore	Yes	No	1400	2600	2000
28	*Myiomela leucura*	MYILEU	White-tailed Robin	25.72	Terrestrial	Insectivore	Yes	No	1200	2200	1700
29	*Niltava grandis*	NILGRA	Large Niltava	36.03	Midstorey	Insectivore	Yes	No	1600	2200	1900
30	*Niltava macgrigoriae*	NILMAC	Small Niltava	11.31	Understorey	Insectivore	Yes	No	800	2000	1400
31	*Niltava sundara*	NILSUN	Rufous-bellied Niltava	20.2	Understorey	Insectivore	Yes	No	1600	3000	2300
32	*Parus monticolus*	PARMON	Green-backed Tit	18.65	Midstorey	Insectivore	Yes	Yes	1200	3000	2100
33	*Phyllergates cuculatus*	ORTCUC	Mountain Tailorbird	6.31	Understorey	Insectivore	Yes	Yes	1400	2200	1800
34	*Phylloscopus chloronotus*	PHYCHL	Lemon-rumped Warbler	5.01	Midstorey	Insectivore	Yes	Yes	2400	3600	3000
35	*Phylloscopus reguloides*	PHYREG	Blyth's Leaf Warbler	7.05	Midstorey	Insectivore	Yes	Yes	1600	3200	2400
36	*Pnoepyga pusilla*	PNOPUS	Pygmy Wren-babbler	12.25	Understorey	Insectivore	No	Yes	1000	2600	1800
37	*Pomatorhinus ferruginosus*	POMFER	Coral-billed Scimitar Babbler	40.65	Understorey	Insectivore	No	Yes	1000	2000	1500
38	*Pomatorhinus ruficollis*	POMRUF	Streak-breasted Scimitar Babbler	29.76	Midstorey	Insectivore	Yes	Yes	1600	2600	2100
39	*Pomatorhinus superciliaris*	XIPSUP	Slender-billed Scimitar Babbler	30.12	Understorey	Omnivore	Yes	No	1600	3400	2500
40	*Psarisomus dalhousiae*	PSADAL	Long-tailed Broadbill	55.06	Midstorey	Insectivore	Yes	Yes	400	2000	1200

41	*Psittiparus ruficeps*	PARRUF	White-breasted Parrotbill	31.54	Midstorey	Insectivore	No	Yes	1000	2000	1500
42	*Rhipidura albicollis*	RHIALB	White-throated Fantail	12.02	Midstorey	Insectivore	Yes	Yes	1200	2400	1800
43	*Rimator malacoptilus*	RIMMAL	Long-billed Wren-Babbler	19.69	Understorey	Insectivore	Yes	Yes	900	2700	1800
44	*Sasia ochracea*	SASOCH	White-browed Piculet	10.03	Midstorey	Insectivore	Yes	Yes	400	2000	1200
45	*Seicercus affinis*	SEIAFF	White-spectacled Warbler	7.58	Midstorey	Insectivore	Yes	Yes	1000	2800	1900
46	*Seicercus poliogenys*	SEIPOL	Grey-cheeked Warbler	6.57	Midstorey	Insectivore	Yes	Yes	1400	2400	1900
47	*Spelaeornis caudatus*	SPECAU	Rufous-throated Wren-Babbler	10.53	Terrestrial	Insectivore	No	Yes	1600	2400	2000
48	*Spelaeornis formosus*	ELAFOR	Spotted Wren-Babbler	10.79	Terrestrial	Insectivore	No	No	800	2200	1500
49	*Sphenocichla humei*	SPHHUM	Sikkim Wedge-billed Babbler	34.1	Understorey	Insectivore	No	Yes	1400	2000	1700
50	*Stachyridopsis chrysaea*	STACHR	Golden Babbler	7.85	Understorey	Insectivore	No	Yes	800	2000	1400
51	*Stachyridopsis ruficeps*	STARUF	Rufous-capped Babbler	9.85	Understorey	Insectivore	Yes	Yes	1200	1800	1500
52	*Stachyris nigriceps*	STANIG	Grey-throated Babbler	14.17	Understorey	Insectivore	No	Yes	400	1800	1100
53	*Suthora nipalensis*	PARNIP	Black-throated Parrotbill	6.31	Understorey	Insectivore	No	Yes	1600	2800	2200
54	*Tesia castaneocoronata*	TESCAS	Chestnut-headed Tesia	8.31	Terrestrial	Insectivore	Yes	No	2200	3400	2800

55	*Tesia cyaniventer*	TESCYA	Grey-bellied Tesia	9.89	Terrestrial	Insectivore	Yes	No	1400	2600	2000
56	*Tesia olivea*	TESOLI	Slaty-bellied Tesia	8.79	Terrestrial	Insectivore	Yes	Yes	1400	2000	1700
57	*Tickellia hodgsoni*	TICHOD	Broad-billed Warbler	5.88	Understorey	Insectivore	Yes	Yes	1600	2600	2100
58	*Trochalopteron erythrocephalum*	GARERY	Chestnut-crowned Laughingthrush	71.63	Understorey	Omnivore	No	Yes	1600	3400	2500
59	*Trochalopteron squamatum*	GARSQU	Blue-winged Laughingthrush	81.48	Understorey	Omnivore	No	Yes	1000	2000	1500
60	*Trochalopteron subunicolor*	GARSUB	Scaly Laughingthrush	66	Understorey	Omnivore	No	Yes	2000	3400	2700
61	*Yuhina flavicollis*	YUHFLA	Whiskered Yuhina	14.97	Midstorey	Omnivore	Yes	No	1600	2800	2200

511 **Supp. Table S2: Raw counts of species in primary and logged forest. The first ten columns**

512 **are from primary, followed by logged.**

	Species Code	2011.P	2012.P	2013.P	2014.P	2015.P	2016.P	2017.P	2018.P	2019.P	2021.P	2011.L	2012.L	2013.L	2014.L	2015.L	2016.L	2017.L	2018.L	2019.L	2021.L
1	ABRSCH	3	5	4	10	3	2	5	0	1	5	3	2	0	2	1	6	5	3	3	3
2	ACTEGE	4	9	13	4	8	4	6	2	4	6	1	7	5	6	6	13	6	14	5	10
3	ACTNIP	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0
4	ACTWAL	1	0	2	0	0	1	0	0	0	0	0	2	0	0	1	1	0	0	0	0
5	AEGCON	0	1	0	0	1	0	3	0	0	0	0	0	0	0	0	0	0	0	0	2
6	ALCCAS	16	32	14	17	16	20	17	11	7	29	41	23	10	12	19	22	21	27	17	48

7	ALCCHR	11	10	10	4	8	16	13	6	6	29	22	23	11	14	21	27	24	16	16	48
8	ALCCIN	16	50	28	10	24	32	16	16	6	42	50	30	26	7	25	18	26	23	7	70
9	BLYPYR	1	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0
10	BRALEU	4	1	0	2	4	1	5	2	4	3	3	2	7	1	2	5	7	3	5	5
11	BRAMON	0	0	0	0	0	0	0	1	0	2	0	1	0	0	0	0	0	0	0	1
12	CERDIS	0	1	2	0	2	2	0	1	0	1	0	0	0	0	0	0	0	0	0	0
13	CETFOR	0	1	0	1	1	4	0	1	1	3	1	0	1	2	2	2	1	4	2	2
14	CISCHI	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0
15	DENFRO	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1
16	ELAFOR	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1
17	FICHYP	18	13	8	13	11	14	10	7	6	18	19	13	4	2	15	10	10	16	4	12
18	FICMON	2	3	5	3	5	6	6	1	2	7	1	0	0	2	6	2	4	3	3	9
19	FICWES	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0
20	GARCAE	3	2	1	1	3	3	5	3	1	11	4	6	1	0	2	4	3	4	1	2
21	GARERY	3	7	6	2	2	5	2	2	6	8	4	4	3	4	2	6	2	8	4	11
22	GARSQU	0	0	1	0	0	0	0	0	0	1	0	0	0	0	4	0	2	3	0	0
23	GARSUB	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0
24	HARERY	0	2	0	0	1	0	0	0	2	1	1	0	0	0	0	0	0	0	0	0
25	LEIARG	0	0	0	0	0	0	0	0	0	0	1	0	0	0	1	2	0	0	1	0
26	LEILUT	0	0	0	0	0	0	0	2	0	0	0	0	1	0	0	2	0	0	0	0
27	LIOBUG	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	2	0	0	0	0
28	MUSFER	0	0	0	0	0	0	0	0	0	0	2	2	0	0	1	0	0	0	0	0

29	MUSHOD	1	0	0	3	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30	MYILEU	1	3	5	1	6	6	5	4	5	8	3	4	2	1	5	5	9	9	9	6
31	NILGRA	7	10	8	3	7	4	6	7	5	13	5	8	3	1	3	5	4	7	2	10
32	NILMAC	0	1	0	0	0	0	0	0	0	1	1	5	0	0	0	0	1	0	0	6
33	NILSUN	16	1	10	4	11	12	16	4	2	17	11	0	4	1	6	6	5	4	1	2
34	ORTCUC	0	2	0	0	1	0	4	0	0	0	0	4	1	3	3	2	4	3	0	0
35	PARNIP	17	33	14	8	7	17	7	6	11	26	33	7	8	1	16	18	32	19	13	26
36	PARRUF	0	0	9	1	2	5	2	0	2	0	5	1	0	0	1	0	1	1	0	4
37	PARSPI	2	0	5	0	1	0	1	0	2	1	0	0	1	0	0	1	0	0	0	3
38	PHYCHL	0	1	0	0	0	0	0	0	0	1	3	0	0	0	0	0	0	1	0	1
39	PHYREG	7	7	6	4	3	2	9	2	1	14	6	9	1	3	0	9	10	5	2	9
40	PITNIP	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
41	PN0PUS	0	0	0	0	0	0	0	0	0	2	0	0	0	0	0	0	0	0	0	0
42	PNOPUS	5	2	2	0	1	0	1	0	1	2	1	1	0	0	0	1	1	0	0	1
43	POMFER	0	1	2	1	0	3	1	4	0	2	1	2	2	2	3	0	1	0	0	1
44	POMRUF	3	4	0	2	2	1	3	2	1	2	6	3	0	1	4	3	8	5	2	7
45	PSADAL	0	0	0	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	0	0
46	RHIALB	2	2	2	1	0	4	3	2	2	9	8	3	3	1	0	1	5	5	4	3
47	RIMMAL	2	0	0	0	2	1	0	0	0	0	0	0	1	0	1	0	0	0	0	0
48	SASOCH	0	0	0	0	0	2	1	2	1	2	0	1	1	0	1	3	2	2	2	6
49	SEIAFF	9	8	8	2	6	6	7	5	6	15	12	9	3	2	6	6	4	4	2	13
50	SEIPOL	10	13	4	7	4	11	11	4	8	24	10	9	4	7	5	14	8	5	3	22

#	Species code																				
51	SPECAU	3	2	1	1	7	5	4	2	1	7	5	1	0	0	2	0	3	3	2	3
52	SPHHUM	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0
53	STACHR	8	15	10	4	6	14	9	8	3	19	9	8	7	7	16	15	14	11	10	19
54	STANIG	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0
55	STARUF	12	16	12	10	8	16	16	7	7	19	33	31	17	16	38	31	26	29	13	37
56	TESCAS	1	0	2	3	0	0	0	0	0	3	5	6	0	1	0	0	0	0	0	8
57	TESCYA	4	4	2	1	3	2	3	3	0	1	11	3	2	2	4	2	1	1	0	6
58	TESOLI	2	1	2	0	3	3	2	3	1	2	3	2	0	2	2	3	0	1	1	5
59	TICHOD	0	1	0	2	0	0	0	0	0	3	4	3	1	1	2	2	1	1	0	6
60	XIPSUP	0	0	2	0	2	1	0	2	2	5	3	5	0	0	1	0	1	1	2	5
61	YUHFLA	12	0	0	1	0	1	2	2	2	9	6	3	8	2	2	3	9	1	1	14

515
516
517

Supp. Table S3: Similarity percentage (SIMPER) routines of species that contribute to 50%
cumulative dissimilarity between the two habitats.
Note: *Schoeniparus cinereus* **and** *Tesia castaneocoronata* **have not been included in Figure 2**
(NMDS)

Species	Species code	average	sd	ratio	ava	avb	cumsum	*p* value	significance
Niltava sundara	NILSUN	0.009681	0.006145	1.5754	0.2031	0.1182	0.038	0.008	**
Stachyris ruficeps	STARUF	0.009032	0.004001	2.2573	0.249	0.3419	0.074	0.001	***
Yuhina flavicollis	YUHFLA	0.007835	0.005886	1.3311	0.0912	0.1336	0.105	0.227	
Lioparus chrysotis	ALCCHR	0.007678	0.004217	1.8208	0.2292	0.3063	0.136	0.001	***
Phyllergates cucullatus	ORTCUC	0.007159	0.005321	1.3454	0.0304	0.0818	0.164	0.048	*
Psittiparus ruficeps	PARRUF	0.006938	0.005376	1.2905	0.0792	0.0472	0.192	0.239	
Schoeniparus cinereus	ALCCIN	0.00692	0.004932	1.4032	0.3324	0.3265	0.219	0.879	
Tickellia hodgsoni	TICHOD	0.006553	0.003706	1.7682	0.0277	0.0825	0.245	0.005	**

Suthora nipalensis	PARNIP	0.00652	0.005306	1.2288	0.2616	0.2528	0.271	0.944	
Spelaeornis caudatus	SPECAU	0.006052	0.004935	1.2263	0.1216	0.07	0.295	0.033	*
Abroscopus schisticeps	ABRSCH	0.005949	0.005781	1.029	0.1275	0.1021	0.318	0.685	
Tesia castaneocoronata	TESCAS	0.005933	0.005626	1.0546	0.0413	0.0508	0.342	0.811	
Pomatorhinus superciliaris	XIPSUP	0.005774	0.004417	1.3074	0.0651	0.0655	0.365	0.842	
Pomatorhinus ferruginosus	POMFER	0.00576	0.004343	1.3263	0.0688	0.0625	0.388	0.922	
Brachypteryx leucophris	BRALEU	0.005728	0.004386	1.3059	0.1109	0.1312	0.41	0.433	
Anthipes monileger	FICMON	0.005618	0.004838	1.1611	0.1383	0.0966	0.433	0.121	
Machlolophus spilonotus	PARSPI	0.005486	0.004884	1.1234	0.0589	0.023	0.455	0.183	
Sasia ochracea	SASOCH	0.005414	0.004066	1.3315	0.0459	0.0739	0.476	0.234	
Ficedula hyperythra	FICHYP	0.005386	0.004097	1.3149	0.245	0.2015	0.497	0.038	*
Niltava grandis	NILGRA	0.005301	0.003144	1.6863	0.1875	0.1365	0.518	0.001	***

524   Significance codes: '***' = p < 0.001, '**' = p < 0.01, '*' = p < 0.05

525   average = average contribution of the species to between-group dissimilarity

526   ava, avb = average abundances of each habitat (primary and logged)

542   **Figure S1: Highest breeding elevation range of the species that contributed to 50%**

543   **dissimilarity in each habitat. The box shows the middle 50% of the data, with the horizontal**

544   **line inside the box indicating the median value. The whiskers of the boxplot extend from the**
545   **box to show the minimum and maximum values of the data. Each dot represents individual**
546   **data points and is plotted with a jitter to avoid overlap.**
547

562   **Figure S2: Lowest breeding elevation range of the species that contributed to 50%**
563   **dissimilarity in each habitat. The box shows the middle 50% of the data, with the horizontal**
564   **line inside the box indicating the median value. The whiskers of the boxplot extend from the**

565 **box to show the minimum and maximum values of the data. Each dot represents individual**
566 **data points and is plotted with a jitter to avoid overlap.**
567

582 **Figure S3: Abundance weighted mass index over time of the species that contributed to 50%**
583 **dissimilarity in each habitat.**
584

Figure S4: Abundance weighted mean breeding elevation index over time of the species that contributed to 50% dissimilarity in each habitat.