Atoh1 is required for the formation of lateral line electroreceptors and hair cells, whereas Foxg1 represses an electrosensory fate

Martin Minařík[1], Alexander S. Campbell[1], Roman Franěk[2#], Michaela Vazačová[2], Miloš Havelka[2], David Gela[2], Martin Pšenička[2] and Clare V. H. Baker[1*]

[1]Department of Physiology, Development & Neuroscience, University of Cambridge, Cambridge, UK
[2]Faculty of Fisheries and Protection of Waters, Research Institute of Fish Culture and Hydrobiology, University of South Bohemia in České Budějovice, Vodňany, Czech Republic

*For correspondence: cvhb1@cam.ac.uk

#Current address: Silberman Institute of Life Sciences, The Hebrew University of Jerusalem, Jerusalem, Israel

Abstract

In electroreceptive jawed fishes and amphibians, individual lateral line placodes form lines of neuromasts on the head containing mechanosensory hair cells, flanked by fields of ampullary organs containing electroreceptors - modified hair cells that respond to weak electric fields. Extensively shared gene expression suggests that conserved molecular mechanisms are involved in their development, but a few transcription factor genes are restricted either to the developing electrosensory or mechanosensory lateral line. Here, we used CRISPR/Cas9-mediated mutagenesis in F0-injected sterlet embryos (*Acipenser ruthenus*, a sturgeon) to test the function of three such genes. We found that the 'hair cell' transcription factor gene *Atoh1* is required for both hair cell and electroreceptor differentiation in sterlet, and for *Pou4f3* and *Gfi1* expression in both neuromasts and ampullary organs. These data support the conservation of developmental mechanisms between hair cells and electroreceptors. Targeting ampullary organ-restricted *Neurod4* did not yield any phenotype, potentially owing to redundancy with other *Neurod* genes that we found to be expressed in sterlet ampullary organs. After targeting mechanosensory-restricted *Foxg1*, ampullary organs formed within neuromast lines, suggesting that Foxg1 normally represses their development. We speculate that electrosensory organs may be the 'default' fate of lateral line primordia in electroreceptive vertebrates.

Introduction

Mechanosensory hair cells in different compartments of the inner ear transduce fluid movements for hearing and balance (Moser et al., 2020; Caprara and Peng, 2022; Mukhopadhyay and Pangrsic, 2022). In fishes and aquatic-stage amphibians, hair cells are also found in lateral line neuromasts in the skin, which are stimulated by local water movement (Mogdans, 2019; Webb, 2021). Supporting cells in neuromasts differentiate into hair cells for homeostasis and repair (Kniss et al., 2016; Lush et al., 2019) and neuromasts are easily accessible in lines over the head and trunk, making the zebrafish lateral line an excellent model for hair cell development and regeneration (Kniss et al., 2016; Nicolson, 2017; Pickett and Raible, 2019).

The lateral line system of zebrafish (a cyprinid teleost) is purely mechanosensory, as is the lateral line system of the other main anamniote lab model, the frog *Xenopus*. However, in many other vertebrates, the lateral line system also has an electrosensory division (Bullock et al., 1983; Baker et al., 2013; Crampton, 2019). In electroreceptive non-teleost jawed vertebrates, some or all of the neuromast lines on the head are flanked by fields of ampullary organs containing electroreceptors, which respond to weak cathodal stimuli such as the electric fields surrounding other animals in water (Bodznick and Montgomery, 2005; Crampton, 2019; Leitch and Julius, 2019; Chagnaud et al., 2021). Electroreception is mediated by voltage-gated calcium channels in the apical membrane (Bodznick and Montgomery, 2005; Leitch and Julius, 2019). The voltage sensor was recently identified in cartilaginous fishes as the L-type voltage-gated calcium channel $Ca_v1.3$ (Bellono et al., 2017; Bellono et al., 2018), whose pore-forming alpha subunit is encoded by *Cacna1d*. $Ca_v1.3$ is also required for synaptic transmission at hair-cell ribbon synapses (Moser et al., 2020; Mukhopadhyay and Pangrsic, 2022).

Electroreceptors, like hair cells, have an apical primary cilium and basolateral ribbon synapses with lateral line afferents (Jørgensen, 2005; Baker, 2019). However, in contrast to the highly ordered, stepped array of apical microvilli ('stereocilia') that forms the 'hair bundle' critical for hair-cell mechanotransduction (Caprara and Peng, 2022), electroreceptors in many species (e.g., cartilaginous fishes; ray-finned paddlefishes and sturgeons) lack apical microvilli altogether (Jørgensen, 2005; Baker, 2019). Electroreceptors in other species have a few apical microvilli, while the electroreceptors of the amphibian axolotl have around 200 microvilli surrounding an eccentrically positioned primary cilium (Jørgensen, 2005; Baker, 2019). Indeed, axolotl electroreceptors were described as "remarkably similar to immature hair cells" (Northcutt et al., 1994). Thus, despite their shared function, the apical surface of electroreceptors (where voltage-sensing occurs; Bodznick and Montgomery, 2005; Leitch and Julius, 2019) varies considerably across different vertebrate groups.

Fate-mapping experiments have shown that neuromasts, ampullary organs (where present) and their afferent neurons all develop from a series of pre-otic and post-otic lateral line placodes on the embryonic head (Northcutt, 1997; Piotrowski and Baker, 2014; Baker, 2019). In electroreceptive jawed vertebrates, lateral line placodes elongate to form sensory ridges that eventually fragment: neuromasts differentiate first, in a line along the centre of each ridge, and ampullary organs (if present) form later, in fields on the flanks of the ridge (Northcutt, 1997; Piotrowski and Baker, 2014; Baker, 2019). The lateral line primordia of electroreceptive vertebrates therefore provide a fascinating model for studying the formation of different sensory cell types and organs. What molecular mechanisms control the formation within the same primordium of mechanosensory neuromasts containing hair cells, versus electrosensory ampullary organs containing electroreceptors?

To gain molecular insight into electroreceptor development, we originally took a candidate-gene approach, based on genes known to be important for neuromast and/or hair cell development. This enabled us to identify a variety of genes expressed in developing ampullary organs as well as neuromasts, in embryos from the three major jawed-vertebrate groups, i.e., cartilaginous fishes (lesser-spotted catshark, *Scyliorhinus canicula*, and little skate, *Leucoraja erinacea*; O'Neill et al., 2007; Gillis et al., 2012); lobe-finned bony fishes/tetrapods (a urodele amphibian, the axolotl, *Ambystoma mexicanum*; Modrell and Baker, 2012); and ray-finned bony fishes (a chondrostean, the Mississippi paddlefish, *Polyodon spathula*; Modrell et al., 2011a; Modrell et al., 2011b; Modrell et al., 2017b). We also took an unbiased discovery approach using differential bulk RNA-seq in late-larval paddlefish, which yielded a dataset of almost 500 genes that were putatively enriched in lateral line organs (Modrell et al., 2017a). Validation by *in situ* hybridization of a subset of candidates from this dataset suggested that conserved molecular mechanisms were involved in hair cell and electroreceptor development, and that hair cells and electroreceptors were closely related physiologically (Modrell et al., 2017a). For example, developing ampullary organs express the key 'hair cell' transcription factor genes *Atoh1* and *Pou4f3* (see Roccio et al., 2020; Iyer and Groves, 2021), and genes essential for the function of hair cell ribbon synapses, including the voltage-gated calcium channel gene *Cacna1d*, encoding $Ca_v1.3$ (Modrell et al., 2017a). We also identified a handful of genes expressed in developing ampullary organs but not neuromasts, including two electroreceptor-specific voltage-gated potassium channel subunit genes (*Kcna5* and *Kcnab3*) and a single transcription factor gene, *Neurod4* (Modrell et al., 2017a).

Up to that point, we had reported the shared expression of fifteen transcription factor genes in both ampullary organs and neuromasts, but only one transcription factor gene with restricted expression, namely, electrosensory-restricted *Neurod4* (Modrell et al., 2011a; Modrell et al., 2011b; Modrell et al., 2017b). More recently (preprint: Minařík et al., 2023), we

used the late-larval paddlefish lateral line organ-enriched dataset (Modrell et al., 2017b), as
well as a candidate gene approach, to identify 23 more transcription factor genes expressed
within developing lateral line organs in paddlefish and/or in a related, more experimentally
tractable chondrostean, the sterlet (*Acipenser ruthenus*, a small sturgeon; e.g., Chen et al.,
2018; Baloch et al., 2019; Stundl et al., 2022). Twelve of these transcription factor genes -
including *Gfi1* - were expressed in both ampullary organs and neuromasts (preprint: Minařík
et al., 2023). Thus, developing ampullary organs, as well as neuromasts, express the three
'hair cell' transcription factor genes - *Atoh1*, *Pou4f3* and *Gfi1* - whose co-expression is
sufficient to drive postnatal mouse cochlear supporting cells to adopt a 'hair cell-like' fate, albeit
not to form fully mature hair cells (Roccio et al., 2020; Iyer and Groves, 2021; Chen et al.,
2021; Iyer et al., 2022). We also identified six novel ampullary organ-restricted transcription
factor genes and the first-reported mechanosensory-restricted transcription factor genes
(preprint: Minařík et al., 2023). One of the five mechanosensory-restricted transcription factor
genes was *Foxg1*, which was expressed and maintained in the central region of sensory ridges
where lines of neuromasts form, although excluded from hair cells (preprint: Minařík et al.,
2023).

Here, we used CRISPR/Cas9-mediated mutagenesis in F0-injected sterlet embryos to
investigate the function in lateral line organ development of *Atoh1*, electrosensory-restricted
Neurod4 and mechanosensory-restricted *Foxg1* (for reference, Supplementary Figure S1
shows the normal expression patterns of these genes). We report that *Atoh1* is required for
the formation of electroreceptors, as well as hair cells. We did not see any phenotype after
targeting ampullary organ-restricted *Neurod4*, potentially owing to redundancy with other
Neurod family members that we found to be expressed in sterlet ampullary organs (and
neuromasts). Targeting mechanosensory-restricted *Foxg1* resulted in a striking phenotype:
the ectopic formation of ampullary organs within neuromast lines, and in some cases the
fusion of ampullary organ fields that normally develop either side of a line of neuromasts. This
suggests the unexpected but intriguing hypothesis that ampullary organs may be the 'default'
fate for lateral line sensory ridges, and that this is repressed by Foxg1, allowing neuromasts
to form instead.
139
140 **<u>Results</u>**
141
CRISPR/Cas9-mediated mutagenesis in F0-injected sterlet embryos
To test gene function during lateral line organ development, we optimised CRISPR/Cas9-
mediated mutagenesis in F0-injected sterlet embryos, building on established protocols for
axolotl (*Ambystoma mexicanum*; Flowers et al., 2014; Fei et al., 2018), newt (*Pleurodeles*
waltl; Elewa et al., 2017), and sea lamprey (*Petromyzon marinus*; Square et al., 2015; York et

al., 2019; Square et al., 2020), whose eggs are all large (1-2 mm in diameter) and easy to

microinject at the 1-2 cell stage. Ovulated sterlet eggs are very large: roughly 2.5 mm in

diameter (Lenhardt et al., 2005). (Since this project started, CRISPR/Cas9-mediated

mutagenesis in F0-injected sterlet embryos has been reported, including by two of us, R.F.

and M.P.; Chen et al., 2018; Baloch et al., 2019; Stundl et al., 2022.)

We targeted the melanin-producing enzyme *tyrosinase* (*Tyr*) as a positive control,

using eight different single-guide (sg) RNAs. Table 1 shows the sgRNA target sequences

(including two that were designed and recently published by Stundl et al., 2022) and the

various combinations in which they were injected. Supplementary Figure S2A shows the

position of each sgRNA relative to the exon structure of the *tyrosinase* gene. Our sgRNAs

were designed prior to publication of the chromosome-level sterlet genome (Du et al., 2020).

Searching the genome for *Tyr* showed that both *Tyr* ohnologs (i.e., gene paralogs originating

from the whole-genome duplication in the sterlet lineage; Du et al., 2020) have been retained,

on chromosomes 8 and 9, with 99.21% nucleotide identity in the coding sequence (98.81%

amino acid identity). However, the copy on chromosome 8 has a 7 base-pair (bp) deletion

(c.344_350delGAAGAGA) that would result in a frameshift early in the sequence, upstream of

the tyrosinase domain. Our sgRNAs target the full-length *Tyr* gene on chromosome 9. The *Tyr*

sgRNA 1 (Table 1; Supplementary Figure S2A) has a one-base mismatch to the *Tyr* gene on

chromosome 8, in position 19 of the target sequence (just 2 bp upstream of the PAM), thus

lying within the expected seed sequence and likely to prevent successful edits (Wu et al.,

2014). Nevertheless, the frameshift-causing deletion in this ohnolog suggests it does not

encode functional protein.

Three different combinations of six of the eight *Tyr* sgRNAs, when injected at the 1-2-

cell stage (2-4 sgRNAs pre-complexed with Cas9), each generated at least four embryos

(hereafter 'crispants') with pigmentation phenotypes at stage 45 (Dettlaff et al., 1993), the

onset of the transition to independent feeding. The other two *Tyr* sgRNAs failed to generate

any phenotypes (Supplementary Table 1). Excluding the *Tyr* sgRNAs that failed, at least some

degree of pigment loss was seen in 63/111 *Tyr* crispants (56.8%) across nine independent

batches. Examples of *Tyr* crispants with pigmentation phenotypes, plus a control, are shown

in Supplementary Figure 2B-D. The most efficient results were obtained by injecting 1-cell

embryos with a preassembled mix of Cas9 protein plus two chemically modified sgRNAs

(purchased from Synthego) against the target gene, and subsequently maintaining the

embryos at room temperature for around 6 hours. The time from fertilization to completion of

the first cleavage is around 2-3 hours at room temperature, giving plenty of time for the

Cas9/sgRNA complex to act before returning the embryos to colder temperatures for

subsequent development. (Sterlet are cold-water fish and the optimum temperature for

maintaining embryos for normal development is 16°C.)

Following embryo injection at the 1-2-cell stage with pre-complexed sgRNAs/Cas9 and

fixation at stage 45, genomic DNA was extracted from the trunk/tail prior to analysis of the

heads by *in situ* hybridization (ISH). The sgRNA-targeted region from trunk/tail genomic DNA

was amplified by PCR for direct Sanger sequencing and *in silico* analysis using Synthego's

online 'Inference of CRISPR Edits' (ICE) tool (Conant et al., 2022) (also see e.g., Uribe-

Salazar et al., 2022) to analyse the identity and frequency of edits of the target gene.

Supplementary Figure S2E-I show examples of successful disruption of *Tyr*.

We note that our genotyping results have shown that most crispants analysed, across

all genes targeted, have some degree of targeted mutagenesis in the trunk/tail, with a range

of deletion sizes. Although phenotypes from the initial spawning seasons were almost always

highly mosaic, suggesting mutations occurred later in development, following optimization a

proportion of embryos showed complete unilateral and occasionally bilateral phenotypes.

Such phenotypes suggest that mutation occurred in one cell at the 2-cell stage (unilateral

phenotype) or even as early as the 1-cell stage (bilateral phenotype). Some degree of

mosaicism can be useful, however, as the unaffected tissue provides an internal control for

the normal expression of the gene being examined by ISH.

201   Targeting *Atoh1* resulted in the loss of hair cells and electroreceptors

We targeted *Atoh1* for CRISPR/Cas9-mediated mutagenesis by injecting sterlet embryos at

the 1-2 cell stage with Cas9 protein pre-complexed with two sgRNAs targeting *Atoh1* (Table

1; Supplementary Figure S2A). Our sgRNAs were designed prior to publication of the

chromosome-level sterlet genome (Du et al., 2020). Searching the genome for *Atoh1* showed

that both *Atoh1* ohnologs have been retained, on chromosomes 1 and 2, with 92.41%

nucleotide identity (and 85.48% amino acid identity) in the coding sequence. The copy on

chromosome 2 encodes a shorter version of the protein with a four amino acid deletion near

the N-terminus (E14_G17del). Our sgRNAs target the full-length *Atoh1* gene on chromosome

1. One sgRNA has a one-base mismatch to the shorter *Atoh1* gene on chromosome 2, in

position 3 of the target sequence (PAM-distal), which is unlikely to prevent successful targeting

(Wu et al., 2014). Thus, we expect our sgRNAs to target both *Atoh1* ohnologs.

The *Atoh1* crispants, together with *Tyr*-targeted siblings as controls, were raised to

stage 45 (the onset of independent feeding, around 14 days post-fertilization). ISH for the hair

cell and electroreceptor marker *Cacna1d* (Modrell et al., 2017a) (also see preprint: Minařík et

al., 2023), which was recently shown to be a direct Atoh1 target gene in mouse cochlear hair

cells (Jen et al., 2022), revealed no phenotype in *Tyr* control crispants (Figure 1A-D; n=0/17

across four independent batches; Supplementary Table 1). However, *Cacna1d* expression

was absent mosaically in neuromast lines and ampullary organ fields in *Atoh1* crispants

(Figure 1E-J; n=13/22, i.e., 59%, across six independent batches; Supplementary Table 1).

Figure 1. *Atoh1* is required for the differentiation of lateral line hair cells and electroreceptors.
Sterlet crispants at stage 45 after *in situ* hybridization (ISH) for the hair cell and electroreceptor marker
Cacna1d, or the electroreceptor-specific marker *Kcnab3*. Black arrowheads indicate examples of
neuromasts; white arrowheads indicate examples of ampullary organs. (**A-D**) In a control *Tyr* crispant,
Cacna1d expression shows the normal distribution of hair cells in lines of neuromasts, and
electroreceptors in fields of ampullary organs flanking the neuromast lines (lateral view: A,B; ventral
view: C,D). (**E-J**) In an *Atoh1* crispant, *Cacna1d* expression is absent on the left side of the head (E,F),
except for a few isolated organs in the otic region and on the operculum. Post-ISH Sox2 immunostaining
(E[1],F[1]) shows that neuromast supporting cells are still present. A ventral view (G,H) and a lateral view
of the right side of the head (I,J; image flipped horizontally for ease of comparison) reveal a unilateral
phenotype, with *Cacna1d*-expressing hair cells and electroreceptors mostly absent from the left side
only of the ventral rostrum (asterisk in G,H) and present on the right side of the head (I,J). (**K,L**) Lateral
view of a control *Tyr* crispant after ISH for *Kcnab3*, showing the position of electroreceptors in ampullary
organs. (**M,N**) Ventral view of another *Tyr* crispant showing *Kcnab3* expression in ampullary organs.
(**O-P[1]**) Lateral view of an *Atoh1* crispant in which *Kcnab3* expression is absent from ampullary organs.

Post-ISH Sox2 immunostaining (O[1],P[1]) shows that supporting cells are still present in neuromasts (strong staining) and can also be detected in ampullary organs (much weaker staining). (**Q-T**) A different *Atoh1* crispant after ISH for *Kcnab3*, shown in ventral view (Q,R: compare with M,N) and lateral view (S,T). *Kcnab3* expression reveals a unilateral phenotype: *Kcnab3*-expressing electroreceptors are mostly absent from the right side (asterisk) but present on the left side. Abbreviations: b, barbel; di, dorsal infraorbital ampullary organ field; e, eye; m, mouth; n, naris; S, stage; vi, ventral infraorbital ampullary organ field. Scale bars: 200 μm.

This suggested that disruption of the *Atoh1* gene in sterlet resulted in the failure of hair cell differentiation (as expected from zebrafish; Millimaki et al., 2007) and also electroreceptors. Post-ISH immunostaining for the supporting-cell marker Sox2 (Hernández et al., 2007; Modrell et al., 2017a) (also see preprint: Minařík et al., 2023) confirmed that neuromasts had formed (Figure 1E[1],F[1]), so the phenotype was specific to receptor cells. Ampullary organs were not reliably detected because Sox2 immunostaining labels neuromasts much more strongly than ampullary organs in sterlet (Supplementary Figure S1A,B; also preprint: Minařík et al., 2023), as it does in paddlefish (Modrell et al., 2017a). ISH for electroreceptor-specific *Kcnab3* (Modrell et al., 2017a) (also see preprint: Minařík et al., 2023) similarly showed no effect in *Tyr* control crispants (Figure 1K-N; n=0/27 across six independent batches; Supplementary Table 1), but the mosaic absence of *Kcnab3* expression in ampullary organ fields in *Atoh1* crispants (Figure 1O-T; n=8/13, i.e., 62%, across three independent batches; Supplementary Table 1). Post-*Kcnab3* ISH immunostaining for Sox2 confirmed that ampullary organs were still present, as well as neuromasts (Figure 1O[1],P[1]).

In mouse cochlear hair cells, the 'hair cell' transcription factor genes *Pou4f3* and *Gfi1* are direct Atoh1 targets (Yu et al., 2021; Jen et al., 2022). ISH for *Pou4f3* showed no phenotype in *Tyr* controls (Figure 2A,B; n=0/9 across three batches; Supplementary Table 1) but the mosaic absence of *Pou4f3* in ampullary organ fields and neuromast lines of *Atoh1* crispants (Figure 2C-D[1]; n=13/15 embryos, i.e., 87%, across three batches; Supplementary Table 1). Similarly, ISH for *Gfi1* showed no phenotype in *Tyr* controls (Figure 2E,F; n=0/17 across five batches; Supplementary Table 1), but mosaic absence of *Gfi1* in ampullary organ fields and neuromast lines of *Atoh1* crispants (Figure 2G-H[1]; n=9/14 embryos, i.e., 64%, across four batches; Supplementary Table 1). Post-ISH immunostaining for Sox2 confirmed that neuromasts and ampullary organs were still present in *Atoh1* crispants (Figure 2C[1],D[1],G[1],H[1]).

Genotyping and ICE analysis (Conant et al., 2022) of tails from individual *Atoh1* crispants targeted with this pair of sgRNAs, showed successful disruption of the *Atoh1* gene (Supplementary Figure S3 shows examples). Taken together, these data suggest that Atoh1 lies upstream of *Pou4f3* and *Gfi1* in ampullary organs as well as neuromasts, and is required for the differentiation of electroreceptors as well as hair cells.

Figure 2. *Atoh1* is required for *Pou4f3* and *Gfi1* expression in ampullary organs and neuromasts. Sterlet crispants at stage 45 after *in situ* hybridization for transcription factor genes expressed by developing hair cells. Black arrowheads indicate examples of neuromasts; white arrowheads indicate examples of ampullary organs. (**A,B**) In a control *Tyr* crispant, *Pou4f3* expression is detected in both neuromasts and ampullary organs. (**C-D**^{1}) In an *Atoh1* crispant, *Pou4f3* expression is absent from both neuromasts and ampullary organs, except for a few isolated organs in the postorbital region. Post-ISH Sox2 immunostaining (C^1,D^1) shows that neuromast supporting cells are still present. Most ampullary organs are not visible as Sox2 expression in ampullary organs is significantly weaker than in neuromasts (Supplementary Figure S1A,B) and often not detectable after post-ISH immunostaining. (**E,F**) In a control *Tyr* crispant, *Gfi1* expression is detected in both neuromasts and ampullary organs. (**G-H**^{1}) In an *Atoh1* crispant, *Gfi1* expression is absent from both neuromasts and ampullary organs, except for a few isolated organs in the supra- and infraorbital region. Post-ISH Sox2 immunostaining (G^1,H^1) shows that neuromast supporting cells are still present. (Most ampullary organs are not visible due to weaker Sox2 immunostaining.) Abbreviations: b, barbel; di, dorsal infraorbital ampullary organ field; e, eye; n, naris; S, stage; vi, ventral infraorbital ampullary organ field. Scale bars: 200 μm.

Targeting electrosensory-restricted *Neurod4* had no obvious effect

We previously identified *Neurod4* in paddlefish as the first-reported ampullary organ-restricted transcription factor in the developing lateral line system (Modrell et al., 2017a). We confirmed that sterlet *Neurod4* is similarly expressed by ampullary organs but not neuromasts (Supplementary Figure S1I,J). We targeted *Neurod4* in sterlet embryos using nine different sgRNAs (Table 1; Supplementary Figure S4A), injected in eight different combinations of 1-4 different sgRNAs across 10 independent batches of 1-2 cell-stage embryos (Supplementary Table 1). This had no detectable effect on expression of electroreceptor-specific *Kcnab3* (n=0/44 across nine batches, Supplementary Table 1) or the hair cell/electroreceptor marker *Cacna1d* (n=0/4 across two batches, Supplementary Table 1). Examples of *Neurod4* crispants after ISH for *Kcnab3*, plus a *Tyr* control crispant for comparison, are shown in Supplementary Figure S4B-D. Our sgRNAs were designed prior to publication of the chromosome-level sterlet

genome (Du et al., 2020). Surprisingly, we were unable to find *Neurod4* by BLAST search of the published genome. However, a single copy of *Neurod4* is annotated in the independent Vertebrate Genomes Project paternal haplotype genome assembly (publicly available via the Ensembl Rapid Release website; https://rapid.ensembl.org/Acipenser_ruthenus_GCA_902713435.1), with a prediction score of 99.96% (ENSARTG00010019470.1). This is also a match for the *Neurod4* sequence from our *de novo*-assembled sterlet transcriptome and the cDNA fragment cloned from this. Therefore, it seems that only one copy of *Neurod4* has been retained in the sterlet genome. Genotyping and ICE analysis (Conant et al., 2022) showed successful disruption of the *Neurod4* gene arising from six different combinations of the sgRNAs (Supplementary Table 1; examples are shown in Supplementary Figure S4E-I). Not all sgRNA combinations were genotyped, but all *Neurod4* crispants counted (n=44) included at least one sgRNA confirmed to disrupt the *Neurod4* gene via genotyping of other embryos (Supplementary Table 1).

The lack of phenotype in *Neurod4* crispants, despite successful disruption of the *Neurod4* gene, suggested either that Neurod4 is not required for electroreceptor differentiation, despite its restriction to ampullary organs in both paddlefish and sterlet (Modrell et al., 2017a; this paper), or that it acts redundantly with other transcription factors. In paddlefish, *Neurod1* expression was restricted to developing lateral line ganglia (Modrell et al., 2011b). We cloned sterlet *Neurod1*, *Neurod2* and *Neurod6*. (Unlike *Neurod4*, these three *Neurod* family members are all direct Atoh1 targets in mouse hair cells; Jen et al., 2022.) These three genes all proved to be expressed in sterlet ampullary organs, as well as neuromasts (Supplementary Figure S5). Thus, it seems likely that the lack of effect of CRISPR/Cas9-mediated targeting of sterlet *Neurod4* is due to redundancy with other Neurod family transcription factors co-expressed in ampullary organs. (Our results also show there is at least some variation in *Neurod* family gene expression within the developing lateral line systems of paddlefish and sterlet.)

Targeting mechanosensory-restricted *Foxg1* led to the formation of ectopic ampullary organs within neuromast lines

We recently identified *Foxg1* as a mechanosensory lateral line-restricted transcription factor gene in paddlefish and sterlet (preprint: Minařík et al., 2023). *Foxg1* is expressed in the central zones of lateral line sensory ridges where neuromasts form, though excluded from hair cells themselves (preprint: Minařík et al., 2023). Our sgRNAs against *Foxg1* (Table 1; Supplementary Figure S6A) were designed prior to publication of the chromosome-level sterlet genome (Du et al., 2020). Searching the genome for *Foxg1* showed that both *Foxg1* ohnologs have been retained, on chromosomes 16 and 24, with 99.35% nucleotide identity in

the coding sequence (99.51% amino acid identity). Our sgRNAs target both *Foxg1* ohnologs without mismatches.

When compared with *Tyr* control crispants (Figure 3A-D), a striking phenotype was seen mosaically after targeting mechanosensory-restricted *Foxg1* with sgRNAs 1 and 2 (Table 1; Supplementary Figure S6A): neuromast lines were often interrupted by ectopic ampullary organs. These were defined as *Cacna1d*-expressing cells present within neuromast lines in larger clusters than expected for neuromasts, resembling ampullary organs (Figure 3E-L; n=9/18, i.e., 50%, across two independent batches; Supplementary Table 1), or by expression of electroreceptor-specific *Kcnab3* within neuromast lines, not seen in *Tyr* control crispants (Figure 3M-V; n=7/24, i.e., 29%, across both batches; Supplementary Table 1). This sometimes led to the apparent fusion of ampullary organ fields across a missing neuromast line (e.g., Figure 3E-H,S-V). Post-*Kcnab3* ISH immunostaining for Sox2, to reveal neuromasts, also confirmed that the ectopic electroreceptors in *Foxg1* crispants were within neuromast lines (Figure 3Q[1],R[1]). Targeting *Foxg1* with a different pair of sgRNAs (Table 1; Supplementary Figure 6A) in a different batch of embryos generated similar phenotypes in 3/21 embryos (14%) overall (*Cacna1d*: n=1/6, i.e., 17%; *Kcnab3*: n=2/15, i.e., 13%; Supplementary Table 1).

We also investigated this phenotype by performing ISH for two ampullary organ-restricted transcription factor genes, *Mafa* (preprint: Minařík et al., 2023) and *Neurod4* (Modrell et al., 2017a; this study), followed by Sox2 immunostaining in some embryos to reveal the location of neuromasts. Relative to the normal ampullary organ expression of *Mafa* in uninjected siblings (sufficient *Tyr* control crispants were not available to test; Figure 4A-D; n=0/6 within one batch), ISH for *Mafa* showed the mosaic presence of ampullary organs in neuromast lines and/or merging of ampullary organ fields in *Foxg1* crispants (Figure 4E-J; n=3/17 embryos i.e., 18%, across two independent batches; Supplementary Table 1). Similarly, relative to the usual ampullary organ expression of *Neurod4* in uninjected siblings (sufficient *Tyr* control crispants were not available to test; Figure 4K-N; n=0/2 embryos within one batch; Supplementary Table 1), ISH for *Neurod4* (more weakly expressed than *Mafa*) revealed the same ectopic ampullary organ phenotype in *Foxg1* crispants as seen for *Cacna1d*, *Kcnab3* and *Mafa* (Figure 4O-T; n=3/6 embryos, i.e., 50%, within one batch; Supplementary Table 1). Genotyping and ICE analysis (Conant et al., 2022) of tails from individual *Foxg1* crispants showed successful disruption of the *Foxg1* gene (Supplementary Figure S6 shows examples). Overall, these data suggest that mechanosensory-restricted Foxg1 acts to repress the formation of ampullary organs and electroreceptors within neuromast lines.

Figure 3. *Foxg1* **represses ampullary organ and electroreceptor formation within neuromast lines.** Sterlet crispants at stage 45 after *in situ* hybridization (ISH) for the hair cell and electroreceptor marker *Cacna1d*, or the electroreceptor-specific marker *Kcnab3*. Black arrowheads indicate examples of neuromasts; white arrowheads indicate examples of ampullary organs. (**A,B**) Lateral view of a control *Tyr* crispant. *Cacna1d* expression shows the normal distribution of hair cells and electroreceptors. Note that ampullary organs have significantly more *Cacna1d*-expressing receptor cells than neuromasts. (**C,D**) Ventral view of a second control *Tyr* crispant. *Cacna1d* expression reveals the infraorbital neuromast line on both sides of the ventral rostrum, flanked by the dorsal infraorbital (di) and ventral infraorbital (vi) ampullary organ fields. (**E,F**) Lateral view of a *Foxg1* crispant. *Cacna1d* expression reveals that distinct neuromast lines are missing and the corresponding space is filled by putative ectopic ampullary organs, based on the large, ampullary organ-like clusters of *Cacna1d*-expressing cells. The dorsal and ventral infraorbital ampullary organ fields seem to have fused across the missing neuromast line (compare with A,B). (**G,H**) Ventral view of a second *Foxg1* crispant. *Cacna1d* expression reveals an apparent fusion of the dorsal infraorbital (di) and ventral infraorbital (vi) ampullary organ

fields across the missing infraorbital neuromast lines on both sides (compare with C,D). (**I-K**) In a third *Foxg1* crispant, *Cacna1d* expression on the left side (I,J) shows that distinct supraorbital and infraorbital neuromast lines are still present. However, some organs within the supraorbital line and most organs within the infraorbital line have large clusters of *Cacna1d*-expressing cells, suggesting they are ectopic ampullary organs (green arrowheads in J show examples). On the right side (K,L; image flipped horizontally for ease of comparison), this phenotype is not seen. (**M,N**) Lateral view of a third control *Tyr* crispant. Electroreceptor-specific *Kcnab3* expression shows the distribution of ampullary organs only. (**O,P**) Ventral view of a fourth control *Tyr* crispant. *Kcnab3* expression shows the distribution of ampullary organ fields. Note the lack of staining where the infraorbital neuromast lines run on either side of the ventral rostrum, flanked by the dorsal infraorbital (di) and ventral infraorbital (vi) ampullary organ fields (compare with *Cacna1d* expression in C,D). (**Q-R[1]**) Lateral view of a fourth *Foxg1* crispant. *Kcnab3* expression shows two ectopic ampullary organs (green arrowheads) within the infraorbital neuromast line (compare with M,N). Post-ISH Sox2 immunostaining for supporting cells (Q[1],R[1]) confirms the position of the neuromast lines. (**S,T**) Ventral view of a fifth *Foxg1* crispant. On the left side, ectopic *Kcnab3*-expressing electroreceptors fill the space where the left infraorbital neuromast line would normally run, such that the dorsal and ventral infraorbital ampullary organ fields seem to have fused (compare with O,P). On the right side, a single ectopic *Kcnab3*-expressing ampullary organ is visible (green arrowhead) where the right infraorbital neuromast line runs (compare with O,P). (**U,V**) Lateral view of a sixth *Foxg1* crispant. *Kcnab3* expression shows fusion of the dorsal infraorbital ampullary organ field with the dorsal part of the ventral infraorbital ampullary organ field (compare with M,N). Abbreviations: b, barbel; di, dorsal infraorbital ampullary organ field; di/vi, fused dorsal infraorbital and ventral infraorbital ampullary organ fields; e, eye; m, mouth; n, naris; S, stage; vi, ventral infraorbital ampullary organ field. Scale bars: 200 µm.

416
Figure 4. Ectopic ampullary organs in *Foxg1* crispants express ampullary organ-specific
transcription factor genes *Mafa* and *Neurod4*. Sterlet crispants at stage 45 after *in situ* hybridization
(ISH) for ampullary organ-restricted transcription factor genes. Black arrowheads indicate examples of
neuromasts; white arrowheads indicate examples of ampullary organs. (**A-D**) In an uninjected sibling,
Mafa expression is restricted to ampullary organs (lateral view: A,B; ventral view: C,D). (**E-J**) A *Foxg1*
crispant. On the left side of the head (E,F), several *Mafa*-expressing ectopic ampullary organs are
present in the space where the infraorbital neuromast line would normally run, such that the dorsal and
ventral infraorbital ampullary organ fields seem to have fused (compare with A,B). Post-ISH Sox2
immunostaining (E[1],F[1]) shows that neuromasts are still present both proximally and distally to the sites
of ampullary organ field fusion. In ventral view (G,H), ectopic ampullary organs (green arrowheads) are
seen bilaterally, within the spaces where the infraorbital neuromast lines run on either side of the ventral
rostrum (compare with C,D). On the right side in lateral view (I,J; image flipped horizontally for ease of
comparison), a single *Mafa*-expressing ectopic ampullary organ (green arrowhead) is also present in
the space where the infraorbital neuromast line runs (compare with A,B). (**K-N**) In an uninjected sibling,

Neurod4 expression is restricted to ampullary organs (lateral view: K,L; ventral view: M,N). (**O-T**) A *Foxg1* crispant. On the left side of the head (O,P), *Neurod4*-expressing ectopic ampullary organs are present in the space where the infraorbital neuromast line would normally run, such that the dorsal and ventral infraorbital ampullary organ fields seem to have fused (compare with K,L). Post-ISH Sox2 immunostaining (O[1],P[1]) suggests that neuromasts are absent from the site of infraorbital ampullary organ field fusion, although neuromasts can be seen in the preopercular and trunk lines (black arrowheads, compare with O,P). In ventral view (Q,R), ectopic *Neurod4*-expressing ampullary organs are seen where the right infraorbital neuromast line would normally run on the ventral rostrum (green arrowhead indicates an example), resulting in partial fusion of the dorsal and ventral infraorbital fields on this side (the left side is unaffected). On the right side in lateral view (S,T; image flipped horizontally for ease of comparison), ectopic ampullary organs are also present in the space where the infraorbital neuromast line runs (green arrowhead in T indicates an example), resulting in the apparent partial fusion of the dorsal and ventral infraorbital ampullary organ fields (compare with K,L). Abbreviations: b, barbel; di, dorsal infraorbital ampullary organ field; di/vi, fused dorsal infraorbital and ventral infraorbital ampullary organ fields; e, eye; m, mouth; n, naris; S, stage; vi, ventral infraorbital ampullary organ field. Scale bars: 200 μm.

Discussion

Conserved molecular mechanisms underlie lateral line electroreceptor and hair cell formation

Here, we aimed to test the function in lateral line electroreceptor and/or hair cell formation of three transcription factor genes that we had previously identified as expressed in developing electrosensory ampullary organ and/or mechanosensory lateral line organs in ray-finned chondrostean fishes - paddlefish and sterlet (Butts et al., 2014; Modrell et al., 2017a; preprint: Minařík et al., 2023). The first gene we investigated was *Atoh1*, which is required for the formation of lateral line hair cells in zebrafish (Millimaki et al., 2007), as well as hair cells in the inner ear (Bermingham et al., 1999; Millimaki et al., 2007). In paddlefish (Butts et al., 2014; Modrell et al., 2017a) and sterlet (preprint: Minařík et al., 2023), *Atoh1* is expressed in ampullary organs as well as neuromasts. Targeting *Atoh1* for CRISPR/Cas9-mediated mutagenesis in F0 sterlet embryos showed that Atoh1 is required for the formation not only of *Cacna1d*-expressing neuromast hair cells, as expected from zebrafish (Millimaki et al., 2007), but also of *Cacna1d*-expressing, *Kcnab3*-expressing electroreceptors. These experiments also showed that Atoh1 is required for the expression of the 'hair cell' transcription factor genes *Gfi1* and *Pou4f3* in developing ampullary organs, as well as neuromasts. This is consistent with both of these genes being direct Atoh1 targets in mouse cochlear hair cells (Yu et al., 2021; Jen et al., 2022).

In both inner-ear hair cells and Merkel cells (epidermal mechanoreceptors found in all vertebrates; see e.g., Whitear, 1989; Brown et al., 2023), Atoh1 acts with Pou4f3 in a conserved 'feed-forward circuit', with Atoh1 directly activating *Pou4f3* expression, and Pou4f3 then acting as a pioneer factor to open a significant subset of Atoh1 target enhancers (some shared and some divergent between hair cells and Merkel cells), enabling mechanosensory

differentiation (Yu et al., 2021). In hair cells, *Gfi1* is one of the Pou4f3-dependent Atoh1 targets
(Yu et al., 2021). Together with the striking conservation of transcription factor gene
expression between developing ampullary organs and neuromasts (Modrell et al., 2011b;
Modrell et al., 2011a; Modrell et al., 2017b; Modrell et al., 2017a; preprint: Minařík et al., 2023),
the phenotypes seen in *Atoh1*-targeted F0 sterlet crispant embryos suggest that the molecular
mechanisms underlying electroreceptor formation are highly conserved with those underlying
hair cell formation. Indeed, the requirement of Atoh1 for *Pou4f3* and *Gfi1* expression in
ampullary organs, as well as neuromasts, suggests that the Atoh1-Pou4f3 'feedforward circuit'
in mechanosensory cells - i.e., hair cells and epidermal Merkel cells (Yu et al., 2021) - may
also be conserved, at least partly, in developing electroreceptors. Taken together, these data
support the hypothesis that electroreceptors evolved as a transcriptionally related "sister cell
type" to lateral line hair cells (Arendt et al., 2016; Baker and Modrell, 2018; Baker, 2019).
485

Electrosensory-restricted Neurod4 is likely redundant with other Neurod family
members in sterlet
Paddlefish *Neurod4* was the first-reported ampullary organ-restricted transcription factor gene
(Modrell et al., 2017a), with conserved expression in sterlet (this study). We were unable to
detect a lateral line organ phenotype in *Neurod4*-targeted sterlet crispants. However, we found
that *Neurod1*, *Neurod2* and *Neurod6* are all expressed in sterlet ampullary organs (as well as
neuromasts), suggesting that Neurod4 may act redundantly with one or more of these factors
in developing ampullary organs. (In paddlefish, however, *Neurod1* expression is restricted to
developing lateral line ganglia; Modrell et al., 2011b.) Targeting multiple *Neurod* genes for
CRISPR/Cas9-mediated mutagenesis in the future may shed light on the role played by this
transcription factor family in ampullary organ development.
497

Foxg1 represses electroreceptor formation in the neuromast-forming central zone of
lateral line sensory ridges
We also targeted *Foxg1*, a mechanosensory-restricted transcription factor gene that we
recently identified in the developing lateral line system of paddlefish and sterlet (preprint:
Minařík et al., 2023). *Foxg1* is expressed in the central zones of lateral line sensory ridges
where neuromasts form, though excluded from the central domains of neuromasts where hair
cells differentiate (preprint: Minařík et al., 2023). Targeting *Foxg1* for CRISPR/Cas9-mediated
mutagenesis in F0 sterlet embryos led to a striking phenotype: the formation within neuromast
lines of ectopic electroreceptors, often in the large clusters normally seen in ampullary organs.
In some cases, ampullary organ fields, which normally flank neuromast lines, effectively
'merged' across missing neuromast lines. This phenotype was revealed by examining
expression of the electroreceptor-specific marker *Kcnab3*, and two ampullary organ-restricted

transcription factor genes: *Mafa* (preprint: Minařík et al., 2023) and *Neurod4* (Modrell et al., 2017a). Thus, *Foxg1* seems to repress an ampullary organ fate within the central domain of lateral line sensory ridges where neuromasts form.

In the mouse inner ear, *Foxg1* is expressed in the prospective cochlea and all sensory patches, in hair cell progenitors and supporting cells (Pauley et al., 2006; Tasdemir-Yilmaz et al., 2021), plus a subset of hair cells (Pauley et al., 2006). Knockout leads to a shorter cochlea with extra rows of hair cells, and to loss or reduction of vestibular end organs (Pauley et al., 2006; Hwang et al., 2009). More hair cells and fewer supporting cells were seen after conditional knockdown of *Foxg1* in neonatal cochlear supporting cells, possibly via the transdifferentiation of supporting cells (Zhang et al., 2019; Zhang et al., 2020). This suggests the possibility that Foxg1 may act in the central zone of lateral line sensory ridges to maintain a proliferative progenitor state, as it does in the mouse olfactory epithelium (Kawauchi et al., 2009).

Furthermore, Fox family members can act as pioneer factors as well as transcription factors (Golson and Kaestner, 2016; Lukoseviciute et al., 2018). A pioneer factor role has been proposed for Foxi3 in otic placode development (see Singh and Groves, 2016). In the developing neural crest, Foxd3 acts early as a pioneer factor, opening enhancers and repositioning nucleosomes to prime genes controlling neural crest specification and migration and, concurrently, to repress the premature differentiation of, e.g., melanocytes (Lukoseviciute et al., 2018). Later in neural crest development, Foxd3 represses enhancers associated with mesenchymal, neuronal and melanocyte lineages (Lukoseviciute et al., 2018). In cortical progenitors, Foxg1 suppresses the adoption at later stages of an early-born cell fate, namely, Cajal-Retzius cells (Hanashima et al., 2004). In the developing chicken otic placode, Foxg1 represses markers of other lineages, such as the olfactory and lens placodes, and epidermis (Anwar et al., 2017). Hence, it is possible that Foxg1 acts in the central zone of lateral line sensory ridges in electroreceptive fishes as a pioneer factor for neuromast/hair cell formation, and/or that it represses ampullary organ/electroreceptor formation. Such a role would be consistent with lateral line expression of *Foxg1* only in electroreceptive fishes: *Foxg1* is not expressed in the developing lateral line of zebrafish or *Xenopus* (e.g., Dirksen and Jamrich, 1995; Papalopulu and Kintner, 1996; Toresson et al., 1998; Eagleson and Dempewolf, 2002; Duggan et al., 2008; Zhao et al., 2009).

Overall, these data lead us to propose the speculative hypothesis that electrosensory organs may be the 'default' fate within lateral line sensory ridges in electroreceptive vertebrates, and that Foxg1 represses this fate to enable mechanosensory neuromasts and hair cells to form. To test these hypotheses directly, it will be important in the future to identify global changes in gene expression and chromatin accessibility in the absence of Foxg1.

Summary and Perspective

Overall, we have found that the 'hair cell' transcription factor Atoh1 is required for the formation of lateral line electroreceptors as well as hair cells, consistent with a close developmental relationship between these putative 'sister cell' types. Electrosensory-restricted Neurod4 may act redundantly with other Neurod family members expressed in developing ampullary organs. Mechanosensory-restricted Foxg1 represses the formation of electroreceptors within neuromast lines, suggesting the surprising possibility that electroreceptors are the 'default' fate within lateral line sensory ridges and raising interesting developmental and evolutionary questions for future investigation.

<u>Materials and Methods</u>

Animals

Fertilized sterlet (*Acipenser ruthenus*) eggs were obtained from the breeding facility at the Research Institute of Fish Culture and Hydrobiology, Faculty of Fisheries and Protection of Waters, University of South Bohemia in České Budějovice, Vodňany, Czech Republic, and staged according to Dettlaff et al. (1993). For detailed information about sterlet husbandry, *in vitro* fertilization and the rearing of embryos and yolk-sac larvae, see Stundl et al. (2022). Upon reaching the desired developmental stages, embryos and yolk-sac larvae were euthanized by overdose of MS-222 (Sigma-Aldrich) and fixed in modified Carnoy's fixative (6 volumes 100% ethanol: 3 volumes 37% formaldehyde: 1 volume glacial acetic acid) for 3 hours at room temperature, dehydrated stepwise into 100% ethanol and stored at -20°C.

All experimental procedures were approved by the Animal Research Committee of the Faculty of Fisheries and Protection of Waters in Vodňany, University of South Bohemia in České Budějovice, Czech Republic, and by the Ministry of Agriculture of the Czech Republic (reference number: MSMT-12550/2016-3). Experimental fish were maintained according to the principles of the European Union (EU) Harmonized Animal Welfare Act of the Czech Republic, and Principles of Laboratory Animal Care and National Laws 246/1992 "Animal Welfare" on the protection of animals.

CRISPR guide RNA design and synthesis

Target gene sequences were identified using the National Center for Biotechnology Information (NCBI) Basic Local Alignment Search Tool BLAST (https://blast.ncbi.nlm.nih.gov/Blast.cgi; McGinnis and Madden, 2004) to search sterlet transcriptomic data (available at DDBJ/EMBL/GenBank under the accessions GKLU00000000 and GKEF01000000; see preprint: Minařík et al., 2023) or draft genomic sequence data (M.H., unpublished) with the relevant paddlefish sequence (Modrell et al.,

2017a). Transcriptomic sequence data were searched for *Tyr*, *Atoh1* and *Neurod4*; genomic sequence data were searched for *Foxg1*. Since the project started, a chromosome-level sterlet genome has been published (Du et al., 2020) and an independent high-quality genome assembly from the Vertebrate Genomes Project has been made publicly available at RapidEnsembl (https://rapid.ensembl.org/Acipenser_ruthenus_GCA_902713435.1). Open reading frames were identified using the NCBI ORF Finder tool (https://www.ncbi.nlm.nih.gov/orffinder/) and exons annotated by comparison with reference anamniote species (*Lepisosteus oculatus*, *Danio rerio*, *Xenopus tropicalis*) available via Ensembl (https://www.ensembl.org; Cunningham et al., 2022). Conserved domains were identified using NCBI BLASTX (https://blast.ncbi.nlm.nih.gov/Blast.cgi; McGinnis and Madden, 2004). gRNAs were preferentially designed to target 5' exons, ideally upstream of or within regions encoding known functional domains, to increase the probability of disrupting gene function. gRNAs were designed using the CRISPR Guide RNA Design Tool from Benchling (https://benchling.com) and selected for synthesis based on the following criteria: (1) a high on-target score, ideally no less than 0.5; (2) no off-target matches identified within coding sequences in transcriptome and genome data, unless there were at least two mismatches in the 3' seed sequence (8-10 bp upstream of the protospacer adjacent motif [PAM], or in the PAM itself); (3) if multiplexing, the gRNA pair were ideally within 50-150 bases of each other to increase the probability of fragment deletion.

DNA templates for CRISPR gRNAs were synthesized using plasmid pX335-U6-Chimeric_BB-CBh-hSpCas9n(D10A) (Addgene, plasmid #42335; Cong et al., 2013) containing the gRNA scaffold. The gRNA scaffold was amplified using a specific forward primer for each gRNA, with an overhang containing the gRNA target sequence and T7 promoter, and a reverse primer that was identical for all reactions. For gRNAs that did not start with G, an additional G was added at the start to ensure efficient transcription. The DNA template was amplified using Q5 polymerase (New England Biolabs, NEB) and purified using the Monarch PCR & DNA Cleanup Kit (NEB). The gRNAs were synthesized using the HiScribe T7 High Yield RNA Synthesis Kit (NEB) and purified using the Monarch RNA Cleanup Kit (NEB) and stored at -80°C before use. Alternatively, chemically modified synthetic gRNAs were ordered directly from Synthego (CRISPRevolution sgRNA EZ Kit).

Embryo injection

On the day of injection, 1200 ng gRNA were mixed with 2400 ng Cas9 protein with NLS (PNA Bio) in 5 µl nuclease-free water and incubated for 10 minutes at room temperature to form ribonucleoprotein (RNP) complexes. For gRNA multiplexing, two RNP mixes were combined 1:1 to a final volume of 5 µl, and 0.5 µl of 10x 10,000 MW rhodamine dextran (Invitrogen) added to better visualize the injection mixture. Injection mixtures were kept on ice throughout

the injection session. 50 µl glass microcapillaries (Drummond Microcaps) were pulled in a capillary needle puller (PC-10, Narishige) set to 58°C with two light and one heavy weights, in single-stage pulling mode. Fertilized sterlet eggs were manually dechorionated using Dumont #5 forceps. A 1000 µl pipette tip cut to the same diameter as a dechorionated sterlet egg was used to prepare a series of wells in an agar plate to allow ideal egg positioning during injection using an automatic microinjector (FemtoJet 4x, Eppendorf), set to 100 hPa. Approximately 20 nl of the injection mixture (corresponding to approximately 4.8 ng gRNAs and 9.6 ng Cas9) were injected into fertilized eggs or two-cell stage embryos, targeting the animal pole at a 45° angle. Injected embryos were moved to a clean Petri dish and, for optimum Cas9 efficiency, kept at room temperature until the end of the injecting session or until at least the 32-cell stage, then moved to a 16°C incubator. No more than 30 eggs were kept per 90 mm Petri dish. Unfertilized eggs and dead embryos were removed at the end of the injection day. Petri dishes were checked regularly for dead embryos and the water was changed at least twice a day before gastrulation was completed, and once daily post-gastrulation. Hatched larvae were kept for approximately 16 days post fertilisation until stage 45, then euthanized by MS-222 overdose and fixed with modified Carnoy's fixative (see above). Fixed larvae were then dehydrated stepwise into 100% ethanol and stored at -20°C.

Gene cloning, *in situ* hybridization and immunohistochemistry

Total RNA was isolated from sterlet embryos using Trizol (Invitrogen, Thermo Fisher Scientific), following the manufacturer's protocol, and cDNA synthesized using the Superscript III First Strand Synthesis kit (Invitrogen, Thermo Fisher Scientific). We used our sterlet transcriptome assemblies (from pooled yolk-sac larvae at stages 40-45; preprint: Minařík et al., 2023), which are available at DDBJ/EMBL/GenBank under the accessions GKLU00000000 and GKEF01000000, to design primers for *Neurod4* (forward: GAGAGAGCCCCAAAGAGACGAG; reverse: CTGCTTGAGCGAGAAGTTGACG). cDNA fragments amplified under standard PCR conditions were cloned into the pDrive cloning vector (Qiagen). Individual clones were verified by sequencing (Department of Biochemistry Sequencing Facility, University of Cambridge, UK) and sequence identity verified using NCBI BLAST (https://blast.ncbi.nlm.nih.gov/Blast.cgi; McGinnis and Madden, 2004). For *Neurod1*, *Neurod2* and *Neurod6*, synthetic gene fragments based on sterlet transcriptome data, with added M13 forward and reverse primer adaptors, were ordered from Twist Bioscience. GenBank accession numbers are as follows: *Neurod1* OQ808944, *Neurod2* OQ808945, *Neurod4* OQ808946, *Neurod6* OQ808947. The other genes used in this study have been published (preprint: Minařík et al., 2023).

Digoxigenin-labelled riboprobes were synthesized as previously described (preprint: Minařík et al., 2023). Whole-mount *in situ* hybridization (ISH) was performed as previously

described (Modrell et al., 2011a). Whole-mount immunostaining post-ISH for Sox2 (rabbit monoclonal, 1:200; ab92494, Abcam) was performed as previously described (Metscher and Müller, 2011), using a horseradish peroxidase-conjugated goat anti-rabbit antibody (1:300, Jackson ImmunoResearch) and the metallographic peroxidase substrate EnzMet kit (Nanoprobes) as per the manufacturer's instructions.

Genotyping

To confirm successful mutation in targeted regions, genotyping was performed on trunk and tail tissue that had been removed before ISH and stored in 100% ethanol at -20°C. The tissue was digested using Rapid Extract Lysis Kit (PCR Biosystems) and the target region amplified using HS Taq Mix Red (PCR Biosystems) according to the manufacturer's instructions. Primers used for genotyping are listed in Supplementary file 1. After agarose gel electrophoresis, PCR products were extracted using MinElute Gel Extraction Kit (Qiagen) and submitted for sequencing (Genewiz by Azenta Life Sciences). To analyze CRISPR editing efficiency, Sanger trace files were uploaded to the Synthego Inference of CRISPR Edits (ICE) tool (https://ice.synthego.com; Conant et al., 2022).

Data availability statement

The publication and associated supplementary figures include representative example images of embryos from each experiment. Additional raw data underlying this publication consist of further images of these and other embryos from each experiment. Public sharing of these images is not cost-efficient but they are available from the corresponding author upon reasonable request.

Ethics statement

Sterlet animal work was reviewed and approved by The Animal Research Committee of Research Institute of Fish Culture and Hydrobiology, Faculty of Fisheries and Protection of Waters, University of South Bohemia in České Budějovice, Vodňany, Czech Republic and Ministry of Agriculture of the Czech Republic (MSMT-12550/2016-3).

Author contributions

CB conceived and designed the project and wrote the manuscript together with MM. MM performed most of the experiments, prepared all the manuscript figures and made a significant contribution to the writing of the manuscript. AC and RF performed some experiments. MV made essential contributions to the experiments. MH provided draft genome sequence data. MP and DG were instrumental in enabling all work with sterlet embryos. All authors read and commented on the manuscript.

Funding

This work was supported by the Biotechnology and Biological Sciences Research Council (BBSRC: grant BB/P001947/1 to CB). Additional support for MM was provided by the Cambridge Isaac Newton Trust (grant 20.07[c] to CB) and by the School of the Biological Sciences, University of Cambridge. AC was supported by a PhD research studentship from the Anatomical Society. The work of RF and MP was supported by the Ministry of Education, Youth and Sports of the Czech Republic, projects CENAKVA (LM2018099), Biodiversity (CZ.02.1.01/0.0/0.0/16_025/0007370) and Czech Science Foundation (20-23836S).

Rights Retention Statement

This work was funded by a grant from the Biotechnology and Biological Sciences Research Council (BBSRC: BB/P001947/1). For the purpose of open access, the author has applied a Creative Commons Attribution (CC BY) licence to any Author Accepted Manuscript version arising.

Acknowledgements

Thanks to Melinda Modrell for pilot work on CRISPR/Cas9 in paddlefish and for her advice and helpful discussions in planning the project. Thanks to Marek Rodina and Martin Kahanec for their help with sterlet spawns. Thanks to David Jandzik, András Simon, Alberto Joven Araus, Ahmed Elewa, and Mustafa Khokha for invaluable advice on CRISPR/Cas9 approaches and genotyping. Thanks to Jan Stundl and David Jandzik for sharing their sterlet *Tyr* gRNA sequences prior to publication. Thanks to Christine Hirschberger and Rolf Ericsson for their help with some of the *in situ* hybridization rounds.

References

Anwar, M., Tambalo, M., Ranganathan, R., Grocott, T., Streit, A., 2017. A gene network regulated by FGF signalling during ear development. Sci. Rep. 7, 6162.

Arendt, D., Musser, J.M., Baker, C.V.H., Bergman, A., Cepko, C., Erwin, D.H., Pavlicev, M., Schlosser, G., Widder, S., Laubichler, M.D., Wagner, G.P., 2016. The origin and evolution of cell types. Nat. Rev. Genet. 17, 744-757.

Baker, C.V.H., 2019. The development and evolution of lateral line electroreceptors: insights from comparative molecular approaches., in: B.A. Carlson, J.A. Sisneros, A.N. Popper, R.R. Fay (Eds.), Electroreception: Fundamental Insights from Comparative Approaches. Springer, Cham, pp. 25-62.

Baker, C.V.H., Modrell, M.S., 2018. Insights into electroreceptor development and evolution from molecular comparisons with hair cells. Integr. Comp. Biol. 58, 329-340.

Baker, C.V.H., Modrell, M.S., Gillis, J.A., 2013. The evolution and development of vertebrate lateral line electroreceptors. J. Exp. Biol. 216, 2515-2522.

Baloch, A.R., Franěk, R., Tichopád, T., Fučíková, M., Rodina, M., Pšenička, M., 2019. *Dnd1* knockout in sturgeons by CRISPR/Cas9 generates germ cell free host for surrogate production. Animals (Basel) 9, 174.

Bellono, N.W., Leitch, D.B., Julius, D., 2017. Molecular basis of ancestral vertebrate electroreception. Nature 543, 391-396.

Bellono, N.W., Leitch, D.B., Julius, D., 2018. Molecular tuning of electroreception in sharks and skates. Nature 558, 122-126.

Bermingham, N.A., Hassan, B.A., Price, S.D., Vollrath, M.A., Ben-Arie, N., Eatock, R.A., Bellen, H.J., Lysakowski, A., Zoghbi, H.Y., 1999. *Math1*: an essential gene for the generation of inner ear hair cells. Science 284, 1837-1841.

Bodznick, D., Montgomery, J.C., 2005. The physiology of low-frequency electrosensory systems, in: T.H. Bullock, C.D. Hopkins, A.N. Popper, R.R. Fay (Eds.), Electroreception. Springer, New York, pp. 132-153.

Brown, T.L., Horton, E.C., Craig, E.W., Goo, C.E.A., Black, E.C., Hewitt, M.N., Yee, N.G., Fan, E.T., Raible, D.W., Rasmussen, J.P., 2023. Dermal appendage-dependent patterning of zebrafish *atoh1a+* Merkel cells. eLife 12, e85800.

Bullock, T.H., Bodznick, D.A., Northcutt, R.G., 1983. The phylogenetic distribution of electroreception: evidence for convergent evolution of a primitive vertebrate sense modality. Brain Res. Rev. 287, 25-46.

Butts, T., Modrell, M.S., Baker, C.V.H., Wingate, R.J.T., 2014. The evolution of the vertebrate cerebellum: absence of a proliferative external granule layer in a non-teleost ray-finned fish. Evol. Dev. 16, 92-100.

Caprara, G.A., Peng, A.W., 2022. Mechanotransduction in mammalian sensory hair cells. Mol.
Cell. Neurosci. 120, 103706.

Chagnaud, B.P., Wilkens, L.A., Hofmann, M., 2021. The ampullary electrosensory system –a
paddlefish case study, in: B. Fritzsch (Ed.), The Senses: A Comprehensive Reference.
Elsevier, pp. 215-227.

Chen, J., Wang, W., Tian, Z., Dong, Y., Dong, T., Zhu, H., Zhu, Z., Hu, H., Hu, W., 2018.
Efficient gene transfer and gene editing in sterlet. Front. Genet. 9, 117.

Chen, Y., Gu, Y., Li, Y., Li, G.L., Chai, R., Li, W., Li, H., 2021. Generation of mature and
functional hair cells by co-expression of *Gfi1*, *Pou4f3*, and *Atoh1* in the postnatal mouse
cochlea. Cell Rep. 35, 109016.

Conant, D., Hsiau, T., Rossi, N., Oki, J., Maures, T., Waite, K., Yang, J., Joshi, S., Kelso, R.,
Holden, K., Enzmann, B.L., Stoner, R., 2022. Inference of CRISPR edits from Sanger
trace data. CRISPR J. 5, 123-130.

Cong, L., Ran, F.A., Cox, D., Lin, S., Barretto, R., Habib, N., Hsu, P.D., Wu, X., Jiang, W.,
Marraffini, L.A., Zhang, F., 2013. Multiplex genome engineering using CRISPR/Cas
systems. Science 339, 819-823.

Crampton, W.G.R., 2019. Electroreception, electrogenesis and electric signal evolution. J.
Fish Biol. 95, 92-134.

Cunningham, F., Allen, J.E., Allen, J., Alvarez-Jarreta, J., Amode, M.R., Armean, I.M., Austine-
Orimoloye, O., Azov, A.G., Barnes, I., Bennett, R., Berry, A., Bhai, J., Bignell, A., Billis,
K., Boddu, S., Brooks, L., Charkhchi, M., Cummins, C., Da Rin Fioretto, L., Davidson,
C., Dodiya, K., Donaldson, S., El Houdaigui, B., El Naboulsi, T., Fatima, R., Giron, C.G.,
Genez, T., Martinez, J.G., Guijarro-Clarke, C., Gymer, A., Hardy, M., Hollis, Z., Hourlier,
T., Hunt, T., Juettemann, T., Kaikala, V., Kay, M., Lavidas, I., Le, T., Lemos, D.,
Marugán, J.C., Mohanan, S., Mushtaq, A., Naven, M., Ogeh, D.N., Parker, A., Parton,
A., Perry, M., Piližota, I., Prosovetskaia, I., Sakthivel, M.P., Salam, A.I.A., Schmitt, B.M.,
Schuilenburg, H., Sheppard, D., Pérez-Silva, J.G., Stark, W., Steed, E., Sutinen, K.,
Sukumaran, R., Sumathipala, D., Suner, M.M., Szpak, M., Thormann, A., Tricomi, F.F.,
Urbina-Gómez, D., Veidenberg, A., Walsh, T.A., Walts, B., Willhoft, N., Winterbottom,
A., Wass, E., Chakiachvili, M., Flint, B., Frankish, A., Giorgetti, S., Haggerty, L., Hunt,
S.E., Ilsley, G.R., Loveland, J.E., Martin, F.J., Moore, B., Mudge, J.M., Muffato, M.,
Perry, E., Ruffier, M., Tate, J., Thybert, D., Trevanion, S.J., Dyer, S., Harrison, P.W.,
Howe, K.L., Yates, A.D., Zerbino, D.R., Flicek, P., 2022. Ensembl 2022. Nucleic Acids
Res. 50, D988-D995.

Dettlaff, T.A., Ginsburg, A.S., Schmalhausen, O.I., 1993. Sturgeon Fishes: Developmental
Biology and Aquaculture. Springer-Verlag, Berlin.

Dirksen, M.-L., Jamrich, M., 1995. Differential expression of *fork head* genes during early
Xenopus and zebrafish development. Dev. Genet. 17, 107-116.
Du, K., Stöck, M., Kneitz, S., Klopp, C., Woltering, J.M., Adolfi, M.C., Feron, R., Prokopov, D.,
Makunin, A., Kichigin, I., Schmidt, C., Fischer, P., Kuhl, H., Wuertz, S., Gessner, J.,
Kloas, W., Cabau, C., Iampietro, C., Parrinello, H., Tomlinson, C., Journot, L.,
Postlethwait, J.H., Braasch, I., Trifonov, V., Warren, W.C., Meyer, A., Guiguen, Y.,
Schartl, M., 2020. The sterlet sturgeon genome sequence and the mechanisms of
segmental rediploidization. Nat. Ecol. Evol. 4, 841-852.
Duggan, C.D., Demaria, S., Baudhuin, A., Stafford, D., Ngai, J., 2008. Foxg1 is required for
development of the vertebrate olfactory system. J. Neurosci. 28, 5229-5239.
Eagleson, G.W., Dempewolf, R.D., 2002. The role of the anterior neural ridge and *Fgf-8* in
early forebrain patterning and regionalization in *Xenopus laevis*. Comp. Biochem.
Physiol. B Biochem. Mol. Biol. 132, 179-189.
Elewa, A., Wang, H., Talavera-López, C., Joven, A., Brito, G., Kumar, A., Hameed, L.S.,
Penrad-Mobayed, M., Yao, Z., Zamani, N., Abbas, Y., Abdullayev, I., Sandberg, R.,
Grabherr, M., Andersson, B., Simon, A., 2017. Reading and editing the *Pleurodeles waltl*
genome reveals novel features of tetrapod regeneration. Nat. Commun. 8, 2286.
Fei, J.-F., Lou, W.P.-K., Knapp, D., Murawala, P., Gerber, T., Taniguchi, Y., Nowoshilow, S.,
Khattak, S., Tanaka, E.M., 2018. Application and optimization of CRISPR-Cas9-
mediated genome engineering in axolotl (*Ambystoma mexicanum*). Nat. Protoc. 13,
2908-2943.
Flowers, G.P., Timberlake, A.T., McLean, K.C., Monaghan, J.R., Crews, C.M., 2014. Highly
efficient targeted mutagenesis in axolotl using Cas9 RNA-guided nuclease.
Development 141, 2165-2171.
Gillis, J.A., Modrell, M.S., Northcutt, R.G., Catania, K.C., Luer, C.A., Baker, C.V.H., 2012.
Electrosensory ampullary organs are derived from lateral line placodes in cartilaginous
fishes. Development 139, 3142-3146.
Golson, M.L., Kaestner, K.H., 2016. Fox transcription factors: from development to disease.
Development 143, 4558-4570.
Hanashima, C., Li, S.C., Shen, L., Lai, E., Fishell, G., 2004. *Foxg1* suppresses early cortical
cell fate. Science 303, 56-59.
Hernández, P.P., Olivari, F.A., Sarrazin, A.F., Sandoval, P.C., Allende, M.L., 2007.
Regeneration in zebrafish lateral line neuromasts: expression of the neural progenitor
cell marker sox2 and proliferation-dependent and-independent mechanisms of hair cell
renewal. Dev. Neurobiol. 67, 637-654.

Hwang, C.H., Simeone, A., Lai, E., Wu, D.K., 2009. *Foxg1* is required for proper separation and formation of sensory cristae during inner ear development. Dev. Dyn. 238, 2725-2734.

Iyer, A.A., Groves, A.K., 2021. Transcription factor reprogramming in the inner ear: turning on cell fate switches to regenerate sensory hair cells. Front. Cell. Neurosci. 15, 660748.

Iyer, A.A., Hosamani, I., Nguyen, J.D., Cai, T., Singh, S., McGovern, M.M., Beyer, L., Zhang, H., Jen, H.I., Yousaf, R., Birol, O., Sun, J.J., Ray, R.S., Raphael, Y., Segil, N., Groves, A.K., 2022. Cellular reprogramming with ATOH1, GFI1, and POU4F3 implicate epigenetic changes and cell-cell signaling as obstacles to hair cell regeneration in mature mammals. eLife 11, e79712.

Jen, H.-I., Singh, S., Tao, L., Maunsell, H.R., Segil, N., Groves, A.K., 2022. GFI1 regulates hair cell differentiation by acting as an off-DNA transcriptional co-activator of ATOH1, and a DNA-binding repressor. Sci. Rep. 12, 7793.

Jørgensen, J.M., 2005. Morphology of electroreceptive sensory organs, in: T.H. Bullock, C.D. Hopkins, A.N. Popper, R.R. Fay (Eds.), Electroreception. Springer, New York, pp. 47-67.

Kawauchi, S., Santos, R., Kim, J., Hollenbeck, P.L.W., Murray, R.C., Calof, A.L., 2009. The role of *Foxg1* in the development of neural stem cells of the olfactory epithelium. Ann. N. Y. Acad. Sci. 1170, 21-27.

Kniss, J.S., Jiang, L., Piotrowski, T., 2016. Insights into sensory hair cell regeneration from the zebrafish lateral line. Curr. Opin. Genet. Dev. 40, 32-40.

Leitch, D.B., Julius, D., 2019. Electrosensory transduction: comparisons across structure, afferent response properties, and cellular physiology., in: B.A. Carlson, J.A. Sisneros, A.N. Popper, R.R. Fay (Eds.), Electroreception: Fundamental Insights from Comparative Approaches. Springer, Cham, pp. 63-90.

Lenhardt, M., Finn, R.N., Cakic, P., Kolarevic, J., Krpocetkovic, J., Radovic, I., Fyhn, H.J., 2005. Analysis of the post-vitellogenic oocytes of three species of Danubian Acipenseridae. Belg. J. Zool. 135, 205.

Lukoseviciute, M., Gavriouchkina, D., Williams, R.M., Hochgreb-Hagele, T., Senanayake, U., Chong-Morrison, V., Thongjuea, S., Repapi, E., Mead, A., Sauka-Spengler, T., 2018. From pioneer to repressor: bimodal foxd3 activity dynamically remodels neural crest regulatory landscape *in vivo*. Dev. Cell 47, 608-628.

Lush, M.E., Diaz, D.C., Koenecke, N., Baek, S., Boldt, H., St Peter, M.K., Gaitan-Escudero, T., Romero-Carvajal, A., Busch-Nentwich, E.M., Perera, A.G., Hall, K.E., Peak, A., Haug, J.S., Piotrowski, T., 2019. scRNA-Seq reveals distinct stem cell populations that drive hair cell regeneration after loss of Fgf and Notch signaling. eLife 8, e44431.

McGinnis, S., Madden, T.L., 2004. BLAST: at the core of a powerful and diverse set of
sequence analysis tools. Nucleic Acids Res. 32, W20-5.

Metscher, B.D., Müller, G.B., 2011. MicroCT for molecular imaging: quantitative visualization
of complete three-dimensional distributions of gene products in embryonic limbs. Dev.
Dyn. 240, 2301-2308.

Millimaki, B.B., Sweet, E.M., Dhason, M.S., Riley, B.B., 2007. Zebrafish *atoh1* genes: classic
proneural activity in the inner ear and regulation by Fgf and Notch. Development 134,
295-305.

Minařík, M., Modrell, M.S., Gillis, J.A., Campbell, A.S., Fuller, I., Lyne, R., Micklem, G., Gela,
D., Pšenička, M., Baker, C.V.H., 2023. Identification of multiple transcription factor genes
potentially involved in the development of electrosensory versus mechanosensory lateral
line organs. bioRxiv https://www.biorxiv.org/content/10.1101/2023.04.14.536701v1

Modrell, M.S., Baker, C.V.H., 2012. Evolution of electrosensory ampullary organs:
conservation of *Eya4* expression during lateral line development in jawed vertebrates.
Evol. Dev. 14, 277-285.

Modrell, M.S., Bemis, W.E., Northcutt, R.G., Davis, M.C., Baker, C.V.H., 2011a.
Electrosensory ampullary organs are derived from lateral line placodes in bony fishes.
Nat. Commun. 2, 496.

Modrell, M.S., Buckley, D., Baker, C.V.H., 2011b. Molecular analysis of neurogenic placode
development in a basal ray-finned fish. genesis 49, 278-294.

Modrell, M.S., Lyne, M., Carr, A.R., Zakon, H.H., Buckley, D., Campbell, A.S., Davis, M.C.,
Micklem, G., Baker, C.V.H., 2017a. Insights into electrosensory organ development,
physiology and evolution from a lateral line-enriched transcriptome. eLife 6, e24197.

Modrell, M.S., Tidswell, O.R.A., Baker, C.V.H., 2017b. Notch and Fgf signaling during
electrosensory versus mechanosensory lateral line organ development in a non-teleost
ray-finned fish. Dev. Biol. 431, 48-58.

Mogdans, J., 2019. Sensory ecology of the fish lateral-line system: Morphological and
physiological adaptations for the perception of hydrodynamic stimuli. J. Fish Biol. 95, 53-
72.

Moser, T., Grabner, C.P., Schmitz, F., 2020. Sensory processing at ribbon synapses in the
retina and the cochlea. Physiol. Rev. 100, 103-144.

Mukhopadhyay, M., Pangrsic, T., 2022. Synaptic transmission at the vestibular hair cells of
amniotes. Mol. Cell. Neurosci. 121, 103749.

Nicolson, T., 2017. The genetics of hair-cell function in zebrafish. J. Neurogenet. 31, 102-112.

Northcutt, R.G., 1997. Evolution of gnathostome lateral line ontogenies. Brain Behav. Evol.
50, 25-37.

Northcutt, R.G., Catania, K.C., Criley, B.B., 1994. Development of lateral line organs in the
axolotl. J. Comp. Neurol. 340, 480-514.

O'Neill, P., McCole, R.B., Baker, C.V.H., 2007. A molecular analysis of neurogenic placode
and cranial sensory ganglion development in the shark, *Scyliorhinus canicula*. Dev. Biol.
304, 156-181.

Papalopulu, N., Kintner, C., 1996. A posteriorising factor, retinoic acid, reveals that
anteroposterior patterning controls the timing of neuronal differentiation in *Xenopus*
neuroectoderm. Development 122, 3409-3418.

Pauley, S., Lai, E., Fritzsch, B., 2006. *Foxg1* is required for morphogenesis and histogenesis
of the mammalian inner ear. Dev. Dyn. 235, 2470-2482.

Pickett, S.B., Raible, D.W., 2019. Water waves to sound waves: using zebrafish to explore
hair cell biology. J. Assoc. Res. Otolaryngol. 20, 1-19.

Piotrowski, T., Baker, C.V.H., 2014. The development of lateral line placodes: Taking a
broader view. Dev. Biol. 389, 68-81.

Roccio, M., Senn, P., Heller, S., 2020. Novel insights into inner ear development and
regeneration for targeted hearing loss therapies. Hear. Res. 397, 107859.

Singh, S., Groves, A.K., 2016. The molecular basis of craniofacial placode development.
WIREs Dev. Biol. 5, 363-376.

Square, T., Romášek, M., Jandzik, D., Cattell, M.V., Klymkowsky, M., Medeiros, D.M., 2015.
CRISPR/Cas9-mediated mutagenesis in the sea lamprey *Petromyzon marinus*: a
powerful tool for understanding ancestral gene functions in vertebrates. Development
142, 4180-4187.

Square, T.A., Jandzik, D., Massey, J.L., Romášek, M., Stein, H.P., Hansen, A.W.,
Purkayastha, A., Cattell, M.V., Medeiros, D.M., 2020. Evolution of the endothelin
pathway drove neural crest cell diversification. Nature 585, 563-568.

Stundl, J., Soukup, V., Franěk, R., Pospisilova, A., Psutkova, V., Pšenička, M., Cerny, R.,
Bronner, M.E., Medeiros, D.M., Jandzik, D., 2022. Efficient CRISPR mutagenesis in
sturgeon demonstrates its utility in large, slow-maturing vertebrates. Front. Cell Dev.
Biol. 10, 750833.

Tasdemir-Yilmaz, O.E., Druckenbrod, N.R., Olukoya, O.O., Dong, W., Yung, A.R., Bastille, I.,
Pazyra-Murphy, M.F., Sitko, A.A., Hale, E.B., Vigneau, S., Gimelbrant, A.A.,
Kharchenko, P.V., Goodrich, L.V., Segal, R.A., 2021. Diversity of developing peripheral
glia revealed by single-cell RNA sequencing. Dev. Cell 56, 2516-2535.

Toresson, H., Martinez-Barbera, J.P., Bardsley, A., Caubit, X., Krauss, S., 1998. Conservation
of *BF-1* expression in amphioxus and zebrafish suggests evolutionary ancestry of
anterior cell types that contribute to the vertebrate telencephalon. Dev. Genes Evol. 208,
431-439.

Uribe-Salazar, J.M., Kaya, G., Sekar, A., Weyenberg, K., Ingamells, C., Dennis, M.Y., 2022.
Evaluation of CRISPR gene-editing tools in zebrafish. BMC Genomics 23, 12.
Webb, J.F., 2021. Morphology of the mechanosensory lateral line system of fishes, in: B.
Fritzsch (Ed.), The Senses: A Comprehensive Reference. Elsevier, pp. 29-46.
Whitear, M., 1989. Merkel cells in lower vertebrates. Arch. Histol. Cytol. 52 Suppl., 415-422.
Wu, X., Kriz, A.J., Sharp, P.A., 2014. Target specificity of the CRISPR-Cas9 system. Quant.
Biol. 2, 59-70.
York, J.R., Zehnder, K., Yuan, T., Lakiza, O., McCauley, D.W., 2019. Evolution of Snail-
mediated regulation of neural crest and placodes from an ancient role in bilaterian
neurogenesis. Dev. Biol. 453, 180-190.
Yu, H.V., Tao, L., Llamas, J., Wang, X., Nguyen, J.D., Trecek, T., Segil, N., 2021. POU4F3
pioneer activity enables ATOH1 to drive diverse mechanoreceptor differentiation through
a feed-forward epigenetic mechanism. Proc. Natl. Acad. Sci. U.S.A. 118, e2105137118.
Zhang, S., Zhang, Y., Dong, Y., Guo, L., Zhang, Z., Shao, B., Qi, J., Zhou, H., Zhu, W., Yan,
X., Hong, G., Zhang, L., Zhang, X., Tang, M., Zhao, C., Gao, X., Chai, R., 2019.
Knockdown of *Foxg1* in supporting cells increases the trans-differentiation of supporting
cells into hair cells in the neonatal mouse cochlea. Cell. Mol. Life Sci. 77, 1401-1419.
Zhang, Y., Zhang, S., Zhang, Z., Dong, Y., Ma, X., Qiang, R., Chen, Y., Gao, X., Zhao, C.,
Chen, F., He, S., Chai, R., 2020. Knockdown of *Foxg1* in Sox9+ supporting cells
increases the trans-differentiation of supporting cells into hair cells in the neonatal mouse
utricle. Aging (Albany NY) 12, 19834-19851.
Zhao, X.-F., Suh, C.S., Prat, C.R., Ellingsen, S., Fjose, A., 2009. Distinct expression of two
foxg1 paralogues in zebrafish. Gene Expr. Patterns 9, 266-272.
959

Target gene	sgRNA	Target sequence	PAM	Combinations used
Tyr	1	GGTGCCAAGGCAAAAACGCT	GGG	1+2, 1+2+3+4
	2	GATATCCCTCCATACATTAT	TGG	1+2, 1+2+3+4
	3	GATGTTTCTAAACATTGGGG	TGG	1+2+3+4
	4	GCTATGAATTTATTTTTTTC	AGG	1+2+3+4
	5	GCAAGGTATACGAAAGTTGA	CGG	5+6
	6	GATTGCAAGTTCGGCTTCTT	TGG	5+6
	7*	GGTTAGAGACTTTATGTAAC	GGG	7+8
	8*	GGCTCCATGTCTCAAGTCCA	AGG	7+8
Atoh1	1	GACCTTGTAAAAGATCGGAA	AGG	1+2
	2	GCTTGTCATTGTCAAATGAC	GGG	1+2
Neurod4	1	GGAGCGTTTCAAGGCCAGGC	GGG	1, 1+2+3+4, 1+6
	2	GTGAGCGTTCTCGCATGCAC	GGG	2, 1+2+3+4
	3	GCCTGGCCCACAACTACATC	TGG	1+2+3+4, 3+4+5+6
	4	GAGGGGCCCCGAGAAGCTGC	AGG	1+2+3+4, 3+4+5+6
	5	GTCTCCCCAGCCCTCCCTAC	GGG	5, 3+4+5+6
	6	GACAACCACTCCCCGGATTG	CGG	1+6, 3+4+5+6
	7	GACCCTGCGCAGGCTCTCCA	GGG	7+9, 7+8+9
	8	GCAGCTGGGTCCCCTGCTGA	CGG	7+8+9
	9	GGGGCCGTGTGCTCAGGGAT	GGG	7+9, 7+8+9
Foxg1	1	GAAACATCTTTTGCCCAACC	CGG	1+2
	2	TCTTCCGAGCAAGGTAACTC	GGG	1+2
	3	TGATGCTGAAGGACGACTTG	GGG	3+4
	4	CTGGCTCGTCCTCGGGCCGG	TGG	3+4

Table 1. sgRNAs used in this study. List of the genes targeted for CRISPR/Cas9-mediated mutagenesis, together with the target sequences and combinations of sgRNAs reported in this study. *Note:* Tyr sgRNAs 7 and 8 (marked with an asterisk) were designed and recently published by Stundl et al. (2022) as their *tyr* sgRNA 3 and *tyr* sgRNA 4, respectively.