Atoh1 is required for the formation of lateral line electroreceptors and hair cells, whereas Foxg1 represses an electrosensory fate

Martin Minařík[1], Alexander S. Campbell[1], Roman Franěk[2], Michaela Vazačová[2], Miloš Havelka[2], David Gela[2], Martin Pšenička[2] and Clare V. H. Baker[1]*

[1]Department of Physiology, Development & Neuroscience, University of Cambridge, Cambridge, UK

[2]Faculty of Fisheries and Protection of Waters, Research Institute of Fish Culture and Hydrobiology, University of South Bohemia in České Budějovice, Vodňany, Czech Republic

*For correspondence: cvhb1@cam.ac.uk

Abstract

In electroreceptive jawed fishes and amphibians, individual lateral line placodes form lines of neuromasts on the head containing mechanosensory hair cells, flanked by fields of ampullary organs containing electroreceptors - modified hair cells that respond to weak electric fields. Extensively shared gene expression between neuromasts and ampullary organs suggests that conserved molecular mechanisms are involved in their development, but a few transcription factor genes are restricted either to the developing electrosensory or mechanosensory lateral line. Here, we used CRISPR/Cas9-mediated mutagenesis in F0-injected sterlet embryos (*Acipenser ruthenus*, a sturgeon) to test the function of three such genes. We found that the 'hair cell' transcription factor gene *Atoh1* is required for both hair cell and electroreceptor differentiation in sterlet, and for *Pou4f3* and *Gfi1* expression in both neuromasts and ampullary organs. These data support the conservation of developmental mechanisms between hair cells and electroreceptors. Targeting ampullary organ-restricted *Neurod4* did not yield any phenotype, potentially owing to redundancy with other *Neurod* genes that we found to be expressed in sterlet ampullary organs. After targeting mechanosensory-restricted *Foxg1*, ampullary organs formed within neuromast lines, suggesting that Foxg1 normally represses their development. We speculate that electrosensory organs may be the 'default' fate of lateral line primordia in electroreceptive vertebrates.

Introduction

Mechanosensory hair cells in different compartments of the inner ear transduce fluid movements for hearing and balance (Moser et al., 2020; Caprara and Peng, 2022; Mukhopadhyay and Pangrsic, 2022). In fishes and aquatic-stage amphibians, hair cells are also found in lateral line neuromasts in the skin, which are stimulated by local water movement (Mogdans, 2019; Webb, 2021). Supporting cells in neuromasts differentiate into hair cells for homeostasis and repair (Kniss et al., 2016; Lush et al., 2019) and neuromasts are easily accessible in lines over the head and trunk, making the zebrafish lateral line an excellent model for hair cell development and regeneration (Kniss et al., 2016; Nicolson, 2017; Pickett and Raible, 2019).

The lateral line system of zebrafish (a cyprinid teleost) is purely mechanosensory, as is the lateral line system of the other main anamniote lab model, the frog *Xenopus*. However, in many other vertebrates, the lateral line system also has an electrosensory division (Bullock et al., 1983; Baker et al., 2013; Crampton, 2019). In electroreceptive non-teleost jawed vertebrates, some or all of the neuromast lines on the head are flanked by fields of ampullary organs containing electroreceptors, which respond to weak cathodal stimuli such as the electric fields surrounding other animals in water (Bodznick and Montgomery, 2005; Crampton, 2019; Leitch and Julius, 2019; Chagnaud et al., 2021). Electroreception is mediated by voltage-gated calcium channels in the apical membrane (Bodznick and Montgomery, 2005; Leitch and Julius, 2019). The voltage sensor was recently identified in cartilaginous fishes as the L-type voltage-gated calcium channel $Ca_v1.3$ (Bellono et al., 2017; Bellono et al., 2018), whose pore-forming alpha subunit is encoded by *Cacna1d*. $Ca_v1.3$ is also required for synaptic transmission at hair-cell ribbon synapses (Moser et al., 2020; Mukhopadhyay and Pangrsic, 2022).

Electroreceptors, like hair cells, have an apical primary cilium and basolateral ribbon synapses with lateral line afferents (Jørgensen, 2005; Baker, 2019). However, in contrast to the highly ordered, stepped array of apical microvilli ('stereocilia') that forms the 'hair bundle' critical for hair-cell mechanotransduction (Caprara and Peng, 2022), electroreceptors in many species (e.g., cartilaginous fishes; ray-finned paddlefishes and sturgeons) lack apical microvilli altogether (Jørgensen, 2005; Baker, 2019). Electroreceptors in other species have a few apical microvilli, while the electroreceptors of the amphibian axolotl have around 200 microvilli surrounding an eccentrically positioned primary cilium (Jørgensen, 2005; Baker, 2019). Indeed, axolotl electroreceptors were described as "remarkably similar to immature hair cells" (Northcutt et al., 1994). Thus, despite their shared function, the apical surface of electroreceptors (where voltage-sensing occurs; Bodznick and Montgomery, 2005; Leitch and Julius, 2019) varies considerably across different vertebrate groups.

71   Fate-mapping experiments have shown that neuromasts, ampullary organs (where 72 present) and their afferent neurons all develop from a series of pre-otic and post-otic lateral 73 line placodes on the embryonic head (Northcutt, 1997; Piotrowski and Baker, 2014; Baker, 74 2019). In electroreceptive jawed vertebrates, lateral line placodes elongate to form sensory 75 ridges that eventually fragment: neuromasts differentiate first, in a line along the centre of each 76 ridge, and ampullary organs (if present) form later, in fields on the flanks of the ridge (Northcutt, 77 1997; Piotrowski and Baker, 2014; Baker, 2019). The lateral line primordia of electroreceptive 78 vertebrates therefore provide a fascinating model for studying the formation of different 79 sensory cell types and organs. What molecular mechanisms control the formation within the 80 same primordium of mechanosensory neuromasts containing hair cells, versus electrosensory 81 ampullary organs containing electroreceptors?

82   To gain molecular insight into electroreceptor development, we originally took a 83 candidate-gene approach, based on genes known to be important for neuromast and/or hair 84 cell development. This enabled us to identify a variety of genes expressed in developing 85 ampullary organs as well as neuromasts, in embryos from the three major jawed-vertebrate 86 groups, i.e., cartilaginous fishes (lesser-spotted catshark, *Scyliorhinus canicula*, and little 87 skate, *Leucoraja erinacea*; O'Neill et al., 2007; Gillis et al., 2012); lobe-finned bony 88 fishes/tetrapods (a urodele amphibian, the axolotl, *Ambystoma mexicanum*; Modrell and 89 Baker, 2012); and ray-finned bony fishes (a chondrostean, the Mississippi paddlefish, 90 *Polyodon spathula*; Modrell et al., 2011a; Modrell et al., 2011b; Modrell et al., 2017b). We also 91 took an unbiased discovery approach using differential bulk RNA-seq in late-larval paddlefish, 92 which yielded a dataset of almost 500 genes that were putatively enriched in lateral line organs 93 (Modrell et al., 2017a). Validation by *in situ* hybridization of a subset of candidates from this 94 dataset suggested that conserved molecular mechanisms were involved in hair cell and 95 electroreceptor development, and that hair cells and electroreceptors were closely related 96 physiologically (Modrell et al., 2017a). For example, developing ampullary organs express the 97 key 'hair cell' transcription factor genes *Atoh1* and *Pou4f3* (see Roccio et al., 2020; Iyer and 98 Groves, 2021), and genes essential for the function of hair cell ribbon synapses, including the 99 voltage-gated calcium channel gene *Cacna1d*, encoding $Ca_v1.3$ (Modrell et al., 2017a). We 100 also identified a handful of genes expressed in developing ampullary organs but not 101 neuromasts, including two electroreceptor-specific voltage-gated potassium channel subunit 102 genes (*Kcna5* and *Kcnab3*) and a single transcription factor gene, *Neurod4* (Modrell et al., 103 2017a).

Up to that point, we had reported the shared expression of fifteen transcription factor 105 genes in both ampullary organs and neuromasts, but only one transcription factor gene with 106 restricted expression, namely, electrosensory-restricted *Neurod4* (Modrell et al., 2011a; 107 Modrell et al., 2011b; Modrell et al., 2017b). More recently (preprint: Minařík et al., 2023), we

used the late-larval paddlefish lateral line organ-enriched dataset (Modrell et al., 2017b), as
well as a candidate gene approach, to identify 23 more transcription factor genes expressed
within developing lateral line organs in paddlefish and/or in a related, more experimentally
tractable chondrostean, the sterlet (*Acipenser ruthenus*, a small sturgeon; e.g., Chen et al.,
2018; Baloch et al., 2019; Stundl et al., 2022). Twelve of these transcription factor genes -
including *Gfi1* - were expressed in both ampullary organs and neuromasts (preprint: Minařík
et al., 2023). Thus, developing ampullary organs, as well as neuromasts, express the three
'hair cell' transcription factor genes - *Atoh1*, *Pou4f3* and *Gfi1* - whose co-expression is
sufficient to drive postnatal mouse cochlear supporting cells to adopt a 'hair cell-like' fate, albeit
not to form fully mature hair cells (Roccio et al., 2020; Iyer and Groves, 2021; Chen et al.,
2021; Iyer et al., 2022). We also identified six novel ampullary organ-restricted transcription
factor genes and the first-reported mechanosensory-restricted transcription factor genes
(preprint: Minařík et al., 2023). One of the five mechanosensory-restricted transcription factor
genes was *Foxg1*, which was expressed and maintained in the central region of sensory ridges
where lines of neuromasts form, although excluded from hair cells (preprint: Minařík et al.,
2023).

Here, we used CRISPR/Cas9-mediated mutagenesis in F0-injected sterlet embryos to
investigate the function in lateral line organ development of *Atoh1*, electrosensory-restricted
Neurod4 and mechanosensory-restricted *Foxg1* (for reference, Supplementary Figure S1
shows the normal expression patterns of these genes). We report that *Atoh1* is required for
the formation of electroreceptors, as well as hair cells. We did not see any phenotype after
targeting ampullary organ-restricted *Neurod4*, potentially owing to redundancy with other
Neurod family members that we found to be expressed in sterlet ampullary organs (and
neuromasts). Targeting mechanosensory-restricted *Foxg1* resulted in a striking phenotype:
the ectopic formation of ampullary organs within neuromast lines, and in some cases the
fusion of ampullary organ fields that normally develop either side of a line of neuromasts. This
suggests the unexpected but intriguing hypothesis that ampullary organs may be the 'default'
fate for lateral line sensory ridges, and that this is repressed by Foxg1, allowing neuromasts
to form instead.

Results

CRISPR/Cas9-mediated mutagenesis in F0-injected sterlet embryos
To test gene function during lateral line organ development, we optimised CRISPR/Cas9-
mediated mutagenesis in F0-injected sterlet embryos, building on established protocols for
axolotl (*Ambystoma mexicanum*; Flowers et al., 2014; Fei et al., 2018), newt (*Pleurodeles
waltl*; Elewa et al., 2017), and sea lamprey (*Petromyzon marinus*; Square et al., 2015; York et

al., 2019; Square et al., 2020), whose eggs are all large (1-2 mm in diameter) and easy to
microinject at the 1-2 cell stage. Ovulated sterlet eggs are very large: roughly 2.5 mm in
diameter (Lenhardt et al., 2005). (Since this project started, CRISPR/Cas9-mediated
mutagenesis in F0-injected sterlet embryos has been reported, including by two of us, R.F.
and M.P.; Chen et al., 2018; Baloch et al., 2019; Stundl et al., 2022.) Analysis of microsatellite
data had originally suggested that although a whole-genome duplication had occurred in the
sterlet lineage, the sterlet was likely to be a functional diploid (Ludwig et al., 2001). Our
sgRNAs were designed before the first chromosome-level sterlet genome was published (Du
et al., 2020). Analysis of this genome showed that approximately 70% of ohnologs (i.e., gene
paralogs originating from the whole-genome duplication) had in fact been retained, suggesting
functional tetraploidy (Du et al., 2020). We comment on this in relation to our experiments at
the relevant points below.

We targeted the melanin-producing enzyme *tyrosinase* (*Tyr*) as a positive control,
using eight different single-guide (sg) RNAs. Table 1 shows the sgRNA target sequences
(including two that were designed and recently published by Stundl et al., 2022) and the
various combinations in which they were injected. Supplementary Figure S2A shows the
position of each sgRNA relative to the exon structure of the *tyrosinase* gene. Our sgRNAs
were designed before chromosome-level sterlet genomes were available (Du et al., 2020 and
the   2022   reference   genome:
https://www.ncbi.nlm.nih.gov/datasets/genome/GCF_902713425.1). Searching the reference
genome for *Tyr* showed that both *Tyr* ohnologs have been retained, on chromosomes 8 and
9, with 99.11% nucleotide identity in the coding sequence (98.81% amino acid identity). Our
sgRNAs target both *Tyr* ohnologs equally.

Three different combinations of six of the eight *Tyr* sgRNAs, when injected at the 1-2-
cell stage (2-4 sgRNAs pre-complexed with Cas9), each generated at least four embryos
(hereafter 'crispants') with altered pigmentation phenotypes evaluated at stage 45 (Dettlaff et
al., 1993), the onset of the transition to independent feeding. The other two *Tyr* sgRNAs failed
to generate any phenotypes (Supplementary Table 1). Excluding the *Tyr* sgRNAs that failed,
at least some degree of pigment loss was seen in 63/111 *Tyr* crispants (56.8%) across nine
independent batches. Examples of *Tyr* crispants with pigmentation phenotypes, plus a control,
are shown in Supplementary Figure S2B-D. The most efficient results were obtained by
injecting 1-cell embryos with a preassembled mix of Cas9 protein plus two chemically modified
sgRNAs (purchased from Synthego) against the target gene, and subsequently maintaining
the embryos at room temperature for around 6 hours. The time from fertilization to completion
of the first cleavage is around 2-3 hours at room temperature, giving plenty of time for the
Cas9/sgRNA complex to act before returning the embryos to colder temperatures for

subsequent development. (Sterlet are cold-water fish and the optimum temperature for
maintaining embryos for normal development is 16 °C.)

Following embryo injection at the 1-2-cell stage with pre-complexed sgRNAs/Cas9 and
fixation at stage 45, genomic DNA was extracted from the trunk/tail prior to analysis of the
heads by *in situ* hybridization (ISH). The sgRNA-targeted region from trunk/tail genomic DNA
was amplified by PCR for direct Sanger sequencing and *in silico* analysis using Synthego's
online 'Inference of CRISPR Edits' (ICE) tool (Conant et al., 2022) (also see e.g., Uribe-
Salazar et al., 2022) to analyse the identity and frequency of edits of the target gene. Although
our genotyping primers were designed before chromosome-level sterlet genomes were
available, comparison with the reference genome showed no mismatches against either of the
two ohnologs. Genotyping and ICE analysis (Conant et al., 2022) of tails from individual *Tyr*
crispants showed successful disruption of the *Tyr* gene (Supplementary Figure S2E-I show
examples of successful disruption of *Tyr*; the genotyping data were consistent with the primers
amplifying both ohnologs).

We note that our genotyping results have shown that most crispants analysed, across
all genes targeted, have shown some degree of targeted mutagenesis in the trunk/tail, with a
range of deletion sizes. Although phenotypes from the initial spawning seasons were almost
always highly mosaic, suggesting mutations occurred later in development, following
optimization a proportion of embryos showed complete unilateral and occasionally bilateral
phenotypes. Such phenotypes suggest that mutation occurred in one cell at the 2-cell stage
(unilateral phenotype) or even as early as the 1-cell stage (bilateral phenotype). Some degree
of mosaicism can be useful, however, as the unaffected tissue provides an internal control for
the normal expression of the gene being examined by ISH.

Targeting *Atoh1* resulted in the loss of hair cells and electroreceptors
We targeted *Atoh1* for CRISPR/Cas9-mediated mutagenesis by injecting sterlet embryos at
the 1-2 cell stage with Cas9 protein pre-complexed with two sgRNAs targeting *Atoh1* (Table
1; Supplementary Figure S2A). Our sgRNAs were designed before chromosome-level sterlet
genomes were available (Du et al., 2020 and the 2022 reference genome:
https://www.ncbi.nlm.nih.gov/datasets/genome/GCF_902713425.1/). Searching the
reference genome for *Atoh1* showed that both *Atoh1* ohnologs have been retained, on
chromosomes 1 and 2, with 91.41% nucleotide identity (and 84.02% amino acid identity) in
the coding sequence. The copy on chromosome 2 encodes a shorter version of the protein
with a four amino acid deletion near the N-terminus (E14_G17del). Our *Atoh1* sgRNA 1 (Table
1; Supplementary Figure S3A) has a one-base mismatch to the shorter *Atoh1* gene on
chromosome 2, in position 3 of the target sequence (PAM-distal), which is unlikely to prevent
successful targeting (Wu et al., 2014). Our *Atoh1* sgRNA 2 (Table 1; Supplementary Figure

S3A) has a two-base mismatch to the longer *Atoh1* gene on chromosome 1, in positions 1 and

2 of the target sequence (PAM-distal), which is also unlikely to prevent successful targeting.

Thus, we expect our sgRNAs to target both *Atoh1* ohnologs.

The *Atoh1* crispants were raised to stage 45 (the onset of independent feeding, around

14 days post-fertilization), together with *Tyr*-targeted siblings/half-siblings as controls (eggs

were fertilized *in vitro* with a mix of sperm from three different males). ISH for the hair cell and

electroreceptor marker *Cacna1d* (Modrell et al., 2017a) (also see preprint: Minařík et al.,

2023), which was recently shown to be a direct Atoh1 target gene in mouse cochlear hair cells

(Jen et al., 2022), revealed no obvious lateral line organ phenotype in *Tyr* control crispants

(Figure 1A-D; n=0/17 across four independent batches; Supplementary Table 1). Even in

wildtype larvae, the number of ampullary organs in individual fields varies considerably at

stage 45, so ampullary organ number was not in itself scored as a phenotype. However,

Cacna1d expression was absent mosaically in neuromast lines and ampullary organ fields in

Atoh1 crispants (Figure 1E-J; n=13/22, i.e., 59%, across five independent batches;

Supplementary Table 1). This suggested that disruption of the *Atoh1* gene in sterlet resulted

in the failure of hair cell differentiation (as expected from zebrafish; Millimaki et al., 2007) and

also electroreceptors. Post-ISH immunostaining for the supporting-cell marker Sox2

(Hernández et al., 2007; Modrell et al., 2017a) (also see preprint: Minařík et al., 2023)

confirmed that neuromasts had formed (Figure 1E[1],F[1]), so the phenotype was specific to

receptor cells. Ampullary organs were not reliably detected because Sox2 immunostaining

labels neuromasts much more strongly than ampullary organs in sterlet (Supplementary

Figure S1A,B; also preprint: Minařík et al., 2023), as it does in paddlefish (Modrell et al.,

2017a). ISH for electroreceptor-specific *Kcnab3* (Modrell et al., 2017a) (also see preprint:

Minařík et al., 2023) similarly showed no effect in *Tyr* control crispants (Figure 1K-N; n=0/27

across six independent batches; Supplementary Table 1), but the mosaic absence of *Kcnab3*

expression in ampullary organ fields in *Atoh1* crispants (Figure 1O-T; n=8/13, i.e., 62%, across

two independent batches; Supplementary Table 1). Post-*Kcnab3* ISH immunostaining for

Sox2 confirmed that ampullary organs were still present, as well as neuromasts (Figure

1O[1],P[1]).

In mouse cochlear hair cells, the 'hair cell' transcription factor genes *Pou4f3* and *Gfi1*

are direct Atoh1 targets (Yu et al., 2021; Jen et al., 2022). ISH for *Pou4f3* showed no

phenotype in *Tyr* controls (Figure 2A,B; n=0/9 across three batches; Supplementary Table 1)

but the mosaic absence of *Pou4f3* in ampullary organ fields and neuromast lines of *Atoh1*

crispants (Figure 2C-D[1]; n=13/15 embryos, i.e., 87%, across three batches; Supplementary

Table 1). Similarly, ISH for *Gfi1* showed no phenotype in *Tyr* controls (Figure 2E,F; n=0/17

across five batches; Supplementary Table 1), but mosaic absence of *Gfi1* in ampullary organ

fields and neuromast lines of *Atoh1* crispants (Figure 2G-H[1]; n=9/14 embryos, i.e., 64%,

Figure 1. *Atoh1* is required for the differentiation of lateral line hair cells and electroreceptors. Sterlet crispants at stage 45 after *in situ* hybridization (ISH) for the hair cell and electroreceptor marker *Cacna1d*, or the electroreceptor-specific marker *Kcnab3*. Black arrowheads indicate examples of neuromasts; white arrowheads indicate examples of ampullary organs. (**A-D**) In a control *Tyr* crispant, *Cacna1d* expression shows the normal distribution of hair cells in lines of neuromasts, and electroreceptors in fields of ampullary organs flanking the neuromast lines (lateral view: A,B; ventral view: C,D). (**E-J**) In an *Atoh1* crispant (from a different batch to the *Tyr* crispant shown in A-D), *Cacna1d* expression is absent on the left side of the head (E,F), except for a few isolated organs in the otic region and on the operculum. Post-ISH Sox2 immunostaining (E¹,F¹) shows that neuromast supporting cells are still present. A ventral view (G,H) and a lateral view of the right side of the head (I,J; image flipped horizontally for ease of comparison) reveal a unilateral phenotype, with *Cacna1d*-expressing hair cells and electroreceptors mostly absent from the left side only of the ventral rostrum (asterisk in G,H) and present on the right side of the head (I,J). (**K,L**) Lateral view of a control *Tyr* crispant after ISH for *Kcnab3*, showing the position of electroreceptors in ampullary organs. (**M,N**) Ventral view of another *Tyr* crispant showing *Kcnab3* expression in ampullary organs. (**O-P¹**) Lateral view of an *Atoh1* crispant

in which *Kcnab3* expression is absent from ampullary organs. Post-ISH Sox2 immunostaining (O[1],P[1]) shows that supporting cells are still present in neuromasts (strong staining) and can also be detected in ampullary organs (much weaker staining). (**Q-T**) A different *Atoh1* crispant after ISH for *Kcnab3*, shown in ventral view (Q,R: compare with M,N) and lateral view (S,T). *Kcnab3* expression reveals a unilateral phenotype: *Kcnab3*-expressing electroreceptors are mostly absent from the right side (asterisk) but present on the left side. Abbreviations: b, barbel; di, dorsal infraorbital ampullary organ field; e, eye; m, mouth; n, naris; S, stage; vi, ventral infraorbital ampullary organ field. Scale bars: 200 μm.

Figure 2. *Atoh1* is required for *Pou4f3* and *Gfi1* expression in ampullary organs and neuromasts. Sterlet crispants at stage 45 after *in situ* hybridization for transcription factor genes expressed by developing hair cells. Black arrowheads indicate examples of neuromasts; white arrowheads indicate examples of ampullary organs. (**A,B**) In a control *Tyr* crispant, *Pou4f3* expression is detected in both neuromasts and ampullary organs. (**C-D[1]**) In an *Atoh1* crispant, *Pou4f3* expression is absent from both neuromasts and ampullary organs, except for a few isolated organs in the postorbital region. Post-ISH Sox2 immunostaining (C[1],D[1]) shows that neuromast supporting cells are still present. Most ampullary organs are not visible as Sox2 expression in ampullary organs is significantly weaker than in neuromasts (Supplementary Figure S1A,B) and often not detectable after post-ISH immunostaining. (**E,F**) In a control *Tyr* crispant, *Gfi1* expression is detected in both neuromasts and ampullary organs. (**G-H[1]**) In an *Atoh1* crispant, *Gfi1* expression is absent from both neuromasts and ampullary organs, except for a few isolated organs in the supra- and infraorbital region. Post-ISH Sox2 immunostaining (G[1],H[1]) shows that neuromast supporting cells are still present. (Most ampullary organs are not visible due to weaker Sox2 immunostaining.) Abbreviations: b, barbel; di, dorsal infraorbital ampullary organ field; e, eye; n, naris; S, stage; vi, ventral infraorbital ampullary organ field. Scale bars: 200 μm.

across four batches; Supplementary Table 1). Post-ISH immunostaining for Sox2 confirmed that neuromasts and ampullary organs were still present in *Atoh1* crispants (Figure 2C[1],D[1],G[1],H[1]).

We performed genotyping and ICE analysis (Conant et al., 2022) of tails from individual *Atoh1* crispants targeted with this pair of sgRNAs (Supplementary Figure S3 shows examples). Our genotyping primers were designed before chromosome-level sterlet genomes

were available; comparison with the reference genome showed two mismatches in the forward

primer against the ohnolog on chromosome 2, and the genotyping data were consistent with

the primers amplifying the chromosome 1 ohnolog only. Thus, we could not determine whether

the chromosome 2 ohnolog was disrupted. However, the genotyping data showed successful

disruption of the *Atoh1* gene on chromosome 1 (Supplementary Figure S3).

Taken together, these data suggest that Atoh1 lies upstream of *Pou4f3* and *Gfi1* in

ampullary organs as well as neuromasts, and is required for the differentiation of

electroreceptors as well as hair cells.

Targeting electrosensory-restricted *Neurod4* had no obvious effect on lateral line

development

We previously identified *Neurod4* in paddlefish as the first-reported ampullary organ-restricted

transcription factor in the developing lateral line system (Modrell et al., 2017a). We confirmed

that sterlet *Neurod4* is similarly expressed by ampullary organs but not neuromasts

(Supplementary Figure S1I,J). We targeted *Neurod4* in sterlet embryos using nine different

sgRNAs (Table 1; Supplementary Figure S4A), injected in eight different combinations of 1-4

different sgRNAs across 10 independent batches of 1-2 cell-stage embryos (Supplementary

Table 1). This had no detectable effect on expression of electroreceptor-specific *Kcnab3*

(n=0/44 across nine batches, Supplementary Table 1) or the hair cell/electroreceptor marker

Cacna1d (n=0/4 across two batches, Supplementary Table 1). Examples of *Neurod4* crispants

after ISH for *Kcnab3*, plus a *Tyr* control crispant for comparison, are shown in Supplementary

Figure S4B-D. Our sgRNAs were designed before chromosome-level sterlet genomes were

available   (Du   et   al.,   2020   and   the   2022   reference   genome:

https://www.ncbi.nlm.nih.gov/datasets/genome/GCF_902713425.1/).   Searching   the

reference genome for *Neurod4* showed that both ohnologs have been retained: one on

chromosome 45 and the other annotated on an unplaced genomic scaffold, with 99.45%

nucleotide identity (and 99.46% amino acid identity) in the coding sequence. Our sgRNAs

target both *Neurod4* ohnologues without mismatches. Our genotyping primers were designed

before chromosome-level sterlet genomes were available; comparison with the reference

genome showed five mismatches in the forward primer used for genotyping larvae targeted

with sgRNAs 1 and 2 against the chromosome 45 ohnolog (Supplementary Table 1). It was

not possible to tell from our genotyping data whether the primers amplified both ohnologs or

only one, however, as the remaining sequence targeted by the primers is identical between

the two ohnologs. Genotyping and ICE analysis (Conant et al., 2022) showed successful

disruption of the *Neurod4* gene arising from six different combinations of the sgRNAs

(Supplementary Table 1; examples are shown in Supplementary Figure S4E-I). Not all sgRNA

combinations were genotyped, but all *Neurod4* crispants counted (n=44) included at least one

sgRNA confirmed to disrupt the *Neurod4* gene via genotyping of other embryos (Supplementary Table 1).

The lack of phenotype in *Neurod4* crispants, despite successful disruption of the *Neurod4* gene, suggested either that Neurod4 is not required for electroreceptor differentiation, despite its restriction to ampullary organs in both paddlefish and sterlet (Modrell et al., 2017a; this paper), or that it acts redundantly with other transcription factors. In paddlefish, *Neurod1* expression was restricted to developing lateral line ganglia (Modrell et al., 2011b). We cloned sterlet *Neurod1*, *Neurod2* and *Neurod6*. (Unlike *Neurod4*, these three *Neurod* family members are all direct Atoh1 targets in mouse hair cells; Jen et al., 2022.) These three genes all proved to be expressed in sterlet ampullary organs, as well as neuromasts (Supplementary Figure S5). Thus, it seems likely that the lack of effect of CRISPR/Cas9-mediated targeting of sterlet *Neurod4* is due to redundancy with other Neurod family transcription factors co-expressed in ampullary organs. (Our results also show there is at least some variation in *Neurod* family gene expression within the developing lateral line systems of paddlefish and sterlet.)

Targeting mechanosensory-restricted *Foxg1* led to the formation of ectopic ampullary organs within neuromast lines

We recently identified *Foxg1* as a mechanosensory lateral line-restricted transcription factor gene in paddlefish and sterlet (preprint: Minařík et al., 2023). *Foxg1* is expressed in the central zones of lateral line sensory ridges where neuromasts form, though excluded from hair cells themselves (preprint: Minařík et al., 2023). Our sgRNAs against *Foxg1* (Table 1; Supplementary Figure S6A) were designed before chromosome-level sterlet genomes were available (Du et al., 2020 and the 2022 reference genome (https://www.ncbi.nlm.nih.gov/datasets/genome/GCF_902713425.1/). Searching the reference genome for *Foxg1* showed that both *Foxg1* ohnologs have been retained, on chromosomes 15 and 18, with 96.43% nucleotide identity in the coding sequence (99.51% amino acid identity). Our sgRNAs target both *Foxg1* ohnologs without mismatches.

When compared with *Tyr* control crispants (Figure 3A-D), a striking phenotype was seen mosaically after targeting mechanosensory-restricted *Foxg1* with sgRNAs 1 and 2 (Table 1; Supplementary Figure S6A): neuromast lines were often interrupted by ectopic ampullary organs. These were defined as *Cacna1d*-expressing cells present within neuromast lines in larger clusters than expected for neuromasts, resembling ampullary organs (Figure 3E-L; n=9/18, i.e., 50%, across two independent batches; Supplementary Table 1), or by expression of electroreceptor-specific *Kcnab3* within neuromast lines, not seen in *Tyr* control crispants (Figure 3M-V; n=7/24, i.e., 29%, across both batches; Supplementary Table 1). This sometimes led to the apparent fusion of ampullary organ fields across a missing neuromast

Figure 3. *Foxg1* represses ampullary organ and electroreceptor formation within neuromast lines. Sterlet crispants at stage 45 after *in situ* hybridization (ISH) for the hair cell and electroreceptor marker *Cacna1d*, or the electroreceptor-specific marker *Kcnab3*. Black arrowheads indicate examples of neuromasts; white arrowheads indicate examples of ampullary organs. (**A,B**) Lateral view of a control *Tyr* crispant. *Cacna1d* expression shows the normal distribution of hair cells and electroreceptors. Note that ampullary organs have significantly more *Cacna1d*-expressing receptor cells than neuromasts. (**C,D**) Ventral view of a second control *Tyr* crispant. *Cacna1d* expression reveals the infraorbital neuromast line on both sides of the ventral rostrum, flanked by the dorsal infraorbital (di) and ventral infraorbital (vi) ampullary organ fields. (**E,F**) Lateral view of a *Foxg1* crispant. *Cacna1d* expression reveals that distinct neuromast lines are missing and the corresponding space is filled by putative ectopic ampullary organs, based on the large, ampullary organ-like clusters of *Cacna1d*-expressing cells. The dorsal and ventral infraorbital ampullary organ fields seem to have fused across the missing neuromast line (compare with A,B). (**G,H**) Ventral view of a second *Foxg1* crispant. *Cacna1d* expression reveals an apparent fusion of the dorsal infraorbital (di) and ventral infraorbital (vi) ampullary organ fields across the missing infraorbital neuromast lines on both sides (compare with C,D). (**I-K**) In a third

Foxg1 crispant, *Cacna1d* expression on the left side (I,J) shows that distinct supraorbital and infraorbital
neuromast lines are still present. However, some organs within the supraorbital line and most organs
within the infraorbital line have large clusters of *Cacna1d*-expressing cells, suggesting they are ectopic
ampullary organs (green arrowheads in J show examples). On the right side (K,L; image flipped
horizontally for ease of comparison), this phenotype is not seen. (**M,N**) Lateral view of a third control
Tyr crispant. Electroreceptor-specific *Kcnab3* expression shows the distribution of ampullary organs
only. (**O,P**) Ventral view of a fourth control *Tyr* crispant. *Kcnab3* expression shows the distribution of
ampullary organ fields. Note the lack of staining where the infraorbital neuromast lines run on either
side of the ventral rostrum, flanked by the dorsal infraorbital (di) and ventral infraorbital (vi) ampullary
organ fields (compare with *Cacna1d* expression in C,D). (**Q-R[1]**) Lateral view of a fourth *Foxg1* crispant.
Kcnab3 expression shows two ectopic ampullary organs (green arrowheads) within the infraorbital
neuromast line (compare with M,N). Post-ISH Sox2 immunostaining for supporting cells (Q[1],R[1])
confirms the position of the neuromast lines. (**S,T**) Ventral view of a fifth *Foxg1* crispant. On the left
side, ectopic *Kcnab3*-expressing electroreceptors fill the space where the left infraorbital neuromast line
would normally run, such that the dorsal and ventral infraorbital ampullary organ fields seem to have
fused (compare with O,P). On the right side, a single ectopic *Kcnab3*-expressing ampullary organ is
visible (green arrowhead) where the right infraorbital neuromast line runs (compare with O,P). (**U,V**)
Lateral view of a sixth *Foxg1* crispant. *Kcnab3* expression shows fusion of the dorsal infraorbital
ampullary organ field with the dorsal part of the ventral infraorbital ampullary organ field (compare with
M,N). Abbreviations: b, barbel; di, dorsal infraorbital ampullary organ field; di/vi, fused dorsal infraorbital
and ventral infraorbital ampullary organ fields; e, eye; m, mouth; n, naris; S, stage; vi, ventral infraorbital
ampullary organ field. Scale bars: 200 μm.
415

line (e.g., Figure 3E-H,S-V). Post-*Kcnab3* ISH immunostaining for Sox2, to reveal neuromasts,

also confirmed that the ectopic electroreceptors in *Foxg1* crispants were within neuromast

lines (Figure 3Q[1],R[1]). Targeting *Foxg1* with a different pair of sgRNAs (Table 1;

Supplementary Figure S6A) in a different batch of embryos generated similar phenotypes in

3/21 embryos (14%) overall (*Cacna1d*: n=1/6, i.e., 17%; *Kcnab3*: n=2/15, i.e., 13%;

Supplementary Table 1).

We also investigated this phenotype by performing ISH for two ampullary organ-

restricted transcription factor genes, *Mafa* (preprint: Minařík et al., 2023) and *Neurod4* (Modrell

et al., 2017a; this study), followed by Sox2 immunostaining in some embryos to reveal the

location of neuromasts. Relative to the normal ampullary organ expression of *Mafa* in

uninjected siblings/half-siblings (sufficient *Tyr* control crispants were not available to test;

Figure 4A-D; n=0/6 within one batch), ISH for *Mafa* showed the mosaic presence of ampullary

organs in neuromast lines and/or merging of ampullary organ fields in *Foxg1* crispants (Figure

4E-J; n=3/17 embryos i.e., 18%, across two independent batches; Supplementary Table 1).

Similarly, relative to the usual ampullary organ expression of *Neurod4* in uninjected

siblings/half-siblings (sufficient *Tyr* control crispants were not available to test; Figure 4K-N;

n=0/2 embryos within one batch; Supplementary Table 1), ISH for *Neurod4* (more weakly

expressed than *Mafa*) revealed the same ectopic ampullary organ phenotype in *Foxg1*

crispants as seen for *Cacna1d*, *Kcnab3* and *Mafa* (Figure 4O-T; n=3/6 embryos, i.e., 50%,

within one batch; Supplementary Table 1). Although our genotyping primers were designed

before chromosome-level sterlet genomes were available, comparison with the reference

437
Figure 4. Ectopic ampullary organs in *Foxg1* crispants express ampullary organ-specific
transcription factor genes *Mafa* and *Neurod4*. Sterlet crispants at stage 45 after *in situ* hybridization
(ISH) for ampullary organ-restricted transcription factor genes. Black arrowheads indicate examples of
neuromasts; white arrowheads indicate examples of ampullary organs. (**A-D**) In an uninjected
sibling/half-sibling (eggs were fertilized *in vitro* with a mix of sperm from three different males), *Mafa*
expression is restricted to ampullary organs (lateral view: A,B; ventral view: C,D). (**E-J**) A *Foxg1*
crispant. On the left side of the head (E,F), several *Mafa*-expressing ectopic ampullary organs are
present in the space where the infraorbital neuromast line would normally run, such that the dorsal and
ventral infraorbital ampullary organ fields seem to have fused (compare with A,B). Post-ISH Sox2
immunostaining (E¹,F¹) shows that neuromasts are still present both proximally and distally to the sites
of ampullary organ field fusion. In ventral view (G,H), ectopic ampullary organs (green arrowheads) are
seen bilaterally, within the spaces where the infraorbital neuromast lines run on either side of the ventral
rostrum (compare with C,D). On the right side in lateral view (I,J; image flipped horizontally for ease of
comparison), a single *Mafa*-expressing ectopic ampullary organ (green arrowhead) is also present in
the space where the infraorbital neuromast line runs (compare with A,B). (**K-N**) In an uninjected

sibling/half-sibling, *Neurod4* expression is restricted to ampullary organs (lateral view: K,L; ventral view:
M,N). (**O-T**) A *Foxg1* crispant. On the left side of the head (O,P), *Neurod4*-expressing ectopic ampullary
organs are present in the space where the infraorbital neuromast line would normally run, such that the
dorsal and ventral infraorbital ampullary organ fields seem to have fused (compare with K,L). Post-ISH
Sox2 immunostaining (O[1],P[1]) suggests that neuromasts are absent from the site of infraorbital
ampullary organ field fusion, although neuromasts can be seen in the preopercular and trunk lines
(black arrowheads, compare with O,P). In ventral view (Q,R), ectopic *Neurod4*-expressing ampullary
organs are seen where the right infraorbital neuromast line would normally run on the ventral rostrum
(green arrowhead indicates an example), resulting in partial fusion of the dorsal and ventral infraorbital
fields on this side (the left side is unaffected). On the right side in lateral view (S,T; image flipped
horizontally for ease of comparison), ectopic ampullary organs are also present in the space where the
infraorbital neuromast line runs (green arrowhead in T indicates an example), resulting in the apparent
partial fusion of the dorsal and ventral infraorbital ampullary organ fields (compare with K,L).
Abbreviations: b, barbel; di, dorsal infraorbital ampullary organ field; di/vi, fused dorsal infraorbital and
ventral infraorbital ampullary organ fields; e, eye; m, mouth; n, naris; S, stage; vi, ventral infraorbital
ampullary organ field. Scale bars: 200 μm.
469
genome showed no mismatches against either of the two ohnologs. Genotyping and ICE
analysis (Conant et al., 2022) of tails from individual *Foxg1* crispants showed successful
disruption of the *Foxg1* gene (Supplementary Figure S6 shows examples; the genotyping data
were consistent with the primers amplifying both ohnologs). Overall, these data suggest that
mechanosensory-restricted Foxg1 acts to repress the formation of ampullary organs and
electroreceptors within neuromast lines.
476
477 **Discussion**
478
Conserved molecular mechanisms underlie lateral line electroreceptor and hair cell
formation
Here, we aimed to test the function in lateral line electroreceptor and/or hair cell formation of
three transcription factor genes that we had previously identified as expressed in developing
electrosensory ampullary organ and/or mechanosensory lateral line organs in ray-finned
chondrostean fishes - paddlefish and sterlet (Butts et al., 2014; Modrell et al., 2017a; preprint:
Minařík et al., 2023). The first gene we investigated was *Atoh1*, which is required for the
formation of lateral line hair cells in zebrafish (Millimaki et al., 2007), as well as hair cells in
the inner ear (Bermingham et al., 1999; Millimaki et al., 2007). In paddlefish (Butts et al., 2014;
Modrell et al., 2017a) and sterlet (preprint: Minařík et al., 2023), *Atoh1* is expressed in
ampullary organs as well as neuromasts. Targeting *Atoh1* for CRISPR/Cas9-mediated
mutagenesis in F0 sterlet embryos showed that Atoh1 is required for the formation not only of
Cacna1d-expressing neuromast hair cells, as expected from zebrafish (Millimaki et al., 2007),
but also of *Cacna1d*-expressing, *Kcnab3*-expressing electroreceptors. These experiments
also showed that Atoh1 is required for the expression of the 'hair cell' transcription factor genes
Gfi1 and *Pou4f3* in developing ampullary organs, as well as neuromasts. This is consistent

with both of these genes being direct Atoh1 targets in mouse cochlear hair cells (Yu et al.,
2021; Jen et al., 2022).

In both inner-ear hair cells and Merkel cells (epidermal mechanoreceptors found in all
vertebrates; see e.g., Whitear, 1989; Brown et al., 2023), Atoh1 acts with Pou4f3 in a
conserved 'feed-forward circuit', with Atoh1 directly activating *Pou4f3* expression, and Pou4f3
then acting as a pioneer factor to open a significant subset of Atoh1 target enhancers (some
shared and some divergent between hair cells and Merkel cells), enabling mechanosensory
differentiation (Yu et al., 2021). In hair cells, *Gfi1* is one of the Pou4f3-dependent Atoh1 targets
(Yu et al., 2021). Together with the striking conservation of transcription factor gene
expression between developing ampullary organs and neuromasts (Modrell et al., 2011b;
Modrell et al., 2011a; Modrell et al., 2017b; Modrell et al., 2017a) (also preprint: Minařík et al.,
2023), the phenotypes seen in *Atoh1*-targeted F0 sterlet crispant embryos suggest that the
molecular mechanisms underlying electroreceptor formation are highly conserved with those
underlying hair cell formation. Indeed, the requirement of Atoh1 for *Pou4f3* and *Gfi1*
expression in ampullary organs, as well as neuromasts, suggests that the Atoh1-Pou4f3
'feedforward circuit' in mechanosensory cells - i.e., hair cells and epidermal Merkel cells (Yu
et al., 2021) - may also be conserved, at least partly, in developing electroreceptors. Taken
together, these data support the hypothesis that electroreceptors evolved as a transcriptionally
related "sister cell type" to lateral line hair cells (Arendt et al., 2016; Baker and Modrell, 2018;
Baker, 2019).

Electrosensory-restricted Neurod4 is likely redundant with other Neurod family
members in sterlet
Paddlefish *Neurod4* was the first-reported ampullary organ-restricted transcription factor gene
(Modrell et al., 2017a), with conserved expression in sterlet (this study). We were unable to
detect a lateral line organ phenotype in *Neurod4*-targeted sterlet crispants. However, we found
that *Neurod1*, *Neurod2* and *Neurod6* are all expressed in sterlet ampullary organs (as well as
neuromasts), suggesting that Neurod4 may act redundantly with one or more of these factors
in developing ampullary organs. (In paddlefish, however, *Neurod1* expression is restricted to
developing lateral line ganglia; Modrell et al., 2011b.) Targeting multiple *Neurod* genes for
CRISPR/Cas9-mediated mutagenesis in the future may shed light on the role played by this
transcription factor family in ampullary organ development.

Foxg1 represses electroreceptor formation in the neuromast-forming central zone of
lateral line sensory ridges
We also targeted *Foxg1*, a mechanosensory-restricted transcription factor gene that we
recently identified in the developing lateral line system of paddlefish and sterlet (preprint:

Minařík et al., 2023). *Foxg1* is expressed in the central zones of lateral line sensory ridges
where neuromasts form, though excluded from the central domains of neuromasts where hair
cells differentiate (preprint: Minařík et al., 2023). Targeting *Foxg1* for CRISPR/Cas9-mediated
mutagenesis in F0 sterlet embryos led to a striking phenotype: the formation within neuromast
lines of ectopic electroreceptors, often in the large clusters normally seen in ampullary organs.
In some cases, ampullary organ fields, which normally flank neuromast lines, effectively
'merged' across missing neuromast lines. This phenotype was revealed by examining
expression of the electroreceptor-specific marker *Kcnab3*, and two ampullary organ-restricted
transcription factor genes: *Mafa* (preprint: Minařík et al., 2023) and *Neurod4* (Modrell et al.,
2017a). Thus, *Foxg1* seems to repress an ampullary organ fate within the central domain of
lateral line sensory ridges where neuromasts form.

In the mouse inner ear, *Foxg1* is expressed in the prospective cochlea and all sensory
patches, in hair cell progenitors and supporting cells (Pauley et al., 2006; Tasdemir-Yilmaz et
al., 2021), plus a subset of hair cells (Pauley et al., 2006). Knockout leads to a shorter cochlea
with extra rows of hair cells, and to loss or reduction of vestibular end organs (Pauley et al.,
2006; Hwang et al., 2009). More hair cells and fewer supporting cells were seen after
conditional knockdown of *Foxg1* in neonatal cochlear supporting cells, possibly via the
transdifferentiation of supporting cells (Zhang et al., 2019; Zhang et al., 2020). This suggests
the possibility that Foxg1 may act in the central zone of lateral line sensory ridges to maintain
a proliferative progenitor state, as it does in the mouse olfactory epithelium (Kawauchi et al.,
2009).

Furthermore, Fox family members can act as pioneer factors as well as transcription
factors (Golson and Kaestner, 2016; Lukoseviciute et al., 2018). A pioneer factor role has
been proposed for Foxi3 in otic placode development (see Singh and Groves, 2016). In the
developing neural crest, Foxd3 acts early as a pioneer factor, opening enhancers and
repositioning nucleosomes to prime genes controlling neural crest specification and migration
and, concurrently, to repress the premature differentiation of, e.g., melanocytes (Lukoseviciute
et al., 2018). Later in neural crest development, Foxd3 represses enhancers associated with
mesenchymal, neuronal and melanocyte lineages (Lukoseviciute et al., 2018). In cortical
progenitors, Foxg1 suppresses the adoption at later stages of an early-born cell fate, namely,
Cajal-Retzius cells (Hanashima et al., 2004). In the developing chicken otic placode, Foxg1
represses markers of other lineages, such as the olfactory and lens placodes, and epidermis
(Anwar et al., 2017). Hence, it is possible that Foxg1 acts in the central zone of lateral line
sensory ridges in electroreceptive fishes as a pioneer factor for neuromast/hair cell formation,
and/or that it represses ampullary organ/electroreceptor formation. Such a role would be
consistent with lateral line expression of *Foxg1* only in electroreceptive fishes: *Foxg1* is not
expressed in the developing lateral line of zebrafish or *Xenopus* (e.g., Dirksen and Jamrich,

1995; Papalopulu and Kintner, 1996; Toresson et al., 1998; Eagleson and Dempewolf, 2002;
Duggan et al., 2008; Zhao et al., 2009).

Overall, these data lead us to propose the speculative hypothesis that electrosensory
organs may be the 'default' fate within lateral line sensory ridges in electroreceptive
vertebrates, and that Foxg1 represses this fate to enable mechanosensory neuromasts and
hair cells to form. To test these hypotheses directly, it will be important in the future to identify
global changes in gene expression and chromatin accessibility in the absence of Foxg1.

Summary and Perspective

Overall, we have found that the 'hair cell' transcription factor Atoh1 is required for the formation
of lateral line electroreceptors as well as hair cells, consistent with a close developmental
relationship between these putative 'sister cell' types. Electrosensory-restricted Neurod4 may
act redundantly with other Neurod family members expressed in developing ampullary organs.
Mechanosensory-restricted Foxg1 represses the formation of electroreceptors within
neuromast lines, suggesting the surprising possibility that electroreceptors are the 'default'
fate within lateral line sensory ridges and raising interesting developmental and evolutionary
questions for future investigation.

<u>Materials and Methods</u>

Animals

Fertilized sterlet (*Acipenser ruthenus*) eggs were obtained from the breeding facility at the
Research Institute of Fish Culture and Hydrobiology, Faculty of Fisheries and Protection of
Waters, University of South Bohemia in České Budějovice, Vodňany, Czech Republic, and
staged according to Dettlaff et al. (1993). For detailed information about sterlet husbandry, *in*
vitro fertilization and the rearing of embryos and yolk-sac larvae, see Stundl et al. (2022). Each
fertilization used a mix of sperm from three different males, so each batch was a mix of siblings
and half-siblings. Upon reaching the desired developmental stages, embryos and yolk-sac
larvae were euthanized by overdose of MS-222 (Sigma-Aldrich) and fixed in modified Carnoy's
fixative (6 volumes 100% ethanol: 3 volumes 37% formaldehyde: 1 volume glacial acetic acid)
for 3 hours at room temperature, dehydrated stepwise into 100% ethanol and stored at -20
°C.

All experimental procedures were approved by the Animal Research Committee of the
Faculty of Fisheries and Protection of Waters in Vodňany, University of South Bohemia in
České Budějovice, Czech Republic, and by the Ministry of Agriculture of the Czech Republic
(reference number: MSMT-12550/2016-3). Experimental fish were maintained according to
the principles of the European Union (EU) Harmonized Animal Welfare Act of the Czech

Republic, and Principles of Laboratory Animal Care and National Laws 246/1992 "Animal
Welfare" on the protection of animals.

CRISPR guide RNA design and synthesis

Target gene sequences were identified using the National Center for Biotechnology
Information (NCBI) Basic Local Alignment Search Tool BLAST
(https://blast.ncbi.nlm.nih.gov/Blast.cgi; McGinnis and Madden, 2004) to search sterlet
transcriptomic data (available at DDBJ/EMBL/GenBank under the accessions
GKLU00000000 and GKEF01000000; see preprint: Minařík et al., 2023) or draft genomic
sequence data (M.H., unpublished) with the relevant paddlefish sequence (Modrell et al.,
2017a). Transcriptomic sequence data were searched for *Tyr*, *Atoh1* and *Neurod4*; genomic
sequence data were searched for *Foxg1*. Chromosome-level sterlet genomes became
available only after the project started: Du et al. (2020) and the 2022 reference genome
(https://www.ncbi.nlm.nih.gov/datasets/genome/GCF_902713425.1/). Open reading frames
were identified using the NCBI ORF Finder tool (https://www.ncbi.nlm.nih.gov/orffinder/) and
exons annotated by comparison with reference anamniote species (*Lepisosteus oculatus*,
Danio rerio, *Xenopus tropicalis*) available via Ensembl (https://www.ensembl.org;
Cunningham et al., 2022). Conserved domains were identified using NCBI BLASTX
(https://blast.ncbi.nlm.nih.gov/Blast.cgi; McGinnis and Madden, 2004). gRNAs were
preferentially designed to target 5' exons, ideally upstream of or within regions encoding
known functional domains, to increase the probability of disrupting gene function. gRNAs were
designed using the CRISPR Guide RNA Design Tool from Benchling (https://benchling.com)
and selected for synthesis based on the following criteria: (1) a high on-target score, ideally
no less than 0.5; (2) no off-target matches identified within coding sequences in transcriptome
and genome data, unless there were at least two mismatches in the 3' seed sequence (8-10
bp upstream of the protospacer adjacent motif [PAM], or in the PAM itself); (3) if multiplexing,
the gRNA pair were ideally within 50-150 bases of each other to increase the probability of
fragment deletion.

DNA templates for CRISPR gRNAs were synthesized using plasmid pX335-U6-
Chimeric_BB-CBh-hSpCas9n(D10A) (Addgene, plasmid #42335; Cong et al., 2013)
containing the gRNA scaffold. The gRNA scaffold was amplified using a specific forward
primer for each gRNA, with an overhang containing the gRNA target sequence and T7
promoter, and a reverse primer that was identical for all reactions. For gRNAs that did not start
with G, an additional G was added at the start to ensure efficient transcription. The DNA
template was amplified using Q5 polymerase (New England Biolabs, NEB) and purified using
the Monarch PCR & DNA Cleanup Kit (NEB). The gRNAs were synthesized using the HiScribe
T7 High Yield RNA Synthesis Kit (NEB) and purified using the Monarch RNA Cleanup Kit

(NEB) and stored at -80 °C before use. Alternatively, chemically modified synthetic gRNAs
were ordered directly from Synthego (CRISPRevolution sgRNA EZ Kit).

Embryo injection

On the day of injection, 1200 ng gRNA were mixed with 2400 ng Cas9 protein with NLS (PNA
Bio) in 5 µl nuclease-free water and incubated for 10 minutes at room temperature to form
ribonucleoprotein (RNP) complexes. For gRNA multiplexing, two RNP mixes were combined
1:1 to a final volume of 5 µl, and 0.5 µl of 10% 10,000 MW rhodamine dextran (Invitrogen)
added to better visualize the injection mixture and allow selection of properly injected embryos
using rhodamine fluorescence. Injection mixtures were kept on ice throughout the injection
session. 10 µl glass microcapillaries (Drummond Microcaps) were pulled in a capillary needle
puller (PC-10, Narishige) set to 58 °C with two light and one heavy weights, in single-stage
pulling mode. Fertilized sterlet eggs were manually dechorionated using Dumont #5 forceps.
A 1000 µl pipette tip cut to the same diameter as a dechorionated sterlet egg was used to
prepare a series of wells in an agar plate to allow ideal egg positioning during injection using
an automatic microinjector (FemtoJet 4x, Eppendorf), set to 100 hPa. Approximately 20 nl of
the injection mixture (corresponding to approximately 4.8 ng gRNAs and 9.6 ng Cas9) were
injected into fertilized eggs or two-cell stage embryos, targeting the animal pole at a 45° angle.
Injected embryos were moved to a clean Petri dish and, for optimum Cas9 efficiency, kept at
room temperature until the end of the injecting session or until at least the 32-cell stage, then
moved to a 16°C incubator. No more than 30 eggs were kept per 90 mm Petri dish. Unfertilized
eggs and dead embryos were removed at the end of the injection day. Petri dishes were
checked regularly for dead embryos and the water was changed at least twice a day before
gastrulation was completed, and once daily post-gastrulation. Hatched larvae were kept for
approximately 16 days post fertilisation until stage 45, then euthanized by MS-222 overdose
and fixed with modified Carnoy's fixative (see above). Fixed larvae were then dehydrated
stepwise into 100% ethanol and stored at -20°C.

Gene cloning, *in situ* hybridization and immunohistochemistry

Total RNA was isolated from sterlet embryos using Trizol (Invitrogen, Thermo Fisher
Scientific), following the manufacturer's protocol, and cDNA synthesized using the Superscript
III First Strand Synthesis kit (Invitrogen, Thermo Fisher Scientific). We used our sterlet
transcriptome assemblies (from pooled yolk-sac larvae at stages 40-45; preprint: Minařík et
al., 2023), which are available at DDBJ/EMBL/GenBank under the accessions
GKLU00000000 and GKEF01000000, to design primers for *Neurod4* (forward:
GAGAGAGCCCCAAAGAGACGAG; reverse: CTGCTTGAGCGAGAAGTTGACG). cDNA
fragments amplified under standard PCR conditions were cloned into the pDrive cloning vector

(Qiagen). Individual clones were verified by sequencing (Department of Biochemistry Sequencing Facility, University of Cambridge, UK) and sequence identity verified using NCBI BLAST (https://blast.ncbi.nlm.nih.gov/Blast.cgi; McGinnis and Madden, 2004). For *Neurod1*, *Neurod2*, *Neurod4* and *Neurod6*, synthetic gene fragments based on sterlet transcriptome data, with added M13 forward and reverse primer adaptors, were ordered from Twist Bioscience. GenBank accession numbers are as follows: *Neurod1* OQ808944, *Neurod2* OQ808945, *Neurod4* OQ808946, *Neurod6* OQ808947. The other genes used in this study have been published (preprint: Minařík et al., 2023).

The sterlet riboprobe template sequences were designed before chromosome-level genome assemblies for sterlet were available (Du et al., 2020 and the 2022 reference genome: https://www.ncbi.nlm.nih.gov/datasets/genome/GCF_902713425.1/). Genome analysis showed that an independent whole-genome duplication occurred in the sterlet lineage, from which approximately 70% of ohnologs (i.e., gene paralogs arising from the whole-genome duplication) have been retained (Du et al., 2020). Supplementary File 1 shows the percentage match for each *Neurod* family riboprobe with the two ohnologs, obtained by performing a nucleotide BLAST search against the reference genome (https://www.ncbi.nlm.nih.gov/datasets/genome/GCF_902713425.1/). Equivalent data for the other riboprobes used in this study are available in the preprint Minařík et al. (2023). The percentage match with the 'targeted' *Neurod* family ohnolog ranged from 98.9-100%; the percentage match with the second ohnolog was also high, ranging from 96.2-100% (Supplementary File 1), suggesting that transcripts from the second ohnolog are likely to be targeted by each of these riboprobes.

Digoxigenin-labelled riboprobes were synthesized as previously described (preprint: Minařík et al., 2023). Whole-mount *in situ* hybridization (ISH) was performed as previously described (Modrell et al., 2011a). Whole-mount immunostaining post-ISH for Sox2 (rabbit monoclonal, 1:200; ab92494, Abcam) was performed as previously described (Metscher and Müller, 2011), using a horseradish peroxidase-conjugated goat anti-rabbit antibody (1:300, Jackson ImmunoResearch) and the metallographic peroxidase substrate EnzMet kit (Nanoprobes) as per the manufacturer's instructions.

Genotyping

To confirm successful mutation in targeted regions, genotyping was performed on trunk and tail tissue that had been removed before ISH and stored in 100% ethanol at -20°C. The tissue was digested using Rapid Extract Lysis Kit (PCR Biosystems) and the target region amplified using HS Taq Mix Red (PCR Biosystems) according to the manufacturer's instructions. Primers used for genotyping are listed in Supplementary Table 1. After agarose gel electrophoresis, PCR products were extracted using MinElute Gel Extraction Kit (Qiagen) and

submitted for sequencing (Genewiz by Azenta Life Sciences). To analyze CRISPR editing efficiency, Sanger trace files were uploaded to the Synthego Inference of CRISPR Edits (ICE) tool (https://ice.synthego.com; Conant et al., 2022).

Imaging and image processing

Larvae were placed in a slit in an agar-coated Petri dish with PBS and imaged using a Leica MZFLIII dissecting microscope equipped with either a MicroPublisher 5.0 RTV camera (QImaging) controlled by QCapture Pro 6.0 or 7.0 software (QImaging), or a MicroPublisher 6 color CCD camera (Teledyne Photometrics) controlled by Ocular software (Teledyne Photometrics). For most larvae, a stack of images was acquired by manually focusing through the sample, then Helicon Focus software (Helicon Soft Limited) was used for focus stacking. Adobe Photoshop (Adobe Systems Inc.) was used to process images.

Data availability statement

The publication and associated supplementary figures include representative example images of embryos from each experiment. Additional raw data underlying this publication consist of further images of these and other embryos from each experiment. Public sharing of these images is not cost-efficient but they are available from the corresponding author upon reasonable request.

Ethics statement

Sterlet animal work was reviewed and approved by The Animal Research Committee of Research Institute of Fish Culture and Hydrobiology, Faculty of Fisheries and Protection of Waters, University of South Bohemia in České Budějovice, Vodňany, Czech Republic and Ministry of Agriculture of the Czech Republic (MSMT-12550/2016-3).

Author contributions

CB conceived and designed the project and wrote the manuscript together with MM. MM performed most of the experiments, prepared all the manuscript figures and made a significant contribution to the writing of the manuscript. AC and RF performed some experiments. MH provided draft genome sequence data. MP and DG were instrumental in enabling all work with sterlet embryos. All authors read and commented on the manuscript.

Funding

This work was supported by the Biotechnology and Biological Sciences Research Council (BBSRC: grant BB/P001947/1 to CB). Additional support for MM was provided by the Cambridge Isaac Newton Trust (grant 20.07[c] to CB) and by the School of the Biological

754 Sciences, University of Cambridge. AC was supported by a PhD research studentship from
755 the Anatomical Society. The work of RF and MP was supported by the Ministry of Education,
756 Youth and Sports of the Czech Republic, projects CENAKVA (LM2018099), Biodiversity
757 (CZ.02.1.01/0.0/0.0/16_025/0007370) and Czech Science Foundation (20-23836S).
758

Rights Retention Statement

This work was funded by a grant from the Biotechnology and Biological Sciences Research
Council (BBSRC: BB/P001947/1). For the purpose of open access, the author has applied a
Creative Commons Attribution (CC BY) licence to any Author Accepted Manuscript version
arising.

Acknowledgements

Thanks to Melinda Modrell for pilot work on CRISPR/Cas9 in paddlefish and for her advice
and helpful discussions in planning the project. Thanks to Marek Rodina and Martin Kahanec
for their help with sterlet spawns. Thanks to David Jandzik, András Simon, Alberto Joven
Araus, Ahmed Elewa, and Mustafa Khokha for invaluable advice on CRISPR/Cas9
approaches and genotyping. Thanks to Jan Stundl and David Jandzik for sharing their sterlet
Tyr gRNA sequences prior to publication. Thanks to Christine Hirschberger and Rolf Ericsson
for their help with some of the *in situ* hybridization rounds.

References

Anwar, M., Tambalo, M., Ranganathan, R., Grocott, T., Streit, A., 2017. A gene network
regulated by FGF signalling during ear development. Sci. Rep. 7, 6162.
Arendt, D., Musser, J.M., Baker, C.V.H., Bergman, A., Cepko, C., Erwin, D.H., Pavlicev, M.,
Schlosser, G., Widder, S., Laubichler, M.D., Wagner, G.P., 2016. The origin and
evolution of cell types. Nat. Rev. Genet. 17, 744-757.
Baker, C.V.H., 2019. The development and evolution of lateral line electroreceptors: insights
from comparative molecular approaches., in: B.A. Carlson, J.A. Sisneros, A.N. Popper,
R.R. Fay (Eds.), Electroreception: Fundamental Insights from Comparative
Approaches. Springer, Cham, pp. 25-62.
Baker, C.V.H., Modrell, M.S., 2018. Insights into electroreceptor development and evolution
from molecular comparisons with hair cells. Integr. Comp. Biol. 58, 329-340.
Baker, C.V.H., Modrell, M.S., Gillis, J.A., 2013. The evolution and development of vertebrate
lateral line electroreceptors. J. Exp. Biol. 216, 2515-2522.
Baloch, A.R., Franěk, R., Tichopád, T., Fučíková, M., Rodina, M., Pšenička, M., 2019. *Dnd1*
knockout in sturgeons by CRISPR/Cas9 generates germ cell free host for surrogate
production. Animals (Basel) 9, 174.

Bellono, N.W., Leitch, D.B., Julius, D., 2017. Molecular basis of ancestral vertebrate
792   electroreception. Nature 543, 391-396.

793   Bellono, N.W., Leitch, D.B., Julius, D., 2018. Molecular tuning of electroreception in sharks
794   and skates. Nature 558, 122-126.

795   Bermingham, N.A., Hassan, B.A., Price, S.D., Vollrath, M.A., Ben-Arie, N., Eatock, R.A.,
796   Bellen, H.J., Lysakowski, A., Zoghbi, H.Y., 1999. *Math1*: an essential gene for the
797   generation of inner ear hair cells. Science 284, 1837-1841.

798   Bodznick, D., Montgomery, J.C., 2005. The physiology of low-frequency electrosensory
799   systems, in: T.H. Bullock, C.D. Hopkins, A.N. Popper, R.R. Fay (Eds.),
800   Electroreception. Springer, New York, pp. 132-153.

801   Brown, T.L., Horton, E.C., Craig, E.W., Goo, C.E.A., Black, E.C., Hewitt, M.N., Yee, N.G.,
802   Fan, E.T., Raible, D.W., Rasmussen, J.P., 2023. Dermal appendage-dependent
803   patterning of zebrafish *atoh1a+* Merkel cells. eLife 12, e85800.

804   Bullock, T.H., Bodznick, D.A., Northcutt, R.G., 1983. The phylogenetic distribution of
805   electroreception: evidence for convergent evolution of a primitive vertebrate sense
806   modality. Brain Res. Rev. 287, 25-46.

807   Butts, T., Modrell, M.S., Baker, C.V.H., Wingate, R.J.T., 2014. The evolution of the
808   vertebrate cerebellum: absence of a proliferative external granule layer in a non-teleost
809   ray-finned fish. Evol. Dev. 16, 92-100.

810   Caprara, G.A., Peng, A.W., 2022. Mechanotransduction in mammalian sensory hair cells.
811   Mol. Cell. Neurosci. 120, 103706.

812   Chagnaud, B.P., Wilkens, L.A., Hofmann, M., 2021. The ampullary electrosensory system –a
813   paddlefish case study, in: B. Fritzsch (Ed.), The Senses: A Comprehensive Reference.
814   Elsevier, pp. 215-227.

815   Chen, J., Wang, W., Tian, Z., Dong, Y., Dong, T., Zhu, H., Zhu, Z., Hu, H., Hu, W., 2018.
816   Efficient gene transfer and gene editing in sterlet. Front. Genet. 9, 117.

817   Chen, Y., Gu, Y., Li, Y., Li, G.L., Chai, R., Li, W., Li, H., 2021. Generation of mature and
818   functional hair cells by co-expression of *Gfi1*, *Pou4f3*, and *Atoh1* in the postnatal
819   mouse cochlea. Cell Rep. 35, 109016.

820   Conant, D., Hsiau, T., Rossi, N., Oki, J., Maures, T., Waite, K., Yang, J., Joshi, S., Kelso, R.,
821   Holden, K., Enzmann, B.L., Stoner, R., 2022. Inference of CRISPR edits from Sanger
822   trace data. CRISPR J. 5, 123-130.

823   Cong, L., Ran, F.A., Cox, D., Lin, S., Barretto, R., Habib, N., Hsu, P.D., Wu, X., Jiang, W.,
824   Marraffini, L.A., Zhang, F., 2013. Multiplex genome engineering using CRISPR/Cas
825   systems. Science 339, 819-823.

826   Crampton, W.G.R., 2019. Electroreception, electrogenesis and electric signal evolution. J.
827   Fish Biol. 95, 92-134.

Cunningham, F., Allen, J.E., Allen, J., Alvarez-Jarreta, J., Amode, M.R., Armean, I.M.,
Austine-Orimoloye, O., Azov, A.G., Barnes, I., Bennett, R., Berry, A., Bhai, J., Bignell,
A., Billis, K., Boddu, S., Brooks, L., Charkhchi, M., Cummins, C., Da Rin Fioretto, L.,
Davidson, C., Dodiya, K., Donaldson, S., El Houdaigui, B., El Naboulsi, T., Fatima, R.,
Giron, C.G., Genez, T., Martinez, J.G., Guijarro-Clarke, C., Gymer, A., Hardy, M.,
Hollis, Z., Hourlier, T., Hunt, T., Juettemann, T., Kaikala, V., Kay, M., Lavidas, I., Le, T.,
Lemos, D., Marugán, J.C., Mohanan, S., Mushtaq, A., Naven, M., Ogeh, D.N., Parker,
A., Parton, A., Perry, M., Piližota, I., Prosovetskaia, I., Sakthivel, M.P., Salam, A.I.A.,
Schmitt, B.M., Schuilenburg, H., Sheppard, D., Pérez-Silva, J.G., Stark, W., Steed, E.,
Sutinen, K., Sukumaran, R., Sumathipala, D., Suner, M.M., Szpak, M., Thormann, A.,
Tricomi, F.F., Urbina-Gómez, D., Veidenberg, A., Walsh, T.A., Walts, B., Willhoft, N.,
Winterbottom, A., Wass, E., Chakiachvili, M., Flint, B., Frankish, A., Giorgetti, S.,
Haggerty, L., Hunt, S.E., IIsley, G.R., Loveland, J.E., Martin, F.J., Moore, B., Mudge,
J.M., Muffato, M., Perry, E., Ruffier, M., Tate, J., Thybert, D., Trevanion, S.J., Dyer, S.,
Harrison, P.W., Howe, K.L., Yates, A.D., Zerbino, D.R., Flicek, P., 2022. Ensembl
2022. Nucleic Acids Res. 50, D988-D995.
Dettlaff, T.A., Ginsburg, A.S., Schmalhausen, O.I., 1993. Sturgeon Fishes: Developmental
Biology and Aquaculture. Springer-Verlag, Berlin.
Dirksen, M.-L., Jamrich, M., 1995. Differential expression of *fork head* genes during early
Xenopus and zebrafish development. Dev. Genet. 17, 107-116.
Du, K., Stöck, M., Kneitz, S., Klopp, C., Woltering, J.M., Adolfi, M.C., Feron, R., Prokopov,
D., Makunin, A., Kichigin, I., Schmidt, C., Fischer, P., Kuhl, H., Wuertz, S., Gessner, J.,
Kloas, W., Cabau, C., Iampietro, C., Parrinello, H., Tomlinson, C., Journot, L.,
Postlethwait, J.H., Braasch, I., Trifonov, V., Warren, W.C., Meyer, A., Guiguen, Y.,
Schartl, M., 2020. The sterlet sturgeon genome sequence and the mechanisms of
segmental rediploidization. Nat. Ecol. Evol. 4, 841-852.
Duggan, C.D., Demaria, S., Baudhuin, A., Stafford, D., Ngai, J., 2008. Foxg1 is required for
development of the vertebrate olfactory system. J. Neurosci. 28, 5229-5239.
Eagleson, G.W., Dempewolf, R.D., 2002. The role of the anterior neural ridge and *Fgf-8* in
early forebrain patterning and regionalization in *Xenopus laevis*. Comp. Biochem.
Physiol. B Biochem. Mol. Biol. 132, 179-189.
Elewa, A., Wang, H., Talavera-López, C., Joven, A., Brito, G., Kumar, A., Hameed, L.S.,
Penrad-Mobayed, M., Yao, Z., Zamani, N., Abbas, Y., Abdullayev, I., Sandberg, R.,
Grabherr, M., Andersson, B., Simon, A., 2017. Reading and editing the *Pleurodeles*
waltl genome reveals novel features of tetrapod regeneration. Nat. Commun. 8, 2286.
Fei, J.-F., Lou, W.P.-K., Knapp, D., Murawala, P., Gerber, T., Taniguchi, Y., Nowoshilow, S.,
Khattak, S., Tanaka, E.M., 2018. Application and optimization of CRISPR-Cas9-

mediated genome engineering in axolotl (*Ambystoma mexicanum*). Nat. Protoc. 13,

2908-2943.

Flowers, G.P., Timberlake, A.T., McLean, K.C., Monaghan, J.R., Crews, C.M., 2014. Highly

efficient targeted mutagenesis in axolotl using Cas9 RNA-guided nuclease.

Development 141, 2165-2171.

Gillis, J.A., Modrell, M.S., Northcutt, R.G., Catania, K.C., Luer, C.A., Baker, C.V.H., 2012.

Electrosensory ampullary organs are derived from lateral line placodes in cartilaginous

fishes. Development 139, 3142-3146.

Golson, M.L., Kaestner, K.H., 2016. Fox transcription factors: from development to disease.

Development 143, 4558-4570.

Hanashima, C., Li, S.C., Shen, L., Lai, E., Fishell, G., 2004. *Foxg1* suppresses early cortical

cell fate. Science 303, 56-59.

Hernández, P.P., Olivari, F.A., Sarrazin, A.F., Sandoval, P.C., Allende, M.L., 2007.

Regeneration in zebrafish lateral line neuromasts: expression of the neural progenitor

cell marker sox2 and proliferation-dependent and-independent mechanisms of hair cell

renewal. Dev. Neurobiol. 67, 637-654.

Hwang, C.H., Simeone, A., Lai, E., Wu, D.K., 2009. *Foxg1* is required for proper separation

and formation of sensory cristae during inner ear development. Dev. Dyn. 238, 2725-

2734.

Iyer, A.A., Groves, A.K., 2021. Transcription factor reprogramming in the inner ear: turning

on cell fate switches to regenerate sensory hair cells. Front. Cell. Neurosci. 15,

660748.

Iyer, A.A., Hosamani, I., Nguyen, J.D., Cai, T., Singh, S., McGovern, M.M., Beyer, L., Zhang,

H., Jen, H.I., Yousaf, R., Birol, O., Sun, J.J., Ray, R.S., Raphael, Y., Segil, N., Groves,

A.K., 2022. Cellular reprogramming with ATOH1, GFI1, and POU4F3 implicate

epigenetic changes and cell-cell signaling as obstacles to hair cell regeneration in

mature mammals. eLife 11, e79712.

Jen, H.-I., Singh, S., Tao, L., Maunsell, H.R., Segil, N., Groves, A.K., 2022. GFI1 regulates

hair cell differentiation by acting as an off-DNA transcriptional co-activator of ATOH1,

and a DNA-binding repressor. Sci. Rep. 12, 7793.

Jørgensen, J.M., 2005. Morphology of electroreceptive sensory organs, in: T.H. Bullock,

C.D. Hopkins, A.N. Popper, R.R. Fay (Eds.), Electroreception. Springer, New York, pp.

47-67.

Kawauchi, S., Santos, R., Kim, J., Hollenbeck, P.L.W., Murray, R.C., Calof, A.L., 2009. The

role of *Foxg1* in the development of neural stem cells of the olfactory epithelium. Ann.

N. Y. Acad. Sci. 1170, 21-27.

Kniss, J.S., Jiang, L., Piotrowski, T., 2016. Insights into sensory hair cell regeneration from
the zebrafish lateral line. Curr. Opin. Genet. Dev. 40, 32-40.

Leitch, D.B., Julius, D., 2019. Electrosensory transduction: comparisons across structure,
afferent response properties, and cellular physiology., in: B.A. Carlson, J.A. Sisneros,
A.N. Popper, R.R. Fay (Eds.), Electroreception: Fundamental Insights from
Comparative Approaches. Springer, Cham, pp. 63-90.

Lenhardt, M., Finn, R.N., Cakic, P., Kolarevic, J., Krpocetkovic, J., Radovic, I., Fyhn, H.J.,
2005. Analysis of the post-vitellogenic oocytes of three species of Danubian
Acipenseridae. Belg. J. Zool. 135, 205.

Ludwig, A., Belfiore, N.M., Pitra, C., Svirsky, V., Jenneckens, I., 2001. Genome duplication
events and functional reduction of ploidy levels in sturgeon (*Acipenser*, *Huso* and
Scaphirhynchus). Genetics 158, 1203-1215.

Lukoseviciute, M., Gavriouchkina, D., Williams, R.M., Hochgreb-Hagele, T., Senanayake, U.,
Chong-Morrison, V., Thongjuea, S., Repapi, E., Mead, A., Sauka-Spengler, T., 2018.
From pioneer to repressor: bimodal foxd3 activity dynamically remodels neural crest
regulatory landscape *in vivo*. Dev. Cell 47, 608-628.

Lush, M.E., Diaz, D.C., Koenecke, N., Baek, S., Boldt, H., St Peter, M.K., Gaitan-Escudero,
T., Romero-Carvajal, A., Busch-Nentwich, E.M., Perera, A.G., Hall, K.E., Peak, A.,
Haug, J.S., Piotrowski, T., 2019. scRNA-Seq reveals distinct stem cell populations that
drive hair cell regeneration after loss of Fgf and Notch signaling. eLife 8, e44431.

McGinnis, S., Madden, T.L., 2004. BLAST: at the core of a powerful and diverse set of
sequence analysis tools. Nucleic Acids Res. 32, W20-5.

Metscher, B.D., Müller, G.B., 2011. MicroCT for molecular imaging: quantitative visualization
of complete three-dimensional distributions of gene products in embryonic limbs. Dev.
Dyn. 240, 2301-2308.

Millimaki, B.B., Sweet, E.M., Dhason, M.S., Riley, B.B., 2007. Zebrafish *atoh1* genes: classic
proneural activity in the inner ear and regulation by Fgf and Notch. Development 134,
295-305.

Minařík, M., Modrell, M.S., Gillis, J.A., Campbell, A.S., Fuller, I., Lyne, R., Micklem, G., Gela,
D., Pšenička, M., Baker, C.V.H., 2023. Identification of multiple transcription factor
genes potentially involved in the development of electrosensory versus
mechanosensory lateral line organs. bioRxiv doi:
https://doi.org/10.1101/2023.04.14.536701.

Modrell, M.S., Baker, C.V.H., 2012. Evolution of electrosensory ampullary organs:
conservation of *Eya4* expression during lateral line development in jawed vertebrates.
Evol. Dev. 14, 277-285.

Modrell, M.S., Bemis, W.E., Northcutt, R.G., Davis, M.C., Baker, C.V.H., 2011a.
Electrosensory ampullary organs are derived from lateral line placodes in bony fishes.
Nat. Commun. 2, 496.
Modrell, M.S., Buckley, D., Baker, C.V.H., 2011b. Molecular analysis of neurogenic placode
development in a basal ray-finned fish. genesis 49, 278-294.
Modrell, M.S., Lyne, M., Carr, A.R., Zakon, H.H., Buckley, D., Campbell, A.S., Davis, M.C.,
Micklem, G., Baker, C.V.H., 2017a. Insights into electrosensory organ development,
physiology and evolution from a lateral line-enriched transcriptome. eLife 6, e24197.
Modrell, M.S., Tidswell, O.R.A., Baker, C.V.H., 2017b. Notch and Fgf signaling during
electrosensory versus mechanosensory lateral line organ development in a non-teleost
ray-finned fish. Dev. Biol. 431, 48-58.
Mogdans, J., 2019. Sensory ecology of the fish lateral-line system: Morphological and
physiological adaptations for the perception of hydrodynamic stimuli. J. Fish Biol. 95,
53-72.
Moser, T., Grabner, C.P., Schmitz, F., 2020. Sensory processing at ribbon synapses in the
retina and the cochlea. Physiol. Rev. 100, 103-144.
Mukhopadhyay, M., Pangrsic, T., 2022. Synaptic transmission at the vestibular hair cells of
amniotes. Mol. Cell. Neurosci. 121, 103749.
Nicolson, T., 2017. The genetics of hair-cell function in zebrafish. J. Neurogenet. 31, 102-
112.
Northcutt, R.G., 1997. Evolution of gnathostome lateral line ontogenies. Brain Behav. Evol.
50, 25-37.
Northcutt, R.G., Catania, K.C., Criley, B.B., 1994. Development of lateral line organs in the
axolotl. J. Comp. Neurol. 340, 480-514.
O'Neill, P., McCole, R.B., Baker, C.V.H., 2007. A molecular analysis of neurogenic placode
and cranial sensory ganglion development in the shark, *Scyliorhinus canicula*. Dev.
Biol. 304, 156-181.
Papalopulu, N., Kintner, C., 1996. A posteriorising factor, retinoic acid, reveals that
anteroposterior patterning controls the timing of neuronal differentiation in *Xenopus*
neuroectoderm. Development 122, 3409-3418.
Pauley, S., Lai, E., Fritzsch, B., 2006. *Foxg1* is required for morphogenesis and histogenesis
of the mammalian inner ear. Dev. Dyn. 235, 2470-2482.
Pickett, S.B., Raible, D.W., 2019. Water waves to sound waves: using zebrafish to explore
hair cell biology. J. Assoc. Res. Otolaryngol. 20, 1-19.
Piotrowski, T., Baker, C.V.H., 2014. The development of lateral line placodes: Taking a
broader view. Dev. Biol. 389, 68-81.

Roccio, M., Senn, P., Heller, S., 2020. Novel insights into inner ear development and
regeneration for targeted hearing loss therapies. Hear. Res. 397, 107859.

Singh, S., Groves, A.K., 2016. The molecular basis of craniofacial placode development.
WIREs Dev. Biol. 5, 363-376.

Square, T., Romášek, M., Jandzik, D., Cattell, M.V., Klymkowsky, M., Medeiros, D.M., 2015.
CRISPR/Cas9-mediated mutagenesis in the sea lamprey *Petromyzon marinus*: a
powerful tool for understanding ancestral gene functions in vertebrates. Development
142, 4180-4187.

Square, T.A., Jandzik, D., Massey, J.L., Romášek, M., Stein, H.P., Hansen, A.W.,
Purkayastha, A., Cattell, M.V., Medeiros, D.M., 2020. Evolution of the endothelin
pathway drove neural crest cell diversification. Nature 585, 563-568.

Stundl, J., Soukup, V., Franěk, R., Pospisilova, A., Psutkova, V., Pšenička, M., Cerny, R.,
Bronner, M.E., Medeiros, D.M., Jandzik, D., 2022. Efficient CRISPR mutagenesis in
sturgeon demonstrates its utility in large, slow-maturing vertebrates. Front. Cell Dev.
Biol. 10, 750833.

Tasdemir-Yilmaz, O.E., Druckenbrod, N.R., Olukoya, O.O., Dong, W., Yung, A.R., Bastille,
I., Pazyra-Murphy, M.F., Sitko, A.A., Hale, E.B., Vigneau, S., Gimelbrant, A.A.,
Kharchenko, P.V., Goodrich, L.V., Segal, R.A., 2021. Diversity of developing peripheral
glia revealed by single-cell RNA sequencing. Dev. Cell 56, 2516-2535.

Toresson, H., Martinez-Barbera, J.P., Bardsley, A., Caubit, X., Krauss, S., 1998.
Conservation of *BF-1* expression in amphioxus and zebrafish suggests evolutionary
ancestry of anterior cell types that contribute to the vertebrate telencephalon. Dev.
Genes Evol. 208, 431-439.

Uribe-Salazar, J.M., Kaya, G., Sekar, A., Weyenberg, K., Ingamells, C., Dennis, M.Y., 2022.
Evaluation of CRISPR gene-editing tools in zebrafish. BMC Genomics 23, 12.

Webb, J.F., 2021. Morphology of the mechanosensory lateral line system of fishes, in: B.
Fritzsch (Ed.), The Senses: A Comprehensive Reference. Elsevier, pp. 29-46.

Whitear, M., 1989. Merkel cells in lower vertebrates. Arch. Histol. Cytol. 52 Suppl., 415-422.

Wu, X., Kriz, A.J., Sharp, P.A., 2014. Target specificity of the CRISPR-Cas9 system. Quant.
Biol. 2, 59-70.

York, J.R., Zehnder, K., Yuan, T., Lakiza, O., McCauley, D.W., 2019. Evolution of Snail-
mediated regulation of neural crest and placodes from an ancient role in bilaterian
neurogenesis. Dev. Biol. 453, 180-190.

Yu, H.V., Tao, L., Llamas, J., Wang, X., Nguyen, J.D., Trecek, T., Segil, N., 2021. POU4F3
pioneer activity enables ATOH1 to drive diverse mechanoreceptor differentiation
through a feed-forward epigenetic mechanism. Proc. Natl. Acad. Sci. U.S.A. 118,
e2105137118.

Zhang, S., Zhang, Y., Dong, Y., Guo, L., Zhang, Z., Shao, B., Qi, J., Zhou, H., Zhu, W., Yan,
X., Hong, G., Zhang, L., Zhang, X., Tang, M., Zhao, C., Gao, X., Chai, R., 2019.
Knockdown of *Foxg1* in supporting cells increases the trans-differentiation of
supporting cells into hair cells in the neonatal mouse cochlea. Cell. Mol. Life Sci. 77,
1401-1419.
Zhang, Y., Zhang, S., Zhang, Z., Dong, Y., Ma, X., Qiang, R., Chen, Y., Gao, X., Zhao, C.,
Chen, F., He, S., Chai, R., 2020. Knockdown of *Foxg1* in Sox9+ supporting cells
increases the trans-differentiation of supporting cells into hair cells in the neonatal
mouse utricle. Aging (Albany NY) 12, 19834-19851.
Zhao, X.-F., Suh, C.S., Prat, C.R., Ellingsen, S., Fjose, A., 2009. Distinct expression of two
foxg1 paralogues in zebrafish. Gene Expr. Patterns 9, 266-272.

1021

1022

Target gene	sgRNA	Target sequence	PAM	Combinations used
Tyr	1	GGTGCCAAGGCAAAAACGCT	GGG	1+2, 1+2+3+4
	2	GATATCCCTCCATACATTAT	TGG	1+2, 1+2+3+4
	3	GATGTTTCTAAACATTGGGG	TGG	1+2+3+4
	4	GCTATGAATTTATTTTTTTC	AGG	1+2+3+4
	5	GCAAGGTATACGAAAGTTGA	CGG	5+6
	6	GATTGCAAGTTCGGCTTCTT	TGG	5+6
	7*	GGTTAGAGACTTTATGTAAC	GGG	7+8
	8*	GGCTCCATGTCTCAAGTCCA	AGG	7+8
Atoh1	1	GACCTTGTAAAAGATCGGAA	AGG	1+2
	2	GCTTGTCATTGTCAAATGAC	GGG	1+2
Neurod4	1	GGAGCGTTTCAAGGCCAGGC	GGG	1, 1+2+3+4, 1+6
	2	GTGAGCGTTCTCGCATGCAC	GGG	2, 1+2+3+4
	3	GCCTGGCCCACAACTACATC	TGG	1+2+3+4, 3+4+5+6
	4	GAGGGGCCCCGAGAAGCTGC	AGG	1+2+3+4, 3+4+5+6
	5	GTCTCCCCAGCCCTCCCTAC	GGG	5, 3+4+5+6
	6	GACAACCACTCCCCGGATTG	CGG	1+6, 3+4+5+6
	7	GACCCTGCGCAGGCTCTCCA	GGG	7+9, 7+8+9
	8	GCAGCTGGGTCCCCTGCTGA	CGG	7+8+9
	9	GGGGCCGTGTGCTCAGGGAT	GGG	7+9, 7+8+9
Foxg1	1	GAAACATCTTTTGCCCAACC	CGG	1+2
	2	TCTTCCGAGCAAGGTAACTC	GGG	1+2
	3	TGATGCTGAAGGACGACTTG	GGG	3+4
	4	CTGGCTCGTCCTCGGGCCGG	TGG	3+4

1023

1024 **Table 1. sgRNAs used in this study.** List of the genes targeted for CRISPR/Cas9-mediated
1025 mutagenesis, together with the target sequences and combinations of sgRNAs reported in this
1026 study. *Note: Tyr* sgRNAs 7 and 8 (marked with an asterisk) were designed and recently
1027 published by Stundl et al. (2022) as their *tyr* sgRNA 3 and *tyr* sgRNA 4, respectively.