Title: Ecological genomics in the Northern krill uncovers loci for local adaptation across ocean basins

Authors: Per Unneberg[1], Mårten Larsson[2,3], Anna Olsson[2], Ola Wallerman[2], Anna Petri[4], Ignas Bunikis[4], Olga Vinnere Pettersson[4], Chiara Papetti[5], Ástþór Gíslason[6], Henrik Glenner[7a,b], Joan E. Cartes[8], Leocadio Blanco-Bercial[9], Elena Eriksen[10], Bettina Meyer[11a,b,c], Andreas Wallberg[2]*

Affiliations:

[1]Department of Cell and Molecular Biology, National Bioinformatics Infrastructure Sweden, Science for Life Laboratory, Uppsala University; Uppsala, Sweden.

[2]Department of Medical Biochemistry and Microbiology, Uppsala University; Husargatan 3, 751 23 Uppsala, Sweden.

[3]Department of Pharmaceutical Biosciences, Uppsala University; Uppsala, Sweden.

[4]Uppsala Genome Center, Department of Immunology, Genetics and Pathology, Uppsala University, National Genomics Infrastructure hosted by SciLifeLab; Uppsala, Sweden.

[5]Biology Department, University of Padova; Padova, Italy.

[6]Marine and Freshwater Research Institute, Pelagic Division; Reykjavik, Iceland.

[7a]Department of Biological Sciences, University of Bergen; Bergen, Norway.

[7b]Center for Macroecology, Evolution and Climate Globe Institute, University of Copenhagen; Copenhagen, Denmark.

[8]Instituto de Ciencias del Mar (ICM); Barcelona, Spain.

[9]Bermuda Institute of Ocean Sciences, Arizona State University; St. George's, Bermuda.

[10]Institute of Marine Research (IMR); Bergen, Norway.

[11a]Section Polar Biological Oceanography, Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, Bremerhaven, Germany.
[11b]Institute for Chemistry and Biology of the Marine Environment, Carlvon Ossietzky University of Oldenburg, Oldenburg, Germany.
[11c]Helmholtz Institute for Functional Marine Biodiversity (HIFMB), University of Oldenburg, Oldenburg, Germany.

*Corresponding author. Email: andreas.wallberg@imbim.uu.se (A.W.)

Abstract:

Krill is a vital food source for many marine animals but also strongly impacted by climate change. Genetic adaptation could support populations, but remains uncharacterized. We assembled the 19 Gb Northern krill genome and compared genome-scale variation among 74 specimens from the colder Atlantic Ocean and warmer Mediterranean Sea. The genome is dominated by methylated transposable elements and contains many duplicated genes implied in molting and vision. Analysis of 760 million SNPs indicates extensive homogenizing gene-flow among populations. Nevertheless, we detect extreme divergence across hundreds of genes, governing ecophysiological functions like photoreception, circadian regulation, reproduction and thermal tolerance. Such standing variation may be essential for resilience in zooplankton, necessitating insight into adaptive variation to forecast their roles in future marine ecosystems and support ocean conservation.

One-Sentence Summary:

Genome-scans of Northern krill link genes for photoreception, reproduction and thermal tolerance to ecological adaptation.

Main Text:

Climate change is affecting all life on Earth and forcing species to move or adapt (*1*, *2*). Ocean plankton are crucial to maintaining food webs and fisheries but face many challenges including increased temperatures and acidification (*3–5*). Many planktonic species are shifting toward higher latitudes (*5*, *6*), and continued warming is expected to impact marine communities and ecosystem services (*7*). The long-term responses to these changes are unclear, but evolutionary adaptation may be important to sustain populations, particularly when physiological and geographical limits have otherwise been reached (*8*). This warrants the need to better understand adaptation in key zooplankton that strongly influence marine ecosystems. Krill (Euphausicea; 86 spp.), or euphausiids, are macrozooplankton crustaceans inhabiting all world oceans. Some species include trillions of individuals and are among the most abundant animals on Earth (*9–11*). As grazers of smaller plankton and food for fish and mammals, krill are critical links between primary production and higher trophic levels (*4*). However, polar krill of both hemispheres have declined in recent decades (*12*, *13*), while boreal species such as the Northern krill *Meganyctiphanes norvegica* spread into new areas (*14*), impacting native biodiversity (*15*, *16*).

The Northern krill is the largest and most abundant North Atlantic krill species, possibly structured into 3–4 basin-scale gene pools (*10*, *17*). While many krill species are stenothermal and have narrow latitudinal ranges, it has unusually broad thermal tolerance and range (*18*, *19*). It occurs across a 2–15°C temperature gradient (Fig. 1A) and breeds within 5–15°C (*10*), much wider thermal envelopes than for example the Antarctic krill *Euphausia superba* that is constrained within -2.0°C to +4.0°C and reproductively challenged already at +1.5°C (*20*). Northern krill from different climates vary in metabolism, nutrition, maturation and timing of reproduction that track local seasonal cycles (*18*, *21*), ranging from spawning in late winter–early

spring in the Mediterranean Sea to summer in the Atlantic Ocean. These phenotypic variations could have genetic bases, making *M. norvegica* an attractive model for environmental adaptation.

Zooplankton generally have large populations with considerable genetic variation, suggesting they have high potential to adapt to changing environments through natural selection ([22, 23]). Many also exhibit extensive larval dispersal and gene flow, which could either interfere with local adaptation or help introduce adaptive variants ([22]). Genomic evidence for mechanisms of adaptation is still critically missing for zooplankton ([24]). In particular, we lack insight into adaptation in euphausiids, due to their large and repetitive genomes ranging between 11–48 Gb (4–16× the human genome), which have hindered genetic analysis until now ([25]). The recently published Antarctic krill genome represented a major step forward but revealed extensive genetic homogeneity and limited adaptive variation in that circumpolar and panmictic species ([26]).

Here we use long-read technologies to assemble and characterize the huge genome of the Northern krill. We perform genome-scale population re-sequencing to map genetic variation and uncover the evolutionary history of this widespread species. Our ultimate aim is to identify genetic adaptations to different environments across its range. Insight into adaptive variation can help monitoring evolutionary processes including migration and adaptation, identify genetic diversity hotspots or offsets and support forecasting resilience of krill stocks under climate change ([27]).

Results

A highly repeated genome

The Northern krill is diploid and its nuclear karyotype consists of 19 large metacentric chromosome pairs without heteromorphy between sexes ([28]). We assembled and scaffolded the nuclear and mitochondrial genomes with Nanopore long-reads, Chromium linked-reads and RNA data from mostly a single specimen (tables S1–3; figs. S1–3; materials and methods). We tracked contiguity, completeness and accuracy by mapping coding transcripts (figs. S4–5). The finished genome assembly spanned 19.2 Gb (n=216,568 scaffolds/contigs) with ~90% of BUSCO genes present and ~0.5 base-level errors/kb (Fig. 1B, table S4) . The genome is GC-poor (%GC=29.9) and repeat-rich. Low-complexity sequences and simple repeats span 15% of the genome and AT-microsatellites alone cover 6% (table S3). Using a custom transposable element (TE) library, we find that about 74% of the Northern krill genome is repetitive (Fig. 1C; table S5), similar to the recently characterized Antarctic krill genome (72–92%) ([26]). While we are unable to classify all TEs, retrotransposons (LINEs+LTRs) outweigh DNA-transposons 4:1, similarly to the American lobster or black tiger shrimp ([29, 30]). This differs from the repeatome of the Antarctic krill, which appears to be dominated by DNA-transposons ([26]), and hints at divergent genome composition and evolution among krill. Furthermore, TE divergence in the Northern krill is comparably unimodal, not indicating multiple bursts of proliferation as seen in the Antarctic krill.

Fig. 1. Sampling, genome assembly and genome analyses of the Northern krill *Meganyctiphanes norvegica*. (**A**) Photo: Adult specimen from the Norwegian Lysefjord (©Rudolf Svensen; approximate scale). Map: Atlantic and Mediterranean sample locations (n=sample-sizes). Sea Surface Temperatures from Climate Reanalyzer (*31*). (**B**) Genome assembly statistics in kilobases (kb), megabases (Mb) or gigabases (Gb). Top: completeness and duplications of BUSCOv5 genes (n=1,013) across the genome assembly and protein models. Bottom: sizes of genomic regions (UTR=untranslated exonic regions; cds=coding). (**C**) The repeat landscape. (a) Proportions of repeat-masked and unmasked bases (%). (b) The divergence landscape of interspersed repeats. (**D**) The coding and full sequence lengths of 7,150 single-copy orthologs between *M. norvegica* and the lobster *H. americanus*. (**E**) Time-calibrated species tree (inferred from 1,011 single-copy orthologs; 100% bootstrap support for all nodes) and gene family evolution. (**F**) The proportion of methylated cytosines across 75 M CpG-dinucleotides in the genome (>10× coverage).

Expansion of cuticular and opsin gene families

We used RNA and comparative data to annotate 25,301 protein-coding genes along with 2,283 TEs (mostly expressed retrotransposons), and also detected another 14,643 potential yet unannotated gene models (tables S6–8; fig. S6A). Gene bodies (introns+exons) span 50,276 bp on average and occupy 10% of the genome, while coding sequence covers only 0.22% (Fig. 1B). About ~15% of the coding sequence appears repetitive (fig. S6B) and the bimodal repeat-content among unannotated genes hints at thousands of additional genes and TEs (fig. S6C). Gene bodies of orthologs between the krill and crustaceans with smaller genomes (table S9) are 3–8× longer in krill, while having similar amounts of coding sequence (Fig. 1D; fig. S7, table S10). Compared to the Antarctic krill (*26*), genes are ~2.5× longer in the Northern krill, consistent with lineage-specific proliferation of retrotransposable elements in its highly repeated introns (fig. S6B). We measured high synonymous divergence (*dS*=0.46) between the two species (n=13,371

genes). Using a decapod molecular clock (*32*) (table S10), this divergence suggests they split
from a common ancestor ~130 MYA, underscoring separate evolution over long time-scales.

We built a crustacean species tree and analyzed gene family evolution. We found 104 rapidly
expanding gene families in the Northern krill (p<0.05; Fig. 1E; fig. S8A–B; table S11), including
those related to chitin, cuticular metabolism, regulation of the molting cycle, which are important
processes for growth and reproduction in crustaceans. This is notable as renewal of the
exoskeleton is unusually frequent and plastic in euphausiids (*10, 33*), and similar expansions
were independently detected in the Antarctic krill genome (*26*). Moreover, we detected
expansions of the opsin gene repertoire, which encodes the light-sensitive receptors in
ommatidia. Fourteen opsins have previously been identified from RNA in the Antarctic krill *E.*
superba (*34*), which are thought to enable vision under the divergent light conditions
experienced throughout its life cycle and vertical migrations (*35*), while 16 opsins have recently
been inferred from *M. norvegica* RNA (*36*). We queried our *M. norvegica* gene-set and the
KrillDB^2 *E. superba* RNA database against the curated crustacean opsin dataset in ref 35. We
detected 19 genes in the former species and 15 putative genes in the latter (fig. S9–10), including
new visual middle wavelength-sensitive (MWS) opsins and non-visual arthropsins. All *E.*
superba opsins have homologues in the *M. norvegica* genome and all but one previously
identified *M. norvegica* transcripts can now be anchored unambiguously 1:1 to our gene models
(fig. S10). Our findings expand the known opsins in both species and suggest that opsin and
molting-gene duplications could be common to all euphausiids.

Ancient whole-genome duplication (WGD) may explain the evolutionary origins of large
genomes but is not commonly reported for crustaceans. To test for WGD, we interrogated
divergences between gene-paralogs and searched for Hox-gene duplications, but found no
supporting evidence (Supplementary text). Instead, the large genome of Northern krill is likely
due to TE proliferation and multiple small-scale duplications.

An active DNA methylation system
Epigenetic regulation of the genome may contribute to genome evolution, function and plastic
responses to environmental change (*37*). For example, CpG-methylation can silence the
expression of harmful transposable elements (TEs), paradoxically allowing them to persist and
contribute to genome expansion (*38*). DNA methylation (DNAm) of both TEs and protein-
coding genes is ancestral in Arthropoda, but has frequently been lost (*39*). Using DNA from
muscle, we here characterize, for the first time, the DNAm toolkit and genomic patterns in a
euphausiid. The genome has low CpG content (CpG$_{O/E}$=0.53), indicating DNAm. We find all
genes encoding the canonical methyltransferases and genes responsible for repair or
demethylation (*Dnmt1–3, AlkB2, TET2*; fig. S12–16; table S12), hallmarks of functional DNAm
(*39*). We scanned the Nanopore-reads for signals of CpG-methylation at 75 million CpG-sites.
Overall, DNAm rates are higher in genes than intergenic regions (75% vs. 62% of reads being
methylated; Fig. 1F; fig. S17) and positively associated with splice isoform variation
(n=2.5±0.05 isoforms/gene with >95% methylation rates vs. 1.8±0.03 isoforms/gene with <5%
methylation rates; fig. S17E). In contrast, we observed a lack of methylation in the mitochondrial
chromosome (4%). DNAm rates are higher across repeats vs. non-repeated DNA (69% vs. 50%),
and appear to target young retrotransposons with similar LTRs, that may have been recently
active (S17F). Gene-body methylation is similar to observations in marbled crayfish (*40*), while
repeat-oriented methylation reminds of the myriapod *Strigamia maritima* (*39*). The krill

methylome thus spans both genes and repeats, suggesting dual roles in gene regulation and
silencing TEs.

Genome-scale variation is shaped by linked selection and pervasive gene flow

To uncover patterns of genetic variation, we collected 74 Northern krill specimens from eight
geographical regions separated by up to 5,800 km, covering a range of environmental conditions
(Fig 1A; Supplementary text). We re-sequenced whole genomes to ~3×/specimen
(20–30×/population), mapped reads and called 760 million quality-filtered SNPs across
accessible sites determined with similar filters (8.4 of 19.2 Gb; fig. S18). We estimate genome-
wide nucleotide diversity (π; the average differences between pairs of sequences) to 1.31% per-
base, intermediate among arthropods and low compared to marine broadcast spawners like
oysters and sea-squirts (*41*). The population mutation rate (θ_w) is 1.62%/bp. Assuming mutation-
drift equilibrium and a mutation rate from snapping shrimp (*32*), we estimate the long-term
effective population size (N_E) to be 1.53 million, far below the expected census population size
of trillions (*10*). However, Tajima's D is negative (-0.53), indicating excess of low-frequency
variants compared to expectation under equilibrium, consistent with population expansion or
selection shaping genetic variation. In accordance, we observe ~30% reduction of variation over
genes (π=1.15%), and more so at coding and non-synonymous sites (Fig. 2Aa–b). This effect
extends up to 50–100 kb around genes (Fig. 2Ac), suggesting widespread impact of linked
selection. Using heterozygous genotypes in the reference specimen (~5,000 scaffolds >500 kb),
we applied the Sequentially Markovian Coalescent to model demographic history (*42, 43*). The
results indicate populations expanded half way through the last glacial period (*44*) (Fig. 2B).
Levels of variation are similar among populations (fig. S19A). The limited structure we find
among populations recapitulates geography (the fixation index F_{ST} is ≈0.06 on average; Fig. 2C)
and previously detected mitochondrial gene pools (*10*). Genetic distances (d_{XY}) only increase
marginally with geographic distance (fig. S19B–D; S20). The Mediterranean Sea sample is the
most divergent, but average d_{XY} is only 1.04× higher compared to distances among Atlantic
populations (1.71% vs. 1.64%). The majority of non-singleton variants are polymorphic in most
populations (fig. S21), indicating extensive gene flow among stocks.

Fig. 2. Genome-scale patterns of genetic variation, demographic history and population structure.
(**A**) The mean levels of genetic variation across: (a) the genome; (b) at non-synonymous (N) and
synonymous sites (S); or (c) across 1 kb non-overlapping windows upstream or downstream of putative
protein-coding gene bodies. Shaded regions are 95% confidence intervals (200 bootstrap replicates). (**B**)
Historical fluctuations in effective population size (N_E) inferred using PSMC and MSMC (y1-axis;
green=100 PSMC bootstrap replicates; KYA=thousands of years ago). The dashed line is long-term N_E.
The LR04 benthic $\delta^{18}O$ isotope (gray line, y2-axis) indicates ice volume and sea temperature (arrows=last
three interglacials). (**C**) Sample and population interrelationships. (a) Main clustering of genetic variation
in a principal component analysis. (b) Admixture analyses indicating levels of shared ancestry. (c)
NeighborNet network of pairwise F_{ST}-distances.

Signatures of ancient adaptive divergence across hundreds of genes

To reveal genomic signatures of adaptation, we partitioned the dataset into two major contrasts:
i) Atlantic vs. Mediterranean samples (at/me); and ii) North-Eastern vs. South-Western North
Atlantic samples (ea/we; Fig. 3A). For each contrast, we computed pairwise divergence in allele
frequencies. While most variation segregates at low differences ($F_{ST(at/me)}$=0.056; $F_{ST(ea/we)}$=0.017;
Fig. 3B; fig. S20A), we also detect about 8× as many highly divergent variants compared to
expectations from simulations of neutral drift (figs. S22B–C). Divergent regions (F_{ST}>0.4) span
<1% of the genome and are about 2× enriched for gene sequences and for extended haplotypes,
compared to undifferentiated regions (F_{ST}<0.2; Fig 3B), consistent with gene-centered signatures
of selective sweeps (*45*). We compared F_{ST} between genes and similarly sized 50 kb flanking
regions 50–100 kb away from genes, which may more often evolve neutrally. At extreme levels

of divergence (the top 0.1% most divergent flanking regions), genes outweigh flanking regions
by 7× (at/me) or 2× (ea/we), respectively, consistent with natural selection driving divergence
across many genes (Fig. 3C; table S13; fig. S23).

We analyzed the distribution of putatively adaptive variation across ocean basins by defining
gene-level haplotypes (having at least four diagnostic SNPs with $F_{ST}>0.5$). At many divergent
genes (exon-wide $F_{ST}>0.4$), both haplotypes were often present in both groups ($n_{at/me}$=319/660;
$n_{ea/we}$=34/34), indicating widespread standing variation (Fig. 3D). Southern or Scandinavian
populations were more polymorphic than Barents Sea and Svalbard populations (Fig. 3D), which
could reflect genetic drift margin of the Arctic species range (*46*), or ongoing selection. We
estimated the ages of minor alleles on the divergent haplotypes to learn how long they may have
been segregating in the species. We first used iSMC (*47*) to estimate a genome-average
recombination rate (r=0.32cM/Mb, n=652 scaffolds) and then applied the Genealogical
Estimation of Variant Age (GEVA) (*48*). This coalescent method infers the time to the most
recent common ancestor using mutation and recombination rates, without requiring *a priori*
assumptions about demographic history. We found that most variation originated over 1 MYA,
predating multiple glacial cycles, and that adaptive variation between Atlantic and Mediterranean
populations may predate that segregating in the Atlantic Ocean (Fig. 3E).

Fig. 3. Genetic divergence among Northern krill sampled across the Atlantic Ocean and Mediterranean Sea. (A) The two contrasts used to measure divergence. **(B)** Features of genomic regions at increasing divergence (mean F_{ST} across 1 kb windows). (a–b) The proportion of the genome or genic sequence (cds+intron+UTRs). (c) The absolute extended haplotype statistic |XP-nSL| scores (95% confidence intervals from 2,000 bootstrap replicates). **(C)** (a–c) Atlantic-Mediterranean contrast: (a) The distribution of per-gene F_{ST}-values computed across the exons of genes (cds+UTRs; n=36,997) vs. flanking intergenic regions (n=31,101). Gray lines are 1% and 0.1% percentiles of flanking F_{ST}. Numbers and proportions of genes are indicated at each percentile. (b) The proportion of genes vs. flanking regions at each F_{ST}-level. (c–d) Eastern-Western contrast: statistics as in a–b (n=36,397 vs. n=31,103). **(D)** Haplotype distribution for divergent genes (exon-wide F_{ST}>0.4). (a–c) Atlantic-Mediterranean contrast. (a) Numbers of genes with shared or private haplotypes (n=660). Mean frequencies of common and rare haplotypes in each group (b) and population (c) (95% confidence intervals as in (B)). (d–f) as in (a–c) but for 34 genes in the Eastern-Western contrast. **(E)** Age distributions of minor alleles across (at/me: ~22–25 K SNPs in 660 genes; we/ea: ~2.5 K in 34 genes).

Candidate genes for ecological adaptation associated with ecophysiological functions

For each contrast, we ranked all genes by exon-wide F_{ST}. The top gene between Atlantic and Mediterranean krill is *nose resistant to fluoxetine protein 6* (*nrf-6*) (table S13), encoding a membrane protein facilitating lipid transport. The gene is located within a high-divergence region and the Mediterranean samples are fixed for an *nrf-6* haplotype with low frequency among Atlantic samples (6%). Exon-wide F_{ST} is >0.8 and high XP-nSL scores indicate loss of variation consistent with a selective sweep among Mediterranean samples (Fig 4Aa–b). A comparative analysis indicates accelerated protein evolution on the Mediterranean haplotype (*dN/dS*: 0.54 vs. 0.31; fig. S24A–B), consistent with positive selection. We predicted the protein topology and detected 1.7× enrichment of missense variants in its extracellular part compared to synonymous variants (9/16 vs. 5/15; Fig 4Ac; fig. S24C-D; table S14), which may alter its function. This is noteworthy as *nrf-6* is important for yolk transport into eggs in worm (*49*) and ovary development and oogenesis in fly (*50*), and overexpressed in the ovaries of sexually precocious crabs (*51*). In the Northern krill, Mediterranean stocks depend on a short spring phytoplankton bloom to accumulate lipid stores and trigger vitellogenesis and reproduction in early spring (*10, 52*). The Mediterranean *nrf-6* variant could contribute to advanced reproductive timing, making it a strong candidate for local adaptation.

We next used fly homologues to query the ranked lists for shared gene ontologies. We find that Atlantic-Mediterranean divergence is significantly enriched for genes involved in oogenesis, muscle function, phototransduction and eye development, regulation of circadian rhythm and heterochronic development (fig. S25A; table S15). These include homologs for *ninaB* that synthesize visual pigment (*53*), *S* that regulate sleep/wake cycles in animals and the transcription factor *Kr-h1* that acts in the juvenile hormone signaling pathway to govern vitellogenesis and reproduction in arthropods (*54*) (Fig 4B). Overall, we find 30 genes in the 0.1%-percentile (F_{ST}=0.60–0.72; n=264) to be associated with vision-related ontologies, 1.5× more than expected by chance (p=0.033), and 112 genes in the 1%-percentile (F_{ST}=0.36–0.72; n=1,024), including four MWS opsins (table S15). Photoreception candidates often involve gene family paralogs (fig. S26) and belong to rapidly expanded gene families, which are otherwise underrepresented in the 1%-percentile (2.98× vs. 0.6×; table S15).

Across the South-Western and North-Eastern North Atlantic basins, the most divergent gene encodes a larval cuticular protein (homologue of *Cpr49Ac*; Fig 4C), while this contrast is enriched for signaling ontologies (fig. S25A–B). Among the top 20 genes we identify three heat-

shock chaperones/co-chaperones, including *CCT8*, that fold proteins and promote proteostasis under thermal stress (Fig 4C). Heat-shock proteins contribute to protective thermal tolerance in krill and many other species (*55*). Chaperonins (*CCT-n*) influence cold shock response in yeast and crustaceans (*56*, *57*), and *CCT8* evolution is implied in freeze-tolerance in Amur sleeper fish (*58*). We detect a sweep-like signature of extended *CCT8* haplotypes in North-Eastern krill, suggesting cold tolerance may have been naturally selected among these stocks.

Fig. 4. Adaptive divergence and candidates for local adaptation. (**A**) High divergence between Atlantic and Mediterranean samples across the gene *nrf-6*. (a) Top: per-SNP F_{ST} along the locus (green line=0.1% percentile for SNPs; dark-blue line=F_{ST} for 1 kb windows; black=gene model). Bottom: per-SNP XP-nSL (dark-gray bars=mean window-based XP-nSL). Positive values imply a selective sweep in the Mediterranean sample, negative values mark the Atlantic Ocean sample. (b) Magnified view of *nrf-6*. Exons colored by UTR (purple) or NRF-6 protein topology. (c) Modeled protein structure and topology (including signaling peptide; boxes=N/C terminals; green circles=non-synonymous/missense variants). (**B**) Examples of highly differentiated genes between Atlantic and Mediterranean samples. (a) Photoreception: *retinal degeneration B* (*rdgB*), *neither inactivation nor afterpotential B* (*ninaB*) and *cacophony* (*cac*). (b) Heterochronic development: *Kruppel homolog 1* (*Kr-h1*). (c) Circadian regulation: *Star* (*S*), *clockwork orange* (*cwo*), *Angiotensin-converting enzyme-related* (*Acer*) and *narrow* (*na*). Statistics as in (A). (**C**) (a) The top gene in the Atlantic comparison was *Cuticular protein 49Ac* (*Cpr49Ac*) (b) Three chaperon genes implied thermal tolerance: *Chaperonin containing TCP1 subunit 8* (*CCT8*), *Hsc/Hsp70-interacting protein related* (*HIP-R*) and *Heat shock protein 83* (*Hsp83*).

Conclusions and implications

We here provide novel insight into the evolutionary history of the Northern krill. Our genome assembly reveals a highly methylated repeatome and many expanded gene families. Paralogous genes may have arisen from ectopic recombination between non-homologous repeated loci, which is more likely to occur when genomes accumulate transposable elements (*59*). The krill genome appears to continuously have evolved new genes, including those involved in molting and vision.

Molting is a crucial process in krill, being interlinked with growth and reproduction, and controlled by environmental cues including light (*10*, *60*). Some of the top gene candidates for local adaptation belong to expanded gene families, including two *ninaB* paralogs and four *ninaE* paralogs that synthesize visual pigment or encode MWS opsins. We detect elevated divergence in fifteen genes encoding cuticular proteins, several belonging to expanded families. Expanded cuticular and opsin gene families and functionally diverged paralogs may have enabled development and photoreception under diverse conditions in ancestral krill and provide substrate for adaptation also today (*61*).

We have detected signatures of polygenic adaptation involving small-to-moderate shifts in allele frequencies across many genes (*62*). Evolve-and-resequence experiments in copepods have similarly found polygenic signals of adaptation to temperature and acidity in laboratory conditions (*63*). Genetic adaptation in zooplankton could commonly involve numerous loci, warranting genome-scale assays to map adaptive variation. Many of our candidate genes have roles in photoreception, circadian rhythm, and oogenesis, ecophysiological functions also implied in adaptation in other widespread pelagic species, such as the Atlantic herring (*64*). The variants may help krill respond to light, temperature and resources in different environments.

Photoreception varies among *M. norvegica* populations: krill from turbid waters around the Gulf of Maine are more light sensitive than those from clearer waters (*65*). Moreover, light sensitivities in the North Pacific krill *E. pacifica* are tuned to local conditions. Individuals inhabiting shallow green water in the Saanich Inlet are more sensitive to green light compared to those from the deeper blue water of the San Diego Trough (*66*). Water clarity and light penetration influences behavior in *M. norvegica*. To avoid predators, stocks prefer deeper depths in more clear Norwegian fjords (*67*), while the deepest daytime depths (400–800 m) are known from the oligotrophic Mediterranean Sea (*10*). Eye traits are generally fast-evolving among krill species and associated with ecological niche (*68*). At least thirty genes involved in eye function diverge strongly between Atlantic and Mediterranean krill, and another six genes segregate across the Atlantic Ocean, suggesting heritable variation could contribute to these phenotypes, and that our gene candidates could help reconstruct the genomic architecture of vision and behavior in krill.

Seasonal and daily cycles of ambient light are central to zooplankton and influence diapause, vertical migrations and entrainment of endogenous circadian clocks that control the daily rhythm of physiological processes (*69*). *Euphausia superba* and *Thysanoessa inermis* krill have endogenous circadian rhythms (ECR) that oscillate faster than 24 h in the absence of light, or respond to minute irradiance. This may reflect adaptation to extreme photoperiodic variability at high-latitudes where the sun is either below or above the horizon for extended periods (*70*, *71*), although the genetic mechanisms of these adaptations are unknown. *Meganyctyphanes norvegica* also shows ECR and expresses a full set of circadian clock genes (*72*). We find that genes likely involved in regulating its circadian clock diverge across its range, including homologs of *narrow abdomen*, *glass* and *clockwork orange*. Functional assessments of the variants could illuminate how biological clocks are set in different environments.

Marine ecosystems are at unprecedented rates, causing redistribution of organisms and impacting food webs (*6*, *16*). In the Southern Ocean, the Antarctic krill is predicted to decline further, which could severely impact the Antarctic ecosystem (*73*), which could severely impact the Antarctic ecosystem. The Northern krill could be declining around Iceland (*74*), but increasing in

11   11

the Barents Sea (*14*). Where will they thrive in the future? We found many variants that could
help krill adapt to new or changing environments, some of which are widely distributed, old and
possibly maintained by long-term balancing selection under slowly fluctuating conditions.
Genotypes that are adaptive in warmer conditions could be key to cope with rapid climate
change (*27*). These adaptive variants could be important for coping with rapid climate change,
and Scandinavian stocks in particular may serve as sources of genetic diversity typical of the
Mediterranean Sea or other southern populations. The next frontier for the Northern krill is the
Arctic Ocean (*14, 16*). Standing variants supporting physiological processes under darker or
colder conditions may help establish it there. The genetically most divergent population is that of
the Mediterranean Sea, which is close to its southern limit. This population appears to lack
variation at many adaptive loci, which might limit its evolutionary potential, although our
analysis is limited by a small sample size.

With the exception of the Antarctic krill (*9*), long-term monitoring of krill abundance is typically
performed or reported in aggregate (*13*), obscuring how individual species fare under climate
change. Our results suggest that krill may commonly be genetically fine-tuned to their
environments, while previous research underscore limited thermal tolerances in many species
(*19, 20*). Genetic adaptation could be the major process determining whether krill will persist or
perish. The many candidate genes reported here can be used as biomarkers to diagnose and
monitor change of adaptive variation also in other species and climates, in order to better forecast
their future distributions and the risks of the great many species and ecosystems that depend on
them.

References and Notes

1. G. T. Pecl, M. B. Araújo, J. D. Bell, J. Blanchard, T. C. Bonebrake, I.-C. Chen, T. D. Clark, R. K. Colwell, F.
Danielsen, B. Evengård, L. Falconi, S. Ferrier, S. Frusher, R. A. Garcia, R. B. Griffis, A. J. Hobday, C.
Janion-Scheepers, M. A. Jarzyna, S. Jennings, J. Lenoir, H. I. Linnetved, V. Y. Martin, P. C. McCormack, J.
McDonald, N. J. Mitchell, T. Mustonen, J. M. Pandolfi, N. Pettorelli, E. Popova, S. A. Robinson, B. R.
Scheffers, J. D. Shaw, C. J. B. Sorte, J. M. Strugnell, J. M. Sunday, M.-N. Tuanmu, A. Vergés, C. Villanueva,
T. Wernberg, E. Wapstra, S. E. Williams, Biodiversity redistribution under climate change: Impacts on
ecosystems and human well-being. *Science*. **355**, eaai9214 (2017).
2. C. Bellard, C. Bertelsmeier, P. Leadley, W. Thuiller, F. Courchamp, Impacts of climate change on the future
of biodiversity. *Ecol. Lett.* **15**, 365–377 (2012).
3. *IPCC, 2019: IPCC Special Report on the Ocean and Cryosphere in a Changing Climate [H.-O. Pörtner, D.C.*
Roberts, V. Masson-Delmotte, P. Zhai, M. Tignor, E. Poloczanska, K. Mintenbeck, A. Alegría, M. Nicolai, A.
Okem, J. Petzold, B. Rama, N.M. Weyer (eds.)]. In press.
4. M. M. McBride, P. Dalpadado, K. F. Drinkwater, O. R. Godø, A. J. Hobday, A. B. Hollowed, T. Kristiansen,
E. J. Murphy, P. H. Ressler, S. Subbey, E. E. Hofmann, H. Loeng, Krill, climate, and contrasting future
scenarios for Arctic and Antarctic fisheries. *ICES J. Mar. Sci.* **71**, 1934–1955 (2014).
5. A. J. Richardson, In hot water: zooplankton and climate change. *ICES J. Mar. Sci.* **65**, 279–295 (2008).
6. E. S. Poloczanska, C. J. Brown, W. J. Sydeman, W. Kiessling, D. S. Schoeman, P. J. Moore, K. Brander, J. F.
Bruno, L. B. Buckley, M. T. Burrows, C. M. Duarte, B. S. Halpern, J. Holding, C. V. Kappel, M. I. O'Connor,
J. M. Pandolfi, C. Parmesan, F. Schwing, S. A. Thompson, A. J. Richardson, Global imprint of climate change
on marine life. *Nat. Clim. Change*. **3**, 919–925 (2013).
7. F. Benedetti, M. Vogt, U. H. Elizondo, D. Righetti, N. E. Zimmermann, N. Gruber, Major restructuring of
marine plankton assemblages under global warming. *Nat. Commun.* **12**, 5226 (2021).
8. A. A. Hoffmann, C. M. Sgrò, Climate change and evolutionary adaptation. *Nature*. **470**, 479–485 (2011).
9. A. Atkinson, V. Siegel, E. A. Pakhomov, M. J. Jessopp, V. Loeb, A re-appraisal of the total biomass and
annual production of Antarctic krill. *Deep Sea Res. Part Oceanogr. Res. Pap.* **56**, 727–740 (2009).
10. G. Tarling, *Biology of Northern Krill* (Academic Press, 2010).
11. Y. M. Bar-On, R. Phillips, R. Milo, The biomass distribution on Earth. *Proc. Natl. Acad. Sci. U. S. A.* **115**,

6506–6511 (2018).

12.   A. Atkinson, S. L. Hill, E. A. Pakhomov, V. Siegel, C. S. Reiss, V. J. Loeb, D. K. Steinberg, K. Schmidt, G.
A. Tarling, L. Gerrish, S. F. Sailley, Krill (Euphausia superba) distribution contracts southward during rapid
regional warming. *Nat. Clim. Change.* **9**, 142 (2019).

13.   M. Edwards, P. Hélaouët, E. Goberville, A. Lindley, G. A. Tarling, M. T. Burrows, A. Atkinson, North
Atlantic warming over six decades drives decreases in krill abundance with no associated range shift.
Commun. Biol. **4**, 1–10 (2021).

14.   E. Eriksen, H. Rune Skjoldal, H. Gjøsæter, R. Primicerio, Spatial and temporal changes in the Barents Sea
pelagic compartment during the recent warming. *Prog. Oceanogr.*, doi:10.1016/j.pocean.2016.12.009.

15.   H. Flores, A. Atkinson, S. Kawaguchi, B. A. Krafft, G. Milinevsky, S. Nicol, C. Reiss, G. A. Tarling, R.
Werner, E. B. Rebolledo, V. Cirelli, J. Cuzin-Roudy, S. Fielding, J. J. Groeneveld, M. Haraldsson, A.
Lombana, E. Marschoff, B. Meyer, E. A. Pakhomov, E. Rombola, K. Schmidt, V. Siegel, M. Teschke, H.
Tonkes, J.-Y. Toullec, P. N. Trathan, N. Tremblay, A. P. V. de Van de Putte, J. A. van Van Franeker, T.
Werner, Impact of climate change on Antarctic krill. *Mar. Ecol. Prog. Ser.* **458**, 1–19 (2012).

16.   R. B. Ingvaldsen, K. M. Assmann, R. Primicerio, M. Fossheim, I. V. Polyakov, A. V. Dolgov, Physical
manifestations and ecological implications of Arctic Atlantification. *Nat. Rev. Earth Environ.* **2**, 874–889
(2021).

17.   C. Papetti, L. Zane, E. Bortolotto, A. Bucklin, T. Patarnello, Genetic differentiation and local temporal
stability of population structure in the euphausiid Meganyctiphanes norvegica. *Mar. Ecol. Prog. Ser.* **289**,
225–235 (2005).

18.   R. Saborowski, F. Buchholz, Metabolic properties of Northern krill, Meganyctiphanes norvegica, from
different climatic zones: Enzyme characteristics and activities. *Mar. Biol.* **140**, 557–565 (2002).

19.   K. Hünerlage, F. Buchholz, Thermal limits of krill species from the high-Arctic Kongsfjord (Spitsbergen).
Mar. Ecol. Prog. Ser. **535**, 89–98 (2015).

20.   F. A. Perry, S. Kawaguchi, A. Atkinson, S. F. Sailley, G. A. Tarling, D. J. Mayor, C. H. Lucas, R. King, A.
Cooper, Temperature–Induced Hatch Failure and Nauplii Malformation in Antarctic Krill. *Front. Mar. Sci.* **7**
(2020) (available at https://www.frontiersin.org/article/10.3389/fmars.2020.00501).

21.   E. Albessard, P. Mayzaud, Influence of tropho-climatic environment and reproduction on lipid composition of
the euphausiid Meganyctiphanes norvegica in the Ligurian Sea, the Clyde Sea and the Kattegat. *Mar. Ecol.
Prog. Ser.* **253**, 217–232 (2003).

22.   K. T. C. A. Peijnenburg, E. Goetze, High evolutionary potential of marine zooplankton. *Ecol. Evol.* **3**, 2765–
2781 (2013).

23.   J. M. Lourenço, S. Glémin, N. Galtier, The Rate of Molecular Adaptation in a Changing Environment. *Mol.
Biol. Evol.* **30**, 1292–1301 (2013).

24.   A. Bucklin, K. R. DiVito, I. Smolina, M. Choquet, J. M. Questel, G. Hoarau, R. J. O'Neill, "Population
Genomics of Marine Zooplankton" in *Population Genomics: Marine Organisms*, M. F. Oleksiak, O. P.
Rajora, Eds. (Springer International Publishing, Cham, 2018;
http://link.springer.com/10.1007/13836_2017_9), *Population Genomics*, pp. 61–102.

25.   N. W. Jeffery, The first genome size estimates for six species of krill (Malacostraca, Euphausiidae): large
genomes at the north and south poles. *Polar Biol.* **35**, 959–962 (2012).

26.   C. Shao, S. Sun, K. Liu, J. Wang, S. Li, Q. Liu, B. E. Deagle, I. Seim, A. Biscontin, Q. Wang, X. Liu, S.
Kawaguchi, Y. Liu, S. Jarman, Y. Wang, H.-Y. Wang, G. Huang, J. Hu, B. Feng, C. De Pittà, S. Liu, R.
Wang, K. Ma, Y. Ying, G. Sales, T. Sun, X. Wang, Y. Zhang, Y. Zhao, S. Pan, X. Hao, Y. Wang, J. Xu, B.
Yue, Y. Sun, H. Zhang, M. Xu, Y. Liu, X. Jia, J. Zhu, S. Liu, J. Ruan, G. Zhang, H. Yang, X. Xu, J. Wang, X.
Zhao, B. Meyer, G. Fan, The enormous repetitive Antarctic krill genome reveals environmental adaptations
and population insights. *Cell* (2023), doi:10.1016/j.cell.2023.02.005.

27.   J. C. Teixeira, C. D. Huber, The inflated significance of neutral genetic diversity in conservation genetics.
Proc. Natl. Acad. Sci. **118** (2021), doi:10.1073/pnas.2015096118.

28.   C. Thiriot-Quiévreux, J. Cuzin-Roudy, Karyological Study of the Mediterranean Krill Meganyctiphanes
norvegica (Euphausiacea). *J. Crustac. Biol.* **15**, 79–85 (1995).

29.   J. M. Polinski, A. V. Zimin, K. F. Clark, A. B. Kohn, N. Sadowski, W. Timp, A. Ptitsyn, P. Khanna, D. Y.
Romanova, P. Williams, S. J. Greenwood, L. L. Moroz, D. R. Walt, A. G. Bodnar, The American lobster
genome reveals insights on longevity, neural, and immune adaptations. *Sci. Adv.* **7**, eabe8290 (2021).

30.   T. Uengwetwanit, W. Pootakham, I. Nookaew, C. Sonthirod, P. Angthong, K. Sittikankaew, W.
Rungrassamee, S. Arayamethakorn, T. Wongsurawat, P. Jenjaroenpun, D. Sangsrakru, R. Leelatanawit, J.
Khudet, J. J. Koehorst, P. J. Schaap, V. M. dos Santos, F. Tangy, N. Karoonuthaisiri, A chromosome-level
assembly of the black tiger shrimp (Penaeus monodon) genome facilitates the identification of growth-

associated genes. *Mol. Ecol. Resour.* **21**, 1620–1640 (2021).

31.   S. D. Birkel, "Monthly Sea Surface Temperature", Climate Reanalyzer (https://ClimateReanalyzer.org),
Climate Change Institute, University of Maine, USA. Accessed on 2017-02-28 (2017).

32.   K. Silliman, J. L. Indorf, N. Knowlton, W. E. Browne, C. Hurt, Base-substitution mutation rate across the
nuclear genome of Alpheus snapping shrimp and the timing of isolation by the Isthmus of Panama. *BMC*
Ecol. Evol. **21**, 104 (2021).

33.   P. J. Seear, G. A. Tarling, G. Burns, W. P. Goodall-Copestake, E. Gaten, Ö. Özkaya, E. Rosato, Differential
gene expression during the moult cycle of Antarctic krill (Euphausia superba). *BMC Genomics*. **11**, 582
(2010).

34.   I. Urso, A. Biscontin, D. Corso, C. Bertolucci, C. Romualdi, C. De Pittà, B. Meyer, G. Sales, A thorough
annotation of the krill transcriptome offers new insights for the study of physiological processes. *Sci. Rep.* **12**,
11415 (2022).

35.   A. Biscontin, E. Frigato, G. Sales, G. M. Mazzotta, M. Teschke, C. De Pittà, S. Jarman, B. Meyer, R. Costa,
C. Bertolucci, The opsin repertoire of the Antarctic krill Euphausia superba. *Mar. Genomics*. **29**, 61–68
(2016).

36.   S. Palecanda, T. Iwanicki, M. Steck, M. L. Porter, Crustacean conundrums: a review of opsin diversity and
evolution. *Philos. Trans. R. Soc. B Biol. Sci.* **377**, 20210289 (2022).

37.   A. Ashe, V. Colot, B. P. Oldroyd, How does epigenetics influence the course of evolution? *Philos. Trans. R.*
Soc. B Biol. Sci. **376**, 20200111 (2021).

38.   W. Zhou, G. Liang, P. L. Molloy, P. A. Jones, DNA methylation enables transposable element-driven genome
expansion. *Proc. Natl. Acad. Sci.* **117**, 19359–19366 (2020).

39.   S. H. Lewis, L. Ross, S. A. Bain, E. Pahita, S. A. Smith, R. Cordaux, E. A. Miska, B. Lenhard, F. M. Jiggins,
P. Sarkies, Widespread conservation and lineage-specific diversification of genome-wide DNA methylation
patterns across arthropods. *PLOS Genet.* **16**, e1008864 (2020).

40.   F. Gatzmann, C. Falckenhayn, J. Gutekunst, K. Hanna, G. Raddatz, V. C. Carneiro, F. Lyko, The methylome
of the marbled crayfish links gene body methylation to stable expression of poorly accessible genes.
Epigenetics Chromatin. **11**, 57 (2018).

41.   E. M. Leffler, K. Bullaughey, D. R. Matute, W. K. Meyer, L. Ségurel, A. Venkat, P. Andolfatto, M.
Przeworski, Revisiting an Old Riddle: What Determines Genetic Diversity Levels within Species? *PLOS Biol.*
10, e1001388 (2012).

42.   H. Li, R. Durbin, Inference of human population history from individual whole-genome sequences. *Nature*.
475, 493–496 (2011).

43.   S. Schiffels, R. Durbin, Inferring human population size and separation history from multiple genome
sequences. *Nat. Genet.* **46**, 919–925 (2014).

44.   L. E. Lisiecki, M. E. Raymo, A Pliocene-Pleistocene stack of 57 globally distributed benthic δ18O records.
Paleoceanography. **20** (2005), doi:10.1029/2004PA001071.

45.   Z. A. Szpiech, T. E. Novak, N. P. Bailey, L. S. Stevison, Application of a novel haplotype-based scan for
local adaptation to study high-altitude adaptation in rhesus macaques. *Evol. Lett.* **5**, 408–421 (2021).

46.   C. G. Eckert, K. E. Samis, S. C. Lougheed, Genetic variation across species' geographical ranges: the central–
marginal hypothesis and beyond. *Mol. Ecol.* **17**, 1170–1188 (2008).

47.   G. V. Barroso, N. Puzović, J. Y. Dutheil, Inference of recombination maps from a single pair of genomes and
its application to ancient samples. *PLOS Genet.* **15**, e1008449 (2019).

48.   P. K. Albers, G. McVean, Dating genomic variants and shared ancestry in population-scale sequencing data.
PLOS Biol. **18**, e3000586 (2020).

49.   R. K. M. Choy, J. H. Thomas, Fluoxetine-Resistant Mutants in C. elegans Define a Novel Family of
Transmembrane Proteins. *Mol. Cell*. **4**, 143–152 (1999).

50.   S. Dzitoyeva, N. Dimitrijevic, H. Manev, Identification of a novel Drosophila gene, beltless, using injectable
embryonic and adult RNA interference (RNAi). *BMC Genomics*. **4**, 33 (2003).

51.   X. Chen, J. Wang, X. Hou, W. Yue, S. Huang, C. Wang, Tissue expression profiles unveil the gene interaction
of hepatopancreas, eyestalk, and ovary in the precocious female Chinese mitten crab, Eriocheir sinensis. *BMC*
Genet. **20**, 12 (2019).

52.   P. Mayzaud, P. Virtue, E. Albessard, Seasonal variations in the lipid and fatty acid composition of the
euphausiid Meganyctiphanes norvegica from the Ligurian Sea. *Mar. Ecol. Prog. Ser.* **186**, 199–210 (1999).

53.   O. Voolstra, V. Oberhauser, E. Sumser, N. E. Meyer, M. E. Maguire, A. Huber, J. von Lintig, NinaB is
essential for Drosophila vision but induces retinal degeneration in opsin-deficient photoreceptors. *J. Biol.*
Chem. **285**, 2130–2139 (2010).

54.   X. Xie, M. Liu, Q. Jiang, H. Zheng, L. Zheng, D. Zhu, Role of Kruppel homolog 1 (Kr-h1) in methyl

farnesoate-mediated vitellogenesis in the swimming crab Portunus trituberculatus. *Gene.* **679**, 260–265 (2018).

55. N. A. Yusof, M. Masnoddin, J. Charles, Y. Q. Thien, F. N. Nasib, C. M. V. L. Wong, A. M. Abdul Murad, N. M. Mahadi, I. Bharudin, Can heat shock protein 70 (HSP70) serve as biomarkers in Antarctica for future ocean acidification, warming and salinity stress? *Polar Biol.* **45**, 371–394 (2022).

56. L. Somer, O. Shmulman, T. Dror, S. Hashmueli, Y. Kashi, The eukaryote chaperonin CCT is a cold shock protein in Saccharomyces cerevisiae. *Cell Stress Chaperones.* **7**, 47–54 (2002).

57. K. Yu, J. Gong, C. Huang, H. Huang, H. Ye, G. Wang, C. Zeng, Characterization of CCTα and evaluating its expression in the mud crab Scylla paramamosain when challenged by low temperatures alone and in combination with high and low salinity. *Cell Stress Chaperones.* **20**, 853–864 (2015).

58. W. Lv, H. Jiang, Y. Qian, M. Meng, C. Wang, N. Sun, Y. Lu, H. Bing, C. Fang, D. M. Irwin, S. He, L. Yang, Multiple omic investigations of freeze tolerance adaptation in the aquatic ectothermic vertebrate, the Amur sleeper (2022), p. 2022.05.10.491133, , doi:10.1101/2022.05.10.491133.

59. I. R. Arkhipova, Neutral Theory, Transposable Elements, and Eukaryotic Genome Evolution. *Mol. Biol. Evol.* **35**, 1332–1337 (2018).

60. P. Seear, G. A. Tarling, M. Teschke, B. Meyer, M. A. S. Thorne, M. S. Clark, E. Gaten, E. Rosato, Effects of simulated light regimes on gene expression in Antarctic krill (Euphausia superba Dana). *J. Exp. Mar. Biol. Ecol.* **381**, 57–64 (2009).

61. W. Qian, J. Zhang, Genomic evidence for adaptation by gene duplication. *Genome Res.* **24**, 1356–1362 (2014).

62. J. K. Pritchard, J. K. Pickrell, G. Coop, The genetics of human adaptation: hard sweeps, soft sweeps, and polygenic adaptation. *Curr. Biol. CB.* **20**, R208-215 (2010).

63. R. S. Brennan, J. A. deMayo, H. G. Dam, M. Finiguerra, H. Baumann, V. Buffalo, M. H. Pespeni, Experimental evolution reveals the synergistic genomic mechanisms of adaptation to ocean warming and acidification in a marine copepod. *Proc. Natl. Acad. Sci.* **119**, e2201521119 (2022).

64. F. Han, M. Jamsandekar, M. E. Pettersson, L. Su, A. P. Fuentes-Pardo, B. W. Davis, D. Bekkevold, F. Berg, M. Casini, G. Dahle, E. D. Farrell, A. Folkvord, L. Andersson, Ecological adaptation in Atlantic herring is associated with large shifts in allele frequencies at hundreds of loci. *eLife.* **9**, e61076 (2020).

65. T. J. Myslinski, T. M. Frank, E. A. Widder, Correlation between photosensitivity and downwelling irradiance in mesopelagic crustaceans. *Mar. Biol.* **147**, 619–629 (2005).

66. F. S. Russell, C. M. Yonge, J. Mauchline, L. R. Fisher, Adv. Mar. Biol. 7: The biology of euphausiids. *Adv. Mar. Biol.* (1969).

67. M. S. R. Onsrud, S. Kaartvedt, Diel vertical migration of the krill Meganyctiphanes norvegica in relation to physical environment, food and predators. *Mar. Ecol. Prog. Ser.* **171**, 209–219 (1998).

68. A. L. Vereshchaka, D. N. Kulagin, A. A. Lunina, A phylogenetic study of krill (Crustacea: Euphausiacea) reveals new taxa and co-evolution of morphological characters. *Cladistics.* **35**, 150–172 (2019).

69. K. Bandara, Ø. Varpe, L. Wijewardene, V. Tverberg, K. Eiane, Two hundred years of zooplankton vertical migration research. *Biol. Rev.* **96**, 1547–1589 (2021).

70. A. Biscontin, T. Wallach, G. Sales, A. Grudziecki, L. Janke, E. Sartori, C. Bertolucci, G. Mazzotta, C. De Pittà, B. Meyer, A. Kramer, R. Costa, Functional characterization of the circadian clock in the Antarctic krill, Euphausia superba. *Sci. Rep.* **7**, 17742 (2017).

71. J. H. Cohen, K. S. Last, C. L. Charpentier, F. Cottier, M. Daase, L. Hobbs, G. Johnsen, J. Berge, Photophysiological cycles in Arctic krill are entrained by weak midday twilight during the Polar Night. *PLOS Biol.* **19**, e3001413 (2021).

72. A. E. Christie, A. Yu, M. G. Pascual, Circadian signaling in the Northern krill Meganyctiphanes norvegica: In silico prediction of the protein components of a putative clock system using a publicly accessible transcriptome. *Mar. Genomics.* **37**, 97–113 (2018).

73. S. Kawaguchi, A. Ishida, R. King, B. Raymond, N. Waller, A. Constable, S. Nicol, M. Wakita, A. Ishimatsu, Risk maps for Antarctic krill under projected Southern Ocean acidification. *Nat. Clim. Change.* **3**, 843–847 (2013).

74. T. Silva, A. Gislason, P. Licandro, G. Marteinsdóttir, A. S. A. Ferreira, K. Gudmundsson, O. S. Astthorsson, Long-term changes of euphausiids in shelf and oceanic habitats southwest, south and southeast of Iceland. *J. Plankton Res.* **36**, 1262–1278 (2014).

Acknowledgments: We thank Stéphane Plourde, Geneviève Perrin and Katharina Michael for providing samples, the staff at Kristineberg Center for Marine Research and Innovation, Zhen Li for constructive discussion and Jessica Heinze, Sarah Demirkale and Ylva Jondelius for lab assistance. The computational infrastructure was enabled by project SNIC 2022/5-472 provided by the Swedish National Infrastructure for Computing (SNIC) at UPPMAX and the PDC Center for High Performance Computing, KTH Royal Institute of Technology, partially funded by the Swedish Research Council through grant agreement no. 2018-05973.

Funding:

FORMAS Future research leaders grant 2017-00413 (A.W.)

NSF OCE 1316040, 1948162 (L.B.B.)

Author contributions:

Conceptualization: AW

Data curation: AW PU IB

Formal analysis: AW PU ML IB

Funding acquisition: AW

Investigation: AW AO AP OW

Methodology: AW PU AP

Project administration: AW OVP

Resources: AW OVP EE AG HG JC LBB

Software: AW PU

Supervision: AW

Visualization: AW PU ML

Writing – original draft: AW PU

Writing – review & editing: AW PU BM LBB EE AG HG JC CP

Competing interests: Authors declare that they have no competing interests.

Data and materials availability:

Supplementary Materials

Materials and Methods

Supplementary Text

Figs. S1 to S26

Tables S1 to S14

References (##–##)

DNA extraction protocol