Oncolytic vesicular stomatitis virus is safe and provides a survival benefit for dogs with naturally occurring osteosarcoma

Kelly M. Makielski[1-3], Aaron L. Sarver[1,2,4], Michael S. Henson[3], Kathleen M. Stuebner[3], Antonella Borgatti[3], Lukkana Suksanpaisan[5], Caitlin Preusser[6], Alexandru-Flaviu Tabaran[7], Ingrid Cornax[7], M. Gerard O'Sullivan[7], Andrea Chehadeh[3], Donna Groschen[3], Kelly Bergsrud[3], Sara Pracht[3], Amber Winter[3], Lauren J. Mills[1,8], Marc D. Schwabenlander[9], Melissa Wolfe[9], Michael A. Farrar[10,11], Gary R. Cutter[12], Joseph S. Koopmeiners[1,13], Stephen J. Russell[6,14], Jaime F. Modiano[1-3,10,15,16], Shruthi Naik[6,14]

[1]Masonic Cancer Center, University of Minnesota, Minneapolis, Minnesota 55455, USA

[2]Animal Cancer Care and Research Program, University of Minnesota, St. Paul, Minnesota 55108, USA

[3]Department of Veterinary Clinical Sciences, University of Minnesota College of Veterinary Medicine, St. Paul, Minnesota 55108, USA

[4]Institute for Health Informatics, University of Minnesota, Minneapolis, Minnesota 55455, USA

[5]Imanis Life Sciences, LLC, Rochester MN

[6]Department of Molecular Medicine, Mayo Clinic, 200 First street SW, Rochester, MN, 55905

[7]Department of Veterinary Population Medicine, St. Paul, Minnesota 55108, USA

[8]Department of Pediatrics, University of Minnesota School of Medicine, Minneapolis, Minnesota 55455, USA

[9]Veterinary Diagnostic Laboratory, University of Minnesota College of Veterinary Medicine, St. Paul, Minnesota 55108, USA

[10]Center for Immunology, University of Minnesota, Minneapolis, Minnesota 55455, USA

[11]Department of Laboratory Medicine and Pathology, University of Minnesota Medical School, Minneapolis, Minnesota 55455, USA

[12]University of Alabama at Birmingham, Birmingham, AL, USA

[13]Division of Biostatistics, University of Minnesota Medical School, Minneapolis, Minnesota 55455, USA

[14]Vyriad, Inc. Rochester MN, 2900 37th St NW, Rochester MN 55901

[15]Stem Cell Institute, University of Minnesota, Minneapolis, Minnesota 55455, USA

[16]Center for Engineering and Medicine, University of Minnesota, Minneapolis, Minnesota 55455, USA

Dr. Tabaran's present address is: Department of Pathology, University of Agricultural Science and Veterinary Medicine, Cluj- Napoca, Romania.

Dr. Cornax's present address is: The Janssen Pharmaceutical Companies of Johnson & Johnson, San Diego, California, United States.

Mr. Schwabenlander's present address is Department of Veterinary and Biological Sciences, University of Minnesota College of Veterinary Medicine, St. Paul, Minnesota 55108, USA.

Abstract

VSV-IFNβ-NIS is a recombinant oncolytic vesicular stomatitis virus (VSV) that is being evaluated clinically for the treatment of advanced malignancies. As with other cancer immunotherapies, identifying biomarkers of response will be critical to the clinical advancement of this treatment approach. Here we describe the first evaluation of neoadjuvant intravenous oncolytic VSV therapy in naturally occurring cancer, specifically appendicular osteosarcoma in companion dogs, a disease with similar natural history as its human counterpart. VSV-IFNβ-NIS was administered prior to standard of care surgical resection, permitting microscopic and genomic analysis of tumors pre- and post-treatment. Alterations in the tumor microenvironment (micronecrosis, fibrosis, and inflammation) were more pronounced in VSV-treated dogs than in placebo-treated dogs. A "tail" of seven long-term survivors (35%) was apparent in the VSV-treated group. RNAseq analysis showed that virtually all the long-term responders had increased expression of a CD8 T-cell anchored immune gene cluster. We conclude that neoadjuvant VSV-IFNβ-NIS has an excellent safety profile and may provide a survival benefit for dogs with osteosarcoma whose tumors are permissive for immune infiltration. These data support ongoing translation of neoadjuvant VSV-IFNβ-NIS to human cancer patients. Strategies to further enhance clinical benefit include dose escalation or combination with other immunomodulatory agents.

Introduction

Osteosarcoma is a devastating bone cancer that primarily affects children and young adults, occurring most commonly in the long bones of the limbs.[1] Standard treatment for patients diagnosed with localized osteosarcoma is definitive tumor resection with limb-sparing surgery elected where possible, combined with multi-agent chemotherapy administered in the neoadjuvant and adjuvant settings. These protocols were developed over 40 years ago, achieving a 5-year survival rate of approximately 60%.[1,2] Patients whose tumors are refractory to frontline treatment and those who develop metastatic disease have a very poor prognosis.[3] Treatment modifications tested to date have failed to have a major impact on clinical outcomes. Thus, there is an unmet clinical need for treatments that improve outcomes for osteosarcoma patients. Development of new treatments is hampered by the rarity of this malignancy. Naturally occurring canine cancer represents a clinically relevant model that can recapitulate the spontaneous development and heterogeneity present in human cancer.[4] Osteosarcoma is a common malignancy in dogs with similar clinical presentation and natural history as in humans. Osteosarcoma treatment in dogs includes surgical amputation of the affected limb and adjuvant chemotherapy.[5] Also as in humans, most dogs develop pulmonary metastases and succumb to the disease; however, in dogs this typically occurs in less than 1 year.[6]

Vesicular stomatitis virus (VSV) expressing interferon-beta (IFNβ) and the sodium iodide symporter (NIS), referred to henceforth as VSV-IFNβ-NIS, is a clinical-stage recombinant oncolytic virus that is currently being tested as a systemic treatment for patients with advanced solid tumors and hematologic malignancies both as a monotherapy and in combination with immune checkpoint inhibitors.[7,8] Preclinical studies showing potent efficacy of systemic VSV-IFNβ-NIS therapy in murine tumor models led to translational studies in dogs with naturally occurring cancer.[9–11] These studies demonstrated that intravenous VSV-IFNβ-NIS therapy is well tolerated in cancer-bearing dogs with clinical responses observed in some dogs with

lymphoma, and stable disease in a dog with metastatic osteosarcoma.[12] Building on these preliminary findings, we launched the VIGOR study (VSV Immunotherapy and Genomics of Osteosarcoma Research). The aims of the VIGOR study were to test the safety, efficacy, and immunomodulatory effects of neoadjuvant intravenous VSV-IFNβ-NIS therapy in dogs with localized appendicular osteosarcoma. We hypothesized that neoadjuvant VSV-IFNβ-NIS would have an excellent safety profile and provide a survival benefit for dogs with osteosarcoma.

Results

Oncolytic VSV kills canine osteosarcoma cells *in vitro*

Oncolytic VSV has been shown to infect, replicate within, and kill a variety of tumor cell lines. To confirm the oncolytic activity of in canine osteosarcoma cells, we tested the ability of recombinant VSV vectors, including VSV-GFP, VSV-hIFNβ-NIS, and VSV-cIFNβ-NIS, to infect three canine osteosarcoma cell lines (OSCA-78, OSCA-08, and OSCA-40) *in vitro*. The VSV vectors replicated in all three canine osteosarcoma cell lines tested resulting in decreased viability between 24 to 48 hours post infection (Figure 1). VSV expressing canine or human IFNβ was able to replicate and kill canine osteosarcoma cell lines with similar potency. These data confirm that canine osteosarcoma cell lines are susceptible to VSV infection, replication, and oncolysis.

Screening and enrollment to the VIGOR study

The VIGOR (**V**SV **I**mmunotherapy and **G**enomics for **O**steosarcoma **R**esearch) study outline is shown in Figure 2. A total of 41 dogs were fully screened for enrollment in the VIGOR study of the 144 inquiries received (Supplemental Figure S1). Of the dogs screened, 13 were excluded. Twelve of these did not meet eligibility criteria due to documentation of metastatic disease on screening diagnostics (n=4), clinical suspicion or diagnosis of a different cancer (n=4), evidence of a pathologic fracture (n=3), or concurrent metastatic disease and pathologic fracture (n=1). One screened dog was not enrolled because the owner declined to participate. The study enrolled a total of 28 dogs over a period of 30 months (June 2016 to January 2019) with demographic characteristics of the dogs enrolled in the VIGOR study, as well as the comparison populations, shown in Table 1. All dogs had an initial diagnosis of sarcoma of an appendicular bone based on a pre-treatment bone biopsy. The first 15 dogs received a single dose of

neoadjuvant VSV-cIFNβ-NIS under an open label design. After the safety of systemic

neoadjuvant VSV-cIFNβ-NIS was documented, the next 13 dogs were randomized in a double-

blinded fashion to receive a single dose of either intravenous VSV-cIFNβ-NIS (n=7) or placebo

(PBS, n=6). All dogs were subsequently treated with standard of care and underwent

amputation of the affected limb and removal of the associated lymph node(s) 10 days after

receiving VSV-cIFNβ-NIS or placebo. Faxitron imaging was performed on the amputated limb to

guide sample collection (Supplemental Figure S2). Pathological analysis of resected tumors

confirmed primary osteosarcoma in 26 dogs, but the diagnoses in two dogs were subsequently

revised: one (treated with VSV in the open label portion of the study) was diagnosed with

intramedullary hemangiosarcoma at amputation, and the other (randomized to receive placebo)

was diagnosed with intramedullary rhabdomyosarcoma at necropsy.

VSV-cIFNB-NIS treatment is safe and is associated with prolonged survival in a subset of dogs with non-metastatic appendicular osteosarcoma

No clinically significant laboratory abnormalities were observed following VSV administration.

About half the treated dogs developed a transient fever (> 1.5°C increase in body temperature)

that resolved without intervention within 72 hours post infusion (not shown). Transient

hepatotoxicity and lymphopenia are commonly observed following intravenous VSV infusion.[7,12]

As expected, mild, transient elevation in liver enzymes and lymphopenia that resolved

spontaneously was observed in treated dogs (Supplemental Figure 3). There were no severe

adverse events (AEs) directly attributable to VSV-cIFNβ-NIS. Three dogs that were treated with

VSV (two open-label and one randomized to receive VSV) had reportable AEs that were

considered independent from the expected side effects of chemotherapy (Supplemental Table

S1), including one case each of pneumonia, facial nerve paralysis, and hypovolemic shock after

amputation. The latter were post-surgical complications that resulted in death of the dog and likely not related to VSV-cIFNβ-NIS administration.

Clinical efficacy endpoints evaluated included event free survival (EFS) and overall survival. Overall survival for dogs in the VIGOR study roughly segregated into 3 populations with a "tail" of 7 dogs that were "long-term" survivors (defined as > 479 days, greater than the 75th percentile of EFS in the COTC cohort). The group of long-term survivors includes 4 dogs that are still alive at the time of this report including three dogs with osteosarcoma (two treated with VSV and one randomized to receive placebo) and one with hemangiosarcoma of bone (treated with VSV). Three dogs were excluded from the survival analysis: two dogs that eventually received a histopathologic diagnosis other than osteosarcoma (one rhabdomyosarcoma of bone (placebo-treated) and one hemangiosarcoma of bone (VSV-treated)) and one dog that died immediately after surgery due to hypovolemic shock (VSV-treated), yielding a cohort of 20 evaluable VSV-cIFNβ-NIS treated dogs. The clinical information, treatment categories, and survival data for each individual dog enrolled in VIGOR are shown in Supplementary Table S2. EFS and overall survival for these evaluable dogs were compared to two contemporary control cohorts of dogs with appendicular osteosarcoma with no evidence of metastasis. The first cohort (n = 57) included dogs seen at the U of M VMC between July 2011 and July 2018, where there was intent to treat with standard-of-care surgery and adjuvant carboplatin chemotherapy, and that had a successful limb amputation and completed at least one cycle of adjuvant chemotherapy. The second control cohort was from a study recently published by the National Cancer Institute's Comparative Oncology Trial Consortium (NCI-COTC) and included 157 dogs enrolled between November 2015 and February 2018 that were treated with standard-of-care surgery and adjuvant carboplatin chemotherapy.[13] The NCI-COTC cohort was made available as a control dataset for evaluation of novel neoadjuvant or adjuvant treatments for osteosarcoma. Comparison to control cohorts showed that neoadjuvant VSV-cIFNβ-NIS therapy did not markedly improve EFS or

overall survival (Figure 3A), but importantly did not worsen survival outcomes. Long-term survivorship of evaluable VSV-cIFNβ-NIS treated dogs was compared by tabulating the proportion of dogs with survival that exceeded the 75th percentile value of the NCI-COTC cohort (479 days). The U of M VMC control cohort had, as expected, 26% of dogs that exceeded an overall survival of 479 days. The VIGOR cohort had a higher-than-expected proportion of long-term survivorship with 35% of dogs exceeding an overall survival of 479 days (Figure 3B), though the sample size lacked the power to establish that this difference was not due to random chance with a high level of confidence.

Histopathological assessment of osteosarcoma tumors shows an increase in tumor inflammation following VSV treatment

 Pre-treatment tumor biopsies and post-treatment amputation-resected tumor samples were scored based on tumor necrosis, inflammation, and fibrosis (Supplemental Table S3). Tumor histopathologic evaluation showed areas of moderate to severe ischemic necrosis, as is typically observed in rapidly growing osteosarcoma lesions, in both VSV-cIFNβ-NIS-treated and placebo-treated cases; however, the tumors of 10 of 22 VSV-cIFNβ-NIS-treated dogs had distinctive areas of micronecrosis that were not consistently present in placebo-treated tumors (Supplemental Figure S4). Inflammatory infiltrates were scored in pre-treatment and post-treatment tumor specimens. Higher tumor infiltration score (TIS) was observed in post-treatment tumor specimens from VSV treated dogs compared to pre-treatment tumor biopsies, while similar increases were not observed in tumor specimens from placebo treated dogs (Figure 4). Matched pre- vs post-treatment tumor specimens were available for assessment of intratumoral inflammatory infiltrate for half of the enrolled dogs showing a significant increase in TIS in post-treatment tumors compared to baseline (P=0.0027). These results are encouraging but are interpreted with caution given the small cohort size and the small number of dogs enrolled in the placebo group.

Regional lymph nodes were evaluated in 27/28 dogs; the associated lymph node could not be found after amputation in one case (from a dog randomized to receive VSV). Two of the 27 cases evaluated had evidence of osteosarcoma metastasis in the lymph node, and the remaining 25 were negative for metastasis. One of the two dogs with lymph node metastasis at the time of amputation had shortened survival and was euthanized 64 days after surgery, with metastasis present throughout the abdomen and thorax on necropsy. Interestingly, the other dog with lymph node metastasis had prolonged survival and was still alive at the time of this report. Both cases with documented lymph node metastasis were treated with VSV.

Virus pharmacokinetics

Monitoring viremia, virus shedding, and antiviral antibodies showed similar trends to previous studies evaluating intravenous VSV therapy in dogs.[12] Viral RNA was detectable in whole blood samples at high quantities for the first 24 hours following infusion, with virus localizing primarily to PBMCs (Figure 5). Decay of viral RNA in blood coincided with an increase in detection of anti-VSV neutralizing antibodies, indicative of antibody-mediated virus clearance (Supplemental Figure S5). Infectious virus was detectable in PBMC samples at low quantities (below the limit of quantification) only at 1 hour post systemic VSV administration and no infectious virus was detected in PBMC samples collected after the 1h timepoint (data not shown). Virus shedding studies showed no detectable infectious virus in urine, rectal swab, or buccal swab samples, though viral RNA was detectable at low levels in some samples as shown (Supplemental Figure S6). Viremia, virus shedding and neutralizing antibodies were measured in samples from two placebo-treated dogs showing no detectable viral RNA in blood or shedding samples, infectious virus, or antiviral antibodies providing relevant negative controls in our pharmacokinetic analyses (data not shown). The dose dependent efficacy of systemic VSV therapy,[12] the low amount and brief period that infectious virus was detectable, the absence of infectious virus

shedding, and the excellent safety profile all suggest higher doses of VSV-cIFNβ-NIS can be safely administered in this treatment setting to potentially improve clinical response.

Pre-treatment immune infiltration is associated with longer overall survival

Acute cytokine responses were measured in the first 15 dogs enrolled and showed a transient elevation in several pro-inflammatory cytokines between 3 and 6 hours following VSV administration (Supplemental Figure S7). Three dogs that were long-term responders to VSV treatment, including the one diagnosed with hemangiosarcoma of bone, had high GM-CSF, IL-2, IL-7, and IL-15 levels at baseline.

Samples were examined for evidence of an immune component in the tumor response using next-generation RNA sequencing. Specifically, we used the gene cluster expression summary score (GCESS) method to summarize co-regulated gene clusters in tumor samples and their association with outcome.[13] Pre- and post-treatment samples from primary osteosarcoma and metastatic osteosarcoma tumors grouped together, separate from both the cell lines derived from the pre-treatment tumors as well as from normal skin biopsy samples obtained from the same dogs at the time of amputation (Figure 6A). Fourteen gene clusters were identified and were apparent in unsupervised hierarchical clustering heatmaps (Figure 6B), including the immune and cell cycle gene clusters previously identified in canine and human osteosarcoma.[14] The distributions of cell cycle and immune GCESS across tumor samples, cell lines, and skin biopsy samples were as predicted, with no immune transcripts identified in osteosarcoma cell lines. VSV treatment did not notably impact the GCESS score of pre- versus post-treatment tumor specimens (Figure 6C). When we analyzed the association between GCESS and survival, we observed that the dogs with the highest pre-treatment CD8-related immune GCESS had longer survival times (Figure 7). This relationship was not observed for the CD37-positive monocyte-related GCESS, and somewhat surprisingly, it was also not evident for the cell cycle

GCESS (Figure 7B). Upon comparison of the VIGOR cohort to this CCOGC control cohort with available RNAseq data (n=23, Supplemental Figure S8), we noted that a higher immune GCESS was associated with a longer progression-free survival in both groups, but that this effect was larger in the VSV-treated group. In the CCOGC cohort, all the dogs with longer survival had a higher CD8 GCESS, but not all the dogs with higher CD8 GCESS had longer survival (Supplemental Figure S8C – see overlapping box plots); in the VIGOR group, all of the dogs with higher CD8 GCESS had longer survival (Figure 7B – see non-overlapping VSV high and VSV low box plots). The Kaplan-Meier Survival curve, showing increased overall survival in the VSV-treated dogs in the VIGOR group when compared with CCOGC cohort where RNA sequencing was available, is shown in Supplemental Figure S9.

Discussion

Oncolytic viruses, including VSV, can selectively infect, replicate within and kill tumor cells, with the potential to promote release of antigens specific to an individual tumor to drive robust anti-tumor immune responses.[15,16] This approach is of particular interest in osteosarcoma, a genomically chaotic cancer, albeit with a low overall mutational burden and where recurrent mutations across the patient population are uncommon.[17–19] The use of oncolytic viruses, therefore, can be used to promote the release of antigens specific to the individual tumor.[15] Clinical studies evaluating oncolytic viruses to date have limited evaluation of the immune responses following treatment and highlight the need to identify and investigate biomarkers of clinical response. The use of oncolytic viruses, therefore, can be used to promote the release of antigens specific to the individual tumor.[15] Here we describe the safety and efficacy of intravenous VSV-cIFNβ-NIS therapy in naturally occurring canine appendicular osteosarcoma, a model utilized to inform development of new treatments for human osteosarcoma.[5] Studies in spontaneous canine cancer supported clinical translation of VSV-IFNβ-NIS, a clinical-stage

oncolytic virus that is being evaluated for the treatment of both solid tumors and hematologic malignancies.[7,8] The major goals of this study were to establish the safety of VSV-IFNβ-NIS in dogs with osteosarcoma, and to utilize correlatives studies in this spontaneous model to identify and evaluate potential determinants of efficacy to oncolytic VSV therapy.

In this study, we documented the safety profile of systemic therapy with VSV-cIFNβ-NIS in canine osteosarcoma. Intravenous VSV-cIFNβ-NIS, administered at 1×10^9 $TCID_{50}$/kg, was remarkably well-tolerated, with no clinically significant changes in any of the monitored parameters. Systemic viral distribution was documented, as live virus was detected (below the limit of quantification) in PBMCs within one-hour post-infusion, but not at later timepoints. Evidence of a systemic response was noted, as a subset of dogs examined showed acute elevations of inflammatory cytokines. No clinically significant adverse events were documented that were attributable to VSV therapy. The low quantities and short duration of infectious virus detected in blood, paired with the absence of adverse events, suggests there is scope for increasing the systemic dose to potentially improve clinical outcomes.

We evaluated changes in GCESS and tumor inflammation scores on histopathology before and after treatment with VSV-cIFNβ-NIS in this study. An increased tumor inflammation score (based on pathology assessment) was noted in tumor samples collected 10 days post treatment compared to baseline samples. Clinical outcomes with neoadjuvant VSV therapy added to standard treatments for dogs with osteosarcoma were compared with two published control cohorts with dogs treated with only standard of care, including the NCI-COTC osteosarcoma cohort, a prospectively enrolled cohort designed to enable assessment and comparison of new osteosarcoma therapies in this clinically relevant model.[12] While we did not observe a major difference in survival between the groups, there was a tail of long-term survivors noted in each population. In the VIGOR cohort, 35% of the dogs were considered long-term survivors, as they

had survival times exceeding 479 days, the 75th percentile overall survival from the NCI-COTC osteosarcoma cohort. Additionally, when we evaluated the association between immune GCESS and survival in both the VIGOR population and the CCOGC comparison group that had available RNAseq data, a higher CD8 immune GCESS was associated with prolonged survival in both groups. However, this effect was more pronounced in the VSV-treated dogs. Our interpretation of these data is that the presence of a CD8-cell signature in osteosarcoma tumors prior to treatment increases the probability of longer survival, and the addition of VSV in the neoadjuvant setting enhances this effect.

In conclusion, these results show that systemically administered VSV-IFNβ-NIS can be used safely to treat dogs with spontaneous osteosarcoma. Additionally, VSV-IFNβ-NIS is biologically active, and can target the tumor environment. The intrinsic tumor immune status at baseline appears to influence the activity and therapeutic benefit of VSV, as dogs with higher immune GCESS at baseline had prolonged survivals. The clinical safety and PK data, when viewed in the context of previous findings indicating dose dependent efficacy of systemic VSV therapy,[20] suggest that there is scope to increase the systemic dose to improve clinical outcomes. Additionally, the observed therapeutic benefit in dogs with a pro-immune environment suggests that combinations with agents that promote immune infiltration or that amplify the immune response, such as immune checkpoint inhibitors, may further enhance the therapeutic benefits of VSV-IFNβ-NIS. The observed exceptional safety and immunological activity of VSV-IFNβ-NIS support further translation to human cancer patients.

Materials & Methods

Manufacturing of VSV

Vesicular stomatitis virus (VSV) expressing canine interferon-beta (IFNβ) and the sodium iodide symporter (NIS), VSV-cIFNβ-NIS, was constructed as previously described.[10] The VSV-cIFNβ-NIS lot used in veterinary studies was provided by the Mayo Clinic Viral Vector Production laboratory (VVPL). Virus product is stored at ≤ -65°C. Virus titers were determined by 50% tissue culture infective dose ($TCID_{50}$) assays on Vero cells. The virus product was tested to confirm sterility (Mayo Clinic Department of Laboratory Medicine and Pathology) and absence of endotoxin using the LAL-Kinetic QCL Kit (Lonza). Whole genome sequencing was performed by utilizing reverse transcription of purified viral RNA to cDNA, polymerase chain reaction (PCR) of overlapping regions of viral genome, sequencing of PCR fragments (Eurofins Genomics), and assembly of sequencing results.

In vitro efficacy of oncolytic VSV in canine osteosarcoma

Recombinant VSV vectors, including VSV expressing green fluorescent protein (GFP) or VSV expressing canine or human IFNβ and NIS (hIFNβ-NIS or cIFNβ-NIS), were used to infect three canine osteosarcoma cell lines, OSCA-78, OSCA-08, and OSCA-40, to evaluate *in vitro* efficacy of oncolytic VSV in canine osteosarcoma. Cells were grown in DMEM media containing 5% glucose and L-glutamine supplemented with 10% fetal bovine serum (FBS), 10mM HEPES buffer, and 0.1% Primocin, and cultured at 37°C in a humidified atmosphere of 5% CO_2. Cell lines were authenticated based on short tandem repeats at regular intervals (IDEXX BioAnalytics). Virus titers were determined by $TCID_{50}$ assay on BHK-21 cells at several timepoints after infection (12h, 24h, 36h, 48h, 60h, and 72h). Cell viability, as a percent of control, was determined by flow cytometry using a live/dead marker (Invitrogen) at several

timepoints after infection (12h, 24h, 36h, 48h, 72h, and 96h). Cell lines are available through Kerafast, Inc (Boston, MA).

Enrollment and eligibility criteria

Dogs with a diagnosis of appendicular bone sarcoma, based on radiographic appearance of the lesion and results of bone aspirate or biopsy, were considered for enrollment in the study (see CONSORT diagram, Supplementary Figure S1). All cases were enrolled at the University of Minnesota (UMN) College of Veterinary Medicine between June 2016 and January 2019. Screening diagnostics were performed prior to inclusion in the study, including physical examination, radiographs of the affected limb and thorax, general blood work (complete blood count, serum chemistry profile), computed tomography of the thorax and abdomen, and biopsy of the bone lesion. Inclusion criteria included cytologic or histologic diagnosis of sarcoma in an appendicular bone and body weight greater than 20kg (to facilitate safe collection of blood for the study). Exclusion criteria included evidence of metastatic lesions, evidence of pathologic fracture, intact reproductive status, clinically significant co-morbidities, treatment with other medications, and residence on a farm or other contact with farm animals (to reduce risk of VSV transmission to susceptible hosts).

VSV dosing

Previous studies have demonstrated the safety of intravenous VSV-hIFNβ-NIS and VSV-cIFNβ-NIS therapy in dogs with cancer at doses up to 10^{10} $TCID_{50}/0.5m^2$, which converts to approximately 1×10^9/kg.[12,21] The VIGOR study enrolled a total of 28 dogs over 30 months (June 2016 to January 2019). The first 15 dogs received a single dose (1×10^9 $TCID_{50}$/kg) of neoadjuvant VSV-cIFNβ-NIS under an open label design. Because VSV at this dose was not

associated with clinically significant adverse events and a preliminary efficacy signal was noted, the next 13 dogs were randomized in a double-blinded fashion to receive a single dose of either intravenous VSV-cIFNβ-NIS or placebo (phosphate buffered saline solution (PBS)). The virus was diluted in sterile PBS to a volume of 10.5ml, which was administered by intravenous infusion over a period of 2-5 minutes. Dogs were housed for up to 24 hours in a BSL-2 housing facility. All procedures were carried out with approval of the University of Minnesota Institutional Animal Care and Use Committee (IACUC #1504-32486A and #1803-35759A) and Institutional Biosafety Committee (IBC #1504-32487H and # 1805-35967H).

Standard of care therapy

All enrolled dogs were treated with standard of care therapy after treatment with intravenous VSV-cIFNβ-NIS or placebo. This consisted of amputation of the affected limb and removal of the associated lymph node(s), which occurred at day 10 following treatment, and initiation of chemotherapy approximately 14 days after amputation. Chemotherapy consisted of six cycles of carboplatin at a dose of 300 mg/m^2, administered intravenously every 3 weeks, with standard monitoring for adverse chemotherapy effects.

Biopsy procedures

Pre-treatment biopsies were obtained from the lesion using 4mm Michele trephine bone biopsy instruments. One pre-treatment tumor biopsy section was placed in 10% neutral buffered formalin for histopathology. A second pre-treatment section was further divided, with one piece used to establish a cancer cell line and a second piece frozen at -80°C for later RNA extraction. Ten days post-treatment with VSV-cIFNβ-NIS, dogs underwent surgical amputation of the affected limb and removal of the associated lymph node(s). A skin punch biopsy was performed,

and the tissue frozen at -80°C for later RNA extraction. After amputation, the soft tissues of the limb were removed, and imaging of the affected bone was performed using a Faxitron (Supplementary Figure S2). The amputated limbs were sectioned longitudinally and tumor sections of approximately 5mm thickness were obtained and placed in 10% neutral buffered formalin. Tumor sections were placed in 10% ethanol for slow decalcification, or in acid for rapid decalcification. Separate sections of the tumor were frozen at -80°C for later RNA extraction. Cell lines derived as part of this study can be made available to the scientific community for research purposes under materials transfer agreements negotiated by the Regents of the University of Minnesota and Mayo Clinic.

Monitoring virus pharmacokinetics

Assays to monitor viremia, virus shedding, and antiviral antibodies following intravenous VSV infusion were performed by Imanis Life Sciences using previously described protocols.[12,21] Briefly, viremia was monitored by detection of infectious virus and viral RNA in blood samples collected at baseline and at specific time points after VSV treatment. For detection of viral RNA, whole blood was collected in RNAprotect animal blood tubes (Qiagen). For detection of infectious virus, peripheral blood mononuclear cells (PBMCs) were isolated from whole blood. Urine samples, fecal samples, and buccal mucosal swabs were collected after treatment with VSV-cIFNβ-NIS to test for presence of live infectious virus and viral RNA. Detection of infectious virus was performed by overlay of sample supernatants on susceptible BHK-21 cells. Viral RNA was detected in shedding and blood samples by quantitative reverse transcription-polymerase chain reaction (qRT-PCR) as previously described. Tumor specimens were stored at < -65°C at the time of surgical tumor resection (10 days post VSV therapy) for subsequent processing to isolate RNA and detect VSV RNA by qRT-PCR. Serum samples were obtained at baseline, and 3, 7, 14, 21, and 28 days post VSV infusion for detection of anti-VSV neutralizing antibodies.

Serum cytokine monitoring

Serum samples were collected at baseline, 1, 3, 6 hours and 1-, 3-, 7- and 14-days post VSV infusion. Acute cytokine responses were determined as previously described by the COTC Pharmacodynamic Core at Colorado State University (CSU).[22] The acute cytokines evaluated were: GM-CSF, IL-6, MCP-1, KC, IL-2, IP-10, IL-8, TNF-alpha, IL-10, IL-7, IL-1, and IL-18, all of the canine species.

Clinical monitoring

Body temperature was monitored after treatment with VSV-cIFNβ-NIS, every 3 hours for the first 24 hours following treatment, then daily. Blood work to evaluate alanine aminotransferase (ALT), aspartate aminotransferase (AST), total white blood cell count, lymphocyte count, and clotting times (PT and aPTT) was completed at baseline, and at days 3, 7, 14, 21, and 28 after treatment.

Histopathology

Pre-treatment core tumor biopsies and post-treatment amputation-resected tumor samples were evaluated with H&E staining to obtain a definitive histologic diagnosis. Samples were also scored based on tumor inflammation, fibrosis, and necrosis. Inflammation was graded on a 4-point scale, where 0 = absence of inflammation, 1 = minimal inflammation, 2 = mild inflammation, 3 = moderate inflammation, and 4 = marked inflammation. Fibrosis was graded on a 4-point scale, where 0 = absence of fibrosis, 1 = minimal fibrosis, 2 = mild fibrosis, 3 = moderate fibrosis, and 4 = marked fibrosis. Necrosis was graded on a 3-point scale, where 0 = no necrosis, 1 = < 15%, 2 = 15 - 50%, and 3 = > 50% of the section had apparent necrosis. The

draining lymph node was evaluated for presence or absence of metastatic osteosarcoma cells. During the placebo-arm of the study, the pathologist was blinded to the treatment assignment of cases (VSV or placebo).

Collection of PBMCs for immune monitoring

Blood samples were obtained for isolation of PBMCs and plasma for immune monitoring. Blood was collected in a commercially available tube for PBMC isolation (BD Mononuclear Cell Preparation Tube), and the manufacturer's protocol was followed. PBMCs were frozen in FBS with 10% DMSO and stored in liquid nitrogen until use.

Diagnostic imaging

Cases were monitored with diagnostic imaging to evaluate for metastatic disease throughout the study period. Thoracic radiographs were performed at study evaluation (d-1), and on days 60, 120, 180, and 360 post-treatment. Computed tomography (CT) was performed at study evaluation (d-1) and at day 90 post-treatment.

RNA extraction from tumor specimens

VIGOR: RNA was extracted from tumor tissue at baseline, and from tumor tissue samples obtained at amputation, post-treatment. RNA was also extracted from cell lines that were established pre-treatment from each case, and from normal skin tissue from each case as a non-malignant control. When available, RNA was also extracted from lung metastases. Tissue samples were disrupted with a tissue homogenizer, and RNA was extracted using a

commercially available kit (QIAGEN RNeasy mini kit). RNA was stored at -80°C until submission to the University of Minnesota Genomics Center (UMGC).

CCOGC: A comparison population (n = 23), consisted of tissue samples provided by the Pfizer CCOGC Biospecimen Repository. Samples with a definitive, histological diagnosis of osteosarcoma were selected for RNA sequencing based on the availability of sufficient frozen tissue (>50 mg), treatment information, RNA integrity numbers (RIN) >6, and available follow-up information. These were included in the final selection for next-generation RNA sequencing.

Library preparation and next-generation sequencing

Unique dual-indexed sequencing libraries were prepared using the Clontech Stranded Total RNA-Seq Kit v2 - Pico Input Mammalian kit. RNA sequencing (2x50-bp paired-end, on a NovaSeq S2) was performed at the UMGC. Approximately 3,100M pass filter reads were generated for the run. Mean quality scores were above Q30 for all libraries. Successful libraries were generated from 64 samples (15 pre-treatment tumors, 20 post-treatment tumors, 18 cancer cell lines generated from pre-treatment tumors, and 11 skin samples) from 24 dogs. Additionally, two dogs had pulmonary metastatic lesions obtained at necropsy from which libraries were generated from tissue and from cell lines. A total of 23 osteosarcoma tissue samples were included in the CCOGC dataset.

Gene cluster expression summary score (GCESS) and Bioinformatics analysis

Initial quality control analysis of RNA sequencing FASTQ data was performed using FastQC software (v0.11.5). FASTQ data were trimmed with Trimmomatic (v0.33.0). Kallisto (v0.43.0) was used for pseudoalignment and quantifying transcript abundance. Sequencing reads were

aligned to the canine reference genome (CanFam3.1). Gene cluster expression summary score, or GCESS, analyses were performed as previously described.[14] Briefly, clusters of genes are identified based on patterns of similar expression, such as those associated with immune cell infiltration or cell cycling. After mean-centering and log_2-transformation, individual genes in each cluster are added together resulting in a single summary score for the cluster that is reflective of the overall degree of gene expression.

Outcome assessment

Formal clinical follow up, including physical examinations, blood work, and diagnostic imaging, occurred for one year after treatment with VSV-cIFNβ-NIS. Thereafter, informal follow up occurred until relapse or death. Necropsies were obtained when possible, with owner consent. Outcomes assessed included event-free survival (EFS, time to relapse) and overall survival (time to death). The dogs enrolled in the VIGOR study (n = 28) were compared with three control populations. The first population consisted of dogs with appendicular osteosarcoma treated at the same institution (UMN, n = 57) between January 2006 and December 2018. The second population consisted of the control arm of a Comparative Oncology Trials Consortium multicenter clinical trial in dogs with appendicular osteosarcoma (COTC, n = 157) enrolled between November 2015 and February 2018.[13] Both comparison populations were treated with standard of care therapy, consisting of amputation of the affected limb and carboplatin chemotherapy administered every 21 days at 300mg/m². A third comparison population (CCOGC, n = 23), consisted of dogs treated with standard of care (chemotherapy and amputation) with known outcome data, enrolled in a multicenter study between April 2008 and March 2011 for which samples were available for tumor next-generation RNA sequencing. This population was used for comparison of survival and tumor gene expression.

Statistical analyses

Tumor inflammation scores were compared between pre-treatment and post-treatment tumor samples using paired student's t-test. P-values are reported without inference to significance, as recommended by the American Statistical Association.[23]

Kaplan-Meier survival curves were generated for the dogs treated with VSV-cIFNβ-NIS in the VIGOR study and the comparison populations. The EFS was calculated as the time between limb amputation and the first detection of metastasis. Dogs were censored if they did not have metastatic disease at the time of last follow-up. Overall survival was calculated as the time between limb amputation and death. Dogs were censored from survival analyses if they were found to have tumors of bone that were not osteosarcoma, if they died of non-osteosarcoma related causes, or if they were alive at last follow up.

Author contributions

SN, SJR, KMS, and JFM were involved in overall study design and execution including attaining regulatory approvals and obtaining funding to implement the described clinical studies. AC, DG, KB, SP, AW, and KMM were responsible for sample collection and processing. AC, KMS, AB, and MSH were responsible for clinical follow-up. LS and SN oversaw and executed correlative pharmacokinetic studies (assays to measure viremia, virus shedding, antibodies, etc.). CP assisted in data processing and analysis. ALS and LJM analyzed sequencing data. MDS and MW completed methodological design for gross sample collection, imaging with the Faxitron, and gross lesion identification, assessment, and sampling for optimal downstream analysis. MW organized pathological images, maintained supplies, and facilitated communication regarding samples. IC, AFT, and MGOS were responsible for analyzing histopathology specimens. SN, ALS, LJM, JFM, and KMM were responsible for additional data analysis. GRC and JSK were

responsible for statistical design and analysis. SN, KMM, ALS, and JFM were responsible for data compilation and manuscript preparation. All authors were responsible for manuscript review and revision.

Acknowledgements

The authors would like to acknowledge Milcah Scott for assistance in training, including cell culture and sequencing techniques. The authors would like to acknowledge Mitzi Lewellen for oversight of sample databases. The authors would also like to acknowledge Katalin Kovacs and Paula Overn for their assistance in preparation of histology specimens. The authors acknowledge the Clinical Investigation Center (CIC) of the University of Minnesota College of Veterinary Medicine, the Veterinary Diagnostic Laboratory (VDL) of the University of Minnesota, the University of Minnesota Genomics Center (UMGC), and the Minnesota Supercomputing Institute (MSI) at the University of Minnesota for providing resources that contributed to the research results reported within this paper. The authors wish to thank the VDL Pathologists for their assistance in sample preparation, including Dr. Arno Wuenschmann, Dr. Erik Olson, and Dr. Jaclyn Dykstra. The authors also wish to thank the CCOGC, Inc. for providing a set of control tumor samples for this study, as well as Dr. Subbaya Subramanian for help with procuring the CCOGC samples. Additionally, the authors wish to thank Dr. Amy LeBlanc and Christina Mazcko from the NCI COTC for providing the raw survival data for the control cohort treated with the standard of care.

The authors wish to thank the clinical teams at the University of Minnesota College of Veterinary Medicine Lewis Small Animal Teaching Hospital for the clinical care of enrolled dogs, including Oncology, Soft Tissue Surgery, Diagnostic Imaging, the Veterinary Diagnostic Laboratory, Clinical Pathology, and associated support staff.

Funding

The work described in this manuscript was supported by Vyriad Pharma, by grant MNP #15.25 from the Minnesota Partnership for Biotechnology and Medical Genomics, grant R21 CA208529 from the National Cancer Institute (NCI) of the National Institutes of Health (NIH), United States Public Health Service, and grants CA170218 and CA190276 from the US Department of Defense Peer Reviewed Cancer Research Program. NCI Comprehensive Cancer Center Support Grant P30 CA077598, to the Masonic Cancer Center, University of Minnesota, provided support for genomics and bioinformatics. Tumor samples and relevant metadata from the CCOGC cohort were provided *pro bono* by the CCOGC, Inc. KMM was supported in part by institutional training grant in Molecular, Genetic, and Cellular Targets of Cancer T32 CA009138 from the NCI. ALS was supported in part by grant R50 CA211249 from the NCI. JFM is supported in part by the Alvin and June Perlman Endowed Chair in Animal Oncology. The authors acknowledge support from individual donors to the Animal Cancer Care and Research Program, University of Minnesota. The content of this manuscript is solely the responsibility of the authors and does not necessarily represent the official views of any of the funding agencies listed above.

References

1.   Mirabello L, Troisi RJ, Savage SA. Osteosarcoma incidence and survival rates from 1973 to 2004: data from the Surveillance, Epidemiology, and End Results Program. *Cancer*. 2009;115(7):1531-1543. doi:10.1002/cncr.24121

2.   Cole S, Gianferante DM, Zhu B, Mirabello L. Osteosarcoma: a Surveillance, Epidemiology, and End Results program-based analysis from 1975 to 2017. *Cancer*. n/a(n/a). doi:10.1002/cncr.34163

3.   Smeland S, Bielack SS, Whelan J, et al. Survival and prognosis with osteosarcoma: outcomes in more than 2000 patients in the EURAMOS-1 (European and American Osteosarcoma Study) cohort. *Eur J Cancer*. 2019;109:36-50. doi:10.1016/j.ejca.2018.11.027

4.   Weiss MC, Eulo V, Van Tine BA. Truly Man's Best Friend: Canine Cancers Drive Drug Repurposing in Osteosarcoma. *Clin Cancer Res*. 2022;28(4):571-572. doi:10.1158/1078-0432.CCR-21-3471

5.   Makielski KM, Mills LJ, Sarver AL, et al. Risk Factors for Development of Canine and Human Osteosarcoma: A Comparative Review. *Vet Sci*. 2019;6(2). doi:10.3390/vetsci6020048

6.   Poon AC, Matsuyama A, Mutsaers AJ. Recent and current clinical trials in canine appendicular osteosarcoma. *Can Vet J*. 2020;61(3):301-308.

7.   Cook J, Peng KW, Witzig TE, et al. Clinical activity of single-dose systemic oncolytic VSV virotherapy in patients with relapsed refractory T-cell lymphoma. *Blood Adv*. 2022;6(11):3268-3279. doi:10.1182/bloodadvances.2021006631

8.   Lutzky J, Marron TU, Powell SF, et al. Optimization of Voyager V1 (VV1) oncolytic virus systemic delivery in combination with cemiplimab and ipilimumab in patients with melanoma and non–small cell lung cancer (NSCLC). *JCO*. 2022;40(16_suppl):TPS9595-TPS9595. doi:10.1200/JCO.2022.40.16_suppl.TPS9595

9.   Naik S, Nace R, Barber GN, Russell SJ. Potent systemic therapy of multiple myeloma utilizing oncolytic vesicular stomatitis virus coding for interferon-β. *Cancer Gene Ther*. 2012;19(7):443-450. doi:10.1038/cgt.2012.14

10.   Naik S, Nace R, Federspiel MJ, Barber GN, Peng KW, Russell SJ. Curative one-shot systemic virotherapy in murine myeloma. *Leukemia*. 2012;26(8):1870-1878. doi:10.1038/leu.2012.70

11.   Zhang L, Steele MB, Jenks N, et al. Safety Studies in Tumor and Non-Tumor-Bearing Mice in Support of Clinical Trials Using Oncolytic VSV-IFNβ-NIS. *Hum Gene Ther Clin Dev*. 2016;27(3):111-122. doi:10.1089/humc.2016.061

12.   Naik S, Galyon GD, Jenks NJ, et al. Comparative Oncology Evaluation of Intravenous Recombinant Oncolytic Vesicular Stomatitis Virus Therapy in Spontaneous Canine Cancer. *Mol Cancer Ther*. 2018;17(1):316-326. doi:10.1158/1535-7163.MCT-17-0432

13.   LeBlanc AK, Mazcko CN, Cherukuri A, et al. Adjuvant Sirolimus Does Not Improve Outcome in Pet Dogs Receiving Standard-of-Care Therapy for Appendicular Osteosarcoma: A Prospective, Randomized Trial of 324 Dogs. *Clin Cancer Res*. 2021;27(11):3005-3016. doi:10.1158/1078-0432.CCR-21-0315

14.   Scott MC, Temiz NA, Sarver AE, et al. Comparative Transcriptome Analysis Quantifies Immune Cell Transcript Levels, Metastatic Progression, and Survival in Osteosarcoma. *Cancer Res*. 2018;78(2):326-337. doi:10.1158/0008-5472.CAN-17-0576

15.   Russell SJ, Barber GN. Oncolytic Viruses as Antigen-Agnostic Cancer Vaccines. *Cancer Cell*. 2018;33(4):599-605. doi:10.1016/j.ccell.2018.03.011

16.   Durham NM, Mulgrew K, McGlinchey K, et al. Oncolytic VSV Primes Differential Responses to Immuno-oncology Therapy. *Mol Ther*. 2017;25(8):1917-1932. doi:10.1016/j.ymthe.2017.05.006

17.   Xie L, Yang Y, Guo W, et al. The Clinical Implications of Tumor Mutational Burden in Osteosarcoma. *Front Oncol*. 2020;10:595527. doi:10.3389/fonc.2020.595527

18.   Sayles LC, Breese MR, Koehne AL, et al. Genome-Informed Targeted Therapy for Osteosarcoma. *Cancer Discov*. 2019;9(1):46-63. doi:10.1158/2159-8290.CD-17-1152

19.   Sakthikumar S, Elvers I, Kim J, et al. SETD2 Is Recurrently Mutated in Whole-Exome Sequenced Canine Osteosarcoma. *Cancer Res*. 2018;78(13):3421-3431. doi:10.1158/0008-5472.CAN-17-3558

20.   Bailey K, Kirk A, Naik S, et al. Mathematical model for radial expansion and conflation of intratumoral infectious centers predicts curative oncolytic virotherapy parameters. *PLoS One*. 2013;8(9):e73759. doi:10.1371/journal.pone.0073759

21.   LeBlanc AK, Naik S, Galyon GD, et al. Safety studies on intravenous administration of oncolytic recombinant vesicular stomatitis virus in purpose-bred beagle dogs. *Hum Gene Ther Clin Dev*. 2013;24(4):174-181. doi:10.1089/humc.2013.165

22.   Paoloni M, Mazcko C, Selting K, et al. Defining the Pharmacodynamic Profile and Therapeutic Index of NHS-IL12 Immunocytokine in Dogs with Malignant Melanoma. *PLoS One*. 2015;10(6):e0129954. doi:10.1371/journal.pone.0129954

23.   Wasserstein RL, Lazar NA. The ASA Statement on p-Values: Context, Process, and Purpose. *The American Statistician*. 2016;70(2):129-133. doi:10.1080/00031305.2016.1154108

Figure 1

Figure 1. Vesicular stomatitis virus replicates in and kills three established canine osteosarcoma cell lines.

Virus titers and cell viability following infection of canine osteosarcoma cell lines: **A.** OSCA-8, **B.** OSCA-40, and **C.** OSCA-78 with recombinant VSVs: VSV-GFP (green), VSV-hIFNβ-NIS (blue), or VSV-cIFNβ-NIS (orange) at an MOI of 0.03.

Figure 2

Figure 2. Design schematic of the VIGOR study, evaluating systemic vesicular stomatitis virus (VSV) in dogs with spontaneous appendicular osteosarcoma.

Tumor samples were obtained pre- and post-treatment with VSV for histopathology. Diagnostic imaging was performed at most visits to evaluate for pulmonary metastases, with either computed tomography (CT) or thoracic radiographs (CXR). Chemotherapy, consisting of intravenous carboplatin every 3 weeks for 6 doses, was started at day 21 after VSV treatment. At each visit, peripheral blood mononuclear cell samples were also obtained for immune monitoring.

Table 1

		VIGOR (n = 28)	UMN (n = 57)	COTC (n = 157)
Breed n (%)	Golden retriever	5 (17.9)	12 (21.2)	7 (4.5)
	Labrador retriever	3 (10.7)	7 (12.3)	23 (14.6)
	Mastiff	2 (7.1) *	2 (3.5)	4 (2.5)
	Rottweiler	1 (3.6)	2 (3.5)	12 (7.6)
	Great Dane	1 (3.6)	3 (5.3)	12 (7.6)
	Greyhound	1 (3.6)	1 (1.8)	10 (6.4)
	Great Pyrenees	1 (3.6)	0 (0)	8 (5.1)
	Other purebreed	6 (21.4)	15 (26.3)	40 (30.6)
	Mixed breed	8 (28.6) **	15 (26.3)	33 (21.0)
Age (years)	Mean (± SD)	7.7 (± 2.5)	8.8 (± 2.5)	8.1 (± 2.4)
	Median	8.1	9.0	8.3
	Range	2.1 - 11.7	2.0 - 16.0	(1.4 - 15.6)
Body weight (kg)	Mean (± SD)	38.3 (± 15.5)	37.9 (± 11.3)	41.2 (± 12.3)
	Median	32.7	37.1	38.8
	Range	20 - 81.6	15.3 - 70.3 ***	25 - 94.5
Sex	Spayed female	16 (57) †	28	64
	Intact female	0	2	3
	Neutered male	12 (43) †	25	83
	Intact male	0	2	7
Tumor location n (%)	L proximal humerus	3 (10.7)	6 (10.5)	17 (10.8)
	L distal radius	5 (17.9)	6 (10.5)	34 (21.7)
	Other L forelimb	3 (10.7)	1 (1.8)	5 (3.2)
	R proximal humerus	7 (25)	4 (7)	16 (10.2)
	R distal radius	2 (7.1)	12 (21.1)	23 (14.6)
	Other R forelimb	0 (0)	3 (5.3)	4 (2.5)
	L distal femur	1 (3.6)	1 (1.8)	8 (5.1)
	L proximal tibia	2 (7.1)	2 (3.5)	4 (2.5)
	L distal tibia	1 (3.6)	6 (10.5)	16 (10.2)
	Other L hindlimb	0 (0)	2 (3.5)	2 (1.3)
	R distal femur	1 (3.6)	5 (8.8)	11 (7)
	R proximal tibia	0 (0)	3 (5.3)	4 (2.5)
	R distal tibia	3 (10.7)	5 (8.8)	11 (7)
	Other R hindlimb	0 (0)	1 (1.8)	2 (1.3)

Table 1. The clinical characteristics of the dogs enrolled in the VIGOR study are similar to historical comparison cohorts with appendicular osteosarcoma

The demographic characteristics of the study population show that the dogs enrolled in VIGOR were generally large breed middle-aged dogs. Only cases with tumors of the appendicular skeleton were enrolled in VIGOR.

VIGOR: test population

UMN: comparison cohort at same site

COTC: national multicenter comparison cohort

VIGOR, UMN, and COTC were all treated with the same standard of care chemotherapy. All cases were treated with amputation.

* One Mastiff in VIGOR was diagnosed with intramedullary rhabdomyosarcoma.
** One mixed breed enrolled in VIGOR was diagnosed with intramedullary hemangiosarcoma.
*** Body weight was not available for 5 of the dogs in the UMN comparison group.
† All dogs were neutered and weighed > 20kg as conditions of enrollment in VIGOR.

Figure 3. Intravenous VSV therapy is safe, and a subset of VSV-treated dogs with osteosarcoma are long term responders.

A. Kaplan-Meier survival curve with event free survival (EFS) of dogs treated with systemic VSV in the VIGOR study, compared to two historical control cohorts. **B.** Long-term response was defined as overall survival (OS) greater then 75th percentile of survival from the COTC cohort (479 days). Similar proportion (26%) of dogs from the VMC cohort were long-term responders. 35% of dogs from the VSV-treated dogs from the VIGOR cohort were long term responders (OS > 479 days).

Figure 4

Figure 4. Increased tumor Inflammation following VSV treatment in resected osteosarcoma tissues

Inflammatory infiltrates were scored by a pathologist in pre-treatment tumor biopsies and post-treatment amputation-resected tumor samples. **A.** Comparison of Tumor Inflammation Score (TIS) in pre- vs post- treatment specimens significantly higher TIS in post-treatment samples from VSV-treated dogs. **B.** Where matched paired pre- and post-treatment tumor samples were available for assessment, we observed a significant increase in TIS in tissues from VSV-treated dogs between baseline and 10 days post treatment (P=0.0027, paired t-test).

Figure 5. Acute viremia following systemic VSV infusion.

qRT-PCR analysis of whole blood, peripheral blood mononuclear cells (PBMCs), and plasma samples collected at indicated time points following VSV infusion shows detection of viral RNA in blood and viral localization primarily to PBMCs.

Figure 6. RNAseq analysis of canine tissues does not show significant changes immune gene signatures

A. RNAseq analysis of osteosarcoma tumor samples (pre- and post-treatment), lung metastases (where available from necropsy samples), skin biopsy, and osteosarcoma cell lines (derived from osteosarcoma tumors) clustered by tissue type; **B.** Unsupervised hierarchical clustering resulted in 14 gene clusters including cell cycle and immune clusters. **C.** Gene cluster expression summary score (GCESS) in osteosarcoma cell lines, skin, osteosarcoma tumors, and metastases were evaluated. Immune or cell cycles GCESS scores were not significantly different in pre- vs. post-treatment VSV and placebo treated tumor samples.

Figure 7. Survival outcomes correlate with pre-treatment CD8 T-cell GCESS

A. Survival outcomes in placebo versus VSV treated dogs, **B.** Correlation of survival outcomes with CD37 monocyte, CD8 T-cell, and cell cycle GCESS indicates a correlation CD8 T-cell GCESS in pre-treatment tumor biopsy samples.