Heterologous Expression and Characterization of *ToxA1* Haplotype from India and its

Interaction with *Tsn1* for Spot Blotch Susceptibility in Spring Wheat

Ranjan Kumar Chaubey[1], Dharamsheela Thakur[2], Sudhir Navathe[3*], Sandeep Sharma[1],

Vinod Kumar Mishra[1], Pawan Kumar Singh[4], Ramesh Chand[1*]

[1] Institute of Agricultural Sciences, Banaras Hindu University Varanasi 221005, India

[2] Department   of Molecular   Biology and Genetic   Engineering, Bihar   Agricultural

University, Sabour 813210, Bhagalpur, India;

[3] Agharkar Research Institute, G. G. Agarkar Road, Pune 411004, India

[4] International Maize and Wheat Improvement Center (CIMMYT), Texcoco 56237, Mexico.

* Correspondence:

Ramesh Chand, Banaras Hindu University, Varanasi, India, Email: rc_vns@yahoo.co.in;

Sudhir Navathe, Agharkar Research Institute, Pune, India, Email: snavathe@aripune.org

Abstract

Background: *ToxA*, a necrotrophic effector protein, is present in the genome of fungal species like *Parastagnospora nodorum*, *Pyrenophora tritici-repentis* and *Bipolaris sorokiniana*. *Tsn1* is the sensitivity gene in the host whose presence indicates more susceptibility to *ToxA* carrying pathogen, and *ToxA-Tsn1* interaction follows an inverse gene-for-gene relationship.

Methods and Results: The present study involved cloning and expressing the *ToxA1* haplotype from *B. sorokiniana*. It was found that the amplicon exhibited an expected product size of 471 bp. Sequence analysis of the *ToxA1* nucleotide sequence revealed the highest identity, 99.79%, with *P. tritici-repentis*. The protein expression analysis showed peak expression at 16.5kDa. Phylogenetic analysis of the *ToxA1* sequence from all the *Bipolaris* isolates formed an independent clade along with *P. tritici-repentis* and diverged from *P. nodorum*. *ToxA-Tsn1* interaction was studied in 18 wheat genotypes (11 *Tsn1* and 7 *tsn1*) at both seedling and adult stages, validating the inverse gene-for-gene relationship, as the toxin activity was highest in the K68 genotype (*Tsn1*) and lowest in WAMI280 (*tsn1*).

Conclusion: The study indicates that the haplotype *ToxA1* is prevailing in the Indian population of *B. sorokiniana*. It would be desirable for wheat breeders to select genotypes with *tsn1* locus for making wheat resistant to spot blotch.

Keywords: *Bipolaris sorokiniana*, effector-triggered susceptibility, inverse gene-for-gene relationship, sensitivity genes, *ToxA1*, *Triticum aestivum*

Introduction

Host selective toxins (HSTs) are the effector molecule(s) released during the invasion of necrotrophic pathogens and act as major determinants of pathogenicity (1). These necrotrophic effectors are specific biomolecules that are structurally complex, chemically diverse, and responsible for causing some key diseases in important crops. HSTs are very important in host-pathogen interaction but are produced by limited genera of fungi involving multiple toxins (2). Spot blotch of wheat is a damaging disease caused by *Bipolaris sorokiniana* (Sacc.) Shoemaker, syn. *Cochliobolus sativus* (S. Ito & Kurib.) a hemi-biotrophic pathogen. This disease is mainly prevalent in South Asia's warm, humid region, especially in the eastern Indo-Gangetic plains of India. Wheat productivity can decline by 40% due to spot blotch alone (3). Spot blotch has been reported to cause up to 17.5% yield loss in the Indian sub-continent, and about 25mha of wheat is affected by spot blotch worldwide (4). Plant pathogens have their genetic makeup for causing host infection, which can decided by the presence of a single gene that confers host susceptibility or resistance (5, 6). The proteinaceous necrotrophic effector ToxA a major HST produced by *B. sorokininana*. Additionally, ToxA is known to be produced from two necrotrophic pathogens, *Parastagonospora nodorum* (Berk.) Quaedvl., Verkley & Crous, syn. *Phaeosphaeria nodorum* (E. Müll.) Hedjar. and *Pyrenophora tritici-repentis* (Died.) Drechsler and cause Septoria nodorum blotch and tan spot, respectively (7, 8). ToxA protein was reported initially in a culture filtrate of the wheat pathogen *P. tritici-repentis* (8). The ToxA protein was first discovered by Tomás and Bockus (9) from the culture filtrate of *P. tritici-repentis*. The *ToxA* manifested high nucleotide diversity in *P. nodorum,* whereas it showed a lack of variation in *P. tritici-repentis*. This pattern of nucleotide diversity and the recent emergence of tan spot suggests the recent introduction of *ToxA* into *P. tritici- repentis* genome (6). The *ToxA* sequences from *P. nodorum* and *P. tritici-repentis* depicted differences at four fixed nucleotide sites resulting in two predicted amino acid changes consistent with the

interspecific transfer of this gene from *S. nodorum* into *P. tritici-repentis* (6), and this *ToxA*

gene jumped horizontally from *P. tritici-repentis* to another pathogen *B. sorokiniana,* (10).

ToxA and its surrounding 14 kb suggest that this gene was horizontally transferred, facilitated

by a type II DNA transposon (11). *ToxA* is a single-copy gene, and it was found to be embedded

within a 12-kb genomic element nearly identical to the corresponding regions in *P. nodorum*

and *P. tritici-repentis* (10). The mature ToxA protein is 13.2 kDa, and it encodes a signal

peptide of 23 amino acids and a pro-domain of 4.3 kDa (required for protein folding), and both

are cleaved before the secretion of mature ToxA protein (1, 12, 13). Studies revealed that the

host sensitivity of ToxA was mapped on wheat chromosome arm 5BL, and the sensitivity gene

was named *Tsn1* (14). Further, it was confirmed that this gene was significant for both tan spot

and SNB, and in both cases, the pathogen had a sensitivity gene such as *ToxA* (6, 15). For the

wheat-*B. sorokiniana* pathosystem, ToxA confers virulence, and it is the only known virulence

gene whose corresponding sensitivity gene (*Tsn1*) in the host has been identified. Further, it

was also demonstrated that *Tsn1*-carrying wheat cultivars are more sensitive to *B. sorokiniana*

isolates that harboured the *ToxA* gene. Cloning of the *Tsn1* gene unfolded characteristics

similar to classical disease resistance nucleotide-binding leucine-rich repeat (NLR) gene that

included serine/threonine kinase, leucine-rich repeats, nucleotide binding and protein kinase

domains (14). The ToxA gene is transcribed as a necrotrophic effector protein responsible for

infection in the host, with *Tsn1* being a resistance gene (16). A resistance-like gene *Tsn1* in

wheat confers susceptibility to *ToxA* in *P. tritici-repentis* and *P. nodorum* isolates (6). This

ToxA–Tsn1 interaction system has been proved as an example of the inverse gene-for-gene

relationship. It was realizing the occurrence of spot blotch in South Asia and the emergence of

ToxA1 as a key virulence gene driven us towards studying this host-pathogen relation. Keeping

this in mind, we cloned the *ToxA1* gene from *B. sorokiniana* and conducted expression analysis

for the ToxA1 protein. We also validated the pathogenicity of *ToxA1* on the allelic status of

Tsn1 by using 18 *Tsn1/tsn1* genotypes.

Materials and Methods

Primer designing:

ToxA is a 534 bp-long gene sequence, and primers were designed using the web-based tool

Primer-BLAST (Basic Local Alignment Search Tool) from NCBI. The *ToxA* sequence from *P.*

tritici-repentis (U79662.1) was obtained in FASTA format from NCBI (National Centre for

Biotechnology Information) GenBank database (https://www.ncbi.nlm.nih.gov/genbank/). The

primers were utilized to amplify *ToxA* in *B. sorokiniana* isolate MCC 1533 (Pusa2). The isolate

MCC1533 (Pusa2) has been identified and deposited at National Center for Microbial

Resources (NCMR), Pune, India. The primer sequence used is given as Forward Primer (FP) -

CGCGGATCCGATCCCGGTTACGAAATCG and Reverse Primer (RP) - CGC

CTCGAGCTAATTTTCTAGCTGCATTCTC. Revised nomenclature of ToxA haplotypes

corresponds the

RNA extraction and cDNA synthesis

The B. sorokiniana isolate MCC 1533 (Pusa2) was cultured for 10 days in a Potato Dextrose

Broth. Mycelium was harvested, and the total RNA was isolated using Invitrogen PureLink^{TM}

RNA Mini kit following the manufacturer's protocol. The integrity and size distribution of total

RNA purified with RNeasy Kit was checked by 1.2% agarose gel electrophoresis and ethidium

bromide staining, visualizing two sharp bands of 28S rRNA to 18S RNA. RNA was quantified

using Nanodrop2000 instrument (Thermo Scientific) and 1µg RNA reverse transcribed to

cDNA was synthesized using a Thermo cDNA synthesis kit following the manufacturer's

protocol.

An additional treatment of RNase was included during cDNA synthesis as supplied and

instructed with the kit and manufacturer's protocol to prevent chances of contamination of

genomic DNA during isolation.

ToxA gene cloning

The expression construct was produced in the bacterial expression vector pET28a that directs

the expression of HIS-tagged fusion proteins in *E. coli*. BamH1 and XhoI restriction sites were

introduced in forward and reverse primer for cloning in pET vector. Primers were designed

such that the PCR amplified portion of the *ToxA1* gene was generated with a 5' BamHI site and

a 3' XhoI site. The PCR product was digested with BamHI and XhoI, gel-purified with the

QIAquick Gel Extraction Kit (Qiagen, Valencia, CA) and cloned into the BamHI and XhoI

sites of pET28a. Primers FP-CGCGGATCCGATCCCGGTTACGAAATCG and RP- CGC

CTCGAGCTAATTTTCTAGCTGCATTCTC were used for the PCR amplification. PCR

amplification was performed using cDNA as a template in an Eppendorf Mastercycler using

the following cycles; initial denaturation of 95°C for 7 min followed by 35 cycles of 94°C (30

s), 55°C (45 s) and 72°C (1 min) and a final extension of 10 min at 72°C. The PCR product

was gel-purified, digested with BamHI and XhoI, and ligated into the respective restriction

sites of pET28a. The resulting construct contained the predicted effector protein gene fused to

a 6X histidine tag at the N-terminus. The ligated product was used to transform *E. coli*

DH5alpha and BL21 (DE3) strains (17) for cloning and protein expression, respectively.

Sequencing and characterization

Cloned fragment of *ToxA* from *B. sorokiniana* isolate MCC1533 (Pusa2) was sequenced by

Sanger sequencing. The sequences obtained in two frames were merged, and Bioedit Sequence

Alignment Editor (v 7.2) was used to identify overlapping sequences in both orientations (18).

Integrated sequences were then prepared in a single frame. The sequence has been deposited

to the National Centre for Biotechnology Information (NCBI) database with Gene Bank

accession number (OP289649). The processed *ToxA* sequence was used as a query for BLASTn

(https://blast.ncbi.nlm.nih.gov/Blast.cgi) for finding homologous sequences with reference to

the NCBI nucleotide database (19). The processed *ToxA1* nucleotide sequence was then

translated to the amino acid sequence in BioEdit 7.2. Sequences of *P. tritici-repentis, P.*

nodorum and different isolates of *B. sorokiniana* were retrieved from the NCBI protein

database. The amino acid sequences retrieved using BLASTp searches were then used to create

multiple sequence alignments of 27 ToxA sequences. MEGA X (Molecular Evolution Genetics

Analysis) software (19) was then used to generate a phylogeny of *ToxA* amino acid sequences

of 49 isolates of *P. tritici-repentis, P. nodorum* and *B. sorokiniana* using the CLUSTAL W

alignment algorithm (20). All alignment parameters were left at their default values; 178 and

114 positions with and without gaps were obtained, respectively. Sequences were analyzed

using substitutions selected with complete deletion of gaps or absent data. The phylogenetic

tree was inferred using the JTT model and the maximum likelihood method (21). Using the

default setting and the Neighbor Joining method, the initial tree was inferred, and Nearest-

Neighbor Interchange was used as the ML heuristic search method. Phylogenetic tree

reconstructions were performed using the Maximum Likelihood Method. For the Poisson

correction model, 1000 bootstrap replicates were used to assess node support (22).

Protein modelling and topographical properties

The predicted protein sequence of cloned Pusa2 *ToxA* was subjected to the SWISS-MODEL

homology modelling online tool (23). Proteins were modelled utilizing Protein Data Bank

(PDB) entry 1zld.1.A (*P. tritici-repentis*) as reference. The quality of modelled protein was

judged based on GMQE (Global Model Quality Estimation) and QMEAN (Qualitative Model

Energy Analysis) Z-score parameters (24, 25). The best model was selected based on the

GMQE score, which ranges from 0 to 1, with a higher value indicating better reliability. On a

global scale, the QMEAN Z-score estimates the "degree of nativeness" of the structural features

observed in the model. QMEAN Z-scores around zero indicate good agreement between the

model structure and experimental structures. Scores of -4.0 or below are an indication of

models with low quality. The quality of the resulting models was also monitored with

PROCHECK and ProsSA-WEB (26). Geometrical and topographical properties of the

modelled proteins were identified using the Computed Atlas of Surface Topography of Proteins

(CASTp) 3.0 online server (27). The UCSF Chimera software was used to create images of

molecular graphics (28). The alpha helix and the beta-sheet percentage were analyzed using

PDBMD2CD.

Isolation of proteins from E. coli culture

E. coli colony harboring the expression construct was inoculated in 5 ml of LB (Sambrook et

al. 2006) with 100μg of kanamycin per ml (LBA) and grown overnight at 37°C and shaking at

250 rpm (G24 Incubator Shaker, New Brunswick Scientific, Edison, NJ). The overnight culture

was used to inoculate 25 ml of fresh LBA, which was then incubated with shaking at 37°C for

one hour. The expression of fusion proteins was induced by the addition of isopropyl-beta-

Dthiogalactopyranoside (IPTG) to a final concentration of 1 mM, followed by an additional 4

hours of incubation under the same conditions. After 20 minutes of centrifugation at 4,000 g

and 4°C, cell particles were resuspended in a cold sonication buffer(SB; 50 mM Na2HPO4, 5

mM Tris-base, 300 mM NaCl, pH 8.0) with 0.5% Triton-x-100 and 10 mM DTT. Resuspended

pellets were sonicated three times, 30 s each, with 2 min on ice between periods. Insoluble

material was then pelleted by centrifugation at $10,000 \times g$, 4°C, for 10 min. Pellets were washed

thrice with SB with 0.5% Triton-× -100 and 10 mM DTT. The final washed pellets, containing

inclusion bodies, were either stored frozen at –80°C or resuspended in 1 ml of 8 M guanidine

HCl (Sigma) in refolding buffer (250 mM NaCl, 100 mM sodium phosphate buffer, 10 mM

Tris-base, 4% glycerol, 1 mM EDTA, and 0.005 % Tween 20, pH 8.0) with freshly prepared

DTT added to a final concentration of 10 mM. Solubilized proteins were sonicated briefly twice

(for 5 to 10 s) and then heated to 37°C for 10 min and 50°C for an additional 5 min to assist in

the solubilization of proteins. Insoluble material was removed from the samples by centrifuging

at 12,000 ×g for 10 min. Then, solubilized proteins were quantitated with Bradford assay

reagent (Bio-Rad, California, USA) with bovine serum albumin used as a standard. Fusion

proteins solubilized in guanidine HCl were used immediately.

Screening of ToxA1 –Tsn1 interaction on different wheat genotypes

The screening of 18 wheat genotypes was conducted for *Tsn1/tsn1,* out of which 11 genotypes

harboring *Tsn1* and the rest 7 genotypes with *tsn1* (see Table 1) were sown in 3 replications at

polyhouse, Plant Pathology, Institute of Agricultural Sciences, Banaras Hindu University in

1m rows at a distance of 22.5cm between two rows. Infiltration of ToxA1 protein, as well as

control (Uninduced protein), was done at both the seedling stage (3-5 leaf stage) and adult stage

(Zadoks growth stage 55) (29). 250µg of ToxA protein was infiltrated by a sterilized syringe,

and symptoms were recorded 7 days after infiltration. The 5th emerged leaf was infiltrated for

seedlings, while the flag leaf was infiltrated with toxin ToxA1 for adult plants. The temperature

was maintained at 25±2°C. The light/dark period was maintained at 14hrs light and 10hrs of

darkness. Photographic data was captured using a Nikon camera. The lesion size was measured

in length, width, and area of lesions using ImageJ software (Version 1.53s) (30).

Statistical Analysis

The statistical analysis was conducted for phenotypic traits, including length, width, and area

of necrotic lesions of wheat genotypes with the *Tsn/tsn1* gene at both seedling and adult stages.

The point biserial correction coefficients between disease components and the

presence/absence of *Tsn1* were calculated (32). The analysis in   R software V 4.2.1 was

performed using packages' *phytools', 'stats', 'agricolae', 'ggplot2', 'ggtree'* (31).

Results

Cloning and expression study of ToxA1

The *ToxA* gene from *B. sorokiniana* was amplified using primers designed from *PtrToxA* through RT-PCR. The amplicon exhibited the expected product size of 471bp, which was then cloned in the pET28a vector. The cloned gene was validated using colony PCR of the transformed cell. The *ToxA1* gene, after validation, was subjected to sequencing, and the sequence data after processing revealed successful cloning of the *ToxA* gene (now *ToxA1- B. sorokininana*). This *ToxA1* nucleotide sequence was further used for BLASTn analysis, revealing the highest identity, 99.79%, with *P. tritici-repentis* and E-value close to zero. Expression of ToxA1 was done by inducing the transformed *E.coli* cell through IPTG of different concentration, namely, at 0.4mM, 0.6mM, 0.8mM and 1.0mM. Maximum expression of ToxA1 was obtained at 1.0mM concentration with an expected band size of 16.5kDa in sodium dodecyl sulfate (SDS)-polyacrylamide gels (Fig. 1).

ToxA1 characterization and Phylogenetic analysis

The identification of *ToxA1* after showing sequence similarity with *P. tritici-repentis ToxA1* using BLASTn analysis on NCBI was conducted. Further characterization of ToxA1 was conducted by translating the nucleotide sequence to amino acid sequence using the web-based tool Expasy (https://www.expasy.org/resources/translate). The amino acid sequence of ToxA1 was subjected to BLASTp analysis on NCBI, which depicted 100% similarity with *P. tritici-repentis* showing E value $9e^{-109}$. Multiple sequence alignment of 25 ToxA protein sequences and *ToxA1* MCC1533 (Pusa2) translated sequence in Bioedit revealed that amino acid differences were found in most *P. nodorum* strains. However, the cloned Pusa2_*ToxA1* sequence showed complete similarity to *P. tritici-repentis* ToxA sequence. *P. nodorum ToxA* sequence depicted amino acid substitution for Glutamate, Aspartate, Leucine, Valine and Asparagine (Fig. 2).

The phylogenetic analysis depicts the evolutionary relatedness of the *ToxA1* amino acid sequence with other ToxA sequences. The tree indicates that *ToxA1* sequences originate from

a common ancestor but form different clades during evolution. The *ToxA1* sequence from all

Bipolaris isolates formed an independent clade along with *P. tritici-repentis* and diverged from

P. nodorum. The phylogeny indicates sequence similarity among *B. sorokiniana* and *P. tritici-*

repentis, whereas it differed from *P. nodorum* (Fig. 3).

Protein modelling and Topographical properties

The structure of the ToxA1 protein of a cloned gene of *B. sorokiniana* isolate MCC1533

(Pusa2) was predicted using the SWISS-MODEL homology modelling online server. For

checking the reliability of models GMQE (ToxA1: 0.56), QMEAN Z scores are included

QMEAN -0.33, Cbeta –1.79, torsion 0.53 and these scores indicated that these models are

reliable and good quality. The quality of the resulting models was monitored with

PROCHECK. Ramachandran plot analysis revealed that 88.1% of the ToxA1 protein structure

residues fell within the most favoured regions, with a further 11.9% in the additionally allowed

region (Fig. 4). No residues were found in the generously allowed or disallowed regions. The

number of non-glycine and non-proline residues was 100%. The number of end residues

(excluding Gly and Pro) is 2, and the number of glycine and proline residues was 12 and 3,

respectively.

Predicted ToxA1 protein structures of *B. sorokiniana* MCC1533 (Pusa2) isolate based

on homology modelling are depicted in Fig. 4. The protein sequences of ToxA1 protein showed

sequence identity of 100% with template PDB entry 1zld.1.A (*P. tritici-repentis*, *PtrToxA*).

When generated structures were compared in UCSF Chimera, it was observed that the RGD

motif (toxic region) was present in the 3D structure of ToxA1 protein and found that arginine

is present in the alpha helix of the loop at 118 positions. At the same time, Glycine and Aspartic

acid is present in the β-sheet of the protein structure at 119 and 120 positions of ToxA1 protein

(Fig. 5). The percentage of the alpha helix was 3.39%. At the same time, antiparallel sheet 1

was 35.55%, antiparallel sheet 2 was 12.76%, the parallel sheet was 0.85%, the turn was 7.63%,

and the other was 39.83%.

Bioactivity of ToxA1 effector protein and its interaction with Tsn1 at both seedling and adult

stage

The bioactivity of the *ToxA1* effector was studied on necrotic lesions symptoms caused due to

ToxA1 protein and was found to be more prominent at the adult stage. At the seedling stage,

the maximum symptomatic area (0.977 cm^2) was recorded in genotype K68(*Tsn1*) and the

minimum (0.265 cm^2) in WAMI280 (*tsn1*). At the same time, the length varied from 2.872 cm

(K68, *Tsn1*) to 0.860cm (PBW343, *tsn1*). However, the maximum width recorded was 0.432

cm for genotype WAMI288 (*Tsn1*), and the minimum in genotype harbouring *tsn1*(LOK1) was

0.307cm.

At the adult stage, the length of the symptom was maximum (4.613cm) in genotypes

harbouring*Tsn1*(K68) and minimum (1.305 cm) WAMI280 (*tsn1*), and the width also had the

same trend ranging from 0.633cm (C306, *Tsn1*) to 0.327cm (WAMI280, *tsn1*). The area was

measured for quantifying the necrotrophic symptom, and it found that the maximum area in

the genotype harboring *Tsn1* was 2.326 cm^2 (K68), and the minimum was 0.411

cm^2(WAMI280, *tsn1*). A significant difference was found between the seedling and adult

stages and for *Tsn1* and *tsn1* (Table 1, Fig. S2, S3, S4).

Discussion

Studies on necrotrophic effectors producing HSTs have provided a new dimension to host-

pathogen interaction and established an inverse gene-for-gene relationship (6, 34). The HSTs

are effectors proteins and induce a susceptible response in the host harbouring the dominant

susceptibility gene. ToxA is a necrotrophic effector protein, one of the key factors for causing

the most devastating disease, spot blotch in wheat, prevalent in warm, humid regions of South

Asia. The present study focuses on cloning of *ToxA1* gene, followed by its characterization and

purification of the resultant protein and its expression analysis. This study successfully cloned

and characterized the *ToxA1* gene from *B. sorokiniana* isolate MCC1533 (Pusa2). This gene

was found initially in *P. nodorum* and horizontally transferred to *P. tritici-repentis* and, more

recently, to *B. sorokiniana* (6, 35). The cloned gene and its expression was studied in the

present study, and the purified protein showed peak expression at a concentration of 1.0 mM

IPTG with band migration at 16.5kDa in size. Manning et al. (36) also reported similar findings

where lysates with His-ToxA1 expressed a phosphorylated band that migrated at 16 kDa

compared to lysates without His-ToxA1 (control). However, mass spectral analyses have

indicated a 13.2 kDa mass of purified mature ToxA1 protein from *P. tritici-repentis* (1, 8). The

importance of a protein purification study for ToxA1, which has selective host toxicity for

genetic analysis of pathogenicity in the *Pyrenophora*-wheat interaction, was also suggested by

Touri et al. (8).

The *ToxA1* gene was truncated in the present study to remove the signalling sequences

comprising 22 amino acids, as the mature ToxA1 gene is released after the proteolytic cleavage

of this signal sequence. Previous studies have also described these signalling peptides in ToxA

protein that involve the first 22 amino acids, which are proteolytically cleaved after import into

the endomembrane system (1). The release of mature toxin involves additional proteolytic

processing, and as the ToxA1 protein is blocked at the N terminus, such modifications would

incur a mass increase (1). The ToxA1 cloned sequence obtained in our study has been compared

with the already reported ToxA1 sequences from *P. nodorum*, *P. tritici-repentis* and *B.*

sorokiniana. Multiple sequence alignment was performed on the amino acid sequences of 26

isolates. The results indicate that the ToxA regions of *B. sorokiniana* and *P. tritici-repentis* are

more closely related to one another than they are to *P. nodorum*. This conclusion is supported

by the identical ToxA1 sequence shared by *B. sorokiniana* and *P. tritici-repentis* and the six-

position difference between *P. nodorum* haplotypes. (Fig. 2). The sequence identity between

the three fungal species accounts for the common evolutionary origin of *ToxA* and the potential of exchanging DNA between these species (10). The significance of phylogenetic studies in establishing evolutionary relationships among the different pathogen races or deriving the specific gene evolution between different genera is well known. The sequencing data for *ToxA1* showed a high similarity between *B. sorokiniana* and *P. tritici-repentis* (10, 37, 38). Previous studies have also suggested that the ToxA1 sequence did not differ for *B. sorokiniana* and *P. tritici-repentis* but differed from the ToxA sequence of *P. nodorum* (10, 34, 37, 38). This evolutionary relatedness of *P. tritici-repentis* and *B. sorokiniana* sequences have also been reported earlier (10, 34). The earliest report suggested that ToxA has been present in *P. nodorum* for a long time as it exhibited greater sequence diversity and was interspecifically transferred to *P. tritici-repentis* quite later. This is evident from monomorphism or lesser sequence diversity in the ToxA1 sequence from *P. tritici-repentis* and the emergence of tan spot symptoms more recently in the near past (38).

Further insights into the ToxA structural analysis were performed by predicting the 3D protein structure using an amino acid sequence of ToxA1 (Fig.4). This is the first report for the 3D protein structure ToxA1 haplotype from *B. sorokiniana*. The predicted protein structure was similar to a single-domain protein with a β-sandwich fold of PtrToxA. This β-fold represents a solvent-exposed loop containing arginyl-glycol-aspartic acid (RGD)- functionally active motif (39, 40). ToxA1 has been known to resemble with fibronectin III domain which retains a solvent-exposed loop containing RGD motif for cell wall-plasma membrane interaction in plants (40, 41). It has been proposed that ToxA with RGD-containing motif has been known to interact with wheat mesophyll cells and thus establish disease (42). The mesophyll cells have a high-affinity matrix to recognize these motifs, and these all suggest the importance of the RGD motif in receptor binding and internalization into wheat cells (40).

ToxA is the only known virulence gene for which a host sensitivity gene (*Tsn1*) has been known

in wheat-*B. sorokiniana* pathosystem. For the first time, the present study reports a cloned and

purified protein ToxA1 infiltration in both seedling and adult stages of wheat lines harboring

Tsn1/tsn1. The severity was enhanced in the adult stage compared to the seedling stage in *Tsn1*

wheat lines. The present data also reveals that *Tsn1*-carrying genotypes exhibited more disease

in seedlings than in the adult stage. The toxin activity was highest in the K68 genotype (*Tsn1*)

and lowest in WAMI280 (*tsn1*) at both the seedling and adult stages. This could be validated

by past studies suggesting that ToxA causes severe necrotic lesions on *Tsn1,* harboring wheat

genotypes.

In contrast, less or no lesion was observed for wheat genotypes lacking the sensitivity gene

(43). Navathe et al. (34) also reported one such possibility of the involvement of a recessive

allele of *Sb2* towards an unknown HST. *Tsn1* has been reported attributing susceptibility

towards both *P. nodorum*-causing SNB (6, 44) and *P. tritici-repentis*, causing tan spots (45).

However, the environmental requirements for *Bipolaris* are different from these two pathogens,

which are not prevalent in the South-Asian region.

Conclusions:

Nevertheless, it would be desirable for wheat breeders to select *tsn1* for making crops resistant

to *B. sorokiniana, P. tritici repentis* and *P. nodorum* in nearest future. The study in this horizon

should focus on loci other than *Tsn1* and *Sb2*. It would be interesting to study other quantitative

trait loci involved to understand the complexity of this *ToxA-Tsn1* interaction and spot blotch

disease resistance.

Acknowledgements: Authors acknowledge Banaras Hindu University and Bihar Agricultural
University for the infrastructure facility
Funding: SN acknowledges SERB, New Delhi, for CRG/2021/007211.

Author Contributions: Conceptualization, RC, VKM, PKS, SN, and SK; methodology, RKC
and DT; validation, DT and SS; formal analysis, DT and SS; investigation, RKC; resources,
RC, DT; data curation, DT, SS; writing—original draft preparation, RKC, SN; writing—review

and editing, RC, PKS; supervision, RC, VKM, SS; project administration, RC; funding
acquisition, RC, PKS. All authors have read and approved the manuscript.

Availability of data and materials: Not applicable.

Declarations:

Ethics approval and consent to participate: Not applicable.

Consent for publication: Not applicable.

Competing interests: The authors declare that they have no competing interests.

References:

1. Ciuffetti, L. M., Tuori, R. P. Gaventa, J. M. (1997). A single gene encodes a selective toxin causal to the development of tan spot of wheat. Plant Cell, *9*, 135-144. https://doi.org/10.1105/tpc.9.2.135

2. Wolpert, T.J., Dunkle, L.D., Ciuffetti, L.M. (2002). Host-selective toxins and avirulence determinants: what's in a name? Annu. Rev. Phytopathol. 40, 251–285. https://doi.org/10.1146/annurev.phyto.40.011402.114210

3. Sharma, P., Duveiller, E., Sharma, R. C. (2006). Effect of mineral nutrients on spot blotch severity in wheat and associated increases in grain yield. Field Crops Res, *95*, 426-430. https://doi.org/10.1016/j.fcr.2005.04.015

4. Pandey AK, Mishra VK, Chand R, Navathe S, Budhlakoti N, Srinivasa J, Sharma S, Joshi AK (2021). Crosses with spelt improve tolerance of South Asian spring wheat to spot blotch, terminal heat stress, and their combination. Sci Rep, *11*, 1-12. https://doi.org/10.1038/s41598-021-85238-x

5. Ballance, G.M., Lamari, L., Bernier, C.C. (1989). Purification and characterization of a host-selective necrosis toxin from *Pyrenophora tritici-repentis*. Physiol. Mol. Plant Pathol. *35*, 203–213 https://doi.org/10.1016/0885-5765(89)90051-9

6. Friesen, T. L., Stukenbrock, E. H., Liu, Z., Meinhardt, S., Ling, H., Faris, J. D., ... Oliver, R. P. (2006). Emergence of a new disease as a result of interspecific virulence gene transfer. Nat. Genet. *38*: 953-956. https://doi.org/10.1038/ng1839

7. McDonald, M. C., & Solomon, P. S. (2018) Just the surface: advances in the discovery and characterization of necrotrophic wheat effectors. Curr. Opin. Microbiol. *46*, 14-18.

8. Solomon, P., Lowe, R., Tan, K.C., Waters, O., Oliver, R. (2006) *Stagonospora nodorum*; cause of Stagonospora nodorum blotch of wheat. Mol. Plant Pathol. *7*, 147–156 https://doi.org/10.1111/j.1364-3703.2006.00326.x

9. Tuori, R. P., Wolpert, T. J. Ciuffetti, L. M. (1995). Purification and immunological characterization of toxic components from cultures of *Pyrenophora tritici-repentis*. Mol. Plant Microbe Interact, *8*, 41-48. https://doi.org/10.1094/mpmi-8-0041

10. Tomás, A., Bockus, W. W. (1987). Cultivar specific toxicity of culture filtrate of *Pyrenophora tritici-repentis*. Phytopathology, *77*,1337–1366. https://doi.org/10.1094/phyto-77-1337

11. McDonald, M. C., Ahren, D., Simpfendorfer, S., Milgate, A. Solomon, P. S. (2018). The discovery of the virulence gene ToxA in the wheat and barley pathogen *Bipolaris sorokiniana*. Mol. Plant Pathol., *19*, 432-439. https://doi.org/10.1111/mpp.12535

12. Ballance, G. M., Lamari, L., Kowatsch, R., Bernier, C. C. (1996). Cloning, expression and
occurrence of the gene encoding the Ptr necrosis toxin from *Pyrenophora tritici-repentis*.
Mol. Plant Pathol. https://doi.org/10.1007/978-94-011-5218-1_21
13. Tuori, R. P., Wolpert, T. J. Ciuffetti, L. M. (2000). Heterologous expression of functional
PtrToxA. Mol. Plant Microbe Interact, *13*, 456-464.
https://doi.org/10.1094/mpmi.2000.13.4.456.
14. Faris, J. D., Zhang, Z., Lu, H., Lu, S., Reddy, L., Cloutier, S., ... Friesen, T. L. (2010). A
unique wheat disease resistance-like gene governs effector-triggered susceptibility to
necrotrophic pathogens. PNAS, *107*, 13544-13549.
https://doi.org/10.1073/pnas.1004090107
15. Lamari, L., Bernier, C. C. (1989). Evaluation of wheat lines and cultivars to tan spot
[*Pyrenophora tritici-repentis*] based on lesion type. Can. J. Plant Pathol. 11, 49-56.
https://doi.org/10.1080/07060668909501146
16. Tan, K. C., Ferguson-Hunt, M., Rybak, K., Waters, O. D., Stanley, W. A., Bond, C. S.,
Stukenbrock, E. H., Friesen, T. L., Faris, J. D., McDonald, B. A., Oliver, R. P. (2012).
Quantitative variation in effector activity of ToxA isoforms from *Stagonospora nodorum*
and *Pyrenophora tritici-repentis*. Mol. Plant Microbe Interact. *25*, 515-522.
https://doi.org/10.1094/mpmi-10-11-0273
17. Sambrook, J., Russell, D. W., Sambrook, J. (2006). The condensed protocols: from
molecular cloning: a laboratory manual, (No. Sirsi) i9780879697723). Cold Spring
Harbor, NY: Cold spring harbour laboratory press. https://doi.org/10.1101/pdb.prot4018
18. Hall, T. A. (1999). A user-friendly Biological Sequence Alignment Editor and Analysis
program for Windows 95/98/NT. Nucleic Acids Symp Ser, *41*, 95–98
19. Altschul S. F., Madden T.L., Schäffer A.A., Zhang J., Zhang Z., Miller W., Lipman D. J.
(1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search
programs. Nucleic Acids Res. *25*, 3389–3402. https://doi.org/10.1093/nar/25.17.3389
20. Kumar S., Stecher G., Li M., Knyaz C., Tamura K. (2018). MEGA X: Molecular
Evolutionary Genetics Analysis across computing platforms. Mol Biol Evol. *35*, 1547–
1549. https://doi.org/10.1093/molbev/msy096
21. Larkin M.A., Blackshields G., Brown N.P., Chenna R., McGettigan P.A., McWilliam H.,
Valentin F., Wallace I.M., Wilm A., Lopez R., Thompson J.D. (2007). Clustal W and
Clustal X version 2.0. Bioinformatics, *23*, 2947–2948.
https://doi.org/10.1093/bioinformatics/btm404
22. Jones DT, Taylor WR, Thornton JM. (1992). The rapid generation of mutation data
matrices from protein sequences. Comput. Appl. Biosci. *8*, 275-282.
https://doi.org/10.1093/bioinformatics/8.3.275
23. Felsenstein (1985). Confidence limits on phylogenies: an approach using the bootstrap.
Evolution *39*, 783–791. https://doi.org/10.1111/j.1558-5646.1985.tb00420.x
24. Arnold K, Bordoli L, Kopp J, Schwede T. (2006). The SWISS-MODELworkspace: a web-
based environment for protein structure homology modelling. Bioinformatics *22*, 95–201.
https://doi.org/10.1093/bioinformatics/bti770
25. Benkert P, Tosatto SCE, Schwede T. (2009) Global and local model quality estimation at
CASP8 using the scoring functions QMEAN and QMEANclust. Proteins Struct Funct
Bioinforma. *77*, 173–180. https://doi.org/10.1002/prot.22532

26. Studer G, Rempfer C, Waterhouse AM, Gumienny R, Haas J, Schwede T. (2020).
QMEAND is Co—distance constraints applied on model quality estimation.
Bioinformatics, *36*,1765–1771. https://doi.org/10.1093/bioinformatics/btz828
27. Laskowski RA, MacArthur MW, Moss DS, Thornton JM. (1993). PROCHECK: a
program to check the stereochemical quality of protein structures. J. Appl. Crystallogr. 26,
283–291 https://doi.org/10.1107/s0021889892009944
28. Tian W, Chen C, Lei X, Zhao J, Liang J. (2018). CASTp 3.0: computed atlas of surface
topography of proteins. Nucleic Acids Res. *46*, W363–W367.
https://doi.org/10.1093/nar/gky473
29. Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, Ferrin TE.
(2004). UCSF Chimera. A visualization system for exploratory research and analysis. J
Comput Chem. *25*, 1605–1612 https://doi.org/10.1002/jcc.20084
30. Zadoks JC, Chang TT, Konzak CF (1974). A decimal code for the growth stages of
cereals. Weed Res, 14, 415-421, https://doi.org/10.1111/j.1365-3180.1974.tb01084.x
31. Abramoff, M.D., Magalhaes, P.J., Ram, S.J. (2004) Image Processing with ImageJ.
Biophotonics Int, 11, 36-42.
32. Tate, R. F. (1954). Correlation between a discrete and a continuous variable. Point-biserial
correlation. Ann. Math. Stat., 25, 603-607. https://www.jstor.org/stable/2236844.
33. R Core Team (2022). R: A language and environment for statistical computing. R
Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/.
34. Navathe S, Yadav PS, Chand R, Mishra VK, Vasistha NK, Meher PK, Joshi AK, Gupta
PK. (2020) ToxA–Tsn1 interaction for spot blotch susceptibility in Indian wheat: An
example of inverse gene-for-gene relationship. Plant Dis, *104,* 71-81.
https://doi.org/10.1094/pdis-05-19-1066-re
35. McDonald, M.C., Razavi, M., Friesen, T.L., Brunner, P.C., McDonald, B.A. (2012).
Phylogenetic and population genetic analyses of *Phaeosphaeria nodorum* and its close
relatives indicate cryptic species and an origin in the Fertile Crescent. Fungal Genet. Biol.
49, 882–895. https://doi.org/10.1016/j.fgb.2012.08.001
36. Manning, V. A., Hamilton, S. M., Karplus, P. A., Ciuffetti, L. M. (2008). The Arg-Gly-
Asp–containing, solvent-exposed loop of PtrToxA is required for internalization. Mol.
Plant Microbe Interact. *21*, 315-325. https://doi.org/10.1094/mpmi-21-3-0315
37. Hafez, M., Despins, T., Nakajima, K., Aboukhaddour, R. (2022). Identification of a novel
ToxA haplotype of *Pyrenophora tritici-repentis* from Japan. Phytopathology, *112*, 1597-
1602 https://doi.org/10.1094/phyto-01-22-0001-sc
38. Aboukhaddour, R., Abdel-Fattah, M. H., McDonald, M., Moffat, C., Navathe, S., Friesen,
T. L., Strelkov, S. E., Oliver, R. P., Tan, K.-C., Liu, Z., Moolhuijzen, P., Phan, H., See, P.
T., Solomon, P. (2023). A revised nomenclature for ToxA haplotypes across multiple
fungal species. Phytopathology, https://doi.org/10.1094/phyto-01-23-0017-sc
39. Sarma GN, Manning VA, Ciuffetti LM, Karplus PA (2005). Structure of PtrToxA: an
RGD-containing host-selective toxin from *Pyrenophora tritici-repentis*. Plant Cell, *17*,
3190–3202. https://doi.org/10.1105/tpc.105.034918
40. Ciuffetti, L. M., Manning, V. A., Pandelova, I., Betts, M. F., Martinez, J. P. (2010). Host-
selective toxins, PtrToxA and PtrToxB, as necrotrophic effectors in the *Pyrenophora*

tritici-repentis‑wheat   interaction.   New   Phytol.   *187,*   911-919.
https://doi.org/10.1111/j.1469-8137.2010.03362.x

41. McDonald, M. C., Solomon, P. S. (2018). Just the surface: advances in the discovery and
characterization of necrotrophic wheat effectors. Curr. Opin. Microbiol. , 46, 14-18.
https://doi.org/10.1016/j.mib.2018.01.019

42. Manning, V. A., Hardison, L. K., Ciuffetti, L. M. (2007). PtrToxA interacts with a
chloroplast-localized   protein.   Mol.   Plant   Microbe   Interact.   *20,*   168-177.
https://doi.org/10.1094/mpmi-20-2-0168

43. Aggarwal, R., Agarwal, S., Sharma, S., Gurjar, M. S., Bashyal, B. M., Rao, A. R., Saharan,
M. S. (2021) Whole Genome Sequencing and Expression Analysis of ToxA in *Bipolaris*
sorokiniana Provides Discernment of Pathogenicity Causing Spot Blotch of Wheat.
PREPRINT (Version 1) available at Research Square. https://doi.org/10.21203/rs.3.rs-
677095/v1

44. Virdi, S., Liu, Z., Overlander, M., Zhang, Z., Xu, S. S., Friesen, T., Faris, J. D. (2016).
New insights into the roles of host gene-necrotrophic effector interactions in governing
susceptibility of durum wheat to tan spot and Septoria nodorum blotch. Genes Genomes
Genet. *6,* 4139-4150. https://doi.org/10.1534/g3.116.036525

45. Faris, J. D., Liu, Z. Xu, S. S. (2013). Genetics of tan spot resistance in wheat. Theor. Appl.
Genet., *126,* 2197-2217. https://doi.org/10.1007/s00122-013-2157-y.

Figure Legends

Figure 1. Expressed ToxA1 protein showing 16.5kd band size on SDS-PAGE

Figure 2 (a.) The multiple sequence alignments with a phylogenetic tree, Colors both on the
phylogenetic tree and the branch scale represent genetic distance. ToxA sequences have
been coloured by four colours shown in the 'seq' column for each nucleotide. Colour
changes on the aligned sequences represent nucleotide differences. (b) Based on the
number of nucleotide variations between ToxA haplotypes, a heat map was created.
The relevant ToxA haplotype in the matrix is represented by each branch of the
phylogenetic tree. The near ties to "dark red" and the distance ties to "white" colour. (c)
Neighbor-joining (NJ) tree for ToxA haplotypes based on the nucleotide sequence
differences. Coloured internal nodes represent the bootstrap confidence level.

Figure 3 Phylogenetic tree using Maximum Likelihood algorithm of different ToxA1 of
Pyrenophora tritici-repentis, Bipolaris sorokiniana, and varipus haplotypes of
Parastagnospora nodorum. The percentage of bootstrap support (1000 replicates) is
indicated at corresponding nodes. The nomenclature of the haplotypes is based on the
Aboukhaddour et al. (2023)

Figure 4. Ramachandran plot and ProsSA z-score analysis of modelled ToxA1 protein from
B. sorokiniana. Ramachandran distributions of amino acids in the model were
calculated using pro check (a), and the z-score plot was plotted using ProsSA-WEB (b).

Figure 5. *ToxA1* protein structure of *B. sorokiniana* isolate MCC 1533 (Pusa2) depicting
conserved RGD motif (ARG-GLY-ASP)

Figure 6. Symptom of leaf necrosis on *Tsn1/tsn1* harbouring genotypes after infiltration ToxA1 protein at seedling (3-5 leaf stage) and on flag leaf of adult plant (Zadoks growth stage 55).

Table Legends

Table1. Response of wheat genotypes (*Tsn1/tsn1*) in terms of leaf necrosis after infiltration of *ToxA1* protein at both seedling and adult stage of wheat genotypes harboring *Tsn1/tsn1* allele.

Supplementary Materials:

Figure S1. Correlation coefficients estimated by point biserial correlation for phenotypic traits associated with*Tsn1/tsn1* gene effect in wheat genotypes with ToxA1 isolate (MCC1533/Pusa2)**.**

Figure S2. The necrotrophic area developed due to ToxA1–Tsn1 interaction at seedling (3-5 leaf stage) and adult stage (Zadoks growth stage 55) of 11 wheat genotypes with *Tsn1* and 7 wheat genotypes with *tsn1***.**

Figure S3. Symptom of leaf necrosis after infiltration ToxA1 protein at both seedling (3-5 leaf stage) and adult stage (Zadoks growth stage 55) of wheat genotypes harbouring *Tsn1*.

Figure S4. Symptom of leaf necrosis after infiltration ToxA1 protein at both seedling (3-5 leaf stage) and adult stage (Zadoks growth stage 55) of wheat genotypes harbouring *tsn1*.

Fig.1: Expressed ToxA1 protein showing 16.5kd band size on SDS-PAGE

Fig. 2 (a.) **The multiple sequence alignments** with a phylogenetic tree, Colors both on the phylogenetic tree and the branch scale represent genetic distance. ToxA sequences have been coloured by four colours shown in the 'seq' column for each nucleotide. Colour changes on the aligned sequences represent nucleotide differences. (b) Based on the number of nucleotide variations between ToxA haplotypes, a heat map was created. The relevant ToxA haplotype in the matrix is represented by each branch of the phylogenetic tree. The near ties to "dark red" and the distance ties to "white" colour. (c) Neighbor-joining (NJ) tree for ToxA haplotypes based on the nucleotide sequence differences. Coloured internal nodes represent the bootstrap confidence level.

Fig. 3: Phylogenetic tree using Maximum Likelihood algorithm of different ToxA1 of *Pyrenophora tritici-repentis, Bipolaris sorokiniana,* and varipus haplotypes of *Parastagnospora nodorum*. The percentage of bootstrap support (1000 replicates) is indicated at corresponding nodes. The nomenclature of the haplotypes is based on the Aboukhaddour et al. (2023)

Fig. 4: Ramachandran plot and ProsSA z-score analysis of modelled ToxA1 protein from *B. sorokiniana*. Ramachandran distributions of amino acids in the model were calculated using pro check (a), and the z-score plot was plotted using ProsSA-WEB (b).

Fig. 5: *ToxA1* protein structure of *B. sorokiniana* isolate MCC 1533 (Pusa2) depicting conserved RGD motif (ARG-GLY-ASP)

Fig. 6 Symptom of leaf necrosis on Tsn1/tsn1 harbouring genotypes after infiltration ToxA1 protein at seedling (3-5 leaf stage) and on flag leaf of adult plant (Zadoks growth stage 55).

Table1: Response of wheat genotypes (*Tsn1/tsn1*) in terms of leaf necrosis after infiltration of ToxA1 protein at both seedling and adult stage of wheat genotypes harboring Tsn1/tsn1 allele.

Genotypes	Length(cm)		Width(cm)		Area(cm2)		Tsn1/tsn1
	Seedling	Adult	Seedling	Adult	Seedling	Adult	
K 68	2.872	4.613	0.404	0.530	0.977	2.326	*Tsn1*
C 306	1.921	4.226	0.329	0.633	0.678	1.737	*Tsn1*
HUW468	2.057	3.472	0.402	0.438	0.718	1.407	*Tsn1*
Yangmai 6	0.860	1.406	0.311	0.381	0.316	0.450	*Tsn1*
K65	1.513	2.582	0.380	0.566	0.388	1.096	*Tsn1*
WAMI160	1.615	3.461	0.345	0.506	0.646	1.521	*Tsn1*
WAMI188	1.693	2.582	0.310	0.500	0.585	1.036	*Tsn1*
WAMI224	1.643	2.857	0.330	0.488	0.476	1.298	*Tsn1*
WAMI226	1.782	2.320	0.405	0.503	0.628	1.090	*Tsn1*
WAMI277	1.464	2.174	0.360	0.485	0.464	1.024	*Tsn1*
WAMI288	1.591	2.054	0.432	0.575	0.622	1.017	*Tsn1*
Max.	2.872	4.613	0.432	0.633	0.977	2.326	
Min.	0.860	1.406	0.310	0.381	0.316	0.450	
HUW234	0.964	1.894	0.335	0.408	0.345	0.561	*tsn1*
PBW343	0.860	1.446	0.349	0.408	0.319	0.423	*tsn1*
LOK1	1.098	1.761	0.307	0.432	0.278	0.493	*tsn1*
Sonalika	1.919	2.923	0.361	0.492	0.584	1.321	*tsn1*
WAMI280	0.972	1.305	0.319	0.327	0.265	0.411	*tsn1*
WAMI289	1.065	1.471	0.319	0.344	0.342	0.529	*tsn1*
WAMI290	1.206	1.467	0.327	0.360	0.365	0.493	*tsn1*
Max.	1.919	2.923	0.361	0.492	0.584	1.321	
Min.	0.860	1.305	0.307	0.327	0.265	0.411	

Figure S1: Correlation coefficients estimated by point biserial correlation for phenotypic traits associated with*Tsn1/tsn1* gene effect in wheat genotypes with ToxA1 isolate (MCC1533/Pusa2)

A. Seedling Stage   B. Adult stage

Figure S2: The necrotrophic area developed due to ToxA1–Tsn1 interaction at seedling (3-5 leaf stage) and adult stage (Zadoks growth stage 55) of 11 wheat genotypes with *Tsn1* and 7 wheat genotypes with *tsn1*.

Figure S3. Symptom of leaf necrosis after infiltration ToxA1 protein at both seedling (3-5 leaf stage) and adult stage (Zadoks growth stage 55) of wheat genotypes harbouring *Tsn1*.

Figure S4. Symptom of leaf necrosis after infiltration ToxA1 protein at both seedling (3-5 leaf stage) and adult stage (Zadoks growth stage 55) of wheat genotypes harbouring *tsn1*