Heterologous Expression and Characterization of *ToxA1* haplotype from India and its

interaction with *Tsn1* for Spot Blotch Susceptibility in spring wheat

Ranjan Kumar Chaubey[1], Dharamsheela Thakur[2], Sudhir Navathe[3*], Sandeep Sharma[1],

Vinod Kumar Mishra[1], Pawan Kumar Singh[4*], Ramesh Chand*

[1] Institute of Agricultural Sciences, Banaras Hindu University Varanasi 221005, India

[2] Department of Molecular Biology and Genetic Engineering, Bihar Agricultural

University, Sabour 813210, Bhagalpur, India;

[3] Agharkar Research Institute, G. G. Agarkar Road, Pune 411004, India

[4] International Maize and Wheat Improvement Center (CIMMYT), Texcoco 56237, Mexico.

* Correspondence:

Ramesh Chand, Banaras Hindu University, Varanasi, India, Email: rc_vns@yahoo.co.in;

Sudhir Navathe, Agharkar Research Institute, Pune, India, Email: snavathe@aripune.org

Abstract

ToxA a necrotrophic effector protein, is known to be present in the genome of fungal species like *Parastagnospora   nodorum*, *Pyrenophora   tritici-repentis* and *Bipolaris sorokiniana*. *Tsn1* is the sensitivity gene in the host whose presence indicates more susceptibility to *ToxA* carrying pathogen, and *ToxA-Tsn1* delivers an inverse gene-for-gene relationship. The present study involved cloning and expressing the *ToxA1* gene from *B. sorokiniana*. It was found that the amplicon exhibited an expected product size of 471 bp. Sequence analysis of the *ToxA1* nucleotide sequence revealed the highest identity, 99.79%, with *Py. tritici-repentis*. The protein expression analysis showed peak expression at 16.5kDa. Phylogenetic analysis of *ToxA1* sequence from all the *Bipolaris* isolates formed independent clade along with *Py. tritici-repentis* and diverged from *P. nodorum*. *ToxA-Tsn1* interaction was studied in 18 wheat genotypes (11 *Tsn1* and 7 *tsn1*) at both the seedling and adult stage, validating the inverse gene-for-gene relationship, as the toxin activity was highest in the K68 genotype (*Tsn1*) and lowest in WAMI280 (*tsn1*). This shows a path towards more genomic regions involved in resistance to spot blotch disease, which could be further explored.

Keywords: *Bipolaris sorokiniana*, effector-triggered susceptibility, the inverse gene for gene relationship, sensitivity genes, *ToxA1*, *Triticum aestivum*

Introduction

Host selective toxins (HSTs) are the effector molecule released during the invasion of

necrotrophic pathogens and act as major determinants of pathogenicity (Ciuffetti et al. 1997).

These necrotrophic effectors are specific biomolecules that are structurally complex,

chemically diverse, and responsible for causing some key diseases in important crops. HSTs

are very important in host-pathogen interaction but are produced by limited genera of fungi

involving multiple toxins (Wolpert et al. 2002). Spot blotch of wheat is a threatening disease

caused by *Bipolaris sorokiniana*, a hemi-biotrophic pathogen. This disease is mainly prevalent

in South Asia's warm, humid region, especially in the eastern Indo-Gangetic plains of India.

Wheat productivity can decline by 40% due to spot blotch alone (Sharma et al. 2006). Spot

blotch has been reported to cause up to 17.5% yield loss in the Indian sub-continent, and about

25mha of wheat is affected by spot blotch worldwide (Pandey et al. 2021). Plant pathogens

have their genetic makeup for causing host infection, which is decided by the presence of a

single gene that confers host susceptibility or resistance (Balance et al. 1989, Friesen et al.

2006). The proteinaceous necrotrophic effector ToxA a major HST produced by *B.*

sorokininana. Apart from this, ToxA is known to be produced from two necrotrophic

pathogens, *P. nodorum* and *Py. tritici-repentis* and cause *Parastagnospora* nodorum blotch

and tan spot, respectively (Solomon et al. 2006, Tuori et al. 1995). ToxA protein was originally

reported in a culture filtrate of the wheat pathogen *Py. tritici-repentis* (Tuori et al. 1995). The

ToxA protein was first discovered by Tomás and Bockus (1987) from the culture filtrate of *Py.*

tritici-repentis. The *ToxA* manifested high nucleotide diversity in *S. nodorum,* whereas it

showed a lack of variation in *Py. tritici-repentis*. This pattern of nucleotide diversity and the

recent emergence of tan spot suggests the recent introduction of *ToxA* into *Py. tritici- repentis*

genome (Friesen et al. 2006). The *ToxA* sequences from *S. nodorum* and *Py. tritici-repentis*

depicted differences at four fixed nucleotide sites resulting in two predicted amino acid changes

consistent with the interspecific transfer of this gene from *S. nodorum* into *Py. tritici-repentis*

(Friesen et al. 2006), and this *ToxA* gene jumped horizontally from *Py. tritici-repentis* to *B.*

sorokiniana, (McDonald et al. 2018). *ToxA* and its surrounding 14 kb suggest that this gene

was horizontally transferred, facilitated by a type II DNA transposon (McDonald et al. 2019).

ToxA is a single-copy gene, and it was found to be embedded within a 12-kb genomic element

nearly identical to the corresponding regions in *P. nodorum* and *Py. tritici-repentis* (McDonald

et al. 2018). The mature ToxA protein is 13.2 kDa, and it encodes a signal peptide of 23 amino

acids and a pro-domain of 4.3kDa (required for protein folding), and both are cleaved before

the secretion of mature ToxA protein (Balance at al. 1996, Ciuffetti et al. 1997, Tuori et al.

2000). Studies revealed that the host sensitivity of ToxA was mapped on wheat chromosome

arm 5BL, and the sensitivity gene was named *Tsn1* (Faris et al. 2010). Further, it was confirmed

that this gene was significant for both tan spot and SNB, and in both cases, the pathogen had a

sensitivity gene such as *ToxA* (Lamari and Bernier 1989, Friesen et al. 2006). It is considering

the wheat-*B. sorokiniana* pathosystem, *ToxA* confers virulence, and it is the only known

virulence gene whose corresponding sensitivity gene (*Tsn1*) in the host is identified. Further,

it was also demonstrated that *Tsn1*-carrying wheat cultivars are more sensitive to *B.*

sorokiniana isolates that harboured the *ToxA* gene. Cloning of the *Tsn1* gene unfolded

characteristics similar to classical disease resistance nucleotide-binding leucine-rich repeat

NLR gene that included serine/threonine kinase, leucine-rich repeats, nucleotide binding and

protein kinase domains (Faris et al. 2010). The ToxA gene is transcribed as a necrotrophic

effector protein responsible for infection in the host, with *Tsn1* being a resistance gene (Tan et

al. 2016). A resistance-like gene *Tsn1* in wheat is responsible for conferring susceptibility to

ToxA in *Py. tritici-repentis* and *P. nodorum* isolates (Friesen et al. 2006). This *ToxA–Tsn1*

interaction system has been proved as an example of the inverse gene-for-gene relationship. It

was realizing the occurrence of spot blotch in South Asia and the emergence of *ToxA1* as a key

virulence gene driven us towards studying this host-pathogen relation. Keeping this in mind,

we cloned the *ToxA1* gene from *B. sorokiniana* and conducted expression analysis for the

ToxA1 protein. We also validated the pathogenicity of *ToxA1* on the allelic status of *Tsn1* by

using 18 *Tsn1/tsn1* genotypes.

Materials and Methods

Primer designing:

The *ToxA* sequence from *Py. tritici-repentis* (U79662.1) was obtained in FASTA format from

NCBI (National Centre for Biotechnology Information) GenBank database

(https://www.ncbi.nlm.nih.gov/genbank/). *ToxA* is a 534bp-long gene sequence, and primers

were designed using the web-based tool Primer-BLAST (Basic Local Alignment Search Tool)

from NCBI. The primers were utilized to amplify *ToxA1* in *B. sorokiniana* isolate MCC 1533

(Pusa2). The isolate MCC1533 (Pusa 2) has been identified and deposited at National Center

for Microbial Resources (NCMR), Pune, India. The primer sequence used is given as Forward

Primer (FP) -CGCGGATCCGATCCCGGTTACGAAATCG and Reverse Primer (RP) - CGC

CTCGAGCTAATTTTCTAGCTGCATTCTC

RNA extraction and cDNA synthesis

B. sorokiniana isolate MCC 1533 (Pusa2) was cultured for 10 days in a Potato Dextrose Broth.

Mycelium was harvested, and the total RNA was isolated using Invitrogen PureLink^{TM} RNA

Mini kit following the manufacturer's protocol. The integrity and size distribution of total RNA

purified with RNeasy Kit was checked by 1.2% agarose gel electrophoresis and ethidium

bromide staining visualizing two sharp bands of 28S rRNA to 18S RNA. The nanodrop results

were obtained with the Nanodrop2000 instrument (Thermo Scientific), which showed the

acceptable purity of RNA isolated from each biological replicate (A_{260}/A_{280} of 1.9 to 2.1). The

isolated RNA was stored immediately at -80°Cin nuclease-free water (supplied with the kit)

for further analysis. cDNA was synthesized using a Thermo cDNA synthesis kit following the

manufacturer's protocol.

RNA previously isolated from the biological sample was taken as a template for cDNA

synthesis. An additional treatment of RT enhancer was included during cDNA synthesis as

supplied and instructed with the kit and manufacturer's protocol to prevent chances of

contamination of genomic DNA during isolation. The template RNA sample was added to the

cDNA reaction mixture, which was obtained by diluting the biological samples of RNA.

ToxA gene cloning

The expression constructs were produced in the bacterial expression vector pET28a that directs

the expression of HIS-tagged fusion proteins in *E. coli*. BamH1 and Xho restriction sites were

introduced in forward and reverse primer for cloning in PET vector. Primers were designed

such that the PCR amplified portion of the *ToxA* gene was generated with a 5' *Bam*HI site and

a 3' *Xho* site. The PCR products were digested with *Bam*HI and *Xho*, gel-purified with the

QIAquick Gel Extraction Kit (Qiagen, Valencia, CA) and cloned into the *Bam*HI and *XhoI*

sites of pET28a. Primers FP-CGCGGATCCGATCCCGGTTACGAAATCG and RP- CGC

CTCGAGCTAATTTTCTAGCTGCATTCTC were used for the PCR amplification. PCR

amplification was performed using cDNA as a template in an Eppendorf Mastercycler using

the following cycles; initial denaturation of 95°C for 7 min followed by 35 cycles of 94°C (30

s), 55°C (45 s) and 72°C (1 min) and a final extension of 10 min at 72°C. The PCR product

was gel-purified, digested with *Bam*HI and *XhoI*, and ligated into the respective restriction sites

of pET28a. The resulting construct contained the predicted effector protein gene fused to a 6X

histidine tag at the N-terminus. The ligated product was used to transform *E. coli* DH5alpha

and BL21 (DE3) strains (Sambrook et al. 2006) for cloning and protein expression.

Sequencing and characterization

Cloned fragment of *ToxA1* from *B. sorokiniana* isolate MCC1533 (Pusa2) was sequenced by

Sanger sequencing. The sequences obtained in two frames were merged, and overlapping

sequences were identified in both directions using Bioedit Sequence Alignment Editor (Hall

1999). Merged sequences were then prepared in a single frame. The sequences have been

deposited to the National Centre for Biotechnology Information (NCBI) database with Gene

Bank accession number (OP289649). The processed *ToxA1* sequence was used as a query for

BLASTn (https://blast.ncbi.nlm.nih.gov/Blast.cgi) for finding homologous sequences with

reference to the NCBI nucleotide database (Altschul et al. 1997). The processed *ToxA1*

nucleotide sequence was then translated to the amino acid sequence in BioEdit 7.2. Sequences

of *Py. tritici-repentis, P. nodorum* and different isolates of *B. sorokiniana* were retrieved from

the NCBI protein database. The amino acid sequences retrieved using BLASTp searches were

then used to create Multiple sequence alignments of 27ToxA sequences. MEGA X (Molecular

Evolution Genetics Analysis) software (Kumar et al. 2018) was then used to create a phylogeny

of *ToxA* amino acid sequences of 49 isolates of *Py. tritici-repentis, P. nodorum* and *B.*

sorokiniana using the CLUSTAL W alignment algorithm (Larkin et al. 2007). All the

alignment settings were employed at default values; 178 and 114 positions with and without

gaps were obtained, respectively. The substitutions selected with complete deletion of gaps or

missing data were used to analyze sequences. The phylogenetic tree was inferred using the

maximum likelihood method based on the JTT model (Jones et al. 1992). The initial tree was

inferred with the default setting using the Neighbour Joining method, and Nearest-Neighbour

Interchange was used as an ML heuristic search method. The reconstructions of phylogenetic

trees were conducted using the Maximum Likelihood Method. Bootstrap with 1000 replicates

for the Poisson correction model was performed to assess node support (Felsenstein 1985).

Protein modelling and topographical properties

The predicted protein sequence of cloned Pusa-2 *ToxA1* was subjected to the SWISS-MODEL

homology modelling online tool (Arnold et al. 2006). Proteins were modelled utilizing Protein

Data Bank (PDB) entry 1zld.1.A (*Py. tritici-repentis*) as reference. The quality of modelled

protein was judged based on GMQE (Global Model Quality Estimation) and QMEAN

(Qualitative Model Energy Analysis) Z-score parameters (Benkert et al. 2009, Studer et al.

2020). The best model was selected based on the GMQE score, which ranges from 0 to 1, with

a higher value indicating better reliability. The QMEAN Z-score estimates the "degree of

nativeness" of the structural features observed in the model on a global scale. QMEAN Z-

scores around zero indicate good agreement between the model structure and experimental

structures. Scores of -4.0 or below is an indication of models with low quality. The quality of

the resulting models was also monitored with PROCHECK and ProSSA-WEB (Laskowski et

al. 1993). Geometrical and topographical properties of the modelled proteins were identified

using the Computed Atlas of Surface Topography of proteins (CASTp) 3.0 online server (Tian

et al. 2018). Molecular graphics images were produced using the UCSF Chimera package

(Pettersen et al. 2004). The alpha helix and the beta-sheet percentage were analyzed using

PDBMD2CD.

Isolation of proteins from E. coli cultures

E. coli cultures harbouring all expression constructs were grown overnight in 5 ml of LB

(Sambrook et al. 2006) with 100μg of kanamycin per ml (LBA) at 37°C and shaking at 250

rpm (G24 Incubator Shaker, New Brunswick Scientific, Edison, NJ). Two hundred and fifty

microliters of the overnight culture were used to inoculate 25 ml of fresh LBA and incubated

with shaking at 37°C for 1 h. Expression of fusion proteins was induced by the addition of

isopropyl-beta-Dthiogalactopyranoside (IPTG) to a final concentration of 1 mM, and the

cultures were incubated for an additional 4hrs under the same conditions. Cells were harvested

by centrifugation at 4,000× *g,* 4°C for 20 min and the pellets were resuspended in cold

sonication buffer (SB; 50 mM Na2HPO4, 5 mM Tris-base, 300 mM NaCl, pH 8.0) with 0.5%

Triton-x-100 and 10 mM DTT. Resuspended pellets were sonicated three times, 30 s each, with

2 min on ice between periods. Insoluble material was then pelleted by centrifugation at 10,000

$\times g$, 4°C, for 10 min. Pellets were washed thrice with SB with 0.5% Triton-$\times$ -100 and 10 mM

DTT. The final washed pellets, containing inclusion bodies, were either stored frozen at –80°C

or resuspended in 1 ml of 8 M guanidine HCl (Sigma) in refolding buffer (250 mM NaCl, 100

mM sodium phosphate buffer, 10 mM Tris-base, 4% glycerol, 1 mM EDTA, and 0.005 %

Tween 20, pH 8.0) with freshly prepared DTT added to a final concentration of 10 mM.

Solubilized proteins were sonicated briefly twice (for 5 to 10 s) and then heated to 37°C for 10

min and 50°C for an additional 5 min to assist in the solubilization of proteins. Insoluble

material was removed from the samples by centrifuging at 12,000 $\times g$ for 10 min. Then,

solubilized proteins were quantitated with Bradford assay reagent (Bio-Rad, California, USA)

with bovine serum albumin used as a standard. Fusion proteins solubilized in guanidine HCl

were used immediately.

Screening of ToxA1 –Tsn1 interaction on different wheat genotypes

The screening of 18 wheat genotypes was conducted for *Tsn1/tsn1,* out of which 11 genotypes

harbouring Tsn1 and the rest 7 genotypes with *tsn1* were sown in 3 replications at polyhouse,

Plant Pathology, Institute of Agricultural Sciences, Banaras Hindu University in 1m rows at a

distance of 22.5cm. Infiltration of ToxA1 protein, as well as control (Uninduced protein), was

done at both the seedling stage (3-5 leaf stage) and adult stage (Zadoks 55). 250µg of ToxA

protein was infiltrated by a sterilized syringe, and symptoms were recorded 7 days after

infiltration. Photographic data was captured using a Nikon camera. The size of the lesion was

measured using ImageJ software.

Statistical Analysis

The lesion size was measured in length, width, and area of lesions. The statistical analysis was

conducted for phenotypic traits, including length, width, and area of necrotic lesions of wheat

genotypes with the *Tsn/tsn1* gene at both seedling and adult stages. The correction coefficients

between disease components were calculated using R software V 4.2.1 (R Core Team, 2022).

Results

Cloning and Expression study of ToxA1

The *ToxA* gene from *B. sorokiniana* was amplified using primers designed from *PtrToxA*

through RT-PCR. The amplicon exhibited the expected product size of 471bp, which was then

cloned in the pET28avector. The cloned gene was validated using colony PCR of the

transformed cell. The *ToxA1* gene, after validation, was subjected to sequencing, and the

sequence data after processing revealed successful cloning of the *ToxA* gene (now *ToxA1- B.*

sorokininana). This ToxA1 nucleotide sequence was further used for BLASTn analysis,

revealing the highest identity, 99.79%, with *Py. tritici-repents* and E-value close to zero. The

cloned gene was then subjected to their expression at different concentrations of IPTG, namely,

at 0.4mM, 0.6mM, 0.8mM and 1.0mM, and expression of the *ToxA1* gene was highest at

1.0mM concentration with an expected band size of 16.5kDa in sodium dodecyl sulfate (SDS)-

polyacrylamide gels (Fig. 1).

ToxA1 characterisation and Phylogenetic analysis

The identification of *ToxA1* after showing sequence similarity with *Py. Tritici-repentis* ToxA

using BLASTn analysis on NCBI was conducted. Further characterisation of ToxA1 was

conducted by translating the nucleotide sequence to amino acid sequence using the web-based

tool Expasy (https://www.expasy.org/resources/translate). The amino acid sequence of ToxA1

was subjected to BLASTp analysis on NCBI, which depicted 100% similarity with *Py. tritici-*

repentis showing E value 9e-109. Multiple sequence alignment of 25ToxA protein sequences

and *ToxA1* MCC1533 (Pusa2) translated sequence in Bioedit revealed that amino acid

differences were found in most *P. nodorum* strains. However, the cloned Pusa2_*ToxA1*

sequence showed complete similarity to *Py. tritici-repentis*. *P. nodorum* depicted amino acid

substitution for Glutamate, Aspartate, Leucine, Valine and Asparagines (Fig. 2).

The phylogenetic analysis depicts the evolutionary relatedness of the ToxA1 amino

acid sequence with other ToxA sequences. The tree indicates that ToxA sequences originate

from a common ancestor but form different clades during evolution. The ToxA sequence from

all the *Bipolaris* isolates formed an independent clade along with *Py. tritici-repentis* and

diverged from *P. nodorum*. This indicates sequence similarity among *B. sorokiniana* and *Py.*

tritici-repentis, whereas it differed from *P. nodorum* (Fig. 3).

Protein modelling and Topographical properties

The structure of the ToxA1 protein of a cloned gene of *B. sorokiniana* isolate MCC1533

(Pusa2) was predicted using the SWISS-MODEL homology modelling online server. For

checking the reliability of models GMQE (ToxA1: 0.56), QMEAN Z scores are included

QMEAN -0.33, Cbeta –1.79, torsion 0.53 and these scores indicated that these models are

reliable and good quality. The quality of the resulting models was monitored with

PROCHECK. Ramachandran plot analysis revealed that 88.1% of the ToxA1 protein structure

residues fell within the most favoured regions, with a further 11.9% in the additionally allowed

region (Fig. 4). No residues were found in the generously allowed or disallowed regions. The

number of non-glycine and non-proline residues was 100%. The number of end residues

(excluding Gly and Pro) is 2, and the number of glycine and proline residues was 12 and 3,

respectively.

Predicted ToxA1 protein structures of *B. sorokiniana* MCC1533 (Pusa2) isolate based

on homology modelling are depicted in Fig. 4. The protein sequences of ToxA1 protein showed

sequence identity of 100% with template PDB entry 1zld.1.A (*Py. tritici-repentis*, *PtrToxA*).

When generated structures were compared in UCSF Chimera, it was observed that the RGD motif (Toxic region) was present in the 3D structure of ToxA1 protein and found that arginine is present in the alpha helix of the loop at 118 positions. At the same time, Glycine and Aspartic acid is present in the β-sheet of the protein structure at 119 and 120 positions of ToxA1 protein (Fig. 5). The percentage of the alpha helix was 3.39%. At the same time, antiparallel sheet 1 was 35.55%, antiparallel sheet 2 was 12.76%, the parallel sheet was 0.85%, the turn was 7.63%, and the other was 39.83%.

Bioactivity of ToxA1 effector protein and its interaction with Tsn1 at both seedling and adult stage

The bioactivity of the *ToxA1* effector was studied on necrotic lesions symptoms caused due to *ToxA1* protein and was found to be more prominent at the adult stage. At the seedling stage, the maximum area (0.977 cm^2) was recorded in genotype K68(*Tsn1*) and the minimum 0.265 cm^2 in WAMI280 (*tsn1*). At the same time, the length varies from 2.872 cm (K68, *Tsn1*) to 0.860cm (PBW343, *tsn1*). However, the maximum width recorded was 0.432 cm for genotype WAMI288 (*Tsn1*), and the minimum in genotype harbouring *tsn1*(LOK1) was 0.307cm.

At the adult stage, the length of the symptom was maximum (4.613cm) in genotypes harbouring*Tsn1*(K68) and minimum (1.305 cm) WAMI280 (*tsn1*), and the width also showed the same trend varies from 0.633cm (C306, *Tsn1*) to 0.327cm (WAMI280, *tsn1*). The area was measured for quantifying the necrotrophic symptom, and it found that the maximum area in the genotype harboring *Tsn1* was 2.326 cm^2 (K68), and the minimum was 0.411 cm^2(WAMI280, *tsn1*). A significant difference was found between the seedling and adult stages and for *Tsn1* and *tsn1* (Table 1, Fig. S2, S3, S4).

Discussion

Studies on necrotrophic effectors producing HSTs have provided a new dimension to host-pathogen interaction and established an inverse gene-for-gene relationship (Friesen et al. 2006,

Navathe et al. 2020). The HSTs are effectors proteins and induce a susceptible response in the

host harbouring the dominant susceptibility gene. ToxA is a necrotrophic effector protein, one

of the key factors for causing the most devastating disease, spot blotch in wheat, prevalent in

warm, humid regions of South Asia. The present study focuses on cloning of *ToxA1* gene,

followed by its characterization and purification of the resultant protein and its expression

analysis. In this study, we have successfully cloned and characterized the *ToxA1* gene from *B.*

sorokiniana isolate MCC1533 (Pusa2). This gene was originally found in *P. nodorum* and

horizontally transferred to *Py. tritici-repentis* and, more recently, to *B. sorokiniana* (Friesen et

al. 2006, McDonald et al. 2012) . The cloned gene and its expression were studied in the present

study, and the purified protein showed peak expression at a concentration of 1.0 mM IPTG

with band migration at 16.5kDa in size. Manning et al. (2008) also reported similar findings

where lysates with His-ToxA1 or without (control) have a phosphorylated band that migrates

at 16 kDa. However, Mass spectral analyses have indicated a 13.2 kDa mass of purified mature

ToxA1 protein from *Py. tritici-repentis* (Ciuffetti et al. 1997, Tuori et al. 1995). The importance

of a protein purification study for ToxA1, which has selective host toxicity for genetic analysis

of pathogenicity in the *Pyrenophora*-wheat interaction, was also suggested by Touri et al.

(1995).

The *ToxA1* gene was truncated in the present study to remove the signalling sequences, which

comprise 22 amino acids, as the mature ToxA1 gene is released after the proteolytic cleavage

of this signal sequence. Previous studies have also described these signalling peptides in ToxA

protein that involve the first 22 amino acids, which are proteolytically cleaved after import into

the endomembrane system (Ciuffetti et al. 1997). The release of mature toxin involves

additional proteolytic processing, and as the ToxA1 protein is blocked at the N terminus, such

modifications would incur a mass increase (Ciuffetti et al. 1997). The ToxA1 cloned sequence

obtained in our study has been compared with the already reported ToxA1 sequences from *P.*

nodorum, *Py. tritici-repentis* and *B. sorokiniana*. Multiple sequence alignment of amino acid

sequences from 26 isolates was performed. The results suggest that the ToxA region from *B.*

sorokiniana and *Py. tritici-repentis* show evolutionary origin relatedness to each other than to

P. nodorum. This is based on the identical ToxA1 sequence shared between *B. sorokiniana* and

Py. tritici-repentis and the difference at six positions in *P. nodorum* haplotypes (Fig. 2). The

sequence identity between the three fungal species accounts for the common evolutionary

origin of ToxA and the potential of exchanging DNA between these species (McDonald et al.

2018). The significance of phylogenetic studies in establishing evolutionary relationships

among the different pathogen races or deriving the specific gene evolution between different

genera is well known. The sequencing data for ToxA1 showed a high degree of similarity

between *B. sorokiniana* and *Py. tritici-repentis* (Hafez et al. 2022, Aboukhaddour et al. 2023).

Previous studies have also suggested that the ToxA1 sequence did not differ for *B. sorokiniana*

and *Py. tritici-repentis* but differed from the ToxA sequence of *P. nodorum* (Navathe et al.

2020, Hafez et al. 2022, Aboukhaddour et al. 2023). This evolutionary relatedness of *Py. tritici-*

repentis and *B. sorokiniana* sequences have also been reported earlier (McDonald et al. 2018,

Navathe et al. 2020). The earlier report suggested that ToxA has been present in *P. nodorum*

for a long time as it exhibited greater sequence diversity and was interspecifically transferred

to *Py. tritici-repentis* quite later. This is evident from monomorphism or lesser sequence

diversity in the ToxA1 sequence from *Py. tritici-repentis* and the emergence of tan spot

symptoms more recently in the near past (Aboukhaddour et al. 2023).

Further insights into the ToxA structural analysis were performed by predicting the 3D protein

structure using an amino acid sequence of ToxA1 (Fig.4). This is the first report for the 3D

protein structure ToxA1 haplotype from *B. sorokiniana*. The predicted protein structure was

similar to a single-domain protein with a β-sandwich fold of PtrToxA. This β-fold represents a

solvent-exposed loop containing arginyl-glycol-aspartic acid (RGD)- functionally active motif

[(Sarma et al 2005, Ciuffetti et al. 2010). ToxA1 has been known to resemble with fibronectin

III domain which retains a solvent-exposed loop containing RGD motif for cell wall-plasma

membrane interaction in plants (Ciuffetti et al. 2010, McDonald and Solomon 2018). It has

been proposed that ToxA with RGD-containing motif has been known to interact with wheat

mesophyll cells and thus establish disease (Manning et al. 2007) The mesophyll cells have a

high-affinity matrix to recognise these motifs, and these all suggest the importance of the RGD

motif in receptor binding and internalisation into wheat cells (Ciuffetti et al. 2010).

ToxA is the only known virulence gene for which a host sensitivity gene (*Tsn1*) has been known

in wheat-*B. sorokiniana* pathosystem. For the first time, the present study reports a cloned and

purified protein ToxA1 infiltration in both seedling and adult stages of wheat lines harbouring

Tsn1/tsn1. The severity was found to be enhanced in the adult stage compared to the seedling

stage in *Tsn1* wheat lines. The present data also reveals that *Tsn1*-carrying genotypes exhibited

more disease in seedlings than in the adult stage. The toxin activity was highest in the K68

genotype (*Tsn1*) and lowest in WAMI280 (*tsn1*) at both the seedling and adult stages. This

could be validated by past studies where it has been suggested that ToxA causes severe necrotic

lesions on *Tsn1,* harbouring wheat genotypes, while less or no lesion was observed for wheat

genotypes lacking the sensitivity gene (Aggarwal et al 2021). Navathe et al. (2020) also

reported one such possibility of the involvement of a recessive allele of *Sb2* towards an

unknown HST. *Tsn1* has been reported attributing susceptibility towards both *P. nodorum*-

causing SNB (Friesen et al 2006, Virdi et al. 2016) and *Py. tritici-repentis* are causing tan spots

(Faris et al. 2013). However, the environmental requirements for *Bipolaris* are different from

these two pathogens, which are not found in the South-Asian region.

Nevertheless, it would be desirable for wheat breeders to select *tsn1* for making crops resistant

to these three pathogens. The study in this horizon should focus on the presence of loci other

than *Tsn1* and *Sb2*. It would be interesting to study other quantitative trait loci involved to understand the complexity of this *ToxA-Tsn1* interaction.

Acknowledgements: Authors acknowledge Banaras Hindu University and Bihar Agricultural University for the infrastructure facility
Funding: SN acknowledges SERB, New Delhi, for CRG/2021/007211.

Author Contributions: Conceptualization, RC, VKM, PKS, SN, and SK; methodology, RKC and DT; validation, DT and SKS; formal analysis DT and SKS; investigation, RKC; resources, RC, DT; data curation, DT, SKS; writing—original draft preparation, RKC, SN; writing—review and editing, RC, PKS; supervision, RC, VKM, SKS; project administration, RC; funding acquisition, RC, PKS. All authors have read and agreed to the published version of the manuscript.

Availability of data and materials: Not applicable.
Declarations
Ethics approval and consent to participate: Not applicable.
Consent for publication: Not applicable.
Competing interests: The authors declare that they have no competing interests.

Ciuffetti, L. M., Tuori, R. P. Gaventa, J. M. A single gene encodes a selective toxin causal to the development of tan spot of wheat. Plant Cell, 1997, *9*, 135-144. https://doi.org/10.1105/tpc.9.2.135

Wolpert, T.J., Dunkle, L.D., Ciuffetti, L.M. Host-selective toxins and avirulence determinants: what's in a name? Annu. Rev. Phytopathol. 2002, 40, 251–285. https://doi.org/10.1146/annurev.phyto.40.011402.114210

Sharma, P., Duveiller, E., Sharma, R. C. Effect of mineral nutrients on spot blotch severity in wheat and associated increases in grain yield. Field Crops Res, 2006, *95*, 426-430. https://doi.org/10.1016/j.fcr.2005.04.015

Pandey AK, Mishra VK, Chand R, Navathe S, Budhlakoti N, Srinivasa J, Sharma S, Joshi AK. Crosses with spelt improve tolerance of South Asian spring wheat to spot blotch, terminal heat stress, and their combination. Sci Rep, 2021, *11*, 1-12. https://doi.org/10.1038/s41598-021-85238-x

Ballance, G.M., Lamari, L., Bernier, C.C. Purification and characterization of a host-selective necrosis toxin from Pyrenophora tritici-repentis. Physiol. Mol. Plant Pathol. 1989, *35*, 203–213 https://doi.org/10.1016/0885-5765(89)90051-9

Friesen, T. L., Stukenbrock, E. H., Liu, Z., Meinhardt, S., Ling, H., Faris, J. D., ... Oliver, R. P. Emergence of a new disease as a result of interspecific virulence gene transfer. Nat. Genet. 2006, *38*: 953-956. https://doi.org/10.1038/ng1839

McDonald, M. C., & Solomon, P. S. Just the surface: advances in the discovery and characterization of necrotrophic wheat effectors. Curr. Opin. Microbiol. 2018, *46*, 14-18.

Solomon, P., Lowe, R., Tan, K.C., Waters, O., Oliver, R. *Stagonospora nodorum*; cause of Stagonospora nodorum blotch of wheat. Mol. Plant Pathol. 2006, *7*, 147–156 https://doi.org/10.1111/j.1364-3703.2006.00326.x

Tuori, R. P., Wolpert, T. J. Ciuffetti, L. M. Purification and immunological characterization of toxic components from cultures of *Pyrenophora tritici-repentis*. Mol. Plant Microbe Interact, 1995, *8*, 41-48. https://doi.org/10.1094/mpmi-8-0041

Tomás, A., Bockus, W. W. Cultivar specific toxicity of culture filtrate of *Pyrenophora tritici-repentis*. Phytopathology, 1987, *77*,1337–1366. https://doi.org/10.1094/phyto-77-1337

McDonald, M. C., Ahren, D., Simpfendorfer, S., Milgate, A. Solomon, P. S. The discovery of the virulence gene ToxA in the wheat and barley pathogen *Bipolaris sorokiniana*. Mol. Plant Pathol., 2018, *19*, 432-439. https://doi.org/10.1111/mpp.12535

Ballance, G. M., Lamari, L., Kowatsch, R., Bernier, C. C. Cloning, expression and occurrence of the gene encoding the Ptrnecrosistoxin from *Pyrenophora tritici-repentis*. Mol. Plant Pathol. 1996, https://doi.org/10.1007/978-94-011-5218-1_21

Tuori, R. P., Wolpert, T. J. Ciuffetti, L. M. Heterologous expression of functional PtrToxA. Mol. Plant Microbe Interact, 2000, *13*, 456-464. https://doi.org/10.1094/mpmi.2000.13.4.456.

Faris, J. D., Zhang, Z., Lu, H., Lu, S., Reddy, L., Cloutier, S., ... Friesen, T. L. A unique wheat disease resistance-like gene governs effector-triggered susceptibility to necrotrophic pathogens. PNAS, 2010, *107*, 13544-13549. https://doi.org/10.1073/pnas.1004090107

Lamari, L., Bernier, C. C. Evaluation of wheat lines and cultivars to tan spot [*Pyrenophora tritici-repentis*] based on lesion type. Comput. Appl. Biosci. 1989, 11, 49-56. https://doi.org/10.1080/07060668909501146

Tan, K. C., Ferguson-Hunt, M., Rybak, K., Waters, O. D., Stanley, W. A., Bond, C. S., Stukenbrock, E. H., Friesen, T. L., Faris, J. D., McDonald, B. A., Oliver, R. P. Quantitative variation in effector activity of ToxA isoforms from *Stagonospora nodorum* and *Pyrenophora tritici-repentis*. Mol. Plant Microbe Interact. 2012, *25*, 515-522. https://doi.org/10.1094/mpmi-10-11-0273

Sambrook, J., Russell, D. W., Sambrook, J. The condensed protocols: from molecular cloning: a laboratory manual, 2006, (No. Sirsi) i9780879697723). Cold Spring Harbor, NY: Cold spring harbour laboratory press. https://doi.org/10.1101/pdb.prot4018

Hall, T. A. A user-friendly Biological Sequence Alignment Editor and Analysis program for Windows 95/98/NT. Nucleic Acids Symp Ser, 1999, *41*, 95–98

Altschul S. F., Madden T.L., Schäffer A.A., Zhang J., Zhang Z., Miller W., Lipman D. J. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. Nucleic Acids Res. 1997, *25*, 3389–3402. https://doi.org/10.1093/nar/25.17.3389

Kumar S., Stecher G., Li M., Knyaz C., Tamura K. MEGA X: MolecularEvolutionary Genetics Analysis across computing platforms. Mol Biol Evol. 2018, *35*, 1547–1549. https://doi.org/10.1093/molbev/msy096

Larkin M.A., Blackshields G., Brown N.P., Chenna R., McGettigan P.A., McWilliam H., Valentin F., Wallace I.M., Wilm A., Lopez R., Thompson J.D. Clustal W and Clustal X version 2.0. Bioinformatics, 2007, *23*, 2947–2948. https://doi.org/10.1093/bioinformatics/btm404

Jones DT, Taylor WR, Thornton JM. The rapid generation of mutation data matrices from protein sequences. Comput. Appl. Biosci. 1992, *8*, 275-282. https://doi.org/10.1093/bioinformatics/8.3.275

Felsenstein. Confidence limits on phylogenies: an approach using the bootstrap. Evolution
1985, *39*, 783–791. https://doi.org/10.1111/j.1558-5646.1985.tb00420.x

Arnold K, Bordoli L, Kopp J, Schwede T. The SWISS-MODELworkspace: a web-based
environment for protein structure homology modelling. Bioinformatics 2006, *22*, 95–
201. https://doi.org/10.1093/bioinformatics/bti770

Benkert P, Tosatto SCE, Schwede T. Global and local model quality estimation at CASP8
using the scoring functions QMEAN and QMEANclust. Proteins Struct Funct
Bioinforma. 2009, *77*, 173–180. https://doi.org/10.1002/prot.22532

Studer G, Rempfer C, Waterhouse AM, Gumienny R, Haas J, Schwede T. QMEAND is Co—
distance constraints applied on model quality estimation. Bioinformatics, 2020, *36*,1765–
1771. https://doi.org/10.1093/bioinformatics/btz828

Laskowski RA, MacArthur MW, Moss DS, Thornton JM. PROCHECK: a program to check
the stereochemical quality of protein structures. J. Appl. Crystallogr. 1993, 26, 283–291
https://doi.org/10.1107/s0021889892009944

Tian W, Chen C, Lei X, Zhao J, Liang J. CASTp 3.0: computed atlas of surface topography of
proteins. Nucleic Acids Res. 2018, *46*, W363–W367. https://doi.org/10.1093/nar/gky473

Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, Ferrin TE. UCSF
Chimera. A visualization system for exploratory research and analysis. J Comput Chem.
2004, *25*, 1605–1612 https://doi.org/10.1002/jcc.20084

R Core Team. R: A language and environment for statistical computing. 2022, R Foundation
for Statistical Computing, Vienna, Austria. https://www.R-project.org/.

Navathe S, Yadav PS, Chand R, Mishra VK, Vasistha NK, Meher PK, Joshi AK, Gupta PK.
ToxA–Tsn1 interaction for spot blotch susceptibility in Indian wheat: An example of
inverse gene-for-gene relationship. Plant dis, 2020, *104,* 71-81.
https://doi.org/10.1094/pdis-05-19-1066-re

McDonald, M.C., Razavi, M., Friesen, T.L., Brunner, P.C., McDonald, B.A. Phylogenetic and
population genetic analyses of *Phaeosphaeria nodorum* and its close relatives indicate
cryptic species and an origin in the Fertile Crescent. Fungal Genet. Biol. 2012, *49*, 882–
895. https://doi.org/10.1016/j.fgb.2012.08.001

Manning, V. A., Hamilton, S. M., Karplus, P. A., Ciuffetti, L. M. The Arg-Gly-Asp–
containing, solvent-exposed loop of PtrToxA is required for internalization. Mol. Plant
Microbe Interact. 2008, *21*, 315-325. https://doi.org/10.1094/mpmi-21-3-0315

McDonald, M. C., Solomon, P. S.   Just the surface: advances in the discovery and
characterization of necrotrophic wheat effectors. Curr. Opin. Microbiol. 2018, 46, 14-18.

Hafez, M., Despins, T., Nakajima, K., Aboukhaddour, R. Identification of a novel ToxA
haplotype of *Pyrenophora tritici-repentis* from Japan. Phytopathology, 2022, *112*, 1597-
1602 https://doi.org/10.1094/phyto-01-22-0001-sc

Aboukhaddour, R., Abdel-Fattah, M. H., McDonald, M., Moffat, C., Navathe, S., Friesen, T.
L., Strelkov, S. E., Oliver, R. P., Tan, K.-C., Liu, Z., Moolhuijzen, P., Phan, H., See, P.
T., Solomon, P. A revised nomenclature for ToxA haplotypes across multiple fungal
species. Phytopathology, 2023, https://doi.org/10.1094/phyto-01-23-0017-sc

Sarma GN, Manning VA, Ciuffetti LM, Karplus PA. Structure of PtrToxA: an RGD-containing
host-selective toxin from *Pyrenophora tritici-repentis*. Plant Cell, 2005, *17*, 3190–3202.
https://doi.org/10.1105/tpc.105.034918

499 Ciuffetti, L. M., Manning, V. A., Pandelova, I., Betts, M. F., Martinez, J. P. Host‑selective
500   toxins, PtrToxA and PtrToxB, as necrotrophic effectors in the *Pyrenophora tritici*‑
501   *repentis*‑wheat interaction. New Phytol. 2010, *187*, 911-919.
502   https://doi.org/10.1111/j.1469-8137.2010.03362.x

503 Baluska, F., Samaj, J., Wojtaszek, P., Volkmann, D., Menzel, D. Cytoskeleton-plasma
504   membrane-cell wall continuum in plants. Emerging links revisited. Plant Physiol. 2003,
505   *133*, 482-491. https://doi.org/10.1104/pp.103.027250

506 Manning, V. A., Hardison, L. K., Ciuffetti, L. M. PtrToxA interacts with a chloroplast-
507   localized protein. Mol. Plant Microbe Interact. 2007, *20*, 168-177.
508   https://doi.org/10.1094/mpmi-20-2-0168

509 Aggarwal, R., Agarwal, S., Sharma, S., Gurjar, M. S., Bashyal, B. M., Rao, A. R., Saharan, M.
510   S. Whole Genome Sequencing and Expression Analysis of ToxA in *Bipolaris*
511   *sorokiniana* Provides Discernment of Pathogenicity Causing Spot Blotch of Wheat.
512   2021, PREPRINT (Version 1) available at Research Square.
513   https://doi.org/10.21203/rs.3.rs-677095/v1

514 Virdi, S., Liu, Z., Overlander, M., Zhang, Z., Xu, S. S., Friesen, T., Faris, J. D. New insights
515   into the roles of host gene-necrotrophic effector interactions in governing susceptibility
516   of durum wheat to tan spot and Septoria nodorum blotch. Genes Genomes Genet. 2016,
517   *6*, 4139-4150. https://doi.org/10.1534/g3.116.036525

518 Faris, J. D., Liu, Z. Xu, S. S. Genetics of tan spot resistance in wheat. Theor. Appl. Genet.,
519   2013, *126*, 2197-2217. https://doi.org/10.1007/s00122-013-2157-y

Figure Legends

522 **Figure 1**. Expressed ToxA1 protein showing 16.5kd band size on SDS-PAGE

523 **Figure 2** (a.) The multiple sequence alignments with a phylogenetic tree, Colors both on the
524   phylogenetic tree and on the branch, scale represent genetic distance. ToxA sequences
525   have been colored by four different colors shown in the 'seq' column for each
526   nucleotide. Color changes on the aligned sequences represent nucleotide differences.
527   (b) Heat map based on the number of nucleotide differences between the ToxA
528   haplotypes. Each branch of the phylogenetic tree represents the corresponding ToxA
529   haplotype in the matrix. The close relationships with "darkred" color and far
530   relationships with "white" color. (c) Neighbor-joining (NJ) tree for ToxA haplotypes of
531   based on the nucleotide sequence differences. Colored internal nodes represent the
532   bootstrap confidence level.

533 **Figure 3** Phylogenetic tree using Maximum Likelihood algorithm of different ToxA1 of
534   *Pyrenophora tritici-repentis, Bipolaris sorokiniana,* and varipus haplotypes of
535   *Parastagnospora nodorum*. The percentage of bootstrap support (1000 replicates) is
536   indicated at corresponding nodes. The nomenclature of the haplotypes is based on the
537   Aboukhaddour et al. (2023)

Figure 4. Ramachandran plot and ProsSA z-score analysis of modelled ToxA1 protein from *B. sorokiniana*. Ramachandran distributions of amino acids in the model were calculated using pro check (a), and the z-score plot was plotted using ProsSA-WEB (b).

Figure 5. *ToxA1* protein structure of *B. sorokiniana* isolate MCC 1533 (Pusa2) depicting conserved RGD motif (ARG-GLY-ASP)

Figure 6. Symptom on *Tsn1/tsn1* harboring genotypes after infiltration ToxA1 protein at seedling and adult plant stage.

Table Legends

Table1. Symptom of spot blotch after infiltration of *ToxA1* protein at both seedling and adult stage of wheat genotypes harboring *Tsn1/tsn1* allele.

Supplementary Materials:

Figure S1. Correlation matrix of phenotypic traits Tsn1/tsn1 gene effect on inoculation with ToxA1 isolate**;**

Figure S2. Necrotrophic area developed due to ToxA1–Tsn1 interaction at both seedling and adult stage of wheat genotypes**;**

Figure S3. Symptom of spot blotch after infiltration ToxA1 protein at both seedling and adult stage of wheat genotypes harbouring *Tsn1*;

Figure S4. Symptom of spot blotch after infiltration ToxA1 protein at both seedling and adult stage of wheat genotypes harbouring *tsn1*

Table1: Symptom of spot blotch after infiltration of *ToxA1* protein at both seedling and adult stage of wheat genotypes harboring *Tsn1/tsn1* allele

Genotypes	Length(cm)		Width(cm)		Area(cm2)		Tsn1/tsn1
	Seedling	Adult	Seedling	Adult	Seedling	Adult	
K 68	2.872	4.613	0.404	0.530	0.977	2.326	*Tsn1*
C 306	1.921	4.226	0.329	0.633	0.678	1.737	*Tsn1*
HUW468	2.057	3.472	0.402	0.438	0.718	1.407	*Tsn1*
Yangmai 6	0.860	1.406	0.311	0.381	0.316	0.450	*Tsn1*
K65	1.513	2.582	0.380	0.566	0.388	1.096	*Tsn1*
WAMI160	1.615	3.461	0.345	0.506	0.646	1.521	*Tsn1*
WAMI188	1.693	2.582	0.310	0.500	0.585	1.036	*Tsn1*
WAMI224	1.643	2.857	0.330	0.488	0.476	1.298	*Tsn1*
WAMI226	1.782	2.320	0.405	0.503	0.628	1.090	*Tsn1*
WAMI277	1.464	2.174	0.360	0.485	0.464	1.024	*Tsn1*
WAMI288	1.591	2.054	0.432	0.575	0.622	1.017	*Tsn1*
Max.	2.872	4.613	0.432	0.633	0.977	2.326	
Min.	0.860	1.406	0.310	0.381	0.316	0.450	
HUW234	0.964	1.894	0.335	0.408	0.345	0.561	*tsn1*
PBW343	0.860	1.446	0.349	0.408	0.319	0.423	*tsn1*
LOK1	1.098	1.761	0.307	0.432	0.278	0.493	*tsn1*
Sonalika	1.919	2.923	0.361	0.492	0.584	1.321	*tsn1*
WAMI280	0.972	1.305	0.319	0.327	0.265	0.411	*tsn1*
WAMI289	1.065	1.471	0.319	0.344	0.342	0.529	*tsn1*
WAMI290	1.206	1.467	0.327	0.360	0.365	0.493	*tsn1*
Max.	1.919	2.923	0.361	0.492	0.584	1.321	
Min.	0.860	1.305	0.307	0.327	0.265	0.411	

Fig.1: Expressed ToxA1 protein showing 16.5kd band size on SDS-PAGE

Fig2 (a.) The multiple sequence alignments with a phylogenetic tree, Colors both on the phylogenetic tree and on the branch scale represent genetic distance. *ToxA* sequences have been colored by four different colors shown in the 'seq' column for each nucleotide. Color changes on the aligned sequences represent nucleotide differences. **(b) Heat map based on the number of nucleotide differences between the ToxA haplotypes.** Each branch of the phylogenetic tree represents the corresponding ToxA haplotype in the matrix. The close relationships with "darkred" color and far relationships with "white" color. **(c) Neighbor-joining (NJ) tree for ToxA haplotypes of based on the nucleotide sequence differences.** Colored internal nodes represent the bootstrap confidence level.

Fig. 3: Phylogenetic tree using Maximum Likelihood algorithm of different ToxA1 of *Pyrenophora tritici-repentis, Bipolaris sorokiniana,* and varipus haplotypes of *Parastagnospora nodorum*. The percentage of bootstrap support (1000 replicates) is indicated at corresponding nodes. The nomenclature of the haplotypes is based on the Aboukhaddour et al. (2023)

Fig. 4: Ramachandran plot and ProsSA z-score analysis of modelled ToxA1 protein from *B. sorokiniana*. Ramachandran distributions of amino acids in the model were calculated using pro check (a), and the z-score plot was plotted using ProsSA-WEB (b).

Fig. 5: *ToxA1* protein structure of *B. sorokiniana* isolate MCC 1533 (Pusa2) depicting conserved RGD motif (ARG-GLY-ASP)

Fig. 6 Symptom on *Tsn1/tsn1* harboring genotypes after infiltration ToxA1 protein at seedling and adult plant stage.