A new 'CFS tracking' paradigm reveals uniform suppression depth regardless
of target complexity or salience

David Alais[*1], Jacob Coorey[*1], Randolph Blake[2], Matthew J. Davidson[1]

*Equal contribution.

Affiliations:

1) School of Psychology, The University of Sydney, Sydney, Australia
2) Department of Psychology, Vanderbilt University, Nashville, TN, USA

Abstract

When the eyes view separate and incompatible images, the brain suppresses one image
and promotes the other into visual awareness. Periods of interocular suppression can be
prolonged during continuous flash suppression (CFS) - when one eye views a static 'target'
while the other views a complex dynamic stimulus. Measuring the time needed for a
suppressed image to break CFS (bCFS) has been widely used to investigate unconscious
processing, and the results have generated controversy regarding the scope of visual
processing without awareness. Here, we address this controversy with a new 'CFS tracking'
paradigm (tCFS) in which the suppressed monocular target steadily increases in contrast
until breaking into awareness (as in bCFS) after which it decreases until it again disappears
(reCFS), with this cycle continuing for many reversals. Unlike bCFS, tCFS provides a
measure of suppression depth by quantifying the difference between breakthrough and
suppression thresholds. tCFS confirms that: (i) breakthrough thresholds indeed differ across
target types (e.g., faces vs gratings, as bCFS has shown) – but (ii) suppression depth does
not vary across target types. Once the breakthrough contrast is reached for a given stimulus,
all stimuli require a strikingly uniform reduction in contrast to reach the corresponding
suppression threshold. This uniform suppression depth points to a single mechanism of CFS
suppression, one that likely occurs early in visual processing because suppression depth
was not modulated by target salience or complexity. More fundamentally, it shows that
variations in bCFS thresholds alone are insufficient for inferring whether the barrier to
achieving awareness exerted by interocular suppression is weaker for some categories of
visual stimuli compared to others.

Significance statement

Research on unconscious vision has proliferated recently, often employing the continuous
flash suppression (CFS) method in which rapidly changing shapes viewed by one eye
suppress the other eye's image from awareness. If that suppressed image is progressively
strengthened in contrast it will eventually break suppression and emerge into visible
awareness. Low breakthrough thresholds are claimed to indicate unconscious processing
during suppression. We introduce a method that quantifies breakthrough thresholds and
suppression thresholds, thus providing a lower bound missing from previous CFS research.
Comparing various image types, including those claimed to undergo unconscious
processing, all images show equal suppression when both thresholds are measured. We
thus find no evidence of differential unconscious processing and conclude reliance on
breakthrough thresholds is misleading without considering suppression thresholds and leads
to spurious claims about unconscious processing.

44
Introduction
46

The quest to understand visual processing outside of awareness is tantalising but

notoriously challenging to implement (Breitmeyer, 2015; Hesselmann & Moors, 2015;

Holender, 1986; Logothetis, 1998; Newell & Shanks, 2014; Schmidt, 2015). There are

controversial claims of higher-level semantic processing (Lanfranco et al., 2022; Mudrik et

al., 2014; Stein & Sterzer, 2014; Sterzer et al., 2014), object categorization (Kouider &

Dehaene, 2007; Rees, 2007; Sterzer et al., 2014), and abstract reasoning (Hassin, 2013;

Sklar et al., 2012) occurring outside of awareness. A number of methods exist for

manipulating visual awareness (Kim & Blake, 2005), with two of the most popular in recent

years being binocular rivalry (Alais & Blake, 2014) and continuous flash suppression (CFS:

(Fang & He, 2005; Tsuchiya & Koch, 2005). Both rely on interocular suppression induced by

dissimilar monocular images independently presented to each eye, the result being that only

one eye's image is perceived (i.e., dominates) at any given moment, with the other

suppressed from awareness. For images approximately matched in salience, as often is the

case in binocular rivalry (Alais & Blake, 2014; Wang et al., 2022), monocular suppression

lasts for just a few seconds before switching to suppress the other eye (and so on,

alternating irregularly over time). With CFS, a highly salient stream of dynamic images seen

by one eye suppresses a smaller, weaker target presented to the other for considerably

longer periods of suppression that can last tens of seconds. The potency of CFS has great

appeal when it comes to assessing residual effectiveness of different categories of visual

stimuli blocked from awareness by CFS.

A commonly used variant known as "breaking CFS" (bCFS) was introduced by (Jiang

et al., 2007) in which the suppressed target slowly ramps up from low contrast until it

becomes sufficiently strong to break suppression and achieve visibility. Time to

breakthrough has become a popular measure, and differences in breakthrough times among

various image types have been used to support claims for unconscious processing of certain

visual images. According to this line of reasoning, image types that consistently emerge from

suppression more rapidly than do other image types are construed to be ones that receive

prioritized processing outside of awareness. To give three examples of this purported effect,

faces portraying emotional expressions break suppression faster than do neutral faces

(Alais, 2012; Jiang et al., 2007; Tsuchiya et al., 2009; Yang et al., 2007), complex scenes

containing incongruent objects break suppression faster than do congruent scenes (Mudrik

et al., 2011), and native words break suppression faster than do unfamiliar foreign words

(Jiang et al., 2007). Sceptics argue, however, that differences in breakthrough times can be

attributed to low-level factors such as spatial frequency, orientation and contrast that vary

between images (Gayet et al., 2014; Moors, 2019; Moors et al., 2016, 2017; Moors &

Hesselmann, 2018; Stuit et al., 2023), and more fundamentally, that breakthrough times

alone are insufficient to measure differential unconscious processing (Stein, 2019).

84   Conclusions based on a comparison of bCFS breakthrough times between different

image categories suffer from a problem of unidirectionality and a false assumption of image

equivalence[1]. Images producing faster breakthrough times (equivalently, lower breakthrough

contrasts) are interpreted as undergoing residual processing outside of awareness, adding

to their salience and weakening their interocular suppression (Mudrik et al., 2011; Sterzer et

al., 2011; G. Zhou et al., 2010; W. Zhou et al., 2010)). An implicit assumption here is that as

all images were initially invisible, the depth of interocular suppression was thus weaker for

images that break suppression more quickly (i.e., produce faster reaction times, as

sometimes stated). Yet, the depth of interocular suppression is rarely measured in CFS

paradigms (see Tsuchiya et al,. 2006 for an exception), and to our knowledge, has not been

explicitly compared between image categories.

95   One method for measuring interocular suppression is to compare the threshold for

change-detection in a target when it is monocularly suppressed and when it is dominant, an

established strategy in binocular rivalry research (Alais, 2012; Alais et al., 2010; Alais &

Melcher, 2007; Nguyen et al., 2003). Probe studies using contrast as the dependent variable

for thresholds measured during dominance and during suppression can advantageously

standardise suppression depth in units of contrast within the same stimulus (e.g., Alais &

Melcher, 2007; Ling et al., 2010). Ideally, the change should be a temporally smoothed

contrast increment to the rival image being measured (Alais, 2012), a tactic that preludes

abrupt onset transients and, moreover, provides a natural complement to the linear contrast

ramps that are standard in bCFS research. In this study, we measure bCFS thresholds as

the analogue of change-detection during suppression, and as their complement, record

thresholds for returns to suppression (reCFS). We do this by including a bidirectional

measure in which the target contrast steadily decreases (ramps down) over time, eventually

culminating in the target's transition from dominance to suppression. By comparing the

thresholds for a target to transition into (reCFS) and out of suppression (bCFS), we

recognise that the criterion for change is much higher, and as a result expect larger

suppression depths (on average) than those recorded in rivalry research. More importantly,

however, by averaging the difference between bCFS and reCFS thresholds, a key analogue

---

[1] An exception here is when the same image is compared under different contexts, such as after associative learning of value (Lunghi & Pooresmaeili, 2023) or fear conditioning (Gayet et al., 2016).

of suppression depth that has been missing from the CFS literature can be provided and

compared between image categories. Figure 1 outlines hypothetical results that can be

obtained when recording reCFS thresholds as a complement to bCFS thresholds in order to

measure suppression depth.

Here, we introduce a novel method termed 'tracking CFS' (tCFS) which combines

alternating down-ramped and up-ramped targets, to provide a measure of suppression depth

that allows a rigorous test of claims that certain image types undergo less suppression than

others (see Figure 2). The participant views a visible target as it declines in contrast until it

becomes suppressed. When suppression is reported the contrast change reverses direction

and the participant indicates when the target emerges out of dominance and is seen once

again (triggering a contrast decrease again, and so on over many reversals). By continually

tracking thresholds over time for breakthroughs (bCFS) and succumbing (reCFS) to

suppression, tCFS is very efficient, makes no assumptions of image equivalence, and

provides inherently bidirectional measures of CFS.

To preview the results, across three experiments we find no evidence using the tCFS

paradigm for differences in suppression depth among several image categories (i.e., faces,

objects, several types of gratings, and visual noise). We do replicate the common

observation that breakthrough thresholds differ across image categories, but now we find

that reCFS thresholds covaried with bCFS thresholds across all image categories, meaning

that suppression depth was constant for faces, objects, gratings and visual noise. In other

words, we find no evidence to support differential unconscious processing among these

particular, diverse categories of suppressed images.

Figure 1. The critical measure of suppression depth remains untested in CFS.
a,b) Example stimulus displays for the discrete trials, during which (a) a low-contrast target steadily increases in contrast until visibility is reported (bCFS), and (b) discrete reCFS trials, with a high-contrast target decreasing until target invisibility is reported. **c)** A typical bCFS result, in which a target image (here a face or non face) is initially weak and steadily increases until it breaks suppression. Faster breakthrough times (lower contrast) for face stimuli are often interpreted as evidence that faces undergo expedient processing that counteracts their susceptibility to suppression relative to other visual stimuli. Without also measuring suppression thresholds for each stimulus, this conclusion is premature. **d-f)** To define the magnitude of suppression during CFS, it is necessary to measure the contrast thresholds at which stimuli enter and exit awareness, with the difference indicating suppression depth. Red bars display hypothetical results measuring the contrast at which an initially visible stimulus with decreasing contrast becomes suppressed by the mask (reCFS). The results of panel **c** are reproduced in d-f in light blue. **d)** If the reCFS thresholds for face and non-face images are the same, this would support reduced suppression depth for faces (as ☐ face < ☐ non-face). **e)** Alternatively, the reCFS thresholds for faces and non-faces might differ, with the face remaining visible at a lower contrast than non-face images (lower reCFS threshold), indicating more suppression for faces than non-faces (as ☐ face > ☐ non-face). **f)** Finally, the reCFS thresholds might differ between faces and non-faces but by an amount equivalent to their bCFS differences, indicating the same suppression depth for both image types (as ☐ face = ☐ non-face). Such a result would argue against enhanced unconscious processing of face stimuli.

Results

Experiment 1

Many previous bCFS studies have shown that *increasing* contrast eventually causes
a target image to overcome interocular suppression (i.e., emerge into awareness). To our
knowledge, however, none has investigated the contrast at which an initially visible image
succumbs to suppression as its contrast is *decreased.* Experiment 1 uses the tCFS method
to continuously track changes in target visibility as it rises and falls in contrast. As shown in
Figure 2b, this provides a series of bCFS thresholds (upper turning points) as well as
thresholds for the target's re-suppression (lower turning points: reCFS thresholds). We
compare these thresholds to those obtained with a discrete procedure in which contrast
either increased steadily from a low starting point until breakthrough (standard bCFS
measure) or decreased steadily from a high starting point until suppression was achieved.
On each trial the target image was either a face or a familiar object, with the images all
matched in size, RMS contrast and mean luminance.

A repeated-measures ANOVA revealed a significant main effect of threshold, such
that bCFS thresholds were significantly higher than reCFS thresholds ($F(1,19) = 50.39$, $p <$
$.001$, $\eta_p^2 = .73$). There was also a significant interaction between threshold and condition
($F(1,19) = 41.19$, $p < .001$, $\eta_p^2 = .68$), indicating that the difference between bCFS and
reCFS thresholds was influenced by whether they were recorded using the discrete trial
procedure or the continuous trial procedure. Subsequent post-hoc tests revealed that bCFS
and reCFS thresholds differed within each type of procedure (Discrete: $t(19) = 2.89$, $p =$
$.009$, $d = 0.65$ Continuous: $t(19) = 12.12$, $p < .001$, $d = 2.7$). Overall, the suppression depth
(i.e., the difference between bCFS and reCFS thresholds) was larger with the continuous
procedure ($M = -15.40$ dB, $SD = 5.68$) compared to the discrete procedure ($M = -5.74$ dB,
$SD = 8.89$), $t(19) = 6.42$, $p < .001$, $d = 1.3$), a finding we return to in the Discussion. Figure
2c displays a summary of these results.
After confirming a difference between bCFS and reCFS thresholds, we next
compared suppression depth measured using different image types. We repeated the
analysis with the additional exploratory factor of image type (face vs object) in a 2 x 2 x 2
repeated-measures design (threshold, procedure, image type). We again found significant
main effects of threshold ($F(1,19) = 50.39$, $p < .001$, $\eta_p^2 = .73$), and a threshold x condition
interaction ($F(1,19) = 41.19$, $p < .001$, $\eta_p^2 = .68$), but non-significance for the effect of image
type ($p = .26$).
We quantified the evidence for this null-effect on suppression depth with a

subsequent Bayesian model comparison. A Bayesian repeated-measures ANOVA (2 x 2;

procedure x image type on suppression depth) found that the best model to explain

suppression depth included the main effect of procedure (BF_{10} = 3231.74), and weak

evidence/data insensitivity for image type (BF_{10} = 0.37). This indicates that the data was

insensitive as to whether image-type was better at predicting suppression depth than the null

model. In other words, there is a large disparity in contrast between targets breaking and re-

entering CFS, but the magnitude of this disparity is most likely the same for objects and for

faces. This potentially important finding is the focus of our next Experiment.

205
206

Figure 2. Example tCFS trial and comparison to discrete conditions**. a-b)** Example tCFS trial
from one participant, showing the change in contrast over time. Red markers indicate the
level at which a target with decreasing contrast became suppressed (reCFS) and blue
markers indicate traditional bCFS responses (breakthrough of contrast-increasing target).
The same trial is shown with target contrast in decibel scale in b. Because the human visual
system has a logarithmic contrast response, it is appropriate to increase/decrease target
contrast logarithmically as in b, to create a contrast change that is perceptually linear. **c)**

Interaction between thresholds and condition type. Individual blue and red dots display
participant means for bCFS and reCFS respectively. Grey lines link thresholds per
participant, per condition. Blue and red diamonds display the mean across participants, and
error bars plot ± 1SEM corrected for within participant comparisons (Cousineau, 2005).
218

Experiment 2

Experiment 1 disclosed that bCFS thresholds were not equivalent to reCFS
thresholds, and demonstrated that the suppression depth of images could be quantified in
an image-specific manner. Furthermore, Experiment 1 showed that suppression depth could
be measured rapidly using the tCFS procedure and did not differ between faces and objects.
In Experiment 2, we tested whether the constant suppression depth obtained in Experiment
1 for two image types would replicate across a larger variety of image categories.

We measured suppression depth for faces, objects, linear gratings, phase
scrambled images, and radial/concentric patterns using the tCFS method. All have been
used to investigate bCFS thresholds before (Stein, 2019). We reasoned that faces and
objects are familiar, meaningful stimuli processed widely throughout the visual hierarchy.
Gratings and phase-scrambled images, on the other hand, comprise simpler, meaningless
stimuli devoid of perceptual relevance. Phase-scrambled images were created from the
object stimuli by randomising their phase spectra, rendering the images unrecognizable
(e.g., Piotrowski & Campbell, 1982; Yu & Blake, 1992). These images thus served as
controls for the regular stimuli, since both ordinary and phase-scrambled images have
approximately the same Fourier energy spectra. Finally, polar patterns (radial and
concentric gratings) were included because they have robust structural coherence that is
registered by extrastriate visual areas in the ventral stream (Wilkinson et al., 2000), but are
lacking in semantic meaning (Hong, 2015).

A 2 x 5 repeated-measures ANOVA revealed a significant main effect of threshold
(bCFS vs reCFS; $F(1,17) = 133.79$, $p < .001$, $\eta_p^2 = .89$), and image type ($F(4,68) = 13.45$, $p <$
$.001$, $\eta_p^2 = .44$). Critically however, there was no significant interaction between thresholds
and image type ($p = .98$), indicating that the relationship between bCFS and reCFS
thresholds was invariant across image categories. A Bayesian repeated-measures ANOVA
(1 x 5, effect of image categories on suppression depth), confirmed strong evidence in
favour of the null hypothesis ($BF_{01} = 20.30$).

This result is plotted in Figure 3, which clearly shows differences in bCFS thresholds
(blue symbols in Figure 3a) over image type (1 x 5 repeated-measures ANOVA; $F(4,68) =$
16.29, $p < .001$, $\eta_p^2 = .49$), as has been reported in many studies. Scrambled noise images,
for example, yield breakthrough thresholds 5.3 dB higher than do face images, and linear
gratings breakthrough thresholds are 3.0 dB higher than polar gratings. Critically, Figure 3

also shows that reCFS thresholds exhibit the same pattern of differences over image type

(red symbols in Figure 3a): and as confirmed by our Bayesian model comparison, the depth

of suppression is constant across all image categories.

Replicating the result of Experiment 1, approximately 15 dB of suppression depth

was observed in Experiment 2 (Fig. 3b), which was practically identical across image types

(Faces, $M = 14.35$ ($SD = 5.64$); Objects, $M = 14.61$ ($SD = 6.07$); Gratings, $M = 14.88$ ($SD =$

5.55); phase-scrambled images, $M = 14.70$ ($SD = 5.39$); polar patterns, M = 14.69 (SD =

6.50). A repeated-measures ANOVA confirmed that suppression depth did not differ across

image categories ($F(4,68) = 0.1$, $p = .98$).

Figure 3. Suppression depth is uniform across image categories. a) bCFS and reCFS
thresholds by image type. Blue and red diamonds display the mean across participants, and
error bars plot ±1 SEM corrected for within participant comparisons (Cousineau, 2005). Grey
lines link thresholds per participant, per image type. For visualization, we have linked the
bCFS and reCFS thresholds with broken lines, to better indicate that both do vary according
to image category. **b)** The relative difference in contrast between bCFS and reCFS
thresholds is the same across image categories.

Experiment 3

  Experiments 1 and 2 introduced the tCFS method and, using that new method,
demonstrated that bCFS thresholds vary depending on target type, as many previous bCFS
studies have shown. Importantly, reCFS thresholds varied in parallel with bCFS thresholds
which, when expressed in terms of suppression depth, reveals that depth of suppression is

strikingly uniform across target image categories in CFS. This uniformity of suppression depth

could indicate that neural events mediating CFS suppression transpire within a common visual

mechanism, one that is not selective for image type, complexity or semantic meaning. This view

is compatible with popular models of binocular rivalry built around the concept of reciprocal

inhibition and neural adaptation (Alais et al., 2010; Kang & Blake, 2010; McDougall, 1901), as

well as with more recent Bayesian-inspired inference-based models in which perceptual

alternations in dominance are triggered by accumulating residual error signal associated with

competing stimulus interpretations (Hohwy et al., 2008). As pointed out elsewhere (Blake, 2022),

steadily increasing error signal effectively plays the same role as does steadily decreasing

inhibition strength caused by neural adaptation in reciprocal inhibition models.

In the context of the tCFS method, the steady increases and decreases in the target's

actual strength (i.e., its contrast) should, respectively, boost its emergence from suppression

(bCFS) and facilitate its reversion to suppression (reCFS) as it competes against the mask.

Whether construed as a consequence of neural adaptation or error signal, we surmise that these

cycling state transitions defining suppression depth should be sensitive to the rate of contrast

change of the monocular target. Specifically, the slower the contrast change, the greater the

amount of accrued adaptation, which will contract the range between breakthrough and

suppression thresholds according to an adapting reciprocal inhibition model. For fast contrast

change, there will be less accrual of adaptation meaning that the range between breakthrough

and suppression thresholds will exhibit less contraction. Expressed in operational terms, the

depth of suppression should be positively related to the rate of target change. Experiment 3

tested this supposition using three rates of contrast change.

A 3 x 2 x 4 repeated measures ANOVA compared three rates of contrast change

(slow, medium, fast), on both thresholds (bCFS, reCFS) across four image categories (face,

object, grating, phase scrambled) using the tCFS paradigm. There was a significant main

effect of threshold ($F(1,16) = 116.56$, $p < .001$, $\eta_p^2 = .88$) again indicating that bCFS and

reCFS contrasts differ. There was also a significant main effect of image type ($F(3,48) =$

9.40, $p < .001$, $\eta_p^2 = .37$), again with no interaction with threshold ($p = .19$). This result

indicates that bCFS and reCFS thresholds vary in tandem regardless of image type. This

null-effect of image-type was again confirmed with a Bayesian model comparison (3 speed x

4 image categories on suppression depth), demonstrating moderate support for the null

effect of image category ($BF_{01} = 4.06$). Critically, there was a significant interaction between

rate of contrast change and thresholds ($F(2,32) = 128.60$, $p < .001$, $\eta_p^2 = .89$), as expected,

indicating that the difference between bCFS and reCFS thresholds (i.e., suppression depth)

depended on the target's rate of contrast change. **Figure 4** displays a summary of these

results (averaged across image types), showing that as the rate of contrast change

increases, so does suppression depth (Slow $M = 9.64$ dB, $SD = 4.37$, Medium $M = 14.6$ dB,

SD = 5.43; Fast *M* = 18.97 dB, *SD* = 6.93).

We performed additional analyses to rule out an alternative explanation for the

increased suppression durations demonstrated in Experiment 3. We reasoned that if

alternations between bCFS and reCFS were happening with a regular periodicity, such that

responses were made every 1 second (for example), then these consistent responses could

result in smaller suppression thresholds when the rate of contrast change was slow, as we

have observed. Similarly, larger suppression thresholds would be measured if the same 1

second interval had elapsed while the rate of contrast change was fast. Inspection of the raw

tCFS time-series qualitatively indicated that perceptual durations were varying with the rate

of target contrast change (Figure 4c). To test this possibility, we compared the group

average perceptual durations across all experiments, to test whether the average duration of

percepts was the same despite different rates of contrast change. Figure 4 - figure

supplement 1 displays the results of this analysis. The histograms of perceptual durations for

Experiments 1 (all tCFS median M = 3.83, SD = 1.66) and Experiment 2 (M = 3.48, SD =

1.36) show similar means and distributions, with no significant difference between them

(t(36) = 0.72, p = .48). This is unsurprising given their similar design and matched rate of

contrast change. In Experiment 3, however, the distribution of percept durations is shown to

vary by rate of contrast change. With shorter median percept durations for fast rates of

contrast change (M = 3.08, SD = 1.07), and slower percept durations for slow rates of

contrast change (M = 4.61, SD = 2.05) compared to medium (M = 3.45, SD = 1.29). A

repeated measures ANOVA confirmed that median percept durations varied by rate of

contrast change (F(2,32) = 30.89, p < .001, ηp2 = .66).

This **analysis** thus confirmed that these differences in suppression depth were not

driven by fixed rates of perceptual alternation across the three levels of rate of contrast

change (Figure 4c and Figure 4 - figure supplement 1)

Figure 4. Suppression depth is greater with less time for adaptation. a) bCFS (blue) and reCFS (red) thresholds collected during tCFS with three rates of contrast change (RCC). Figure elements are the same as in Figure 3a. **b)** Suppression depth increases when the rate of contrast change increases during tCFS. All error bars correspond to ±1 SEM corrected for within participant comparisons (Cousineau, 2005). **c)** Example trials at each rate of contrast change from a single participant. Red markers indicate reCFS responses, blue markers show bCFS.

Perceptual switches during tCFS are described by a damped harmonic oscillator

The results of Experiment 3 demonstrated that when the opportunity for target adaptation is increased, as when the target's rate of contrast change was slow, that suppression depth is reduced and a smaller contrast decrease is needed for a visible target

to reenter CFS (see Figure 4a). One possible account for this relates to the balance of

excitation/inhibition in neural systems, which have been particularly fruitful models of

interocular competition (Alais et al., 2010; Li et al., 2017). In these models, adaptation over

time is a critical parameter governing changes in visual consciousness (Alais et al., 2010),

which motivated us to explore in our final analysis whether suppression depths also

fluctuated over time. Accordingly, our final analysis sought to model the temporal nature of

perceptual switches during tCFS, to understand whether a balance of excitation and

inhibition may be contributing to the sequential contrast thresholds that govern target

visibility.

For this analysis, a key dependent variable is the contrast difference between

sequential thresholds (i.e., from bCFS to reCFS, or reCFS to bCFS). As the number of

thresholds was the same in each trial, we averaged each threshold over observers to obtain

a sequence of mean thresholds that preserved the order across the trial (e.g. average reCFS

threshold at response 1, average bCFS threshold at response 2, average reCFS threshold at

response 3 etc). Importantly, pooling across the threshold order rather than each

participant's time series avoids smearing the data due to observers differing in their

perceptual durations (**cf. Figure 4 - figure supplement 1**). **Figure 5a** displays the mean

result separately for each target rate of change condition. To investigate the balance

between bCFS and reCFS thresholds, we calculated the absolute change in contrast

between sequential thresholds, as plotted in **Figure 5b**. These sequential estimates of

suppression depth show marked fluctuations early in each trial, followed by a stabilisation to

the grand average suppression depth for each rate of contrast change (Slow ~10 dB,

Medium ~ 15 dB, Fast ~ 19 dB).

We next tested whether these sequential changes in suppression depth could be

described by models of excitation and inhibition. For each rate of change condition, we

compared the goodness of fit of three models: a simple harmonic oscillator, a damped

harmonic oscillator, and a cubic polynomial model. We assessed the relative goodness of fit

using the change in Bayesian Information Criterion (BIC) between models (see **Methods**).

As can be appreciated in **Figure 5c,** sequential changes in suppression depth were

well described by the damped harmonic oscillator model of excitation and inhibition, which

well captured the modulations early in the trial. For each rate of target contrast change, the

damped harmonic oscillator model was the superior fit to the data (Slow contrast change, R^2

= 0.68, BIC = -1847.2; Medium contrast change, R^2 = 0.63, BIC= -2125.29; Fast contrast

change, R^2 = 0.36, BIC= -1041.29). In each case, the change in Bayesian Information

Criterion between the damped harmonic oscillator and the next best fitting cubic polynomial

model was large (Slow contrast change, ΔBIC = -1072.88; Medium contrast change, ΔBIC =

-781.5; Fast contrast change, ΔBIC = -97.06). Table 1 displays the assessments of each

model fit for each rate of change condition.

The damped harmonic oscillator has been proposed to capture the response of

neural populations governed by excitation/inhibition balance after external perturbation, with

a given rate of decay. We return to the interpretation of these models in the Discussion.

Figure 5. Suppression depth over time is well described by a damped harmonic

oscillator. a) Top row: Mean bCFS (blue) and reCFS (red) thresholds averaged by

threshold order over the trial. Each column displays the average for slow, medium and fast

rate of contrast change (RCC), respectively. **b) Middle row.** Absolute change in target

contrast between successive thresholds (i.e., suppression depth). Red bars indicate the

decrease in target contrast needed for a visible target to reenter CFS. Blue bars indicate the

increase in contrast for an invisible target to break CFS. **c) Bottom row:** Yellow lines and

shading plot the average change in suppression depth between successive thresholds. The

data in each RCC condition is best fit by a damped harmonic oscillator (DHO) model shown

in black. The change in Bayesian Information Criterion relative to the next best (cubic

polynomial) model is displayed in each panel. All error bars and shading denote ±1 SEM

corrected for within participant comparisons (Cousineau, 2005).

		Slow RCC	Medium RCC	Fast RCC
Model	**Measure**			
(cubic) 1	R^2	.07	.19	.29
	BIC	-774.31	1343.79	-944.23
(harmonic) 2	R^2	.005	.006	0
	BIC	-706.85	1138.595	-600.34
(dHO) 3	R^2	.684	.632	.361
	BIC	-1847.19	-2125.29	-1041.29
Comparison				
2-1	ΔBIC	67.46	205.195	343.89
3-1	ΔBIC	-1072.88	-781.5	-97.06
3-2	ΔBIC	-1140.34	-986.695	-440.95

412   **Table 1.** Summary of results from model fits and comparisons.

Discussion

This study introduces a new CFS methodology that indexes suppression depth based on a participant's continuous report of the visibility status of a monocularly viewed target that steadily waxes and wanes in contrast, while the other eye views a CFS mask that forces the target into and out of suppression. The key to the technique, dubbed tracking CFS (tCFS), is that the participant's responses control when the reversals in the direction of contrast change occur: for example, a visible target initially decreases from high contrast until the participant reports the target's disappearance from awareness, at which time the target begins increasing in contrast until it again achieves awareness, and so on in an ongoing cycle. By tracking target visibility in this way, the method produces breakthrough thresholds (as in bCFS), but also yields measures of suppression thresholds(reCFS). By measuring the relative difference between these two thresholds, one can derive a rational index of suppression depth that, in turn, allows critical evaluation of whether suppression depth varies for different image categories (e.g., faces vs gratings). Experiment 1 introduced and validated the tCFS method for measuring suppression depth. Experiment 2 applied tCFS to images from different categories and found that suppression depth was invariant across a set of different types of images, regardless of their complexity, familiarity or salience. Experiment 3 manipulated suppression depth by varying the rate of target contrast change, providing presumptive evidence consistent with particular interpretation of the neural bases of the tCFS procedure. The following sections elaborate on each of these points.

The tCFS method offers two important advances for studying visual processing outside of awareness. First, it is fast and efficient. The data in Figure 2b, for example, show that 90 seconds is sufficient to collect a robust set of data comprising 16 thresholds, eight each for bCFS and reCFS. The second advantage is that by providing easy quantification of breakthrough and suppression thresholds, tCFS allows the calculation of a rational index of strength of suppression for evaluating whether certain categories of images (e.g., faces) receive prioritized processing, a controversial issue within the CFS literature. We will have more to say about claims of preserved visual processing despite suppression, and claims of unconscious processing more generally in a moment (see ***No effect of image category on suppression depth***).

Thresholds for breaking and entering suppression quantify suppression depth

In Experiment 1, we used the tracking CFS method to measure bCFS and reCFS contrast thresholds, and compared them to those obtained using discrete unidirectional contrast changes – a series of measures with increasing target contrast to obtain bCFS

thresholds as conventionally done, and also a series with decreasing target contrast to

measure suppression (reCFS) thresholds. For both conditions, significant differences

between bCFS and reCFS thresholds were observed. This difference indicates that

suppression is not a passive process where a given contrast threshold determines a narrow

awareness/suppression border. Instead, there is a contrast range (i.e., suppression depth)

between awareness and suppression. This is consistent with models of interocular

suppression involving mutually inhibitory left- and right-eye processes. The fact that

binocular rivalry produces a significant degree of suppression depth is well known in that

literature (Blake & Camisa, 1979; Nguyen et al., 2003) but is rarely considered in CFS

studies[2] where the focus is typically on the breakthrough threshold for awareness.

Intuitively, the average contrast threshold for reCFS should be lower than for bCFS,

yet in the discrete conditions there were some participants who showed the inverse effect (*n*

= 4; **Figure 2c**). One possibility is that saccades, blinks, or other oculomotor reflexes that

differentially impact transitions during interocular rivalry (van Dam & van Ee, 2006a, 2006b)

might have influenced the discrete bCFS and reCFS trials at separate times. Another

possibility is that by using a blocked design in the discrete trials, a participant's motivation or

attention to task might have changed in the time elapsed between recording thresholds in

bCFS and reCFS blocks. These possibilities highlight inherent advantages and controls of

the tCFS procedure, for which no participant showed an inverse effect: bCFS and reCFS

thresholds are recorded in alternation, thus each threshold judgment is relative to the

previous judgment on the same image, recorded a matter of seconds (rather than minutes)

prior. During tCFS, bCFS and reCFS tasks alternate, which helps maintain a level of

engagement as the task is never repeated. Alternating between bCFS/reCFS tasks also

means that any adaptation occurring over the trial will occur in close proximity to each

threshold, as will any waning of attention. The benefit being that bCFS and reCFS thresholds

are measured under closer conditions in continuous trials, compared to discrete ones. Long

periods of target invisibility that can occur in typical bCFS studies when the target is raised

from near-zero contrast are also avoided, as target contrast hovers around the

visibility/invisibility thresholds. As Figure 2c shows, tCFS produces an increased size of

suppression depth compared to discrete trials. Moreover, it can be administered quickly, and

provides threshold measures with less variance.

As an aside, the existence of distinctly different, complementary transition thresholds

for bCFS and reCFS is reminiscent of the behavior termed hysteresis: a property of

dynamical systems wherein output values, rather than being solely governed by

---

[2] An exception to this rule is the study by Tsuchiya et al. (2006) who used a briefly presented test probe to compare depth of suppression associated with CFS and with BR; suppression depth was considerably larger for the former than for the latter.

corresponding input values, also exhibit lags or delays based on the valence of continuous

changes in the input values. Examples of hysteresis in visual perception include transitions

between binocular fusion and binocular rivalry (Anderson, 1992; Buckthought et al., 2008;

Julesz & Tyler, 1976), perception of motion direction in random-dot cinematograms (Williams

et al., 1986), and repetition priming in perception of bistable configurations (Pastukhov et al.,

2015). To paraphrase Maglio and Polman (Maglio & Polman, 2016), hysteresis can be

construed as a form of memory whereby prior states influence the persistence of current

states into the future.

No effect of image category on suppression depth

Having demonstrated a difference between bCFS and reCFS thresholds using the

tCFS procedure, Experiment 2 compared suppression depth across five different image

categories. As noted in the Introduction, a number of previous studies have interpreted

differences in bCFS thresholds (or suppression durations) for different categories of images

without taking into account thresholds for *inducing* suppression, which we believe is an

important piece of information necessary for calculating suppression depth. In this paper, we

applied the tCFS method to assess whether bCFS thresholds differed among several image

categories and, as well, whether category-specific differences in suppression depth would be

obtained when reCFS thresholds were also included in the equation. As an important

prerequisite for asking this question, our bCFS thresholds indeed did replicate the often

reported finding that certain image types break out of suppression into awareness at lower

contrasts than do others (see Figure 3a). For example, bCFS thresholds for faces were

lower than bCFS thresholds for phase-scrambled images by 5.3 dB. Critically, however,

while bCFS thresholds varied with image type, the reCFS thresholds for those and, indeed,

for all images was approximately 15 dB lower than bCFS, *regardless of image type*. In other

words, all five of the image categories we tested, regardless of complexity or familiarity,

produced a constant depth of suppression averaging about 15 dB (see Fig. 3b), even though

their bCFS thresholds varied.

As a reminder, we explicitly tested image types that in other studies have shown

differential susceptibility to CFS attributed to some form of expedited unconscious

processing. Nevertheless, one could argue that our failure to obtain evidence for category

specific suppression depth is based on the limited range of image categories sampled in this

study. We agree it would be informative to broaden the range of image types tested using

tCFS to include images varying in familiarity, congruence and affect. We can also foresee

value in deploying tCFS to compare bCFS and reCFS thresholds for visual targets

comprising physically meaningless 'tokens' whose global configurations can synthesise

recognizable perceptual impressions. To give a few examples, dynamic configurations of

small dots varying in location over time can create the compelling impression of rotational

motion of a rigid, 3D object (structure from motion) or of a human engaged in given activity

(biological motion) (Grossmann & Dobbins, 2006; Watson et al., 2004). These kinds of visual

stimuli are associated with neural processing in higher-tier visual areas of the human brain,

including the superior occipital lateral region (e.g., (Vanduffel et al., 2002)) and the posterior

portion of the superior temporal sulcus (e.g., (Grossman et al., 2000)). These kinds of

perceptually meaningful impressions of objects from rudimentary stimulus tokens are

capable of engaging binocular rivalry. Such stimuli would be particularly useful in assessing

high-level processing in CFS because they can be easily manipulated using phase-

scrambling to remove the global percept without altering low-level stimulus properties. In a

similar vein, small geometric shapes can be configured so as to resemble human or human-

like faces, such as those used by (Zhou et al., 2021)[3]. These kinds of faux faces could be

used in concert with tCFS to compare suppression depth with that associated with actual

faces.

Next we turn to another question raised about our conclusion concerning invariant

depth of suppression. If a certain image type had overall lower bCFS and reCFS contrast

thresholds relative to another image type (despite equivalent suppression depth), would that

imply the former image enjoyed "preferential processing" relative to the latter? And, what

would determine the differences in bCFS and reCFS thresholds? Figure 3 shows that polar

patterns tend to emerge from suppression at slightly lower contrasts than do gratings and

that polar patterns, once dominant, tend to maintain dominance to lower contrasts than do

gratings and this happens even though the rate of contrast change is identical for both types

of stimuli. But while rate of contrast change is identical, the neural responses to those

contrast changes may not be the same: neural responses to changing contrast will depend

on the neural contrast response functions (CRFs) of the cells responding to each of those

two types of stimuli, where the CRF defines the relationship between neural response and

stimulus contrast. CRFs rise monotonically with contrast and typically exhibit a steeply rising

initial response as stimulus contrast rises from low to moderate values, followed by a

reduced growth rate for higher contrasts. CRFs can vary in how steeply they rise and at what

contrast they achieve half-max response. CRFs for neurons in mid-level vision areas such

as V4 and FFA (which respond well to polar stimuli and faces, respectively) are generally

steeper and shifted towards lower contrasts than CRFs for neurons in primary visual cortex

---

[3] Zhou et al. (2021) derived dominance and suppression durations with fixed-contrast images. In their study, genuine face images and faux faces remained suppressed for equivalent durations whereas genuine faces remained dominant significantly longer than did faux faces. The technique used by those investigators - interocular flash suppression (Wolfe, 1994) - is quite different from CFS in that it involves abrupt, asynchronous presentation of dissimilar stimuli to the two eyes. It would be informative to repeat their experiment using the tCFS procedure.

(which respond well to gratings). Therefore, the effective strength of the contrast changes in

our tCFS procedure will depend on the shape and position of the underlying CRF, an idea

we develop in more detail in **Appendix 1**, comparing the case of V1 and V4 CRFs.

Interestingly, the comparison of V1 and V4 CRFs shows two interesting points: (i) that V4

CRFs should produce much lower bCFS and reCFS thresholds than V1 CRFs, and (ii) that

V4 CRFs should produce much more suppression than V1 CRFs. Our data do not support

either prediction: bCFS and reCFS thresholds for the polar shape are not 'much lower' than

those for gratings (Fig. 3) and neither is there 'much more' suppression depth for the polar

form. There is no room in these results to support the claim that certain images are special

and receive "preferential processing" or processing outside of awareness. Instead, the

similar data patterns for all image types is most parsimoniously explained by a single

mechanism processing all images (see Appendix 1), although there are many other kinds of

images still to be tested in tCFS and exceptions may yet be found. As a first step in exploring

this idea, one could use standard psychophysical techniques (e.g., (Ling & Carrasco, 2006))

to derive CRFs for different categories of patterns and then measure suppression depth

associated with those patterns using tCFS.

In a similar vein, it's natural to wonder whether this non-selectivity of CFS depth of

suppression applies to binocular rivalry suppression, too. Blake and Fox (Blake & Fox, 1974)

concluded that rivalry suppression is non-selective based on a task where observers were

unable to notice large changes in the spatial frequency or orientation of a suppressed

grating. On the other hand, Alais and Melcher (Alais & Melcher, 2007) found that the

detectability of a brief, monocular probe presented to an eye during rivalry varied depending

on the 'complexity' (e.g., grating vs face) of the stimulus being probed, with suppression

being greater for complex images. Tsuchiya et al. (Tsuchiya et al., 2006) found that

detection of a brief test probe was much more difficult to detect when presented to an eye

during suppression phases of CFS compared to binocular rivalry. In a similar vein, durations

of suppression phases associated with CFS are considerably longer than those associated

with rivalry (~15x longer, in the study by Blake et al., 2019). A clear next step will be to apply

a variant of the tCFS paradigm to binocular rivalry to assess the uniformity of rivalry

suppression depth based on stimulus complexity.

Turning to a methodological issue, an important caveat for interpreting suppression

depth during tCFS is that bCFS thresholds will be slightly overestimated due to the brief

reaction time delay between the perceived transition and execution of the motor response

signalling that event (a problem for most bCFS studies). By the same token, reCFS

thresholds will be slightly underestimated because of this response delay. These disparities

between transition times and the responses to them lead to a slight inflation of suppression

depth. For example, if we assume an average reaction time of 500 ms for appearance and

disappearance events, then suppression depth will be inflated by ~4.2 dB at the rate of

contrast change used in Experiments 1 and 2 (.07 dB per frame at 60 fps). This cannot

account for suppression depth in its entirety, which was many times larger at approximately

14 dB across image categories. We have no reason to suspect that reaction-times would

differ when reporting on the appearance or disappearance of different image categories

under CFS, and indeed found no significant evidence for an interaction between thresholds

and image categories in our analyses. This leaves a constant suppression depth of

approximately 14 dB to be explained, and the value of comparing between image categories

remains. In Experiment 3, the rate of contrast change varied which led to corresponding

changes in suppression depth, which we note could also not be attributed to a reaction-time

delay (Figure 4 - figure supplement 1). Using the same assumptions of a 500 ms response

time delay, this would predict a suppression depth of 2.1 dB, 4.2 dB and 6.3 dB for the slow,

medium and fast ramp speeds respectively. However, this difference cannot account for the

size of the effects (*Slow* 9.64 dB, *Medium* 14.6 dB, *Fast* 18.97 dB). The difference in

suppression depth based on reaction-time delays (± 2.1 dB) also does not match with our

empirical data (*Medium - Slow* = 4.96 dB; *Fast - Medium* = 4.37 dB).

Previous research has attributed faster CFS breakthrough (equivalently, lower

contrast) to unconscious processing of suppressed images (Gayet et al., 2014; Mudrik et al.,

2011). As the current study found uniform suppression depth for all tested images, even

though bCFS thresholds varied, it is clear that differences in bCFS thresholds alone should

not be interpreted in terms of expeditious unconscious processing of semantically relevant

images. Indeed, if image categories such as faces were processed unconsciously, they

reasonably should be harder to re-suppress, and thus have a smaller suppression depth

compared to neutral stimuli (see Figure 1) – which was not the case.

As an alternative to lower bCFS thresholds being due to unconscious processing of

images with relevant semantic content, such images (here, faces and objects) may break

suppression at lower contrasts because they are particularly rich in low-level feature content

that promotes image salience (e.g., (Itti & Koch, 2000)). For example, faces and objects

could be more salient due to peaks in local image contrast (Parkhurst & Niebur, 2004),

contour integration (Kapadia et al., 2000), closed curvilinear form (concavity: (Schmidtmann

et al., 2015), and phase aligned spatial frequency spectra (Maehara et al., 2009). The

combination of these image properties plausibly render images such as faces and objects

particularly salient within early stages of visual processing and thus lead to lower bCFS

thresholds than those measured for less salient stimuli such as gratings or scrambled noise.

Moreover, this derivation of salience is purely stimulus-based and arises without regard to

semantic implications. This possibility, which has been raised by others (Gayet et al., 2014;

Moors, 2019; Moors et al., 2016, 2017; Moors & Hesselmann, 2018), has appeal on grounds

of parsimony. With the advent of tCFS, it is now possible to measure bCFS and reCFS for

salience-indexed stimuli to learn the extent to which stimulus-based salience impacts

suppression depth.

Some caution is warranted here, however. It is not clear that all variation in bCFS

thresholds can be explained by low-level image properties. There may be important high-

level factors that also contribute to the salience of a given target image that make it visible at

lower contrasts than other images (Gayet et al., 2014; Jiang et al., 2007; Mudrik et al., 2011;

Yang et al., 2007). For example, faces provide essential social information and we are very

highly attuned to them. Face images may therefore be salient at lower contrasts than other

images such as random noise for high-level reasons specific to face processing. However,

given there is a parsimonious account of low bCFS thresholds for faces in terms of low-level

image properties, this would need to be established experimentally, perhaps using face

inversion (Matsuyoshi et al., 2015) or contrast reversal (Liu-Shuang et al., 2022),

manipulations that differentially impact visual processing of faces compared to other object

categories This matter is still open and ripe for further experimental evidence, but our

current findings add support to the argument that faces may become visible in CFS at lower

contrasts for reasons unrelated to special access to awareness or partial processing during

suppression..

A hybrid-model might therefore be needed for a full account of CFS, similar to those

proposed for binocular rivalry (Cao et al., 2021; Wilson, 2003). Based on our results here,

we imagine a hybrid model in which relative suppression depth for a given image arises from

low-level interocular mutual inhibition acting equivalently on any kind of image (yielding the

uniform suppression depth we observe), and the absolute level of breakthrough threshold

could be impacted by high-level salience (e.g., (Hesse & Tsao, 2020)). Careful manipulation

of low- and high-level properties, in combination with the tCFS method, should be able to

test this model, as discussed earlier.

Suppression depth is modulated by rate of contrast change

Experiment 3 varied the rate of target contrast change with the expectation that this

would alter the magnitude of adaptation during tCFS. We predicted that a faster contrast

change would reduce the opportunity for adaptation to accrue, thereby requiring a greater

change in contrast to overcome suppression during CFS. Similarly, a slower rate of change

should increase the opportunity for adaptation, resulting in the inverse effect. We observed

strong modulations of suppression depth based on the rate of contrast change, confirming

these predictions (**Figure 4**). Follow-up analyses confirmed that all three rates of contrast

change had distinct percept duration times (Figure 4 - figure supplement 1), indicating that

the differences in suppression depth we observed were not due to an artefact such as

participants responding with a fixed inter-response interval, which would spuriously increase
suppression depth for a fast rate of change.
667
Damped harmonic oscillator model
We modelled the changes in contrast between successive bCFS and reCFS
thresholds over a trail and found that a damped harmonic oscillator (DHO) provided an
excellent fit to these sequential estimates of suppression depth. The applicability of this
model is noteworthy for a number of reasons. First, in neuroscience, the DHO model
provides a valuable mathematical framework for understanding the dynamics of neural
systems and their responses to external stimuli, particularly with regard to the interplay
between excitation and inhibition (Freeman, 1961; Hodgkin & Huxley, 1952; Spyropoulos et
al., 2022). In the present context, the high starting contrast of the suprathreshold target in
tCFS trials is analogous to the external perturbation. The asymptotic differences in
thresholds over time are reminiscent of both earlier (Wilson, 2003), and more recent
computational models (Cao et al., 2021) of interocular competition, models proposing that
changes to visual awareness are driven by an out-of-equilibrium cortical network. We note
that the locus of competing neural ensembles could reside in early visual stages (Alais et al.,
2010; Lankheet, 2006; Li et al., 2017), late stages (Hohwy et al., 2008) or across hierarchical
(Cao et al., 2021; Wilson, 2003) levels of visual processing. Although it is beyond the scope
of the present work, future studies could vary the starting conditions of the tCFS procedure,
or manipulate higher-order influences such as attention and expectation to examine whether
the return to equilibrium we have revealed conforms to the predictions of competing models.
687
688
Future Directions
The tCFS method equips researchers with a convenient method to measure bCFS and
reCFS thresholds, and thus suppression depth. We have used tCFS here to establish that a
uniform suppression depth exists across image categories, and that differences in bCFS
thresholds alone cannot provide strong evidence for unconscious processing. Many
substantive questions remain. For example, the depth of interocular suppression is reported
to partially depend on spatial feature similarity between the competing images (Alais &
Melcher, 2007; Drewes et al., 2023) and their temporal frequency (Han et al., 2018; Han &
Alais, 2018). These factors could be parametrically varied to examine specifically whether
they modulate bCFS thresholds alone, or whether they also cause a change in suppression
depth by asymmetrically affecting reCFS thresholds. Previous findings can easily be
revisited, such as results showing that bCFS varies with manipulations of semantic content
(e.g., face inversion, or manipulating a face's emotion), results which form part of the

claimed evidence for expedited unconscious processing of certain suppressed images.

Conclusion

Across three experiments we have introduced the tCFS method and shown that

traditional evidence for unconscious processing – based on differences in the threshold to

reach awareness (bCFS threshold) – provide only half the story. Misleading conclusions

about unconscious processing must be supported by measures of suppression depth, which

can be calculated as the difference between both breakthrough (bCFS) and suppression

(reCFS) thresholds. Using the tCFS method we have measured these thresholds, and found

uniform suppression depth across five image five categories. Notably, this uniform

suppression depth is increased with reduced opportunity for target image adaptation, as is

the case when target contrast changes rapidly. Collectively, the three tCFS experiments

refute existing claims of high-level semantic information or target complexity influencing the

depth of unconscious processing during interocular suppression. Future findings may yet

confirm differences in suppression depth in certain circumstances, yet this will require

measurement of both breakthrough and suppression thresholds to demonstrate the requisite

changes in suppression depth.

General Materials and Methods

Participants

A total of 36 undergraduate psychology participants volunteered in exchange for course credit. All participated with informed consent and had normal or corrected-to-normal vision. Our sample size for Experiment 1 was based on power estimates to detect a moderate sized effect in a 2 x 2 repeated-measures design (Faul et al., 2009), while also exceeding the typical sample size used in bCFS studies to compensate for our novel paradigm (e.g., n = 10, Cha et al., 2019, n = 10–16, Han et al., 2021). We adjusted our power analysis after observing a strong effect size for the difference between bCFS and reCFS thresholds when using the tCFS method, resulting in fewer participants in Experiments 2 and 3. Experiment 1: N = 20, (15 females), Experiment 2: N = 18 (12 females), Experiment 3, N = 17 (11 females). All participants in Experiment 3 also did Experiment 2, and one author (JC) also completed experiments 2 and 3. This study was approved by the University of Sydney Human Research Ethics Committee (HREC 2021/048).

Apparatus

Visual stimuli were displayed on a Mac Pro (2013; 3.7 GHz Quad-Core Intel Xeon E5) computer, displayed on an Apple LED Cinema monitor (24 inch, 1920 x 1200 pixel resolution, 60 Hz refresh rate), running OS X El Capitan (10.11.6). All experiments were programmed using custom MATLAB code, and displayed using Matlab (ver R2017b) and Psychtoolbox (ver 3.0.13; Brainard et al., 1997). Responses were collected via the left mouse button of the right hand. A mirror stereoscope was used to partition participant's vision into separate left- and right-eye views, located approximately 51 cm from the screen, with a total optical path length of 57 cm.

Stimuli

Participants dichoptically viewed a high-contrast Mondrian mask pattern (400 x 400 pixels, 7° x 7°) with one eye and a small target stimulus (130 x 130 pixels, 2.2° x 2.2°) with the other eye. Two binocularly presented white squares surrounded the mask and served as a fusion lock to maintain stable fusion, and each eye had a central fixation cross (18 x 18 pixels; 0.3° x 0.3°). The Mondrian pattern was greyscale and consisted of overlapping circles of various sizes and intensities and was updated every fifth video frame (12 Hz). The mask's RMS contrast ranged between 0.07 and 0.09. As previous research has indicated that achromatic masks may be optimal to suppress achromatic targets (reviewed in (Pournaghdali & Schwartz, 2020), we opted for grayscale Mondrian patterns to match our targets.

In all experiments targets were viewed by the right eye and were selected at random

from a set that was standardised in RMS contrast (20%) and mean luminance (before

contrast ramping was applied). Target contrast was ramped up or down by scaling the target

image's standard contrast within a range of .02 to 1.0. Importantly, all contrast scaling was

done on a logarithmic scale in decibel units (i.e., conDb = 20 x log10(con)) to make the

changing contrast effectively linear, thus coinciding with the visual system's logarithmic

contrast response function. Minimum (.02) and maximum (1.0) contrast values were thus -

33.98 and 0 dB, respectively, and contrast steps were .07 dB per video frame. In experiment

3, where the rate of target contrast change was manipulated, the contrast steps were .035,

.07, or .105 dB units per video frame. Target location varied between trials, drawn from a

uniform distribution of 200 x 200 pixels centred on the fixation cross.

Procedure

Participants were given practice trials until they were familiar with CFS and the task used in

these experiments. Participants were instructed to respond via mouse click the moment their

subjective visibility of the target stimulus changed (either when a visible target became

suppressed, or when a suppressed target became visible). Realizing that phenomenological

quality of reversals may differ among participants (Moors et al., 2017; Zadbood et al., 2011),

we encouraged participants in our study to establish a criteria for target

appearance/disappearance in the practice session and to maintain it throughout the

experiment. In all experiments, the dependent variable was the target contrast at the

moment when a change in target visibility was judged to have occurred (either breaking

suppression or succumbing to suppression).

Experiment 1 - Using tCFS to measure suppression depth

Our first experiment tested two different procedures for measuring reCFS and bCFS

thresholds: a continuous tracking procedure and a discrete trials procedure similar to that

used in previous bCFS research. We hypothesised that a difference between bCFS and

reCFS thresholds would provide evidence for a contrast range (i.e., suppression depth)

between awareness and suppression, in contrast to the possibility of a given contrast

threshold determining a narrow awareness/suppression border. By comparing the results

between discrete and continuous methods, we sought to establish the feasibility of collecting

multiple thresholds within a single-trial, allowing the rapid quantification of suppression depth

in CFS paradigms. Experiment 1 compared CFS thresholds for targets (faces and objects)

increasing in contrast and decreasing in contrast (i.e., bCFS and reCFS thresholds) in

discrete trials and in continuous tracking trials, in a 2 x 2 repeated-measures, within-subjects

design. In the discrete conditions, participants completed eight blocks of eight trials, during

which target contrast always changed in one direction - either increasing from low to high as

is typically done in bCFS studies to measure breakthrough thresholds, or decreasing from

high contrast to low to measure a suppression threshold. In the continuous condition, the

target contrast tracked down and up continuously, reversing direction after each participant

response. Continuous trials always began with the target decreasing from maximum contrast

so that the participant's first response was to report when it disappeared, which caused

target contrast to increase until breakthrough was reported, which caused it to decrease

again until suppression, etc.. Continuous trials terminated after 16 reports of change in target

visibility. When the contrast time series is plotted as shown in Figure 2b, the plot shows eight

upper turning points where the target broke into awareness (bCFS thresholds) and eight

lower turning points where the target became re-suppressed (reCFS thresholds). The order

of discrete and continuous blocks was counterbalanced and randomized across participants.

Images for the eight trials of each block type were drawn from the same set of four faces and

four naturalistic objects, with no repetitions within a block. Before each block began,

participants completed a series of practice trials that utilised an independent set of six

images. They were able to complete practice trials as many times as they wished until they

had confidently established interocular fusion and were comfortable with the requirements of

the task.

Experiment 2 - The effect of image category

Experiment 2 tested the suppression depth obtained for different image categories.

Experiment 2 used only the continuous tracking 'tCFS' method and compared five image

types (faces, familiar objects, linear gratings, phase scrambled images, and polar patterns

that were radial lines or concentric circles). The trials contained 20 reports (10 bCFS and 10

reCFS thresholds) and the data were analysed in a 5 (image type) x 2 (bCFS vs reCFS

thresholds) within-subjects, repeated-measures ANOVA. There were 10 tracking trials, with

each trial containing a single target image from a subset of ten (two of each image category,

randomly ordered for each participant).

Experiment 3 - Rate of contrast change on suppression depth

Experiments 1 and 2 introduced the tCFS method and demonstrated a uniformity of

suppression depth across multiple target image categories. This uniformity of suppression

depth could indicate that neural events mediating CFS suppression are not selective for

complexity or semantic meaning, and like popular models of binocular rivalry, could instead

be based on low-level reciprocal inhibition and neural adaptation processes (Alais et al.,

2010; Kang & Blake, 2010; McDougall, 1901), in which case suppression depth should vary

with the rate of contrast change of the monocular target. More specifically, based on the

adapting mutual inhibition model, we predicted that at a slower rate of contrast change

neural adaptation for the monocular target would increase, lowering the amount of contrast

change necessary to transition between visibility states. Similarly, a faster rate of target

contrast change would reduce the time for neural adaptation of the monocular target,

resulting in an increase in the required change in contrast necessary to transition a target

into and out of awareness. Expressed in operational terms, the depth of suppression should

increase with the rate of target change, which was the focus of Experiment 3.

841   Experiment 3 used the tCFS method and compared the rate of target contrast

change (slow, medium, fast) across four image categories (faces, objects, linear gratings,

and phase scrambled images). The trials contained 20 reports (10 bCFS and 10 reCFS

thresholds) and the data were analysed in a 3 (contrast change rate) x 4 (image type) x 2

(bCFS vs reCFS thresholds) repeated-measures, within-subjects design. The medium rate

of change was the same as used in Experiments 1 and 2, and the slow and fast rates were

0.5 and 1.5 times the medium rate, respectively. There were 12 tracking trials, given by the

factorial combination of four target image types repeated at the three rates of target contrast

change (in a randomised order for each participant).

Data Analysis

852   Data analysis was performed in Matlab (ver R2022a), and SPSS/JASP (ver 28).

853   Initial inspection identified 1 participant for exclusion (from Experiment 3), based on failure to

854   follow task instructions. For visualization and analysis, all contrast thresholds are expressed

855   in decibel units.

856   We performed Bayesian model comparison to quantify evidence for and against the

null in JASP, using Bayesian repeated measures ANOVAs (uninformed prior with equal

weight to all models). We report Bayes factors (B) for main effects of interest (e.g. effect of

image type on suppression depth), as evidence in favour compared to the null model ($BF_{10}=$

B). Following the guidelines recommended in (Dienes, 2021), B values greater than 3

indicate moderate evidence for H1 over H0, and B values less than 1/3 indicate moderate

evidence in favour of the null. B values residing between 1/3 and 3 are interpreted as weak

evidence, or an insensitivity of the data to distinguish between the null and alternative

models.

Data Availability

867   Analysis code to reproduce the main results and figures are available in a public

868   repository (github.com/Davidson-MJ/tCFS-elife-VOR). Raw data and analysis code are also

869   available at (https://osf.io/nzp9v/).

Model Fitting

We additionally quantified the change in relative contrast over time, and evaluated a
series of model fits to describe these data. For this analysis, bCFS and reCFS thresholds
were first averaged within their respective response number, enabling a comparison of
thresholds over the course of each trial. The modelling used the absolute change in contrast
between each sequential threshold in the tracking series as its dependent variable, which we
modelled after detrending, per participant, and at the group level.
We compared three basic models to this data. All models were fit with a non-linear
least-squares approximation using a maximum of 400 iterations (lsqcurvefit.m in MATLAB)
The first was a simple cubic polynomial with three free parameters:

1) $a\square^3 + b\square^2 + c\square$

where a, b and c are coefficients for the cubic, quadratic and linear term. We also fit a simple
harmonic oscillator with three free parameters:

2) $a \times \sin(b \times t + c)$

Where a is amplitude, b is frequency, c is a phase offset, and t is time. We also fit a damped
harmonic oscillator model with four free parameters:

3) $a \times e^{(-b \times t)} \times \sin(c \times t + d)$.

Where a and b describe the amplitude and damping coefficient of decay, and c and d
describe the frequency and phase shift of the oscillatory response.
To fit each model, we linearly interpolated between the turning points (thresholds) in
the tCFS time series of each trial to increase the observations to 1000 samples, and
estimated the goodness of each fit through a series of steps. First, we calculated the sum of
squared residual errors for each fit (SSE), and calculated the Bayesian Information Criterion
(BIC) using equation 4:

4) $BIC = n \times \log(SSE/n) + k \times \log(n)$;

Where n represents the number of observations in the dataset, SSE is the sum of squared
errors, and k is the number of parameters in the model. The BIC allows a comparison of

model fits while taking into account the goodness of fit and complexity of each model. It

includes a penalty on the number of parameters in the model by including a term that scales

with the logarithm of sample size. When comparing two models, the model with a lower BIC

is considered favourable, with a change of 0 to 2 BIC as weak evidence in favour, and 6 to

10 as strong evidence in favour (Kass & Raftery, 1995),

Figure 4 - figure supplement 1. Perceptual durations in Experiments 1 to 3. a) Average

histogram across participants for all percept durations by experimental condition. Solid lines

show tCFS conditions, broken lines show discrete (unidirectional) conditions. Shading

corresponds to ±1 SEM corrected for within participant comparisons (Cousineau, 2005). **b-c)**

Average histograms for tCFS in Experiment 2, and Experiment 3, respectively. Figure

conventions as in a). **d)** tCFS durations of Experiment 3 when split by rate of contrast

change. Vertical lines and arrows indicate the mean of median percept durations across

participants.

Appendix 1: Difference thresholds as a measure of suppression depth and the role of
contrast response functions

One question raised in review concerned using a decibel contrast scale and the calculation
of suppression depth as the difference between breakthrough and suppression thresholds.
The justification for this is twofold. First, neurons in early visual cortex exhibit a logarithmic-
like response to contrast that grows rapidly at low contrasts and then saturates at higher
contrasts. Data from single V1 neurons show that these contrast response functions (CRFs)
to linear contrast can be very adequately described by fitting a logarithmic curve, but not by
linear or power functions (Albrecht & Hamilton, 1982). Consistent with a logarithmic
response, perceptual thresholds for contrast increments tend to be a constant proportion of
the base contrast (i.e, Weber's law holds). For these reasons, logarithmically scaled contrast
changes should therefore produce approximately linear changes in perceived contrast and
should be used when ramping contrast in CFS procedures. Second, once contrast is log-
transformed to decibels, the difference calculation used to obtain suppression depth is
effectively a ratio. It is edifying to learn, as we show here, that suppression depth is constant
as this indicates that whatever value the bCFS threshold has, the reCFS threshold will be *n*
times lower. For example, a difference of -12 dB is a factor of four (e.g., bCFS = .2, reCFS =
.05). It is important, though, to plot bCFS and reCFS values as well as suppression depth
because these thresholds could be higher or lower for a different image category and yet still
produce the same suppression depth (e.g., values of bCFS = .6 and reCFS = .15 also
produce a suppression depth of -12 dB but are very different in absolute terms). Plotting
bCFS and reCFS thresholds and suppression depth ensures absolute shifts are not
obscured.

It is interesting to consider that the shapes of CRFs vary at different stages of cortical
processing. CRFs tend to be shallower in V1 than in subsequent areas such as V2, V4 and
FFA where they are considerably steeper. Appendix 1 - figure 1 plots two CRFs, modelled
with Naka-Rushton functions, showing a steep and a shallow version. The Naka-Rushton
equation provides a very good description of the CRF (Albrecht & Hamilton, 1982) and is the
preferred way to model them. The shallower CRF typifies that measured from neurons in
primary visual cortex. It has a C50 (contrast giving half-max response) of 0.24 and an
exponent of 3.4 – the mean values of a sample of 98 primate V1 CRFs reported by (Albrecht
& Hamilton, 1982). The steeper CRF is typical of a V4 CRF and has a C50 of 0.15 and an
exponent of 1.9 – the mean values of 131 V4 CRFs reported by (Cheng et al., 1994; Williford
& Maunsell, 2006). For the sake of illustrating a key point, we have added horizontal lines
labelled bCFS and reCFS threshold. We make the simple assumption that when contrast is

ramping up, a target will break through from suppression to visibility when response level
reaches the 'bCFS threshold'. Conversely, a visible target ramping down in contrast will
become suppressed at the 'reCFS threshold'.
966
The vertical lines in Appendix 1 - figure 1 show the bCFS/reCFS threshold on the contrast
axis, revealing a smaller range of 0.24 – 0.33 for the V1 CRF than for the V4 CRF (0.15 –
0.265). When expressed in decibels ($SD_{dB} = 20 * log10(reCFS/bCFS)$) these thresholds
produce suppression depths of 2.77 dB for the V1 CRF and 4.94 dB for the V4 CRF. This
implies that CFS suppression should be deeper if it were occurring at the level of V4, and
shallower if it were occurring in V1.
973
The suppression values calculated from the figure are smaller than those we report here in
our data. However, it is important to note that the example CRFs in Appendix 1- figure 1 are
taken from recordings of single neurons, whereas an individual percept, such as a grating
emerging into visibility, will be the result of a whole population of neurons. This is significant
because when CRFs are derived from pooled neural responses using brain imaging
techniques such as EEG or MEG population CRFs are much broader (Avidan et al., 2002;
Hall et al., 2005). Importantly they broaden yet still maintain the same pattern as plotted in
Appendix 1 - figure 1 where V1 CRFs show a more graded response to contrast and CRFs
for areas beyond V1 show left-shifted and steeper CRFs. Therefore, the broader CRFs at
the population level would produce much bigger bCFS/reCFS threshold differences and
consequently a greater suppression depth that would be closer to the values we report here.

Outlining a realistic model of CRFs not only adds clarity about one of the key mechanisms
underlying the contrast ramping procedure used in tCFS, it also allows us to make
predictions and test hypotheses. For example: does CFS suppression depend on the type of
target image? Neurons in area V4 are selective for polar form (Gallant et al., 1996) but V1
neurons are not. V1 neurons, on the other hand, are highly selective for linear gratings.
Neither area is selective for faces, yet area FFA is specialised for this (Bao & Tsao, 2018)
and shares similar CRFs to V4. If CFS suppression were to occur in the area specialised for
a given target stimulus, our CRF model makes two predictions: (i) that the bCFS and reCFS
thresholds should be very different when gratings are compared to polar gratings or faces
(V1 bCFS/reCFS thresholds should be higher), and (ii) suppression depth should be greater
for polar gratings and faces than for gratings. However, neither of these predictions is
supported by our data. Figure 3a shows clearly that across different kinds of target images
bCFS and reCFS thresholds tend to be very similar. Indeed, post hoc t-tests comparing
gratings with objects, faces, and polar gratings show no significant differences between

bCFS thresholds (all ps > .572) or reCFS thresholds (all ps > .285). Moreover, the level of

suppression depth is the same (Fig. 3b). This pattern is not consistent with target images

being suppressed in their preferred cortical areas but with suppression occurring in a single

area governed by a common CRF. Based on our findings, we suggest the critical single area

for CFS suppression is most likely area V1 because: (i) this is the first cortical area where

interocular mismatches are detected, and (ii) regardless of image type, all target images

tested in our study are very much mismatched with the Mondrian mask and would likely

trigger interocular suppression at this first binocular stage. It remains a possibility that future

work will uncover evidence of higher-level areas feeding into CFS suppression in a kind of

hybrid model that has found support in studies of binocular rivalry suppression (van Boxtel et

al., 2008; Wilson, 2003). One way to do that might be to create targets and masks that are

congruent in as many features as possible to circumvent a front-end suppression triggered

by an initial interocular mismatch.

1012

1013

Appendix 1 - figure 1 **. Contrast response functions and CFS thresholds.** Two Naka-

Rushton functions modelling contrast response functions (CRFs) are shown, one has

parameters typical of primary visual cortical neurons (blue) and the other is typical of CRFs

found in area V4 (red). The C50 and exponent values are taken from single-unit

neurophysiological studies of primate areas V1 and V4. We make the simple assumption

that when contrast is ramping up, a target will break through from suppression to visibility

when response level reaches the 'bCFS threshold'. Conversely, a visible target ramping

down in contrast will become suppressed at the 'reCFS threshold'. Note that the V4 CRF has

bCFS/reCFS thresholds that are much lower than those from the V1 CRF, and that the

bCFS/reCFS range is greater for the V4 CRF, implying greater suppression depth (see

appendix text). Our finding of constant suppression depth and very similar bCFS/reCFS

thresholds across diverse image types suggests that CFS suppression for all our tested

target images occurs in a common mechanism likely to be early in binocular processing.

1027

References

Alais, D. (2012). Binocular rivalry: competition and inhibition in visual perception. *Wiley Interdisciplinary Reviews. Cognitive Science*, *3*(1), 87–103.

Alais, D., & Blake, R. (2014). Binocular rivalry and perceptual ambiguity. In J. Wagemans (Ed.), *Oxford Handbook of Perceptual Organization.* Oxford University Press.

Alais, D., Cass, J., O'Shea, R. P., & Blake, R. (2010). Visual sensitivity underlying changes in visual consciousness. *Current Biology: CB*, *20*(15), 1362–1367.

Alais, D., & Melcher, D. (2007). Strength and coherence of binocular rivalry depends on shared stimulus complexity. *Vision Research*, *47*(2), 269–279.

Albrecht, D. G., & Hamilton, D. B. (1982). Striate cortex of monkey and cat: contrast response function. *Journal of Neurophysiology*, *48*(1), 217–237.

Anderson, B. L. (1992). Hysteresis, cooperativity, and depth averaging in dynamic random-dot stereograms. *Perception & Psychophysics*, *51*(6), 511–528.

Avidan, G., Harel, M., Hendler, T., Ben-Bashat, D., Zohary, E., & Malach, R. (2002). Contrast sensitivity in human visual areas and its relationship to object recognition. *Journal of Neurophysiology*, *87*(6), 3102–3116.

Bao, P., & Tsao, D. Y. (2018). Representation of multiple objects in macaque category-selective areas. *Nature Communications*, *9*(1), 1774.

Blake, R. (2022). The Perceptual Magic of Binocular Rivalry. *Current Directions in Psychological Science*, *31*(2), 139–146.

Blake, R., & Camisa, J. (1979). On the inhibitory nature of binocular rivalry suppression. *Journal of Experimental Psychology. Human Perception and Performance*, *5*(2), 315–323.

Blake, R., & Fox, R. (1974). Binocular rivalry suppression: insensitive to spatial frequency and orientation change. *Vision Research*, *14*(8), 687–692.

Breitmeyer, B. G. (2015). Psychophysical "blinding" methods reveal a functional hierarchy of unconscious visual processing. *Consciousness and Cognition*, *35*, 234–250.

1056 Buckthought, A., Kim, J., & Wilson, H. R. (2008). Hysteresis effects in stereopsis and

1057   binocular rivalry. *Vision Research*, *48*(6), 819–830.

1058 Cao, R., Pastukhov, A., Aleshin, S., Mattia, M., & Braun, J. (2021). Binocular rivalry reveals

1059   an out-of-equilibrium neural dynamics suited for decision-making. *eLife*, *10*, e61581.

1060 Cheng, K., Hasegawa, T., Saleem, K. S., & Tanaka, K. (1994). Comparison of neuronal

1061   selectivity for stimulus speed, length, and contrast in the prestriate visual cortical areas

1062   V4 and MT of the macaque monkey. *Journal of Neurophysiology*, *71*(6), 2269–2280.

1063 Cousineau, D. (2005). Confidence intervals in within-subject designs: A simpler solution to

1064   Loftus and Masson's method. *Tutorials in Quantitative Methods for Psychology*, *1*(1),

1065   42–45.

1066 Dienes, Z. (2021). How to use and report Bayesian hypothesis tests. *Psychology of*

1067   *Consciousness: Theory, Research, and Practice*, *8*(1), 9–26.

1068 Drewes, J., Witzel, C., & Zhu, W. (2023). Feature-based interaction between masks and

1069   target in continuous flash suppression. *Scientific Reports*, *13*(1), 4696.

1070 Fang, F., & He, S. (2005). Cortical responses to invisible objects in the human dorsal and

1071   ventral pathways. *Nature Neuroscience*, *8*(10), 1380–1385.

1072 Faul, F., Erdfelder, E., Buchner, A., & Lang, A.-G. (2009). Statistical power analyses using

1073   G*Power 3.1: tests for correlation and regression analyses. *Behavior Research*

1074   *Methods*, *41*(4), 1149–1160.

1075 Freeman, W. J. (1961). Harmonic Oscillation as Model for Cortical Excitability Changes with

1076   Attention in Cats. *Science*, *133*(3470), 2058–2059.

1077 Gallant, J. L., Connor, C. E., Rakshit, S., Lewis, J. W., & Van Essen, D. C. (1996). Neural

1078   responses to polar, hyperbolic, and Cartesian gratings in area V4 of the macaque

1079   monkey. *Journal of Neurophysiology*, *76*(4), 2718–2739.

1080 Gayet, S., Paffen, C. L. E., Belopolsky, A. V., Theeuwes, J., & Van der Stigchel, S. (2016).

1081   Visual input signaling threat gains preferential access to awareness in a breaking

1082   continuous flash suppression paradigm. *Cognition*, *149*, 77–83.

1083 Gayet, S., Van der Stigchel, S., & Paffen, C. L. E. (2014). Breaking continuous flash

suppression: competing for consciousness on the pre-semantic battlefield. *Frontiers in*

Psychology, *5*, 460.

Grossman, E., Donnelly, M., Price, R., Pickens, D., Morgan, V., Neighbor, G., & Blake, R.

(2000). Brain areas involved in perception of biological motion. *Journal of Cognitive*

Neuroscience, *12*(5), 711–720.

Hall, S. D., Holliday, I. E., Hillebrand, A., Furlong, P. L., Singh, K. D., & Barnes, G. R. (2005).

Distinct contrast response functions in striate and extra-striate regions of visual cortex

revealed with magnetoencephalography (MEG). *Clinical Neurophysiology: Official*

Journal of the International Federation of Clinical Neurophysiology, *116*(7), 1716–1722.

Han, S. 'er, & Alais, D. (2018). Strength of continuous flash suppression is optimal when

target and masker modulation rates are matched. *Journal of Vision*, *18*(3), 3.

Han, S. 'er, Blake, R., & Alais, D. (2018). Slow and steady, not fast and furious: Slow

temporal modulation strengthens continuous flash suppression. *Consciousness and*

Cognition, *58*, 10–19.

Hassin, R. R. (2013). Yes It Can: On the Functional Abilities of the Human Unconscious.

Perspectives on Psychological Science: A Journal of the Association for Psychological

Science, *8*(2), 195–207.

Hesse, J. K., & Tsao, D. Y. (2020). Representation of conscious percept without report in the

macaque face patch network. In *bioRxiv* (p. 2020.04.22.047522).

https://doi.org/10.1101/2020.04.22.047522

Hesselmann, G., & Moors, P. (2015). Definitely maybe: can unconscious processes perform

the same functions as conscious processes? *Frontiers in Psychology*, *6*, 584.

Hodgkin, A. L., & Huxley, A. F. (1952). A quantitative description of membrane current and

its application to conduction and excitation in nerve. *The Journal of Physiology*, *117*(4),

500–544.

Hohwy, J., Roepstorff, A., & Friston, K. (2008). Predictive coding explains binocular rivalry:

an epistemological review. *Cognition*, *108*(3), 687–701.

Holender, D. (1986). Semantic activation without conscious identification in dichotic listening,

parafoveal vision, and visual masking: A survey and appraisal. *The Behavioral and Brain Sciences*, *9*(1), 1–23.

Hong, S. W. (2015). Radial bias for orientation and direction of motion modulates access to visual awareness during continuous flash suppression. *Journal of Vision*, *15*(1), 15.1.3.

Itti, L., & Koch, C. (2000). A saliency-based search mechanism for overt and covert shifts of visual attention. *Vision Research*, *40*(10-12), 1489–1506.

Jiang, Y., Costello, P., & He, S. (2007). Processing of Invisible Stimuli: Advantage of Upright Faces and Recognizable Words in Overcoming Interocular Suppression. *Psychological Science*, *18*(4), 349–355.

Julesz, B., & Tyler, C. W. (1976). Neurontropy, an entropy-like measure of neural correlation, in binocular fusion and rivalry. *Biological Cybernetics*, *23*(1), 25–32.

Kang, M.-S., & Blake, R. (2010). What causes alternations in dominance during binocular rivalry? *Attention, Perception & Psychophysics*, *72*(1), 179–186.

Kapadia, M. K., Westheimer, G., & Gilbert, C. D. (2000). Spatial distribution of contextual interactions in primary visual cortex and in visual perception. *Journal of Neurophysiology*, *84*(4), 2048–2062.

Kass, R. E., & Raftery, A. E. (1995). Bayes Factors. *Journal of the American Statistical Association*, *90*(430), 773–795.

Kim, C.-Y., & Blake, R. (2005). Psychophysical magic: rendering the visible "invisible." *Trends in Cognitive Sciences*, *9*(8), 381–388.

Kouider, S., & Dehaene, S. (2007). Levels of processing during non-conscious perception: a critical review of visual masking. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, *362*(1481), 857–875.

Lanfranco, R. C., Rabagliati, H., & Carmel, D. (2022). The importance of awareness in face processing: A critical review of interocular suppression studies. *Behavioural Brain Research*, 114116.

Lankheet, M. J. M. (2006). Unraveling adaptation and mutual inhibition in perceptual rivalry. *Journal of Vision*, *6*(4), 304–310.

Li, H.-H., Rankin, J., Rinzel, J., Carrasco, M., & Heeger, D. J. (2017). Attention model of

binocular rivalry. *Proceedings of the National Academy of Sciences of the United States*

of America, *114*(30), E6192–E6201.

Ling, S., & Carrasco, M. (2006). When sustained attention impairs perception. *Nature*

Neuroscience, *9*(10), 1243–1245.

Liu-Shuang, J., Yang, Y.-F., Rossion, B., & Goffaux, V. (2022). Natural Contrast Statistics

Facilitate Human Face Categorization. *eNeuro*, *9*(5).

https://doi.org/10.1523/ENEURO.0420-21.2022

Logothetis, N. K. (1998). Single units and conscious vision. *Philosophical Transactions of the*

Royal Society of London. Series B, Biological Sciences, *353*(1377), 1801–1818.

Lunghi, C., & Pooresmaeili, A. (2023). Learned value modulates the access to visual

awareness during continuous flash suppression. *Scientific Reports*, *13*(1), 756.

Maehara, G., Huang, P.-C., & Hess, R. F. (2009). Importance of phase alignment for

interocular suppression. *Vision Research*, *49*(14), 1838–1847.

Maglio, S. J., & Polman, E. (2016). Revising probability estimates: Why increasing likelihood

means increasing impact. *Journal of Personality and Social Psychology*, *111*(2), 141–

158.

Matsuyoshi, D., Morita, T., Kochiyama, T., Tanabe, H. C., Sadato, N., & Kakigi, R. (2015).

Dissociable cortical pathways for qualitative and quantitative mechanisms in the face

inversion effect. *The Journal of Neuroscience: The Official Journal of the Society for*

Neuroscience, *35*(10), 4268–4279.

McDougall, W. (1901). Some New Observations in Support of Thomas Young's Theory of

Light- and Colour-Vision. *Mind; a Quarterly Review of Psychology and Philosophy*,

10(37), 52–97.

Moors, P. (2019). What's Up with High-Level Processing During Continuous Flash

Suppression? In *Transitions between Consciousness and Unconsciousness* (pp. 39–

70). https://doi.org/10.4324/9780429469688-2

Moors, P., Boelens, D., van Overwalle, J., & Wagemans, J. (2016). Scene Integration

Without Awareness: No Conclusive Evidence for Processing Scene Congruency During

Continuous Flash Suppression. *Psychological Science*, *27*(7), 945–956.

Moors, P., & Hesselmann, G. (2018). A critical reexamination of doing arithmetic

nonconsciously. In *Psychonomic Bulletin & Review* (Vol. 25, Issue 1, pp. 472–481).

https://doi.org/10.3758/s13423-017-1292-x

Moors, P., Hesselmann, G., Wagemans, J., & van Ee, R. (2017). Continuous Flash

Suppression: Stimulus Fractionation rather than Integration. *Trends in Cognitive*

Sciences, *21*(10), 719–721.

Mudrik, L., Breska, A., Lamy, D., & Deouell, L. Y. (2011). Integration without awareness:

expanding the limits of unconscious processing. *Psychological Science*, *22*(6), 764–770.

Mudrik, L., Faivre, N., & Koch, C. (2014). Information integration without awareness. *Trends*

in Cognitive Sciences, *18*(9), 488–496.

Newell, B. R., & Shanks, D. R. (2014). Unconscious influences on decision making: a critical

review. *The Behavioral and Brain Sciences*, *37*(1), 1–19.

Nguyen, V. A., Freeman, A. W., & Alais, D. (2003). Increasing depth of binocular rivalry

suppression along two visual pathways. *Vision Research*, *43*(19), 2003–2008.

Parkhurst, D. J., & Niebur, E. (2004). Texture contrast attracts overt visual attention in

natural scenes. *The European Journal of Neuroscience*, *19*(3), 783–789.

Pastukhov, A., Vivian-Griffiths, S., & Braun, J. (2015). Transformation priming helps to

disambiguate sudden changes of sensory inputs. *Vision Research*, *116*(Pt A), 36–44.

Piotrowski, L. N., & Campbell, F. W. (1982). A demonstration of the visual importance and

flexibility of spatial-frequency amplitude and phase. *Perception*, *11*(3), 337–346.

Pournaghdali, A., & Schwartz, B. L. (2020). Continuous flash suppression: Known and

unknowns. *Psychonomic Bulletin & Review*, *27*(6), 1071–1103.

Rees, G. (2007). Neural correlates of the contents of visual awareness in humans.

Philosophical Transactions of the Royal Society of London. Series B, Biological

Sciences, *362*(1481), 877–886.

Schmidtmann, G., Jennings, B. J., & Kingdom, F. A. A. (2015). Shape recognition:

convexities, concavities and things in between. *Scientific Reports*, *5*(1), 1–11.

Schmidt, T. (2015). Invisible Stimuli, Implicit Thresholds: Why Invisibility Judgments Cannot

be Interpreted in Isolation. *Advances in Cognitive Psychology / University of Finance*

and Management in Warsaw, *11*(2), 31–41.

Sklar, A. Y., Levy, N., Goldstein, A., Mandel, R., Maril, A., & Hassin, R. R. (2012). Reading

and doing arithmetic nonconsciously. *Proceedings of the National Academy of*

Sciences, *109*(48), 19614–19619.

Spyropoulos, G., Saponati, M., Dowdall, J. R., Schölvinck, M. L., Bosman, C. A., Lima, B.,

Peter, A., Onorato, I., Klon-Lipok, J., Roese, R., Neuenschwander, S., Fries, P., &

Vinck, M. (2022). Spontaneous variability in gamma dynamics described by a damped

harmonic oscillator driven by noise. *Nature Communications*, *13*(1), 2019.

Stein, T. (2019). The breaking continuous flash suppression paradigm. In G. Hesselmann

(Ed.), *Transitions between consciousness and unconsciousness* (Vol. 1). Routledge1.

Stein, T., & Sterzer, P. (2014). Unconscious processing under interocular suppression:

getting the right measure. *Frontiers in Psychology*, *5*, 387.

Sterzer, P., Hilgenfeldt, T., Freudenberg, P., Bermpohl, F., & Adli, M. (2011). Access of

emotional information to visual awareness in patients with major depressive disorder.

Psychological Medicine, *41*(8), 1615–1624.

Sterzer, P., Stein, T., Ludwig, K., Rothkirch, M., & Hesselmann, G. (2014). Neural

processing of visual information under interocular suppression: a critical review.

Frontiers in Psychology, *5*(MAY), 1–12.

Stuit, S. M., Paffen, C. L. E., & Van der Stigchel, S. (2023). Prioritization of emotional faces

is not driven by emotional content. *Scientific Reports*, *13*(1), 1–9.

Tsuchiya, N., & Koch, C. (2005). Continuous flash suppression reduces negative

afterimages. *Nature Neuroscience*, *8*(8), 1096–1101.

Tsuchiya, N., Koch, C., Gilroy, L. A., & Blake, R. (2006). Depth of interocular suppression

associated with continuous flash suppression, flash suppression, and binocular rivalry.

Journal of Vision, *6*(10), 1068–1078.

Tsuchiya, N., Moradi, F., Felsen, C., Yamazaki, M., & Adolphs, R. (2009). Intact rapid detection of fearful faces in the absence of the amygdala. *Nature Neuroscience*, *12*(10), 1224–1225.

van Boxtel, J. J. A., Alais, D., & van Ee, R. (2008). Retinotopic and non-retinotopic stimulus encoding in binocular rivalry and the involvement of feedback. *Journal of Vision*, *8*(5), 17.1–10.

van Dam, L. C. J., & van Ee, R. (2006a). Retinal image shifts, but not eye movements per se, cause alternations in awareness during binocular rivalry. *Journal of Vision*, *6*(11), 1172–1179.

van Dam, L. C. J., & van Ee, R. (2006b). The role of saccades in exerting voluntary control in perceptual and binocular rivalry. *Vision Research*, *46*(6-7), 787–799.

Vanduffel, W., Fize, D., Peuskens, H., Denys, K., Sunaert, S., Todd, J. T., & Orban, G. A. (2002). Extracting 3D from motion: differences in human and monkey intraparietal cortex. *Science*, *298*(5592), 413–415.

Wang, G., Alais, D., Blake, R., & Han, S. 'er. (2022). CFS-crafter: An open-source tool for creating and analyzing images for continuous flash suppression experiments. *Behavior Research Methods*. https://doi.org/10.3758/s13428-022-01903-7

Williams, D., Phillips, G., & Sekuler, R. (1986). Hysteresis in the perception of motion direction as evidence for neural cooperativity. *Nature*, *324*(6094), 253–255.

Williford, T., & Maunsell, J. H. R. (2006). Effects of spatial attention on contrast response functions in macaque area V4. *Journal of Neurophysiology*, *96*(1), 40–54.

Wilson, H. R. (2003). Computational evidence for a rivalry hierarchy in vision. *Proceedings of the National Academy of Sciences of the United States of America*, *100*(24), 14499–14503.

Yang, E., Zald, D. H., & Blake, R. (2007). Fearful expressions gain preferential access to awareness during continuous flash suppression. *Emotion* , *7*(4), 882–886.

Yu, K., & Blake, R. (1992). Do recognizable figures enjoy an advantage in binocular rivalry? *Journal of Experimental Psychology. Human Perception and Performance*, *18*(4), 1158–

1173.

Zadbood, A., Lee, S.-H., & Blake, R. (2011). Stimulus fractionation by interocular

suppression. *Frontiers in Human Neuroscience*, *5*, 135.

Zhou, G., Zhang, L., Liu, J., Yang, J., & Qu, Z. (2010). Specificity of face processing without

awareness. *Consciousness and Cognition*, *19*(1), 408–412.

Zhou, L.-F., Wang, K., He, L., & Meng, M. (2021). Twofold advantages of face processing

with or without visual awareness. *Journal of Experimental Psychology. Human*

Perception and Performance, *47*(6), 784–794.

Zhou, W., Jiang, Y., He, S., & Chen, D. (2010). Olfaction modulates visual perception in

binocular rivalry. *Current Biology: CB*, *20*(15), 1356–1358.

Blake, R., Goodman, R., Tomarken, A., & Kim, H. W. (2019). Individual differences in

continuous flash suppression: Potency and linkages to binocular rivalry dynamics.

Vision Research, *160*, 10-23.

Piotrowski, L. N., & Campbell, F. W. (1982). A demonstration of the visual Importance and

flexibility of spatial-frequency amplitude and phase. *Perception*, *11*(3), 337-346.

https://doi.org/10.1068/p110337

1268

1269   **ACKNOWLEDGEMENTS**
1270   RB was supported by funds from a Centennial Professorship Research Award, Vanderbilt
1271   University, with thanks to Jeffrey Schall. DA and MD were supported by an Australian
1272   Research Council grant (DP210101691).