Integrin β4 promotes DNA damage drug resistance in triple-negative breast cancer via TNFAIP2/IQGAP1/Rac1

Huan Fang[1, 2#], Wenlong Ren[1,3#], Qiuxia Cui[4, 5, 1#], Huichun Liang[1#], Chuanyu Yang[1], Wenjing Liu[1], Xinye Wang[1], Xue Liu[6], Yujie Shi[7*], Jing Feng[8, 6*], Ceshi Chen[9,10,1*]

1. Key Laboratory of Animal Models and Human Disease Mechanisms, Kunming Institute of Zoology, Chinese Academy of Sciences. Kunming, Yunnan, China.

2. Kunming College of Life Sciences, University of Chinese Academy Sciences, Kunming, Yunnan, China.

3. School of Life Science, University of Science & Technology of China. Hefei, Anhui, China.

4. Affiliated Hospital of Guangdong Medical University. Zhanjiang, Guangdong, China.

5. Department of Breast Surgical Oncology, National Cancer Center/National Clinical Research Center for Cancer/Cancer Hospital & Shenzhen Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College, Shenzhen, Guangdong, China.

6. Shanghai University of Medicine & Health Sciences, Affiliated Sixth People's Hospital South Campus, Shanghai, China.

7. Department of Pathology, Henan Provincial People's Hospital, Zhengzhou University, Zhengzhou, Henan, 450003, China.

8. The Second Affiliated Hospital of the Chinese University of Hong Kong (Shenzhen), Shenzhen, 518172, China.

9. Academy of Biomedical Engineering, Kunming Medical University, Kunming 650500, China.

10. The Third Affiliated Hospital, Kunming Medical University, Kunming 650118,

China.

Short Title: ITGB4 promotes drug resistance in TNBC through TNFAIP2

# These authors contribute equally

* Corresponding authors: Ceshi Chen, E-mail: chenc@kmmu.edu.cn, Jing Feng,

E-mail: fengjing71921@163.com, or Yujie Shi, E-mail: shiyujie523@126.com

Summary

Anti-tumor drug resistance is a challenge for triple-negative breast cancer

treatment. Our previous work demonstrated that TNFAIP2 activates Rac1 to

promote triple-negative breast cancer cell proliferation and migration. However,

the mechanism by which TNFAIP2 activates Rac1 is unknown. In this study,

we found that TNFAIP2 interacts with IQGAP1 and Integrin β4. Integrin β4

activates Rac1 through TNFAIP2 and IQGAP1 and confers DNA damage drug

resistance in triple-negative breast cancer. These results indicate that the

Integrin β4/TNFAIP2/IQGAP1/Rac1 axis provides potential therapeutic targets

to overcome DNA damage drug resistance in triple-negative breast cancer.

Keywords

ITGB4, TNFAIP2, IQGAP1, Rac1, drug resistance, DNA damage repair,

triple-negative breast cancer

Introduction

Breast cancer is the most common malignant tumor in women, and the

morbidity and mortality rates of female malignant tumors are ranked first in the

world[1]. Although the diagnosis and treatment of breast cancer has entered the

era of molecular typing,35% of breast cancers still experience recurrence,

metastasis and treatment failure[2]. According to the expression of estrogen

receptor (ERα), progesterone receptor (PR) and human epidermal growth

factor receptor (HER2), breast cancer is divided into ER/PR-positive,

HER2-positive and triple-negative breast cancer (TNBC)[3].For ER/PR-positive

and HER2-positive breast cancer, endocrine therapies such as tamoxifen and

anti-HER2 targeted therapy such as trastuzumab have achieved good efficacy.

Targeted drugs for TNBC patients with BRCA1/2 mutations include two PARP

inhibitors, olaparib and talazoparib. These targeted drugs cannot fully meet the

clinical needs of patients with various TNBC subtypes[4]. Currently, DNA

damage chemotherapy drugs, including epirubicin and cisplatin, are widely

used for TNBC treatment.

TNBC often recurs and metastasizes due to the development of

chemoresistance, although it is initially responsive to chemotherapeutic drugs[5].

Chemoresistance severely impacts the clinical outcomes of patients. Tumor

cells become resistant to chemotherapeutic agents through several

mechanisms, such as improving DNA damage repair, changing the

intracellular accumulation of drugs, or increasing anti-apoptotic mechanisms[6].

Therefore, characterization of the underlying molecular mechanisms by which

resistance occurs will provide opportunities to develop precise therapies to

enhance the efficacy of standard chemotherapy regimens [7,8].

TNFAIP2 is abnormally highly expressed in a variety of tumor cells, including

TNBC[9], nasopharyngeal carcinoma [10], malignant glioma[11], uroepithelial

carcinoma[12]and esophageal squamous cell carcinoma[13], and is associated

with poor prognosis. Our previous work [9,14]showed that TNFAIP2, as a KLF5

downstream target protein, can interact with Rac1[15], a member of the Rho

small GTP enzyme family, and activate Rac1 to alter the cytoskeleton, thereby

inducing filopodia and lamellipodia formation and promoting the adhesion,

migration and invasion of TNBC cells. After activation, Rac1 can activate AKT,

PAKs, NADPH oxidase and other related signaling pathways to promote cell

survival, proliferation, adhesion, migration and invasion[16]. Activation of Rac1

can reduce the therapeutic response to trastuzumab in breast cancer and

increase the resistance of TNBC cells to paclitaxel[17], but the specific

mechanism of action is not completely clear.

Rac1 has been shown to play an important role in DNA damage repair.

Activated Rac1 can promote the phosphorylation of the DNA damage

response-related molecules ATM/ATR, CHK1/2 and H2AX by activating the

activity of protein kinases such as ERK1/2, JNK and p38 [18,19], thus improving

the DNA damage repair ability and inhibiting tumor cell apoptosis[20-22]. Rac1

also promotes aldolase release and activation by changing the cytoskeleton

and activates the ERK pathway to increase thepentose phosphate pathway to

promote nucleic acid synthesis, providing more raw materials for DNA damage

repair[23,24]. At the same time, the interaction of Rac1 and PI3K promoted AKT

phosphorylation and glucose uptake[25,26]. Therefore, Rac1 is well established

to promote the chemoresistance of breast cancer by promoting DNA damage

repair.

Integrin β4 (ITGB4) is a major component of hemidesmosomes and a receptor

molecule of laminin. Studies have shown that laminin-5 interacts with ITGB4 to

activate Rac1 activity and promote cell migration[27] and polarization[28] by

altering the cytoskeleton. Since ITGB4-positive cancer stem cell

(CSC)-enriched mesenchymal cells were found to reside in an intermediate

epithelial/mesenchymal phenotypic state, ITGB4 can be used to enable

stratification of mesenchymal-like TNBC cells[29]. In addition, the expression of

ITGB4 on ALDH[high] breast cancer and head and neck cancer cells was

significantly greater than that on ALDH[low] cells, proving the effects that ITGB4

targets on both bulk and CSC populations[30]. Furthermore,

ITGB4-overexpressing TNBC cells provided cancer-associated fibroblasts

(CAFs) with ITGB4 proteins via exosomes, and ITGB4-overexpressing

CAF-conditioned medium promoted the proliferation,

epithelial-to-mesenchymal transition, and invasion of breast cancer cells[31].

ITGB4 also promotes breast cancer cell resistance to tamoxifen-induced

apoptosis by activating the PI3K/AKT signaling pathway and promotes breast

cancer cell resistance to anoikis by activating Rac1[32]. However, how ITGB4

activates Rac1 is not completely clear.

Rac1 activity is regulated by guanylate exchange factors (GEFs), GTPase

activation proteins (GAPs), and guanine separation inhibitors (GDIs)[33]. GAPs

typically provide the necessary catalytic groups for GTP hydrolysis, but not all

GAPs function as hydrolases. IQGAP1 lacks an arginine in the GTPase

binding domain and cannot exert the hydrolysis effect of GAPs[34]. IQGAP1 can

increase the activity of Rac1 and CDC42[35,36].

In this study, we demonstrated that TNFAIP2 interacts with IQGAP1 and

ITGB4. ITGB4 promotes TNBC drug resistance via the

TNFAIP2/IQGAP1/Rac1 axis by promoting DNA damage repair. Our results

suggest that ITGB4 and TNFAIP2 might serve as promising therapeutic targets

for TNBC.

Results

TNFAIP2 promotes TNBC DNA damage drug resistance

To explore the functional significance of TNFAIP2 in TNBC drug resistance, we constructed stable TNFAIP2 overexpression and TNFAIP2 knockdown HCC1806 and HCC1937 cells. As shown in Figure 1A-E, overexpression of TNFAIP2 significantly increased cell viability when treated with EPI and BMN. Additionally, knockdown of TNFAIP2 significantly decreased cell viability when treated with EPI and BMN (Fig. 1F-J). We then examined the effects of TNFAIP2 on DNA damage repair and found that TNFAIP2 promotes DNA damage repair in response to EPI and BMN. TNFAIP2 overexpression decreased the protein expression levels of γH2AX, a marker of DNA damage, and cleaved-PARP, a marker of apoptosis (Fig. 1K). Additionally, knockdown of TNFAIP2 significantly increased γH2AX and cleaved-PARP protein expression levels in response to EPI and BMN in both cell lines (Fig. 1L).

The function of TNFAIP2 was further validated by using two other DNA damage drugs, DDP and AZD (Fig. S1A-L). These results suggested that TNFAIP2 enhances TNBC cell drug resistance by promoting DNA damage repair.

TNFAIP2 confers TNBC drug resistance *in vivo*

To test whether TNFAIP2 also decreases the sensitivity of TNBC cells to EPI

and BMN *in vivo*, we performed animal experiments in nude mice. HCC1806

cells with stable TNFAIP2 knockdown were orthotopically inoculated into the

fat pad of 7-week-old female mice (n=8 or 12/group). When the tumor mass

reached approximately 50 mm^3, each group was divided into two subgroups to

receive either EPI (2.5 mg/kg, twice a week) or vehicle control for 23 days and

either BMN (1 mg/kg, twice a week) or vehicle control for 29 days. We

observed that depletion of TNFAIP2 suppressed breast cancer cell growth *in*

vivo. This is consistent with our previous report[9]. More importantly, TNFAIP2

depletion further decreased tumor volume when mice were treated with EPI

and BMN (Fig. 2A-F). Meanwhile, BMN treatment had no effect on the body

weight of mice (Fig. S2F). Consistently, EPI and DDP generated similar results

but decreased mouse body weight due to their high toxicity (Fig.S2D-E). These

results suggest that inhibition of TNFAIP2 expression can overcome HCC1806

breast cancer cell drug resistance in animals.

TNFAIP2 promotes TNBC drug resistance and DNA damage repair via

Rac1

Since chemotherapeutic agents and PRAP inhibitors induce DNA damage

directly or indirectly, DNA damage repair ability profoundly affects the

sensitivity of cancer cells to these therapies[37,38]. Since TNFAIP2 can activate

Rac1, a well-known drug resistance protein, we investigated whether TNFAIP2

induces chemotherapeutic resistance through Rac1. We found that Rac1

knockdown rescued the effects of TNFAIP2 overexpression-induced drug

resistance to EPI and BMN in HCC1806 and HCC1937 cells (Fig. 3A-F). We

also found that γH2AX and cleaved-PARP protein levels were up-regulated

again in Rac1 knockdown and TNFAIP2-overexpressing HCC1806 and

HCC1937 cells in response to EPI and BMN (Fig. 3G-J). We obtained similar

results by using DDP and AZD treatment (Fig. 3SA-J). Collectively, these

results suggest that TNFAIP2 promotes DNA damage repair and drug

resistance via Rac1.

IQGAP1 mediates Rac1 activation by TNFAIP2 and promotes TNBC drug

resistance

To characterize the mechanism by which TNFAIP2 activates Rac1, we

performed an IP-MS experiment. We found that TNFAIP2 interacts with

IQGAP1 (Fig. 4A). To validate whether TNFAIP2 interacts with IQGAP1, we

constructed HCC1806 cells with stable Flag-TNFAIP2 overexpression and

collected Flag-tagged TNFAIP2 cell lysates for immunoprecipitation assays

using Flag-M2 beads (Fig. S4A). We performed immunoprecipitation using

anti-IQGAP1 antibody and found that endogenous IQGAP1 protein

interacted with endogenous TNFAIP2 protein inHCC1806 cells (Fig. 4B). Next,

we mapped the regions of TNFAIP2 and IQGAP1 proteins responsible for the

interaction by generating a series of Flag-TNFAIP2 deletion mutants and

transfected them into HEK293T cells together with full-length IQGAP1. Then,

we performed immunoprecipitation assays using Flag-M2 beads (Figure S4B),

and the N-terminus (1-79 aa) of the TNFAIP2 protein interacted with IQGAP1.

To explore the function of IQGAP1 in TNBC drug resistance, we knocked down

IQGAP1 in HCC1806 and HCC1937 cells. As shown in Figure 4C-G,

knockdown of IQGAP1 significantly decreased cell viability in the presence of

EPI and BMN in both cell lines. We also examined the effects of IQGAP1 on

DNA damage repair and found that IQGAP1 promotes DNA damage repair.

IQGAP1 knockdown increased γH2AX and cleaved-PARP protein expression

levels when HCC1806 and HCC1937 cells were treated with EPI and BMN

(Fig. 4H). Next, we found that IQGAP1 knockdown abrogated the effects of

TNFAIP2 overexpression on resistance to EPI and BMN (Fig. 4I-K, Fig.

S4C-E). We also found that γH2AX and cleaved-PARP protein levels were

up-regulated in IQGAP1 knockdown and TNFAIP2-overexpressing HCC1806

and HCC1937 cells (Fig. 4L). In addition, we found that the TNFAIP2

overexpression-induced increase in Rac1 activity was abolished by IQGAP1

knockdown (Fig. 4M).

ITGB4 interacts with TNFAIP2 and promotes TNBC drug resistance and

DNA damage repair

In addition to IQGAP1, TNFAIP2 may interact with ITGB4 (Fig. 4A). To validate

whether TNFAIP2 interacts with ITGB4, we immunoprecipitated exogenous

Flag-tagged TNFAIP2 proteins from HCC1806 cells by using Flag-M2 beads

and detected endogenous ITGB4 proteins(Fig. 5A).To further confirm the

protein–protein interaction between endogenousTNFAIP2 and ITGB4proteins,

we collected HCC1806 cell lysates and performed immunoprecipitation using

an anti-TNFAIP2/ITGB4 antibody and found that endogenous TNFAIP2/ITGB4

protein interacted with endogenous ITGB4/TNFAIP2 protein (Fig. 5B-C).We

further mapped the regions of TNFAIP2 and ITGB4 proteins responsible for the

interaction(Fig. S5I) by generating a series of Flag-TNFAIP2/GST-fused

TNFAIP2 deletion mutants and transfected them into HEK293T cells together

with full-length GST-fused ITGB4/ITGB4. Then, we performed

immunoprecipitation assays using Flag-M2 beads and glutathione beads. As

shown in Figure S5J-K, the N-terminus (218-287 aa) of the TNFAIP2

(TNFAIP2-S-N1-3) protein interacted with ITGB4. To map the domains of

ITGB4 that interact with TNFAIP2, we transfected Flag-tagged full-length

TNFAIP2 into HEK293T cells with full-length or truncated ITGB4. We found

that the C-terminus (710-740 aa) of the ITGB4 protein interacted with TNFAIP2

(Fig. S5L-M). Taken together, these results suggest that TNFAIP2 interacts

with ITGB4 and that their interaction is mediated through the N-terminus of

TNFAIP2 and the C-terminus of ITGB4.

To explore the function of ITGB4 in TNBC drug resistance, we knocked down

ITGB4 in HCC1806 and HCC1937cells. As shown in Figure 5D-I, knockdown

of ITGB4 significantly decreased cell viability in the presence of EPI and BMN

in both cell lines. Knockdown of ITGB4 also suppressed HCC1806 xenograft

growth *in vivo*. More importantly, ITGB4 knockdown further decreased tumor

volume when mice were treated with EPI and BMN (Figure. 5J-N). Meanwhile,

BMN treatment had no effect on the body weight of mice, but EPI treatment

decreased mouse body weight due to its toxicity (Fig. S5H). We then

examined the effects of ITGB4 on DNA damage repair and found that ITGB4

promotes DNA damage repair in response to EPI and BMN. ITGB4 knockdown

increased γH2AX and cleaved-PARP protein expression levels when

HCC1806 and HCC1937 cells were treated with EPI and BMN (Fig. 5O).

Furthermore, the function of ITGB4 was validated by using two other drugs,

DDP and AZD (Fig. S5A-G). These results suggested that ITGB4 increases

TNBC drug resistance and promotes DNA damage repair.

ITGB4 activates Rac1 through TNFAIP2 and IQGAP1

It is well known that ITGB4 can activate Rac1[27] and that TNFAIP2 interacts

with Rac1 and activates it[9]. To test whether ITGB4 activates Rac1 through

TNFAIP2, we measured the levels of GTP-bound Rac1 in

ITGB4-overexpressing and ITGB4-knockdown cells. Overexpression of ITGB4

significantly increased the levels of GTP-bound Rac1 in bothHCC1806 and

HCC1937 cells (Fig. 6A). In agreement with this observation, knockdown of

ITGB4 significantly decreased the levels of GTP-bound Rac1 in both cell lines

(Fig. 6B). Next, we knocked down TNFAIP2 in ITGB4-overexpressing

HCC1806 and HCC1937 cells and found that ITGB4-increased Rac1 activity

was blocked by TNFAIP2 knockdown (Fig. 6C-D). Collectively, these results

demonstrate that ITGB4 activates Rac1 through TNFAIP2.

It has been reported that Rac1 activity is promoted by IQGAP1[34] and that

TNFAIP2 activates Rac1 through IQGAP1 (Fig. 4P). We wondered whether

ITGB4 activates Rac1 through IQGAP1; therefore, we knocked down IQGAP1

in HCC1806 and HCC1937 cells with stable overexpression of ITGB4 and

found that the ITGB4-induced increase in Rac1 activity was abolished by

IQGAP1 knockdown (Fig. 6E-F). These results suggest thatITGB4 activates

Rac1 through TNFAIP2 and IQGAP1.

ITGB4 promotes TNBC drug resistance via TNFAIP2/IQGAP1/Rac1

Since ITGB4, TNFAIP2, and IQGAP1 promote drug resistance by promoting

DNA damage repair in TNBC, we wondered whether ITGB4 promoted drug

resistance through the TNFAIP2/IQGAP1/Rac1 axis. We knocked down

TNFAIP2, IQGAP1, and Rac1in ITGB4-overexpressing cells and found that

blocking the TNFAIP2/IQGAP1/Rac1 axis increased the sensitivity of

ITGB4-overexpressing HCC1806 (Fig. 7A-I) and HCC1937 cells to EPI and

BMN (Fig. S6O-W).We also found that γH2AX and cleaved-PARP levels were

upregulated in TNFAIP2/IQGAP1/Rac1 knockdown HCC1806 and HCC1937

cells stably expressing ITGB4 in the presence of EPI and BMN (Fig. 7J-L, Fig.

S6X-Z). DDP and AZD treatment generated similar results (Fig. S6A-N).

Together, these results suggest that ITGB4 promotes DNA damage repair and

drug resistance via the TNFAIP2/IQGAP1/Rac1 axis.

TNFAIP2 expression levels positively correlated with ITGB4 in TNBC

tissues

To test whether ITGB4 and TNFAIP2 are coexpressed in TNBC, we collected

135 TNBC specimens for IHC (the IQGAP1 antibody did not work for IHC).

Clinical pathological parameters, including patient age, tumor size, lymph node

status at the time of diagnosis, and follow-up status, including adjuvant

treatment and tumor recurrence, were retrospectively obtained from the

Department of Pathology, Henan Provincial People's Hospital, Zhengzhou

University, China. We performed IHC analyses on two breast cancer tissue

chips containing a total of 135 patients with TNBC (Fig. 8A-D).TNFAIP2 and

ITGB4 protein expression levels were significantly positively correlated(Table

1).

Discussion

Chemotherapies, including EPI and DDP, are the main choice for TNBC

patients. Unfortunately, TNBC frequently develops resistance to

chemotherapy[39]. Currently, PARP inhibitors are effective for TNBC with

BRCA1/2 mutation or homologous recombination deficiency (HRD)[40-42]. PARP

inhibitors can cause DNA damage repair defects and have synergistic lethal

effects with HRD. Meanwhile, chemotherapy and PARP inhibitor resistance is

also a major problem in the clinic.

In this study, we first found that TNFAIP2 promotes TNBC drug resistance and

DNA damage repair through Rac1. Next, we found that TNFAIP2 interacts with

IQGAP1 and ITGB4. We verified that ITGB4 promotes TNBC drug resistance

and DNA damage repair through the TNFAIP2/IQGAP1/Rac1 axis.

Interestingly, we discovered for the first time that ITGB4 and TNFAIP2 promote

Rac1 activity through IQGAP1. Our study reveals that ITGB4 promotes TNBC

resistance through TNFAIP2-, IQGAP1-, and Rac1-mediated DNA damage

repair (Figure 7). This study provides new targets for reversing TNBC

resistance.

ITGB4 is well known to promote breast cancer stemness and can be activated

by laminin 5[43]. In addition, ITGB4 is generally in partner with ITGA6, which is

another marker of breast cancer stem cells[44] and drug resistance[43]. Therefore,

whether ITGA6 has similar functions needs further study. It was reported that

ITGB4 activates Rac1[45], but the mechanism is unclear. For the first time, we

revealed that ITGB4 activates Rac1 through TNFAIP2 and IQGAP1. More

importantly, ITGB4 promotes drug resistance through the

TNFAIP2/IQGAP1/Rac1 axis.

TNFAIP2 plays important roles in different cellular and physiological processes,

including cell proliferation, adhesion, migration, membrane TNT formation,

angiogenesis, inflammation and tumorigenesis[14]. We previously found that

TNFAIP2 was regulated by KLF5 and interacted with the small GTPases Rac1

and Cdc42, thereby regulating the actin cytoskeleton and cell morphology in

breast cancer cells[9]. However, the detailed mechanism is not clearly

understood. In this study, we found that IQGAP1 mediates this

process.IQGAP1 is a crucial regulator of cancer development by scaffolding

and facilitating different oncogenic pathways, especially Rac1/Cdc42, thus

affecting proliferation, adhesion, migration, invasion, and metastasis[46]. In

addition, IQGAP1 is increased during the differentiation of ovarian cancer stem

cells and promotes aggressive behaviors[47]. In our study, we found that

TNFAIP2 interacts with IQGAP1 and thus activates Rac1 to induce

chemotherapy and PARP inhibitor drug resistance.

Furthermore, TNFAIP2 was reported to induce epithelial-to-mesenchymal

transition and confer platinum resistance in urothelial cancer cells[12], and in

embryonic stem cell (ESC) differentiation, TNFAIP2 was found to be important

in controlling lipid metabolism, which supports the ESC differentiation process

and planarian organ maintenance[48]. Another study found that TNFAIP2

overexpression enhanced TNT-mediated autophagosome and lysosome

exchange, preventing advanced glycation end product (AGE)-induced

autophagy and lysosome dysfunction and apoptosis[49]. In cancer treatment,

TNFAIP2 was chosen as one of the six signature genes predicting

chemotherapeutic and immunotherapeutic efficacies, with high-senescore

patients benefiting from immunotherapy and low-senescore patients

responsive to chemotherapy[50].

These reports provide apossible explanation for previous studies showing that

ITGB4 is important in EMT and cancer stemness. According to our results that

there isan interaction between ITGB4 and TNFAIP2, ITGB4 might regulate

EMT and stemness through TNFAIP2.TNFAIP2 is one of the important factors

induced by tumor necrosis factor alpha (TNFα). Interestingly, TNFα release

could be induced by therapeutic drugs from multiple tumor cell lines. The

acquisition of docetaxel resistance was accompanied by increased constitutive

production of TNFα[51]. In addition, TNFα is a key tumor-promoting effector

molecule secreted by tumor-associated macrophages. *In vitro* neutralizing

TNFα was observed to inhibit tumor progression and improve the curative

effect of bevacizumab[52]. Therefore, the mechanism by which TNFα plays a

role in chemotherapeutic resistance in breast cancer should be further

investigated. TNFα neutralizing provides novel opinions elucidating drug

resistance.

For future studies, it will be important to develop *Tnfaip2* knockout mice to

investigate the exact role of TNFAIP2 physiologically. According to recent

studies and our findings, agents targeting the interaction among

ITGB4/TNFAIP2/IQGAP1 would be a promising trend for developing drugs to

overcome the resistance phenomenon.

In summary, ITGB4 and TNFAIP2 play important roles in breast cancer

chemoresistance. TNFAIP2 activates Rac1 to promote chemoresistance

through IQGAP1. In addition, ITGB4 activates Rac1 through TNFAIP2 and

IQGAP1 and confer DNA damage drug resistance in TNBC (Figure 8 E).

These results indicate that the ITGB4/TNFAIP2/IQGAP1/Rac1 axis provides

potential therapeutic targets to overcome DNA damage drug resistance in

TNBC.

Data availability statement

The authors confirm that the data supporting the findings of this study are

available within the article and its supplementary materials.

Acknowledgments

This work was supported by National Key R&D Program of China

(2020YFA0112300), National Natural Science Foundation of China (81830087,

U2102203, 81672624, 82102987, and 82203413), the Yunnan Fundamental

Research Projects (202101AS070050), and the Guangdong Foundation

Committee for Basic and Applied Basic Research projects

(2022A1515012420).

Author contributions

H.F., W.R., Q.C., H.L. and C.Y. conducted different parts of the experiments.

W.L., X.W., X.L. and Y.S. provided some advice about this project. C.C. and J.F.

designed the study. C.C., J.F., and Y.S. supervised the study. H.F. wrote the

original draft and W.R., Q.C., H.L., C.C. and J.F. reviewed and edited the draft.

Declaration of interests

The authors declare no conflicts of interest.

Figure titles and legends

Figure1. TNFAIP2 promotes TNBC DNA damage drug resistance

(A-E) Stable TNFAIP2 overexpression in HCC1806 and HCC1937 cells

significantly increased cell viability in the presence of EPI (0-1.6 µM) or BMN

(0-40 µM) treatment for 48h, as measured by the SRB assay. TNFAIP2 protein

expression was detected by WB.

(F-J) Stable TNFAIP2 knockdown in HCC1806 and HCC1937 cells

significantly decreased cell viability in the presence of EPI (0-1.6 µM) or BMN

(0-40 µM) treatment for 48h, as measured by the SRB assay. TNFAIP2 protein

expression was detected by WB.

(K) TNFAIP2 promoted DNA damage repair in the presence of EPI and BMN.

HCC1806 and HCC1937 cells stably overexpressing TNFAIP2 were treated

with 400 or 800 nM EPI for 48h and 10 µM BMN for 24h, respectively.

TNFAIP2, γH2AX, and PARP protein expression was detected by WB.

(L) TNFAIP2 knockdown increased DNA damage in the presence of EPI and

BMN. Stable TNFAIP2 knockdown cells were treated with 400 or 800 nM EPI

for 24 or 48h and 2.5 µM BMN for 24h. TNFAIP2, γH2AX, and PARP protein

expression was detected by WB.

Figure2. TNFAIP2 confers TNBC drug resistance *in vivo*

(A-F) TNFAIP2 knockdown increased the sensitivity of HCC1806 breast

cancer cells to EPI and BMN *in vivo*. HCC1806 cells with stable TNFAIP2

knockdown were transplanted into the fat pad of 7-week-old female nude mice.

When the average tumor size reached approximately 50 mm^3 after inoculation,

mice in each group were randomly divided into two subgroups (n=4/group) to

receive EPI (2.5 mg/kg), BMN (1 mg/kg) or vehicle control for 23 or 29 days

(A-B). Tumor size was measured twice a week (C-D), and tumor masses were

collected and weighed at the end of the experiments (E-F). *: $P<0.05$, **:

$P<0.01$, *t*-test.

Figure3. TNFAIP2 promotes TNBC drug resistance and DNA damage

repair via Rac1

(A-F) Rac1 knockdown abolished TNFAIP2-induced TNBC resistance to EPI

and BMN. HCC1806 (A-C) and HCC1937 (D-F) cells with stable TNFAIP2

overexpression were transfected with Rac1 or control siRNA, followed by

treatment with EPI (0-1600 nM) and BMN (0-40 μM) for 48 or 72 h, respectively.

Cell viability was measured by the SRB assay. Protein expression levels were

analyzed by WB.

(G-J) Rac1 depletion abolished TNFAIP2-induced DNA damage decrease in

response to EPI and BMN. HCC1806 (G-H) and HCC1937 (I-J) cells with

stable TNFAIP2 overexpression were transfected with Rac1 or control siRNA,

followed by treatment with EPI (400 or 800 nM) and BMN (10 μM) for 24h,

respectively. Protein expression levels were analyzed by WB.

Figure4. IQGAP1 mediates Rac1 activation by TNFAIP2 and promotes

TNBC drug resistance

(A) The IP-MS result of TNFAIP2 in HCC1806 cells.

(B) Endogenous TNFAIP2 interacts with IQGAP1 in HCC1806 cells.

Endogenous TNFAIP2 protein was immunoprecipitated using ananti-IQGAP1

antibody. Immunoglobulin (Ig)G served as the negative control. Endogenous

TNFAIP2 was detected by WB.

(C-G) IQGAP1 knockdown in HCC1806 and HCC1937 cells significantly

decreased cell viability in the presence of EPI (0-1600 nM) and BMN (0-40 μM),

as measured by the SRB assay. IQGAP1 protein expression was detected by

WB.

(H) IQGAP1 knockdown in HCC1806 and HCC1937 cells increased DNA

damage of EPI and BMN. HCC1806 and HCC1937 cells with IQGAP1

knockdown were treated with 800 nM EPI for 24h and 10 μM BMN for24h,

respectively. ITGB4, γH2AX, and PARP protein expression was detected by

WB.

(I-K) IQGAP1 knockdown abolished TNFAIP2-confered resistance to EPI and

BMN. HCC1806 cells with stable TNFAIP2 overexpression were transfected

with IQGAP1 or control siRNA, followed by treatment with EPI (0-1600 nM)

and BMN (0-40 µM) for 48 or 72 h, respectively. Cell viability was measured by

the SRB assay. IQGAP1 protein expression was detected by WB.

(L) IQGAP1 knockdown abolished TNFAIP2-confered resistance to EPI and

BMN. HCC1806 and HCC1937 cells with stable TNFAIP2 overexpression

were transfected with IQGAP1 or control siRNA, followed by treatment with

EPI (800 nM) and BMN (10 µM) for 24h, respectively. Protein expression levels

were analyzed by WB.

(M) IQGAP1 knockdown abolished TNFAIP2-confered Rac1 activation.

HCC1806 and HCC1937 cells with stable TNFAIP2 overexpression were

transfected with IQGAP1 or control siRNA. GTP-Rac1 levels were assessed

using PAK-PBD beads.

Figure5. ITGB4 interacts with TNFAIP2 and promotes TNBC drug

resistance and DNA damage repair

(A) TNFAIP2 interacts with ITGB4.HCC1806 cells with stable TNFAIP2

overexpression were collected from flag-tagged TNFAIP2 cell lysates for

immunoprecipitation assays using Flag-M2 beads, and ITGB4 was detected by

WB.

(B) Endogenous TNFAIP2 interacts with ITGB4 in HCC1806 cells.

Endogenous TNFAIP2 protein was immunoprecipitated using an anti-TNFAIP2

antibody. IgG served as the negative control. Endogenous ITGB4 was

detected by WB.

(C) Endogenous ITGB4 interacts with TNFAIP2 in HCC1806 cells.

Endogenous ITGB4 protein was immunoprecipitated using an anti-ITGB4

antibody. IgG served as the negative control. Endogenous TNFAIP2 was

detected by WB.

(D-I) ITGB4 knockdown in HCC1806 and HCC1937 cells significantly

decreased cell viability in the presence of EPI (0-800 nM) and BMN (0-40 µM),

as measured by the SRB assay. ITGB4 protein expression was detected by

WB.

(J-N) ITGB4 depletion promotes HCC1806 breast cancer cell sensitivity to EPI

and BMN treatment *in vivo*. HCC1806 cells with stable ITGB4 knockdown were

transplanted into the fat pad of 7-week-old female nude mice. When the

average tumor size reached approximately 50 mm^3 after inoculation, the mice

in each group were randomly divided into two subgroups (n=4/group) to

receive EPI (2.5 mg/kg), BMN (1 mg/kg) or vehicle control for 22 days(J).

Tumor masses were collected and weighed at the end of the experiments(K),

and tumor size was measured twice a week (L-N).*: *P*<0.05, **: *P*<0.01, *t*-test.

(O) ITGB4 knockdown increased DNA damage of EPI and BMN. HCC1806

and HCC1937 cells with ITGB4 knockdown were treated with 400 nM EPI for

24h and 5 µM BMN for 24h, respectively. ITGB4, γH2AX, and PARP protein

expression was detected by WB.

Figure6. ITGB4 activates Rac1 through TNFAIP2 and IQGAP1

(A) Overexpression of ITGB4 increased GTP-Rac1 levels in HCC1806 and

HCC1937 cells. GTP-Rac1 were assessed using PAK-PBD beads.

(B) Knockdown of ITGB4 by shRNA decreased GTP-Rac1 levels in HCC1806

and HCC1937 cells.

(C-D) ITGB4 activates Rac1 through TNFAIP2. HCC1806 (C) and HCC1937

(D) cells with stable ITGB4 overexpression were transfected with TNFAIP2 or

control siRNA.

(E-F) ITGB4 activates Rac1 through IQGAP1.HCC1806 (E) and HCC1937 (F)

cells with stable ITGB4 overexpression were transfected with IQGAP1 or

control siRNA.

Figure7. ITGB4 promotes TNBC drug resistance via

TNFAIP2/IQGAP1/Rac1

(A-C) ITGB4 promotes TNBC drug resistance through TNFAIP2. TNFAIP2

knockdown abolished ITGB4-induced resistance to EPI and BMN. HCC1806

cells with stable ITGB4 overexpression were transfected with TNFAIP2 or

control siRNA, followed by treatment with EPI (0-400 nM) and BMN (0-30 µM)

for 48 or 72 h, respectively. Cell viability was measured by the SRB assay.

Protein expression levels were analyzed by WB.

(D-F) ITGB4 promotes TNBC drug resistance through IQGAP1. HCC1806

cells with stable ITGB4 overexpression were transfected with IQGAP1 or

control siRNA, followed by treatment with EPI (0-800 nM) and BMN (0-40 µM)

for 48 or 72 h, respectively. Cell viability was measured by the SRB assay.

Protein expression levels were analyzed by WB.

(G-I) ITGB4 promotes TNBC drug resistance through Rac1. HCC1806 cells

with stable ITGB4 overexpression were transfected with Rac1 or control siRNA,

followed by treatment with EPI (0-400 nM) and BMN (0-40 µM) for 48 or 72 h,

respectively. Cell viability was measured by the SRB assay. Protein expression

levels were analyzed by WB.

(J) ITGB4 promotes DNA damage repair through TNFAIP2. HCC1806 cells

with stable ITGB4 overexpression were transfected with TNFAIP2 or control

siRNA, followed by treatment with EPI (400 nM) and BMN (5 µM) for 24h.

Protein expression levels were analyzed by WB.

(K) ITGB4 promotes DNA damage repair through IQGAP1. HCC1806 cells

with stable ITGB4 overexpression were transfected with IQGAP1 or control

siRNA, followed by treatment with EPI (400 nM) and BMN (5 µM) for 24h.

Protein expression levels were analyzed by WB.

(L) ITGB4 promotes DNA damage repair through Rac1. HCC1806 cells with

stable ITGB4 overexpression were transfected with Rac1 or control siRNA,

followed by treatment with EPI (400 nM) and BMN (5 µM) for 24h. Protein

expression levels were analyzed by WB.

Figure8. TNFAIP2 expression levels positively correlated with ITGB4 in

TNBC tissues

Representative IHC images of TNFAIP2 and ITGB4 protein expression in

breast cancer tissues are shown. The final expression assessment was

performed by combining the two scores (0–2=low, 6–7=high). A and B indicate

low scores, C and D indicate high scores, and Figure E is the work model of

this paper.

Table 1. The correlation of TNFAIP2 and ITGB4 protein expression levels
in human TNBC specimens.

STAR Methods

Cell lines and reagents

All cell lines used in this study, including HCC1806, HCC1937, and HEK293T

cells, were purchased from ATCC (American Type Culture Collection,

Manassas, VA, USA) and validated by STR (short tandem repeat) analysis.

HCC1806 and HCC1937 cells were cultured in RPMI 1640 medium

supplemented with 5% FBS. HEK293T cells were cultured in DMEM (Thermo

Fisher, Grand Island, USA) with 5% FBS at 37°C with 5% CO_2. Epirubicin(EPI)

(Cat#HY-13624A),cisplatin(DDP)(Cat#HY-17394), talazoparib (BMN)

(Cat#HY-16106), and olaparib (AZD) (Cat#HY-10162) were purchased from

MCE (New Jersey, USA).

Plasmid construction and stable TNFAIP2 and ITGB4 overexpression

We constructed the full-length *TNFAIP2/ITGB4* gene and then subcloned them

into the pCDH lentiviral vector. The packaging plasmids (including

pMDLg/pRRE, pRSV-Rev, and pCMV-VSV-G) and pCDH-TNFAIP2/ITGB4

expression plasmid were cotransfected into HEK293T cells (2×10^6 in 10 cm

plate) to produce lentivirus. Following transfection for 48 h, the lentivirus was

collected and used to infect HCC1806 and HCC1937 cells. Forty-eight hours

later, puromycin (2 µg/ml) was used to screen the cell populations.

Stable knockdown of TNFAIP2 and ITGB4

The pSIH1-H1-puro shRNA vector was used to expressTNFAIP2, ITGB4 and

luciferase(LUC)shRNAs.TNFAIP2shRNA#1,5'-GACUUGGGCUCACAGAUAA

-3';TNFAIP2shRNA#2,5'-GAUUGAGGUGGCCACUUAU-3';ITGB4shRNA#1,5'

-ACGACAGCTTCCTTATGTA-3';ITGB4shRNA#2,5'-CAGCGACTACACTATTG

GA-3';LuciferaseshRNA,5'-CUUACGCUGAGUACUUCGA-3'; HCC1806 and

HCC1937 cells were infected with lentivirus. Stable populations were selected

using 1 to 2 mg/mL puromycin. The knockdown effect was evaluated by

Western blotting.

RNA interference

The siRNA target sequences used in this study are as follows:

TNFAIP2siRNA#1,5'-GACUUGGGCUCACAGAUAA-3';TNFAIP2siRNA#2,5'-

GAUUGAGGUGGCCACUUAU-3';ITGB4siRNA#1,5'-ACGACAGCTTCCTTAT

GTA-3';ITGB4siRNA#2,5'-CAGCGACTACACTATTGGA-3';Rac1siRNA,5'-CG

GCACCACUGUCCCAACA-3';IQGAP1siRNA#1,5'-GCAGGTGGATTACTATAA

A-3';IQGAP1siRNA#2,5'-CUAGUGAAACUGCAACAGA-3'. All siRNAs were

synthesized by RiboBio (RiboBio, China) and transfected at a final

concentration of 50 nM.

Antibodies and Western blotting (WB)

The WB procedure has been described in our previous study[53]. Anti-TNFAIP2

(sc-28318), anti-ITGB4 (sc-9090) and anti-GAPDH (sc-25778) antibodies were

purchased from Santa Cruz Biotechnology (Santa Cruz, CA, USA). The

anti-PARP (#9542) antibody was purchased from CST. Anti-Rac1 (05–389)

and anti-γH2AX (3475627) antibodies were purchased from Millipore (Billerica,

MA, USA). Anti-β-actin (A5441) and anti-Tubulin (T5168) antibodies were

purchased from Sigma–Aldrich (St Louis, MO, USA).The anti-IQGAP1

(ab86064) antibody was purchased from Abcam.

Immunopurification and silver staining

Immunoprecipitation and silver staining lysates from HCC1806 cells stably

expressing Flag-TNFAIP2 were prepared by incubating the cells in lysis buffer

containing a protease inhibitor cocktail (MCE). Cell lysates were obtained from

approximately 2.5×10^8 cells, and after binding with anti-Flag M2 affinity gel

(Sigma) for 2 h as recommended by the manufacturer, the affinity gel was

washed with cold lysis plus 0.2% NP-40. FLAG peptide (Sigma) was applied to

elute the Flag-labeled protein complex as described by the vendor. The elutes

were collected and visualized on NuPAGE 4%-12% Bis-Trisgels (Invitrogen,

CA, USA) followed by silver staining with a silver staining kit (Pierce, Illinois,

USA). The distinct protein bands were retrieved and analyzed by LC–Mass.

Immunoprecipitation and GST pull-down

For exogenous interaction between ITGB4 and Flag-TNFAIP2, cell lysates

were directly incubated with anti-Flag M2 affinity gel (A2220; Sigm) overnight

at 4°C. For endogenous protein interaction, cell lysates were first incubated

with anti-TNFAIP2/ITGB4/IQGAP1 antibodies or mouse IgG/rabbit IgG

(sc-2028; Santa Cruz Biotechnology, CA, USA) and then incubated with

Protein A/G plusagarose beads (sc-2003; Santa Cruz BiotechnologyFor the

GST pulldown assay, cell lysates were directly incubated with

GlutathioneSepharose 4B (52-2303-00; GE Healthcare) overnight at 4°C. The

precipitates were washed four times with 1 ml of lysis buffer, boiled for 10

minutes with 1×SDS sample buffer, and subjected to WB analysis.

Cell viability assays

Cell viability was measured by SRB assays as described in our previous

study[54].Cell viability was measured by SRB assays. Briefly, cells were seeded

in 96-well plates. Then, the cells were cultured for the indicated timeand fixed

with 10% trichloroacetic acid at room temperature for30 min, followed by

incubation with 0.4% SRB (w/v) solution in 1% acetic acid for 20 min at room

temperature. Finally, SRB was dissolved in 10 mM unbuffered Tris base, and

the absorbance was measured at a wavelength of 530 nm on a plate reader

(Bio Tek, Vermont, USA).

Rac1 activation assays

Rac1 activation was examined using the Cdc42 Activation Assay Biochem Kit

(BK034, Cytoskeleton, Denver, USA) following them anufacturer's instructions.

Cells were harvested with cell lysis buffer, and1 mg of protein lysate in a 1 ml

total volume at 4°C was immediately precipitated with 10 μg of PAK-PBD

beads for 60 min with rotation. After washing three times with wash buffer,

agarose beads were resuspended in 30 μl of 2×SDS sample buffer and boiled

for 5 min. Rac1-GTP was examined by WB with an anti-Rac1 antibody.

Xenograft experiments

We purchased 6- to 7-week-old female BALB/c nude mice from SLACCAS

(Changsha, China). Animal feeding and experiments were approved by the

animal ethics committee of Kunming Institute of Zoology, Chinese Academy of

Sciences. HCC1806-shLuc, HCC1806-shTNFAIP2, or HCC1806-shITGB4

cells (1×10^6 in Matrigel (BD Biosciences, NY, USA)) were implanted into the

mammary fat pads of the mice. When the tumor volume reached

approximately 50 mm^3, the nude mice were randomly assigned to the control

and treatment groups (n = 4/group). EPI, BMN, and DDP were dissolved in

ddH$_2$O. The control group was given vehicle alone, and the treatment group

received EPI (2.5 mg/kg), BMN (1 mg/kg), and DDP (2.5 mg/kg) alone via

intraperitoneal injection every three days for 18 or 27 days. The tumor volume

was calculated as follows: tumor volume was calculated by the formula: ($\pi\times$

length $\times$ width^2)/6.

Immunohistochemical staining

Paraffin-embedded clinical TNBC specimens were obtained from the First

Affiliated Hospital, Zhengzhou University, Zhengzhou, China. Informed

consent was obtained from all subjects. Two tissue microarrays containing 135

TNBC breast cancer tissues were constructed. For the immunohistochemistry

(IHC) assay, the slides were deparaffinized, rehydrated, and pressure cooker

heated for 2.5 min in EDTA for antigen retrieval. Endogenous peroxidase

activity was inactivated by adding an endogenous peroxidase blocker

(OriGene, China) for 15 min at room temperature. Slides were incubated

overnight at 4°Cwith anti-TNFAIP2 (1:200) or anti-ITGB4 (1:500). After 12 h,

the slides were washed three times with PBS and incubated with secondary

antibodies (hypersensitive enzyme-labeled goat anti-mouse/rabbit IgG

polymer (OriGene, China) at room temperature for 20 min, DAB concentrate

chromogenic solution (1:200dilution of concentrated DAB chromogenic

solution), counterstained with 0.5% hematoxylin, dehydrated with graded

concentrations of ethanol for 3 min each (70%–80%–90%–100%),and finally

stained with dimethyl benzene immunostainedslides were evaluated by light

microscopy. The IHC signal was scored using the 'Allred Score' method.

Statistical analysis

All graphs were created using GraphPad Prism software version 5.0.

Comparisons between two independent groups were assessed by two-tailed

Student's t-test. One-way analysis of variance with least significant differences

was used for multiple group comparisons. P-values of <0.05, 0.01 or 0.001

were considered to indicate a statistically significant result, comparisons

significant at the 0.05 level are indicated by*, at the 0.01 level are indicated by

, or at the 0.001 level are indicated by *.

Supplemental information titles and legends

FigureS1. TNFAIP2 promotes TNBC DNA damage drug resistance

(A-E) Stable TNFAIP2 overexpression in HCC1806 and HCC1937 cells

significantly increased cell viability in the presence of DDP (0-40 μM) or AZD

(0-40 μM) treatment for 48h, as measured by the SRB assay. TNFAIP2 protein

expression was detected by WB.

(F-J) Stable TNFAIP2 knockdown in HCC1806 and HCC1937 cells

significantly decreased cell viability in the presence of DDP (0-40 μM) or AZD

(0-40 μM) treatment for 48h, as measured by the SRB assay. TNFAIP2 protein

expression was detected by WB.

(K) TNFAIP2 promoted DNA damage repair in the presence of DDP and AZD.

HCC1806 and HCC1937 cells stably overexpressing TNFAIP2 were treated

with 20 μM DDP for 24 h or 48h and 10 μM AZD for 24h, respectively. TNFAIP2,

γH2AX, and PARP protein expression was detected by WB.

(L) TNFAIP2 knockdown increased DNA damage in the presence of DDP and

AZD. Stable TNFAIP2 knockdown cells were treated with 2.5 or 20 μM DDP for

24h and 2.5 μM AZD for 24h. TNFAIP2, γH2AX, and PARP protein expression

was detected by WB.

FigureS2. TNFAIP2 confers TNBC drug resistance *in vivo*

(A-F) TNFAIP2 knockdown increased the sensitivity of HCC1806 breast

cancer cells to DDP *in vivo*. HCC1806 cells with stable TNFAIP2 knockdown

were transplanted into the fat pad of 7-week-old female nude mice. When the

average tumor size reached approximately 50 mm^3 after inoculation, mice in

each group were randomly divided into two subgroups (n=4/group) to receive

DDP (2.5 mg/kg) or vehicle control for 23 days (A). Tumor size was measured

twice a week (B), tumor masses were collected and weighed at the end of the

experiments (C), and mouse masses were collected and weighed at the

beginning or end of the experiments (D-F). *: *P*<0.05, **: *P*<0.01, *t*-test.

FigureS3. TNFAIP2 promotes TNBC drug resistance and DNA damage

repair via Rac1

(A-F) Rac1 knockdown abolished TNFAIP2-induced TNBC resistance to DDP

and AZD. HCC1806 (A-C) and HCC1937 (D-F) cells with stable TNFAIP2

overexpression were transfected with Rac1 or control siRNA, followed by

treatment with DDP (0-40 μM) and AZD (0-40 μM) for 48 or 72 h, respectively.

Cell viability was measured by the SRB assay. Protein expression levels were

analyzed by WB.

(G-J) Rac1 depletion abolished TNFAIP2-induced DNA damage decrease in

response to DDP and AZD. HCC1806 (G-H) and HCC1937 (I-J) cells with

stable TNFAIP2 overexpression were transfected with Rac1 or control siRNA,

followed by treatment with DDP (20 μM) and AZD (10 μM) for 24h, respectively.

Protein expression levels were analyzed by WB.

FigureS4. IQGAP1 mediates Rac1 activation by TNFAIP2 and promotes TNBC drug resistance

(A) TNFAIP2 interacts with IQGAP1. HCC1806 cells with stable TNFAIP2 overexpression were collected from flag-tagged TNFAIP2 cell lysates for immunoprecipitation assays using Flag-M2 beads, and IQGAP1 was detected by WB.

(B) Mapping the domains of TNFAIP2 that interact with IQGAP1. Flag-tagged full-length or truncated TNFAIP2 was transfected into HEK293T cells with no-tagged full-length IQGAP1. Cell lysates were collected for immunoprecipitation using Flag-M2 beads, and IQGAP1 was detected by WB.

(C-E) IQGAP1 knockdown abolished TNFAIP2-confered resistance to EPI and BMN. HCC1937 cells with stable TNFAIP2 overexpression were transfected with IQGAP1 or control siRNA, followed by treatment with EPI (0-1600 nM) and BMN (0-40 μM) for 48 or 72 h, respectively. Cell viability was measured by the SRB assay, and IQGAP1 was detected by WB.

FigureS5. ITGB4 interacts with TNFAIP2 and promotes TNBC drug resistance and DNA damage repair

(A-F) ITGB4 knockdown in HCC1806 and HCC1937 cells significantly decreased cell viability in the presence of DDP (0-40 μM) and AZD (0-40 μM), as measured by the SRB assay. ITGB4 protein expression was detected by WB.

(G) ITGB4 knockdown increased DNA damage of DDP and AZD. HCC1806 and HCC1937 cells with ITGB4 knockdown were treated with 5μM or 7.5 μM DDP for 24h and 15 μM or 20 μM AZD for 24h, respectively. ITGB4, γH2AX, and PARP protein expression was detected by WB.

(H) Mouse masses were collected and weighed at the end of the experiments.

(I) The model of full-length or truncated TNFAIP2.

(J) Mapping the domains of TNFAIP2 that interact with ITGB4. GST-tagged

Full-length ITGB4 was transfected into HEK293T cells with flag-tagged

full-length or truncated TNFAIP2. TNFAIP2 protein was immunoprecipitated

usingFlag-M2 beads, and ITGB4 was detected by WB.

(K) Mapping the domains of TNFAIP2 that interact with ITGB4. No-tagged

full-length ITGB4 was transfected into HEK293T cells with GST-tagged

full-length or truncated TNFAIP2. Cell lysates were collected for the GST

pull-down assay, and ITGB4 was detected by WB.

(L) The model of full-length or truncated ITGB4.

(M) Mapping the domains of ITGB4 that interact with TNFAIP2. Flag-tagged

full-length TNFAIP2 was transfected into HEK293T cells with no-tagged

full-length or truncated ITGB4. Cell lysates were collected for

immunoprecipitation using Flag-M2 beads, and ITGB4 was detected by WB.

FigureS6. ITGB4 promotes TNBC drug resistance via

TNFAIP2/IQGAP1/Rac1

(A-F) ITGB4 promotes TNBC drug resistance through TNFAIP2. TNFAIP2

knockdown abolished ITGB4-induced resistance to DDP and AZD. HCC1806

(A-C) and HCC1937 (D-F) cells with stable ITGB4 overexpression were

transfected with TNFAIP2 or control siRNA, followed by treatment with DDP

(0-30 μM) and AZD (0-30 μM) for 48 or 72h, respectively. Cell viability was

measured by the SRB assay. Protein expression levels were analyzed by WB.

(G-L) ITGB4 promotes TNBC drug resistance through Rac1. Rac1 knockdown

abolished ITGB4-induced resistance to DDP and AZD. HCC1806 (G-I) and

HCC1937 (J-L) cells with stable ITGB4 overexpression were transfected with

Rac1 or control siRNA, followed by treatment with DDP (0-20 μM) and AZD

(0-40 μM) for 48 or 72h, respectively. Cell viability was measured by the SRB

assay. Protein expression levels were analyzed by WB.

(M) ITGB4 promotes DNA damage repair through TNFAIP2. HCC1806 and

HCC1937 cells with stable ITGB4 overexpression were transfected with

TNFAIP2 or control siRNA, followed by treatment with DDP (7.5 μM or 10 μM)

and AZD (20 µM or 30 µM) for 24h. Protein expression levels were analyzed

by WB.

(N) ITGB4 promotes DNA damage repair through Rac1. HCC1806 and

HCC1937 cells with stable ITGB4 overexpression were transfected with Rac1

or control siRNA, followed by treatment with DDP (7.5 µM) and AZD (20 µM or

30 µM) for 24h. Protein expression levels were analyzed by WB.

(O-Q) ITGB4 promotes TNBC drug resistance through TNFAIP2. TNFAIP2

knockdown abolished ITGB4-induced resistance to EPI and BMN. HCC1937

cells with stable ITGB4 overexpression were transfected with TNFAIP2 or

control siRNA, followed by treatment with EPI (0-800 nM) and BMN (0-30 µM)

for 48 or 72 h, respectively. Cell viability was measured by the SRB assay.

Protein expression levels were analyzed by WB.

(R-T) ITGB4 promotes TNBC drug resistance through IQGAP1. HCC1937

cells with stable ITGB4 overexpression were transfected with IQGAP1 or

control siRNA, followed by treatment with EPI (0-800 nM) and BMN (0-40 µM)

for 48 or 72 h, respectively. Cell viability was measured by the SRB assay.

Protein expression levels were analyzed by WB.

(U-W) ITGB4 promotes TNBC drug resistance through Rac1. HCC1937 cells

with stable ITGB4 overexpression were transfected with Rac1 or control siRNA,

followed by treatment with EPI (0-800 nM) and BMN (0-30 µM) for 48 or 72 h,

respectively. Cell viability was measured by the SRB assay. Protein expression

levels were analyzed by WB.

(X) ITGB4 promotes DNA damage repair through TNFAIP2. HCC1937 cells

with stable ITGB4 overexpression were transfected with TNFAIP2 or control

siRNA, followed by treatment with EPI (800 nM) and BMN (10 µM) for 24h.

Protein expression levels were analyzed by WB.

(Y) ITGB4 promotes DNA damage repair through IQGAP1. HCC1937 cells

with stable ITGB4 overexpression were transfected with IQGAP1 or control

siRNA, followed by treatment with EPI (800 nM) and BMN (10 µM) for 24h.

Protein expression levels were analyzed by WB.

(Z) ITGB4 promotes DNA damage repair through Rac1. HCC1937 cells with

stable ITGB4 overexpression were transfected with Rac1 or control siRNA,

followed by treatment with EPI (800 nM) and BMN (10 μM) for 24h. Protein

expression levels were analyzed by WB.

References

1. Bray, F., Ferlay, J., Soerjomataram, I., Siegel, R.L., Torre, L.A., and Jemal, A. (2018). Global
cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36
cancers in 185 countries. CA: a cancer journal for clinicians *68*, 394-424. 10.3322/caac.21492.

2. Zhao, S., Ma, D., Xiao, Y., Li, X.M., Ma, J.L., Zhang, H., Xu, X.L., Lv, H., Jiang, W.H., Yang, W.T., et
al. (2020). Molecular Subtyping of Triple-Negative Breast Cancers by Immunohistochemistry:
Molecular Basis and Clinical Relevance. *25*, e1481-e1491. 10.1634/theoncologist.2019-0982.

3. Sotiriou, C., and Pusztai, L. (2009). Gene-expression signatures in breast cancer. The New
England journal of medicine *360*, 790-800. 10.1056/NEJMra0801289.

4. Bai, X., Ni, J., Beretov, J., Graham, P., and Li, Y. (2021). Triple-negative breast cancer
therapeutic resistance: Where is the Achilles' heel? Cancer letters *497*, 100-111.
10.1016/j.canlet.2020.10.016.

5. Jamdade, V.S., Sethi, N., Mundhe, N.A., Kumar, P., Lahkar, M., and Sinha, N. (2015).
Therapeutic targets of triple-negative breast cancer: a review. British journal of pharmacology
172, 4228-4237. 10.1111/bph.13211.

6. Hill, D.P., Harper, A., Malcolm, J., McAndrews, M.S., Mockus, S.M., Patterson, S.E., Reynolds, T.,
Baker, E.J., Bult, C.J., Chesler, E.J., and Blake, J.A. (2019). Cisplatin-resistant triple-negative
breast cancer subtypes: multiple mechanisms of resistance. BMC cancer *19*, 1039.
10.1186/s12885-019-6278-9.

7. Longley, D.B., and Johnston, P.G. (2005). Molecular mechanisms of drug resistance. The
Journal of pathology *205*, 275-292. 10.1002/path.1706.

8. Rincón, R., Zazo, S., Chamizo, C., Manso, R., González-Alonso, P., Martín-Aparicio, E., Cristóbal,
I., Cañadas, C., Perona, R., Lluch, A., et al. (2016). c-Jun N-Terminal Kinase Inactivation by
Mitogen-Activated Protein Kinase Phosphatase 1 Determines Resistance to Taxanes and
Anthracyclines in Breast Cancer. Molecular cancer therapeutics *15*, 2780-2790.
10.1158/1535-7163.mct-15-0920.

9. Jia, L., Zhou, Z., Liang, H., Wu, J., Shi, P., Li, F., Wang, Z., Wang, C., Chen, W., Zhang, H., et al.
(2016). KLF5 promotes breast cancer proliferation, migration and invasion in part by
upregulating the transcription of TNFAIP2. Oncogene *35*, 2040-2051. 10.1038/onc.2015.263.

10. Chen, L.C., Chen, C.C., Liang, Y., Tsang, N.M., Chang, Y.S., and Hsueh, C. (2011). A novel role
for TNFAIP2: its correlation with invasion and metastasis in nasopharyngeal carcinoma.
Modern pathology : an official journal of the United States and Canadian Academy of
Pathology, Inc *24*, 175-184. 10.1038/modpathol.2010.193.

11. Cheng, Z., Wang, H.Z., Li, X., Wu, Z., Han, Y., Li, Y., Chen, G., Xie, X., Huang, Y., Du, Z., and Zhou,
Y. (2015). MicroRNA-184 inhibits cell proliferation and invasion, and specifically targets
TNFAIP2 in Glioma. Journal of experimental & clinical cancer research : CR *34*, 27.

879   10.1186/s13046-015-0142-9.

880 12. Niwa, N., and Tanaka, N. (2019). TNFAIP2 expression induces epithelial-to-mesenchymal
881   transition and confers platinum resistance in urothelial cancer cells. *99*, 1702-1713.
882   10.1038/s41374-019-0285-y.

883 13. Xie, Y., and Wang, B. (2017). Downregulation of TNFAIP2 suppresses proliferation and
884   metastasis in esophageal squamous cell carcinoma through activation of the Wnt/β-catenin
885   signaling pathway. Oncology reports *37*, 2920-2928. 10.3892/or.2017.5557.

886 14. Jia, L., Shi, Y., Wen, Y., Li, W., Feng, J., and Chen, C. (2018). The roles of TNFAIP2 in cancers and
887   infectious diseases. *22*, 5188-5195. 10.1111/jcmm.13822.

888 15. Didsbury, J., Weber, R.F., Bokoch, G.M., Evans, T., and Snyderman, R. (1989). rac, a novel
889   ras-related family of proteins that are botulinum toxin substrates. The Journal of biological
890   chemistry *264*, 16378-16382.

891 16. Rul, W., Zugasti, O., Roux, P., Peyssonnaux, C., Eychene, A., Franke, T.F., Lenormand, P., Fort, P.,
892   and Hibner, U. (2002). Activation of ERK, controlled by Rac1 and Cdc42 via Akt, is required for
893   anoikis. Annals of the New York Academy of Sciences *973*, 145-148.
894   10.1111/j.1749-6632.2002.tb04624.x.

895 17. Liu, W., Han, J., Shi, S., Dai, Y., and He, J. (2019). TUFT1 promotes metastasis and
896   chemoresistance in triple negative breast cancer through the TUFT1/Rab5/Rac1 pathway.
897   Cancer cell international *19*, 242. 10.1186/s12935-019-0961-4.

898 18. Yan, Y., Greer, P.M., Cao, P.T., Kolb, R.H., and Cowan, K.H. (2012). RAC1 GTPase plays an
899   important role in γ-irradiation induced G2/M checkpoint activation. Breast cancer research :
900   BCR *14*, R60. 10.1186/bcr3164.

901 19. Wu, M., Li, L., Hamaker, M., Small, D., and Duffield, A.S. (2019). FLT3-ITD cooperates with
902   Rac1 to modulate the sensitivity of leukemic cells to chemotherapeutic agents via regulation
903   of DNA repair pathways. Haematologica *104*, 2418-2428. 10.3324/haematol.2018.208843.

904 20. Hu, H., Juvekar, A., Lyssiotis, C.A., Lien, E.C., Albeck, J.G., Oh, D., Varma, G., Hung, Y.P., Ullas, S.,
905   Lauring, J., et al. (2016). Phosphoinositide 3-Kinase Regulates Glycolysis through Mobilization
906   of Aldolase from the Actin Cytoskeleton. Cell *164*, 433-446. 10.1016/j.cell.2015.12.042.

907 21. Li, Q., Qin, T., Bi, Z., Hong, H., Ding, L., Chen, J., Wu, W., Lin, X., Fu, W., Zheng, F., and Yao, Y.
908   (2020). Rac1 activates non-oxidative pentose phosphate pathway to induce chemoresistance
909   of breast cancer. *11*, 1456. 10.1038/s41467-020-15308-7.

910 22. Hervieu, A., and Heuss, S.F. (2020). A PI3K- and GTPase-independent Rac1-mTOR mechanism
911   mediates MET-driven anchorage-independent cell growth but not migration. *13*.
912   10.1126/scisignal.aba8627.

913 23. Li, Y., Li, P., and Wang, N. (2021). Effect of let-7c on the PI3K/Akt/FoxO signaling pathway in
914   hepatocellular carcinoma. Oncology letters *21*, 96. 10.3892/ol.2020.12357.

915 24. Feng, Y.X., Zhao, J.S., Li, J.J., Wang, T., Cheng, S.Q., Yuan, Y., Wang, F., Wang, X.F., and Xie, D.
916   (2010). Liver cancer: EphrinA2 promotes tumorigenicity through Rac1/Akt/NF-kappaB
917   signaling pathway. Hepatology (Baltimore, Md.) *51*, 535-544. 10.1002/hep.23313.

918 25. Higuchi, M., Onishi, K., Kikuchi, C., and Gotoh, Y. (2008). Scaffolding function of PAK in the
919   PDK1-Akt pathway. Nature cell biology *10*, 1356-1364. 10.1038/ncb1795.

920 26. Hu, F., Li, N., Li, Z., Zhang, C., Yue, Y., Liu, Q., Chen, L., Bilan, P.J., and Niu, W. (2018). Electrical
921   pulse stimulation induces GLUT4 translocation in a Rac-Akt-dependent manner in C2C12
922   myotubes. FEBS letters *592*, 644-654. 10.1002/1873-3468.12982.

923   27.   Hamill, K.J., Hopkinson, S.B., DeBiase, P., and Jones, J.C. (2009). BPAG1e maintains
924   keratinocyte polarity through beta4 integrin-mediated modulation of Rac1 and cofilin
925   activities. Molecular biology of the cell *20*, 2954-2962. 10.1091/mbc.e09-01-0051.

926   28.   Wu, J., Zhao, R., Lin, J., and Liu, B. (2018). Integrin β4 reduces DNA damage-induced p53
927   activation in colorectal cancer. Oncology reports *40*, 2183-2192. 10.3892/or.2018.6628.

928   29.   Bierie, B., Pierce, S.E., Kroeger, C., Stover, D.G., Pattabiraman, D.R., Thiru, P., Liu Donaher, J.,
929   Reinhardt, F., Chaffer, C.L., Keckesova, Z., and Weinberg, R.A. (2017). Integrin-β4 identifies
930   cancer stem cell-enriched populations of partially mesenchymal carcinoma cells. Proceedings
931   of the National Academy of Sciences of the United States of America *114*, E2337-e2346.
932   10.1073/pnas.1618298114.

933   30.   Ruan, S., Lin, M., Zhu, Y., Lum, L., Thakur, A., Jin, R., Shao, W., Zhang, Y., Hu, Y., Huang, S., et al.
934   (2020). Integrin β4-Targeted Cancer Immunotherapies Inhibit Tumor Growth and Decrease
935   Metastasis. Cancer research *80*, 771-783. 10.1158/0008-5472.can-19-1145.

936   31.   Kang, S., Jang, Y., Chae, Y.C., and Kim, B.G.

937   32.   Kim, B.G., Gao, M.Q., Choi, Y.P., Kang, S., Park, H.R., Kang, K.S., and Cho, N.H. (2012). Invasive
938   breast cancer induces laminin-332 upregulation and integrin β4 neoexpression in
939   myofibroblasts to confer an anoikis-resistant phenotype during tissue remodeling. Breast
940   cancer research : BCR *14*, R88. 10.1186/bcr3203.

941   33.   Cherfils, J., and Zeghouf, M. (2013). Regulation of small GTPases by GEFs, GAPs, and GDIs.
942   Physiological reviews *93*, 269-309. 10.1152/physrev.00003.2012.

943   34.   Schmidt, V.A. (2012). Watch the GAP: Emerging Roles for IQ Motif-Containing
944   GTPase-Activating Proteins IQGAPs in Hepatocellular Carcinoma. International journal of
945   hepatology *2012*, 958673. 10.1155/2012/958673.

946   35.   Smith, J.M., Hedman, A.C., and Sacks, D.B. (2015). IQGAPs choreograph cellular signaling
947   from the membrane to the nucleus. Trends in cell biology *25*, 171-184.
948   10.1016/j.tcb.2014.12.005.

949   36.   Gorisse, L., Li, Z., Wagner, C.D., Worthylake, D.K., Zappacosta, F., Hedman, A.C., Annan, R.S.,
950   and Sacks, D.B. (2020). Ubiquitination of the scaffold protein IQGAP1 diminishes its
951   interaction with and activation of the Rho GTPase CDC42. *295*, 4822-4835.
952   10.1074/jbc.RA119.011491.

953   37.   Woods, D., and Turchi, J.J. (2013). Chemotherapy induced DNA damage response:
954   convergence of drugs and pathways. Cancer biology & therapy *14*, 379-389.
955   10.4161/cbt.23761.

956   38.   Cheung-Ong, K., Giaever, G., and Nislow, C. (2013). DNA-damaging agents in cancer
957   chemotherapy: serendipity and chemical biology. Chemistry & biology *20*, 648-659.
958   10.1016/j.chembiol.2013.04.007.

959   39.   Kim, C., Gao, R., Sei, E., Brandt, R., Hartman, J., Hatschek, T., Crosetto, N., Foukakis, T., and
960   Navin, N.E. (2018). Chemoresistance Evolution in Triple-Negative Breast Cancer Delineated by
961   Single-Cell Sequencing. Cell *173*, 879-893.e813. 10.1016/j.cell.2018.03.041.

962   40.   Noordermeer, S.M., and van Attikum, H. (2019). PARP Inhibitor Resistance: A Tug-of-War in
963   BRCA-Mutated Cells. Trends in cell biology *29*, 820-834. 10.1016/j.tcb.2019.07.008.

964   41.   Geenen, J.J.J., Linn, S.C., Beijnen, J.H., and Schellens, J.H.M. (2018). PARP Inhibitors in the
965   Treatment of Triple-Negative Breast Cancer. Clinical pharmacokinetics *57*, 427-437.
966   10.1007/s40262-017-0587-4.

42.   Lee, A., and Djamgoz, M.B.A. (2018). Triple negative breast cancer: Emerging therapeutic
modalities and novel combination therapies. Cancer treatment reviews *62*, 110-122.
10.1016/j.ctrv.2017.11.003.

43.   Campbell, P.S., Mavingire, N., Khan, S., Rowland, L.K., Wooten, J.V., Opoku-Agyeman, A.,
Guevara, A., Soto, U., Cavalli, F., Loaiza-Pérez, A.I., et al. (2018). AhR ligand aminoflavone
suppresses α6-integrin-Src-Akt signaling to attenuate tamoxifen resistance in breast cancer
cells. *234*, 108-121. 10.1002/jcp.27013.

44.   Ali, H.R., Dawson, S.J., Blows, F.M., Provenzano, E., Pharoah, P.D., and Caldas, C. (2011).
Cancer stem cell markers in breast cancer: pathological, clinical and prognostic significance.
Breast cancer research : BCR *13*, R118. 10.1186/bcr3061.

45.   Friedland, J.C., Lakins, J.N., Kazanietz, M.G., Chernoff, J., Boettiger, D., and Weaver, V.M.
(2007). alpha6beta4 integrin activates Rac-dependent p21-activated kinase 1 to drive
NF-kappaB-dependent resistance to apoptosis in 3D mammary acini. Journal of cell science
120, 3700-3712. 10.1242/jcs.03484.

46.   Wei, T., and Lambert, P.F. (2021). Role of IQGAP1 in Carcinogenesis. *13*.
10.3390/cancers13163940.

47.   Huang, L., Xu, S., Hu, D., Lu, W., Xie, X., and Cheng, X. (2015). IQGAP1 Is Involved in Enhanced
Aggressive Behavior of Epithelial Ovarian Cancer Stem Cell-Like Cells During Differentiation.
International journal of gynecological cancer : official journal of the International
Gynecological Cancer Society *25*, 559-565. 10.1097/igc.0000000000000394.

48.   Deb, S., Felix, D.A., and Koch, P. (2021). Tnfaip2/exoc3-driven lipid metabolism is essential for
stem cell differentiation and organ homeostasis. *22*, e49328. 10.15252/embr.201949328.

49.   Barutta, F., and Bellini, S. (2022). Protective effect of the tunneling nanotube-TNFAIP2/M-sec
system on podocyte autophagy in diabetic nephropathy. 1-20.
10.1080/15548627.2022.2080382.

50.   Zhou, L., Niu, Z., Wang, Y., Zheng, Y., Zhu, Y., Wang, C., Gao, X., Gao, L., and Zhang, W. (2022).
Senescence as a dictator of patient outcomes and therapeutic efficacies in human gastric
cancer. *8*, 13. 10.1038/s41420-021-00769-6.

51.   Guo, F., and Yuan, Y. (2020). Tumor Necrosis Factor Alpha-Induced Proteins in Malignant
Tumors: Progress and Prospects. *13*, 3303-3318. 10.2147/ott.s241344.

52.   Liu, Y., Ji, X., Kang, N., Zhou, J., Liang, X., Li, J., Han, T., Zhao, C., and Yang, T. (2020). Tumor
necrosis factor α inhibition overcomes immunosuppressive M2b macrophage-induced
bevacizumab resistance in triple-negative breast cancer. Cell death & disease *11*, 993.
10.1038/s41419-020-03161-x.

53.   Chen, C., Sun, X., Ran, Q., Wilkinson, K.D., Murphy, T.J., Simons, J.W., and Dong, J.T. (2005).
Ubiquitin-proteasome degradation of KLF5 transcription factor in cancer and untransformed
epithelial cells. Oncogene *24*, 3319-3327. 10.1038/sj.onc.1208497.

54.   Chen, C., Zhou, Z., Ross, J.S., Zhou, W., and Dong, J.T. (2007). The amplified WWP1 gene is a
potential molecular target in breast cancer. International journal of cancer *121*, 80-87.
10.1002/ijc.22653.

1007

1008

1009

1010

1011

1012

1013

1014

1015

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8