1

**AHL10 phosphorylation determines RRP6L1 chromatin association
and growth suppression during water stress.**

Min May Wong[1], Xin-Jie Huang, Yu-Chiuan Bau and Paul E. Verslues

Institute of Plant and Microbial Biology, Academia Sinica, Taipei 11528, Taiwan

1.Present Address: Department of Genetics, Development and Cell Biology, Iowa State University, Ames, IA, 50011 USA.

Correspondence to: Paul E. Verslues
                          Institute of Plant and Microbial Biology
                          Academia Sinica
                          No. 128 Sec. 2 Academia Rd, Nankang Dist.
                          Taipei, 11529
                          Taiwan
                          Phone: 886-2-2787-1186
                          FAX: 886-2-2782-7954
                          Email: paulv@gate.sinica.edu.tw

Orchid ID: Min May Wong   0000-0001-7725-1615
              Paul E. Verslues   0000-0001-5340-6010

Abstract

Phosphorylation of AHL10, one of the AT-hook family of plant-specific DNA binding proteins, is critical for growth suppression during moderate severity drought (low water potential, ψ_w) stress. To understand how AHL10 phosphorylation determines drought response, we identified putative AHL10 interacting proteins and further characterized interaction with RRP6L1, a protein involved in epigenetic regulation. RRP6L1 and AHL10 mutants, as well as *ahl10-1rrp6l1-2*, had similar phenotype of increased growth maintenance during low ψ_w. Chromatin precipitation demonstrated that RRP6L1 chromatin association increased during low ψ_w stress and was dependent upon AHL10 phosphorylation. Transcriptome analyses showed that AHL10 and RRP6L1 have concordant effects on expression of stress- and development-related genes. Together these results indicate that stress signaling can act via AHL10 phosphorylation to control the chromatin association of the key regulatory protein RRP6L1. AHL10 and RRP6L1 interaction in meristem cells is part of a mechanism to down-regulate growth during low ψ_w stress. Interestingly, loss of AHL13, which is homologous to AHL10 and phosphorylated at similar C-terminal site, blocked the enhanced growth maintenance of *ahl10-1*. Thus, AHL10 and AHL13, despite their close homology, are not redundant but rather have distinct roles, likely related to the formation of AHL hetero-complexes.

Keywords: drought stress, protein phosphorylation, nuclear localization, chromatin-associated proteins, At-Hook, RRP6 Like protein, growth regulation

Summary Statement: Phosphorylation of *Arabidopsis thaliana* AHL10 is important to control growth during drought stress, a function which is distinct from the closely related AHL13. One way that phosphorylation determines AHL10 function is by altering its ability to mediate chromatin recruitment of RRP6L1.

Introduction

Protein phosphorylation and de-phosphorylation is a central feature of signaling pathways that control plant responses to environmental stimuli. Members of the Clade A Type 2C Protein Phosphatases (PP2Cs) have established roles in stress signaling (Cutler et al., 2010; Danquah et al., 2014; Bhaskara et al., 2019). However, there is diversity of physiological function among the Clade A PP2Cs despite their close phylogenetic relatedness. HAI1, in particular, has relatively weak effect on ABA sensitivity of seed germination, the canonical phenotype used to define the ABA signaling function of other Clade A PP2Cs, but has strong effect on growth and osmotic adjustment during low water potential (low ψ_w) stress (Bhaskara et al., 2012; Wong et al., 2019). At the molecular level, it has also been shown that HAI1, but not the Clade A PP2Cs ABI1 or ABI2, could directly inactivate MPK3 and MPK6 (Mine et al., 2017).

Phosphoproteomic analysis of *hai1-2* identified putative HAI1-regulated phosphorylation sites on nuclear-localized proteins (Wong et al., 2019), consistent with the predominant nuclear localization of HAI1 (Antoni et al., 2011; Bhaskara et al., 2012). These included S313 and S314 of AT-Hook Like10 (AHL10). These AHL10 phosphorylation sites had been observed in a number of previous phosphoproteome profiling studies but the functional significance of AHL10 S313 and S314 phosphorylation was not known. Further experiments demonstrated that HAI1 could directly dephosphorylate the AHL10 C-terminal region containing S313 and S314 (Wong et al., 2019). Analysis of *hai1-2* and *35S:HAI1* plants indicated that HAI1 acts to attenuate low ψ_w-induced AHL10 phosphorylation. Further characterization of AHL10 S313 and S314 phosphomimic and null mutations found that these phosphorylation sites are important for controlling growth during low ψ_w and also important in determining whether AHL10 forms nuclear foci or is more dispersed in the nucleoplasm (Wong et al., 2019).

The phosphoproteomic data of Wong et al. (2019) also contained a phosphopeptide from the equivalent site on one of the most closely related AHL10 homologs (AHL13 S376) (Wong et al., 2019). In a separate study, phosphoproteome analysis of chromatin-associated proteins found that AHL13 S376 was phosphorylated by Mitogen Activated protein kinases (MPKs) in response to the pathogen elicitor flg22 (Rayapuram et al., 2021). They found that AHL10 S314 was also a putative MPK6 phosphorylation site but was not responsive to flg22 treatment. Thus, phosphorylation of the C-terminal region of AHL10 and AHL13 may be regulated by MPKs and HAI1 (both directly and by HAI1 suppression of MPK3 and MPK6

activity); however, the two AHLs differ in how their phosphorylation status is regulated in

response to specific environmental stimuli. AHL10 and AHL13 are Clade B AHLs (AHL1 to

AHL14), which are less studied than the Clade A AHLs (AHL15 to AHL29; Zhao et al.,

2013). The C-terminal regions of the Clade B AHLs are highly variable and the MPK-HAI1

regulated phosphorylation sites identified in AHL10 and AHL13 are not conserved among

other Clade B AHLs (Wong et al., 2019).

AHL10 and AHL13 are members of the AT-Hook family of plant-specific DNA

binding proteins. AHLs bind AT-rich DNA sequences via their N-terminal AT-hook domain

(Fujimoto et al., 2004; Zhao et al., 2013). AHLs also contain a Plant and Prokaryote

Conserved (PPC)/DUF296 domain which is involved in self-interaction, or cross-interaction

among AHLs, to form trimers (Fujimoto et al., 2004; Seo and Lee, 2021). In contrast to

many other types of transcriptional regulators, AHLs bind the minor groove of DNA and it

has been proposed that binding of AHL complexes to AT-rich sites can modify DNA

architecture to bring together loops of DNA and place otherwise distal elements in close

proximity to induce the formation of transcriptional regulatory complexes (Zhao et al., 2013;

Franco-Zorrilla et al., 2014).

AHLs are not thought to have transcriptional regulatory activity of their own. Rather,

the highly variable AHL C-terminal domain may recruit other transcriptional regulators or

may mediate interactions that determine AHL sub-nuclear localization. Varied AHL protein

interactions may explain observations that even though different AHLs bind to essentially

identical AT-rich sequences (Franco-Zorrilla et al., 2014), they can either induce or repress

gene expression. For example, ChIP-sequencing of AHL29 found that binding motifs of TCP

and bHLH transcription factors were more highly enriched than AT-rich sequences directly

bound by AHL29 (Favero et al., 2020), consistent with previous data showing that AHL29

interacts with TCP transcription factors (Zhao et al., 2013). Such observations raise the

question of whether AHLs have sufficient sequence specificity of their own to recognize

specific genomic sites or whether it is the AHLs that are recruited by other proteins and then

interact with nearby AT-rich sequences in a more non-specific manner. For, AHL10

specifically, we are aware of only one study examining its cellular function (other than the

phosphoproteomic studies mentioned above). AHL10 was found to interact with Admetos

(ADM), a protein involved in hybrid seed lethality after interploidy hybridization, and with

the SET domain-containing SU(VAR)3–9 homolog SUVH9, a protein involved in deposition

of the heterochromatin mark H3K9me2 (Jiang et al., 2017).

AHLs have also been associated with nuclear matrix attachment regions (Fujimoto et

al., 2004) and can be involved in histone modifications at these sites (Lee and Seo, 2017;

Morisawa et al., 2000; Xu et al., 2013; Xu et al., 2023). Nuclear matrix-associated AHLs are

typically localized in foci within nucleoplasm. Similar foci were also observed for YFP-

tagged AHL10 and for AHL10 homocomplexes visualized using Bi-molecular Fluorescence

Complementation (BiFC). In both cases these nuclear foci were abolished by the AHL10

S314A phosphonull mutation (Wong et al., 2019). Together these results illustrate how

functional diversity among AHLs is likely due to their interaction with a wide range of

proteins via their variable C-terminal domains and association with many types of AT-rich

regions in the genome. Although AHLs are sometimes referred to as transcription factors,

there is still much unclarity about how they function in gene regulation.

To better understand why AHL10 S314 phosphorylation is so critical for its function

to restrict growth during drought stress, we searched for AHL10-interacting proteins. One of

those proteins, RRP6L1, was found to also be a suppressor of growth during low ψ_w stress.

RRP6L1 (also referred to as Atrimmer1) is an RNA binding protein with several proposed

functions related to small RNA production and RNA-directed DNA Methylation (RdDM)

(Zhang et al., 2014, Ye et al., 2016; Wendte et al., 2017). Both AHL10 and RRP6L1 act by

being recruited to specific sites in the genome; DNA binding sites in the case of AHL10 or

sites where specific RNAs are being transcribed in the case of RRP6L1 (Shin and

Chekanova, 2014; Zhang et al., 2014; Wendte et al., 2017). However, in both cases it is

unclear how this occurs as neither protein is thought to have sequence specific binding.

Surprisingly, we found that RRP6L1 chromatin recruitment depended upon AHL10 and

AHL10 S314 phosphorylation. We also made the unexpected discovery that AHL10 and

AHL13 are not functionally redundant, despite their close homology and common regulation

by HAI1 and MPKs.

Methods

Yeast Two-Hybrid Screening

Yeast two-hybrid screening was performed according to the manufacturer's

instructions using the ProQuest[TM] Two-hybrid system (Invitrogen). *AHL10* full length coding

sequence was cloned into the entry vector pENTR-D-TOPO and recombined into the bait

destination vector pDEST32 with GAL4 DNA binding domain (DBD). The yeast two hybrid

cDNA library was constructed from the mRNA of seedlings exposed to -1.2 MPa stress for

96 h according to the CloneMiner[TM] II cDNA library construction kit (Invitrogen) and then

cloned into prey vector pDEST22. The library used and further details of its construction

have been previously described (Chong et al., 2019). Co-transformed yeast (MaV203 strain)

containing both the bait and the prey constructs using the lithium acetate method were spread

on the selection plate SC-Leu-Trp-His with 50 mM 3-AT which was sufficient to suppress

auto-activation. A total of 20 μg cDNA was used and approximately 3.36×10^5 colony

forming units were screened. Large colonies grown on the selection plates were picked and

re-streaked on the YPAD plates after 4-5 days at 30°C and some smaller colonies were

picked after 8 days. Selected colonies were grown on the YPAD plates overnight at 30°C and

interaction were confirmed by β-galactosidase filter lift assay. Confirmed colonies were then

cultured, DNA isolated by yeast plasmid miniprep (Zymoprep) and sequenced. Gene

sequences of AHL10 and truncated RRP6L1 used for yeast two-hybrid assays were cloned

into the pENTR-D-TOPO and recombined into pDEST32 and pDEST22 for β-galactosidase

filter lift assay. Primers used to clone the different truncated version of RRP6L1are listed in

Supplemental Table S4.

Plant materials, growth conditions and physiological assays

The T-DNA insertion line of AHL10, *ahl10-1* (SALK_073165C) was previously

described (Wong et al, 2019). T-DNA insertion lines of RRP6L1, *rrp6l1-1* (SALK_004432)

and *rrp6l1-2* (SALK_027601) and the *ahl13-1* (SALK_014014) were obtained from the

Arabidopsis Biological Resource Center and the homozygous plants confirmed using primers

designed using the Signal web resource (http://signal.salk.edu/). To generate transgenic plants

expressing *AHL10* under control of its native promoter, the genomic fragment containing the

AHL10 gene body (5' UTR, introns and exons up to the stop codon) and 2.1 kb of promoter

upstream of the transcriptional start site were cloned into pENTR-D-TOPO and recombined

into pGWB540 by Gateway recombination before transformation into *ahl10-1* or *ahl10-*

1rrp6l1-2 (all in the Col-0 background) using *Agrobacterium tumefacians* strain GV3101 and

the floral dip method. To express phosphomimic and phosphonull AHL10 under control of its

native promoter, the entry clone in the pENTR-D-TOPO vector was used as template plasmid

for PCR using site-directed mutagenic primers followed by Dpn1 digestion and cloning as

previously described (Bhaskara et al., 2017). All constructs were verified by sequencing.

Primers used for genotyping, cloning and site-directed mutagenesis are given in

Supplemental Table S4.

Low ψ_w treatment on PEG-infused agar plates and soil drying experiments were

performed as described previously (Verslues et al., 2006; Wong et al., 2019). For plate

experiments to assay plant growth, 5-day-old seedlings were transferred from control media

to the indicated treatment and growth measured 5 (control) or 10 (stress) days after transfer.

For growth assay involving transgenic lines, three independent lines were tested in at least

three independent experiments with two or three plates containing the test genotype and wild

type used as technical repeats per experiment. Similar replication was performed for assays

involving T-DNA mutants. Growth of the test genotype is reported relative to wild type

growing in the same plate.

For soil drying experiments, a standard potting mix was combined with 25% Turface

(Turface MVP, Profile Products LLC, USA). Four genotypes were planted in sectors (two

plants per sector) of 8 cm x 8 cm x 10 cm (LxWxH) plastic pots to ensure that the different

genotypes were exposed to the same extent of soil drying. Seeds placed into pots with fully

watered soil mix, stratified and then transferred to a short-day chamber (8 h light period, 23

C, light intensity of 100-120 μmol m^{-2} sec^{-1}) and Hyponex nutrient solution (1 g liter^{-1})

supplied once per week along with regular watering to maintain full soil hydration. The

position of individual pots within the chamber was rotated every two or three days. On day

19 after planting, pots were watered to saturation, allowed to drain and weighed. Water was

then withheld for 12 days (leading to 50-60 percent reduction in pot weight), each pot re-

watered to 75 percent of the initial pot weight and then allowed to dry another 8-10 day until

pot weight again reached 50-60 percent of the starting weight. At the end of the experiment,

representative rosettes were photographed and the rest weighed (fresh weight) and then

dehydrated in an oven to obtain the dry weight. Growth of the test genotype is reported

relative to wild type growing in the same pot. Each genotype was tested in at least three

independent experiments with two or three pots containing the test genotypes and wild type

used as technical repeats per experiment.

In-vitro Pull-down assay

GST-tagged protein expression vector, pGEX-4T-1, was used to clone the RRP6L1

N-terminal 300 and C-terminal 337. AHL10 non-mutated wild type and phosphomimic of

S313D and S314D was cloned into the pET300NT destination vector (N-terminal HIS tag).

For GST-tagged RRP6L1 N- and C-terminal fusion proteins, the expression plasmids were

transformed into BL21(DE3) pGro7 *E. coli* competent cells and induced by 0.5 mM

isopropyl β-D-thiogalactoside (IPTG) in the presence of 0.5 g/L L-arabinose, and incubated

for an additional 16-20 hours at 20°C. For His-tagged AHL10 (N.M., S313D and S314D),

the expression plasmids were transformed into Rosetta 2(DE3) pLysS *E. coli* competent cells

and induced by 0.3 mM IPTG, and incubated for an additional 22-24 hours at 16°C. The cells

were harvested by centrifugation (6,000xg for 30 min at 4 □C) snap-frozen, and stored in -

80°C until use.

The bacterial pellets harboring GST-tagged RRP6L1 protein were resuspend in

prechilled lysis buffer (20 mM HEPES-KOH at pH7.50, 200 mM NaCl, 0.1 mM $MgCl_2$, 5%

[v/v] glycerol, 5-mM 2-mercaptoethanol, 1x Complete, EDTA-free Protease Inhibitor

Cocktail [Roche]), 1% [v/v] IGEPAL CA-630, 5 mM 2-mercaptoethanol), followed by

sonication on ice using a 1/4-inch microtip controlled by the S-4000 ultrasonic processor

(Misonix Sonicators). The cel lysates were clarified by high-speed centrifugation at 50,000g

for 60 min at 4°C (Avanti J-26 XP centrifuge with JA-25.50 rotor; Beckman Coulter),

followed by a 0.45-mm filtration (Millex-HP, 33 mm, polyethersulfone; Merck Millipore)

before affinity chromatography. The pellets harboring His-tagged AHL10 protein were

resuspend in prechilled lysis buffer (50 mM potassium phosphate at pH7.40, 200 mM NaCl,

10 mM Imidazole-Cl, 10% [v/v] glycerol, 5-mM 2-mercaptoethanol, 1x Complete, EDTA-

free Protease Inhibitor Cocktail [Roche]), 5 mM 2-mercaptoethanol), followed by the same

lysis and clarification procedures as RRP6L1.

For RRP6L1 N- and C-terminal fusion proteins (and free GST), GST affinity

purification was performed using Glutathione Sepharose 4FF resin and proteins eluted using

lysis buffer supplied additional freshly prepared 10 mM reduced glutathione-NaOH. His-

tagged AHL10 proteins were purified using Clontech His60 Ni Superflow resin and eluted

using 400 mM imidazole-Cl. The AHL10 proteins were rapidly diluted with equal volume of

ion exchange buffer (10 mM potassium phosphate at pH7.40, 50 mM NaCl, 0.1 mM $MgCl_2$,

10% [v/v] glycerol, 5-mM 2-mercaptoethanol) before being applied to a HiTrap Heparin HP

(GE Healthcare), and eluted by a linear NaCl gradient up to 1.0 M. Purified proteins were

dialyzed against pull-down buffer (25 mM HEPES [pH 7.5], 150 mM NaCl, 0.1 mM MgCl2,

10% [v/v] glycerol, 0.1% [v/v] IGEPAL CA-630, 5 mM 2-mercaptoethanol) and protein

concentration measured using Bradford method (Bio-Rad Protein Assay Dye Reagent

Concentrate; Bio-Rad Laboratories).

For pull down assays, 20 µl of glutathione magnetic beads (Pierce catalog no. 88821)

were incubated with GST-tagged protein (30 µg free GST or 70 µg RRP6L1 N-300/C337)

and 30 µg of AHL10 (10% of the reaction removed for input detection) for 120 min at 4 C

with gentle rotation. Beads were then washed 5x with 1 ml of pull-down buffer and bound

proteins eluted by adding 40 µl pull-down buffer and 10 µl of 5x SDS-PAGE Laemmli

loading buffer to the washed beads. Proteins were then separated by SDS-PAGE and blotted

onto PVDF membranes. Membranes were blocked using 5% nonfat milk and probed with

anti-poly-His primary antibody (cat. no. H1029; Sigma-Aldrich) at 1:3,000 dilution or anti-

GST primary antibody (cat. no. SAB4200692; Sigma-Aldrich) at 1:2,000 dilution, followed

by secondary antibody (rabbit anti-mouse IgG H&L [HRP], AbCam ab6728) at 1:10,000

dilution). Immunoreactive bands were visualized by the enhanced chemiluminescence

method (Pierce ECL Western Blotting Substrate, Thermo Fisher Scientific) and detected

using Thermo Scientific CL-XPosure Film). For band quantification, images of the film (48-

bit RGB tiff, 720dpi) were converted to 16-bit gray scale images. ImageJ was used to subtract

background, invert the images and then measure intensity of equal sized regions of interest

for each band. The ratio of pulldown band versus input band was calculated for each protein

interaction and the data for unmutated AHL10 compared to the AHLS313D and

AHL10S314D data. Three independent sets of pulldown reactions were conducted and used

for band intensity quantification.

270   Purification of His-tagged RRP6L1 C-terminal fragment and Generation of RRP6L1

271   antisera

The portion of the RRP6L1 cDNA encoding the C-terminal portion of the protein

(amino acid 301-638) was cloned into pENTR-D-TOPO and LR reaction transferred to the

pET300/NT-DEST vector to generate recombinant protein with an N-terminal fusion to

6×HIS tag. Recombination protein was produced in Rosetta 2(DE3) pLysS *E. coli* competent

cells by addition of 1mM IPTG when cultures reached an optical density of 0.4 followed by

incubation for 4 h at 37°C. Cells were harvested and resuspended in 50 mL lysis buffer (50

mM HEPES pH 7.4—adjusted pH with NaOH, 150 mM NaCl, 0.5% Tween-20, 1X protease

inhibitor, 1 mM DTT, 15% glycerol) and lysed using a cell disruptor (Constant Systems).

Lysate was filtered and applied to a 1 mL Mini Profinity IMAC cartridge using BioLogic

low-pressure chromatography system (Bio-Rad). An imidazole gradient was used to elute the

protein in elution buffer (300 mM imidazole; 50 mM HEPES pH 7.4—pH adjusted with

NaOH, 300 mM NaCl, 20% glycerol). Protein purify was checked by SDS-PAGE and

commassie blue staining. For final purification, 2 mg purified protein was separated by SDS-

PAGE and band corresponding to the expected molecular weight of the RRP6L1 301-638

fragment (40 kDa) excised and used for antisera generation in rabbit (performed by LTK

Biolaboratories, Taiwan). Antibody titer and specificity was checked by immunoblot using

protein extracts from the wild type and *rrp6l1* mutants probed with affinity purified RRP6L1

antisera (immunoblot procedures and optimized antisera dilution are described below).

Ratiometric Bi-molecular Fluorescence Complementation (rBiFC) and subcellular
localization

Ratiometric BiFC (rBiFC) was performed using the vector system described by

(Grefen and Blatt, 2012) whereby a constitutively expressed RFP is present on the BiFc

plasmid. AHL10 wild type, phosphomimic, or phosphonull cDNA clones (described in Wong

et al., 2019) were inserted into pDONR221 P1-P4 (Invitrogen) and the RRP6L1 cDNA was

amplified from total mRNA and cloned into pDONR221 P3-P2 (Invitrogen). The AHL27

and AHL29 cDNAs were amplified from plasmid clones generously provided by David

Faverro (Riken) and cloned into pDONR221 P1-P4. Primer sequences used for cloning are

given in Supplemental Table S4. These donor vectors were used for two site Gateway

recombination into the destination vector pBiFCt2in1-NN and transformed into the

Agrobacterium GV3101. Five-day-old seedlings of an Arabidopsis line with Dexamethasone

(Dex) inducible AvrPto expression (Tsuda et al., 2012) were used for transient expression,

followed by confocal microscopy (using a Zeiss LSM 510 META or Zeiss LSM 880), and

quantification of YFP and RFP signal intensities performed as previously described (Wong et

al., 2019). Subcellular localization of *AHL10$_{pro}$:AHL10-EYFP/ahl10-1* and its phosphomimic

and phosphonull variants were visualized using a Zeiss LSM 880 with airyscan.

Collection of meristem-enriched tissue samples and immunoblotting

Seven-day-old seedlings were transferred to control and stress treatments (-0.7 MPa

PEG-infused plates) for 4 days. Samples of the shoot meristem and surrounding tissue were

collected using a 7mm diameter round metal punch to excise all tissue surrounding the

meristem regions. Root tip samples were collected by excising the apical 3-5 mm of the root

tip with a razor blade. Meristem-enriched samples or whole seedlings were ground in liquid

nitrogen and homogenized in cold protein extraction buffer (50 mM Tris HCl pH 7.5, 10%

glycerol, 1 mM DTT, 1% IGEPAL, 1× Roche EDTA-free protease inhibitor and 1× Roche

phosphatase inhibitor). The homogenate was centrifuged and the protein concentration of the

supernatant was measured using a Pierce 660 nm Protein Assay Reagent (Thermo Scientific).

Thirty-five microgram of protein was loaded in each lane. Two SDS-PAGE gels were run

with the same aliquots of all the tissues samples, one was used for anti-GFP (Sigma-Aldrich

Cat# 1181446001) and the other used for blotting with anti-RRP6L1 polyclonal antisera

described above (antibody dilutions and other blotting conditions are described below). The

anti-RRP6L1 blots were stripped using mild stripping buffer (1.5 g glycine, 0.1g SDS, 1 mL

Tween-20 in 100 mL solution, pH 2.2) and re-probed with anti-HSC70 (Enzo # ADI-SPA-

818-D) at 1:5000 dilution, as a loading control.

Isolation of total nuclear, nucleoplasmic and chromatin-associated protein

Seven-day-old seedlings were transferred to -0.7MPa PEG-infused plates for 4 days.

Seedlings were harvested and cross-linked with 1% formaldehyde solution in $1\times$ PBS by

vacuum infiltration (3 times, 10 minutes total infiltration time with total exposure time to

formaldehyde not exceeding 15 minute). Cross-linking was stopped by adding 0.125 M

glycine followed by vacuum infiltration (2 times, 90 seconds each time). The samples were

rinsed twice with cold PBS, dried with a paper tower and 600 mg of tissue frozen in liquid

nitrogen. The same cross-linking procedure was used for assays of chromatin associated

protein as well as ChIP-quantitative PCR assays (described below).

Isolation of total nuclear, nucleoplasmic and chromatin-bound associated was

conducted following the procedure of Wu et al., (2016) with some modifications. Samples

were ground into powder using a mortar and pestle and suspended in 5 ml of Honda Buffer

(20 mM HEPES-KOH pH 7.4, 0.44 M sucrose, 1.25% ficoll, 2.5% Dextran T40, 10 mM

MgCl2, 0.1% Triton X-100, 5 mM DTT, 0.5 mM PMSF, $1\times$ protease inhibitors [Roche]),

filtered through two layers of Miracloth and centrifuged at 2000 x g for 15 min at 4 °C. The

supernatant was removed, and the nuclear pellets were washed twice with 1ml of Honda

buffer, resuspended in 600 μl nuclei lysis buffer (10 mM Tris-HCl pH 7.5, 2 mM EDTA, $1\times$

proteinase inhibitor, $1\times$ phosphatase inhibitor and 1% SDS). 600 μl aliquots of lysed nuclei

solution were aliquoted into two low binding microcentrifuge tubes. One tube was kept on

ice (this is the total nuclear extract), and the other tube vortexed for 2 minutes followed by

centrifugation at 14,000g for 10 mins at 4°C and collection of the supernatant (this is the

soluble fraction of nucleoplasm protein). The remaining pellet was re-suspended in 300 μl of

lysis buffer containing 1% SDS and sonicated using a bioruptor (Diagenode) at 4°C (15 sec

ON/OFF for 5-min and repeated for 5 cycles). The sonicated fragment was centrifuged at

14,000g for 10 mins at 4°C and the supernatant collected (this is the chromatin-associated

protein). The pellet was resuspended and collected (pellet after sonication fraction). From the

collected fractions of total nuclear extract, soluble nucleoplasm, chromatin-associated and

pellet after sonication (each in 300 µl total volume), 50 µl was mixed with 6× loading dye,

heated at 95°C for 10 mins, loaded on 10% SDS-PAGE, and transferred to PVDF membrane

followed by blocking in 5% w/v non-fat milk powder in 1% TBST for overnight. The anti-

GFP antisera (Sigma-Aldrich Cat# 1181446001) was used at 1:2000 dilution followed by

1:10000 secondary anti-mouse HRP. The membrane was stripped using mild stripping buffer

(1.5 g glycine, 0.1g SDS, 1 ml Tween-20 in 100 ml solution, pH to 2.2) and re-probed with

anti-RRP6L1 antibody (1:8000 dilution) followed by 1:10000 anti-rabbit HRP. Separate 50

µl (same aliquot from the total 300 µl) were separated on 13% SDS-PAGE gel for

immunoblot using anti-Histone 3 (Sigma #06-755) at 1:3000 dilution with 1:10000 anti-

rabbit secondary antibody. Quantification of band intensity for the 74 kd RRP6L1-specific

band as well as AHL10-YFP band was performed as described above for the GST pull down

experiments.

ChIP-quantitative PCR assays

ChIP-quantitative PCR (qPCR) assays were performed as previously described

(Yamaguchi et al., 2014). Plant samples were collected, harvested and cross-linked as

described above for the chromatin-associated protein assays. 300 mg tissue from whole

seedlings was used with 30 µl of GFP magnetic trap beads (Chromotek). The plants were

ground to powder in liquid nitrogen and lysed in 2.5 ml cold nuclei extraction buffer. The

lysate was filtered through 2-layers of miracloth (Millipore, Cat. 475855) set up in a 50 ml

centrifuge tube, centrifuged at low speed to collect the pellet (nuclei) and the pellet

resuspended with nuclei lysis buffer. Sonication was performed in an ice bath using a

Bioruptor (Diagenode) set to high power and operated using 15sec on/off cycles for 5-min

and repeated for 3 cycles. The procedures for nuclei isolation, shearing of chromatin,

immunoprecipitation, washing, reverse crosslinking, elution to get the purified DNA using

QIAquick spin columns (Qiagen, Cat: 28106) and ChIP-qPCR DNA reaction composition

were as described in Yamaguchi et al. (2014). A 5-fold dilution series of input was amplified

for each set of primers and the data used to calculate the percent input. Two or three

independent experiments were performed for each ChIP target site analyzed, as described in

the figure legends. Primers used for ChIP-qPCR assays are shown Supplemental Dataset S4.

RNA Sequencing and Quantitative Reverse Transcriptase-PCR analysis of gene

expression

The RNA sequencing of *rrp6l1-2* was conducted in the same set of experiments as
previously reported data of wild type and *ahl10-1* and followed the previously described
procedures for sample processing, and data analysis (Wong et al. 2019). Seven-day-old
seedlings were transferred to fresh control plates or PEG-infused agar plates (-0.7 MPa) and
whole seedlings collected four days after transfer (11 days old). Significant differences in
gene expression were determined using DEseq2. Heat maps with hierarchical clustering
(Euclidian distance clustering) were prepared using Morpheus (Broad Institute, Morpheus
(broadinstitute.org)).

For QPCR, the sample collection procedure was that same as described above for
RNAseq analysis. Total RNA was extracted from control or -0.7 MPa stress-treated
seedlings using RNeasy Plant Mini Kit (Qiagen). 1 µg RNA was reverse transcribed using
SuperScript III (Invitrogen). Real-time quantitative PCR assays were prepared using KAPA
SYBR FAST qPCR kit (Kapa Biosystems) and run on an Applied Biosystems QuantStudio
12K QPCR instrument. The comparative $\Delta\Delta C_T$ method was used to determine relative gene
expression using *EFL1α* as a reference gene (Nicot et al., 2005). Note that for some genes
(*STM, WES1, DFL1, MYB117* and *PRR*) the QPCR data *ahl10-1* was previously reported in
Wong et al. (2019) and is shown again here for comparison to *rrp6l1-2*.

Statistical Analysis

Graph preparation and statistical analyses were performed using GraphPad Prism 9.
Data were analyzed by ANOVA, T-test or one sample T-test as indicted in figure legends
with corrected $P \geq 0.05$ used as a cutoff for significant differences.

Results

AHL10-interacting proteins, including RRP6L1, identified by yeast two-hybrid
screening.

To identify AHL10 interacting proteins, we used full length AHL10 as bait to screen a
yeast two-hybrid library prepared using cDNA isolated from low ψ_w-treated seedlings (the
library has been described in Chong et al., 2019). The screening identified several clones of
AHL1, AHL3, AHL10 and AHL13 that showed strong interaction with AHL10
(Supplemental Fig. S1A). This indicated that these four Clade B AHLs can form complexes
with each other and is consistent with the results of Jiang et al. (2017) where TAP tagging
and immunoprecipitation of ADM also found these four AHLs. Among other nuclear-

localized proteins identified in the screen, AHL10 strongly interacted with RRP6L1 and the

cell-cycle regulator CDC20.2 and had weaker putative interaction with RNA-binding

(RBP45A), DNA-binding (WRKY1, Histone 3.1) and chromatin associated (SGO2) proteins.

We focused further attention on RRP6L1 because of its strong interaction in the yeast-two

hybrid assays (Fig. 1A). The interacting clone detected in the yeast two-hybrid screen

contained the C-terminal 76 amino acids of RRP6L1. Retransformation assays confirmed

AHL10 interaction of this clone as well as a longer clone containing the C-terminal 192

amino acids. However, neither the C-terminal 338 amino acids nor full length RRP6L1gave

a positive interaction signal with AHL10 (Fig 1A). It is possible that the longer RRP6L1

clones interfered with reporter activation in the yeast assays.

AHL10-RRP6L1 interaction

Ratiometric Bi-molecular fluorescence complementation (rBiFC) assays found that

AHL10-RRP6L1 generated a strong fluorescence signal in seedlings exposed to low ψ_w stress

but not in unstressed seedlings (Fig. 1B and C). As a control for the rBiFC interaction assays,

we also tested AHL10 interaction with AHL27 and AHL29. In the low ψ_w treatment, AHL27

and AHL29 had lower rBiFC signal compared to the AHL10-RRP6L1 interaction (Fig. 1B).

This lower rBiFC signal may represent non-specific background, perhaps due to high level of

transient expression of the proteins, or may represent a limited interaction of RRP6L1 with

other AHLs. We also found that AHL10 had a relatively strong rBiFC signal with WRKY1

in both unstressed and low ψ_w-treated plants (Supplemental Fig. S1B), indicating that AHL10

expression was not the cause of the lower AHL10-RRP6L1 rBiFC signal in unstressed plants.

Together these data demonstrated that RRP6L1 interacted with AHL10 in the nucleus and

that this interaction may be stimulated by low ψ_w stress.

AHL10-RRP6L1 interaction was also detected using *in vitro* pull-down assays.

RRP6L1 was divided into the N-terminal 300 amino acids (N300) and C-terminal 337 amino

acids (C337) as full length RRP6L1 could not be expressed in *E. coli*, consistent with

previous report (Zhang et al., 2014). The N300 fragment of RRP6L1 contains the 3'-5'

exonuclease domain while the C337 fragment contains a Helicase and RNase D C-terminal

(HRDC) nucleic acid-binding domain. Interestingly, both halves of RRP6L1 interacted with

AHL10; however, the pull-down band intensity was significantly greater for the C337

fragment of RRP6L1 than for N300 (Supplemental Fig. S2A and C). Phosphomimic

mutations at the S313 and S314 phosphorylation sites previously found to be critical for

AHL10 function (Wong et al., 2019) increased the pull-down band intensity of the N300

fragment but had no significant effect on interaction of C337 (Supplemental Fig. S2B and C).

These data confirmed AHL10-RRP6L1 interaction and indicated that AHL10

phosphorylation at S313 or S314 is unlikely to block interaction with RRP6L1; however, we

cannot rule out that phosphorylation affects the affinity of AHL10-RRP6L1 interaction or

affects how AHL10 interacts with the different domains of RRP6L1.

AHL10 and RRP6L1 accumulate in meristematic tissue

We generated transgenic lines where a genomic fragment containing the AHL10

promoter and gene body (5' UTR, exons, and introns) up to the stop codon was fused with

YFP (*AHL10$_{genomic}$:YFP*) and used to complement *ahl10-1* (Supplemental Fig. S3). Imaging

of these transgenic lines found that AHL10-YFP accumulated in the root quiescent center,

developing stele and columella cells under both control and low ψ_w stress conditions (Fig.

1D). AHL10-YFP also accumulated to high levels in the shoot meristem region as well as in

the meristem region of lateral roots and sites of lateral root emergence (Supplemental Fig.

S4). The same AHL10 native promoter construct was produced for phosphonull AHL10

(AHL10^{S313A}, AHL10^{S314A}) and phosphomimic AHL10 (AHL10^{S313D}, AHL10^{S314D}) and

transformed into *ahl10-1*. Consistent with previous results AHL10^{S313D} and AHL10^{S314D}

complemented *ahl10-1* while AHL10^{S314A} did not complement. AHL10^{S313A} was hyper-

functional and suppressed growth below the wild type level during low ψ_w stress

(Supplemental Fig. S3). We hypothesize that preventing phosphorylation at S313 caused

S314 to become hyperphosphorylated (Wong et al 2019). Hyperphosphorylation of S314

may have larger growth suppression effect than the S314 phosphomimic construct as

aspartate substation, while useful for most functional analyses, is not a perfect mimic of

phosphorylation (it introduces a single negative charge while phosphorylation introduces a -2

charge). AHL10^{S314A} was diffusely localized in the nucleoplasm and lacked the nuclear foci

seen for wild type AHL10, AHL10^{S313A} and phosphomimic AHL10 (Supplemental Fig. S5).

Note that the nucleoplasm localization of AHL10-YFP matches previously reported

localization of RRP6L1 (Lange et al., 2008; Zhang et al., 2014).

To study RRP6L1 protein accumulation, and compare it to that of AHL10, we

generated antisera recognizing the N-terminal domain of RRP6L1. Immunoblotting of wild

type and two previously described RRP6L1 knockout mutants, *rrp6l1-1* (Lange et al., 2008;

Zhang et al., 2014; Ye et al., 2016; Wendte et al., 2017) and *rrp6l1-2* (Hsu et al., 2014),

found that the antisera recognized a 74 kD band which corresponded to the expected

molecular weight of RRP6L1 and which was absent in both of the *rrp6l1* alleles (Fig. 1E). It

was also consistently observed that RRP6L1 protein level was moderately increased by low

ψ_w stress. Samples of shoot and root meristem (and surrounding tissue) had substantially

higher levels of RRP6L1 and AHL10-YFP than whole seedling samples (Fig. 1F). These

data indicated that AHL10 and RRP6L1 proteins accumulate in the same parts of the plant,

consistent with their physical interaction. Immunoblots also showed that RRP6L1 protein

level did not substantially differ between wild type and *ahl10-1*. Likewise, AHL10 protein

level was not substantially affected by *rrp6l1-2* (Fig. 1E and F). Thus, the purpose of the

AHL10-RRP6L1 interaction was not related to one protein affecting the abundance of the

other protein.

RRP6L1 and AHL10, but not AHL13, restrict growth at low ψ_w.

We subjected both *rrp6l1* alleles to a moderate severity low ψ_w (-0.7 MPa) that

decreases, but does not completely inhibit, growth of wild type (Wong et al., 2019). Both

rrp6l1 mutants maintained a higher level of growth than wild type during this moderate

severity low ψ_w stress while having minimal or no effect on growth in the unstressed control,

similar to *ahl10-1* (Fig. 2A, Supplemental Fig. 6A, wild type data used for normalization can

be found in Supplemental Fig. S6B). An *ahl10-1rrp6l1-2* double mutant had similar

increased growth maintenance as either single mutant. *rrp6l1-2* also had enhanced growth

maintenance during controlled soil drying assays where plants were exposed to an extended

period of partial soil drying that slowed rosette growth but did not lead to wilting (Fig 2B, C).

Note that the *ahl10-1* data from these soil drying experiments has been previously reported

(Wong et al., 2019) and is shown again here to allow better comparison between all

genotypes.

AHL13 is closely related to AHL10, including conservation of the C-terminal

phosphorylation site (Wong et al., 2019; Rayapuram et al., 2021), and likely forms

heteromeric complexes with AHL10 (Supplemental Fig. S1). However, *ahl13-1* did not

differ from wild type in growth maintenance at low ψ_w (Fig 2A). Interestingly though, loss

of AHL13 blocked the increased growth of *ahl10-1* as *ahl10-1ahl13-1* grew similarly to wild

type in both PEG-plate (Fig. 2A) and soil drying assays (Fig 2B). Transformation of *ahl10-*

1ahl13-1 with *35S:AHL13* restored the *ahl10-1* phenotype (Supplemental Fig. 6C), validating

that AHL13 is required for the increased growth maintenance of *ahl10-1*. These data indicate

that AHL10 and AHL13 do not have redundant function. Rather AHL13 acts genetically

downstream of AHL10, consistent with the formation AHL10-AHL13 heterocomplexes.

AHL10 S314 phosphorylation is required for RRP6L1 chromatin association.

AHLs and RRP6L1 are both hypothesized to require additional interacting factors to

correctly target and associate with specific genomic sites or transcripts (Zhao et al., 2013;

Wendte et al., 2017; Favero et al., 2020). Thus, we used chromatin precipitation to test

whether AHL10 or RRP6L1 affected one another's partitioning to the nucleoplasm or

chromatin-associated fractions of nuclear extracts and whether this partitioning was affected

by low ψ_w stress. In these assays, centrifugation of nuclear extracts was used to separate

soluble nuclear proteins (Soluble Fraction, labeled as "S.F." in Fig. 3) from chromatin-

associated proteins (labeled as "C.A." in Fig. 3). Sonication then allowed the chromatin-

associated proteins to be solubilized and separated from the remaining insoluble material

(pellet after sonication, labeled as "P.S." in Fig. 3).

RRP6L1 presence in the chromatin-associated fraction was strongly increased by low

ψ_w and was dependent upon AHL10 S314 phosphorylation. RRP6L1 chromatin association

was reduced to background level in *ahl10-1* (compare Fig. 2B and C to Fig. 2E where a line

expressing GFP alone was used as a negative control). In the low ψ_w treatment, RRP6L1

presence in the chromatin-associated fraction was also strongly reduced by the AHL10^{S314A}

phosphonull mutation (Fig 2B and D). This reduced RRP6L1 chromatin association in plants

expressing AHL10^{S314A} correlates with the altered sub-nuclear localization of AHL10^{S314A}

(Supplemental Fig. S5) and with its inability to complement *ahl10-1* (Supplemental Fig. S3).

Interestingly, the opposite effect was observed for AHL10^{S313A} where the portion of RRP6L1

in the chromatin-associated fraction was high and similar to the high level seen in the AHL10

phosphomimic mutants (AHL10^{S313D} and AHL10^{S314D}). The high level of chromatin-

associated RRP6L1 in plants expressing AHL10^{S313A} correlates with the strong growth

suppression activity of AHL10^{S313A} (Supplemental Fig. S3; Wong et al., 2019) and is

consistent with the idea that blocking AHL10 S313 phosphorylation directs more

phosphorylation to the critical S314 site (Wong et al., 2019). These effects of AHL10 on

RRP6L1 chromatin association were more apparent in the low ψ_w stress treatment, consistent

with the low ψ_w-induced increase of AHL10-RRP6L1 association seen in rBiFC assays (Fig.

1B).

Conversely, the proportion of wild type AHL10 (labeled as N.M., for Non-Mutated)

in the chromatin-associated fraction was not affected by *rrp6l1-2* but was significantly

increased by low ψ_w stress (Fig. 3A and C). We also found that the AHL10^{S313A},

AHL10^{S313D}, and AHL10^{S314D} phosphomimic and null mutations did not affect AHL10

presence in the chromatin-associated fraction (Fig. 3A and D), consistent with

phosphorylation of the C-terminal domain having no direct effect on DNA binding activity of

the N-terminal AT-hook domain. AHL10^{S314A} may have been somewhat less present in the

chromatin-associated fraction at low ψ_w, however, the effect was marginally non-significant

(P = 0.06) and thus should be interpreted with caution.

The remaining pellet after chromatin-associated proteins were released by sonication

did not contain detectable AHL10 or RRP6L1, suggesting that neither protein was part of

other, non-chromatin-associated, insoluble protein aggregates in the nucleus. Histone 3 (H3)

was used as a control chromatin-associated protein (Fig. 3C). Note that our assays indicated

that low ψ_w stress led to an increased level of H3 in the soluble nuclear fraction but had little

effect on the chromatin associated fraction. This indicated that the increased chromatin

association of AHL10 and RRP6L1 in the stress treatment was a specific effect and not a

general effect of the low ψ_w treatment on all nuclear proteins.

RRP6L1 and AHL10 affect the expression of an overlapping set of low ψ_w-responsive

genes

We conducted transcriptome analysis of *rrp6l1-2* under unstressed conditions and

after four-day acclimation to moderate severity low ψ_w stress (-0.7 MPa). This relatively

long term low ψ_w treatment was used to find genes that undergo sustained induction or

repression and thus are more likely to be associated with the differences in growth between

rrp6l1 versus wild type at low ψ_w. The *rrp6l1-2* transcriptome data were collected in the

same set of experiments as *ahl10-1* and wild type data previously reported (Wong et al.,

2019). In comparison of unstressed wild type to unstressed *rrp6l1-2*, we found 147 genes

having increased expression and 147 genes having decreased expression, including *RRP6L1*

itself (log2 fold-change ≥ 0.5, adjusted P ≤ 0.05; Supplemental Table S1). Genes up- or

down-regulated in *rrp6l1-2* had the same direction of change in *ahl10-1* in nearly all cases

(Supplemental Fig. S7A), albeit that the *ahl10-1* effect was weaker in many cases. This was

consistent with AHL10 and RRP6L1 acting together in gene regulation. Many of the genes

with altered expression in unstressed *rrp6l1-2* were stress responsive in wild type

(Supplemental Fig. S7A). Particularly, genes with increased expression in unstressed *rrp6l1-*

2 were enriched for GO terms related to abiotic stress or environmental stimuli

(Supplemental Table S2), suggesting that RRP6L1 has a role in preventing aberrant

expression of these genes in unstressed conditions.

In the low ψ_w stress treatment, relatively fewer genes were differentially expressed in

rrp6l1-2 (11 genes had increased expression, 11 genes had decreased expression including

RRP6L1; log2 fold-change $\geq$ 0.5 versus wild type at low ψ_w, adjusted P $\leq$ 0.05; Supplemental

Table S3). There was again substantial concordance between gene expression changes seen

in *rrp6l1-2* and *ahl10-1* (Fig. 4A). This concordance could be seen even more clearly if we

considered a slightly relaxed cutoff for differential expression (log2 fold-change $\geq$ 0.3,

adjusted P $\leq$ 0.05) as shown in Supplemental Fig. S7B. The closest concordance between

rrp6l1-2 and *ahl10-1* was for genes that had decreased expression in both mutants (Fig. 4A;

Supplemental Fig. S7B). We validated the decreased expression of a number of these genes

in low ψ_w-treated *rrp6l1-2* and *ahl10-1* by QPCR (Fig. 4B). Interestingly, given the

proposed functions of *rrp6l1-2* (Wendte et al., 2017), *RNA-Dependent RNA Polymerase 3*

(*RdR3*) and *RdR4* had decreased expression in both *rrp6l1-2* and *ahl10-1*. These two RdRs

are of unclear physiological function (Willmann et al., 2011). The large number of genes

with relatively small change in expression in *rrp6l1-2* and *ahl10-1* (Supplemental Fig. S7B)

was not unexpected given the tissue specific accumulation of both proteins (Fig. 2) such that

localized changes in gene expression may have been diluted in the whole seedling samples

used for RNAseq.

AHL10 has been previously associated with AT-rich transposons (Jiang et al., 2017).

Our transcriptome analysis of *ahl10-1* found altered RNA levels of several transposons

(Wong et al., 2019) and many of these had similarly altered expression in *rrp6l1-2*. For

example, the copia-like retrotransposon *At5g35935* and hAT-like transposase family gene

At3g29120 were the two most highly upregulated genes in the low ψ_w treatment for both

ahl10-1 and *rrp6l1-2* (Fig. 4A; Supplemental Table S3). *At5g35935* is adjacent to the

promoter region of *At5g35940* which was down-regulated in both *rrp6l1-2* and *ahl10-1*.

ChIP assays found low ψ_w-induced enrichment of AHL10 at several AT-rich sites in the

region between *At5g35935* and *At5g35940* (Fig. 4C). This stress-induced association was not

observed for a nearby region in the *At5g35940* gene body or a distal off-site (EIF4a

promoter). A similar enrichment pattern was observed for AHL10[S314A], consistent with the

idea that the phosphonull mutation in the C-terminal region of AHL10 does not interfere with

the DNA binding region in the N-terminal part of the protein. We also conducted ChIP

assays for AT-rich promoter regions within 1 kb of the transcriptional start site for several

other genes having decreased expression in *ahl10-1* and *rrp6l1-2* (Supplemental Fig. S8).

However, we did not find enrichment of AHL10 on those sites. Possibly AHL10 is

associated with more distal sites which influence expression of those genes by formation of

chromatin loops which bring AHL10 (and RRP6L1) in closer proximity to the gene or its

newly transcribed RNA.

Two studies have previously identified regions of reduced methylation in *rrp6l1-2*

(Zhang et al., 2014; Wendte et al., 2017) or conducted transcriptome analysis of *rrp6l1-2*

(Zhang et al., 2014). However, we found little overlap between *rrp6l1-2* differentially

expressed genes identified by Zhang et al. (2014) and those identified in our analysis

(Supplemental Fig. S9A) or between *rrp6l* DEGs identified in any study and *rrp6l1*-related

demethylated regions (Supplemental Fig. S9B). The differences in the results could be

influenced by slight age difference in seedlings between experiments (11 days in our case,

12-15 days for the other studies). However, the very limited overlap in differentially

expressed genes and demethylated regions between these data sets suggests a strong effect of

environmental conditions on RRP6L1 action in transcriptional regulation.

Discussion

Our results demonstrate that RRP6L1 is involved in low ψ_w response, that AHL10

and RRP6L1 interact and, that AHL10 S314 phosphorylation may facilitate AHL10 function

as an auxiliary factor that promotes RRP6L1 chromatin association. Both RRP6L1 and

AHL10 protein levels were high in meristem tissue indicating that this is where AHL10-

RRP6L1 interaction mainly occurs and where AHL10 and RRP6L1 exert their effects on

growth during low ψ_w stress. High levels of AHL10 and RRP6L1 in meristem can explain

their effects on expression of meristem-specific genes such as *STM* and the auxin-related

genes such as *WES1* and *DFL1*. In the context of previous work, our results indicate that

control of AHL10 phosphorylation by competing action of MPKs versus Clade A PP2Cs

(particularly HAI1) in meristems is a new connection of stress-related phospho-signaling to

gene regulation and growth suppression during low ψ_w stress.

While our experiments focused on the effect of AHL10 phosphorylation and did not

seek to investigate the mechanism of RRP6L1-mediated gene regulation, it is of interest to

put our results in the context of previous characterization of RRP6L1.   The lack of RRP6L1

chromatin recruitment in *ahl10-1* during low ψ_w stress is surprising as it indicates that

AHL10 has a controlling influence on RRP6L1 compared with other RRP6L1-interacting

proteins such as the C-Terminal Domain (CTD) of NRPE1, the largest subunit of RNA

Polymerease V (Wendte et al., 2017). In that study the authors speculated that RRP6L1

interaction with the CTD is transient or involves only a small portion of Pol V. Our data

suggest that the converse may also be true: CTD interaction may involve only a small portion

of RRP6L1, particularly during low ψ_w stress. Alternatively, RRP6L1 interaction with

AHL10 could precede, or be required for, CTD interaction (or other interactions) that

position RRP6L1 at specific chromatin locations. Zhang et al. (2014) noted that it is not

known what factors determine RRP6L1 preference for certain sites over others. Determining

whether AHL10 binding is competitive with CTD binding, or whether AHL10 binding can

influence the 3'-5' exonuclease activity of RRP6L1 (Zhang et al., 2014) will be questions of

interest for future studies.   It should also be kept in mind that previous studies of RRP6L1

(Lange et al., 2008; Shin and Chekanova, 2014; Zhang et al., 2014; Ye et al., 2016; Wendte et

al., 2017) have only observed RRP6L1 function under unstressed conditions and mainly

focused on RRP6L1 effects on non-coding RNAs. Thus, it is unclear how their proposed

mechanisms of RRP6L1 function may be altered by low ψ_w stress and whether RRP6L1

affects protein coding genes via distinct mechanisms compared to its effects on non-coding

RNAs.

The lack of RRP6L1 chromatin recruitment in *ahl10-1* is also surprising given that

one may have expected a greater degree of redundancy among the closely related Clade B

AHLs. Also surprising is our observation that AHL10 and AHL13 are not genetically

redundant, despite their overall high similarity and conserved C-terminal phosphorylation site

(Wong et al., 2019; Rayapuram et al., 2021). Instead, *ahl13-1* blocked the *ahl10-1* mutant

phenotype of increased growth maintenance during low ψ_w stress. While they are not

redundant, AHL10 and AHL13 have common regulatory targets. For example, both affect

the regulation of JA-related genes (*AOC1, AOS*) in response to low ψ_w (for *ahl10-1*) or

pathogen infection (for *ahl13-1;* Wong et al., 2019; Rayapuram et al., 2021). These

seemingly contradictory results can be interpreted in light of previous evidence that AHLs act

as trimers (Zhao et al., 2013). It is presumably these trimeric complexes that recruit other

transcriptional regulators; albeit, this hypothesis awaits direct testing *in planta*. AHL13 may

act as an anchor for different AHL complexes, thus explaining why it is genetically

downstream of AHL10. In this case, the combination of AHL13 with AHL10 gives

specificity of recruiting additional regulators such as RRP6L1. Knocking out one of these

genes disrupts the composition of AHL complexes and thus disrupts recruitment of

interacting proteins and downstream phenotypes in ways more complex than simple

redundancy among AHLs.

Data Availability

RNAseq data has been deposited to Gene Expression Omnibus and is available under

accession number GSE101197. All other data supporting the findings of this study are in the

paper and supplemental material.

Supplemental Data

Supplemental Figure S1: AHL10 interactors.

Supplemental Figure S2: *In-vitro* pull down experiments confirm direct interaction of AHL10

and RRP6L1.

Supplemental Figure S3: Complementation of *ahl10-1* with AHL10 phosphonull and

phosphomimic mutants under control of the *AHL10* native promoter.

Supplemental Figure S4: AHL10 accumulates in root and shoot meristems.

Supplemental Figure S5: Effect of phospho-null and phospho-mimic mutations on AHL10

sub-nuclear localization pattern.

Supplemental Figure S6: Effect of *RRP6L1, AHL10* and *AHL13* mutants on growth during

moderate severity low water potential stress.

Supplemental Figure S7: Gene expression changes in *rrp6l1-2* and *ahl10-1* compared to low

water potential stress-induced changes in wild type.

Supplemental Figure S8: ChIP assay of AT-rich promoter sites for other genes with

decreased expression in *ahl10-1* and *rrp6l1-2* under low water potential stress.

Supplemental Figure S9: Comparison of our *rrp6l1-2* transcriptome data to genes in or near

hypomethylated regions in previously reported analyses.

Supplemental Table S1: Genes up or down regulated in *rrp6l1-2* compared to Col-0 wild type

in the unstressed control treatment.

Supplemental Table S2: Gene ontology analysis of genes up or down regulated in *rrp6l1-2* in

the unstressed control treatment.

Supplemental Table S3: Genes up or down regulated in *rrp6l1-2* compared to Col-0 wild type

in the low water potential stress treatment (-0.7 MPa, 96 h).

Supplemental Table S4: Primers used for cloning, genotyping, mutagenesis, gene expression

analysis and ChIP.

Acknowledgements

We thank David Favero (Riken) for providing cDNA clones of AHL27 and AHL29; Ming-

Che Shih (Agricultural Biotechnology Research Center, Academia Sinica) for providing the

35S:GFP transgenic line; Toshisangba Longkumer for experimental assistance, Ji-Ying

Huang and Mei-Jane Fang (Live Cell Imaging core facility, Institute of Plant and Microbial

Biology) for microscopy assistance; Wen-Dar Lin (Bioinformatics Core Facility, Institute of

Plant and Microbial Biology) for assistance with transcriptome data analysis and Shih-Shan

Huang for laboratory assistance. This research was supported by an Academia Sinica

Investigator Award (AS-IA-108-L04) and a National Science and Technology Council of

Taiwan grant (MoST 110-2311-B-001 -019) to P.E.V.

Author Contributions

P.E.V. conceived research, M.M.W., Y.-C.B. and P.E.V. designed experiments, M.M.W., X.-

J..H and Y.-C.B. performed experiments, M.M.W., X.-J..H, Y.-C.B. and P.E.V. analyzed

data, P.E.V. wrote the manuscript with assistance from M.M.W. All authors read and

approved the manuscript.

References

Antoni R, Gonzalez-Guzman M, Rodriguez L, Rodrigues A, Pizzio GA, Rodriguez PL (2011) Selective inhibition of clade A phosphatases type 2C by PYR/PYL/RCAR abscisic acid receptors. Plant Physiol 158: 970-980

Bhaskara GB, Nguyen TT, Verslues PE (2012) Unique Drought Resistance Functions of the Highly ABA-Induced Clade A Protein Phosphatase 2Cs. Plant Physiology 160: 379-395

Bhaskara GB, Wong MM, Verslues PE (2019) The flip side of phospho-signalling: Regulation of protein dephosphorylation and the protein phosphatase 2Cs. Plant Cell and Environment 42: 2913-2930

Chong GL, Foo MH, Lin WD, Wong MM, Verslues PE (2019) Highly ABA-Induced 1 (HAI1)-Interacting protein HIN1 and drought acclimation-enhanced splicing efficiency at intron retention sites. Proceedings of the National Academy of Sciences of the United States of America 116: 22376-22385

Cutler SR, Rodriguez PL, Finkelstein RR, Abrams SR (2010) Abscisic Acid: Emergence of a Core Signaling Network. *In* S Merchant, WR Briggs, D Ort, eds, Annual Review of Plant Biology, Vol 61, Vol 61, pp 651-679

Danquah A, de Zelicourt A, Colcombet J, Hirt H (2014) The role of ABA and MAPK signaling pathways in plant abiotic stress responses. Biotechnology Advances 32: 40-52

Favero DS, Kawamura A, Shibata M, Takebayashi A, Jung JH, Suzuki T, Jaeger KE, Ishida T, Iwase A, Wigge PA, Neff MM, Sugimoto K (2020) AT-Hook Transcription Factors Restrict Petiole Growth by Antagonizing PIFs. Current Biology 30: 1454-+

Franco-Zorrilla JM, Lopez-Vidriero I, Carrasco JL, Godoy M, Vera P, Solano R (2014) DNA-binding specificities of plant transcription factors and their potential to define target genes. Proceedings of the National Academy of Sciences of the United States of America 111: 2367-2372

Fujimoto S, Matsunaga S, Yonemura M, Uchiyama S, Azuma T, Fukui K (2004) Identification of a novel plant MAR DNA binding protein localized on chromosomal surfaces. Plant Molecular Biology 56: 225-239

Grefen C, Blatt MR (2012) A 2in1 cloning system enables ratiometric bimolecular fluorescence complementation (rBiFC). Biotechniques 53: 311-314

Hsu Y-F, Hsiao Y-C, Wang C-S, Zhan X, Zhang H, Wang C-S (2014) AtRRP6L1, a Homolog of Conserved Yeast Exosomal Rrp6p, Plays an Important Role in Transcriptional Gene Silencing in Arabidopsis. Molecular Plant 7: 1490-1493

Jiang H, Moreno-Romero J, Santos-Gonzalez J, De Jaeger G, Gevaert K, Van De Slijke E, Kohler C (2017) Ectopic application of the repressive histone modification H3K9me2 establishes post-zygotic reproductive isolation in Arabidopsis thaliana. Genes & Development 31: 1272-1287

Lange H, Holec S, Cognat V, Pieuchot L, Le Ret M, Canaday J, Gagliardi D (2008) Degradation of a polyadenylated rRNA maturation by-product involves one of the three RRP6-Like proteins in *Arabidopsis thaliana*. Molecular and Cellular Biology 28: 3038-3044

Lee K, Seo PJ (2017) Coordination of matrix attachment and ATP-dependent chromatin remodeling regulate auxin biosynthesis and Arabidopsis hypocotyl elongation. Plos One **12:** e0181804

Mine A, Berens ML, Nobori T, Anver S, Fukumoto K, Winkelmuller TM, Takeda A, Becker D, Tsuda K (2017) Pathogen exploitation of an abscisic acid- and jasmonate-inducible

MAPK phosphatase and its interception by Arabidopsis immunity. Proceedings of the National Academy of Sciences of the United States of America 114: 7456-7461

Morisawa G, Han-yama A, Moda I, Tamai A, Iwabuchi M, Meshi T (2000) AHM1, a Novel Type of Nuclear Matrix–Localized, MAR Binding Protein with a Single AT Hook and a J Domain–Homologous Region. Plant Cell **12:** 1903-1916

Nicot N, Hausman JF, Hoffmann L, Evers D (2005) Housekeeping gene selection for real-time RT-PCR normalization in potato during biotic and abiotic stress. Journal of Experimental Botany 56: 2907-2914

Rayapuram N, Jarad M, Alhoraibi HM, Bigeard J, Abulfaraj AA, Volz R, Mariappan KG, Almeida-Trapp M, Schloffel M, Lastrucci E, Bonhomme L, Gust AA, Mithofer A, Arold ST, Pflieger D, Hirt H (2021) Chromatin phosphoproteomics unravels a function for AT-hook motif nuclear localized protein AHL13 in PAMP-triggered immunity. Proceedings of the National Academy of Sciences of the United States of America 118: e2004670118

Seo M, Lee JY (2021) Dissection of Functional Modules of AT-HOOK MOTIF NUCLEAR LOCALIZED PROTEIN 4 in the Development of the Root Xylem. Frontiers in Plant Science 12

Shin JH, Chekanova JA (2014) Arabidopsis RRP6L1 and RRP6L2 Function in FLOWERING LOCUS C Silencing via Regulation of Antisense RNA Synthesis. Plos Genetics 10

Tsuda K, Qi YP, Nguyen LV, Bethke G, Tsuda Y, Glazebrook J, Katagiri F (2012) An efficient Agrobacterium-mediated transient transformation of Arabidopsis. Plant Journal 69: 713-719

Verslues PE, Agarwal M, Katiyar-Agarwal S, Zhu JH, Zhu JK (2006) Methods and concepts in quantifying resistance to drought, salt and freezing, abiotic stresses that affect plant water status. Plant Journal 45: 523-539

Wendte JM, Haag JR, Singh J, McKinlay A, Pontes OM, Pikaard CS (2017) Functional Dissection of the Pol V Largest Subunit CTD in RNA-Directed DNA Methylation. Cell Reports 19: 2796-2808

Willmann MR, Endres MW, Cook RT, Gregory BD (2011) The Functions of RNA-Dependent RNA Polymerases in Arabidopsis. The Arabidopsis Book 2011

Wong MM, Bhaskara GB, Wen TN, Lin WD, Nguyen TT, Chong GL, Verslues PE (2019) Phosphoproteomics of *Arabidopsis* Highly ABA-Induced1 identifies AT-Hook-Like10 phosphorylation required for stress growth regulation. Proceedings of the National Academy of Sciences of the United States of America 116: 2354-2363

Wu Z, Zhu DL, Lin XY, Miao J, Gu LF, Deng X, Yang Q, Sun KT, Zhu DM, Cao XF, Tsuge T, Dean C, Aoyama T, Gu HY, Qu LJ (2016) RNA Binding Proteins RZ-1B and RZ-1C Play Critical Roles in Regulating Pre-mRNA Splicing and Gene Expression during Development in Arabidopsis. Plant Cell 28: 55-73

Xu L, Zheng S, Witzel K, Slijke EVD, Baekelandt A, Mylle E, Damme DV, Cheng J, Jaeger GD, Inze D, Jiang H (2023) Chromatin attachment to the nuclear matrix represses hypocotyl elongation in *Arabidopsis thaliana*. bioRxiv**:** 2023.2007.2005.547753

Xu Y, Wang Y, Stroud H, Gu X, Sun B, Gan E-S, Ng K-H, Jacobsen SE, He Y, Ito T (2013) A Matrix Protein Silences Transposons and Repeats through Interaction with Retinoblastoma-Associated Proteins. Current Biology **23:** 345-350

Yamaguchi N, Winter CM, Wu M-F, Kwon CS, William DA, Wagner D (2014) PROTOCOL: Chromatin Immunoprecipitation from Arabidopsis Tissues. The Arabidopsis Book 2014

Ye R, Chen Z, Lian B, Rowley MJ, Xia N, Chai J, Li Y, He X-J, Wierzbicki Andrzej T, Qi Y
(2016) A Dicer-Independent Route for Biogenesis of siRNAs that Direct DNA
Methylation in Arabidopsis. Molecular Cell 61: 222-235
Zhang HM, Tang K, Qian WQ, Duan CG, Wang BS, Zhang H, Wang PC, Zhu XH, Lang ZB,
Yang Y, Zhu JK (2014) An Rrp6-like Protein Positively Regulates Noncoding RNA
Levels and DNA Methylation in Arabidopsis. Molecular Cell 54: 418-430
Zhao JF, Favero DS, Peng H, Neff MM (2013) Arabidopsis thaliana AHL family modulates
hypocotyl growth redundantly by interacting with each other via the PPC/DUF296
domain. Proceedings of the National Academy of Sciences of the United States of
America 110: E4688-E4697

Figures and Legends

Figure 1: AHL10 interacts with RRP6l1 and both proteins accumulate in root and shoot meristem regions.

A. β☐galactosidase staining colony lift assays show the interaction of AHL10 with clones encoding the C-terminal portion of RRP6L1. RRP6L1$^{\Delta1-562}$ is the clone found in the original yeast two hybrid library screening. Other truncations and full length *RRP6L1* clones were used to confirm the interaction. AHL10 self-interaction is shown for comparison as another strong interaction detected in the yeast two hybrid screen. Empty prey vector is a negative control. Additional proteins found in the yeast two hybrid screen are listed in Supplemental Fig. S1A.

B. Quantification of interaction intensity for rBIFC assays conducted using unstressed Arabidopsis seedlings or seedlings exposed to low ψ_w stress (-0.7 MPa for 48 h).

Experiments were conducted using transient expression in Arabidopsis seedlings with

constructs driven by the *35S* promoter (see methods). Individual data points represent

the YFP/RFP fluorescence intensity ratio of individual cells. Data are combined from

three independent experiments (n = 13 - 97). Different letters indicate statistically

significant differences (ANOVA, corrected $P \leq 0.05$).

C. Representative images of rBiFC data presented in B. Scale bars indicate 10 μm.

D. Localization of AHL10-YFP expressed under control of its native promoter (genomic

clone of the *AHL10* gene body consisting of 5' UTR, introns and exons up to the stop

codon, and 2.1 kb promoter region; with C-terminal YFP fusion).   Images show the

primary root tip of 9-day-old seedlings growing control media or four days after

transfer to low ψ_w (-0.7 MPa). Scale bars indicate 50 μm.

E. Validation of polyclonal antisera raised against the C-terminal region of RRP6L1.

Red arrow marks major band corresponding to the molecular weight of RRP6L1 (74

kD) which is absent in both *rrp6l1* mutants. 35 μg of protein was loaded in each lane.

A non-specific band was detected at approximately 110 kD. The blot was stripped

and re-probed with anti-HSC70 as a loading control. The experiment was repeated

with essentially identical results. C = Control; S = Stress (-0.7 MPa for four days).

The blot was stripped and re-probed with anti-HSC70 as a loading control.

F. Detection of RRP6L1 and AHL10-YFP in whole seedling, shoot meristem-enriched

(shoot tip), and root meristem-enriched (root tip) samples. 35 μg of protein was

loaded in each lane. The RRP6L1 blot was stripped and re-probed with anti-HSC70

as a loading control. The experiment was repeated three times with consistent results

each time.

Figure 2: *RRP6L1* **is a negative regulator of growth during moderate severity low ψ$_w$**
stress and *ahl10-1* **effect on growth depends upon** *AHL13***.**

A. Seedling dry weight and primary root elongation in control and stress treatments. For control,

five-day-old seedlings were transferred to fresh control agar plates with root length increase

and total seedling dry weight measured five days after transfer. For stress treatment, five-day-

old seedlings were transferred from control media to PEG-infused agar plates (-0.7 MPa) and

dry weight and root elongation measured 10 days after transfer. Data are expressed relative to

wild type grown in the same plate as the test genotype. Pictures of representative seedlings in

the stress treatment are shown in Supplemental Fig. S6A. Wild type dry weight and root

elongation data used for normalization are shown in Supplemental Fig. S6B. Data are means

± S.D. for 3-5 replicate experiments with 2-3 plates of seedlings measured in each experiment.

Asterisks (*) indicate significant difference from 100% (wild type level) by one sample T-test.

B. Rosette fresh and dry weights for soil-grown plants kept under well-watered conditions or

exposed to partial soil drying. Data are expressed relative to wild type plants grown together

in the same pot. Data are means ± S.D. for 3-5 replicate experiments (2-3 biological

replicates per experiment). Asterisks (*) indicate significant difference from 100% (wild type

level) by on sample T-test.

C. Pictures of representative wild type and *rrp6l1-2* plants from the experiments presented in B.

Scale bars indicate 1 cm.

Figure 3: RRP6L1 chromatin association depends upon AHL10 S314 phosphorylation.

A. Portion of AHL10-YFP associated with chromatin, measured as the ratio of chromatin

associated (C.A.) versus total nuclear (T.N.) immunoblot band intensities for *rrp6l1-2*

or *ahl10-1* complemented with non-mutated AHL10 (N.M), phosphonull AHL10

(S313A, S314A) or phosphomimic AHL10 (S313D, S314D). Representative blots

used for quantitation can be seen in C and D. Samples were collected for 11-day-old

seedlings kept under unstressed control conditions or transferred to low ψ_w stress

treatment (-0.7 MPa for four days). Data are means ± S.D. for three or four

independent experiments (one replicate per experiment). Asterisks or indicated P

value at the bars in the graph indicate the results of ANOVA comparing all genotypes

within the stress or control treatments to the wild type AHL10 (N.M.) data for that

treatment. The wild type AHL10 (N.M.) data were further compared between control

and stress treatment using T-test (Asterisk indicates significant difference, $P \leq 0.05$).

B. Portion of RRP6L1 associated with chromatin measured as the ratio of chromatin

associated (C.A.) versus total nuclear (T.N.) immunoblot band intensities for *ahl10-1*

or *ahl10-1* complemented with non-mutated AHL10 (N.M), phosphonull AHL10

(S313A, S314A) or phosphomimic AHL10 (S313D, S314D). Representative blots

used for quantitation can be seen in C and D. Sampling conditions and statistical

analysis are as described in A. Asterisks above bars indicate a significant difference

compared to AHL10 N.M. in that treatment ($P \leq 0.05$).

C. Representative blots used for band intensity quantitation shown in A and B. The

AHL10 genomic fragment with 5' UTR and 2.1 kb promoter region was fused to YFP

and expressed in the *ahl10-1* or *ahl10-1rrp6l1-2* genetic backgrounds to detect

AHL10 chromatin association in the presence of absence of RRP6L1. The *ahl10-1*

mutant was used to detect RRP6L1 chromatin association in the absence of AHL10.

Red arrow indicates the RRP6L1 specific band which matches the predicted RRP6L1

molecular weight (74 kD) and was used for quantification. T.N. = Total Nuclear; S.F.

= Soluble Fraction of nuclear proteins; C.A. = Chromatin Associated; P.S. = Pellet

after Sonication (to check for remaining insoluble protein). Detection of Histone3

(H3) was performed as a control for the purity of the S.F. and C.A fractions.

However, note that we consistently observed that stress treatment caused an increase

in the amount of H3 in the soluble fraction which was not related to any technical

aspects of the chromatin precipitation procedure.

D. Representative blots of lines expressing phosphomimic or phosphonull AHL10 used

for band intensity quantification shown in A and B.

E. Nuclear fractionation and chromatin association of a transgenic line expressing GFP

under control of the constitutive *35S* promoter used to check the background level of a

non-chromatin associated protein in the chromatin pellet. Numbers under the GFP

blot indicate the fraction of chromatin associated GFP based on the ratio of C.A and

T.N. band intensities. Data are representative of two biologically independent

experiments.

Figure 4: *rrp6l1-2* and *ahl10-1* have concordant effects on gene expression and low ψ_w stress can increase association of AHL10 with specific AT-rich regions.

A. Heat map and hierarchical clustering of genes having significantly increased or decreased expression (log2 fold change ≥ 0.5, corrected P-value ≤ 0.05) in stress-treated (transfer to -0.7 MPa for four days) seedlings. Effect of stress treatment on these genes in wild type is also shown for comparison. Underlined genes were used as targets of ChIP assay shown in Fig. 4C. Similar heat map of genes with altered expression in unstressed *rrp6l1-2* or *ahl10-1* is shown in supplemental Fig. S7A. Heat map of genes affected by stress in *rrp6l1-2* or *ahl10-1* at slightly less stringent cutoff (log2 fold change ≥ 0.3, corrected P-value ≤ 0.05) is shown in Supplemental Fig. S7B. Genes labeled as "RNA" are non-coding RNAs. T.E. = Transposable Element.

B. QPCR validation of gene expression differences of *rrp6l1-2* and *ahl10-1* compared to
wild type in the stress treatment. Note that some of the genes selected for validation
did not meet the cutoff for inclusion in the heat map shown in Fig. 4A but are
included in the heat map shown in Supplemental Fig S7B. Data are means ± S.D. for
data from 3 independent biological experiments (one sample per experiment, three
technical PCR replicates per sample). Data are shown as log2 fold change relative to
wild type. Asterisks indicate significant difference between *ahl10-1* and *rrp6l1-2* by
T-test (**, $P \leq 0.01$). Note that for some genes (*STM, WES1, DFL1, MYB117* and
PRR) the QPCR data *ahl10-1* was previously reported in Wong et al. (2019) and is
shown again here to facilitate comparison to *rrp6l1-2*.

C. ChIP assays of AT-rich sites lying in the region between *At5g35935* and *At5g35940*
(*At5g35940* promoter region) as well as an off-site in exon of *At5g35940* and a distal
offsite (*EIF4a* promoter). Samples were collected for unstressed and low ψ_w -treated
(-0.7 MPa for four days) seedlings. Data are means ± S.D. for data from 3
independent biological experiments (one sample per experiment, three technical PCR
replicates per sample). Asterisks indicate significant differences compared to the
EIF1α offsite in the stress treatment (ANOVA, $P \leq 0.05$). No significant differences
were found in the unstressed control.