Running head: Learning dietary specialization

The ontogeny of individual specialization

Anne G. Hertel[a,b], Jörg Albrecht[b], Nuria Selva[c,d], Agnieszka Sergiel[c], Keith A. Hobson[e,f],

David M. Janz[g], Andreas Mulch[b], Jonas Kindberg[h,i], Jennifer E. Hansen[j], Shane C. Frank[j],

Andreas Zedrosser[j,k], Thomas Mueller[b,l]

[a]*Behavioural Ecology, Department of Biology, Ludwig-Maximilians University of Munich,*

Planegg-Martinsried, Germany

[b]*Senckenberg Biodiversity and Climate Research Centre (SBiK-F), Frankfurt (Main),*

Germany

[c]*Institute of Nature Conservation, Polish Academy of Sciences, Adama Mickiewicza 33,*

31120 Krakow, Poland

[d]*Departamento de Ciencias Integradas, Facultad de Ciencias Experimentales, Centro de*

Estudios Avanzados en Física, Matemáticas y Computación, Universidad de Huelva, 21071

Huelva, Spain

[e]*Environment and Climate Change Canada, Science and Technology, Saskatoon, SK, Canada*

[f]*Department of Biology and Advanced Facility for Avian Research (AFAR), University of*

Western Ontario, London, ON, Canada

[g]*Department of Veterinary Biomedical Sciences, Western College of Veterinary Medicine,*

University of Saskatchewan, 52 Campus Drive, Saskatoon, Saskatchewan S7N 5B4, Canada

[h]*Norwegian Institute for Nature Research, Trondheim, Norway*

[i]*Department of Wildlife, Fish, and Environmental Studies, Swedish University of Agricultural*

Sciences, Umeå, Sweden

^{j}*Department of Natural Sciences and Environmental Health, University of South-Eastern*

Norway, Bø, Norway

^{k}*Institute of Wildlife Biology and Game Management, University of Natural Resources and*

Life Sciences, Vienna, Austria

^{l}*Department of Biological Sciences, Goethe University Frankfurt, Max-von-Laue-Straße 9,*

60438 Frankfurt (Main), Germany

Corresponding author: Anne Gabriela Hertel, Ludwig-Maximilian University of Munich,

Planegg-Martinsried, Germany, hertel@biologie.uni-muenchen.de

ABSTRACT

Individual dietary specialization, where individuals occupy a subset of a population's wider

dietary niche, is of key importance for species' resilience towards environmental change.

However, the ontogeny of individual specialization, as well as the underlying social learning,

genetic, and environmental drivers remain poorly understood. Using a multigenerational

dataset of female European brown bears (*Ursus arctos*) followed since birth, we discerned

the relative contributions of social learning, genetic predisposition, environment forcings, and

maternal effects to individual dietary specialization. Individual specialization varied from

omnivorous to carnivorous diets spanning half a trophic position. The main determinants of

this dietary specialization were maternal learning during rearing (11%), environmental

similarity (12%), and maternal effects (11%), whereas the contribution of genetic heritability

was negligible. Importantly, the offspring's trophic position closely resembled the trophic

position of their mother during the first 3-4 years after separation from the mother, but this

relationship ceased with increasing time since separation. Our study reveals that social

learning and maternal effects are as important for individual dietary specialization as

environmental forcings. We propose a tighter integration of social effects into future studies

of range expansion and habitat selection under global change, that to date are mostly

explained by environmental drivers.

KEYWORDS

Dietary specialization, heritability, maternal effects, maternal learning, trophic position,

trophic niche, omnivore, stable isotopes, carbon-13, nitrogen-15, *Ursus arctos*

1. INTRODUCTION

Among individuals of the same species, niche variation is common and may occur when availability of food resources or habitat structure change across the species' range. Ecological generalists, species with a wide niche, also seem to exhibit more individual specialization [1] and are hence particularly well adapted to persist under shifts in resource availability or composition enabling them to occupy larger distributional ranges than ecological specialists [2]. Individual variation is key for making species resilient towards changing resource availabilities in a rapidly changing world and may ultimately determine local persistence or extinction of species [3].

Inter- and intraspecific competition, predation and ecological opportunity, alter resource availability and have been identified as the main ecological drivers explaining variation in the degree of individual specialization between populations [4]. Yet, how individual variation emerges and is maintained within populations has been rarely quantified in the wild. In principle, four potential sources of variation exist: social and individual learning, genetic inheritance, the environment, and maternal effects. Individual differences in resource preference or competence to secure a resource may therefore be determined during early ontogeny through social (e.g., maternal) learning via imitation [5-9] leading to similarities between the offspring's and their mother's dietary phenotype. Effects of maternal learning can be lifelong or are modified through individual experiential learning [10]. Resource preferences have also been suggested to be genetically determined through genes inherited from both mother and father, where closely related individuals have more similar diets than distantly related individuals [11]. In addition, maternal effects account for lifelong similarities in dietary phenotype among offspring of the same mother [12]. Such similarities can arise from social interactions, maternal genotype, or maternal environment. Statistically, maternal effects are quantified as the similarity of repeated samples from siblings of the same

mother but not as the similarity of behavioral expression between mother and offspring (i.e.,

"maternal learning"). In range resident species, where individuals occupy a subset of a

population's range, the environment, in terms of habitat composition or availability of

particular food resources, may differ among home ranges and lead to individual

specialization [11]. Accounting for the environmental heterogeneity when studying the

drivers of individual specialization is therefore essential in range resident species [13, 14].

Attributing variation in diet to the individual level, to isolate its sources and to identify

developmental drivers of diet preferences requires multigenerational datasets of repeated

measures of the diet of individuals throughout their life. We harness a 30-year longitudinal

dataset of 72 female Scandinavian brown bears of known mothers with repeated measures of

annual diets to first assess whether individual specialization occurs. Using information about

their mother's diet, a genetic pedigree, and individual movement data we then aim to attribute

individual variation in diet to its sources: maternal learning, genetic heritability, environment,

and maternal effects.

Brown bears are ecological generalists with a species range spanning the northern

hemisphere from tundra to deserts, paralleled by extensive variation in diet: from populations

tracking food resource pulses, such as spawning fish [15], over ones scavenging on ungulate

carcasses or preying on ungulates neonates [16], feeding extensively on invertebrates, to

populations with primarily fruiting plant based diets [17, 18]. Given this extreme dietary

plasticity, it is not surprising that great dietary variation has been found within populations

[19, 20], however, the determinants and ontogeny of this variation at the individual level

remain largely unknown. In ecology, differences in diet are often primarily attributed to

differences in resource availability and abundance. Even within populations inhabiting a

continuous biome, home range scale variation in habitat composition [21] can lead to

variation in resource availability. The most parsimonious source of variation in diet are,

therefore, differences in the environment. Brown bears maintain non-territorial home ranges

but live a solitary lifestyle except for the period of offspring rearing with up to three years

[22] of maternal care. In their first years of life, bear cubs accompany their mother and it is

therefore reasonable to assume that brown bear offspring learn behaviors such as habitat, den

site, or diet selection from their mothers and hence show similar behavior to their mother. If

mothers differ in their dietary selection, these differences may hence be maintained in the

population through learning by imitation of the mother (hereafter "maternal learning"), even

after offspring gain independence, however such similarities may wane over time [23]. On

the other hand, genetic heritability or maternal effects can have lifelong effects on offspring

phenotype. Body size has been shown to be genetically heritable in the study population [24]

suggesting greater similarity among closely related individuals also in other linked traits,

such as trophic position. Alternatively, maternal effects (i.e., maternal genotype or maternal

environment) alone can shape the phenotype of offspring. For example, milk quantity or

quality [25] can vary among females either due to genetic differences or differences in the

environments, leading to greater similarity among all offspring from the same mother (e.g.

being smaller or larger in body size), which in turn could cause similarities in trophic position

among siblings. To assess individual specialization along a continuum from a more plant-

based to a more meat- or insect-based diet, we analyzed annual trophic positions from

nitrogen stable isotopes ($\delta^{15}N$) stored in bear hair keratin [26]. Stable isotopes reflect

cumulative diet intake and are deposited into the hair during growth with, a delay of

approximately one month (i.e. a growing hair in June reflects the diet intake in May, [27]).

Bear hair is regularly renewed through molting in June, regrows over the summer and fall

and stops growing during winter hibernation (**Fig 1A, [28, 29]**). Guard hair samples collected

in spring and early summer (April - June) therefore reflect an individual's diet during the

previous active season prior to hibernation [28]. Using repeated samples of known mother-

daughter pairs, we fit a spatially explicit Bayesian hierarchical model (i.e. ´animal model´)

[30-32] to disentangle the relative contributions of maternal learning, genetic relatedness, the

environment, and maternal effects as determinants of individual specialization. Specifically,

the model accounted for genetic relatedness with a pedigree and for environmental similarity

of bear home ranges with pairwise habitat similarity encompassing the proportion of mature

habitat such as old and mid-successional forests, disturbed habitat such as clearcuts and

regenerating young forest, and habitat diversity (measured as Simpson's diversity index) in a

bear's home range. The model also accounted for maternal effects by incorporating the

mother's ID as a random effect (i.e. if daughters from the same mother behaved in a similar

fashion throughout life), and for maternal learning as the fixed effect of a mother's trophic

positions on her daughter's trophic position. To this end, we determined maternal trophic

positions from a population-wide model accounting for sexual dimorphism, age, and

individual consistency in diet (**Supplement 3**). Because bears may alter diet selection over

time through individual learning, we allowed the effect of maternal learning to shift with time

since the offspring gained independence. Last, we also accounted for permanent individual

effects that could not be attributed to any of the aforementioned sources, by including a

random effect for bear ID. We focused on the effect of maternal trophic position on female

offspring trophic position, because male offspring were only monitored for a short period

after family breakup. In the supplementary material we provide an additional analysis of the

relationship between maternal and both female and male offspring trophic position in the first

4 years after family breakup and of the relationship between paternal trophic position and

offspring trophic position. We also provide an alternative analysis accounting for spatial

correlation via a spatial distance instead of the habitat similarity matrix. Last, we provide a

reduced model excluding the effect of environmental similarity to test whether spatial and

genetic effects are confounded in philopatric female bears.

2. RESULTS

We analyzed annual trophic positions in 213 hair samples collected from 71 female brown bears born to 33 unique mothers (1 – 7 daughters per mother; median 2 daughters). Repeated sampling (median 3 years; range 1-11 years) revealed that female trophic position was unaffected by age (explained variance = 1% [0 – 4%]) and that individuals showed long-term individual specialization, accounting for 48% [31 – 61%] (median [89% equal tails credible interval]) of the total variance in trophic position (**Fig 2,** *Basic model*). Individual specialization spanned half a trophic position ranging from 2.7 to 3.1 for individual females (**Fig 1B**), which is equivalent to the difference between an omnivore feeding on a mix of plants and animal prey and a carnivore feeding predominantly on animal prey.

Individual specialization was primarily driven by initial maternal learning, the environment, and maternal effects. Maternal trophic position dynamic over the time since separation accounted for 13% [5% - 23%] of variation in trophic position, while environmental similarity accounted for 9% [0.1 - 5%] of the total phenotypic variation in trophic position. Additionally, maternal effects accounted for 11% [0.5% - 30%] of variation in trophic position, indicating that siblings (full and half) of the same mother were more similar in trophic position throughout life as compared to non-siblings. A remaining 8% [0.3 – 26%] of variance in trophic position was attributed to permanent individual effects (**Fig 2**).Genetically more closely related individuals did not share a more similar trophic position (3% [<0.1% – 17%] of variance explained) providing no evidence that dietary specialization could be heritable in this population (**Fig 2**).

After separating from their mother, female offspring initially maintained a similar trophic position as their mother (Pearson's $r = 0.66$ in the first two years after separation), which gradually became more dissimilar over time (Pearson's $r = 0.31$ in year 3 - 4 after separation,

Fig 3). In the first years, offspring of more carnivorous mothers also had a high trophic

position while offspring of less carnivorous mothers had a lower trophic position. About five

years after the separation of the mother, this correlation ceased to exist. Bears inhabiting

home ranges with a similar composition of mature and disturbed forest, as well as a similar

habitat diversity in the home range, also had more similar trophic positions. The distance

between pairwise home range centroids ranged from 0.7 to 172 km with a median pairwise

distance of 48 km and individuals living in closer proximity had a more similar trophic

position than individuals living farther apart (**Supplementary material S5**). Spatial distance

and maternal effects seemed to be confounded in this female philopatric species (**Fig S5**):

After excluding spatial distance, maternal learning and maternal effects but not heritability

explained more variance in trophic position, corroborating that spatial proximity is

confounded with philopatric females forming clusters of mothers and daughter in space, so

called matrilines (**Fig S6**). In a separate analysis (**Fig S7**) we could also show that the

relationship between maternal and offspring trophic position in the first years after family

breakup was not sex-specific. Both male (n = 31, Pearson correlation coefficient = 0.4) and

female (n = 69, Pearson correlation coefficient = 0.45) offspring's trophic positions

resembled their mother's trophic position in the first 4 years of independence, corroborating

our findings that initial maternal learning determines foraging behavior in the early years

after family breakup. Conversely, paternal trophic position had no effect on offspring trophic

position in the first 4 years of independence (Pearson correlation coefficient = 0.13, **Fig S8**).

Figure 1. A) Bear hair generally grows from June until October. Stable isotopes are deposited into the growing hair with a delay of approximately one month. The quiescent phase, when hair ceases growing, lasts through hibernation, followed by emergence from the winter den and molting in late May-early June. Hair samples were taken in April - June and reflect the bears' diet in the previous year; **B**) Posterior distribution of the population trophic niche (bold line) and individual specialization indicated by each individual's posterior trophic position. Scientific illustration by Juliana D. Spahr, SciVisuals.com.

Figure 2. Proportion of variance (median of the posterior distribution) in brown bear trophic position explained by age, age sensitive maternal learning, permanent individual effects,

environmental similarity, permanent maternal effects, genetic heritability, and residual

components.

Figure 3. Relationship between female brown bear trophic position and their mother's

trophic position over number of years since separation (i.e. since the female became

independent, usually at 1.5 years of age). The females' trophic position resembled their

mothers' in the first years after separation but this similarity ceased after 4 years.

3. DISCUSSION

Our multigenerational dataset reveals unique insights into the ontogeny of individual dietary

specialization along a continuum from a more herbivorous to a more carnivorous diet in a

long-lived omnivore. Specifically, the foraging strategy of sons and daughters was intimately

tied to the foraging strategy of their mother, a relationship that lasted up to four years after

independence. We interpret this relationship as evidence that maternal learning plays an

important role in shaping an individual's dietary specialization. Five years into independence,

the similarity between the mothers' and their daughters' trophic position slowly faded, likely

due to individual learning and experience. In addition, siblings of the same mother also

shared lifelong similarities in their trophic position, potentially mediated through maternal

genetic or environmental effects on body size [24]. In general, previous ecological studies

have mainly concentrated on resource availability as the main driver of resource selection

[33] and individual specialization [4], however our results show that within populations, the

environment is only one of several components shaping individual dietary variation. We

conclude that early-life imitation of maternal dietary preferences and maternal effects (i.e.,

maternal genotype and environment), which together explained about 24% of the variation in

trophic position, play a pivotal role in spreading and maintaining feeding strategies within

populations, even in species with otherwise solitary lifestyles. In addition, variation solely

linked to individual variation (in our study 8 %) demonstrates potential for behavioral

innovation and the potential to adapt to changing conditions.

Our findings are particularly relevant for species in which dietary specialization impacts

individual fitness [7, 34, 35]. For example, protein-rich diets may promote greater offspring

survival or mass gain [36]. Maternal and social learning in general therefore present an

important, yet understudied pathway by which alternative behavioral strategies can establish

and spread more rapidly within populations than by genetic evolution alone [37]. Species

more adept in social learning of dietary strategies may therefore show greater behavioral

variability on the population level, which could give them an advantage when adapting to

changing environments due to landscape modification or urbanization, climatic variations or

global change in general. Moreover, there is evidence that the strength of social learning in

shaping individual phenotypes is not only species-specific, but can also vary among

populations or individuals of the same species [38].

Our research also points to several aspects of maternal learning that warrant future research.

First, there is little information on whether maternal care and maternal learning tend to be

more prevalent in species or populations with greater dietary specialization. There is some

evidence that within populations, dietary generalists (i.e. those with a wider dietary niche)

seem to provide more intense parental care [39], than their conspecific dietary specialists (i.e.

ones with a narrower dietary niche), but the links to parental learning of foraging preferences

remains unclear. Second, while generalist species with a wide ecological niche have been

frequently shown to be more successful under changing environmental conditions, such as

urban environments or fragmented landscapes, than specialist species [40-42], it is currently

unknown whether this success could be partially mediated by social or maternal learning.

Last, social learning could alternatively limit behavioral innovation and adaptation due to

adherence to social traditions [43]. We therefore suggest that alternative hypotheses should

be evaluated that consider how social learning impacts individual specialization and in turn

the adaptability of species under global change.

Our findings that dietary specialization can be socially learned and transmitted are

particularly relevant for species where specialization is related to human-wildlife conflict

[44]. For example, the removal of single individuals which are known to cause conflict is an

effective strategy to halt the spread of problematic behavior, increase societal acceptance by

effectively mitigating the conflict, while minimizing the impact for species conservation

goals [44]. Foraging behavior that causes conflict has also been shown to change in ursids

across life time, remarking the crucial role of individuality and plasticity in behavior [45].

Maternal learning of behavior [46], including dietary specialization and foraging on

anthropogenic food resources is commonly observed in ursids [47-50]. However, none of

these studies tracked offspring diet over their lifetimes or were able to simultaneously

account for the mother's diet, genetics, the environment, and other maternal effects, that

could explain similar patterns of dietary specialization. While some of the aforementioned

studies suggest either the environment or maternal learning as primary drivers of individual

specialization, we suggest using caution in assigning causality in dietary specialization, when

potentially confounding alternative sources cannot be accounted for. Specifically, in female-

biased philopatric species, spatial proximity does not only encode for spatial variation in

resource abundance but is also conflated with relatedness and, in particular, with maternal

effects. In brown bears, philopatric daughters settle close to their mother's home range [51]

creating spatial clusters of closely related females, so called matrilinear assemblages [52].

Due to spatial dependence of matrilinear assemblages, it can therefore be difficult to

disentangle maternal learning from other maternal effects (i.e., maternal genotype or maternal

environment) or the ambient environment. Our study population spanned over 170 km with

spatial proximity explaining 59% of the total phenotypic variation in trophic position of

female bears: individuals further apart tended to have more different diets. However, when

replacing spatial proximity with environmental similarity among home ranges, the

explanatory power was attributed to maternal learning and maternal effects along with the

environment. Our results therefore demonstrate that individual dietary specialization is not

caused by a single driver in isolation but the product of many factors, namely: maternal

learning, maternal effects, and the environment.

Our finding that maternal learning has a similar impact on resource selection as the

environment provides important insights for a range of studies on habitat selection, dispersal,

and range expansion. For example, a popular theory known as "natal habitat preference

induction" suggests that dispersing animals select areas for settlement that resemble their

natal habitat, even at fine habitat scales [21]. Our results challenge the notion that habitat

similarity alone drives natal settlement strategies and rather suggest that maternally induced

diet preferences, and hence the selection for food resources themselves, could play an

important role in producing similar patterns of settlement selection like induced natal habitat

preferences. Recent studies of migration and short stopover behavior in whooping cranes

(*Grus americana*) have also observed that social learning rather than environmental

conditions [53] or genetic inheritance [54] led to the emergence and establishment of

alternative migratory behavior. Similar to what our study shows with respect to dietary

specialization, social learning of migration strategies primarily determined behavior in early

life whereas individual-experiential learning shaped behavior later in life [55].

Conclusion

Drivers of dietary specialization are well documented among populations of the same species,

however, systematic studies delineating the sources of individual specialization within

populations are lacking, likely because suitable datasets including multigenerational, genetic,

environmental, and life-history information are rare. We show here that in addition to the

environment, maternal learning and (other) maternal effects can be important sources of

dietary specialization.

4. METHODS

Bear sample collection

We collected brown bear (*Ursus arctos*) hair samples in south-central Sweden (~N61°, E15°)

as part of a long-term, individual-based monitoring project (Scandinavian Brown Bear

Research Project; www.bearproject.info). Hair samples were collected from known

individuals and their offspring during bear captures in spring (April - June) 1993 – 2015 after

bears emerged from hibernation. Bears were immobilized from a helicopter (Arnemo &

Fahlman, 2011). A vestigial premolar tooth was collected from all bears not captured as a

yearling to estimate age based on the cementum annuli in the root [56]. Bears were weighed

in a stretcher suspended beneath a spring scale. Tissue samples (stored in 95% alcohol) were

taken for DNA extraction to assign parentage and construct a genetic pedigree [52]. Guard

hairs and follicles were plucked with pliers from a standardized spot between the shoulder

blades and archived at the Swedish National Veterinary Institute. We used data of adult bears

(solitary or with offspring) and of offspring after separation from their mother. Bear cubs are

born in January or February during winter hibernation and are typically first captured

together with their mother as yearlings at the age of ~ 15 months. Cubs in this population

separate from their mother during the mating season in May or June after 1.5 or 2.5 years

[57]. Only hair samples of solitary, independent offspring taken in spring and early summer

at least 10 months after separation from the mother were included in this study. A hair sample

taken in spring reflects the summer-fall diet of the bear in the previous active season (**Fig**

1A).

Food sample collection

We collected samples of the natural foods most important for brown bear in the study area,

including 21 samples of moose hair (*Alces alces*), the most common meat source in the

brown bears' diet in our study area [58], in the spring-autumn field season of 2014 (**Fig S1**).

Samples were placed in a paper envelope and dried at ambient temperature for 48.

Stable isotope analyses

Hair samples were rinsed with a 2:1 mixture of chloroform:methanol or with pure methanol

to remove surface oils [59]. Dried samples were ground with a ball grinder (Retsch model

MM-301, Haan, Germany).

We weighed 1 mg of ground hair into pre-combusted tin capsules and combusted at 1030°C

in a Carlo Erba NA1500 elemental analyser. N_2 and CO_2 were separated chromatographically

and introduced to an Elementar Isoprime isotope ratio mass spectrometer (Langenselbold,

Germany). Two reference materials were used to normalize the results to VPDB and AIR:

BWB III keratin ($\delta^{13}C$ =- 20.18‰, $\delta^{15}N$ = 14.31‰, respectively) and PRC gel ($\delta^{13}C$ =-

13.64‰, $\delta^{15}N$ = 5.07‰, respectively). Measurement precisions as determined from both

reference and sample duplicate analyses were ± 0.1‰ for both $\delta^{13}C$ and $\delta^{15}N$.

Bear trophic position

We calculated the trophic position of each bear hair sample relative to the average $\delta^{15}N$ value

of moose (mean ± sd = 1.8 ± 1.26 ‰, n = 21, **Fig S1**). Trophic position is calculated as the

the discrepancy of $\delta^{15}N$ in a secondary consumer and its food source divided by the

enrichment of $\delta^{15}N$ per trophic level, plus lambda, the trophic position of the food source

(e.g. 1 for primary producers, 2 for primary consumers, 3 for secondary consumer, 4 for

tertiary consumers) [60]. We used an average trophic enrichment factor of 3.4‰ [60] and

added a lambda of 2 given that the moose baseline trophic position as a strict herbivore.

$$\text{Bear trophic position} = (\delta^{15}N_{Ursus\ arctos} - \text{average}(\delta^{15}N_{Alces\ alces})) / 3.4 + 2$$

Under an omnivorous diet including the consumption of herbivores (in particular moose but

also herbivorous insects), bear trophic position values were expected to fall between 2 and 3.

Values approaching 4 indicate a trophic enrichment through consumption of other

omnivorous or carnivorous animals.

Genetic pedigree and parentage assignment

A genetic pedigree based on 16 microsatellite loci was available for the population including 1614 individual genotypes [61]. Genotyping followed the protocols of Waits, Taberlet [62], Taberlet, Camarra [63], and Andreassen, Schregel [64]. All female offspring in this study were genotyped and included in the population's genetic pedigree. All females included in this study had a known mother that was also captured and followed. We used Cervus 3.0 [65] for assignment of fathers and COLONY [66] for creating putative unknown mother or father genotypes and sibship reconstruction (see [61] for details).

Maternal trophic position

Based on repeated hair samples of 115 female (nfemale = 335) and 98 male (nmale = 219) bears, we fitted a *basic* linear mixed effects model for female and male bears respectively, to estimate sex-specific among individual variation in trophic position (**Supplementary analysis 3**). We modelled trophic position as a function of a quadratic relationship with age and we controlled for individual random intercepts. Female trophic position did not vary with age (**Fig S3A**) but was highly repeatable over multiple years (**Fig S3B**). For all daughters, we extracted their mother's (and father's) trophic position as the median of the posterior distribution of their respective random intercept.

Environmental similarity

Resources, i.e. access to moose and ants, may not be distributed evenly in space. For moose, population density and hunting quotas (which determine availability of slaughter remains) vary across the study area. For ants (*Formica* spp.*, Camponotus herculeanus*), the availability of old forests and clearcuts determine their abundance [67]. Further, brown bear daughters are often philopatric with limited dispersal and settle close to their mother's home range [51].

Genetic, spatial, and maternal learning effects may therefore be confounded with related

bears occupying adjacent ranges with similar environments and resource availability.

Elsewhere, accounting for environmental similarity through spatial autocorrelation in animal

models has revealed that a major portion of variance may be attributed to environmental

similarity rather than genetic heritability [30, 31, 68, but see also 69]. Here, we accounted for

environmental similarity by extracting habitat composition in each bear's lifetime home

range. For individuals with sufficient locations (> 1000 GPS locations or VHF locations on at

least 25 days) we constructed home ranges using a 95% kernel density estimator. We used a

corine landcover map (25 m resolution) which we updated annually with polygons of newly

emerged clearcuts (data obtained from the Swedish Forest Agency). We extracted home

range composition in the year when diet was assessed. When individuals were monitored for

multiple years, we extracted the home range composition for the median year. We calculated

the proportion of mid-aged and old forest and proportion of disturbed forest (clearcuts and

regenerating young forest) within the 95% utilization distribution. Additionally, we

calculated habitat diversity using the Simpson diversity index from the R package

landscapemetrics [70]. Following Thomson et al. [30] we calculated the Euclidean distance

between scaled and centered habitat composition and habitat diversity in multivariate space,

assuming equal importance of each component. Pairwise distances were scaled between 0

and 1, where increasing values indicated more similar habitat composition. In the

supplementary material we provide an alternative analysis accounting for spatial

autocorrelation in dietary specialization with a pairwise spatial distance matrix (S matrix;

Supplementary analysis 5, **Fig S5**).

Statistical analysis

We applied a two-step modelling approach. First, we fitted a *basic* linear mixed effects model

to estimate individual specialization as among individual variation in annual trophic position.

We accounted for a nonlinear effect of age (second order polynomial) and for repeated

measures of the same individual with individual random intercepts. We extracted the variance

in fitted values (variance explained by fixed effects), among-individual, and residual variance

and estimated the proportional contribution of fixed and random effects on the total

phenotypic variance through variance standardization (i.e. repeatability [71], marginal and

conditional R^2-values [72]). Second, we used a spatially explicit Bayesian hierarchical model

(i.e. 'animal model') [30, 32] to partition among-individual variance in trophic position into

environmental similarity (σ^2_{env}), additive genetic (σ^2_a), permanent among-individual (σ^2_{ind}),

maternal ($\sigma2_{mat}$), and residual within-individual effects (σ^2_r). Similar to the basic model, we

accounted for a nonlinear effect of age on trophic position (fitted as time since separation of

mother and daughter scaled by the standard deviation, true age and time since separation

were perfectly correlated: Pearson correlation coefficient > 0.99). We tested for maternal

effects on offspring trophic position by incorporating the mother's trophic position as a

covariate into the model. To account for a potential decrease of the maternal effect over time,

we let maternal trophic position interact with the time since separation of mother and

daughter (both scaled by their standard deviation and centered). We partitioned the variance

explained by the two components of the fixed effect, the effect of maternal learning over time

(i.e. maternal trophic position and the interaction between maternal trophic position and time

since separation) and age (i.e. the main effect of time since separation), respectively, by

calculating the independent contribution of each component to the total variance explained by

the fixed effects, following the approach by Stoffel, Nakagawa [73] adapted to a Bayesian

framework (see code under [74]).

All models were fit using the R package "brms" [75] based on the Bayesian software

Stan [76, 77]. We ran four chains to evaluate convergence which were run for 6,000

iterations, with a warmup of 3,000 iterations and a thinning interval of 10. All estimated

model coefficients and credible intervals were therefore based on 1200 posterior samples and

had satisfactory convergence diagnostics with $\hat{R} < 1.01$, and effective sample sizes > 400

[78]. Posterior predictive checks recreated the underlying Gaussian distribution of trophic

position well. For all parameters, we report the median and 89% credible intervals, calculated

as equal tail intervals, as measure of centrality and uncertainty [79]. We deemed explained

variance proportions as inconclusive when the lower credible interval limit was < 0.001 (i.e.,

$< 0.1\%$) [80]. All statistical analyses were performed in R 4.0.0 [81]. Primary data and code

to reproduce all analyses are provided under (https://doi.org/10.17605/OSF.IO/68B9U, [74]).

ACKNOWLEDGEMENTS

AGH has received funding from the European Union's Horizon 2020 Research and

Innovation Programme under the Marie Skłodowska-Curie Grant agreement No 793077 and

from the German Science Foundation (HE 8857/1-1). Isotope analyses were funded through a

Robert Bosch Foundation grant to TM and analyzed by KH and DJ. The study was further

funded by the Norway Grants under the Polish-Norwegian Research Programme

administered by the National Research Centre for Research and Development in Poland and

the Norwegian Research Council (JA, NS, AS, and AZ; GLOBE No POL-

NOR/198352/85/2013). We thank the Scandinavian Brown Bear Research Project (SBBRP)

for providing access to the data. The SBBRP was funded by the Norwegian Environment

Agency, the Swedish Environmental Protection Agency, the Austrian Science Fund, and the

Norwegian Research Council.

AUTHOR'S CONTRIUTIONS

AH, JA, and TM developed the work. AZ, JK, KAH, NS and AS provided the data. AZ

managed the sample collection. AS managed the hair samples database and prepared samples

for stable isotope analyses by KAH. DMJ provided laboratory space and resources and

supervised preparatory procedures. SF constructed and provided the genetic pedigree. JH

provided home range centroids. AM and JA advised to the analysis and interpretation of

stable isotope data. TM, NS and AZ secure project funding. AH performed the statistical

analyses with input from JA. AH wrote the manuscript with help from TM, JA, and input

from all authors.

1. Bolnick, D.I., et al., *Comparative support for the niche variation hypothesis that more generalized populations also are more heterogeneous.* Proceedings of the National Academy of Sciences of the United States of America, 2007. **104**(24): p. 10075-10079.
2. Huang, S., et al., *Scale-dependent effects of niche specialisation: The disconnect between individual and species ranges.* Ecology Letters, 2021. **24**(7): p. 1408-1419.
3. Forsman, A. and L. Wennersten, *Inter-individual variation promotes ecological success of populations and species: evidence from experimental and comparative studies.* Ecography, 2016. **39**(7): p. 630-648.
4. Araújo, M.S., D.I. Bolnick, and C.A. Layman, *The ecological causes of individual specialisation.* Ecology Letters, 2011. **14**(9): p. 948-958.
5. Estes, J.A., et al., *Individual variation in prey selection by sea otters: patterns, causes and implications.* Journal of Animal Ecology, 2003. **72**(1): p. 144-155.
6. Thornton, A. and K. McAuliffe, *Teaching in wild meerkats.* Science, 2006. **313**(5784): p. 227-229.
7. Annett, C.A. and R. Pierotti, *Long-term reproductive output in western gulls: consequences of alternative tactics in diet choice.* Ecology, 1999. **80**(1): p. 288-297.
8. Altbäcker, V., R. Hudson, and Á. Bilkó, *Rabbit-mothers' Diet Influences Pups' Later Food Choice.* Ethology, 1995. **99**(1-2): p. 107-116.
9. Slagsvold, T. and K.L. Wiebe, *Social learning in birds and its role in shaping a foraging niche.* Philos Trans R Soc Lond B Biol Sci, 2011. **366**(1567): p. 969-77.
10. Reid, A.L., F. Seebacher, and A.J.W. Ward, *Learning to hunt: the role of experience in predator success.* Behaviour, 2010. **147**(2): p. 223-233.
11. Daniel I. Bolnick, et al., *The Ecology of Individuals: Incidence and Implications of Individual Specialization.* The American Naturalist, 2003. **161**(1): p. 1-28.
12. McAdam, A.G., D. Garant, and A.J. Wilson, *The effects of others' genes: maternal and other indirect genetic effects*, in *Quantitative Genetics in the Wild*, A. Charmantier, D. Garant, and L.E.B. Kruuk, Editors. 2014, Oxford University Press. p. 84-103.

13. Hertel, A.G., et al., *A guide for studying among-individual behavioral variation from*
movement data in the wild. Movement Ecology, 2020. **8**(1): p. 30.
14. Sprau, P. and N.J. Dingemanse, *An Approach to Distinguish between Plasticity and*
Non-random Distributions of Behavioral Types Along Urban Gradients in a Wild
Passerine Bird. Frontiers in Ecology and Evolution, 2017. **5**(92).
15. Deacy, W., et al., *Kodiak brown bears surf the salmon red wave: direct evidence from*
GPS collared individuals. Ecology, 2016. **97**(5): p. 1091-1098.
16. Bojarska, K. and N. Selva, *Spatial patterns in brown bear Ursus arctos diet: the role*
of geographical and environmental factors. Mammal Review, 2012. **42**(2): p. 120-
143.
17. Qin, A., et al., *Predicting the current and future suitable habitats of the main dietary*
plants of the Gobi Bear using MaxEnt modeling. Global Ecology and Conservation,
2020. **22**: p. e01032.
18. Rodríguez, C., et al., *Long-term trends in food habits of a relict brown bear*
population in northern Spain: the influence of climate and local factors.
Environmental Conservation, 2007. **34**(01): p. 36-44.
19. Edwards, M.A., et al., *Fast carnivores and slow herbivores: differential foraging*
strategies among grizzly bears in the Canadian Arctic. Oecologia, 2011. **165**(4): p.
877-889.
20. Mangipane, L.S., et al., *Dietary plasticity in a nutrient-rich system does not influence*
brown bear (Ursus arctos) body condition or denning. Polar Biology, 2018. **41**(4): p.
763-772.
21. Merrick, M.J. and J.L. Koprowski, *Evidence of natal habitat preference induction*
within one habitat type. Proceedings of the Royal Society B: Biological Sciences,
2016. **283**(1842).
22. Steyaert, S.M., et al., *The mating system of the brown bear Ursus arctos.* Mammal
Review, 2012. **42**(1): p. 12-34.
23. White, S.J. and A.J. Wilson, *Evolutionary genetics of personality in the Trinidadian*
guppy I: maternal and additive genetic effects across ontogeny. Heredity, 2019.
122(1): p. 1-14.
24. Rivrud, I.M., et al., *Heritability of head size in a hunted large carnivore, the brown*
bear (Ursus arctos). Evolutionary Applications, 2019. **12**(6): p. 1124-1135.
25. Renaud, L.-A., et al., *Causes and short-term consequences of variation in milk*
composition in wild sheep. Journal of Animal Ecology, 2019. **88**(6): p. 857-869.
26. Deniro, M.J. and S. Epstein, *Influence of diet on the distribution of nitrogen isotopes*
in animals. Geochimica et Cosmochimica Acta, 1981. **45**(3): p. 341-351.
27. Rode, K.D., et al., *Isotopic Incorporation and the Effects of Fasting and Dietary Lipid*
Content on Isotopic Discrimination in Large Carnivorous Mammals. Physiol
Biochem Zool, 2016. **89**(3): p. 182-97.
28. Jimbo, M., et al., *Hair Growth in Brown Bears and Its Application to Ecological*
Studies on Wild Bears. Mammal Study, 2020. **45**(4): p. 337-345, 9.
29. Cattet, M., et al., *Can concentrations of steroid hormones in brown bear hair reveal*
age class? Conservation Physiology, 2018. **6**(1).
30. Thomson, C.E., et al., *A guide to using a multiple-matrix animal model to disentangle*
genetic and nongenetic causes of phenotypic variance. PLOS ONE, 2018. **13**(10): p.
e0197720.
31. Gervais, L., et al., *Quantifying heritability and estimating evolutionary potential in*
the wild when individuals that share genes also share environments. Journal of
Animal Ecology, 2022. **n/a**(n/a).

32.   Wilson, A.J., et al., *An ecologist's guide to the animal model.* Journal of Animal
Ecology, 2010. **79**(1): p. 13-26.
33.   MacArthur, R.H. and E.R. Pianka, *On optimal use of a patchy environment.* American
Naturalist, 1966: p. 603-609.
34.   Zango, L., et al., *Year-round individual specialization in the feeding ecology of a*
long-lived seabird. Scientific Reports, 2019. **9**(1): p. 11812.
35.   Balme, G.A., et al., *Ecological opportunity drives individual dietary specialization in*
leopards. Journal of Animal Ecology, 2020. **89**(2): p. 589-600.
36.   Seress, G., et al., *Food availability limits avian reproduction in the city: An*
experimental study on great tits Parus major. Journal of Animal Ecology, 2020.
89(7): p. 1570-1580.
37.   Sutherland, W.J., *Evidence for Flexibility and Constraint in Migration Systems.*
Journal of Avian Biology, 1998. **29**(4): p. 441-446.
38.   Mesoudi, A., et al., *The Evolution of Individual and Cultural Variation in Social*
Learning. Trends in Ecology & Evolution, 2016. **31**(3): p. 215-225.
39.   Nicolaus, M., S.C.Y. Barrault, and C. Both, *Diet and provisioning rate differ*
predictably between dispersing and philopatric pied flycatchers. Behavioral Ecology,
2018. **30**(1): p. 114-124.
40.   Sol, D., S. Timmermans, and L. Lefebvre, *Behavioural flexibility and invasion*
success in birds. Animal Behaviour, 2002. **63**(3): p. 495-502.
41.   Sol, D., O. Lapiedra, and C. González-Lagos, *Behavioural adjustments for a life in*
the city. Animal Behaviour, 2013. **85**(5): p. 1101-1112.
42.   Devictor, V., R. Julliard, and F. Jiguet, *Distribution of specialist and generalist*
species along spatial gradients of habitat disturbance and fragmentation. Oikos,
2008. **117**(4): p. 507-514.
43.   Keith, S.A. and J.W. Bull, *Animal culture impacts species' capacity to realise*
climate-driven range shifts. Ecography, 2017. **40**(2): p. 296-304.
44.   Swan, G.J.F., et al., *Ecology of Problem Individuals and the Efficacy of Selective*
Wildlife Management. Trends in Ecology & Evolution, 2017. **32**(7): p. 518-530.
45.   Berezowska-Cnota, T., et al., *Individuality matters in human–wildlife conflicts:*
Patterns and fraction of damage-making brown bears in the north-eastern
Carpathians. Journal of Applied Ecology, In Press. **n/a**(n/a).
46.   Lillie, K.M., et al., *Development of on-shore behavior among polar bears (Ursus*
maritimus) in the southern Beaufort Sea: inherited or learned? Ecology and
Evolution, 2018. **8**(16): p. 7790-7799.
47.   Morehouse, A.T., et al., *Nature vs. Nurture: Evidence for Social Learning of Conflict*
Behaviour in Grizzly Bears. PLOS ONE, 2016. **11**(11): p. e0165425.
48.   Shimozuru, M., et al., *Maternal human habituation enhances sons' risk of human-*
caused mortality in a large carnivore, brown bears. Scientific Reports, 2020. **10**(1):
p. 16498.
49.   Mazur, R. and V. Seher, *Socially learned foraging behaviour in wild black bears,*
Ursus americanus. Anim Behav, 2008. **75**.
50.   Jimbo, M., et al., *Diet selection and asocial learning: Natal habitat influence on*
lifelong foraging strategies in solitary large mammals. Ecosphere, 2022. **13**(7): p.
e4105.
51.   Hansen, J.E., et al., *Social environment shapes female settlement decisions in a*
solitary carnivore. Behavioral Ecology, 2021. **33**(1): p. 137-146.
52.   Frank, S.C., et al., *Harvest is associated with the disruption of social and fine-scale*
genetic structure among matrilines of a solitary large carnivore. Evolutionary
Applications, 2021. **14**(4): p. 1023-1035.

53. Mendgen, P., et al., *Differential shortstopping behaviour in Whooping Cranes: Habitat or social learning?* Global Ecology and Conservation, 2023. **41**: p. e02365.

54. Mueller, T., et al., *Social Learning of Migratory Performance.* Science, 2013. **341**(6149): p. 999-1002.

55. Abrahms, B., et al., *Ontogenetic shifts from social to experiential learning drive avian migration timing.* Nature Communications, 2021. **12**(1): p. 7326.

56. Matson, G., et al., *A laboratory manual for cementum age determination of Alaska brown bear first premolar teeth.* Matson's Laboratory, Milltown, Montana, USA, 1993.

57. Van de Walle, J., et al., *Hunting regulation favors slow life histories in a large carnivore.* Nature Communications, 2018. **9**(1): p. 1100.

58. Stenset, N.E., et al., *Seasonal and annual variation in the diet of brown bears Ursus arctos in the boreal forest of southcentral Sweden.* Wildlife Biology, 2016. **22**(3): p. 107-116.

59. Sergiel, A., et al., *Compatibility of preparatory procedures for the analysis of cortisol concentrations and stable isotope (δ(13)C, δ(15)N) ratios: a test on brown bear hair.* Conservation physiology, 2017. **5**(1): p. cox021-cox021.

60. Post, D.M., *Using stable isotopes to estimate trophic position: models, methods, and assumptions.* Ecology, 2002. **83**(3): p. 703-718.

61. Frank, S.C., et al., *Harvest is associated with the disruption of social and fine-scale genetic structure among matrilines of a solitary large carnivore.* Evolutionary Applications. **n/a**(n/a).

62. Waits, L., et al., *Nuclear DNA microsatellite analysis of genetic diversity and gene flow in the Scandinavian brown bear (Ursus arctos).* Mol Ecol, 2000. **9**(4): p. 421-31.

63. Taberlet, P., et al., *Noninvasive genetic tracking of the endangered Pyrenean brown bear population.* Mol Ecol, 1997. **6**(9): p. 869-76.

64. Andreassen, R., et al., *A forensic DNA profiling system for Northern European brown bears (Ursus arctos).* Forensic Sci Int Genet, 2012. **6**(6): p. 798-809.

65. Kalinowski, S.T., M.L. Taper, and T.C. Marshall, *Revising how the computer program cervus accommodates genotyping error increases success in paternity assignment.* Molecular Ecology, 2007. **16**(5): p. 1099-1106.

66. Jones, O.R. and J. Wang, *COLONY: a program for parentage and sibship inference from multilocus genotype data.* Molecular Ecology Resources, 2010. **10**(3): p. 551-555.

67. Frank, S.C., et al., *A "clearcut" case? Brown bear selection of coarse woody debris and carpenter ants on clearcuts.* Forest Ecology and Management, 2015. **348**: p. 164-173.

68. Stopher, K.V., et al., *Shared spatial effects on quantitative genetic parameters: accounting for spatial autocorrelation and home range overlap reduces estimates of heritability in wild red deer.* Evolution, 2012. **66**(8): p. 2411-2426.

69. Regan, C.E., et al., *Accounting for female space sharing in St. Kilda Soay sheep (Ovis aries) results in little change in heritability estimates.* Journal of Evolutionary Biology, 2017. **30**(1): p. 96-111.

70. Hesselbarth, M.H.K., et al., *landscapemetrics: Landscape Metrics for Categorical Map Patterns. R package version 1.0.* 2019.

71. Dingemanse, N.J., et al., *Behavioural reaction norms: animal personality meets individual plasticity.* Trends in Ecology & Evolution, 2010. **25**(2): p. 81-89.

72. Nakagawa, S. and H. Schielzeth, *A general and simple method for obtaining R2 from generalized linear mixed-effects models.* Methods in Ecology and Evolution, 2013. **4**(2): p. 133-142.

664   73.   Stoffel, M.A., S. Nakagawa, and H. Schielzeth, *partR2: partitioning R2 in*
665   *generalized linear mixed models.* PeerJ, 2021. **9**: p. e11414.
666   74.   Hertel, A.G., *Data&Code: The ontogeny of individual specialization.* 2023.
667   75.   Bürkner, P.-C., *brms: An R package for Bayesian multilevel models using Stan.*
668   Journal of Statistical Software, 2017. **80**(1): p. 1-28.
669   76.   Stan Development Team, *RStan: the R interface to Stan. R package version 2.17.3.*
670   2018.
671   77.   Carpenter, B., et al., *Stan: A Probabilistic Programming Language.* 2017, 2017.
672   **76**(1): p. 32.
673   78.   Vehtari, A., et al., *Rank-normalization, folding, and localization: An improved R for*
674   *assessing convergence of MCMC.* Bayesian Analysis, 2020.
675   79.   Kruschke, J., *Doing Bayesian data analysis: A tutorial with R, JAGS, and Stan.* 2014.
676   80.   Bonnet, T., et al., *Genetic variance in fitness indicates rapid contemporary adaptive*
677   *evolution in wild animals.* Science, 2022. **376**(6596): p. 1012-1016.
678   81.   R Core Team, *R: A language and environment for statistical computing. R*
679   *Foundation for Statistical Computing.* 2020.
680