Title: High-throughput Automated Muropeptide Analysis (HAMA) Reveals Peptidoglycan

Composition of Gut Microbial Cell Walls

Authors: Ya-Chen Hsu[1], Pin-Rui Su[1,2], Lin-Jie Huang[1], Kum-Yi Cheng[1], Chun-hsien Chen[1],

Cheng-Chih Hsu[1]*

Affiliations:

[1] Department of Chemistry, National Taiwan University, Taipei 10617, Taiwan

[2] Department of Molecular Genetics, Erasmus MC, Rotterdam, Netherlands

Corresponding Author:

* Cheng-Chih Hsu

Department of Chemistry, National Taiwan University, Taipei 10617, Taiwan

+886-2-3366-3844, ccrhsu@ntu.edu.tw

ORCID: Cheng-Chih Hsu: 0000-0002-2892-5326

ABSTRACT

Peptidoglycan (PGN), a net-like polymer constituted by muropeptides, provides protection for microorganisms and has been a major target for antibiotics for decades. Researchers have explored host-microbiome interactions through PGN recognition systems and discovered key muropeptides modulating host responses. However, most common characterization techniques for muropeptides are labor-intensive and require manual analysis of mass spectra due to the complex cross-linked PGN structures. Each species has unique moiety modifications and inter-/intra-bridges, which further complicates the structural analysis of PGN. Here, we developed a high-throughput automated muropeptide analysis (HAMA) platform leveraging tandem mass spectrometry and *in silico* muropeptide MS/MS fragmentation matching to comprehensively identify muropeptide structures, quantify their abundance, and infer PGN cross-linking types. We demonstrated the effectiveness of HAMA platform using well-characterized PGNs from *E. coli* and *S. aureus* and further applied it to common gut bacteria including *Bifidobacterium, Bacteroides, Lactobacillus, Enterococcus,* and *Akkermansia muciiniphila.* Specifically, we found that the stiffness and strength of the cell envelopes may correspond to the lengths and compositions of interpeptide bridges within *Bifidobacterium* species. In summary, the HAMA framework exhibits an automated, intuitive, and accurate analysis of PGN compositions, which may serve as a potential tool to investigate the post-synthetic modifications of saccharides, the variation in interpeptide bridges, and the types of cross-linking within bacterial PGNs.

INTRODUCTION

Peptidoglycan (PGN) is an essential and distinct component of bacterial cell walls that envelop the cytoplasmic membrane.[1] It maintains a defined bacterial shape, preserves cell integrity by counteracting the internal osmotic pressure, and acts as a scaffold for other cell wall-anchored components such as proteins and teichoic acids.[2] Peptidoglycan provides a protective barrier for microorganisms against environmental threats, contributing to its structural rigidity. Generally, PGN is a net-like polymeric structure composed of liner glycan chains cross-linked by short peptides. The PGN-repeat unit (so-called muropeptide) comprises *N*-acetylglucosamine (GlcNAc) and *N*-acetylmuramic acid (MurNAc) linked by β1,4-glycosidic bonds and a peptide stem attached to lactyl moiety of MurNAc. The stem peptide generally contains L-alanine, D-iso-glutamic acid (amidated to D-iso-glutamine in most Gram-positive bacteria), *meso*-diaminopimelic acid (mDAP) in Gram-negative bacteria, L-lysine in Gram-positive bacteria, D-alanine, and D-alanine sequentially. In PGN architecture, portions of the stem peptide stem from adjacent glycan strands are connected through transpeptidation, forming a multi-layered, three-dimensional (3-D) meshwork.[3] Although this main structure is well-conserved among bacteria, the mature peptidoglycan structure remains heterogeneous and species-specific. This structural diversity is attributed to chemical modifications of the sugar backbone (*O*-acetylation, *N*-deacetylation, and *N*-glycolylation), variations in the type of di-amino acid at position 3 of the stem peptide, variations in the interpeptide bridge, and patterns of cross-linking (4-3 linkage, 3-3 linkage, and 2-4 linkages).[4, 5]

In recent years, characterizing muropeptides has become increasingly important, not only because peptidoglycan is a target for antibiotic drug design, but also because of the various roles muropeptides play as signaling molecules involved in microbial interaction, antimicrobial

release, and host innate immunity.[6-9] The gut microbiota, consisting of thousands of bacteria, is a

source of peptidoglycans in organisms. The signaling functions of gut microbiota-derived PGN

fragments in host-microbiota interactions have been extensively studied in inflammation,

metabolism, autoimmune diseases, and brain development.[10-13] Therefore, analyzing the PGN

structural composition and its chemical modifications is critical to understand the PGN

recognition processes and subsequent activation mechanisms in the immune system.[14]

Nowadays, analytical methods for peptidoglycan architecture include HPLC, LC-MS, solid-

state NMR, and AFM imaging, enabling a comprehensive study of PGN structure from chemical

composition to 3-D architecture.[5] Online ultra-performance liquid chromatography coupled with

electrospray ionization– tandem mass spectrometry (UPLC-ESI-MS/MS) is the primary strategy

for muropeptide structural identification, requiring minimal sample amounts and time.[15-17]

However, data processing, such as peak annotation and MS/MS spectra alignment, still relies on

the prior knowledge of experienced biological chemists. Previous work published by Bern et al.

demonstrated the analysis of muropeptides of *Clostridium difficile* with MS/MS fragmentation

spectra by Byonic peptide identification software.[18] Another innovative tool, PGFinder, created

by Patel et al., shows consistent and reproducible data analysis of Gram-negative bacterial PGN

with MS1 library searching.[19] Nevertheless, an automated analysis based on MS/MS

identification is certainly needed for a thorough PGN investigation of complex muropeptide

structures, particularly for gut microbial peptidoglycans.

Here, we developed an in-house, high-throughput automated muropeptide analysis

(HAMA) platform that simplifies muropeptide structures to sequence format, mimicking the

proteomic "bottom-up" approach. We also established a comprehensive *in silico* MS/MS

fragmentation database for muropeptide identification. Using this platform coupled with high-

resolution mass spectrometry, we revealed the peptidoglycan compositions of common gut

bacteria, including *Bifidobacterium, Bacteroides, Lactobacillus, Enterococcus,* and *Akkermansia*

muciiniphila.

MATERIAL AND METHODS

Bacterial strains and cell culture. Bacteria and their respective media used in this work were listed in Supplementary Table 1. Anaerobic bacteria such as *Bifidobacterium*, *Bacteroides, Lactobacillus, Enterococcus,* and *Akkermansia muciiniphila* were cultured anaerobically by using an anaerobic workstation (Whitley DG 250, Don Whitley Scientific Limited, England).

Peptidoglycan extraction and Mutanolysin Digestion. The extraction of peptidoglycan was performed using a previously described method, with some modifications.[9, 16] Ten milliliters of overnight cultures that have reached the stationary phase were harvested and lysed by 0.5% SDS solution in 0.1 M Tris/HCl (pH 7.0). After boiled for 30 min at 100 ℃ and thoroughly washed with H_2O three times, the pellets were resuspended in 750 mL of H_2O and sonicated for 30 min. The suspensions were spiked with 750 mL of 0.1 M Tris/HCl solution (pH 7.0) and treated with DNase and RNase at 37 ℃ in a shaker for 1 hr. After the removal of residual nucleic acids, the suspensions were subsequently treated with pronase to digest cell wall-bound proteins (final concentration of 1 mg/mL for 16 h at the same conditions). The pronase-treated cell walls were washed twice with H_2O before wall teichoic acids were released by 1 mL of 1 N HCl incubation for 4 h. Insoluble peptidoglycans were washed with H_2O until pH 5-6 and resuspended in 200-500 µL of 12.5 mM sodium dihydrogen-phosphate buffer (pH 5.5) to an OD_{600} of 3.0. One hundred microliters of insoluble peptidoglycan suspensions were digested overnight with 1 µL of mutanolysin solution (1.000 U/mL in H_2O). Following mutanolysin inactivation (100 ℃ boiling for 3 min) and centrifugation (5 min at 10,000 rpm), the supernatants were spiked with 50 µL of reduction solution (10 mg/mL sodium borohydride in 250 mM borate buffer at pH 9.0). After 20 min incubation at room temperature, the reduction reaction was stopped by adding 1 µL of

phosphoric acid (98%), resulting in pH 3-4. Reduced muropeptide samples were subsequently

analyzed by UPLC-MS/MS.

UPLC-MS/MS analysis of muropeptides. UPLC-MS/MS analysis was performed by a Dionex

UltiMate 3000 UHPLC system coupled to a Q-Exactive Plus hybrid quadrupole-orbitrap mass

spectrometer (Thermo Scientific). Soluble muropeptides were separated by an Acquity UPLC

CSH C18 column (130 Å, 1.7 μm, 2.1 mm × 100 mm), with a solvent A (0.1% formic acid in

water) and solvent B (0.1% formic acid in acetonitrile) as the mobile phase. A flow rate of 0.25

mL/min and 1% mobile phase B were applied in column condition. The injection volume of each

sample was 5 μL. An elution gradient was run for 25 min: starting with 1% solvent B for 0-2

min; 1-20% B (linear), 2-15 min, 20-95% B (linear), 15-17 min; 95% B, 17-19 min; 95-1% B

(linear), 19-21 min; 1% B, 21-25 min for re-equilibration. The column temperature was

controlled at 52 ℃ throughout the whole analysis program. A Heat electrospray ionization

(HESI) source was operated in positive mode, with parameters automatically optimized under a

flow rate of 250 μL/min (a capillary temperature of 300 ℃, a probe temperature of 300 ℃, and a

spray voltage of 3.5 kV). Data-dependent acquisition (DDA) mode was used in the instrument. A

mass range of MS1 acquisition was from 400 to 2,000 m/z at a resolution of 70,000 and the top 5

most abundant ions were subjected to higher-energy collision-induced dissociation (HCD)

fragmentation with Δm/z 3 isolation window, stepped normalized collision energy (NCE) at 20,

25, and 35 (A.U.), and dynamic exclusion time of 4 sec.

Data processing and data analysis. LC-MS/MS raw data was converted to an mzXML format

using MSConvert (ProteoWizard) and then processed by HAMA for mass deconvolution and

PGN identification.

Mutanolysin digestion assay. Purified peptidoglycan was adjusted to OD_{600} 1.0 in sodium

dihydrogen-phosphate buffer and digested with 5 µL of mutanolysin (1.000 U/mL in H_2O).

Absorbance was measured at OD_{600} of time points at 0, 3, 6, 9, and 16 hr in a Synergy H1

microplate reader (BioTek) with constant orbital shaking at 37 °C.

Immobilization of cultured bacteria for AFM imaging. The method of bacteria

immobilization was performed as described previously, with some modifications.[20] Overnight

cultured bacteria were washed twice and diluted to OD_{600} 0.3 with 1x PBS. Freshly cleaved Si

wafers were coated by 500 µL of 0.1% (w/v) poly-l-lysine in H_2O and left overnight at room

temperature. Afterward, the substrates were rinsed five times with ultrapure H_2O and dried with a

nitrogen flow. One hundred microliters of the bacteria suspension were dropped on a silicon

substrate, left for 10-20 min, and then rinsed three times in H_2O bath. The bacteria were dried

onto the substrate with flowing nitrogen and rehydrated again in 1x PBS bath for liquid imaging.

Atomic force microscopy (AFM) imaging. Cultured bacteria images were acquired in 1x PBS

bath at room temperature by AFM (Dimension Icon, Bruker) with ScanAsyst[TM]. Data were

collected by the PeakForce Quantitative NanoMechanics (QNM) mode with qp-CONT probes

(spring constant: 0.25 N/m, resonant frequency: 30 kHz, NANOSENSORS[TM]). Images were

acquired at a scan rate of 25 µm/s, an applied force of 500 pN, and with a resolution of 256 × 256

pixels per image frame. For mechanical analysis, the approaching part of the curves were fitted

with the Derjaguin-Muller-Toporov (DMT) model. DMT modulus of each bacterium was

recorded with 16 × 16 curves on top of the bacteria with the area of 250 × 250 nm^2.

RESULTS

HAMA platform: a High-throughput Automated Muropeptide Analysis for Identification

of PGN Fragments

The complex composition and structure of PGN make their structural analysis and

identification a time-consuming and challenging process. To address this, we developed a high-

throughput and automated platform for PGN structural analysis based on a proteomic "bottom-

up" approach. Our strategy is to simplify the structure of muropeptides into sequences by which

MS/MS patterns of *b*- and *y*- ions could be generated *in silico* for CID/HCD fragmentation

spectra matching. The overview of peptidoglycan analysis in this work is depicted in Fig. 1a.

Briefly, bacterial peptidoglycan was extracted, purified, and hydrolyzed by mutanolysin, and the

muropeptide products were then analyzed by UPLC-MS/MS. The automated identification of

these muropeptides was accomplished using the HAMA platform, a MATLAB-based software

with a user-friendly graphic user interface (Supplementary Fig. 1) that includes three parts:

DBuilder, *Analyzer*, and *Viewer*.

(i) ***DBuilder*** *for Species-specific muropeptide database construction.* In order to construct

muropeptide databases *in silico*, inspired by the well-developed glyco-proteomics analysis, we

named each residue in disaccharide peptide with a letter and simplified the muropeptide structure

into a sequence format. The representation letters of monosaccharides and amino acids are listed

in Supplementary Table 2. Dashes and parentheses were used to represent peptide/glycosidic

bonds and discriminate certain subunits, respectively. DBuilder combined the input options of

three parts (GlcNAc, MurNAc, and peptide) to generate a comprehensive database that contains

monomeric, dimeric, and trimeric muropeptides, as shown in Fig. 1b. Herein, we constructed

PGN multimers using two types of polymerization events: transglycosylation connected via

glycosidic bonds, and transpeptidation linked through peptide bonds. In DBuilder, the type of

peptide linkage used was the common 4-3 cross-link, which could be achieved either directly

through a covalent bond between the penultimate D-Ala and the third residue in the acceptor

peptide stem, or via an interpeptide bridge. We also considered that the terminal D-Ala (at the

fifth position) in the donor stem was not allowed during the transpeptidation reaction. In

addition, choosing more possible modifications (anhydro MurNAc residues, deacetyl GlcNAc

residues, acetyl MurNAc residues, and amidated iso-Glu) to construct a PGN database could lead

to severe mass coincidence issues for MS1 searching in the Analyzer. Hence, we set a maximum

modification number of six to avoid generating multi-modified muropeptide candidates. Finally,

the species-specific muropeptide database was outputted to a *.csv format file, including

muropeptide sequences and corresponding chemical formula and theoretical monoisotopic mass.

(ii) ***Analyzer*** *for analysis of muropeptide MS raw files.* Analyzer is an integrated tool for

processing MS data and identifying muropeptides. The flowchart is shown in Fig. 1c. First, the

species-specific database, MS raw file (mzXML format),[21] and parameters were loaded into

Analyzer. Parameters were originally set with MS1 range *m/z* 400-2000, retention time range 2-

12 min, mass tolerance of 10 ppm (orbitrap mass spectrometry data), intensity threshold of 1e5,

etc., which could be input by users for different experimental conditions. Then, the mzXML file

was processed sequentially by peak picking, deisotoping, feature selection, and deconvolution.

The observed masses were searched against the loaded database within 10 ppm tolerance.

However, in cases where the observed mass matched more than one inferred sequence in MS1

searching, Analyzer compared the MS/MS spectra of those MS1 matched features to *in silico*

MS/MS fragmentation spectra of corresponding inferred candidates through cosine similarity and

matched peak score (MPS= the number of the matched peaks/the number of the predicted peaks)

calculation. The observed mass was then identified as the sequence with the highest matching

score. Lastly, all identified muropeptides were merged into an Excel spreadsheet reporting

charge state, molecular weight, retention time, peak intensity, peak area, sequence, main scan

number, cosine similarity, score, etc. Analyzer also outputted a base peak chromatogram with

peak annotations and a result file (in Matlab data) for Viewer input. The entire analysis was

completed within a few minutes.

(iii) ***Viewer*** *for annotation and identification display.* Viewer, a visualization tool that

allows users to browse extracted ion chromatograms (XICs) of the identified muropeptides and

visualize the *in-silico* MS/MS matching spectra annotated with *b-* and *y-* ions. In the

muropeptide spectra match (PSM) page of each identified muropeptide, Viewer lists all inferred

candidates and visualizes their individual MS/MS spectral matches to clearly demonstrate the

process of scoring in Analyzer. Additionally, Viewer provides a data sorting function that allows

users to classify the identification list by molecular weight, peak area, score (in ascending or

descending order), and sequence (in alphabetical order).

Demonstration of the HAMA Platform Using Well-characterized PGNs of *E. coli* and *S.

aureus

As a proof-of-concept, we demonstrated the HAMA platform using well-characterized

peptidoglycans from Gram-negative *Escherichia coli* DH5α and Gram-positive *Staphylococcus*

aureus SA113. The typical PGN subunit of *E. coli* is GlcNAc – MurNAc – L-Ala – D-iGlu –

mDAP– D-Ala – D-Ala, represented as a sequence of B-M-l(-A-E-H-A-A) in the HAMA

platform. For *S. aureus*, the classic PGN structure is a disaccharide-pentapeptide with the Gly_5

interpeptide bridge branching from the lysine, which is simplified as B-M-l(-A-Q-(G-G-G-G-G-

)K-A-A. Based on the analyzed input dataset, approximately 70% of the peak area in the base

peak chromatograms was assigned to muropeptide signals, which allowed for a comprehensive

PGN mapping (Fig. 2a, c). The HAMA platform successfully identified *E. coli* and *S. aureus*

muropeptides, and their XICs and MS/MS matching spectra could be visualized in Viewer (Fig.

2b, d). In total, 28 and 89 unique muropeptides were identified in PGNs of *E. coli* and *S. aureus*,

respectively. Table 1 lists overviews of the main muropeptide peaks in the chromatograms. Upon

examining the entire identification results, we found that some muropeptides were eluted in

multiple retention times due to the existence of a few abundant stereoisomers. Additionally, a

few monosaccharide muropeptides identified as the loss of GlcNAc could be recognized as in-

source fragments by exact co-elution.

In the *E. coli* data, the most abundant muropeptides were disaccharide-tetrapeptide (Tetra

monomer) and disaccharide-tetrapeptide-disaccharide-tetrapeptide (Tetra-Tetra dimer). We also

identified two low-abundant muropeptides, B-M-l(-A-E-H-G) and B-M-l(-A-E-H-A-G), in

which the fourth and fifth Ala are individually substituted with a Gly. This unique composition

has been reported in strain *E. coli* Nissle 1917 from the previous analysis,[16] but not for DH5α

strain used in our study. In the output of *S. aureus* peptidoglycan, the elution profile was the

same as what has been previously known: the most abundant monomers were disaccharide-

pentapeptide with a $(Gly)_5$ bridge and disaccharide-pentapeptide with a Gly bridge. The most

abundant cross-linked dimer was disaccharide-tetrapeptide-disaccharide-pentapeptide, which

contains a total of ten Gly residues.[16] We also characterized the known modifications and

structural variations within *S. aureus* peptidoglycan, such as *O*-acetylation of MurNAc, the

presence of D-iGlu (non-amidated) in the stem peptides, and the length variation of the

interpeptide bridge. The high-throughput analysis allowed for the identification of monomeric

muropeptides consisting of one to nine Gly residues and dimeric muropeptides containing a total

of five to fourteen Gly residues in a single analysis. However, due to the insufficient structural

information in the peptide backbone provided by the HCD fragmentation spectra, the exact

number of Gly residues harbored in each PGN unit could not be determined. Hence, further

careful analysis and manual verification are required to confirm the identity of identified

muropeptides, particularly for low-score dimers and trimers. Collectively, these results provide

valuable insights into the PGN compositions and architectures.

Characterizing Gut Bacterial PGN Compositions and Resolving Isomeric Muropeptides

Over the past two decades, an increasing number of gut microbial species have been found

to be associated with human health. In addition, emerging evidence has supported the notion that

gut microbial muropeptides work as signaling molecules that mediate host−microbiome

interactions in metabolism, gut homeostasis, and immunity.[22] However, little has been discussed

about the structures of PGN fragments due to the diversity and complexity of gut bacterial PGNs

and the lack of an efficient analytical tool.[14] To date, most of the structural information on gut

bacterial peptidoglycan comes from reports published between 1970 to 2000,[23-25] whereas that of

the health-promoting gut microbes discovered in the last two decades have been rarely reported.

Therefore, we utilized the HAMA platform to investigate the specific PGN structures of several

common gut bacterial species, including *Bifidobacterium, Bacteroides, Lactobacillus,*

Enterococcus, and *Akkermansia muciiniphila*. We collected LC-MS/MS data of purified

peptidoglycans from ten gut bacterial species and identified those with the self-built species-

specific muropeptide databases. Overall, the base peak chromatograms of each species showed

approximately 70% of peaks were annotated as muropeptides, and these elution profiles reflected

the actual composition and organization within the peptidoglycan architecture. Additionally, the

structural compositions of these purified peptidoglycans are consistent with previous reports,

listed in Table 2. The muropeptide profiles of the species in this study are summarized in

Supplementary Table 3-12.

In the identification results of Gram-positive gut bacterial peptidoglycans, the main

muropeptide sequence is GlcNAc – MurNAc – L-Ala – D-iGln – L-Lys– D-Ala – D-Ala, in

which the second amino acid of the stem peptide is usually amidated to D-iGln from D-iGlu.

This chemical modification is done by the MurT or GatD biosynthetic enzyme and is supposed to

control the PGN cross-linking levels, which has recently been demonstrated using labeled PGN

stem mimics in certain species.[26] Another unique feature is L-ornithine (L-Orn) at the third

position of the stem peptide. Unlike L-lysine, which is a usual feature of Gram-positive

peptidoglycans, L-Orn is a non-proteinogenic amino acid and has been found in the

peptidoglycan of certain *Bifidobacterium* species before.[23] As anticipated, we identified the L-

Orn-harbored muropeptides in *B. bifidum* and *B. longum* peptidoglycans. The most species-

specific variation among Bifidobacterial PGN was the architecture of the interpeptide bridge,

such as Gly, Ser-Asp, and Ser-Ala-Thr-Ala, which corresponded to *B. breve, B. bifidum,* and *B.*

longum species. This structural diversity arouses our interest in the relationship between the

length of an interpeptide bridge and the physical property of the bacterial cell envelope, which

will be discussed in a later section.

In the peptidoglycans of *E. faecium* and *L. acidophilus* (Supplementary Table 4-5)*,* we

observed the monomeric structure of GlcNAc – MurNAc – L-Ala – D-iGln – L-Lys– D-Ala – D-

Ala, which harbored an interpeptide bridge of asparagine (D-Asn) or aspartate (D-Asp),

shortened as B-M-l(-A-Q-(N-)K-A-A) or B-M-l(-A-Q-(D-)K-A-A). In this part, we found that

both B-M-l(-A-E-(N-)K) and B-M-l(-A-Q-(D-)K) had identical molecular weights since the mass

difference (+ 0.984 Da) between Asn (N) and Asp (D) is the same as between Gln (Q) and Glu

(E). This kind of isomeric muropeptides makes identification more complicated, but it can still

be addressed by MS/MS *in silico* fragmentation matching under an appropriate separation

chromatography. For example, two disaccharide-tripeptides separately eluted at 6.18 min and

7.02 min were found as structural isomers that existed in *E. faecium* peptidoglycan (Fig. 3 and

Supplementary Table 4). Through *in silico* MS/MS fragmentation matching, these two isomers

can be discriminated and identified as the sequences of B-M-l(-A-E-(N-)K) and B-M-l(-A-Q-(D-

)K), respectively. To validate the correctness of the automated platform, we also extracted the

experimental data of those isomers and manually inspected the MS/MS spectra (Supplementary

Fig. 2). However, this strategy did not work for the identification of *E. faecalis* peptidoglycan

whose interpeptide bridge is composed of two alanine residues (Supplementary Table 3). In this

case, two structurally isomeric muropeptides, a disaccharide-tripeptide with a bridge (B-M-l(-A-

Q-(A-A-)K)) and a disaccharide-pentapeptide (B-M-l(-A-Q-K-A-A)), have similar *in silico*

MS/MS fragmentation patterns, which often leads to misidentification. Nevertheless, the

retention time still gave us information to identify them. We annotated the peak at a retention

time of 3.66 min as disaccharide-pentapeptide (B-M-l(-A-Q-K-A-A)) by comparing it to an

identical sequence that appeared in *L. acidophilus* peptidoglycan with a similar retention time

(3.61 min). Therefore, the other peak at a retention time of 6.39 min could be determined as

disaccharide-tripeptide with a bridge (B-M-l(-A-Q-(A-A-)K)), which had a longer retention time

than the linear structure one.

Apart from Gram-positive bacteria, we also analyzed the peptidoglycan of several anaerobic

Gram-negative bacteria, including *Bacteroides fragilis*, *Bacteroides ovatus*, *Bacteroides*

thetaiotaomicron, and *Akkermansia muciniphila* (Table 2 and Supplementary Table 9-12).

Structurally, the general stem peptide of Gram-negative peptidoglycan is L-Ala – D-iGlu – m-

DAP– D-Ala – D-Ala, with diaminopimelic acid in the third position being a representative

feature. The peptidoglycan structures we identified were consistent with previously published

ones, with *O*-acetyl-MurNAc found in *Bacteroides* species and *N*-deacetyl-GlcNAc (GlcN) in

Akkermansia muciniphila.[27, 28] The output analysis showed that around 56-66% of the total

muropeptides in *Bacteroides* species' peptidoglycans contained *O*-acetylated MurNAc, while

approximately 87% of the total muropeptides in *A. muciniphila* peptidoglycan contained de-*N*-

acetylated GlcNAc. The high occurrence of *N*-deacetylation in *A. muciniphila* peptidoglycan

suggests that *A. muciniphila* might possess a homolog of oxidative stress-induced PGN

deacetylase (PgdA) found in *Helicobacter pylori*.[29-31] Chemical modifications to the disaccharide

backbone are known to provide resistance to lysozyme and protect bacteria against enzymatic

attack from the host innate immune system.[32, 33]

Inferring PGN Cross-linking Types Based on Identified PGN Fragments

We not only identified peptidoglycan structures but also were dedicated to exploring the

type of peptide cross-linking within those gut bacterial peptidoglycan. The substrates and

catalyzed enzymes involved in peptide cross-linking have been targets for antibiotic development

and antimicrobial resistance studies.[34] In general, there are two types of PGN cross-linkage: 4-3

cross-links generated by D,D-transpeptidases (Ddts) and 3-3 cross-links created by L,D-

transpeptidases (Ldts) (see Fig. 4a). As the enzyme names imply, the substrates that Ddts and

Ldts bind to are terminated as D,D-stereocenters and L,D-stereocenters, which structurally

means pentapeptides and tetrapeptides. During D,D-transpeptidation, Ddts first remove the

terminal (fifth) D-Ala residue of the pentapeptide stem and form an intermediate. Then, the

intermediate cross-links the NH_2 group in the third position of the neighboring acceptor stem,

forming a 4-3 cross-link. Following this distinctive rule of PGN biosynthesis, the possible

combinations of 4-3 cross-linked dimers (Donor-Acceptor) are disaccharide-tetrapeptide–

disaccharide-pentapeptide (D45) and disaccharide-tetrapeptide–disaccharide-tetrapeptide (D44).

During L,D-transpeptidation, Ldts cleave the terminal (fourth) D-Ala residue of the donor

tetrapeptide stem and generate a peptide bond between the third residue and the NH_2 group in the

third position of adjacent peptide.[35] The possible structures of 3-3 cross-linked dimers are

disaccharide-tripeptide–disaccharide-tetrapeptide   (D34)   and   disaccharide-tetrapeptide–

disaccharide-tripeptide (D33). Hence, we could infer the possible PGN cross-linkage types and

the involved enzymes based on the main PGN fragments.

Take *Bifidobacterium breve* ATCC 15700, CSCC 1900, and ATCC 15698 as examples. The

LC-MS profiles showed noticeable differences in muropeptide compositions among the three

strains (Fig. 4b). The relative muropeptide compositions of each strain are presented in a

heatmap (Fig. 4c). Obviously, the main muropeptides of *B. breve* ATCC 15700 are disaccharide-

tripeptide with an interpeptide bridge (M3b), disaccharide-tetrapeptide with an interpeptide

bridge (M4b), and disaccharide-tripeptide–disaccharide-tetrapeptide (D34), whereas the CSCC

1900 strain showed great abundances in disaccharide-tetrapeptide with an interpeptide bridge

(M4b), disaccharide-pentapeptide with an interpeptide bridge (M5b), and disaccharide-

tetrapeptide–disaccharide-tetrapeptide (D44). These results suggested that 3-3 cross-links and 4-

3 cross-links might be predominant in the peptidoglycans of ATCC 15700 and CSCC 1900

strains, respectively. In the case of ATCC 15698 strain, its peptidoglycan likely contains both

types of cross-links since the abundances of M3b and M4b, as well as D34 and D44, are almost

equivalent.

Broad-spectrum β-Lactams are known to inhibit D,D-transpeptidases, such as penicillin-

binding proteins (PBPs). However, L,D-transpeptidases are generally insensitive to β-Lactams

and offer alternative cross-links in the PGNs.[36] Numerous studies have investigated the role of

L,D-transpeptidases in the maintenance and remodeling of mature peptidoglycan in organisms

such as *Enterococcus faecium, Clostridium difficile, Escherichia coli,* and *Mycobacterium*

tuberculosis.[37-42] Based on the muropeptide compositional analysis mentioned above, we found

high abundances of M3/M3b monomer and D34 dimer in the peptidoglycans of *E. faecalis*, *E.*

faecium, L. acidophilus, B. breve, B. longum, and *A. muciniphila*, which may be the PGN

products catalyzed by Ldts. While the homologs of Ldts in *L. acidophilus, B. breve, B. longum,*

and *A. muciniphila* have been supported by genome sequence evidence,[43-46] biochemical

evidence is needed to confirm the existence of L,D-transpeptidases in those species.

Proposed Bridge Length-dependent Cell Envelope Stiffness in *B. longum* and *B. breve*

Peptidoglycan, a protective exoskeleton around cells, provides structural integrity to the

cell. The porosity of the PGN scaffold, which is defined by the degree of cross-link, might affect

the delivery and diffusion of drugs and signaling molecules to the cell membrane. Thus,

modifications to PGN structure are thought to significantly influence bacterial cell mechanics.[47,]

[48] Previous evidence has indicated that a high PGN cross-linking level enhances the stiffness of

the cell wall material in Gram-positive bacteria.[49] The PGN-lattice architecture based on the

interpeptide bridge length has been investigated using solid-state NMR in *S. aureus*.[50] However,

the effect of the interpeptide bridge length variants on the porosity and stiffness of bacterial cell

envelope has little been discussed. Interestingly, the identification table presented above showed

that *Bifidobacterium* peptidoglycans have different architecture of the interpeptide bridges

among species: a tetrapeptide bridge (Ser-Ala-Thr-Ala) found in *Bifidobacterium longum* and a

monopeptide bridge (Gly) found in *Bifidobacterium breve* (Supplementary Fig. 3). We wondered

whether the length of interpeptide bridges may be related to the bacterial cell envelope's

mechanical properties and hypothesized that the cross-linking with shorter bridges may form a

tighter meshwork in peptidoglycan layers (Fig. 5a).

In order to explore the above relevance, we selected three *B. breve* strains and three *B.*

longum strains, whose cell morphology is similar, as research objects. The proportion of

muropeptide composition of these species is summarized in Fig. 5b. A slightly higher abundance

of peptide-linked dimers in *B. breve* than in *B. longum* implies a higher degree of PGN cross-

linking level in *B. breve*. Next, we directly measured the stiffness (defined by Young's modulus,

E) of these two species using the single-cell atomic force microscopy technique (AFM).

Representative data obtained on the different strains are presented in Fig. 5c. For each cell, a

high-resolution elasticity map was recorded on top of the cell in the central region to avoid edge

effects. The values of Young's modulus were extracted and subjected to statistical analysis,

shown in Fig. 5d. Overall for the cell envelope properties, *B. longum* displayed smaller values of

stiffness than *B. breve*, demonstrating softer cell envelope properties with one-fifth lower values

of Young's modulus. As anticipated, these data suggested the *B. breve* harboring shorter

interpeptide bridges corresponded to a higher cell stiffness. Furthermore, a slower hydrolysis rate

was observed in purified PGN of *B. breve* than in that of *B. longum*, implying that the cell wall

of *B. breve* is a harder exoskeleton (Supplementary Fig. 4). Interestingly, among three *B. breve*

strains, the average stiffness of ATCC 15700 is higher than the other strains, likely due to the

more abundant 3-3 cross-linkages in its peptidoglycan. Computational modeling suggested a

more rigid stem peptide in the conformation of L,D-cross-linking, implying the 3-3 cross-

linkages strengthen the PNG layer.[36, 51] These results might support our implication that a tight

peptidoglycan network woven with shorter peptides gives the bacterium a stiffer cell wall,

proposed as an idea of bridge length-dependent cell envelope.

DISCUSSION

The purpose of this study was to develop an automated platform for identifying and analyzing the muropeptide compositions of gut bacterial PGNs. Using high-resolution MS data, we characterized PGN structures of common gut microbes, investigated their PGN cross-linking types, and evaluated the effect of bridge-length variants on the stiffness of bacterial cell walls.

In early 1970s, the amino acid compositions of PGN stem peptides were characterized in many bacteria species, including certain gut microorganisms. At that time, PGNs were purified and hydrolyzed under harsh acidic conditions and then separated by two-dimensional paper chromatography to determine the amino acid sequence. Based on this characterization work, Schleifer and Kandler proposed a bacterial PGN classification system that is still used today.[23] In the *Bifidobacterium* and *Lactobacillu*s genera, only type A PGNs were identified, representing 4-3 cross-linking through interpeptide bridges.[25] However, our study of *B. breve*, *B. longum*, and *L. acidophilus* revealed a significant abundance of M3b and D34 in their muropeptide compositions, implying the incorporation of 3-3 cross-linkage type within PGN network. This cross-linking type is catalyzed by penicillin-insensitive L,D-transpeptidases. We speculate that this finding may be due to the comprehensive mass spectrometric approaches we used. Furthermore, we adopted the enzymatic method using mutanolysin to cleave the β-N-acetylmuramyl-(1,4)-N-acetylglucosamine linkage, which is crucial for obtaining the complete PGN subunit that preserves the original peptide linkage. Importantly, our HAMA platform provides a powerful tool for mapping peptidoglycan architecture and generating structural information on the PGN biosynthesis system, which is closely related to the antibiotic resistance of bacteria.

In most Gram-positive bacteria, the interpeptide bridges vary depending on the species and

are likely associated with the PGN architectures. Although the chemical structure of PGN units

has been extensively studied, the physical structure and 3-D architecture remain open

questions.[52, 53] A previous review by Kim et al. suggested that the length of interpeptide bridges

is a key factor in determining different PGN architectures and proposed three PGN-bridge

length-dependent architectures based on the orientations of cross-linked peptide stems.[50] The

parallel-stem model, which has the smallest pore size and the highest cross-linking level, is

suited for bacteria with long-length bridges. The perpendicular-stem model, with intermediate

pore size, is proposed for bacteria with intermediate-length bridges, while the antiparallel-stem

model, which has the largest pore size, is for bacteria without bridges. Notably, our study

provides observations regarding the relationship between the bridge length and the stiffness of

bacterial cell walls in *Bifidobacterium* species (Fig. 5). *B. longum*, which harbors tetrapeptide

bridges (Ser-Ala-Thr-Ala) corresponds to lower stiffness, whereas *B. breve,* whose PGN is cross-

linked by monopeptide bridges (Gly), has higher stiffness. This finding implies that the PGN

architecture with shorter cross-linked bridges corresponds to a more rigid skeleton. This is not

surprising as *Bifidobacterium* species are known for their thick peptidoglycan layers, which

account for 30-70% of the cell wall.[35] Meanwhile, other biomolecules in the cell wall also

influence bacterial stiffness, such as lipoprotein, wall teichoic acids, and lipoteichoic acids.[54-57]

Future studies on the effect of quantitatively changing the PGN-bridge length on stiffness will

require intensive research, considering the cross-linking densities, glycan strand lengths, and

PGN architecture models. These studies may provide further insight into the physical structure

and architecture of PGN, which would contribute to a better understanding of the factors

determining bacterial cell wall properties.

In the HAMA platform, simplifying muropeptide structures into sequences has facilitated

the generation of *in silico* MS/MS fragments for spectra matching, enabling the structural

resolving of isomeric muropeptides. However, there are several limitations to our study design.

Our species-specific PGN database is built based on known structures and modifications reported

previously. While the identified muropeptides cover approximately 70% of peak area in the base

peak chromatograms, certain low-abundant muropeptides composed of uncoded amino acids or

saccharides required additional MS/MS identification with manual analysis. For peptide-linked

multimers, the current DBuilder only builds up 4-3 cross-linked sequences because including the

3-3 cross-link type results in severe misidentifications, as it produces similar *in silico* MS/MS

fragmentation patterns as the 4-3 cross-links (about 88-90% similarity). However, we can infer

possible PGN cross-linkages based on the type of PGN fragments obtained from hydrolysis.[58]

The primary limitation is that MS/MS spectra generated by HCD fragmentation carry mostly

structural information from disaccharide moiety (glyco-oxonium ions) without providing

sufficient peptide fragment ions to derive peptide sequence,[59] making it difficult to accurately

identify multimers and determine the position of modifications. Nevertheless, accurate

monoisotopic masses acquired from high-resolution mass spectrometry are still informative in

peptidoglycan compositional components. Alternative approaches can be considered to elucidate

branched structures of muropeptides, such as breaking down PGN with cell wall amidases

(cleaving the amide bond between MurNAc and L-Ala residue) to target stem peptide analysis,[60,

61] acquiring MS^3 spectra to obtain fragmentation maps, or using electron transfer dissociation

(ETD) fragmentation mode to leave more peptide fragments.[18] Additionally, strategies

commonly used in glycopeptide identification, such as multi-fragmentation modes (CID, HCD,

and ETD), could also be applied to PGN identification to provide more information on peptide

sequences and disaccharide structures.[62] To achieve a more comprehensive identification of

muropeptides, future maneuvers are proposed for an expanded database, innovative *in-silico*

fragmentation patterns, and improved MS/MS spectra acquisition.

HAMA is an innovative automated platform that constructs species-specific muropeptide

databases, validates structural identification using *in-silico* MS/MS analysis, and provides

visualized results for efficient and reliable identification of muropeptides, greatly reducing the

time-consuming task of manual interpretation of LC-MS/MS data. The HAMA platform has the

potential to be a valuable tool for various research fields, including microbiology, pathology,

molecular biology, and immunology. Its applications in identifying activation ligands for

antimicrobial resistance studies, characterizing key motifs recognized by pattern recognition

receptors for host-microbiota immuno-interaction research, and mapping peptidoglycan in cell

wall architecture studies. With the ease and efficiency offered by HAMA, we believe that

muropeptide analysis will become more accessible and contribute to a deeper understanding of

cell wall biology.

REFERENCES

1. Silhavy, T. J.; Kahne, D.; Walker, S., The bacterial cell envelope. *Cold Spring Harbor Perspect. Biol.* **2010,** *2* (5), a000414.

2. Vollmer, W.; Blanot, D.; de Pedro, M. A., Peptidoglycan structure and architecture. *FEMS Microbiol Rev* **2008,** *32* (2), 149–167.

3. Irazoki, O.; Hernandez, S. B.; Cava, F., Peptidoglycan Muropeptides: Release, Perception, and Functions as Signaling Molecules. *Front Microbiol* **2019,** *10*, 500.

4. Schumann, P., Peptidoglycan structure. In *Methods in microbiology*, Elsevier: 2011; Vol. 38, pp 101-129.

5. Porfírio, S.; Carlson, R. W.; Azadi, P., Elucidating Peptidoglycan Structure: An Analytical Toolset: (Trends in Microbiology 27, 607-622; 2019). *Trends Microbiol* **2019,** *27* (7), 653-654.

6. Wolf, A. J.; Underhill, D. M., Peptidoglycan recognition by the innate immune system. *Nat Rev Immunol* **2018,** *18* (4), 243-254.

7. Nikolaidis, I.; Favini-Stabile, S.; Dessen, A., Resistance to antibiotics targeted to the bacterial cell wall. *Protein Sci* **2014,** *23* (3), 243-259.

8. Girardin, S. E.; Travassos, L. H.; Hervé, M.; Blanot, D.; Boneca, I. G.; Philpott, D. J.; Sansonetti, P. J.; Mengin-Lecreulx, D., Peptidoglycan molecular requirements allowing detection by Nod1 and Nod2. *J Biol Chem* **2003,** *278* (43), 41702-41708.

9. Huang, Y. W.; Wang, Y.; Lin, Y.; Lin, C.; Lin, Y. T.; Hsu, C. C.; Yang, T. C., Impacts of Penicillin Binding Protein 2 Inactivation on β-Lactamase Expression and Muropeptide Profile in Stenotrophomonas maltophilia. *mSystems* **2017,** *2* (4), e00077-00017.

10. Cavallari, J. F.; Fullerton, M. D.; Duggan, B. M.; Foley, K. P.; Denou, E.; Smith, B.

K.; Desjardins, E. M.; Henriksbo, B. D.; Kim, K. J.; Tuinema, B. R.; Stearns, J. C.;

Prescott, D.; Rosenstiel, P.; Coombes, B. K.; Steinberg, G. R.; Schertzer, J. D.,

Muramyl Dipeptide-Based Postbiotics Mitigate Obesity-Induced Insulin Resistance via

IRF4. *Cell Metab* **2017,** *25* (5), 1063-1074.

11.   Inohara, N.; Ogura, Y.; Fontalba, A.; Gutierrez, O.; Pons, F.; Crespo, J.; Fukase, K.;

Inamura, S.; Kusumoto, S.; Hashimoto, M.; Foster, S. J.; Moran, A. P.; Fernandez-

Luna, J. L.; Nuñez, G., Host recognition of bacterial muramyl dipeptide mediated through

NOD2. Implications for Crohn's disease. *J Biol Chem* **2003,** *278* (8), 5509-5512.

12.   Huang, Z.; Wang, J.; Xu, X.; Wang, H.; Qiao, Y.; Chu, W. C.; Xu, S.; Chai, L.;

Cottier, F.; Pavelka, N.; Oosting, M.; Joosten, L. A. B.; Netea, M.; Ng, C. Y. L.;

Leong, K. P.; Kundu, P.; Lam, K. P.; Pettersson, S.; Wang, Y., Antibody neutralization

of microbiota-derived circulating peptidoglycan dampens inflammation and ameliorates

autoimmunity. *Nat Microbiol* **2019,** *4* (5), 766-773.

13.   Arentsen, T.; Qian, Y.; Gkotzis, S.; Femenia, T.; Wang, T.; Udekwu, K.; Forssberg,

H.; Diaz Heijtz, R., The bacterial peptidoglycan-sensing molecule Pglyrp2 modulates

brain development and behavior. *Mol Psychiatry* **2017,** *22* (2), 257-266.

14.   Bersch, K. L.; DeMeester, K. E.; Zagani, R.; Chen, S.; Wodzanowski, K. A.; Liu, S.;

Mashayekh, S.; Reinecker, H. C.; Grimes, C. L., Bacterial Peptidoglycan Fragments

Differentially Regulate Innate Immune Signaling. *ACS Cent Sci* **2021,** *7* (4), 688-696.

15.   Alvarez, L.; Hernandez, S. B.; de Pedro, M. A.; Cava, F., Ultra-Sensitive, High-

Resolution Liquid Chromatography Methods for the High-Throughput Quantitative

Analysis of Bacterial Cell Wall Chemistry and Structure. *Methods Mol Biol* **2016,** *1440*,

11-27.

16.   Kühner, D.; Stahl, M.; Demircioglu, D. D.; Bertsche, U., From cells to muropeptide

structures in 24 h: peptidoglycan mapping by UPLC-MS. *Sci Rep* **2014,** *4*, 7494.

17.   Glauner, B., Separation and quantification of muropeptides with high-performance liquid

chromatography. *Anal Biochem* **1988,** *172* (2), 451-464.

18.   Bern, M.; Beniston, R.; Mesnage, S., Towards an automated analysis of bacterial

peptidoglycan structure. *Anal Bioanal Chem* **2017,** *409* (2), 551-560.

19.   Patel, A. V.; Turner, R. D.; Rifflet, A.; Acosta-Martin, A. E.; Nichols, A.; Awad, M.

M.; Lyras, D.; Gomperts Boneca, I.; Bern, M.; Collins, M. O.; Mesnage, S., PGFinder,

a novel analysis pipeline for the consistent, reproducible, and high-resolution structural

analysis of bacterial peptidoglycans. *Elife* **2021,** *10*, e70597.

20.   Pasquina-Lemonche, L.; Burns, J.; Turner, R. D.; Kumar, S.; Tank, R.; Mullin, N.;

Wilson, J. S.; Chakrabarti, B.; Bullough, P. A.; Foster, S. J.; Hobbs, J. K., The

architecture of the Gram-positive bacterial cell wall. *Nature* **2020,** *582* (7811), 294-297.

21.   Chambers, M. C.; Maclean, B.; Burke, R.; Amodei, D.; Ruderman, D. L.; Neumann,

S.; Gatto, L.; Fischer, B.; Pratt, B.; Egertson, J.; Hoff, K.; Kessner, D.; Tasman, N.;

Shulman, N.; Frewen, B.; Baker, T. A.; Brusniak, M.-Y.; Paulse, C.; Creasy, D.;

Flashner, L.; Kani, K.; Moulding, C.; Seymour, S. L.; Nuwaysir, L. M.; Lefebvre, B.;

Kuhlmann, F.; Roark, J.; Rainer, P.; Detlev, S.; Hemenway, T.; Huhmer, A.;

Langridge, J.; Connolly, B.; Chadick, T.; Holly, K.; Eckels, J.; Deutsch, E. W.;

Moritz, R. L.; Katz, J. E.; Agus, D. B.; MacCoss, M.; Tabb, D. L.; Mallick, P., A cross-

platform toolkit for mass spectrometry and proteomics. *Nat. Biotechnol.* **2012,** *30* (10),

918-920.

22.   Bastos, P. A. D.; Wheeler, R.; Boneca, I. G., Uptake, recognition and responses to

peptidoglycan in the mammalian host. *FEMS Microbiol. Rev.* **2020,** *45* (1), fuaa044.

23. Schleifer, K. H.; Kandler, O., Peptidoglycan types of bacterial cell walls and their taxonomic implications. *Bacteriol Rev* **1972,** *36* (4), 407-477.

24. Veerkamp, J. H., The structure of the cell wall peptidoglycan of Bifidobacterium bifidum var. pennsylvanicus. *Arch. Biochem. Biophys.* **1971,** *143* (1), 204-211.

25. Pyclik, M.; Srutkova, D.; Schwarzer, M.; Górska, S., Bifidobacteria cell wall-derived exo-polysaccharides, lipoteichoic acids, peptidoglycans, polar lipids and proteins - their chemical structure and biological attributes. *Int J Biol Macromol* **2020,** *147*, 333-349.

26. Pidgeon, S. E.; Apostolos, A. J.; Nelson, J. M.; Shaku, M.; Rimal, B.; Islam, M. N.; Crick, D. C.; Kim, S. J.; Pavelka, M. S.; Kana, B. D.; Pires, M. M., L,D-Transpeptidase Specific Probe Reveals Spatial Activity of Peptidoglycan Cross-Linking. *ACS Chem Biol* **2019,** *14* (10), 2185-2196.

27. Weadge, J. T.; Clarke, A. J., Identification and characterization of O-acetylpeptidoglycan esterase: a novel enzyme discovered in Neisseria gonorrhoeae. *Biochemistry* **2006,** *45* (3), 839-851.

28. Garcia-Vello, P.; Tytgat, H. L. P.; Gray, J.; Elzinga, J.; Di Lorenzo, F.; Biboy, J.; Vollmer, D.; De Castro, C.; Vollmer, W.; de Vos, W. M.; Molinaro, A., Peptidoglycan from Akkermansia muciniphila MucT: chemical structure and immunostimulatory properties of muropeptides. *Glycobiology* **2022,** *32* (8), 712-719.

29. Moynihan, P. J.; Sychantha, D.; Clarke, A. J., Chemical biology of peptidoglycan acetylation and deacetylation. *Bioorg Chem* **2014,** *54*, 44-50.

30. Wang, G.; Olczak, A.; Forsberg, L. S.; Maier, R. J., Oxidative stress-induced peptidoglycan deacetylase in Helicobacter pylori. *J Biol Chem* **2009,** *284* (11), 6790-

583  6800.

584 31. Wang, G.; Maier, S. E.; Lo, L. F.; Maier, G.; Dosi, S.; Maier, R. J., Peptidoglycan

585  deacetylation in Helicobacter pylori contributes to bacterial survival by mitigating host

586  immune responses. *Infect Immun* **2010,** *78* (11), 4660-4666.

587 32. Davis, K. M.; Weiser, J. N., Modifications to the peptidoglycan backbone help bacteria to

588  establish infection. *Infect Immun* **2011,** *79* (2), 562-570.

589 33. Sychantha, D.; Brott, A. S.; Jones, C. S.; Clarke, A. J., Mechanistic Pathways for

590  Peptidoglycan O-Acetylation and De-O-Acetylation. *Front. Microbiol.* **2018,** *9*, 2332.

591 34. Srisuknimit, V.; Qiao, Y.; Schaefer, K.; Kahne, D.; Walker, S., Peptidoglycan Cross-

592  Linking Preferences of Staphylococcus aureus Penicillin-Binding Proteins Have

593  Implications for Treating MRSA Infections. *J Am Chem Soc* **2017,** *139* (29), 9791-9794.

594 35. Peltier, J.; Courtin, P.; El Meouche, I.; Lemée, L.; Chapot-Chartier, M. P.; Pons, J. L.,

595  Clostridium difficile has an original peptidoglycan structure with a high level of N-

596  acetylglucosamine deacetylation and mainly 3-3 cross-links. *J Biol Chem* **2011,** *286* (33),

597  29053-29062.

598 36. Aliashkevich, A.; Cava, F., LD-transpeptidases: the great unknown among the

599  peptidoglycan cross-linkers. *Febs j* **2022,** *289* (16), 4718-4730.

600 37. Mainardi, J. L.; Morel, V.; Fourgeaud, M.; Cremniter, J.; Blanot, D.; Legrand, R.;

601  Frehel, C.; Arthur, M.; Van Heijenoort, J.; Gutmann, L., Balance between two

602  transpeptidation mechanisms determines the expression of beta-lactam resistance in

603  Enterococcus faecium. *J Biol Chem* **2002,** *277* (39), 35801-35807.

604 38. Lecoq, L.; Dubée, V.; Triboulet, S.; Bougault, C.; Hugonnet, J. E.; Arthur, M.;

605  Simorre, J. P., Structure of Enterococcus faeciuml,d-transpeptidase acylated by ertapenem

provides insight into the inactivation mechanism. *ACS Chem Biol* **2013,** *8* (6), 1140-1146.

39.   Magnet, S.; Arbeloa, A.; Mainardi, J. L.; Hugonnet, J. E.; Fourgeaud, M.; Dubost, L.; Marie, A.; Delfosse, V.; Mayer, C.; Rice, L. B.; Arthur, M., Specificity of L,D-transpeptidases from gram-positive bacteria producing different peptidoglycan chemotypes. *J Biol Chem* **2007,** *282* (18), 13151-13159.

40.   Sütterlin, L.; Edoo, Z.; Hugonnet, J. E.; Mainardi, J. L.; Arthur, M., Peptidoglycan Cross-Linking Activity of l,d-Transpeptidases from Clostridium difficile and Inactivation of These Enzymes by β-Lactams. *Antimicrob Agents Chemother* **2018,** *62* (1), e01607-01617.

41.   Magnet, S.; Dubost, L.; Marie, A.; Arthur, M.; Gutmann, L., Identification of the L,D-transpeptidases for peptidoglycan cross-linking in Escherichia coli. *J Bacteriol* **2008,** *190* (13), 4782-4785.

42.   Lavollay, M.; Arthur, M.; Fourgeaud, M.; Dubost, L.; Marie, A.; Veziris, N.; Blanot, D.; Gutmann, L.; Mainardi, J. L., The peptidoglycan of stationary-phase Mycobacterium tuberculosis predominantly contains cross-links generated by L,D-transpeptidation. *J Bacteriol* **2008,** *190* (12), 4360-4366.

43.   Jeon, S.; Kim, H.; Choi, Y.; Cho, S.; Seo, M.; Kim, H., Complete Genome Sequence of the Newly Developed Lactobacillus acidophilus Strain With Improved Thermal Adaptability. *Front Microbiol* **2021,** *12*, 697351.

44.   Morita, H.; Toh, H.; Oshima, K.; Nakano, A.; Omori, E.; Hattori, Y.; Arakawa, K.; Suda, W.; Honda, K.; Hattori, M., Complete genome sequence of Bifidobacterium breve JCM 1192(T) isolated from infant feces. *J Biotechnol* **2015,** *210*, 81-82.

45.   Poyet, M.; Groussin, M.; Gibbons, S. M.; Avila-Pacheco, J.; Jiang, X.; Kearney, S. M.;

Perrotta, A. R.; Berdy, B.; Zhao, S.; Lieberman, T. D.; Swanson, P. K.; Smith, M.; Roesemann, S.; Alexander, J. E.; Rich, S. A.; Livny, J.; Vlamakis, H.; Clish, C.; Bullock, K.; Deik, A.; Scott, J.; Pierce, K. A.; Xavier, R. J.; Alm, E. J., A library of human gut bacterial isolates paired with longitudinal multiomics data enables mechanistic microbiome research. *Nat Med* **2019,** *25* (9), 1442-1452.

46. Ottman, N. A. Host immunostimulation and substrate utilization of the gut symbiont Akkermansia muciniphila. PhD thesis, Wageningen University, Wageningen, 2015.

47. Auer, G. K.; Weibel, D. B., Bacterial Cell Mechanics. *Biochemistry* **2017,** *56* (29), 3710-3724.

48. Chang, J. D.; Wallace, A. G.; Foster, E. E.; Kim, S. J., Peptidoglycan Compositional Analysis of Enterococcus faecalis Biofilm by Stable Isotope Labeling by Amino Acids in a Bacterial Culture. *Biochemistry* **2018,** *57* (7), 1274-1283.

49. Loskill, P.; Pereira, P. M.; Jung, P.; Bischoff, M.; Herrmann, M.; Pinho, M. G.; Jacobs, K., Reduction of the peptidoglycan crosslinking causes a decrease in stiffness of the Staphylococcus aureus cell envelope. *Biophys J* **2014,** *107* (5), 1082-1089.

50. Kim, S. J.; Chang, J.; Singh, M., Peptidoglycan architecture of Gram-positive bacteria by solid-state NMR. *Biochim Biophys Acta* **2015,** *1848* (1 Pt B), 350-362.

51. de Pedro, M. A.; Cava, F., Structural constraints and dynamics of bacterial cell wall architecture. *Front Microbiol* **2015,** *6*, 449.

52. Vollmer, W.; Seligman, S. J., Architecture of peptidoglycan: more data and more models. *Trends Microbiol.* **2010,** *18* (2), 59-66.

53. Turner, R. D.; Vollmer, W.; Foster, S. J., Different walls for rods and balls: the diversity of peptidoglycan. *Mol. Microbiol.* **2014,** *91* (5), 862-874.

54.   Mathelié-Guinlet, M.; Asmar, A. T.; Collet, J.-F.; Dufrêne, Y. F., Lipoprotein Lpp

regulates the mechanical properties of the E. coli cell envelope. *Nat. Commun.* **2020,** *11*

(1), 1789.

55.   Auer, G. K.; Lee, T. K.; Rajendram, M.; Cesar, S.; Miguel, A.; Huang, K. C.; Weibel,

D. B., Mechanical Genomics Identifies Diverse Modulators of Bacterial Cell Stiffness.

Cell Syst **2016,** *2* (6), 402-411.

56.   Santa Maria Jr., J. P.; Sadaka, A.; Moussa, S. H.; Brown, S.; Zhang, Y. J.; Rubin, E. J.;

Gilmore, M. S.; Walker, S., Compound-gene interaction mapping reveals distinct roles for

Staphylococcus aureus teichoic acids. *Proc Natl Acad Sci U S A* **2014,** *111* (34), 12510-

12515.

57.   Saar-Dover, R.; Bitler, A.; Nezer, R.; Shmuel-Galia, L.; Firon, A.; Shimoni, E.; Trieu-

Cuot, P.; Shai, Y., D-alanylation of lipoteichoic acids confers resistance to cationic

peptides in group B streptococcus by increasing the cell wall density. *PLoS Pathog* **2012,**

8 (9), e1002891.

58.   Rimal, B.; Senzani, S.; Ealand, C.; Lamichhane, G.; Kana, B.; Kim, S. J.,

Peptidoglycan compositional analysis of Mycobacterium smegmatis using high-

resolution LC-MS. *Sci Rep* **2022,** *12* (1), 11061.

59.   Liu, G.; Cheng, K.; Lo, C. Y.; Li, J.; Qu, J.; Neelamegham, S., A Comprehensive,

Open-source Platform for Mass Spectrometry-based Glycoproteomics Data Analysis. *Mol*

Cell Proteomics **2017,** *16* (11), 2032-2047.

60.   Greene, N. G.; Narciso, A. R.; Filipe, S. R.; Camilli, A., Peptidoglycan branched stem

peptides contribute to Streptococcus pneumoniae virulence by inhibiting pneumolysin

release. *PLoS Pathog.* **2015,** *11* (6), e1004996.

61.   Biswas, R.; Voggu, L.; Simon, U. K.; Hentschel, P.; Thumm, G.; Götz, F., Activity of

the major staphylococcal autolysin Atl. *FEMS Microbiol. Lett.* **2006,** *259* (2), 260-268.

62.   Reiding, K. R.; Bondt, A.; Franc, V.; Heck, A. J. R., The benefits of hybrid

fragmentation methods for glycoproteomics. *TrAC, Trends Anal. Chem.* **2018,** *108*, 260-

268.

ACKNOWLEDGMENTS

This research was supported by Ministry of Science and Technology (MOST), R.O.C.

(Grants MOST 108-2636-M-002-008-, 109-2636-M-002-005-, and 110-2636-M-002-014-). Y.-C.

H. acknowledged the financial support of "The Program of Research Performance Enhancement

via Students Entering Ph.D. Programs Straight from an Undergraduate/Master's Program" from

National Taiwan University. The instrument support from the NTU Mass Spectrometry Platform

was acknowledged.

AUTHOR CONTRIBUTIONS

Y.-C.H., P.-R.S., and C.-C.H. designed the experiments. P.-R.S. and Y.-C.H. built the HAMA

software. Y.-C.H. carried out all the MS experiments and analyzed the data. L.-J.H., K.-Y.C., and

C.-h.C. designed and carried out the AFM experiments. C.-C.H. supervised the study. Y.-C.H.

and C.-C.H. wrote the paper.

AUTHOR INFORMATION

Corresponding Author

* Cheng-Chih Hsu: ccrhsu@ntu.edu.tw

ORCID

Ya-Chen Hsu: 0000-0003-1202-0132

Chun-hsien Chen: 0000-0001-5507-3248

Cheng-Chih Hsu: 0000-0002-2892-5326

COMPETING INTERESTS

705 The authors declare no competing financial interest.

ADDITIONAL INFORMATION

708 Supplementary information includes supplementary figures (MS/MS spectra) and supplementary

709 tables (lists of automated identification of gut bacterial muropeptides). The Source Data

710 underlying Figs. 4–5 is provided as a separate Source Data file. The HAMA software package is

711 available   at: https://drive.google.com/drive/folders/17LVGOm-

712 LEHzNmw7nPULXMEZvg_cnO3k1?usp=share_link

FIGURES

Figure 1. Schematic representation of the high-throughput automated muropeptide analysis (HAMA) framework. (a) The peptidoglycans of bacteria were extracted and purified, followed by mutanolysin digestion. The resulting muropeptide products are analyzed by UHPLC-MS/MS and identified using the HAMA platform. The HAMA strategy involves simplifying muropeptide structures to sequence format, which facilitates the database

construction and *in silico* generation of *b-* and *y-* ion fragmentation spectra for matching.

Muropeptide symbols: B, *N*-acetylglucosamine; M, *N*-acetylmuraminitol (without lactyl group);

l, lactic acid; A, alanine; E, glutamic acid; H, diaminopimelic acid. (b) DBuilder constructs a

muropeptide database containing monomers, dimers, and trimers with two types of linkage:

glycosidic bonds and peptide bonds. For peptide linkages, the direct way is through a direct

covalent bond between the penultimate D-Ala and mDAP residue in the acceptor peptide stem,

and the indirect way is via an interpeptide bridge branching from the lysine. Donors are labeled

in red, and acceptors are labeled in black. (c) The flowchart outlines the LC-MS data processing

in Analyzer.

Figure 2. Automated identification of well-characterized peptidoglycans from *E. coli* and *S. aureus* using the HAMA platform. (a, c) Base peak chromatograms showing the muropeptide analysis of *E. coli* and *S. aureus*. The label content includes retention time (in red), feature index, and muropeptide class (in black). (b, d) Extracted ion chromatograms of the most abundant muropeptide and their MS/MS spectra annotated with *b-* and *y-* fragments were visualized in Viewer. Muropeptide symbols: B, N-acetylglucosamine; M, N-acetylmuraminitol (without lactyl group); l, lactic acid; A, alanine; E, glutamic acid; Q, glutamine; K, lysine; G, glycine.

Figure 3. Resolving isomeric muropeptides by *in silico* MS/MS fragmentation matching. Two isomeric muropeptides with the same parent ion, *m/z* 471.22, were identified as two disaccharide-tripeptides: (a) B-M-l(-A-E-(N-)K eluted at 6.18 min, and (b) B-M-l(-A-Q-(D-)K eluted at 7.02 min. The sequence of each isomer was determined using *in silico* MS/MS fragmentation matching, with the identified sequence having the highest matching score. Muropeptide symbols: B, N-acetylglucosamine; M, N-acetylmuraminitol (without lactyl group); l, lactic acid; A, alanine; E, glutamic acid; Q, glutamine; K, lysine; N, Asparagine; D, Aspartic acid.

Figure 4. Muropeptide composition analysis of *Bifidobacterium breve* strains. (a) Schematic representation of the two possible cross-linking types in the PGN of *B. breve*: Ddt-mediated 4-3 cross-link and Ldt-mediated 3-3 cross-link. Donor peptide stems are labeled in red. The arrow indicates the direction of cross-links catalyzed by transpeptidases. (b) Base peak chromatograms of muropeptide analysis of *B. breve* ATCC 15700, CSCC 1900, and ATCC 15698 strains. The main peaks were annotated with muropeptide symbols. (c) Heatmap showing the muropeptide compositions (% of total) of the PGN of three *B. breve* strains. Symbols: M, monomer; D, dimer; T, trimer (numbers following the letters indicate the number of amino acids in stem peptides). M0, disaccharide; M1, disaccharide-monopeptide; M2, disaccharide-dipeptide; M4, disaccharide-tetrapeptide; M3b, disaccharide-tripeptide with an interpeptide bridge; M3Nb, disaccharide-tripeptide with an anhydro-MurNAc and an interpeptide bridge; M4b, disaccharide-tetrapeptide with an interpeptide bridge; M4Nb, disaccharide-tetrapeptide with an anhydro-MurNAc and an interpeptide bridge; M5b, disaccharide-pentapeptide with an interpeptide

bridge; D34, disaccharide-tripeptide–disaccharide-tetrapeptide with a peptide cross-link; D34N,

disaccharide-tripeptide–disaccharide-tetrapeptide with a peptide cross-link and an anhydro-

MurNAc; D44, disaccharide-tetrapeptide–disaccharide-tetrapeptide with a peptide cross-link;

D44N, disaccharide-tripeptide–disaccharide-tetrapeptide with a peptide cross-link and an

anhydro-MurNAc; D45, disaccharide-tetrapeptide–disaccharide-pentapeptide with a peptide

cross-link; T335, disaccharide-tripeptide–disaccharide-tripeptide-disaccharide-pentapeptide with

two peptide cross-links; T444, disaccharide-tetrapeptide–disaccharide-tetrapeptide-disaccharide-

tetratapeptide with two peptide cross-links; T445, disaccharide-tetrapeptide–disaccharide-

tetrapeptide-disaccharide-pentapeptide with two peptide cross-links.

Figure 5. Bridge length-dependent cell envelope stiffness in *B. longum* and *B. breve*. (a) Schematic illustration of the Gram-positive PGN architecture of *B. longum* (orange tetrapeptide bridges) and *B. breve* (green monopeptide bridges). Glycan strands composed of repeating units of the β-1,4-linked disaccharides are cross-linked by interpeptide bridges, forming 3-D peptidoglycan layers. (b) Heatmap displaying the muropeptide compositions (% of total) of the PGNs in *B. longum* and *B. breve* (three strains each). Symbols: M, monomer; D, dimer; T, trimer

(numbers following the letters indicate the number of amino acids in stem peptides). M0, disaccharide; M1, disaccharide-monopeptide; M2, disaccharide-dipeptide; M4, disaccharide-tetrapeptide; M5, disaccharide-pentapeptide; M3b, disaccharide-tripeptide with an interpeptide bridge; M3Nb, disaccharide-tripeptide with an anhydro-MurNAc and an interpeptide bridge; M4b, disaccharide-tetrapeptide with an interpeptide bridge; M4Nb, disaccharide-tetrapeptide with an anhydro-MurNAc and an interpeptide bridge; M5b, disaccharide-pentapeptide with an interpeptide bridge; D34, disaccharide-tripeptide–disaccharide-tetrapeptide with a peptide cross-link; D34N, disaccharide-tripeptide–disaccharide-tetrapeptide with a peptide cross-link and an anhydro-MurNAc; D44, disaccharide-tetrapeptide–disaccharide-tetrapeptide with a peptide cross-link; D44N, disaccharide-tripeptide–disaccharide-tetrapeptide with a peptide cross-link and an anhydro-MurNAc; D45, disaccharide-tetrapeptide–disaccharide-pentapeptide with a peptide cross-link; T335, disaccharide-tripeptide–disaccharide-tripeptide-disaccharide-pentapeptide with two peptide cross-links; T444, disaccharide-tetrapeptide–disaccharide-tetrapeptide-disaccharide-tetratapeptide with two peptide cross-links; T445, disaccharide-tetrapeptide–disaccharide-tetrapeptide-disaccharide-pentapeptide with two peptide cross-links; Glycan-T, trimers linked by glycosidic bonds. (c) AFM imaging of living *Bifidobacterium*. Topographical images in PBS buffer and the inset shows elasticity images from the top of the cell. Distribution of Young's modulus values corresponding to the elasticity images in the inset. (d) Statistical analysis was performed for each strain, showing the distribution of two groups with the shorter interpeptide bridge corresponded to higher stiffness of cell envelope. Shown here are the mean values (cross), the median, and the 25 and 75% quartiles (boxes) obtained from *N* independent cells over at least three independent experiments. P values were calculated using a one-way ANOVA analysis. *P ≤ 0.05, **P ≤ 0.01, and ****P ≤ 0.0001.

TABLES

800 **Table 1. Muropeptides of *Escherichia coli DH5α* / *Staphylococcus aureus* SA113 analyzed by UPLC-MS/MS.**

	Muropeptide sequence	Retention time (min)	Class[a]	Observed MW (Da)	Theoretical MW (Da)	Δppm	Stem peptides
Escherichia coli	B-M-l(-A-E-H)	2.69	nM3	870.3711	870.3706	-0.60	Tri
	B-M-l(-A-E-H-G)	3.17	nM4	927.3918	927.3921	0.25	Tetra-Gly(4)
	B-M-l(-A-E-H-A)*	**5.13**	**nM4**	**941.4076**	**941.4077**	**0.14**	**Tetra**
	M-l(-A-E-H-A)[b]	5.13	nM4	738.3277	738.3283	0.91	Tetra
	(B-M-l-A-E-H-A-)(B-M-l-A-E-)H	8.53	nDP43	1793.7739	1793.7677	-3.43	Tri-Tetra
	(B-M-l-A-E-H-A-)(B-M-l-A-E-)H-A*	**8.82**	**nDP44**	**1864.8074**	**1864.8048**	**-1.39**	**Tetra-Tetra**
Staphylococcus aureus	B-M-l(-A-Q-K-A-A)	3.39	pM5	967.4721	967.4710	-1.20	Penta
	B-M-l(-A-Q-(G-G-G-G-G-G-)K-A)	5.95	pM10	1238.5641	1238.5626	-1.14	Tetra
	B-M-l(-A-Q-(G-)K-A-A)*	**6.15**	**pM6**	**1024.4925**	**1024.4924**	**-0.02**	**Penta**
	B-M-l(-A-Q-(G-G-G-G-G-G-G-G-)K-A)	6.18	pM12	1352.6041	1352.6056	1.12	Tetra
	B-M-l(-A-Q-(G-G-)K-A-A)	6.41	pM7	1081.5163	1081.5139	-2.26	Penta
	B-M-l(-A-Q-(G-G-G-)K-A-A)	6.48	pM8	1138.5362	1138.5354	-0.75	Penta
	B-M-l(-A-Q-(G-G-G-G-)K-A-A)	6.53	pM9	1195.5577	1195.5568	-0.71	Penta
	B-M-l(-A-Q-(G-G-G-G-G-)K-A-A)*	**6.60**	**pM10**	**1252.5806**	**1252.5783**	**-1.86**	**Penta**
	B-M-l(-A-E-(G-G-G-G-G-)K-A-A)	7.30	pM10	1253.5648	1253.5623	-2.02	Penta
	(B-M-l-A-Q-(G-G-G-G-G-)K-A-G-G-G-G-G-)(B-M-l-A-Q-)K-A-A*	**8.39**	**pDP136**	**2416.1030**	**2416.1089**	**2.44**	**Tetra-Penta**

801 a Muropeptides are cataloged by classes of Gram-negative monomer, nM, Gram-positive monomer, pM, Gram-negative dimer (peptide bond), nDP, Gram-
802 positive dimer (peptide bond), pDP, Gram-negative dimer (glycosidic bond), nDG, Gram-positive dimer (glycosidic bond), pDG, and so on. Numbers following
803 the letters are the total number of amino acids in each peptide strand.

804 [b] Monomer formed in the electrospray source by loss of monosaccharides and recognized as an in-source fragment by exact co-elution.

805 * The main products from PGN hydrolysis were labeled with bold font.

Table 2. The characterized peptidoglycan types of gut bacteria used in this study.

Species	Strains	Glycan backbone		Stem peptide amino acids					Interpeptide bridge (Acceptor ← Donor)
		GlcNAc	MurNAc	1	2	3	4	5	
Bifidobacterium bifidum	DSM 20082	+	+	L-Ala	D-iGln/Glu	L-Orn	D-Ala	D-Ala	D-Ser-D-Asp
Bifidobacterium breve	ATCC 15700 infant intestine	+	+	L-Ala	D-iGln/Glu	L-Lys	D-Ala	D-Ala	Gly
	CSCC 1900 nursling stools	+	+	L-Ala	D-iGln/Glu	L-Lys	D-Ala	D-Ala	Gly
	ATCC 15698 infant intestine	+	+	L-Ala	D-iGln/Glu	L-Lys	D-Ala	D-Ala	Gly
Bifidobacterium longum	ATCC 15707 adult intestine	+	+	L-Ala	D-iGln/Glu	L-Orn	D-Ala	D-Ala	L-Ser-L-Ala-L-Thr-L-Ala
	CSCC 1901 nursling stools	+	+	L-Ala	D-iGln/Glu	L-Orn/	D-Ala	D-Ala	L-Ser-L-Ala-L-Thr-L-Ala
	ATCC 15697 infant intestine	+	+	L-Ala	D-iGln/Glu	L-Orn	D-Ala	D-Ala	L-Ser-L-Ala-L-Thr-L-Ala
Lactobacillus acidophilus	ATCC 4356	+	+	L-Ala	D-iGln/Glu	L-Lys	D-Ala	D-Ala	D-iAsx (Asn+Asp)
Enterococcus faecalis	ATCC 19433	+	+	L-Ala	D-iGln/Glu	L-Lys	D-Ala	D-Ala	L-Ala-L-Ala
Enterococcus faecium	ATCC 19434	+	+	L-Ala	D-iGln/Glu	L-Lys	D-Ala	D-Ala	D-iAsx (Asn+Asp)
Bacteroides fragilis	ATCC 25285	+	*O*-acetylation	L-Ala	D-iGlu	mDAP	D-Ala	D-Ala/ Gly	-
Bacteroides ovatus	ATCC 8483	+	*O*-acetylation	L-Ala	D-iGlu	mDAP	D-Ala	D-Ala/ Gly	-
Bacteroides thetaiotaomicron	ATCC 29741	+	*O*-acetylation	L-Ala	D-iGlu	mDAP	D-Ala	D-Ala/ Gly	-
Akkermansia muciniphila	ATCC BAA-835	*N*-deacetylation	+	L-Ala	D-iGlu	mDAP	D-Ala	D-Ala/ Gly	-