Ustilago maydis Trf2 ensures genome stability by antagonizing Blm-mediated telomere recombination: optimizing DNA repair factor activity through opposing regulations

Authors

Syed S. Zahid[1], Sarah Aloe[1], Jeanette Sutherland[1], William K. Holloman[1], Neal F. Lue [1, 2] *

[1]Department of Microbiology & Immunology, W. R. Hearst Microbiology Research Center, Weill Cornell Medicine, 1300 York Avenue, New York, New York, 10065, United States of America
[2]Sandra and Edward Meyer Cancer Center, Weill Cornell Medicine, 1300 York Avenue, New York, New York, 10065, United States of America

* Corresponding author. Tel: +1 212 746-6506; Fax: +1 212 746-8587; Email: nflue@med.cornell.edu

Running Title: *U. maydis* Trf2 suppresses Blm at telomeres

Abstract

The duplex telomere repeat binding protein TRF2 stabilizes chromosome ends by suppressing DNA damage response and aberrant DNA repair. Herein we report our analysis of the Trf2 ortholog in *Ustilago maydis* (a basidiomycete fungus that bears strong resemblances to mammals with regard to telomere regulation), which underscores the suppression of Blm helicase as a key protective mechanism. Trf2 binds to Blm *in vitro* and inhibits Blm-mediated unwinding of telomeric DNA substrates. Notably, while the loss of Trf2 engenders growth arrest and a multiplicity of telomere aberrations, these defects are fully suppressed by concurrent deletion of *blm* and *mre11* (but not other DNA repair factors). Over-expression of Trf2 induces telomere shortening through a pathway that is epistatic to *blm*Δ . Moreover, over-expression of Blm alone triggers aberrant telomere recombination as evidenced by the accumulation ECTRs and C-circles, and these aberrations can be ameliorated by concurrent over-expression of Trf2. Together, these observations reveal a new mechanism for telomere protection by Trf2, namely through the repression of Blm activity. Interestingly, *U. maydis* harbors a second double-strand telomere repeat-binding protein (Tay1), which promotes Blm function at telomeres to ensure efficient replication. Our results thus point to opposing regulation of Blm helicase by telomere proteins as a strategy for fine-tuning Blm function – such that this helicase is able to promote telomere maintenance without triggering telomere de-protection.

Introduction

Telomeres promote genome integrity by stabilizing chromosome ends against aberrant re-arrangements (Muller, 1938; McClintock, 1941). Stabilization requires the formation of nucleoprotein complexes that prevent the recognition of chromosome ends as double strand breaks (DSBs), which activate DNA damage response (DDR) and DNA repair activities (Arnoult and Karlseder, 2015; de Lange, 2018). Ultimately this concept stipulates that telomeres must directly or indirectly suppress factors in the DDR and DNA repair pathways. However, this relationship is complicated by the fact that DNA repair factors are themselves important for the maintenance of telomere DNA; owing to the propensity of G-rich telomere DNAs to form replication barriers such G-quadruplexes and T-loops, efficient telomere replication requires the participation of DNA repair factors such as Rad51 and Blm to stabilize stalled forks and unwind replication obstacles (Martinez and Blasco, 2015; Brenner and Nandakumar, 2022). Therefore, instead of unconditionally inhibiting DNA repair, telomere proteins must also permit DNA repair factors to act in the appropriate setting to promote telomere maintenance and safeguard genome stability. Understanding the complicated and dichotomous relationships between telomeres and the DNA repair machinery is a central task in telomere research.

The functions of telomeres in mammalian cells are primarily mediated by the shelterin complex, a six-protein assembly that coats double-stranded telomere DNA repeats as well as the terminal 3' overhang (G-overhang) (de Lange, 2018). Among the components of shelterin are two proteins that directly recognize double-stranded telomere repeats (TRF1 and TRF2), one that recognizes the G-strand overhang (POT1), and several associated and bridging factors (RAP1, TIN2, and TPP1). Interestingly, the two double-strand telomere binding proteins, TRF1 and TRF2, most likely arose through gene duplication during metazoan evolution and evidently mediate quite distinct functions in telomere regulation (de Lange, 2018; Myler *et al.*, 2021). TRF1 is crucial for telomere DNA maintenance; it facilitates efficient telomere replication by recruiting or activating the BLM helicase to unwind replication barriers (Zimmermann *et al.*, 2014). In mouse cells, TRF1 also suppresses recombination and BIR through as yet undefined mechanisms (Sfeir *et al.*, 2009; Porreca *et al.*, 2020). By contrast, TRF2 serves essential functions in telomere protection; loss of TRF2 function results in ATM activation and chromosome fusions due to classical non-homologous end joining (cNHEJ), indicating that these DDR and repair systems are the main targets of TRF2 inhibition (van Steensel *et al.*, 1998). The inhibition of cNHEJ by TRF2 is achieved through multiple mechanisms including TRF2-mediated T-loop formation and suppression of 53BP1 accumulation (Doksani *et al.*, 2013; Okamoto *et al.*, 2013). Also potentially playing a role in the inhibition is the binding TRF2 to KU70, an essential cNHEJ factor (Ribes-Zamora *et al.*, 2013). It should be noted, however, that the mechanisms and functions of TRF2 are far from fully understood. For example, overexpression of TRF2 has been shown to cause accelerated telomere shortening as well as telomere anaphase bridges, which lacks clear mechanistic explanation. In addition, TRF2 has been shown to harbor other activities that may be relevant to telomere regulation (Kim *et al.*, 2009; Cicconi *et al.*, 2020). For example, multiple studies have suggested that TRF2 binds directly

to the BLM helicase (Opresko *et al.*, 2002; Lillard-Wetherell *et al.*, 2004), which is implicated in promoting telomere replication and telomere recombination (e.g., in the alternative lengthening of telomeres, or ALT pathway) (Zimmermann *et al.*, 2014; Min *et al.*, 2019; Loe *et al.*, 2020). However, the *in vivo* significance of TRF2-BLM interaction remains unclear. Understanding the full set of interaction partners for TRF2 and the functional consequences of these interactions is necessary for delineating its telomere regulatory mechanisms.

The budding yeast *Saccharomyces cerevisiae* and fission yeast *Schizosaccharomyces pombe* have been used extensively as model systems to investigate telomere regulation. However, both models manifest substantial differences from mammalian telomeres, including the identity and mechanisms of duplex telomere binding proteins (Bianchi and Shore, 2008). As an alternative, we have examined the telomere machinery in the basidiomycete fungus *Ustilago maydis*, originally developed by Robin Holliday to investigate recombinational repair (Holloman *et al.,* 2008). In comparison to budding and fission yeasts, *U. maydis* bears greater resemblances to mammalian cells with respect to the telomere and DNA repair machinery, suggesting that it may be a more informative model (Guzman and Sanchez, 1994; Holloman, 2011). *U. maydis* also harbors the canonical 6-bp telomere repeat sequence ($[TTAGGG]_n/[CCCTAA]_n$), which can better mimic the physical-chemical properties of mammalian telomeres. The greater resemblance of *U. maydis* to mammals is consistent with the more basal position of basidiomycetes in fungal phylogeny, which resulted in a closer evolutionary kinship between this fungal branch and metazoans. In previous studies, we identified two duplex telomere repeat-binding factors in *U. maydis*, named Tay1 and Trf2. These proteins are structurally distinct and mediate non-overlapping functions in telomere maintenance and protection (Yu *et al.*, 2018; Yu *et al.*, 2020). Trf2 is structurally similar to metazoan TRFs and is critical for telomere protection; loss of Trf2 triggers growth arrest and a multiplicity of telomere defects (Yu *et al.*, 2020). Tay1, by contrast, is a fungi-specific telomere protein that promotes telomere replication and recombination by enhancing Blm activity (Yu *et al.*, 2020; Lue, 2021). Thus, even though Tay1 and mammalian TRF1 do not share the same molecular ancestry, they have evidently acquired the same function, thus underscoring the evolutionary advantage of enhancing Blm activity so as to ensure efficient telomere replication.

Herein we report our further investigation of *U. maydis* Trf2, the essential duplex telomere-binding protein that is orthologous to mammalian TRF2. We showed that the telomere deprotection phenotypes of *trf2*-deficienct cells can be greatly suppressed by deleting *blm* and *mre11*, but not several other DNA repair factors. Trf2 inhibits the unwinding of telomeric DNA by Blm *in vitro*. Moreover, while the over-expression of Blm triggers telomere recombination, this phenotype can be suppressed by concurrent overexpression of Trf2. Together with previous findings, our results point to opposing regulation of Blm helicase by telomere proteins as a strategy for fine-tuning Blm function – such that this helicase is able to promote telomere maintenance without triggering telomere de-protection. Previous findings on mammalian TRF1, TRF2 and

BLM suggest that this regulatory strategy may be conserved in evolution, with TRF1 and TRF2 acting in analogous manners as *U. maydis* Tay1 and Trf2.

Results

The growth and telomere defects triggered by Trf2 deficiency are suppressed by deletions of *blm* and *mre11*

Because *U. maydis trf2* is essential, we assessed the consequences of Trf2 loss using *trf2^crg1^*, a mutant in which *trf2* expression is controlled by the arabinose-dependent *crg1* promoter. In this mutant, Trf2 is over-expressed in YPA and transcriptionally repressed in YPD. We had previously shown that loss of Trf2 (*trf2^crg1^* grown in YPD) results in growth arrest and a constellation of telomere defects including telomere length heterogeneity, accumulation of telomere ssDNA, and high levels of C-circles (Yu *et al.*, 2020). To identify the DNA repair factors responsible for triggering the growth defect and telomere aberrations, we constructed and analyzed a series of double mutants that carry both the *trf2^crg1^* allele and a deletion in a DNA repair gene (*blm*, *dna2*, *exo1*, *rad51* and *mre11*). When grown in YPA (*trf2* on), none of the double mutants exhibited significant growth defects (Fig. 1A). However, upon transcriptional repression of *trf2^crg1^* in YPD, all of the double mutants except *trf2^crg1^ blm*Δ and *trf2^crg1^ mre11*Δ manifested growth arrest. Telomere restriction fragment Southern analysis revealed a tight correspondence between suppression of growth defect and maintenance of normal telomere restriction fragment patterns in the double mutants, indicating that *blm*Δ and *mre11*Δ restored viability by suppressing telomere aberrations (Fig. 1B). This is also consistent with the levels of G-strand ssDNA and C-circles in the double mutants; these aberrant structures were present at low or undetectable levels in *trf2^crg1^ blm*Δ and *trf2^crg1^ mre11*Δ, but at high levels in other *trf2^crg1^* double mutants grown in YPD (Fig. 1C and 1D). Together, these results indicate that Blm and Mre11 are both essential for inducing telomere aberrations in the context of Trf2-deficiency and that *U. maydis* Trf2 mediates telomere protection by antagonizing the activities of these DNA repair factors.

Interestingly, while the *trf2^crg1^ blm*Δ and *trf2^crg1^ mre11*Δ mutants are viable on YPD, they manifested reduced growth in comparison to the wild type strain as evidenced by the mutants' small colony size and decreased growth rate (Supp. Fig. 1A and 1B). The basis for this growth defect is unclear given the lack of telomere aberrations. Importantly, the defect does not appear to be the consequence of telomere shortening given that these mutants harbor similarly short telomeres in YPA and YPD and yet exhibit poorer growth in YPD (Fig. 1A). Possibly even without drastic telomere aberration, the lack of Trf2 on telomeric chromatin in Trf2-deficient cells can induce DNA-damage signaling that impairs cell proliferation.

Trf2 overexpression induces telomere shortening

While neither *trf2^crg1^* nor any of the double mutant exhibited growth defect when grown in YPA, they all harbor dramatically shortened but stably maintained telomeres (Fig. 1B). This result suggests that Trf2

over-expression (as a result of growing *trf2^crg1* cells in YPA) may cause telomere shortening. To test the generality of this observation, we generated and analyzed another set of *trf2^crg1* mutants in the FB1 strain background – different from the strain UCM350 used for the majority of this study. Again, these mutants exhibited dramatically shortened telomeres in YPA and manifested growth arrest as well as drastic telomere aberrations in YPD (Supp. Fig. 2A and 2B).

We reasoned that if Trf2 over-expression in YPA triggers telomere shortening, then reducing *trf2* transcription should ameliorate the effects. To titrate the expression of Trf2, we grew *trf2^crg1* in 1.5 % arabinose and increasing concentrations of glucose. (Previous analysis indicates that even in the presence of arabinose, glucose can inhibit *crg1* promoter activity (Bottin *et al.*, 1996).) RT-qPCR analysis confirmed the progressive reduction in *trf2* RNA levels as the glucose concentrations in the media became elevated (Supp. Fig. 3A). As expected, this reduction in *trf2* RNA level is accompanied by a progressive increase in telomere lengths. At 0.125% glucose, the TRF lengths of the *trf2^crg1* mutant resembled those of the parental strain, whereas at lower glucose concentrations, the TRF lengths of the mutant were significantly shorter (Supp. Fig. 3B).

Notably, the telomere shortening phenotype of *trf2^crg1* is reminiscent of several previously characterized DNA repair mutants (e.g., *blm∆* and *rad51∆*), suggesting a defect in telomere replication (Yu *et al.*, 2013; Yu *et al.*, 2018). Given the apparent ability of Trf2 to antagonize Blm and Mre11 activity, a plausible explanation is that Trf2 overexpression may compromise telomere replication by inhibiting the activity of one or more factors required for efficient telomere replication. To assess this hypothesis, we examined the epistasis relationship between the telomere shortening triggered by Trf2 overexpression and those triggered DNA repair gene deletions, e.g., by comparing the telomeres of *trf2^crg1*, *blm∆*, and *trf2^crg1 blm∆* mutants (Fig. 2). Interestingly, this analysis suggests that Trf2 overexpression may act in the same pathway as *blm∆* and *exo1∆* to induce telomere shortening (Fig. 2A). In particular, *trf2^crg1* manifested more severe telomere shortening than *blm∆* or *exo1∆*, but there is no exacerbation of the phenotype in the *trf2^crg1 blm∆* and *trf2^crg1 exo1∆* double mutants (Fig. 2A). We further examined the telomere length distributions of *trf2^crg1* and *trf2^crg1 blm∆* mutants in YPA using STELA and again found no significant difference (Fig. 2B). In contrast, Trf2 over-expression evidently induces telomere shortening through a different pathway than those mediated by *mre11∆* and *dna2∆*; the two double mutants manifested further telomere shortening in comparison to any of the single mutant (Fig. 2C). Together, these results underscore the ability of Trf2 to antagonize DNA repair and highlight an especially interesting interaction between Trf2 and Blm.

Trf2 binds to Blm and inhibits Blm-mediated unwinding of telomeric DNA substrates *in vitro*
The foregoing results suggest that a main function of *Um*Trf2 is to restrain Blm activity at telomeres. We therefore investigated Trf2-Blm interactions as well as the effects of Trf2 on the Blm helicase activity *in

vitro. These assays were performed using an N-terminally truncated Trf2 (Trf2$^{\Delta N}$) because the full length protein could not be expressed in *E. coli* (Yu *et al.*, 2020). Pull down assays using recombinant Blm-FG and His-Trf2$^{\Delta N}$ revealed a modest but detectable interaction between these two proteins (Fig. 3A). To assess the effect of Trf2 on Blm helicase activity, we utilized two 3'-tailed substrates, one with completely non-telomeric sequences and the other with telomere repeats in the duplex region and the 3' tail (Fig. 3B). The substrates were 5' end-labeled on the recessed strands, incubated with Blm and ATP as well as increasing concentrations of Trf2, and then analyzed directly by electrophoresis. Alternatively, the reactions were first terminated with the addition of EDTA and SDS, and then subjected to electrophoresis. In either protocol, the unwinding of substrates by Blm should result in the conversion of the labeled, duplex DNA species into a faster migrating single-stranded DNA band. As shown in Fig. 3B and 3C, Trf2$^{\Delta N}$ inhibited Blm unwinding of telomeric DNA, but had no effect on non-telomeric DNA substrates. At 700 nM Trf2$^{\Delta N}$, the unpairing of telomeric DNA duplex by Blm was almost completely inhibited, whereas that of non-telomeric DNA was unaffected. This result suggests that the DNA-binding activity of Trf2$^{\Delta N}$ is important for Blm inhibition. Indeed, the loss of ssDNA products in telomeric DNA reactions that contained high concentrations of Trf2 is accompanied by the appearance of extremely slowly migrating DNA that likely represents Trf2-DNA complexes (Fig. 3B, marked by an arrow). Together, these results indicate that Trf2 has the capacity to inhibit Blm-mediated unwinding of telomeric DNA.

Overexpression of Blm triggers telomere aberrations

The findings described thus far suggest that stringent control of Blm activity by Trf2 is required for telomere protection. This in turn raises the possibility that over-expression of Blm could override the control and trigger telomere de-protection. We examined this possibility by analyzing conditional *blm* mutants (*blm^{crg1}*) that overexpress *blm* in YPA. RT-qPCR analysis indicates that the *blm* RNA levels in two independently generated mutants grown in YPA are ~7 fold higher than that of the wild type, parental strain (Fig. 4A). Interestingly, this overexpression triggered significant growth deficiency; in comparison to the *UCM350* and *blm*Δ strains, the *blm^{crg1}* mutants exhibited reduced growth on YPA, but not on YPD (Fig. 4B). Analysis of telomere lengths and structures in the *blm*-overexpressing clones revealed multiple aberrations, including telomere length heterogeneity and accumulation extra-short telomeric fragments (Fig. 4C, marked by a bracket), as well as high levels of C-circles (Fig. 4D) and C-strand ssDNA (Fig. 4E and 4F). The extra-short telomeric fragments are extra-chromosomal because they are detectable in the absence of restriction enzyme cleavage (Supp. Fig. 4A) These aberrations were rapidly produced upon Blm overexpression and reached maximal defects following ~ 24 h of Blm induction (Fig. 4C, 4D and 4E). The defects were also reversible upon repression of Blm in YPD (Supp. Fig. 4B). Together, these observations indicate that abnormally high levels of Blm is sufficient to trigger telomere de-protection. Notably, the telomere phenotypes of *blm^{crg1}* grown in YPA share similarities with but are not identical to those of Trf2-deficient cells. For example, the abnormal telomere fragments in the Blm-overexpressing cells were predominantly short, whereas the Trf2-deficient cells accumulate high levels of both long and

short telomeric fragments (compare Fig. 4C and 1B). In addition, the former exhibits high levels of C-strand ssDNA, whereas the latter high levels of G-strand ssDNA. These results suggest that Blm is not the only target of Trf2 inhibition. That is, the phenotypes of Trf2 deficiency may be due to the combined actions of multiple DNA repair factors rather than Blm alone.

Blm overexpression-induced telomere defects are counter-balanced by Trf2 overexpression

The ability of Blm overexpression to trigger telomere defects suggests an *in vivo* assay for assessing Blm inhibition by Trf2. In particular, we reasoned that if Trf2 is indeed a direct Blm inhibitor, then overexpressing Trf2 may counteract the effects of Blm overexpression. To test this possibility, we engineered a series of single and double mutants harboring the blm^{nar1} and $trf2^{crg1}$ allele, which can be independently overexpressed in nitrate-containing and arabinose-containing media, respectively (Fig. 5A). For comparison, we also tested combinations of blm^{nar1} and $pot1^{crg1}$. The strains were grown in YPA to overexpress Trf2/Pot1 only, and in MMA to overexpress Blm and Trf2/Pot1.

The ability of Trf2 overexpression to antagonize Blm overexpression was assessed using C-circle analysis. As shown in Fig. 5B, Blm overexpression alone (blm^{nar1} grown in MMA) induced high levels of C-circles, whereas concurrent overexpression of Trf2 (blm^{nar1} $trf2^{crg1}$ grown in MMA) suppressed C-circle formation. Interestingly, Pot1 overexpression also reduced C-circle formation, although not to the same degree as Trf2 overexpression (Fig. 5B). We also examined the ability of Trf2 overexpression to antagonize telomere length alterations triggered by Blm overexpression. Notably, the telomere length heterogeneity observed in blm^{nar1} grown in nitrate-containing media is considerably milder than that of blm^{crg1} grown in arabinose-containing media, suggesting that *nar1* may overexpress Blm to a lesser degree. Nevertheless, a substantial increase in extra-short telomere fragments can be detected in blm^{nar1} grown in MMA (Fig. 5C, marked by a bracket). Consistent with suppression of Blm by Trf2, these extra-short telomere fragments were largely eliminated in the blm^{nar1} $trf2^{crg1}$ mutant grown in MMA. These results further support inhibition of Blm activity by Trf2 *in vivo*. In addition, given that Blm and Trf2 were the only transcriptionally manipulated factors in the mutant, the inhibitory effect of Trf2 is likely to be direct. As in the case of C-circle analysis, Pot1 overexpression also reduced the level of extra-short telomeres, suggesting that Pot1 can also antagonize Blm activity at telomeres.

Opposing regulation of Blm by the two duplex telomere binding proteins in *U. maydis*

As noted before, in addition to Trf2, *U. maydis* harbors a fungi-specific duplex telomere repeat binding factor named Tay1 (Yu *et al.*, 2020; Lue, 2021). Previous studies suggest that instead of telomere protection, the main function of Tay1 is to stimulate Blm activity at telomeres in order to promote telomere replication and recombination. Thus, *U. maydis* appears to achieve optimal regulation of Blm by utilizing two telomere proteins with opposing effects on this helicase. One potential implication of this is that loss of Tay1 function might dampen Blm activity and partially suppress defects engendered by Trf2 deficiency.

Alternatively, Tay1 might have a secondary function in telomere protection that is only discernible in the absence of Trf2. To further investigate the interplay between Tay1 and Trf2 in telomere regulation, we compared the phenotypes of the *tay1Δ*, *trf2crg1*, and *tay1Δ trf2crg1* mutants (Fig. 6). Because we had previously generated *tay1Δ* using the FB1 parental strain, we proceeded to construct and analyze both *trf2crg1* and *tay1Δ trf2crg1* in the same strain background.

Growth analysis revealed similar growth defects for *trf2crg1* and *tay1Δ trf2crg1* on YPD, suggesting that Tay1 does not alter the toxicity induced by Trf2 deficiency (Fig. 6A). However, TRF Southern analysis uncovered several interesting differences between the single and double mutants (Fig. 6B). First, when grown in YPA, the *tay1Δ trf2crg1* double mutant manifested shorter telomeres than either *tay1Δ* and *trf2crg1* single mutants. Second, when grown in YPD, the *tay1Δ trf2crg1* double mutant manifested less severe telomere aberrations than the *trf2crg1* single mutant, consistent with mild suppression of telomere deprotection by *tay1Δ* – presumably through the reduction of Blm helicase activity (Fig. 6B). We reasoned that this suppression could be more prominent if Trf2 is only partially lost such that Blm is not overactive. Indeed, when spotted on plates containing 1.5% arabinose and 1% glucose (to enable partial expression of the *trf2crg1* allele), *tay1Δ trf2crg1* manifested improved growth and substantially milder telomere aberrations in comparison to *trf2crg1*(Supp. Fig. 5A and 5B). Together, these observations highlight the opposing effects of Tay1 and Trf2 on Blm function at telomeres and suggest that Tay1 does not serve any function in telomere protection even in the absence of Trf2.

Aberrant transcription of G- and C-strand telomere repeats in Blm-overexpressing and Trf2-deficient cells

Our recent analysis of Pot1-deficient cells indicates that upon telomere deprotection, both the G- and C-strand of telomere repeats are transcribed at abnormally high levels, suggesting that deprotected telomeres are substrates for the transcription apparatus (Zahid *et al.*, 2022). There is also growing evidence that DNAs flanking DSBs are actively transcribed, pointing to another resemblance between DSBs and deprotected telomeres (Ohle *et al.*, 2016; Liu *et al.*, 2021; Sharma *et al.*, 2021). To determine if aberrant transcription is associated with deprotected telomeres in general, we examined the levels of G- and C-strand telomere repeat RNAs in Blm-overexpressing mutants (*blmcrg1* grown in YPA) using both dot blot and RT-qPCR analyses (Fig. 7A and 7B). Consistent with elevated transcription at de-protected telomeres, we found high levels of both G- and C-strand RNAs in *blmcrg1* strains grown in YPA but not YPD (Fig. 7A and 7B). Interestingly, quantifications of the results point to higher levels of C-strand RNAs in comparison to G-strand RNAs (by ~ 10 and 50-fold as assessed by dot blot and RT-qPCR, respectively). We also used endpoint RT-PCR to assess the structures of these RNAs, which confirmed the existence of G- and C-strand RNAs that span both subtelomeric and telomeric regions (Fig. 7C). For example, using forward and reverse primers that anneal to the subtelomeric and telomeric regions, respectively, we successfully amplified heterogeneously sized products that could only come from RNAs

containing both regions (Fig. 7C, middle panel). As another test of the hypothesis that telomere deprotection triggers aberrant transcription, we examined the levels of G- and C-strand RNAs in Trf2-deficient cells (Supp. Fig. 6). Indeed, both *trf2^crg1* and *tay1Δ trf2^crg1* manifested high levels of telomere repeat RNAs when grown in YPD. As in the case of Blm-overexpressing cells, the C-strand RNAs appear to be present at significantly higher levels as judged by qRT-PCR analysis (Supp. Fig. 6B) Together, these findings indicate that aberrant transcription of both strands of the telomere repeat region is a shared feature of de-protected telomeres triggered by multiple mutations.

Discussion

Studies of mammalian TRF2, one of two major duplex telomere repeat binding factors, have highlighted its critical roles in suppressing ATM signaling and cNHEJ at chromosome ends. In the current study, we interrogated the functions of *trf2* in the basidiomycete fungus *U. maydis* and uncovered instead, a major function for this protein in suppressing Blm-driven telomere recombination at chromosome ends. Notably, several previous investigations of mammalian TRF2 also hint at a physical and functional connection between TRF2 and BLM. Below we discuss the mechanistic, functional and evolutionary implications of our findings.

The regulation of Blm by Trf2; functional connections between Trf2 and Ku70/Ku80 complex

Our data argue that a primary function of *U. maydis* Trf2 is to suppress Blm activity at telomeres; all of the major telomere aberrations observed in the *trf2*-deficient setting are eliminated by deletion of *blm*. This is quite unexpected and provides another striking illustration of the concept that telomeres have evolved to antagonize aberrant repair. Notably, the phenotypes of *trf2*-deficient cells are quite similar to those triggered by loss of *U. maydis* Ku70 and Ku80, including telomere length heterogeneity, accumulation of ssDNA, and high levels ECTRs. Moreover, these phenotypes of the *trf2* and *ku70/ku80* mutants are both strongly suppressed by *blm* or *mre11* deletion. These findings strongly suggest a physical connection or functional overlap between Trf2 and Ku70/Ku80. Indeed, previous analyses of human telomeres have uncovered an interaction between these proteins and implicated this interaction in the regulation NHEJ at telomeres (Song *et al.*, 2000; Ribes-Zamora *et al.*, 2013). Therefore, the existence of such a direct interaction in *U. maydis* warrants further investigation. However, it is also worth noting that the *trf2* and *ku70/ku80* mutants are not phenotypically identical. In particular, whereas *ku70/ku80* loss results in high levels C-strand ssDNA, Trf2 deficiency leads to the accumulation of G-strand. Therefore, either Trf2 or Ku70/Ku80 must be capable of independently acting on some DNA repair factors to alter resection or recombination. The distinct ssDNA accumulation thus provides a phenotypic assay to dissect the mechanistic distinctions between Trf2 and the Ku complex at telomeres.

The viability of the *trf2^crg1^ blm∆* and *trf2^crg1^ mre11∆* mutants has striking implications and raises some interesting issues. One implication relates to the DNA damage signaling pathway(s) triggered by Trf2 loss: given the viability of *trf2^crg1^ blm∆* and *trf2^crg1^ mre11∆*, this signaling pathway is either inactive or unable to arrest growth in absence of Blm or Mre11. Therefore, DNA damage signaling and DNA repair appear to work in a mutually reinforcing manner in the context of telomere protein deficiency to trigger telomere de-protection and growth arrest. Unfortunately, while a number of DDR kinases have been identified in *U. maydis*, relatively little is known about their activities at *U. maydis* telomeres. In this regard, it is worth noting that both *atr1∆* and *chk1∆* can rescue the lethality of Ku70-deficiency, but only partially suppress the telomere defects of this mutant (de Sena-Tomas *et al.*, 2015; Yu *et al.*, 2015). This result would suggest that Blm acts upstream of *atr1* and *chk1* to instigate telomere deprotection and checkpoint activation.

An unresolved question regarding the *trf2^{crg1} blm*Δ and *trf2^{crg1} mre11*Δ mutants is the nature of the telomere nucleoprotein structure in these mutants when Trf2 is depleted, i.e., when the strains are grown in YPD. Telomeres are substantially shorter (due to inefficient telomere replication in the absence of Blm), but it is unclear if Tay1, the other duplex telomere binding protein, can fully occupy the telomere repeat tracts. Could nucleosome bind to a significant portion of the telomere tracts, and what are the impacts of such an altered telomere structure on telomere protection and maintenance? Also unclear is whether changes at duplex telomere repeats can affect the nucleoprotein structure (e.g., the binding of Pot1 and Tpp1) at the single-stranded telomere termini. Interestingly, even though *trf2^{crg1} blm*Δ and *blm*Δ exhibit similarly short telomeres in YPD, the former manifested slower growth, suggesting that the altered telomeres devoid of Trf2 may impinge on cell cycle regulation. Further analysis of the structure and nucleoprotein composition of telomeres in the *trf2^{crg1} blm*Δ and *trf2^{crg1} mre11*Δ mutants should be informative.

Opposing regulation of Blm by duplex telomere repeat binding proteins

A key observation of the current study is opposing regulation of Blm by two duplex telomere repeat binding proteins (i.e., Tay1 and Trf2), which underscores the complex roles of Blm at telomeres. The rationale for the antagonism between Tay1 and Trf2 is evidently to fine-tune Blm activity such that this helicase is able to promote telomere replication without triggering abnormal repair. The need to balance the activating and repressing function of Tay1 and Trf2 vis-à-vis Blm is further illustrated by the dosage effects of both Trf2 and Blm at telomeres. Excessive suppression of Blm via Trf2-overexpression leads to telomere shortening. In contrast, over expression of Blm can override Trf2-mediated telomere protection, unless Trf2 is concurrently over-expressed. Therefore, the optimal calibration of Blm function is also predicated on expressing appropriate levels of Blm and Trf2 proteins. In short, the elaborate system that has evolved in *U. maydis* to optimize Blm function include proper levels of Tay1, Trf2 and Blm, as well as opposing effects of Tay1 and Trf2 on Blm activity.

Physical and function interactions between mammalian TRF2 and BLM; potential similarities between U. maydis and mammals

Even though mammalian TRF2 has not been reported to suppress BLM activity, this possibility is hinted by several previous findings. First, overexpression of TRF2 in primary cells increased the rate of telomere shortening, which could be due to reduced efficiency of telomere replication as a result of BLM inhibition (Karlseder *et al.*, 2002). Second, overexpression of TRF2 in telomerase-positive cancer cells caused replication fork stalling at telomeres and induced the formation of telomeric ultrafine bridges, which may also be explained by compromised BLM function (Nera *et al.*, 2015). Indeed, anaphase ultrafine bridges are characteristic of BLM-deficient cells (Chan *et al.*, 2007). Interestingly, in contrast to *Um*Trf2, mammalian TRF2 was previously reported to stimulate BLM helicase activity *in vitro* (Opresko *et al.*, 2002; Lillard-Wetherell *et al.*, 2004). This discrepancy could be due to differences in reaction parameters between the

earlier and current studies, e.g., in the lengths of telomere repeats in substrate DNAs or in the reaction conditions. Additional studies in mammalian cells, especially regarding the *in vivo* effect of TRF2 expression on BLM activity, may be informative.

As noted before, the predominant consequence of TRF2 dysfunction at mammalian telomeres is NHEJ, whereas that of Trf2 deficiency at *U. maydis* telomeres is Blm-mediated resection and recombination. This difference, however, does not necessarily argue against a BLM inhibitory function for mammalian TRF2. One possibility is that TRF2 may work redundantly with another factor to restrain BLM activity. Alternatively, NHEJ may be less active at *U. maydis* telomeres, leading to preferential channeling of Trf2-deficient telomeres through the Blm-dependent recombination pathway.

Elevated transcription at de-protected telomeres
We found that similar to the *pot1* mutant, in both Blm-overexpressing and Trf2-deficient cells, the telomere de-protection is accompanied by elevated levels of both G- and C-strand telomere repeat RNAs (Zahid *et al.*, 2022). These findings suggest that increased transcription of both telomere strands is a general consequence of telomere de-protection. This in turn raises a key question regarding the nature and consequences of these transcripts, namely, do they represent elevated levels of RNAs that are normally expressed at low levels, or do they represent fundamentally abnormal RNAs that are produced only in pathologic settings? In this regard, it is worth noting that most studies of cells with normal telomere structures have revealed G-strand RNAs that initiate from subtelomeric promoters and elongate into telomere repeats (i.e., TERRA) (Azzalin and Lingner, 2015; Kwapisz and Morillon, 2020). C-strand RNAs, in contrast, have not been reported or characterized in normal cells except in the case of *S. pombe*; whether these C-strand RNAs perform important physiologic functions remains unclear (Bah *et al.*, 2012; Greenwood and Cooper, 2012). Thus, it seems likely that at least a portion of the RNAs in the mutants represent fundamentally different transcripts that are only synthesized from deprotected telomeres.

While more studies will be required to understand the nature of the telomere transcripts in the mutants, we note that the synthesis of such aberrant RNAs exhibits similarities to two previous described pathways. First, there is growing evidence that regions flanking double strand breaks (DSBs) are transcribed, and that these RNAs form RNA-DNA hybrids that can modulate the repair of DSBs (Ohle *et al.*, 2016; Liu *et al.*, 2021; Sharma *et al.*, 2021). Since dysfunctional telomeres induce similar cellular response as that by double strand breaks, the telomere transcripts may be synthesized through the same pathway(s) as the DSB-associated RNAs. In support of this idea, the DSB-associated RNAs are thought to correspond primarily to the strand that bears the 5' ends, i.e., equivalent to the C-strand (Liu *et al.*, 2021). Therefore, the DSB paradigm provides a plausible explanation for why the *U. maydis* telomere protein mutants harbor higher levels of C-strand RNAs. Alternatively, the abnormal telomere transcripts in the mutants could be a consequence of perturbing telomere chromatin structure, much as global depletion/perturbation of nucleosomes has been shown to trigger promiscuous transcription (Doris *et al.*, 2018; Kornberg and Lorch,

2020). In this paradigm, the synthesis of aberrant telomere RNAs is unrelated to the presence of the DNA ends and is simply governed by the accessibility of DNA to the transcription apparatus. Regardless of the mechanisms of telomere RNA synthesis, once generated, these RNAs have the capacity to form hybrids and R-loops with telomere repeats, thereby altering DNA damage response and DNA repair. Hence a full understanding of telomere dysfunction requires insights not only on the synthesis of aberrant RNAs, but also their impacts on telomere structure and function. The conditional mutants that we have developed in *U. maydis* provides a useful system to interrogate these issues.

Materials and Methods

<u>*Ustilago maydis* strain construction and growth conditions</u>

Standard protocols were employed for the genetic manipulation of *U. maydis* (Kojic *et al.*, 2002; Brachmann *et al.*, 2004; Kamper, 2004). All *U. maydis* strains used in this study were haploid and were derived from the UCM350 or FB1 background (Banuett and Herskowitz, 1989; Kojic *et al.*, 2002) and are listed in S1 Table.

The *blm^{crg1}* strain was constructed by integrating a linearized pRU12-*blm* plasmid containing fragments that correspond to (1) 700 bp of the *blm* promoter and (2) the first 700 bp of the *blm* ORF into the *blm* genomic locus in UCM350. Briefly, the two *blm* fragments were generated by PCR using appropriate oligos with flanking restriction enzyme sites (S2 Table), and then cloned together in between the *Nde*I and *Eco*RI sites of pRU12 (Bottin *et al.*, 1996). Correct recombination of *Xba*I treated pRU12-*blm* into the genomic locus results in the insertion of the *cbx^R* marker and the *crg1* promoter precisely adjacent to and upstream of the *blm* ATG initiation codon. Following transformation, the correct mutant strains (*blm^{crg1}*) were selected on YPA supplemented with carboxin (4 µg/ml) and confirmed by PCR. The *blm^{nar1}* strain was similarly constructed using a linearized (digested by *Nsi*I) pRU2-Hyg-*blm* plasmid carrying identical *blm* promoter and coding regions. The transformants were selected on YPA supplemented with hygromycin (150 µg/ml) and confirmed by PCR. The *trf2^{crg1} blm^{nar1}* and *pot1^{crg1} blm^{nar1}* mutants were made by transforming *trf2^{crg1}* and *pot1^{crg1}* with the linearized pRU2-Hyg-*blm* plasmid and screening for correct transformants. The *blm∆ trf2^{crg1}*, *dna2∆ trf2^{crg1}*, *exo1∆ trf2^{crg1}*, and *mre11∆ trf2^{crg1}* mutants were constructed by transforming XbaI digested pRU12-Trf2 into the respective deletion mutants, and screening for correct transformant by PCR genotyping. The *rad51∆ trf2^{crg1}* mutant was made by transforming *trf2^{crg1}* with a *rad51* disruption cassette (pUC-M2T1-rad51 digested with *Ssp*I).

For growth analysis of the mutants, liquid cultures of *U. maydis* strains were first diluted to OD$_{600}$ of 1.0 (corresponding to ~ 0.5 x 10^6 cells/ml), and subjected to serial 6-fold dilutions. Four microliters of the diluted cultures were then spotted onto YPA or YPD plates and grown at 30°C for 2 to 3 days.

<u>RT-qPCR analysis of *blm* and *trf2* RNA levels</u>

For RT-qPCR analysis *blm* and *trf2* expressions levels, RNAs were extracted and purified from log-phase *U. maydis* cultures using acid phenol extraction (Collart and Oliviero, 1993). The samples were treated with TURBO DNase (Thermo Fisher) and then reverse transcribed in 20 µl reactions that contained 1x SSIV Buffer, 12.5 µM primer (Supp. Table 2), 1 mM dNTP, 5 mM DTT, 20 U NxGen RNase Inhibitor (Lucigen), and 40 U Superscript IV RT (Thermo Fisher). The reaction mixtures were incubated at 55°C for 40 min and then heated at 70°C for 15 min to inactivate the reverse transcriptase. qPCR was carried out in Chai Open qPCR Dual Channel thermocycler using Chai Green Master mix with 8 µM primers

(Supp. Table 2) and 1 µl of cDNA. Cycling parameters were as follows: 45 cycles of 95° C for 30 sec, 53° C for 20 sec, 72° C for 40 sec.

<u>Telomere length and structural analyses</u>

Southern analysis of telomere restriction fragments (TRF Southern) was performed using DNA treated with *Pst*I as described previously (Yu *et al.*, 2013; Yu *et al.*, 2018). The blots were hybridized to an oligonucleotide containing eight copies of the TTAGGG repeats. For quantitation, we used the formula

mean TRF length = Σ (OD$_i$ × L$_i$) / Σ (OD$_i$)

because the majority of TRF length variation stems from differences in subtelomeric lengths (Kimura *et al.*, 2010; Mender and Shay, 2015). The in-gel hybridization assays for detecting ssDNA on the G- and C-strand have been described (de Sena-Tomas *et al.*, 2015). STELA for telomeres bearing UT4 subtelomeric elements was performed essentially as reported before (Swapna *et al.*, 2018). C-circle and G-circles assays were performed as before except that NxGen phi29 DNA Polymerase (Lucigen Corp.) were used for rolling circle amplifications (Yu *et al.*, 2018).

<u>Analysis of telomere repeat-containing RNAs</u>

RNAs were purified from log-phase *U. maydis* cultures using acid phenol extraction (Collart and Oliviero, 1993). For dot blot analysis, 0.2 µg of RNAs (untreated or pre-treated with various nucleases) in 10 µl were diluted with 30 µl of RNA incubation/denaturation solution (65.7% formamide, 7.7 % formaldehyde, 1.3x MOPS buffer), heated at 65°C for 5 min, and then cooled on ices. After adding 40 µl of 20x SSC, the entire samples were applied to Hybond-N using a Bio-Dot Microfiltration Apparatus (Bio-Rad Laboratories, Inc.). After UV crosslinking, the blot was hybridized sequentially to the same probes as those for Northern analysis. For end-point RT-PCR, the RNA samples from acid phenol extraction were first treated with Turbo DNase (Thermo Fisher Inc.). Fifty ng of RNAs were subjected to reverse transcription using Superscript III RT at 55°C for 40 min, followed by heating at 70°C for 15 min to inactivate RT, and then subjected to PCR (33-37 cycles of 95°C for 15 sec, 66°C for 15 sec, and 72°C for 30 sec) with Q5 polymerase (NEB Inc.) and appropriate primers (Supp. Table 2).

<u>Pull down and helicase assays for analyzing Trf2-Blm interactions</u>

Blm-FG was first expressed as His$_6$-SUMO-Blm-FG fusion in the BL21 codon plus strain and purified by a combination of (i) Ni-NTA chromatography, (ii) ULP1 cleavage, and (iii) anti-FLAG affinity chromatography as previously described (Ref.). Following binding to the anti-FLAG resin and washing with FLAG(150) buffer (50 mM Tris-HCl, pH 7.5, 150 mM NaCl, 10 % glycerol, 0.1 % NP-40, 2.5 mM MgCl$_2$, 1 mM DTT), the beads were used directly in pull down assays. The prey for these assays, Trf2$^{\Delta N}$, was likewise expressed as His$_6$-SUMO-Trf2$^{\Delta N}$ fusion in BL21 and purified via Ni-NTA chromatography. The partially purified His$_6$-SUMO-Trf2$^{\Delta N}$ was then mixed with anti-FLAG resin carrying immobilized Blm-FG in FLAG(65) buffer (same as FLAG(150) except that NaCl is at 150 mM). Following incubation with mixing

on a rotator at 4°C for 1 h, the beads were washed 4 times with 0.5 ml of the FLAG(100) buffer (same as FLAG(150) except that NaCl is at 100 mM), and then the bound proteins eluted with 60 µl FLAG(150) containing 0.2 mg/ml FLAG$_3$ peptide. The eluates were analyzed by Western Blot using anti-His and anti-FLAG antibodies.

The helicase assays were performed as described using 25 nM Blm-FG and varying concentrations of Trf2$^{\Delta N}$ (Yu *et al.*, 2020). After 20 min of incubation at 35^{0}C, the reactions were terminated by the addition of 4 µl stop solution (30 % glycerol, 50 mM EDTA, 0.9 % SDS, 0.1 % BPB and 0.1 % xylene cyanol) and 20 ng of unlabeled bottom strand oligo (as trap to prevent reannealing of unwound ssDNA) or directly loaded onto 10% polyacrylamide gels (acrylamide : bis-acrylamide = 29 : 1) containing 1x TBE. The substrates and products were separated via electrophoresis, and the results analyzed using a Phosphor scanner. The ratio of labeled DNA in single-stranded form (bottom species in the gel) to total DNA (all the labeled species) at the end of the reaction is taken as the fraction unwound by Blm helicase.

Acknowledgments

We thank members of our laboratories for discussion. The authors declare no competing interests.

References

Arnoult, N., and Karlseder, J. (2015). Complex interactions between the DNA-damage response and mammalian telomeres. Nat Struct Mol Biol *22*, 859-866.

Azzalin, C.M., and Lingner, J. (2015). Telomere functions grounding on TERRA firma. Trends Cell Biol *25*, 29-36.

Bah, A., Wischnewski, H., Shchepachev, V., and Azzalin, C.M. (2012). The telomeric transcriptome of Schizosaccharomyces pombe. Nucleic acids research *40*, 2995-3005.

Banuett, F., and Herskowitz, I. (1989). Different a alleles of Ustilago maydis are necessary for maintenance of filamentous growth but not for meiosis. Proceedings of the National Academy of Sciences of the United States of America *86*, 5878-5882.

Bianchi, A., and Shore, D. (2008). How telomerase reaches its end: mechanism of telomerase regulation by the telomeric complex. Mol Cell *31*, 153-165.

Bottin, A., Kamper, J., and Kahmann, R. (1996). Isolation of a carbon source-regulated gene from Ustilago maydis. Mol Gen Genet *253*, 342-352.

Brachmann, A., Konig, J., Julius, C., and Feldbrugge, M. (2004). A reverse genetic approach for generating gene replacement mutants in Ustilago maydis. Molecular genetics and genomics : MGG *272*, 216-226.

Brenner, K.A., and Nandakumar, J. (2022). Consequences of telomere replication failure: the other end-replication problem. Trends Biochem Sci *47*, 506-517.

Chan, K.L., North, P.S., and Hickson, I.D. (2007). BLM is required for faithful chromosome segregation and its localization defines a class of ultrafine anaphase bridges. EMBO J *26*, 3397-3409.

Cicconi, A., Rai, R., Xiong, X., Broton, C., Al-Hiyasat, A., Hu, C., Dong, S., Sun, W., Garbarino, J., Bindra, R.S., Schildkraut, C., Chen, Y., and Chang, S. (2020). Microcephalin 1/BRIT1-TRF2 interaction promotes telomere replication and repair, linking telomere dysfunction to primary microcephaly. Nat Commun *11*, 5861.

Collart, M.A., and Oliviero, S. (1993). Preparation of yeast RNA. Curr Protoc Mol Biol *23*, 13.12.11-13.12.15.

de Lange, T. (2018). Shelterin-Mediated Telomere Protection. Annu Rev Genet *52*, 223-247.

de Sena-Tomas, C., Yu, E.Y., Calzada, A., Holloman, W.K., Lue, N.F., and Perez-Martin, J. (2015). Fungal Ku prevents permanent cell cycle arrest by suppressing DNA damage signaling at telomeres. Nucleic acids research.

Doksani, Y., Wu, J.Y., de Lange, T., and Zhuang, X. (2013). Super-resolution fluorescence imaging of telomeres reveals TRF2-dependent T-loop formation. Cell *155*, 345-356.

Doris, S.M., Chuang, J., Viktorovskaya, O., Murawska, M., Spatt, D., Churchman, L.S., and Winston, F. (2018). Spt6 Is Required for the Fidelity of Promoter Selection. Mol Cell *72*, 687-699 e686.

Greenwood, J., and Cooper, J.P. (2012). Non-coding telomeric and subtelomeric transcripts are differentially regulated by telomeric and heterochromatin assembly factors in fission yeast. Nucleic acids research *40*, 2956-2963.

Guzman, P.A., and Sanchez, J.G. (1994). Characterization of telomeric regions from Ustilago maydis. Microbiology *140 (Pt 3)*, 551-557.

Holloman, W.K. (2011). Unraveling the mechanism of BRCA2 in homologous recombination. Nat Struct Mol Biol *18*, 748-754.

Holloman, W.K., Schirawski, J., and Holliday, R. (2008). The homologous recombination system of Ustilago maydis. Fungal Genet Biol *45 Suppl 1*, S31-39.

Kamper, J. (2004). A PCR-based system for highly efficient generation of gene replacement mutants in Ustilago maydis. Molecular genetics and genomics : MGG *271*, 103-110.

Karlseder, J., Smogorzewska, A., and de Lange, T. (2002). Senescence induced by altered telomere state, not telomere loss. Science *295*, 2446-2449.

Kim, H., Lee, O.H., Xin, H., Chen, L.Y., Qin, J., Chae, H.K., Lin, S.Y., Safari, A., Liu, D., and Songyang, Z. (2009). TRF2 functions as a protein hub and regulates telomere maintenance by recognizing specific peptide motifs. Nat Struct Mol Biol *16*, 372-379.

Kimura, M., Stone, R.C., Hunt, S.C., Skurnick, J., Lu, X., Cao, X., Harley, C.B., and Aviv, A. (2010). Measurement of telomere length by the Southern blot analysis of terminal restriction fragment lengths. Nature Protocols *5*, 1596-1607.

Kojic, M., Kostrub, C.F., Buchman, A.R., and Holloman, W.K. (2002). BRCA2 homolog required for proficiency in DNA repair, recombination, and genome stability in Ustilago maydis. Mol Cell *10*, 683-691.

Kornberg, R.D., and Lorch, Y. (2020). Primary Role of the Nucleosome. Mol Cell *79*, 371-375.

Kwapisz, M., and Morillon, A. (2020). Subtelomeric Transcription and its Regulation. J Mol Biol *432*, 4199-4219.

Lillard-Wetherell, K., Machwe, A., Langland, G.T., Combs, K.A., Behbehani, G.K., Schonberg, S.A., German, J., Turchi, J.J., Orren, D.K., and Groden, J. (2004). Association and regulation of the BLM helicase by the telomere proteins TRF1 and TRF2. Hum Mol Genet *13*, 1919-1932.

Liu, S., Hua, Y., Wang, J., Li, L., Yuan, J., Zhang, B., Wang, Z., Ji, J., and Kong, D. (2021). RNA polymerase III is required for the repair of DNA double-strand breaks by homologous recombination. Cell *184*, 1314-1329 e1310.

Loe, T.K., Li, J.S.Z., Zhang, Y., Azeroglu, B., Boddy, M.N., and Denchi, E.L. (2020). Telomere length heterogeneity in ALT cells is maintained by PML-dependent localization of the BTR complex to telomeres. Genes & development *34*, 650-662.

Lue, N.F. (2021). Duplex Telomere-Binding Proteins in Fungi With Canonical Telomere Repeats: New Lessons in the Rapid Evolution of Telomere Proteins. Front Genet *12*, 638790.

Martinez, P., and Blasco, M.A. (2015). Replicating through telomeres: a means to an end. Trends Biochem Sci *40*, 504-515.

McClintock, B. (1941). The Stability of Broken Ends of Chromosomes in Zea Mays. Genetics *26*, 234-282.

Mender, I., and Shay, J.W. (2015). Telomere Restriction Fragment (TRF) Analysis. Bio Protoc *5*.

Min, J., Wright, W.E., and Shay, J.W. (2019). Clustered telomeres in phase-separated nuclear condensates engage mitotic DNA synthesis through BLM and RAD52. Genes & development *33*, 814-827.

Muller, H.J. (1938). The remaking of chromosomes. Collecting Net *8*, 182-198.

Myler, L.R., Kinzig, C.G., Sasi, N.K., Zakusilo, G., Cai, S.W., and de Lange, T. (2021). The evolution of metazoan shelterin. Genes & development.

Nera, B., Huang, H.S., Lai, T., and Xu, L. (2015). Elevated levels of TRF2 induce telomeric ultrafine anaphase bridges and rapid telomere deletions. Nat Commun *6*, 10132.

Ohle, C., Tesorero, R., Schermann, G., Dobrev, N., Sinning, I., and Fischer, T. (2016). Transient RNA-DNA Hybrids Are Required for Efficient Double-Strand Break Repair. Cell *167*, 1001-1013 e1007.

Okamoto, K., Bartocci, C., Ouzounov, I., Diedrich, J.K., Yates, J.R., 3rd, and Denchi, E.L. (2013). A two-step mechanism for TRF2-mediated chromosome-end protection. Nature *494*, 502-505.

Opresko, P.L., von Kobbe, C., Laine, J.P., Harrigan, J., Hickson, I.D., and Bohr, V.A. (2002). Telomere-binding protein TRF2 binds to and stimulates the Werner and Bloom syndrome helicases. J Biol Chem *277*, 41110-41119.

Porreca, R.M., Herrera-Moyano, E., Skourti, E., Law, P.P., Gonzalez Franco, R., Montoya, A., Faull, P., Kramer, H., and Vannier, J.B. (2020). TRF1 averts chromatin remodelling, recombination and replication dependent-break induced replication at mouse telomeres. Elife *9*.

Ribes-Zamora, A., Indiviglio, S.M., Mihalek, I., Williams, C.L., and Bertuch, A.A. (2013). TRF2 interaction with Ku heterotetramerization interface gives insight into c-NHEJ prevention at human telomeres. Cell Rep *5*, 194-206.

Sfeir, A., Kosiyatrakul, S.T., Hockemeyer, D., MacRae, S.L., Karlseder, J., Schildkraut, C.L., and de Lange, T. (2009). Mammalian telomeres resemble fragile sites and require TRF1 for efficient replication. Cell *138*, 90-103.

Sharma, S., Anand, R., Zhang, X., Francia, S., Michelini, F., Galbiati, A., Williams, H., Ronato, D.A., Masson, J.Y., Rothenberg, E., Cejka, P., and d'Adda di Fagagna, F. (2021). MRE11-RAD50-NBS1 Complex Is Sufficient to Promote Transcription by RNA Polymerase II at Double-Strand Breaks by Melting DNA Ends. Cell Rep *34*, 108565.

Song, K., Jung, D., Jung, Y., Lee, S.G., and Lee, I. (2000). Interaction of human Ku70 with TRF2. FEBS Lett *481*, 81-85.

Swapna, G., Yu, E.Y., and Lue, N.F. (2018). Single telomere length analysis in Ustilago maydis, a high-resolution tool for examining fungal telomere length distribution and C-strand 5'-end processing. Microb Cell *5*, 393-403.

van Steensel, B., Smogorzewska, A., and de Lange, T. (1998). TRF2 protects human telomeres from end-to-end fusions. Cell *92*, 401-413.

Yu, E.Y., Hsu, M., Holloman, W.K., and Lue, N.F. (2018). Contributions of recombination and repair proteins to telomere maintenance in telomerase-positive and negative Ustilago maydis. Mol Microbiol *107*, 81-93.

Yu, E.Y., Kojic, M., Holloman, W.K., and Lue, N.F. (2013). Brh2 and Rad51 promote telomere maintenance in Ustilago maydis, a new model system of DNA repair proteins at telomeres. DNA Repair (Amst) *12*, 472-479.

Yu, E.Y., Perez-Martin, J., Holloman, W.K., and Lue, N.F. (2015). Mre11 and Blm-Dependent Formation of ALT-Like Telomeres in Ku-Deficient Ustilago maydis. PLoS Genet *11*, e1005570.

Yu, E.Y., Zahid, S.S., Ganduri, S., Sutherland, J.H., Hsu, M., Holloman, W.K., and Lue, N.F. (2020). Structurally distinct telomere-binding proteins in Ustilago maydis execute non-overlapping functions in telomere replication, recombination, and protection. Commun Biol *3*, 777.

Zahid, S., Aloe, S., Sutherland, J.H., Holloman, W.K., and Lue, N.F. (2022). Ustilago maydis telomere protein Pot1 harbors an extra N-terminal OB fold and regulates homology-directed DNA repair factors in a dichotomous and context-dependent manner. PLoS Genet *18*, e1010182.

Zimmermann, M., Kibe, T., Kabir, S., and de Lange, T. (2014). TRF1 negotiates TTAGGG repeat-associated replication problems by recruiting the BLM helicase and the TPP1/POT1 repressor of ATR signaling. Genes & development *28*, 2477-2491.

S1 Table. *U. maydis* strains used in this study

Alias (Haploids)	*Relevant Genotype*	Reference
FB1[a]	Wild type	(Banuett and Herskowitz, 1989)
USZ111 [ab]	*trf2 ^{crg1}*	This work
UEY25 [ac]	*tay1Δ*	Yu et al., 2020
USZ112 [abc]	*tay1Δ trf2 ^{crg1}*	This work
UCM350 [d]	wild type	Kojic et al., 2002
USZ100 [bd]	*trf2 ^{crg1}*	Yu et al., 2020
USZ121 [bde]	*blmΔ trf2^{crg1}*	This work
USZ122 [bdf]	*dna2Δ trf2^{crg1}*	This work
USZ123 [bdg]	*exo1Δ trf2^{crg1}*	This work
USZ124 [bdh]	*mre11Δ trf2^{crg1}*	This work
USZ125 [bdi]	*trf2^{crg1} rad51Δ*	This work
USZ131 [j]	*blm^{crg1}*	This work
USZ132 [j]	*blm^{crg1}*	This work
USZ141 [k]	*blm^{nar1}*	This work
USZ142 [bk]	*trf2^{crg1} blm^{nar1}*	This work
USZ143 [kl]	*pot1^{crg1} blm^{nar1}*	This work

[a] The genotype of FB1 is *a1b1* for the mating type loci.
[b] *trf2* was placed downstream of the arabinose-dependent *crg1* promoter through the introduction of a Cbx^R-containing cassette.
[c] *tay1* was disrupted by the insertion of *hph* cassette expressing the hygromycin resistance gene *(Hyg^R).*
[d] The genotype of UCM350 *is nar1-6 pan1-1 a1b1*. *nar*, *pan*, and *ab* indicate inability to reduce nitrate, auxotrophic requirement for pantothenate, and mating type loci, respectively.
[e] *blm* was disrupted by the insertion of *hph* cassette expressing the hygromycin resistance gene *(Hyg^R).*
[f] *dna2* was disrupted by the insertion of *hph* cassette expressing the hygromycin resistance gene *(Hyg^R).*
[g] *exo1* was disrupted by the insertion of *hph* cassette expressing the hygromycin resistance gene *(Hyg^R).*
[h] *mre11* was disrupted by the insertion of *hph* cassette expressing the hygromycin resistance gene *(Hyg^R).*
[i] *rad51* was disrupted by the insertion of *hph* cassette expressing the hygromycin resistance gene *(Hyg^R).*
[j] *blm* was placed downstream of the arabinose-dependent *crg1* promoter through the introduction of a Cbx^R-containing cassette.
[k] *blm* was placed downstream of the nitrate-dependent *nar1* promoter through the introduction of a Hyg^R-containing cassette.
[l] *pot1* was placed downstream of the arabinose-dependent *crg1* promoter through the introduction of a Cbx^R-containing cassette.

S2 Table. Oligos used in this study (Trf2 paper new)

Name	Sequence 5' to 3'
Protein Expression	
UmTrf2-416F-SalI	ATTAC GTCGAC CG CAA TCC GAA CAG CGG TTA
UmTrf2-dn-st-NotI	AA GCGGCCGC CTA TTC GTT TGA AAG AGA ACT CGA CG
UmBlm-F-Nco	AAT CCATGG CA ATG CCG CAA TCC GCA CTA ACC CCA
UmBlm-R-FG-NotI	AAT GCGGCCGC CTA CTT GTC ATC GTC ATC CTT GTA ATC ACC CGA ACG AGG TAG ATT GGG C
Strain Construction and Genotyping	
UmBlm(1)Nde	GAAATC CAT ATG CCG CAA TCC GCA CTA
UmBlm(720R)Xba	ATT TCTAGA CAA GAT CTG AAA CGA GCT CG
UmBlm(-720)Xba	ATA TCTAGA TTG GAA CTG CGC AGT AAG CT
UmBlm(-1R)Eco	ATT GAATTC AAC TGA AAG AAG GCA ATC CT
UmBlm(720R)Nsi	AGA ATGCAT CAA GAT CTG AAA CGA GCT CG
UmBlm(-720)Nsi	ATA ATGCAT TTG GAA CTG CGC AGT AAG CT
Helicase assays	
NT – top	TTCTTCCTTTCCCTCTTCCTGATACGGCTGCTTCTCATCTACAACGTGATCCGTCATGGT
NT – bottom	ATGAGAAGCAGCCGTATCAGGAAGAGGGAAAGGAAGAA
Telo – top	TTCTTCCTTTCCCTC**TAGGGTTAGGGTTAGGGTTAGGGTTAGGGTTAGGGTTAGGGTTAG**
Telo – bottom	**CCCTAACCCTAACCCTAACCCTA**GAGGGAAAGGAAGAA
PCR, hybridization, and EMSA assays	
TTAGGG$_4$ (G4)	TTAGGG TTAGGG TTAGGG TTAGGG
CCCTAA$_4$ (C4)	CCCTAA CCCTAA CCCTAA CCCTAA
TTAGGG$_8$ (G8)	TTAGGG TTAGGG TTAGGG TTAGGG TTAGGG TTAGGG TTAGGG TTAGGG
CCCTAA$_8$ (C8)	CCCTAA CCCTAA CCCTAA CCCTAA CCCTAA CCCTAA CCCTAA CCCTAA
UmrRNA_26S_121F	GCTTCGGACCATGCCTAAG
UmrRNA_26S_642R	CTTGGTCCGTGTTTCAAGACG
RT-PCR and RT-qPCR	
UT6-TERRA-F1	GGACGGCAGATATATATTGTGAGTGG
UT6-TERRA-F2	GTGGCAACATTGGGTGAGC
UT6-TERRA-R1	CCGTTGACACATTCAATCCCTC
UT6-TERRA-R2	CTTCAAGCCCTGCAGCC
CCCTAA$_4$ (C4)	CCCTAA CCCTAA CCCTAA CCCTAA
Blm-PCR-180F	CAC TGC TCA AAA GAG ATC GAG GTC TGG
blm-PCR-209F	CAT CGG CCG CTT CCA TCT CAA ACG

blm-PCR-305R	CGA GGT GTC TGC GAT ATC GAG CCA C
Blm-PCR-400R	CAA AAT CGA TCC GAA GCT CGT CTT CG
Blm-PCR-500R	AAG GCT TCG TGC TTT TGG TAC CAA G
Trf2-PCR-1100R	TCT TCC GAG CCT TCA TCC GA
Trf2-PCR-865-F	CGC GAC ACC AAC CAT ACA AGC A
Trf2-PCR-939-R	GGT GAG TGG GGC ACT GTG TC
STELA and fusion assays	
UT4-F	TCGGGCAACGTTCCATGTCG
UT4-subtel-R2375	CCCTCGAAGGCAGTGCATAC
UT6-F	CTACTACACATCGGTTCAGGC
UT6-subtel-R2400	ATGCCAAAGTGGAAATCGTGCAC
C Telorette 1	GCTCCGTGCATCTGGCATCCCCTAAC
C Telorette 2	GCTCCGTGCATCTGGCATCTAACCCT
C Telorette 3	GCTCCGTGCATCTGGCATCCCTAACC
C Telorette 4	GCTCCGTGCATCTGGCATCCTAACCC
C Telorette 5	GCTCCGTGCATCTGGCATCAACCCTA
C Telorette 6	GCTCCGTGCATCTGGCATCACCCTAA
Teltail	GCTCCGTGCATCTGGCATC

Figure captions

Fig. 1. The roles of specific DNA repair factors in causing growth defects and telomere aberrations in Trf2-deficient U. maydis cells.

A. Serial dilutions of the indicated strains were spotted onto YPA and YPD media and incubated at 30°C. Following 2 and 3 days of growth, the plates were imaged.

B. Genomic DNAs were isolated from the indicated strains grown in YPA or YPD, digested with PstI, and subjected to TRF Southern analysis.

C. Genomic DNAs from the indicated strains grown in YPA or YPD were subjected to C-circle analysis.

D. Genomic DNAs were isolated from the indicated strains grown in YPA or YPD, digested with PstI, and subjected to in-gel hybridization to assess the levels of G-strand ssDNA (top left panel) or C-strand ssDNA (top right panel). The gels were then denatured and re-hybridized with the same probes to determine the overall telomere content (bottom panels).

Fig. 2 Epistasis analysis of telomere shortening pathways mediated by Trf2-overexpression and DNA repair factor deficiencies

A. and B. Genomic DNAs were isolated from the indicated strains grown in YPA, digested with PstI, and subjected to TRF Southern analysis.

C. STELA analysis of telomere length distributions in UCM350, $trf2^{crg1}$, and $trf2^{crg1}$ $blm\Delta$ strains grown in YPA. The Southern blot is shown at the top and the estimated STELA fragment lengths summarized at the bottom.

Figure 3. Blm-Trf2 interaction *in vitro* and the effects of Trf2 on Blm helicase activity

A. The elution samples from a Blm-FG/His-SUMO-Trf2$^{\Delta N}$ pull down experiment were analyzed by Western using anti-FLAG (left panel) and anti-His (right panel) antibodies, respectively. For comparison, the His-SUMO-Trf2$^{\Delta N}$ input fraction in the pull down assays were also analyzed. Note that Blm-FG cross-reacted with the anti-His antibodies.

B. The effects of Trf2$^{\Delta N}$ (175 nM, 350 nM and 700 nM) on Blm helicase activity were analyzed using substrates with and without telomeric repeats (left and right panels, respectively). Following incubation, the reaction products were analyzed by electrophoresis with or without prior treatment with SDS/EDTA (to dissociate protein-DNA complex). The bands corresponding to the duplex DNA substrates, ssDNA products and protein-DNA complexes are marked by schematic illustrations to the right of the panels.

C. The assay results in B were quantified and plotted.

Figure 4. Blm overexpression triggers growth defects and telomere deprotection

A. RT-qPCR analysis of Blm RNA levels in the parental strain and blm^{crg1} mutants.

B. Serial dilutions of the indicated *U. maydis* strains were spotted onto YPA or YPD plates and incubated at 30 °C. The plates were imaged following 2 days of growth.

C. Chromosomal DNAs from the indicated strains grown in the specified media were subjected to TRF Southern analysis.

D. Genomic DNAs from the designated strains grown in the specified media were tested for C-circle levels.

E. Genomic DNAs from the designated strains grown in the specified media were assessed for the levels of G- and C-strand ssDNA using in-gel hybridization analysis (left panels). The gels were also denatured and re-probed to characterize the distribution of all telomere-containing fragments.

F. The levels of ssDNA from the in-gel hybridization assays were quantified and plotted.

Figure 5. The consequences of overexpressing Blm, Trf2 and both

A. The transcription activities of the *crg1* and *nar1* promoter in different media are schematically illustrated.

B. Genomic DNAs from the designated strains grown in YPA were subjected to C-circle analysis. All the reactions (with and without φ29 polymerase) were performed in duplicates.

C. Genomic DNAs from the designated strains grown in MMD and MMA were subjected to TRF Southern analysis. The extra-short telomeres triggered by either Blm overexpression of Trf2 and Pot1 repression are marked by a bracket to the right of the blot.

Figure 6. The relationships between Tay1 and Trf2 in telomere protection

A. Serial dilutions of the indicated *U. maydis* strains were spotted onto YPA or YPD plates and incubated at 30 °C. The plates were imaged following 3 days of growth.

B. DNAs from the designated strains grown in the specified media were subjected to TRF Southern analysis.

Fig. 7. Elevated telomere G- and C-strand RNA synthesis in Blm-overexpression mutants

A. RNAs were prepared from UCM and *blm^{crg1}* grown in YPA or YPD and spotted onto nylon filters with and without prior RNase A digestion. The filters were probed sequentially for G-strand RNA, C-strand RNA, and rRNAs.

B. RNAs were isolated from UCM and *blm^{crg1}* grown in YPA or YPD, and subjected to RT-qPCR analysis to quantify the levels of G- and C-strand RNAs. For the G-strand RNA, the primers were designed to amplify TERRA (RNAs spanning subtelomeres and telomeres) from the UT6-bearing telomeres. For C-strand RNA, the primers can in principle amplify both C-TERRA (RNAs spanning subtelomeres and telomeres) and ARRET (RNAs with subtelomere sequences only) from UT6 telomeres.

C. RNAs were isolated from the indicated strains grown in the specified media, and subjected to endpoint RT-PCR analysis to characterize the structures of the G- and C-strand RNAs.

Supplementary Figure Legends

Supp. Fig. 1. Growth defects of the *trf2^{crg1} blmΔ* and *trf2^{crg1} mre11Δ* mutants

A. Serial dilutions of the indicated strains were spotted onto YPD medium. The growth of the strains was imaged after 2 days and 3 days.

B. The indicated strains were inoculated into fresh YPA at an OD_{600} of 0.01 and grown at 30 degree for 17 hours. The fold increases in OD_{600} for the cultures were determined and plotted.

Supp. Fig. 2 Phenotypic analysis of Trf2-deficient cells derived from the FB1 strain background

A. Serial dilutions of the indicated strains were spotted onto YPA and YPD media and incubated at 30°C. Following 2 days of growth, the plates were imaged.

B. Genomic DNAs from the indicated strains were subjected to TRF Southern analysis.

Supp. Fig. 3. A progressive reduction in Trf2 expression level correlates with a gradual increase in telomere lengths

A. UCM and *trf2^{crg1}* were grown in media containing 1.5% arabinose and varying concentrations of glucose (to reduce the activity of the crg1 promoter). RNAs were isolated from these cultures and subjected to RT-qPCR analysis to determine the relative *trf2* RNA levels.

B. Genomic DNAs were isolated from UCM and *trf2^{crg1}* grown in media with the specified arabinose and glucose concentrations and subjected to TRF Southern analysis.

Supp. Fig. 4. Blm overexpression triggers ECTRs in a reversible manner.

A. DNAs were isolated from *blm^{crg1}* strains grown in the specified culture media and subjected to Southern analysis for telomere DNA with and without prior *Pst*I digestion.

B. DNAs were isolated from UCM and *blm^{crg1}* strains grown in the designated culture media and subjected to Southern analysis. To test the reversibility of telomere defects induced by Blm overexpression, the *blm^{crg1}* clones were first grown on a YPA plate and then re-streaked once or twice on YPD plates prior to liquid culture growth and genomic DNA isolation (designated by YPA -> YPD and YPA -> YPD -> YPD, respectively).

Supp. Fig. 5. Partial suppression of Trf2-deficiency phenotypes by *tay1Δ*.

A. Serial dilutions of the designated strains were spotted onto semi-solid media containing the specified combinations of arabinose and glucose, and grown for 2 or 3 days at 30°C.

B. Genomic DNAs were isolated from the indicated strains grown in 1.5% arabinose and 1 % glucose, and subjected to TRF Southern analysis.

Supp. Figure 6. Up-regulation of telomere G- and C-strand RNA synthesis in Trf2-deficient mutants

A. RNAs were isolated from the indicated strains grown in YPD and spotted onto nylon membrane. After crosslinking, the membrane was probed sequentially for G-strand RNA, C-strand RNA, and rRNAs.

B. RNAs were isolated from the indicated strains grown in YPD, and subjected to RT-qPCR analysis to quantify the levels of G- and C-strand RNAs. For the G-strand RNA, the primers were designed to amplify TERRA (RNAs spanning subtelomeres and telomeres) from the UT6-bearing telomeres. For C-strand RNA, the primers can in principle amplify both C-TERRA (RNAs spanning subtelomeres and telomeres) and ARRET (RNAs with subtelomere sequences only) from UT6 telomeres.

Fig. 1

Fig. 1. The roles of specific DNA repair factors in causing growth defects and telomere aberrations in Trf2-deficient U. maydis cells.
A. Serial dilutions of the indicated strains were spotted onto YPA and YPD media and incubated at 30°C. Following 2 and 3 days of growth, the plates were imaged.
B. Genomic DNAs were isolated from the indicated strains grown in YPA or YPD, digested with PstI, and subjected to TRF Southern analysis.
C. Genomic DNAs from the indicated strains grown in YPA or YPD were subjected to C-circle analysis.
D. Genomic DNAs were isolated from the indicated strains grown in YPA or YPD, digested with PstI, and subjected to in-gel hybridization to assess the levels of G-strand ssDNA (top left panel) or C-strand ssDNA (top right panel). The gels were then denatured and re-hybridized with the same probes to determine the overall telomere content (bottom panels).

Fig. 2

Fig. 2 Epistasis analysis of telomere shortening pathways mediated by Trf2-overexpression and DNA repair factor deficiencies
A. and B. Genomic DNAs were isolated from the indicated strains grown in YPA, digested with PstI, and subjected to TRF Southern analysis.
C. STELA analysis of telomere length distributions in UCM350, *trf2crg1*, and *trf2crg1 blmΔ* strains grown in YPA. The Southern blot is shown at the top and the estimated STELA fragment lengths summarized at the bottom.

Fig. 3

Figure 3. Blm-Trf2 interaction *in vitro* and the effects of Trf2 on Blm helicase activity
A. The elution samples from a Blm-FG/His-SUMO-Trf2$^{\Delta N}$ pull down experiment were analyzed by Western using anti-FLAG (left panel) and anti-His (right panel) antibodies, respectively. For comparison, the His-SUMO-Trf2$^{\Delta N}$ input fraction in the pull down assays were also analyzed. Note that Blm-FG cross-reacted with the anti-His antibodies.
B. The effects of Trf2$^{\Delta N}$ (175 nM, 350 nM and 700 nM) on Blm helicase activity were analyzed using substrates with and without telomeric repeats (left and right panels, respectively). Following incubation, the reaction products were analyzed by electrophoresis with or without prior treatment with SDS/EDTA (to dissociate protein-DNA complex). The bands corresponding to the duplex DNA substrates, ssDNA products and protein-DNA complexes are marked by schematic illustrations to the right of the panels.
C. The assay results in B were quantified and plotted.

Fig. 4

Figure 4. Blm overexpression triggers growth defects and telomere deprotection
A. RT-qPCR analysis of Blm RNA levels in the parental strain and *blm^crg1* mutants.
B. Serial dilutions of the indicated *U. maydis* strains were spotted onto YPA or YPD plates and incubated at 30 °C. The plates were imaged following 2 days of growth.
C. Chromosomal DNAs from the indicated strains grown in the specified media were subjected to TRF Southern analysis.
D. Genomic DNAs from the designated strains grown in the specified media were tested for C-circle levels.
E. Genomic DNAs from the designated strains grown in the specified media were assessed for the levels of G- and C-strand ssDNA using in-gel hybridization analysis (left panels). The gels were also denatured and re-probed to characterize the distribution of all telomere-containing fragments.
F. The levels of ssDNA from the in-gel hybridization assays were quantified and plotted.

Figure 5. The consequences of overexpressing Blm, Trf2 and both
A. The transcription activities of the *crg1* and *nar1* promoter in different media are schematically illustrated.
B. Genomic DNAs from the designated strains grown in YPA were subjected to C-circle analysis. All the reactions (with and without φ29 polymerase) were performed in duplicates.
C. Genomic DNAs from the designated strains grown in MMD and MMA were subjected to TRF Southern analysis. The extra-short telomeres triggered by either Blm overexpression of Trf2 and Pot1 repression are marked by a bracket to the right of the blot.

Fig. 6

Figure 6. The relationships between Tay1 and Trf2 in telomere protection
A. Serial dilutions of the indicated *U. maydis* strains were spotted onto YPA or YPD plates and incubated at 30 °C. The plates were imaged following 3 days of growth.
B. DNAs from the designated strains grown in the specified media were subjected to TRF Southern analysis.

Fig. 7

Fig. 7. Elevated telomere G- and C-strand RNA synthesis in Blm-overexpression mutants

A. RNAs were prepared from UCM and *blm^crg1* grown in YPA or YPD and spotted onto nylon filters with and without prior RNase A digestion. The filters were probed sequentially for G-strand RNA, C-strand RNA, and rRNAs.

B. RNAs were isolated from UCM and *blm^crg1* grown in YPA or YPD, and subjected to RT-qPCR analysis to quantify the levels of G- and C-strand RNAs. For the G-strand RNA, the primers were designed to amplify TERRA (RNAs spanning subtelomeres and telomeres) from the UT6-bearing telomeres. For C-strand RNA, the primers can in principle amplify both C-TERRA (RNAs spanning subtelomeres and telomeres) and ARRET (RNAs with subtelomere sequences only) from UT6 telomeres.

C. RNAs were isolated from the indicated strains grown in the specified media, and subjected to endpoint RT-PCR analysis to characterize the structures of the G- and C-strand RNAs.

Supp. Fig. 1

A

Supp. Fig. 1. Growth defects of the *trf2^{crg1} blmΔ* and *trf2^{crg1} mre11Δ* mutants
A. Serial dilutions of the indicated strains were spotted onto YPD medium. The growth of the strains was imaged after 2 days and 3 days.
B. The indicated strains were inoculated into fresh YPA at an OD_{600} of 0.01 and grown at 30 degree for 17 hours. The fold increases in OD_{600} for the cultures were determined and plotted.

Supp Fig. 2

A

B

Supp. Fig. 2 Phenotypic analysis of Trf2-deficient cells derived from the FB1 strain background
A. Serial dilutions of the indicated strains were spotted onto YPA and YPD media and incubated at 30°C.
Following 2 days of growth, the plates were imaged.
B. Genomic DNAs from the indicated strains were subjected to TRF Southern analysis.

Supp. Fig. 3

A

B

Supp. Fig. 3. A progressive reduction in Trf2 expression level correlates with a gradual increase in telomere lengths

A. UCM and *trf2crg1* were grown in media containing 1.5% arabinose and varying concentrations of glucose (to reduce the activity of the crg1 promoter). RNAs were isolated from these cultures and subjected to RT-qPCR analysis to determine the relative *trf2* RNA levels.

B. Genomic DNAs were isolated from UCM and *trf2crg1* grown in media with the specified arabinose and glucose concentrations and subjected to TRF Southern analysis.

Supp. Fig. 4

Supp. Fig. 4. Blm overexpression triggers ECTRs in a reversible manner.
A. DNAs were isolated from *blm^crg1* strains grown in the specified culture media and subjected to Southern analysis for telomere DNA with and without prior *Pst*I digestion.
B. DNAs were isolated from UCM and *blm^crg1* strains grown in the designated culture media and subjected to Southern analysis. To test the reversibility of telomere defects induced by Blm overexpression, the *blm^crg1* clones were first grown on a YPA plate and then re-streaked once or twice on YPD plates prior to liquid culture growth and genomic DNA isolation (designated by YPA -> YPD and YPA -> YPD -> YPD, respectively).

Supp. Fig. 5. Partial suppression of Trf2-deficiency phenotypes by *tay1*Δ.
A. Serial dilutions of the designated strains were spotted onto semi-solid media containing the specified combinations of arabinose and glucose, and grown for 2 or 3 days at 30°C.
B. Genomic DNAs were isolated from the indicated strains grown in 1.5% arabinose and 1 % glucose, and subjected to TRF Southern analysis.

Supp. Fig. 6

A

B

Supp. Figure 6. Up-regulation of telomere G- and C-strand RNA synthesis in Trf2-deficient mutants
A. RNAs were isolated from the indicated strains grown in YPD and spotted onto nylon membrane. After crosslinking, the membrane was probed sequentially for G-strand RNA, C-strand RNA, and rRNAs.

B. RNAs were isolated from the indicated strains grown in YPD, and subjected to RT-qPCR analysis to quantify the levels of G- and C-strand RNAs. For the G-strand RNA, the primers were designed to amplify TERRA (RNAs spanning subtelomeres and telomeres) from the UT6-bearing telomeres. For C-strand RNA, the primers can in principle amplify both C-TERRA (RNAs spanning subtelomeres and telomeres) and ARRET (RNAs with subtelomere sequences only) from UT6 telomeres.