Opportunistic evidence of the impact of bacterial infections on social integration in vampire

bats

Imran Razik[1,2], Sebastian Stockmaier[2,3], Michael Abou-Elias[1], Gerald G. Carter[1,2]

Affiliations

[(1)] Department of Evolution, Ecology and Organismal Biology, The Ohio State University, Columbus

OH, USA

[(2)] Smithsonian Tropical Research Institute, Balboa Ancon, Panama, Republic of Panama

[(3)] Department of Ecology and Evolutionary Biology, University of Tennessee, Knoxville, TN, USA

Corresponding author: Imran Razik (razik.2@buckeyemail.osu.edu)

Abstract

Social integration can affect an individual's susceptibility to infectious disease. Conversely,

infectious disease can reduce an individual's social activity. Yet, it remains unclear to what extent

short-term infections can inhibit aspects of long-term social integration (e.g., formation of new

relationships). During a previous study on relationship formation, we captured female common

vampire bats (*Desmodus rotundus*) from different wild populations and housed them together in

captivity. Upon introduction, we observed an unplanned outbreak of bacterial infections that caused

cutaneous lesions. After treating infected bats with an antibiotic, 5 bats recovered, but 4 others

suffered lasting injuries. We analyzed how allogrooming rates changed over time among familiar and

new pairs consisting of the 9 infected and 12 asymptomatic bats. Infected bats demonstrated

lethargy and reduced social behavior. We found that more severely infected bats gave and received

less allogrooming compared to asymptomatic bats, that this infection effect was initially larger for

new pairs relative to familiar pairs, and that this effect decreased as infected bats recovered and

new pairs became more familiar. These observed effects, although opportunistic and based on only

9 infected bats, were consistent with the hypothesis that short-term infections can impact

relationship formation when new individuals join a group.

Introduction

Social integration, the quantity and quality of an individual's supportive social relationships,

can affect that individual's susceptibility to infection (e.g.,[1–4]) and ability to tolerate the adverse

effects of parasites [5,6]. Conversely, infectious disease can often inhibit social behaviors of infected

individuals [7–9] by diminishing their motivation to interact with others (e.g.,[10–12]), reducing their

movement (e.g., in mice [13] and mongoose [14]), or causing others to avoid them [15–19]. The

effects of infection on social behavior vary across pathogens, hosts, levels of infection severity, and

types of social relationships. For example, in vampire bats, sickness behavior reduces non-kin social

grooming, but does not decrease maternal grooming of offspring [20]. Similarly, mandrills avoid

grooming parasite-infected non-kin but continue to groom infected close maternal kin [21]. More

generally, infections might reduce social interactions disproportionately more within new or weak

social relationships compared to strong social bonds [22]. For example, short-term infections are

likely to have a bigger impact on mate choice (reviewed in [23]) and on the process of pair bonding

than on an existing pair bond [24]. It remains unclear, however, to what extent infections impact

same-sex social bond formation.

New social bonds form when new adults join a social community, which is also when

individuals are exposed to new pathogens. Given that short-term events during relationship

formation can have persistent long-term effects [25], the impacts of infections may have

disproportionately greater long-term consequences on relationship formation than on existing

relationships. To address this hypothesis, we report opportunistic evidence that an acute bacterial

infection can reduce social integration in common vampire bats (*Desmodus rotundus*). In this

species, social bond formation is observable as strangers transition from clustering to social

grooming and regurgitated food sharing—a costly cooperative behavior [26]. Allogrooming is the

best measure of relationship formation, because it is a directed investment of time and energy that

also occurs frequently enough to track over time. During a previous 4-month study, we tracked

changes in allogrooming among 21 female vampire bats before and after a week of forced proximity

in which small groups of 3 bats from 3 distant sites were housed together for one week (25). Shortly

after the 21 bats from different sites were first introduced in captivity, we observed an unplanned

outbreak of a bacterial infection, which we treated with antibiotics (25). In this paper, we analyzed

the social effect of these infections by tracking changes in allogrooming over time among pairs of 9

infected female bats and the 12 asymptomatic female bats that were previously familiar or unfamiliar

with each other.

Methods

Study subjects

For an experiment testing the effects of forced proximity on relationship formation (25), we

captured 8-9 common vampire bats from each of three sites in Panama that were 120-350 km apart.

Upon capture, the bats (23 adult females, 2 juvenile males, and one juvenile female) appeared

healthy with no pathological lesions. We housed the bats together in an outdoor flight cage (2.1 × 1.7

× 2.3 m) for approximately 4 months. During this time, five females gave birth to four males and one

female pup (31 total bats). Details of the study colony can be found in (25,27).

We tracked the allogrooming relationships among 252 new and 126 familiar pairs of 21 adult

females. We define 'pairs' as unique directed combinations of a groomer and receiver. We use the

label 'new' for pairs that were introduced in captivity and 'familiar' for pairs that were captured from

the same site (hollow tree or cave). We excluded from our analysis 42 pairs that were forced into

proximity together to induce social bonding for the previous study; however, including these 42 pairs

in the analysis does not change our conclusions.

Behavior sampling

We used three infrared surveillance cameras to sample social interactions among the

individually marked bats (25). Over 114 days, we sampled 682 h of interactions, recording all visible

allogrooming and mouth-licking interactions (≥ 5 seconds in duration) as well as agonistic

interactions (≥ 1 second in duration). For each interaction, we noted the actor, receiver, and

duration. We used allogrooming to assess social integration because allogrooming was 94% of the

17,592 observed social interactions, while mouth-licking and aggression were only 6% and 0.3% of

interactions, respectively. To estimate dyadic allogrooming rates, we averaged the seconds of

allogrooming per hour across all hours in which the actor and receiver could interact. For further

details, see (25).

Bacterial infection and treatment

Two to three weeks after the 26 wild-caught bats were first introduced, 10 female bats began

to show progressive symptoms of a cutaneous infection (blisters or open lesions with pus, as well as

inflamed tissue), which was often localized around joints. Symptoms progressed for approximately

two weeks without signs of recovery. These infections were unexpected and to our knowledge have

not previously been described in vampire bat colonies. The 10 infected bats varied in the severity of

their symptoms. The most severe symptoms occurred in bats from a hollow tree near La Chorrera,

Panama. The most severely infected bat developed large, pus-filled lesions on its back and had

extreme inflammation around the wrists that led tissue to break apart and expose bone; we

euthanized this bat before it received antibiotic treatment, and we did not include it in our analysis of

social integration. Four other bats developed severe symptoms with ulcers and inflammation on one

or both wrists, leading to tissue scarring and the inability to sustain flight. One bat from the same

roost experienced only minor inflammation and ulcers on both wrists but healed without permanent

tissue scarring and remained able to sustain flight. Three bats from a cave near Lake Bayano,

Panama were also infected. Two bats developed pus-filled abscesses: on the lower abdomen and

upper thigh for one bat, on the lower back for the other bat, and the third bat developed only minor

inflammation on 1-2 toes. Finally, one bat from a hollow tree near Tolé, Panama had a large, round

lesion on its knee, but this was evident at the start of the study and did not change over time. This

bat displayed no other signs of infection. With most cases having been observed in bats from either

La Chorrera or Lake Bayano, it seems the bats from Tolé were most resistant to the pathogen.

Swab and tissue samples from skin lesions were cultured to isolate bacterial pathogens,

which indicated the isolation of *Bacillus sp.* and *Staphylococcus aureus*. A Kirby-Bauer antibiogram

indicated susceptibility of the isolated bacteria to the antibiotic Enrofloxacin. Given the urgency and

results of the antibiogram, we immediately used Enrofloxacin antibiotic to treat the infected female

bats (described above) over the course of seven days via daily subcutaneous injections. We did not

perform additional testing to further identify the bacterial species or strain or test for other potentially

infectious pathogens. After the antibiotic treatment, five bats recovered without injury (henceforth

"treated"), but four bats suffered lasting injuries that rendered them unable to fly for the remainder of

the study (henceforth "injured"). We refer to all other bats in the colony as "asymptomatic".

Confirming sickness behavior

To confirm that bacterial infections reduced individual activity and social behavior, i.e.,

'sickness behavior', we compared the activity of 9 infected versus 9 asymptomatic adult female bats

during a sample period of 1 week before the antibiotic treatment (pre-treatment week) and again

during another 1 week about 11 weeks later near the end of study (post-treatment week). An

observer scored videos using focal sampling for the presence or absence of 4 behavioral states:

moving, resting (including sleeping), allogrooming (including mouth-licking), and self-grooming,

following definitions in (12). These behaviors were scored every 30 seconds for 1 hour (starting one

hour after sunset) per day for each bat. We sampled all 18 bats 120 times on each of 14 days. The

observer could not be entirely blind to the condition because lesions were visible, but videos are

available for confirmation. To test if infection predicted the frequency of these six behavioral states,

we calculated the mean proportion of sampled times where each behavior occurred and estimated

95% confidence intervals (CIs) using bootstrapping with 5000 iterations (percentile method, boot

package in R, 28).

Effects of infection on dyadic allogrooming rates

To assess how dyadic rates of allogrooming changed over time among familiar or new pairs

of bats that were either infected or asymptomatic, we fit a logistic mixed effect model (lme4 R

package, 29). The response was the presence or absence of allogrooming for each groomer-

receiver pair and hour. Fixed effects were prior familiarity of the pair (true/false), infection severity of

the actor (0 = asymptotic, 1 = infected, 2 = injured), infection severity of the receiver (same scale),

time (unit was 10 days), interactions between time and other fixed effects, interactions between

familiarity and infection severity, and two three-way interactions between groomer/receiver infection

severity, familiarity, and time. Groomer and receiver identity were random intercepts. We converted

each model coefficient to an odds ratio (OR) and used the profile likelihood method to estimate 95%

confidence intervals around ORs. When estimating confidence intervals via bootstrapping, the 95%

CIs were very similar and the conclusions were the same..

We predicted a three-way interaction between familiarity, time, and infection severity (for

both groomers and receivers), because we expected that the impact of infection on the slope of

allogrooming rates over time would differ between familiar pairs and new pairs. We expected

infections to have a greater impact on allogrooming rates in new pairs relative to familiar pairs. In

new pairs, we predicted an interaction between infection severity and time as predictors of

allogrooming, because infections should cause allogrooming rates to be initially lower during

relationship formation and then allogrooming rates of infected and asymptotic bats should converge

over time as the infected bats recovered. In familiar pairs, we expected this interaction to be weaker

because relationships were already formed, so infections should have a smaller impact on how

allogrooming changes over time in more established relationships.

Ethics

This work was approved by the Smithsonian Tropical Research Institute Animal Care and

Use Committee (no. 2015-0501-2022) and the Panamanian Ministry of the Environment (no. SEX/A-

67-2019).

Results

Confirming sickness behavior

Consistent with sickness behavior and recovery, the 9 infected bats spent less time moving and allogrooming, but these differences faded with time (Fig. 1). Before antibiotic treatment, infected bats spent 1.3% of their time allogrooming, 95% CI = [0.6, 2.1%], versus 3.9% [2.3, 5.6%] in asymptomatic bats. Infected bats spent 7.4% [4.8, 9.9%] of their time moving versus 17% [11, 22%] in asymptomatic bats. After recovery, however, the previously infected bats no longer showed clear differences from the asymptomatic bats in allogrooming or movement (Fig. 1). The pre-treatment differences from the asymptomatic bats in allogrooming and moving were also greater in the 4 injured bats than in the 5 treated bats (Fig. S1). We did not detect clear effects of infection on resting or self-grooming (Fig. 1).

Figure 1. Bats with bacterial infections spent less time allogrooming and moving pre-treatment but not post-treatment. Dots and error bars show mean proportions of time doing each behavior ± bootstrapped 95% CIs for infected bats (red triangle) and asymptomatic bats (black circle) female bats during the outbreak (pre-treatment) and after they had recovered (post-treatment). Resting and self-grooming did not clearly differ by infection. Fig. S1 shows a version of this plot with infected bats divided into 'treated' and 'injured' bats.

Effects of infection on dyadic allogrooming rates

        In vampire bats, allogrooming is often mutual and highly symmetrical within pairs over time (26,31,32). We found no evidence that infection removed allogrooming symmetry: pairwise allogrooming was bidirectional within the hour in 84% of cases with an infected bat and in 81% of

cases with two asymptomatic bats. This confirms our expectation that any sickness effects on

allogrooming given and allogrooming received will be causally linked (i.e., grooming given causes

grooming received), and that the behavior of both bats should influence the dyadic allogrooming

rate. This justifies our decision to include infection severity of both groomers and receivers as

simultaneous predictors with additive effects in our statistical model. After detecting a three-way

interaction between infection severity, familiarity, and time for both allogrooming given and received

(given: OR = 1.04, 95% CI = [1.03, 1.06], $p < 0.0001$; received: OR = 1.03 [1.01, 1.05], $p = 0.002$),

we fit separate models for new and familiar pairs to compare coefficients (Fig. 2).

In new pairs, an interaction between infection severity and time showed that infections

affected how allogrooming rates changed over time (interaction effect on grooming given: OR = 1.06

[1.05, 1.07], $p < 0.001$; on grooming received: OR = 1.05 [1.04, 1.06], $p < 0.001$). Specifically,

infections initially reduced allogrooming rates during relationship formation, then the allogrooming

rates of infected and asymptomatic bats converged over time as the infected bats recovered (Fig. 3).

Overall, allogrooming rates in new pairs were reduced by infections (grooming given: OR = 0.45

[0.25, 0.83], $p = 0.007$; grooming received: OR = 0.49 [0.26, 0.92], $p = 0.019$).

In familiar pairs, by contrast, the interaction between infection severity and time was weaker

(interaction effect on grooming given: OR = 1.02 [95% CI: 1.00, 1.03], $p = 0.02$; on grooming

received: OR = 1.03 [1.01, 1.04], $p = 0.001$; Fig. 2). Relative to infections in new pairs, infections in

familiar pairs had a smaller impact on how allogrooming rates changed over time (Fig. 3). Unlike the

new pairs, we did not detect a clear overall effect of infections on allogrooming given or received

(grooming given: OR = 0.80 [0.37, 1.82], $p = 0.57$; received: OR = 0.80 [0.38, 1.78], $p = 0.55$).In

sum, we found evidence that bacterial infections decreased allogrooming given and received, that

this effect of sickness behavior was greater for new pairs relative to familiar pairs, and that it also

decreased with time as infected bats recovered (Fig. 2).

Figure 2. The effect of infection severity on allogrooming over time differed between new and familiar pairs. Points show odds ratios, and profile likelihood 95% CIs, from generalized linear mixed effect models fit for both new pairs and unfamiliar pairs.

Figure 3. Social integration by infection status. Left panel shows changes in the probability of

receiving allogrooming over time from a familiar groomer when the receiver was asymptomatic (solid

cyan), treated (dashed orange), or injured (dotted red). Yellow shaded region indicates the week

prior to antibiotic treatment (pre-treatment) and blue shaded region indicates one week at the end of

the experiment (post-treatment). Middle panel shows the same plot but for previously unfamiliar

groomers (new pairs). The probability of receiving allogrooming prior to the antibiotic treatment was

reduced in infected bats, and especially the injured bats. Right panel shows the change in log odds

(pre-treatment vs post-treatment) of allogrooming received for each type of receiver. Large points

with error bars are means and bootstrapped 95% CIs. Effects on allogrooming given and received

are causally linked through reciprocity. These plots assess social integration through allogrooming

received; results for allogrooming *given* are even more dramatic (Fig. 2).

Discussion

While several studies have quantified the effect of pathogens on host social behavior (14,16–

18,46–51), it is unclear to what extent short-term infections can impact long-term social bond

formation. We present opportunistic observational evidence that acute symptoms of bacterial

infection can limit allogrooming relationship formation in vampire bats. Relative to asymptomatic

bats, the initially infected bats (and the injured bats in particular) spent less time moving and

allogrooming. Infected bats gave and received less allogrooming with new partners, but we did not

detect that infection severity predicted less allogrooming with familiar partners. These differences

faded during 3 months of recovery, suggesting they were caused by short-term sickness behavior

rather than by permanent injuries that reduced movement. These observations, albeit opportunistic

and sparse, are valuable for informing future experiments with greater power and control.

Because we did not randomly and experimentally infect the bats, a possible alternative

explanation is the reverse causal path: bats with worse social integration (or worse social

competence) had weaker immune systems and therefore suffered worse infections. However, this

hypothesis does not explain why the gap in social integration disappeared over time after antibiotic

treatment. If the infected bats were already less social or less healthy for reasons besides the

infection, then we would expect their social integration to be worse overall, not just during the

outbreak. If variation in the severity of infection was based on dominance or social competence, we

would expect variation in infection severity within each group. Instead, infection severity was almost

perfectly correlated with capture population, suggesting that infections were a cause, not a

consequence, of how bats formed new relationships.

The sickness behavior we observed was largely similar to the responses of vampire bats to

simulated infections from injections of lipopolysaccharide, LPS (12,20). In vertebrates, quantifying

sickness behaviors in response to a naturally-infecting pathogen is less common (but see

45,47,52,53) compared to sickness behaviors caused by an immune challenge like LPS (8,10–

12,15,20,24,31,54–56). The use of immune challenges allows for controlled experiments on the

connection between the inflammatory and the behavioral response to infection, but naturally

infecting pathogens can have different, pathogen-specific effects. For instance, LPS-injected

vampire bats reduced their self-grooming (12), but these infected bats did not, probably because the

bacterial pathogens caused skin lesions. Another pathogen that infects the skin, the fungus

Pseudogymnoascus destructans (which causes 'white-nose syndrome' in North American bats) also

leads to increased self-grooming (52).

The skin ulcers, abscesses, and inflammation we observed in common vampire bats were

due to an acute bacterial infection. Samples collected from symptomatic bats' lesions indicated the

presence of *Bacillus sp.* and *Staphylococcus aureus*. There are several *Bacillus* species (e.g.,

Bacillus anthracis, Bacillus cereus, and *Bacillus pumilus*) that may infect the skin or colonize

wounds, but these infections are uncommon and often present with classic symptoms (e.g. black

eschars and necrosis, 33–36) that we did not observe in our colony. Alternatively, several species

within the *S. aureus* complex, including *Staphylococcus schweitzeri* and *Staphylococcus argenteus,*

among others (37–39), typically infect the skin and cause symptoms that are consistent with what we

observed. *S. aureus* complex has also been isolated from the feces of Old-World fruit bats (40–43)

and the common vampire bat (44,45).

We view our results as a starting point for future studies that address the effect of infections

on long-term social integration and relationship formation. Our findings raise several questions.

When might vampire bats or other animals suppress their infection-induced behavioral symptoms to

continue social interactions and integration (54,55,57–59)? At what stages of relationship formation

do sickness behaviors have the strongest effect on social interactions? How long must sickness

behavior last for it to cause lingering social effects? To test for potential trade-offs between the

severity of behavioral symptoms and continued social interactions, researchers could use immune

challenges that are dose-dependent (55,60). Immune challenges like LPS could be used to measure

how long sickness behavior would have to last for lingering effects on social integration, to control

the duration and precise onset times of symptomatic periods, and to test what stage of relationship

formation is most impacted by sickness behaviors.

In conclusion, we found opportunistic observational evidence for reduced social integration in

infected vampire bats after an outbreak of a naturally infecting pathogen. Sickness behavior and

other cues of infection can reduce social activity and mate attraction (reviewed in 23), and there may

be similar effects on choice and formation of same-sex social bonds. If there are critical periods of

relationship formation, then infections during this time might have long-term social consequences.

Acknowledgements

The authors thank Rachel Page for lab space and resources, the staff at the Smithsonian Tropical

Research Institute, and the following individuals who assisted with vampire bat care and/or data

collection: D. Aparicio, B. Brown, G. Cohen, L. Dück, D. Girbino, E. Kline, C. Marroquin, and S.

Ripperger.

Funding

I.R. was supported by a short-term fellowship from the Smithsonian Tropical Research Institute, a

student research grant from the Animal Behaviour Society, a Graduate Enrichment Fellowship from

The Ohio State University, a Tinker Foundation Field Research Grant, a Grant-in-aid of Research

from the American Society of Mammalogists, and a Smithsonian Institution Predoctoral Fellowship.

S.S. was supported by the Ohio State University President's Postdoctoral Scholars Program. This

publication is based upon work supported by the National Science Foundation under grant no. IOS-

2015928 to G.C.

Author contributions

Conceptualization: IR, SS, GC

Data Curation: IR, GC

Formal Analysis: IR, SS, GC

Funding Acquisition: IR, GC

Investigation: IR, MA, GC

Methodology: IR, GC

Project Administration: IR, GC

Resources: IR, GC

Software: IR, GC

Supervision: GC

Validation: IR, SS, GC

Visualization: IR, SS, GC

Writing – Original Draft Preparation: IR, SS, GC

Writing – Review & Editing: IR, MA, SS, GC

References

1. Pavez-Fox MA, Negron-Del Valle JE, Thompson IJ, Walker CS, Bauman SE, Gonzalez O, et al.

Sociality predicts individual variation in the immunity of free-ranging rhesus macaques. Physiol

Behav. 2021;241:113560.

2. Snyder-Mackler N, Burger JR, Gaydosh L, Belsky DW, Noppert GA, Campos FA, et al. Social

determinants of health and survival in humans and other animals. Science.

2020;368(6493):eaax9553.

3. Snyder-Mackler N, Sanz J, Kohn JN, Brinkworth JF, Morrow S, Shaver AO, et al. Social status

alters immune regulation and response to infection in macaques. Science.

2016;354(6315):1041–5.

4. Tung J, Barreiro LB, Johnson ZP, Hansen KD, Michopoulos V, Toufexis D, et al. Social

environment is associated with gene regulatory variation in the rhesus macaque immune

system. Proc Natl Acad Sci. 2012;109(17):6490–5.

5. Almberg ES, Cross PC, Dobson AP, Smith DW, Metz MC, Stahler DR, et al. Social living

mitigates the costs of a chronic illness in a cooperative carnivore. Ecol Lett. 2015;18(7):660–7.

6. Loehle C. Social barriers to pathogen transmission in wild animal populations. Ecology.

1995;76(2):326–35.

7. Hart BL. Biological basis of the behavior of sick animals. Neurosci Biobehav Rev.

1988;12(2):123–37.

8. Lopes PC, French SS, Woodhams DC, Binning SA. Sickness behaviors across vertebrate taxa:

proximate and ultimate mechanisms. J Exp Biol. 2021;224(9):jeb225847.

9. Stockmaier S, Stroeymeyt N, Shattuck EC, Hawley DM, Ancel Meyers L, Bolnick DI. Infectious

diseases and social distancing in nature. Science. 2021;371(6533):eabc8881.

10. Fishkin RJ, Winslow JT. Endotoxin-induced reduction of social investigation by mice: interaction

with amphetamine and anti-inflammatory drugs. Psychopharmacol Berl. 1997;132(4):335–41.

11. Moreno KR, Weinberg M, Harten L, Salinas Ramos VB, Herrera M. LG, Czirják GÁ, et al. Sick

bats stay home alone: fruit bats practice social distancing when faced with an immunological

challenge. Ann N Y Acad Sci. 2021;1505(1):178–90.

12. Stockmaier S, Bolnick DI, Page RA, Carter GG. An immune challenge reduces social grooming

in vampire bats. Anim Behav. 2018;140:141–9.

13. Lopes PC, Block P, Konig B. Infection-induced behavioural changes reduce connectivity and the

potential for disease spread in wild mice contact networks. Sci Rep. 2016;6:31790.

14. Fairbanks BM, Hawley DM, Alexander KA. The Impact of Health Status on Dispersal Behavior in

Banded Mongooses (Mungos mungo). EcoHealth. 2014;11(2):258–62.

15. Lopes PC, Woodhams DC, Binning SA. Infection avoidance behaviors across vertebrate taxa:

patterns, processes and future directions. In: Animal Behavior and Parasitism. Oxford University

Press; 2022.

16. Behringer DC, Butler MJ, Shields JD. Ecology: avoidance of disease by social lobsters. Nature.

2006;441(7092):421.

17. Kiesecker JM, Skelly DK, Beard KH, Preisser E. Behavioral reduction of infection risk. Proc Natl

Acad Sci U A. 1999;96(16):9165–8.

18. Poirotte C, Massol F, Herbert A, Willaume E, Bomo PM, Kappeler PM, et al. Mandrills use

olfaction to socially avoid parasitized conspecifics. Sci Adv. 2017;3(4).

19. Boillat M, Challet L, Rossier D, Kan C, Carleton A, Rodriguez I. The Vomeronasal System

Mediates Sick Conspecific Avoidance. Curr Biol. 2015;25(2):251–5.

20. Stockmaier S, Bolnick DI, Page RA, Carter GG. Sickness effects on social interactions depend

on the type of behaviour and relationship. J Anim Ecol. 2020; 89:1387-1394.

21. Poirotte C, Charpentier MJE. Unconditional care from close maternal kin in the face of parasites.

Biol Lett. 2020;16(2):20190869.

22. Stockmaier S, Ulrich Y, Albery GF, Cremer S, Lopes PC. Behavioural defences against

parasites across host social structures. Funct Ecol. 2023; 37, 809– 820.

23. Beltran-Bech S, Richard FJ. Impact of infection on mate choice. Anim Behav. 2014;90:159–70.

24. Love AC, Anthony AC, Nash A, Campos-Melara A, Kodali J, DuRant SE. Simulated infection

alters the behavior of pair bonded songbirds and their asymptomatic neighbors. Behav Ecol.

2023;34(2):251–60.

25. Razik I, Brown BKG, Carter GG. Forced proximity promotes the formation of enduring

cooperative relationships in vampire bats. Biol Lett. 2022;18(4):20220056.

26. Carter GG, Farine DR, Crisp RJ, Vrtilek JK, Ripperger SP, Page RA. Development of New Food-

Sharing Relationships in Vampire Bats. Curr Biol. 2020;30(7):1275-1279.e3.

27. Razik I, Brown BKG, Page RA, Carter GG. Non-kin adoption in the common vampire bat. R Soc

Open Sci. 202;8(2):201927.

28. Canty A, Ripley B. boot: Bootstrap R (S-Plus) functions. R Package Version. 2017;1:3–20.

29. Douglas Bates MM. Fitting Linear Mixed-Effects Models Using lme4. J Stat Softw. 2015;67:1–48.

30. Razik I, Carter G, Stockmaier S, Abou-Elias M. Data and R code for Opportunistic evidence of

the impact of bacterial infections on social integration in vampire bats. 2023; Available from:

https://doi.org/ 10.6084/m9.figshare.22573018

31. Wilkinson GS. Social grooming in the common vampire bat, *Desmodus rotundus*. Anim Behav.

1986;34(6):1880–9.

32. Carter GG, Wilkinson GS. Food sharing in vampire bats: reciprocal help predicts donations more

than relatedness or harassment. Proc Biol Sci. 2013;280(1753):20122573.

33. Cromartie WJ, Bloom WL, Watson DW. Studies on Infection with *Bacillus anthracis*: I.

Histopathological Study of Skin Lesions Produced by *B. anthracis* in Susceptible and Resistant

Animal Species. J. Infect. Dis. 1947;80(1):1–13.

34. Esmkhani M, Shams S. Cutaneous infection due to *Bacillus cereus*: a case report. BMC Infect.

Dis. 2022;22(1);393.

35. Tena D, Martínez-Torres JA, Pérez-Pomata MT, Sáez-Nieto JA, Rubio V, Bisquert J. Cutaneous

Infection Due to *Bacillus pumilus*: Report of 3 Cases. Clin. Infect. Dis. 2007;44(4):e40–e42.

36. Veysseyre F, Fourcade C, Lavigne JP, Sotto A. *Bacillus cereus* infection: 57 case patients and a

literature review. Med Mal Infect 2015;45(11):436-440.

37. Becker K, Schaumburg F, Kearns A, Larsen AR, Lindsay JA, Skov RL, Westh H. Implications of

identifying the recently defined members of the *Staphylococcus aureus* complex *S. argenteus*

and *S. schweitzeri*: a position paper of members of the ESCMID Study Group for Staphylococci

and Staphylococcal Diseases (ESGS). Clin. Microbiol. Infect. 2019;25(9):1064–1070.

38. Chew KL, Octavia S, Lai D, Lin RTP, Teo JWP. *Staphylococcus singaporensis* sp. nov., a new

member of the *Staphylococcus aureus* complex, isolated from human clinical specimens. Int. J.

Syst. Evol. Microbiol. 2021;71(10):005067.

39. Tong SYC, Schaumberg F, Ellington MJ, Corander J, Pichon B, Leendertz F, Bentley SD,

Parkhill J, Holt DC, Peters G, Giffard PM. Novel staphylococcal species that form part of a

Staphylococcus aureus-related complex: the non-pigmented *Staphylococcus argenteus* sp. nov.

and the non-human primate-associated *Staphylococcus schweitzeri* sp. nov. Int. J. Syst. Evol.

Microbiol. 2015;65:15–22.

40. Fountain K, Barbon A, Gibbon MJ, Lloyd DH, Loeffler A, Feil EJ. *Staphylococcus aureus*

lineages associated with a free-ranging population of the fruit bat *Pteropus livingstonii* retained

over 25 years in captivity. Sci. Rep. 2022;12(1):13457.

41. Held J, Gmeiner M, Mordmüller B, Matsiégui P, Schaer J, Eckerle I, Weber N, Matuschewski K,

Bletz S, Schaumburg F. Bats are rare reservoirs of *Staphylococcus aureus* complex in Gabon.

Infect. Genet. Evol. 2017;47:118–120.

42. Monecke S, Schaumburg F, Shittu AO, Scharz S, Mühldorfer K, Brandt C, Braun SD, Collatz M,

Diezel C, Gawlik D, Hanke D, Hotzel H, Müller E, Reinicke M, Feßler AT, Ehricht R. Description

of Staphylococcal Strains from Straw-Coloured Fruit Bat (*Eidolon helvum*) and Diamond Firetail

(*Stagonopleura guttata*) and a Review of their Phylogenetic Relationships to Other

Staphylococci. Front. Cell. Infect. Microbiol. 2022;12:878137.

43. Olatimehin A, Shittu AO, Onwugamba FC, Mellmann A, Becker K, Schaumburg F.

Staphylococcus aureus Complex in the Straw-Colored Fruit Bat (*Eidolon helvum*) in Nigeria.

Front. Microbiol. 2018;9:162.

44. Chaverri G. Aerobic bacterial flora from the digestive tract of the common vampire bat,

Desmodus rotundus (Chiroptera: Phyllostomidae). Rev. Biol. Trop. 2006;54:717–724.

45. Monero M, Lopes CAM, Seabra E, Pavan C, Correa A. Aspectos bacteriologicos da flora

intestinal de *Desmodus rotundus*. Arq. Inst. Biol. São Paulo 1975;42:229.

46. Cárdenas-Canales EM, Stockmaier S, Cronin E, Rocke TE, Osorio JE, Carter GG. Social effects

of rabies infection in male vampire bats (*Desmodus rotundus*). Biol Lett. 2022;18(9):20220298.

47. Fairbanks BM, Hawley DM, Alexander KA. No evidence for avoidance of visibly diseased

conspecifics in the highly social banded mongoose (*Mungos mungo*). Behav Ecol Sociobiol.

2015;69(3):371–81.

48. Bouwman KM, Hawley DM. Sickness behaviour acting as an evolutionary trap? Male house

finches preferentially feed near diseased conspecifics. Biol Lett. 2010;6(4):462–5.

49. Stephenson JF. Parasite-induced plasticity in host social behaviour depends on sex and

susceptibility. Biol Lett. 2019;15(11):20190557.

50. Stroeymeyt N, Grasse AV, Crespi A, Mersch DP, Cremer S, Keller L. Social network plasticity

decreases disease transmission in a eusocial insect. Science. 2018;362(6417):941.

51. Cremer S. Social immunity in insects. Curr Biol. 2019;29(11):R458–63.

52. Brownlee-Bouboulis SA, Reeder DM. White-nose syndrome-affected little brown myotis (Myotis

lucifugus) increase grooming and other active behaviors during arousals from hibernation. J

Wildl Dis. 2013;49(4):850–9.

53. Bos N, Lefevre T, Jensen AB, d'Ettorre P. Sick ants become unsociable. J Evol Biol.

2012;25(2):342–51.

54. Lopes PC, Adelman J, Wingfield JC, Bentley GE. Social context modulates sickness behavior.

Behav Ecol Sociobiol. 2012;66(10):1421–8.

55. Willette AA, Lubach GR, Coe CL. Environmental context differentially affects behavioral,

leukocyte, cortisol, and interleukin-6 responses to low doses of endotoxin in the rhesus monkey.

Brain Behav Immun. 2007;21(6):807–15.

56. Alciatore G, Ugelvig LV, Frank E, Bidaux J, Gal A, Schmitt T, et al. Immune challenges increase

network centrality in a queenless ant. Proc R Soc B. 2021;288(1958):20211456.

57. Aubert A, Goodall G, Dantzer R, Gheusi G. Differential Effects of Lipopolysaccharide on Pup

Retrieving and Nest Building in Lactating Mice. Brain Behav Immun. 1997;11(2):107–18.

58. Lopes PC. When is it socially acceptable to feel sick? Proc Biol Sci. 2014;281(1788):20140218.

59. Weil ZM, Bowers SL, Dow ER, Nelson RJ. Maternal aggression persists following

lipopolysaccharide-induced activation of the immune system. Physiol Behav. 2006;87(4):694–9.

60. Armour EM, Bruner TL, Hines JK, Butler MW. Low-dose immune challenges result in detectable

levels of oxidative damage. J Exp Biol. 2020;223(6).