Opportunistic evidence of the impact of bacterial infections on social integration in vampire

bats

Imran Razik[1,2], Sebastian Stockmaier[2,3], Michael Abou-Elias[1], Gerald G. Carter[1,2]

Affiliations

[(1)] Department of Evolution, Ecology and Organismal Biology, The Ohio State University, Columbus

OH, USA

[(2)] Smithsonian Tropical Research Institute, Balboa Ancon, Panama, Republic of Panama

[(3)] Department of Ecology and Evolutionary Biology, University of Tennessee, Knoxville, TN, USA

Corresponding author: Imran Razik (razik.2@buckeyemail.osu.edu)

Abstract

Social integration can affect an individual's susceptibility to infectious disease. Conversely,

infectious disease can reduce an individual's social activity. Yet, it remains unclear to what extent

short-term infections can inhibit social integration and the formation of new relationships. During a

previous study on relationship formation, we captured 21 female common vampire bats (*Desmodus*

rotundus) from different wild populations and housed them together in captivity. Upon introduction,

we observed an unplanned outbreak of bacterial infections that caused cutaneous lesions. After

treating infected bats with an antibiotic, 5 bats recovered, but 4 others suffered lasting injuries. Given

that observations of how natural infections alter relationship formation are rare, we analyzed how

allogrooming rates changed over time among familiar and new dyads consisting of the 9 infected

and 12 asymptomatic bats. We found that (1) infected bats demonstrated reduced activity and social

behavior, (2) that more severely infected bats gave and received less allogrooming compared to

asymptomatic bats, (3) that the effect of infection was larger for new dyads relative to familiar dyads,

and (4) that this effect decreased as infected bats recovered and new dyads became more familiar.

These opportunistic observations were consistent with the hypothesis that short-term infections can

impact relationship formation when new individuals join a group.

Introduction

Social integration, or the quantity and quality of an individual's social relationships, can affect

an individual's susceptibility to infection (e.g.,1–4) and ability to tolerate or suppress the adverse

effects of parasites (5–7). Conversely, infectious disease can inhibit the social behaviors of infected

individuals (8–10) by diminishing their motivation to interact with others (e.g.,11–13), reducing their

movement (e.g., in mice (14) and mongoose (15)), or causing others to avoid them (16–21). These

feedbacks between social integration and infectious disease are complex in part because the effects

of infection on social behavior vary widely across pathogens, hosts and host sex, levels of infection

severity, and types of social relationships. In particular, stable social bonds should be more resilient

to immune challenges or changes in individual condition compared to weak social relationships (22-

26). For example, in vampire bats, sickness behavior reduces individual activity and social grooming

among non-kin, but it does not decrease maternal grooming of offspring (22). Similarly, mandrills

avoid grooming parasite-infected non-kin but continue to groom infected close maternal kin (23). For

animals that form monogamous pair bonds, short-term infections are likely to have a bigger impact

on choosing mates (reviewed in 25) and forming pair bonds than on existing pair bonds (26). It

remains unclear, however, to what extent infections impact affiliative same-sex social bond

formation.

New social bonds form when individuals join a new social community, which is also when

individuals are exposed to new pathogens. If exposure to new pathogens is more likely when

encountering new social partners, and if infectious disease causes sickness behavior that limits

individual activity and the motivation to interact, then acute infection during the earliest stages of

relationship formation might negatively affect rates of social integration. In a group with both familiar

and new social partners, we therefore predict that the effect of infection on social integration should

be stronger in new, unfamiliar dyads (pairs) compared to familiar dyads with well-established

relationships. In new dyads of unfamiliar bats, infections should initially reduce social interaction

rates for infected bats; as the infected individuals recover and sickness behaviors resolve,

interaction rates of infected and asymptomatic bats should then converge over time. In familiar

dyads, we expect this interaction to be weaker, because infections should have a smaller impact on

how interaction rates change over time in more established relationships.

To address this hypothesis, we report opportunistic evidence that an acute bacterial infection

reduced social integration in common vampire bats (*Desmodus rotundus*). In this species, social

integration is observable as strangers transition from no-cost clustering to low-cost allogrooming and

high-cost regurgitated food sharing (27). Allogrooming is the best measure of relationship formation,

because it is a directed investment of time and energy (unlike clustering) that also occurs frequently

enough to track over time through relationship formation (unlike food sharing). During a previous 4-

month study, we tracked changes in allogrooming among 21 female vampire bats before and after a

week of forced proximity in which small groups of 3 bats from 3 distant sites were housed together

for one week (28). Shortly after the 21 bats from different sites were first introduced in captivity, we

observed an unplanned outbreak of a bacterial infection, which we treated with antibiotics (28). In

this paper, we analyzed the social effect of these infections by tracking changes in allogrooming over

time among the 9 infected female bats and the 12 asymptomatic female bats, comparing dyads that

were previously familiar or unfamiliar with each other.

Methods

Study subjects

For an experiment testing the effects of forced proximity on relationship formation (27), we

captured 8-9 common vampire bats from each of three sites in Panama that were 120-350 km apart.

Upon capture, the 26 bats (23 adult females, 2 juvenile males, and one juvenile female) appeared

healthy with no pathological lesions. We housed the bats together in an outdoor flight cage (2.1 × 1.7

× 2.3 m) for approximately 4 months. During this time, five females gave birth to four males and one

female pup (colony included 31 total bats). Although we caught 23 females, two recaptured females

entered the captive colony several weeks after the colony was created, were not included in all

phases of our experiment, and were therefore not included in this analysis. We analyzed the

allogrooming relationships among 252 new and 126 familiar dyads of 21 adult females. We define

'dyads' as unique directed combinations of a groomer and receiver. We use the label 'new' for dyads

that were introduced in captivity and 'familiar' for dyads that were captured from the same site

(hollow tree or cave). Details of the study colony can be found in (28,29).

Behavior sampling

We used three infrared surveillance cameras to sample social interactions among the

individually marked bats (28). Over 114 days, we sampled 682 h of interactions, recording all visible

allogrooming and mouth-licking interactions (≥ 5 seconds in duration) as well as any aggression. For

each interaction, we noted the actor, receiver, and duration. We used allogrooming to assess social

integration because allogrooming was 94% of the 17,592 observed social interactions, while mouth-

licking and aggression were only 6% and 0.3% of interactions, respectively. For each directed edge,

we summed the total seconds of allogrooming from the groomer to the receiver. We also summed

the total sampled seconds that each undirected dyad had the opportunity to interact. For further

details on behavior sampling, see (28).

Bacterial infection and treatment

Two weeks after the 26 wild-caught bats were first introduced to one another in the flight

cage, 10 female bats began to show progressive symptoms of a cutaneous infection (blisters or

open lesions with pus, as well as inflamed tissue). Symptoms progressed for approximately two

weeks without signs of recovery. These infections were unexpected and to our knowledge have not

previously been described in vampire bat colonies. The 10 infected bats varied in the severity of their

symptoms. The most severe symptoms occurred in bats from a hollow tree near La Chorrera,

Panama. The most severely infected bat developed large, pus-filled lesions on its back and had

extreme inflammation around the wrists that led tissue to break apart and expose bone; we

euthanized this bat before it received antibiotic treatment and did not include it in our analysis of

social integration. Four other bats developed severe symptoms with ulcers and inflammation on one

or both wrists, leading to tissue scarring and the inability to sustain flight. One bat from the same

roost experienced only minor inflammation and ulcers on both wrists but healed without permanent

tissue scarring and continued to fly. Three bats from a cave near Lake Bayano, Panama were also

infected. Two bats developed pus-filled abscesses: on the lower abdomen and upper thigh for one

bat, on the lower back for the other bat, and the third bat developed only minor inflammation on 1-2

toes. Finally, one bat from a hollow tree near Tolé, Panama had a large, round lesion on its knee,

but this was evident at the start of the study and did not change over time. This bat displayed no

other signs of infection. With most cases having been observed in bats from either La Chorrera or

Lake Bayano, it seems the bats from Tolé were most resistant to the pathogen. Swab and tissue

samples from skin lesions were cultured to isolate bacterial pathogens, which indicated the presence

of *Bacillus sp.* and *Staphylococcus aureus*. There are several *Bacillus* species (e.g., *Bacillus*

anthracis, Bacillus cereus, and *Bacillus pumilus*) that may infect the skin or colonize wounds, but

these infections are uncommon and often present with classic symptoms (e.g. black eschars and

necrosis, 30–33) that we did not observe in our colony. Alternatively, several species within the *S.*

aureus complex, including *Staphylococcus schweitzeri* and *Staphylococcus argenteus,* among

others (34–36), typically infect the skin and cause symptoms that are consistent with what we

observed. *S. aureus* complex has also been isolated from the feces of Old-World fruit bats (37–40)

and the common vampire bat (41,42).

A Kirby-Bauer antibiogram indicated susceptibility of the isolated bacteria to the antibiotic

Enrofloxacin. Given the urgency and results of the antibiogram, we immediately used Enrofloxacin

antibiotic to treat the infected female bats (described above) over the course of seven days via daily

subcutaneous injections. We did not perform additional testing to identify the bacterial species or

strain. After the antibiotic treatment, five bats recovered without injury (henceforth "treated"), but four

bats suffered lasting injuries that rendered them unable to fly for the remainder of the study

(henceforth "injured"). We refer to the remaining 12 bats in our analyses as "asymptomatic".

Confirming sickness behavior

To confirm that bacterial infections reduced individual activity and social behavior, i.e.,
'sickness behavior', we compared the activity of 9 infected versus 12 asymptomatic adult female
bats during a sample period of 1 week before the antibiotic treatment (pre-treatment week) and
again during 1 week at the end of the study (post-treatment week). An observer scored videos using
focal sampling for the presence or absence of 3 behavioral states: moving, allogrooming (including
mouth-licking), and self-grooming, following definitions in (13). The observer could not be entirely
blind to the condition of bats in the videos because lesions were visible, but the analyzed behavioral
categories were not subjective. For one hour (starting one hour after sunset), an observer noted at
30-second intervals whether a focal female was engaged in any of the behaviors of interest. We
therefore sampled each of the 21 bats for a total of 14 days, resulting in 1680 samples per bat (120
per day).

To test if infection predicted the frequency of each behavioral state, we fit Bayesian
multilevel binomial regressions in R (brms package, 43) and obtained median posterior probability
estimates and 90% highest posterior density intervals (HPDI). The 90% HPDI is a more stable
interval than a 95% interval, and is often used to prevent readers from interpreting HPDIs as 95%
confidence intervals or denoting "statistical significance". For each behavioral state, model predictors
were infection status (asymptomatic vs. infected), treatment period (pre- vs post-treatment) the
interaction between infection status and period, as well as bat identity as a random intercept. The
response variable was the proportion of observations doing a behavior (allogrooming, self-grooming,
and moving) out of the total number of observations of the bat on camera. To evaluate interactions
between infection and treatment period, we then subset the data into pre-treatment and post-
treatment periods and fit separate models to estimate the effect of infection within each period.
Across models, all parameters had above 1000 effective samples and an R-hat equal to 1.00 with
four chains, indicating successful convergence. Since the effects of acute bacterial infection on
vampire bat behavior have not been observed, we used conservative default priors across all

effects. However, to put our effect sizes (odds ratios) in context, we compared them with effect sizes

from past experimental immune challenges, explained below.

Comparison of results to previous experimentally induced sickness behavior

We present our results alongside reference effect sizes from two past experiments on the

behavioral responses of female vampire bats to a simulated bacterial infection (13, 22). In both

experiments, behaviors of vampire bats were observed after being injected with either

lipopolysaccharide (LPS) or saline. In the first study (13), vampire bats were observed in groups of

2-4 bats within small cages. LPS injection reduced the odds of allogrooming by a factor of 3.73

(binomial mixed effect model, Odds Ratio, 95% confidence interval (CI) = 2.25–6.50) and reduced

the odds of self-grooming by a factor of 16 (Odds Ratio, 95% CI = 12–27). In the second study (22),

vampire bats were fasted to induce food sharing and observed in a larger group (36 bats) in a flight

cage. Under these conditions, LPS injection reduced the odds of allogrooming by a factor of 1.89

(Odds Ratio, 95% CI = 1.84–1.93). This smaller effect on allogrooming could be due to differences in

group size, cage size, the hunger state of the injected bats, or some combination of those factors.

We present reference effects from both studies because the present study shares some features of

both; the bats were unfasted like the first study but held in a large group in the same flight cage as

the second study.

Effects of infection on symmetry of allogrooming given and received

We expected that any sickness effects on allogrooming given would cause at least some

effects on allogrooming received, because allogrooming is highly reciprocal and most allogrooming

occurs bidirectionally (28,44,45). To assess whether infection reduced or removed the symmetry of

directed allogrooming, we compared the proportion of hourly allogrooming rates that were

bidirectional for dyads with at least one infected bat versus dyads with two asymptomatic bats.

Effects of infection and familiarity on dyadic allogrooming rates over time

We assessed how allogrooming changed over time among familiar and new dyads of bats

that were either asymptomatic, infected, or injured. To do this, we fit a Bayesian social relations

model in R (STRAND package, 46). We used an aggregated binomial outcome to model the

proportion of total seconds of observation time with or without allogrooming for each directed

groomer-receiver dyad. Four dyadic covariates included whether the dyad had been forced into

proximity (true/false), as well as the main effects and interaction between familiarity (true/false) and

infection severity (a sum of both individuals' infection severity scores: 0 = asymptomatic, 1 =

infected, 2 = injured). This measure of infection severity makes assumes that these states have

linearly increasing and additive effects, but this simplifying assumption is consistent with

comparisons of allogrooming for infected bats versus all others (Fig. 1) and for injured bats versus all

others (Fig. S1). To explore how these effects change through time, we binned the data by week (15

social networks across 16 weeks, excluding one week when bats were not sampled altogether), and

fit the same model described above to each time bin and extracted posterior probabilities with 90%

HPDIs. Model convergence diagnostics can be found in the online data repository (47).

We predicted a three-way interaction between familiarity, time, and infection severity.

Specifically, we expected that (1) infections would have a greater negative impact on allogrooming

rates in new dyads relative to familiar dyads, and (2) the effect of infection on the slope of dyadic

allogrooming rates over time would be greater for new dyads than for familiar dyads. Over time, both

infection and familiarity effects should converge towards zero because by the end of the study new

dyads are no longer new and infected bats are no longer infected.

Ethics

This work was approved by the Smithsonian Tropical Research Institute Animal Care and

Use Committee (no. 2015-0501-2022) and the Panamanian Ministry of the Environment (no. SEX/A-

67-2019).

Data accessibility

217   Data and R code for all analyses are available on Github (47).

219  **Results**

220  *Confirming sickness behavior*

221   Consistent with sickness behavior and recovery, infected bats spent less time moving and

222  allogrooming than did asymptomatic bats before antibiotic treatment, and these differences faded to

223  zero over time (Fig. 1A, Fig. 1C). Before antibiotic treatment, infected bats spent 2.5% of their time

224  allogrooming, 90% credibility intervals = [<0.001, 5.4%], whereas asymptomatic bats spent 7.3%

225  [3.9, 16.5%] of their time allogrooming. Infected bats spent 13.5% [5.1, 3.3%] of their time moving,

226  compared to 29.7% [22.5, 35.1%] in asymptomatic bats. After recovery, the previously infected bats

227  no longer showed differences from the asymptomatic bats in allogrooming or movement (Fig. 1A,

228  Fig. 1C).

229   In contrast with effects on allogrooming, infections did not clearly reduce self-grooming (Fig.

230  1B). Instead, after recovering (post-treatment period) there was a trend for self-grooming rates

231  among previously infected bats to be greater than self-grooming rates between asymptomatic bats.

232  This pattern was driven by higher self-grooming rates in the injured bats, which did not fully recover

233  (Fig. S1).

Figure 1. Infections reduced allogrooming and moving but not self-grooming. Figure panels show effects of infections on allogrooming (A), self-grooming (B), and moving (C). Left-hand panels show median posterior probability estimates (points) with 90% highest posterior density intervals (HPDI). Vertical dashed lines show previous effect sizes estimated from two LPS-induced immune challenge experiments (purple, 13; orange, 22). Right-hand panels show proportions of time for each behavior (jittered data) for infected bats (red triangles) and asymptomatic bats (black squares), during the outbreak (pre-treatment) and after bats recovered (post-treatment). Points with error bars are predicted median posterior proportions ± 90% HPDIs.

Effects of infection on symmetry of allogrooming given and received

246   We found no evidence that infection reduced or removed allogrooming symmetry. Dyadic

247 allogrooming was bidirectional within the hour in 84% of cases with at least one infected bat and in

248 81% of cases with two asymptomatic bats. The observation that allogrooming with an infected bat

249 was not more likely to be unidirectional than allogrooming among asymptomatic bats confirms our

250 assumption that any effect of infection on allogrooming given would also impact allogrooming

251 received.

253 *Effects of infection and familiarity on dyadic allogrooming rates* over time

254   The interaction between familiarity and infection severity changed over time (Fig. 2). Prior to

255 antibiotic treatment (weeks 1 – 6, Fig. 2), posterior probability estimates of the interaction between

256 familiarity and infection severity were largely negative (i.e. the effect of infection differed between

257 new and familiar dyads). This difference then faded during the post-treatment period (weeks 8 – 16;

258 Fig 2). In familiar dyads, infection did not clearly impact dyadic allogrooming rates over time,

259 whereas in new dyads, there was a clear negative impact of infection severity on allogrooming rates

260 that diminished with time (Fig 3).

Figure 2. The effect of infection on allogrooming initially differs between new and familiar

dyads, but this difference fades with time as bats recover and become familiar. Y-axis shows

posterior probability estimates for the interaction between familiarity and infection severity as

predictors of allogrooming rate, for each week (x-axis). Points with error bars show median posterior

probabilities with 90% highest posterior density intervals.

Figure 3. Observational evidence that infections disproportionately reduce allogrooming in

new dyads (developing social relationships) relative to familiar dyads (existing social

relationships). Y-axis shows the log-transformed (log(x + 1)) mean daily allogrooming rates among

all familiar dyads (left) and new dyads (right) across weeks. Yellow points and solid slopes indicate

dyads with an infection severity of 0-1 (e.g. two asymptomatic bats, or one infected bat), whereas

dark red triangles and dashed slopes indicate dyads with an infection severity of 2-4 (e.g. two

infected bats, two injured bats). Shaded regions indicate 95% CI of slope.

Discussion

We present opportunistic evidence that acute symptoms of bacterial infection can limit the

development of allogrooming relationships in vampire bats. Relative to asymptomatic bats, the

initially infected bats spent less time moving and allogrooming. This sickness behavior, which lasted

approximately 2-3 weeks, led to changes in social integration, whereby infections had a greater

impact on the formation of new allogrooming relationships than on allogrooming in existing

relationships. Allogrooming rates were therefore influenced by a three-way interaction between

familiarity, infection severity, and time. Infections had a more negative impact on allogrooming rates

in new dyads relative to familiar dyads, such that the slope of allogrooming over time was greatest

for new dyads with infected and/or injured bats. Familiar dyads and new dyads without infected bats

were less impacted by infections. As predicted, the effects of infection and familiarity effects, and

their interaction, converged towards zero as new dyads became familiar and infected individuals

recovered over three months. This convergence suggests that sickness effects were not entirely

driven by injuries that reduced movement.

Because we did not randomly and experimentally infect the bats, a possible alternative

explanation for our findings is the reverse causal path to what we suggested above: perhaps bats

with worse social integration (or worse social competence) had weaker immune systems and

therefore suffered worse infections. However, this hypothesis cannot as easily explain why the gap

in social integration faded over time after antibiotic treatment. If the infected bats were already less

social or less healthy for reasons besides the infection, then we would expect their social integration

to be worse throughout the study, not just during the outbreak. Also, if social factors (like dominance

or social competence) caused infection severity, then infection rates should be similar within each

wild population. Instead, infection rate and severity were associated with wild population, suggesting

that infections were a cause rather than a consequence of how bats formed new relationships.

The sickness behavior we observed was similar in some respects to the responses of

vampire bats to simulated infections from injections of lipopolysaccharide, LPS (13,22). The use of

immune challenges like LPS allows for controlled experiments on the connection between the

inflammatory and the behavioral response to infection (9,11–13,16,22,26,33,48–50), but naturally

infecting pathogens can have different, pathogen-specific effects. For instance, LPS-injected

vampire bats significantly reduced their self-grooming, but the naturally infected bats in our study did

not, possibly because the bacterial pathogens caused skin lesions that encouraged self-grooming

even when overall activity was reduced. Another pathogen that infects the skin of bats, the fungus

Pseudogymnoascus destructans (the cause of 'white-nose syndrome') also causes increased self-

grooming (51).

These observations are opportunistic and sparse (only 9 infected bats), but we believe our

results are valuable for two main reasons. First, effect size estimates for sickness behaviors of

vertebrates in response to naturally-infecting pathogens are uncommon compared to experiments

that use immune challenges (but see 51–55). Second, while several studies have quantified the

effect of immune challenges (infection or immunogenic antigens) on host social behavior (15,17–

19,51–59), the effect of short-term infections on the formation and development of long-term social

bonds development remains largely unexplored. Our results can inform future experiments on the

effect of infections on social integration and relationship formation.

Our findings also raise several questions. When might animals suppress their infection-

induced behavioral symptoms to continue social integration (7,48,49,60–62)? At what stages of

relationship formation do sickness behaviors have the strongest effect on social interactions? What

is the relationship between the duration or intensity of sickness behavior and the effects on social

integration? To address these questions experimentally, researchers could use immune challenges

at different doses (49, 63–64), and/or administer them at varying stages of relationship formation.

In conclusion, we found opportunistic observational evidence for reduced social integration in

infected vampire bats after an outbreak of a naturally infecting pathogen. Just as sickness behavior

and other cues of infection can reduce social activity and mate attraction (reviewed in 25), there may

be similar effects on formation of same-sex social bonds. If there are critical periods of relationship

formation, then infections during this time might have lingering social consequences.

Acknowledgements

The authors thank Rachel Page for lab space and resources, the staff at the Smithsonian Tropical Research Institute, and the following individuals who assisted with vampire bat care and/or data collection: D. Aparicio, B. Brown, G. Cohen, L. Dück, D. Girbino, E. Kline, C. Marroquin, and S. Ripperger.

Funding

I.R. was supported by a short-term fellowship from the Smithsonian Tropical Research Institute, a student research grant from the Animal Behaviour Society, a Graduate Enrichment Fellowship from The Ohio State University, a Tinker Foundation Field Research Grant, a Grant-in-aid of Research from the American Society of Mammalogists, and a Smithsonian Institution Predoctoral Fellowship. S.S. was supported by the Ohio State University President's Postdoctoral Scholars Program. This publication is based upon work supported by the National Science Foundation under grant no. IOS-2015928 to G. C.

Author contributions

Conceptualization: IR, SS, GC

Data Curation: IR, GC

Formal Analysis: IR, SS, GC

Funding Acquisition: IR, GC

Investigation: IR, MA, GC

Methodology: IR, GC

Project Administration: IR, GC

Resources: IR, GC

Software: IR, GC

Supervision: GC

Validation: IR, SS, GC

Visualization: IR, SS, GC

Writing – Original Draft Preparation: IR, SS, GC

Writing – Review & Editing: IR, MA, SS, GC

References

1. Pavez-Fox MA, Negron-Del Valle JE, Thompson IJ, Walker CS, Bauman SE, Gonzalez O, et al.

Sociality predicts individual variation in the immunity of free-ranging rhesus macaques. Physiol

Behav. 2021;241:113560.

2. Snyder-Mackler N, Burger JR, Gaydosh L, Belsky DW, Noppert GA, Campos FA, et al. Social

determinants of health and survival in humans and other animals. Science.

2020;368(6493):eaax9553.

3. Snyder-Mackler N, Sanz J, Kohn JN, Brinkworth JF, Morrow S, Shaver AO, et al. Social status

alters immune regulation and response to infection in macaques. Science.

2016;354(6315):1041–5.

4. Tung J, Barreiro LB, Johnson ZP, Hansen KD, Michopoulos V, Toufexis D, et al. Social

environment is associated with gene regulatory variation in the rhesus macaque immune

system. Proc Natl Acad Sci. 2012;109(17):6490–5.

5. Almberg ES, Cross PC, Dobson AP, Smith DW, Metz MC, Stahler DR, et al. Social living

mitigates the costs of a chronic illness in a cooperative carnivore. Ecol Lett. 2015;18(7):660–7.

6. Loehle C. Social barriers to pathogen transmission in wild animal populations. Ecology.

1995;76(2):326–35.

7. Jog MG, Sackett ME, Kisty SD, Hansen JA, Stephenson JF. The behaviour of infected guppies

depends on social context, parasite tolerance and host sex. Anim Behav. 2022;187:97-104.

doi:10.1016/j.anbehav.2022.03.001

8. Hart BL. Biological basis of the behavior of sick animals. Neurosci Biobehav Rev.

1988;12(2):123–37.

9. Lopes PC, French SS, Woodhams DC, Binning SA. Sickness behaviors across vertebrate taxa:
proximate and ultimate mechanisms. J Exp Biol. 2021;224(9):jeb225847.

10. Stockmaier S, Stroeymeyt N, Shattuck EC, Hawley DM, Ancel Meyers L, Bolnick DI. Infectious
diseases and social distancing in nature. Science. 2021;371(6533):eabc8881.

11. Fishkin RJ, Winslow JT. Endotoxin-induced reduction of social investigation by mice: interaction
with amphetamine and anti-inflammatory drugs. Psychopharmacol Berl. 1997;132(4):335–41.

12. Moreno KR, Weinberg M, Harten L, Salinas Ramos VB, Herrera M. LG, Czirják GÁ, et al. Sick
bats stay home alone: fruit bats practice social distancing when faced with an immunological
challenge. Ann N Y Acad Sci. 2021;1505(1):178–90.

13. Stockmaier S, Bolnick DI, Page RA, Carter GG. An immune challenge reduces social grooming
in vampire bats. Anim Behav. 2018;140:141–9.

14. Lopes PC, Block P, Konig B. Infection-induced behavioural changes reduce connectivity and the
potential for disease spread in wild mice contact networks. Sci Rep. 2016;6:31790.

15. Fairbanks BM, Hawley DM, Alexander KA. The Impact of Health Status on Dispersal Behavior in
Banded Mongooses (Mungos mungo). EcoHealth. 2014;11(2):258–62.

16. Lopes PC, Woodhams DC, Binning SA. Infection avoidance behaviors across vertebrate taxa:
patterns, processes and future directions. In: Animal Behavior and Parasitism. Oxford University
Press; 2022.

17. Behringer DC, Butler MJ, Shields JD. Ecology: avoidance of disease by social lobsters. Nature.
2006;441(7092):421.

18. Kiesecker JM, Skelly DK, Beard KH, Preisser E. Behavioral reduction of infection risk. Proc Natl
Acad Sci U A. 1999;96(16):9165–8.

19. Poirotte C, Massol F, Herbert A, Willaume E, Bomo PM, Kappeler PM, et al. Mandrills use
olfaction to socially avoid parasitized conspecifics. Sci Adv. 2017;3(4).

20. Boillat M, Challet L, Rossier D, Kan C, Carleton A, Rodriguez I. The Vomeronasal System
Mediates Sick Conspecific Avoidance. Curr Biol. 2015;25(2):251–5.

21. Hawley DM, Gibson AK, Townsend AK, Craft ME, Stephenson JF. Bidirectional interactions

between host social behaviour and parasites arise through ecological and evolutionary

processes. Parasitology. 2021;148:274-288. doi:10.1017/S0031182020002048

22. Stockmaier S, Bolnick DI, Page RA, Carter GG. Sickness effects on social interactions depend

on the type of behaviour and relationship. J Anim Ecol. 2020; 89:1387-1394.

23. Poirotte C, Charpentier MJE. Unconditional care from close maternal kin in the face of parasites.

Biol Lett. 2020;16(2):20190869.

24. Seyfarth RM, Cheney DL. The evolutionary origins of friendship. Ann Rev Psych. 2012;

63(1):153-177.

25. Beltran-Bech S, Richard FJ. Impact of infection on mate choice. Anim Behav. 2014;90:159–70.

26. Love AC, Anthony AC, Nash A, Campos-Melara A, Kodali J, DuRant SE. Simulated infection

alters the behavior of pair bonded songbirds and their asymptomatic neighbors. Behav Ecol.

2023;34(2):251–60.

27. Carter GG, Farine DR, Crisp RJ, Vrtilek JK, Ripperger SP, Page RA. Development of New Food-

Sharing Relationships in Vampire Bats. Curr Biol. 2020;30(7):1275-1279.e3.

28. Razik I, Brown BKG, Carter GG. Forced proximity promotes the formation of enduring

cooperative relationships in vampire bats. Biol Lett. 2022;18(4):20220056.

29. Razik I, Brown BKG, Page RA, Carter GG. Non-kin adoption in the common vampire bat. R Soc

Open Sci. 202;8(2):201927.

30. Cromartie WJ, Bloom WL, Watson DW. Studies on Infection with *Bacillus anthracis*: I.

Histopathological Study of Skin Lesions Produced by *B. anthracis* in Susceptible and Resistant

Animal Species. J. Infect. Dis. 1947;80(1):1–13.

31. Esmkhani M, Shams S. Cutaneous infection due to *Bacillus cereus*: a case report. BMC Infect.

Dis. 2022;22(1);393.

32. Tena D, Martínez-Torres JA, Pérez-Pomata MT, Sáez-Nieto JA, Rubio V, Bisquert J. Cutaneous

Infection Due to *Bacillus pumilus*: Report of 3 Cases. Clin. Infect. Dis. 2007;44(4):e40–e42.

33. Veysseyre F, Fourcade C, Lavigne JP, Sotto A. *Bacillus cereus* infection: 57 case patients and a

literature review. Med Mal Infect 2015;45(11):436-440.

34. Becker K, Schaumburg F, Kearns A, Larsen AR, Lindsay JA, Skov RL, Westh H. Implications of

identifying the recently defined members of the *Staphylococcus aureus* complex *S. argenteus*

and *S. schweitzeri*: a position paper of members of the ESCMID Study Group for Staphylococci

and Staphylococcal Diseases (ESGS). Clin. Microbiol. Infect. 2019;25(9):1064–1070.

35. Chew KL, Octavia S, Lai D, Lin RTP, Teo JWP. *Staphylococcus singaporensis* sp. nov., a new

member of the *Staphylococcus aureus* complex, isolated from human clinical specimens. Int. J.

Syst. Evol. Microbiol. 2021;71(10):005067.

36. Tong SYC, Schaumberg F, Ellington MJ, Corander J, Pichon B, Leendertz F, Bentley SD,

Parkhill J, Holt DC, Peters G, Giffard PM. Novel staphylococcal species that form part of a

Staphylococcus aureus-related complex: the non-pigmented *Staphylococcus argenteus* sp. nov.

and the non-human primate-associated *Staphylococcus schweitzeri* sp. nov. Int. J. Syst. Evol.

Microbiol. 2015;65:15–22.

37. Fountain K, Barbon A, Gibbon MJ, Lloyd DH, Loeffler A, Feil EJ. *Staphylococcus aureus*

lineages associated with a free-ranging population of the fruit bat *Pteropus livingstonii* retained

over 25 years in captivity. Sci. Rep. 2022;12(1):13457.

38. Held J, Gmeiner M, Mordmüller B, Matsiégui P, Schaer J, Eckerle I, Weber N, Matuschewski K,

Bletz S, Schaumburg F. Bats are rare reservoirs of *Staphylococcus aureus* complex in Gabon.

Infect. Genet. Evol. 2017;47:118–120.

39. Monecke S, Schaumburg F, Shittu AO, Scharz S, Mühldorfer K, Brandt C, Braun SD, Collatz M,

Diezel C, Gawlik D, Hanke D, Hotzel H, Müller E, Reinicke M, Feßler AT, Ehricht R. Description

of Staphylococcal Strains from Straw-Coloured Fruit Bat (*Eidolon helvum*) and Diamond Firetail

(*Stagonopleura guttata*) and a Review of their Phylogenetic Relationships to Other

Staphylococci. Front. Cell. Infect. Microbiol. 2022;12:878137.

40. Olatimehin A, Shittu AO, Onwugamba FC, Mellmann A, Becker K, Schaumburg F.

Staphylococcus aureus Complex in the Straw-Colored Fruit Bat (*Eidolon helvum*) in Nigeria.

Front. Microbiol. 2018;9:162.

41. Chaverri G. Aerobic bacterial flora from the digestive tract of the common vampire bat,

Desmodus rotundus (Chiroptera: Phyllostomidae). Rev. Biol. Trop. 2006;54:717–724.

42. Monero M, Lopes CAM, Seabra E, Pavan C, Correa A. Aspectos bacteriologicos da flora

intestinal de *Desmodus rotundus*. Arq. Inst. Biol. São Paulo 1975;42:229.

43. Burkner P. brms: An R Package for Bayesian Multilevel Models Using Stan. J. Stat. Software.

2017;80:1–28. doi:10.18637/jss.v080.i01.

44. Wilkinson GS. Social grooming in the common vampire bat, *Desmodus rotundus*. Anim Behav.

1986;34(6):1880–9.

45. Carter GG, Wilkinson GS. Food sharing in vampire bats: reciprocal help predicts donations more

than relatedness or harassment. Proc Biol Sci. 2013;280(1753):20122573.

46. Ross CT, McElreath R, Redhead D. Modelling human and non-human animal network data in R

using STRAND. bioRxiv 2022; 2022.05.13.491798.

47. Razik I, Carter G, Stockmaier S, Abou-Elias M. Data and R code for Opportunistic evidence of

the impact of bacterial infections on social integration in vampire bats. 2023; Available from:

https://github.com/imranrazik/infection_vampires

48. Lopes PC, Adelman J, Wingfield JC, Bentley GE. Social context modulates sickness behavior.

Behav Ecol Sociobiol. 2012;66(10):1421–8.

49. Willette AA, Lubach GR, Coe CL. Environmental context differentially affects behavioral,

leukocyte, cortisol, and interleukin-6 responses to low doses of endotoxin in the rhesus monkey.

Brain Behav Immun. 2007;21(6):807–15.

50. Alciatore G, Ugelvig LV, Frank E, Bidaux J, Gal A, Schmitt T, et al. Immune challenges increase

network centrality in a queenless ant. Proc R Soc B. 2021;288(1958):20211456.

51. Brownlee-Bouboulis SA, Reeder DM. White-nose syndrome-affected little brown myotis (Myotis

lucifugus) increase grooming and other active behaviors during arousals from hibernation. J

Wildl Dis. 2013;49(4):850–9.

52. Cárdenas-Canales EM, Stockmaier S, Cronin E, Rocke TE, Osorio JE, Carter GG. Social effects

of rabies infection in male vampire bats (*Desmodus rotundus*). Biol Lett. 2022;18(9):20220298.

53. Fairbanks BM, Hawley DM, Alexander KA. No evidence for avoidance of visibly diseased

conspecifics in the highly social banded mongoose (*Mungos mungo*). Behav Ecol Sociobiol.

2015;69(3):371–81.

54. Kollias GV, Sydenstricker KV, Kollias HW, Ley DH, Hosseini PR, Connolly V, Dhondt AA.

Experimental infection of house finches with Mycoplasma gallisepticum. J Wildl Dis.

2004;40(1):79-86.

55. Bouwman KM, Hawley DM. Sickness behaviour acting as an evolutionary trap? Male house

finches preferentially feed near diseased conspecifics. Biol Lett. 2010;6(4):462–5.

56. Stephenson JF. Parasite-induced plasticity in host social behaviour depends on sex and

susceptibility. Biol Lett. 2019;15(11):20190557.

57. Stroeymeyt N, Grasse AV, Crespi A, Mersch DP, Cremer S, Keller L. Social network plasticity

decreases disease transmission in a eusocial insect. Science. 2018;362(6417):941.

58. Cremer S. Social immunity in insects. Curr Biol. 2019;29(11):R458–63.

59. Bos N, Lefevre T, Jensen AB, d'Ettorre P. Sick ants become unsociable. J Evol Biol.

2012;25(2):342–51.

60. Aubert A, Goodall G, Dantzer R, Gheusi G. Differential Effects of Lipopolysaccharide on Pup

Retrieving and Nest Building in Lactating Mice. Brain Behav Immun. 1997;11(2):107–18.

61. Lopes PC. When is it socially acceptable to feel sick? Proc Biol Sci. 2014;281(1788):20140218.

62. Weil ZM, Bowers SL, Dow ER, Nelson RJ. Maternal aggression persists following

lipopolysaccharide-induced activation of the immune system. Physiol Behav. 2006;87(4):694–9.

512 63. Grigoleit J, Kullmann JS, Wolf OT, Hammes F, Wegner A, Jablonowski S, Engler H, Gizewski E,

513   Oberbeck R, Schedlowski M. Dose–dependent effects of endotoxin on neurobehavioral

514   functions in humans. PLoS One. 2011;6(12):e28330.

515 64. Dantzer R, Bluthe R, Gheusi G, Cremona S, Laye S, Parnet P, Kelley KW. Molecular basis of

516   sickness behavior. Ann N Y Acad Sci. 2006;856(1):132–138.

517