Opportunistic evidence for impacts of *Staphylococcus* infection on social integration in vampire bats

Imran Razik[1,2], Sebastian Stockmaier[3], Michael Abou-Elias[1], Gerald Carter[1,2]

Affiliations

[1] Department of Evolution, Ecology and Organismal Biology, The Ohio State University, Columbus OH, USA
[2] Smithsonian Tropical Research Institute, Balboa Ancon, Panama, Republic of Panama
[3] Department of Ecology and Evolutionary Biology, University of Tennessee, Knoxville, TN, USA

Corresponding author: Imran Razik (razik.2@buckeyemail.osu.edu)

Abstract

Social integration affects an individual's susceptibility to infection, and conversely, infectious disease can reduce their social activity. It remains unclear, however, to what extent short-term infections might affect the formation of enduring social bonds. During a previous study on the formation of new allogrooming relationships among 21 common vampire bats (*Desmodus rotundus*), we used antibiotics to treat unplanned *Staphylococcus* infections in 9 bats, including 5 bats that recovered and 4 bats that suffered lasting injuries that prevented sustained flight. We compared the behaviors of the *Staphylococcus*-infected bats to healthy controls, and analyzed changes in allogrooming over time among infected and healthy bats that were familiar or unfamiliar. Relative to healthy bats, the 9 *Staphylococcus*-infected bats spent less time moving and allogrooming, and the 4 bats that suffered injuries received less grooming from new partners. These differences faded during 3 months of recovery. This pattern is consistent with the hypothesis that acute infections impact the development of social relationships. Our findings raise questions about how short-term infections might impact long-term social integration or partner choice.

Introduction

Social integration, the quantity and quality of beneficial social relationships, can affect an individual's susceptibility to infection (e.g.,1–4) or help individuals tolerate the adverse effects of parasites (5,6). Conversely, infectious disease can inhibit social behaviors of infected individuals (7–9) by reducing their motivation to interact with others (e.g.,10–12), reducing their movement (e.g., mice, (13) and mongoose (14), or causing others to avoid them (15–19). The effects of an infection on social behavior vary with many factors including the pathogen, the host, the infection severity, and the type of social relationship. For example, sickness behavior in a juvenile vampire bat can increase grooming received from its mother and reduce grooming received from other groupmates (20). Similarly, mandrills avoid grooming parasite-infected non-kin but continue to groom their infected close maternal kin (21). In general, one expects that infections might reduce social interactions disproportionately more within new or weak social relationships compared to strong social bonds (22). For example, infections are likely to have a bigger impact on mate choice (reviewed in 23) and on the process of pair bonding than on an existing pair bond (see example of simulated infections in pair-bonded animals, 24). It remains unclear, however, to what extent infections impact same-sex social bond formation.

We expect new social bonds to form when new adults join a social network, which is also when individuals are exposed to new pathogens. Social bonds are relatively stable relationships, so events that influence relationship formation might have persistent long-term effects (25). If so, the impacts of infections on relationship formation may be greater and have disproportionately greater long-term consequences than the impacts on existing relationships. This begs the question: Can short-term sickness behaviors shape the long-term development of social bonds?

To help address this question, we report opportunistic evidence that bacterial infections appeared to reduce social bond formation in common vampire bats (*Desmodus rotundus*). In this

species, social bond formation is observable as strangers transition from clustering to social
grooming and regurgitated food sharing (26). During a previous 4-month study, we tracked changes
in allogrooming among 21 female vampire bats before and after a week of forced proximity in which
3 bats from 3 distant sites were housed together for one week (25). When the bats from different
sites were first housed together, we observed an unplanned outbreak of *Staphylococcus* infections,
which we treated with antibiotics (25). Here, we analyzed changes in allogrooming over time among
the 9 *Staphylococcus*-infected bats and the 12 healthy bats that were familiar or unfamiliar with each
other.
62
63 **Methods**
64
Behavior sampling
We captured 26 bats (8-9 bats from each of three sites in Panama that were 120-350 km
apart) and housed them together in an outdoor flight cage. Four males and one female were born in
captivity. Of these 31 bats, we used 21 females in a study testing the effects of forced proximity on
relationship formation (25). We use the label 'unfamiliar' for pairs of bats that were introduced in
captivity and 'familiar' for pairs that were captured from the same site (hollow tree or cave). We used
three infrared surveillance cameras to sample social interactions among the individually marked
bats. Over 114 days, we sampled 682 h of interactions, mostly bouts of allogrooming (>5 s in
duration) and noted the actor and receiver. To estimate dyadic allogrooming rates, we averaged the
seconds of allogrooming per hour across all hours in which the actor and receiver could interact.
Details of the study colony can be found in (25,27).
76
Staph infection
Soon after bats were housed together, 10 bats developed progressive symptoms of an
infection (open, oozing lesions or inflamed tissue), which also infected their joints. Swab and tissue
samples were sent to a veterinary doctor for bacterial cultures, which confirmed bacterial infection of
Staphylococcus aureus. A Kirby-Bauer antibiotic test indicated strain susceptibility to the antibiotic
Enrofloxacin, which we used to treat 9 infected bats. One bat died prior to receiving antibiotic
treatment. Of the remaining 9 infected bats, five bats recovered without injury ("treated" bats), but
four bats suffered lasting injuries that rendered them unable to fly for the remainder of the study
("injured" bats). We refer to the other 21 bats in the colony as "healthy".
Compared to the treated bats, injured bats were more impacted by the infection prior to the
antibiotic treatment. The injured bats suffered from severe swelling with lesions on one or both wrists
leading to tissue scarring and inability to sustain flight because of their infection. All injured bats were
from a roost near La Chorrera, Panama. One treated bat, also from the La Chorrera roost,
experienced minor swelling and injuries on both wrists, but healed without permanent tissue scarring
and could sustain flight. Two treated bats from a different roost (a cave near Lake Bayano, Panama)
had pus-filled lesions: on the lower abdomen and upper thigh for one bat, and on the lower back for
the other bat. A third bat from Lake Bayano had minor swelling on 1-2 toes which disappeared after
treatment with antibiotic. Finally, one treated bat from a hollow tree near Tolé, Panama had a large
swollen knee, which was evident at the start of the study and did not change after antibiotic
treatment. This bat displayed no other signs of infection and in retrospect may not have been
infected with *Staphylococcus*.
To confirm that *Staphylococcus* infections caused sickness behavior, we compared
behaviors of the 9 infected adult female bats with a control group of 9 healthy adult female bats
during a sample period of 1 week before the antibiotic treatment (pre-treatment week) and again
during another 1 week about 11 weeks later near the end of study (post-treatment week). For each
of the 18 bats, an observer scored the presence or absence of 4 behavioral states: moving, resting
(including sleeping), allogrooming (including mouth-licking), and self-grooming, following definitions
in (12). These behaviors were scored every 30 seconds for 1 hour (starting one hour after sunset)
per day. Overall, we sampled six behaviors across all 18 bats 120 times on each of 14 days. To test
if infection predicted the probability of these six behavioral states, we calculated the mean proportion

of sampled times where each behavior occurred and estimated 95% confidence intervals (CIs) using
bootstrapping with 5000 iterations (percentile method, boot package in R, 28).
109
Effects of infection on dyadic grooming rates
In the previous study (25), a subset of 21 new pairs were forced into proximity together to
induce social bonding. In this study, we excluded these pairs from our analysis to remove any effect
of experimental forced proximity on relationship formation. However, including these pairs in the
analysis does not change our conclusions about infection as a predictor of poor social integration.
We considered *grooming received* as the best response measure of social integration, but we also
tested how infection influenced grooming rates over time when the actor was infected (*grooming*
given) and when either the actor or receiver was infected. To assess if dyadic rates of grooming for
the 9 infected and 12 healthy bats changed over time, we fit three binomial mixed effect models
(lme4 R package, 29) to test which factors predicted the presence or absence of allogrooming in
sampled hours. In all models, actor and receiver identity were random intercepts. In the first model,
fixed effects were prior familiarity of pair (true/false), infection status of either actor or receiver
(true/false), time, and all interactions between those three fixed effects. To scale the β coefficients,
we used 10 days as a unit of time. The second and third model were similar except that we defined
'infection status' based on the grooming *actor* (model 2) or grooming *receiver* (model 3).
In all models, we tested for a three-way interaction, and re-fit the model separately for
familiar and unfamiliar pairs. In addition to Wald Test p-values, we report p-values from the likelihood
ratio test (LRT p) for our three main predicted effects: (i) a three-way interaction between familiarity,
infection status, and time, (ii) an interaction between infection status and time within unfamiliar dyads
(infections have a large impact on how grooming changes over time), and (iii) a weaker interaction or
no interaction between infection status and time within familiar dyads (infections have a smaller
impact on how grooming changes over time).
Finally, we compared the post-recovery increases in grooming rates for bats that were
healthy, treated, or injured. To do this, we first fit a binomial model for each bat to estimate its
change in log odds of grooming per hour (change in grooming received). Then we used
bootstrapping to estimate 95% CIs around the mean change in grooming received for healthy,
treated, and injured bats. Due to our small sample sizes, caution must be taken when interpreting
these CIs and p-values.
138
139 *Ethics*
140   This work was approved by the Smithsonian Tropical Research Institute Animal Care and
141 Use Committee (no. 2015-0501-2022) and the Panamanian Ministry of the Environment (no. SEX/A-
142 67-2019).
143
144 *Data accessibility*
145   Data and R code are available on Figshare (30).
146
147 **Results**
148
Confirming sickness behavior
Consistent with sickness behavior, we found that the 9 *Staphylococcus*-infected bats spent
less time moving and allogrooming, but these differences faded after 3 months of recovery (Figure
1). Before antibiotic treatment, infected bats spent 1.3% (95% CI = [0.6—2.1%]) of their time
allogrooming versus 3.9% [2.3—5.6 %] in healthy bats, and infected bats spent 7.4% [4.8—9.9%] of
their time moving around versus 17% [11—22%] in healthy bats. After recovery, however, the
previously infected bats no longer showed clear differences from the healthy bats in resting or self-
grooming (Figure 1). Further analysis suggested that the pre-treatment differences from the healthy
bats in allogrooming and moving were greater in the 4 injured bats than the 5 treated bats (Figure
S1).
159

Effects of infection on dyadic grooming rates
Overall, we found evidence that infected bats engaged in less allogrooming compared to
healthy bats, that this difference decreased with time, and that this difference was even larger for
new relationships and for bats that were injured. We found no evidence that infection decreased
allogrooming symmetry: allogrooming is often mutual and highly symmetrical within pairs (26,31,32),
and pairwise allogrooming was bidirectional within the hour in 84% of cases with an infected bat and
in 81% of cases with two healthy bats. This confirms our expectation that any sickness effects on
grooming given and grooming received are causally linked.
For pairs of bats with either an infected groomer or receiver, we found a three-way
interaction between familiarity, infection status, and time (β = 0.041, SE = 0.014, LRT p = 0.003). In
familiar pairs, infection did not predict how allogrooming changed over time (β = 0.009, SE = 0.011,
LRT p = 0.412), and familiar allogrooming did not clearly change with time (β = -0.005, SE = 0.009, Z
= -0.522, p = 0.601) or differ by infection (β = 0.086, SE = 0.133, Z = 0.645, p = 0.519). In new
(unfamiliar) pairs, however, allogrooming did increase over time (β = 0.019, SE = 0.007, Z = 2.731, p
= 0.006) and infection predicted both less grooming (β = -1.106, SE = 0.086, Z = -12.855, p < 0.001)
and a larger increase in grooming with time (β = 0.059, SE = 0.009, LRT p < 0.001, Table S2).
The pattern was similar when testing the allogrooming effects of an infected actor. We found
a trend suggesting the same three-way interaction (β = 0.027, SE = 0.014, LRT p = 0.051). In
familiar pairs, infection in the actor did not predict how much grooming changed over time (β = 0.02,
SE = 0.011, LRT p = 0.068) but in new pairs, infection in the actor did influence how grooming
changed with time (β = 0.043, SE = 0.009, LRT p < 0.001, Table S1).
When testing the allogrooming effects of an infected receiver, we did not detect a three-way
interaction (β = 0.009, SE = 0.014, LRT p = 0.498). In both familiar and new pairs, infected receivers
had initially lower levels of allogrooming received leading to greater increases in grooming received
over time compared to healthy receivers (familiar: β = 0.028, SE = 0.011, LRT p = 0.009; unfamiliar:
β = 0.034, SE = 0.009, LRT p < 0.001, Table S1; Figure 2).
186

Figure 1. Sickness behavior: bats with *Staphylococcus* infections spent less time
allogrooming and moving. Plots show mean proportions ± bootstrapped 95% CIs for behaviors of
infected (red triangle) and healthy (black circle) female bats during the outbreak (pre-treatment) and
after they had recovered (post-treatment). We did not detect effects of infection on resting or self-
grooming either before or after antibiotic treatment. Figure S1 shows this plot with infected bats
divided into 'treated' and 'injured' bats.
194
195

196
197
Figure 2. Evidence for effects of infection on social integration. Left panel shows changes in the
probability of receiving grooming over time from a familiar groomer to a receiver bat that was healthy
(solid cyan), treated (dashed orange), or injured (dotted red). Middle panel shows the same plot but
for new (unfamiliar) pairs of bats. In both left and middle panels, yellow shaded region indicates the
week prior to antibiotic treatment (pre-treatment) and blue shaded region indicates one week at the
end of the experiment (post-treatment). Right panel shows the change in log odds (from middle
panel) for each type of receiver. Large points with error bars are means and bootstrapped 95% CIs.
205
Discussion
This study presents opportunistic and observational evidence for reduced social integration
in infected bats. Our results should be treated with caution due to the impossibility of estimating
effect sizes precisely with small sample sizes (9 infected bats with varying levels of infection
severity). However, while several studies have quantified social interactions when animals are
infected with, or exposed to pathogens (14,16–18,33–38), to our knowledge, no studies have
focused on the impact of infections on social bond formation. Our findings and effect size estimates
could therefore be valuable for informing future studies that test this hypothesis in more detail.
We found that, relative to healthy bats, the 9 bats initially infected with *Staphylococcus* (and
the 4 injured bats in particular) exhibited sickness behavior; they spent less time moving and
allogrooming. As expected from an infectious pathogen, infections were biased based on the bat's
origin. The top five most severely infected bats and all 4 injured bats were from the same population.
Several results are consistent with the hypothesis that acute infections impacted the
development of social relationships. Infected bats gave and received less allogrooming with new

partners. Infection did not predict reduced allogrooming as much in familiar partners. These differences faded during 3 months of recovery, suggesting they were caused by short-term sickness behavior rather than by permanent injuries that reduced movement (Figure 2).

A possible alternative explanation is the reverse causal relationship: bats with worse social integration had weakened immune systems and therefore suffered worse infections because of their social traits. However, this hypothesis does not explain why severe infections only occurred in bats from the same population or why the gap in social integration disappeared over time after antibiotic treatment. If the infected bats were already less social or less healthy for other reasons besides the infection, then we would expect their social integration to be worse overall and not just during the outbreak.

The sickness behavior we observed was similar to the responses of vampire bats to simulated infections with injections of lipopolysaccharide, LPS (12,20). We quantified sickness behaviors in response to a naturally-infecting pathogen, which is less often done for vertebrates (but see 33,35,39,40) compared to studies on sickness behavior using an immune challenge (e.g., (8,10–12,15,20,24,31,41–43). The use of immune challenges, like LPS, allows for controlled experiments on the connection between the inflammatory and the behavioral response to infection, but naturally infecting pathogens can have different, pathogen-specific effects**.** For instance, LPS-injected vampire bats reduced their self-grooming (12), but *Staphylococcus*-infected bats did not**,** possibly because *Staphylococcus* causes skin lesions. Similarly, little brown bats (*Myotis lucifugus*) infected with *Pseudogymnoascus destructans* (white-nose syndrome), a fungal skin disease, showed increased self-grooming (39).

Our findings raise several questions about how short-term infections might impact relationship formation and long-term social integration. When might vampire bats or other animals suppress their infection-induced behavioral symptoms to continue social interactions and integration (20,41,42,44–46)? At what stages of relationship formation do sickness behaviors have the strongest effect on social interactions? How long must sickness behavior last for it to cause lingering social effects? To test for potential trade-offs between the severity of behavioral symptoms and continued social interactions, researchers could use immune challenges that are dose-dependent (42,47). Immune challenges like LPS could also be used to test at what stage of relationship formation sickness behaviors have the strongest effects, and for how long they would have to last for lingering effects on social integration. Immune challenges can be used to control the duration and precise onset times of symptomatic periods.

In conclusion, social integration benefits health, immune function, and susceptibility to disease (1–4). Conversely, however, infections might also reduce social integration in several ways. Sickness behavior can cause individuals to be less socially active. Cues of infection have clear impacts on mate choice (reviewed in 23) and there may be similar effects on choice of other kinds of social partners (e.g., same-sex social bonds). If there are critical periods of social integration, then infections during this time might have long-term social consequences. Consistent with this possibility, we presented opportunistic and observational evidence for reduced social integration in vampire bats after an outbreak of a naturally infecting pathogen. While our sample sizes are small and do not allow us to estimate precise effect sizes, we view our results as a starting point for future studies that address the effect of infections on long-term social integration and relationship formation.

Acknowledgements

The authors thank Rachel Page for lab space and resources, the staff at the Smithsonian Tropical Research Institute, and the following individuals who assisted with vampire bat care and/or data collection: D. Aparicio, B. Brown, G. Cohen, L. Dück, D. Girbino, E. Kline, C. Marroquin, and S. Ripperger.

Funding

I.R. was supported by a short-term fellowship from the Smithsonian Tropical Research Institute, a student research grant from the Animal Behaviour Society, a Graduate Enrichment Fellowship from The Ohio State University, a Tinker Foundation Field Research Grant, a Grant-in-aid of Research from the American Society of Mammologists, and a Smithsonian Institution Predoctoral Fellowship. S.S. was supported by the Ohio State University President's Postdoctoral Scholars Program. This publication is based upon work supported by the National Science Foundation under grant no. IOS-2015928 to G.G.C.

Author contributions
Data Curation: IR, GC
Formal Analysis: IR, SS, GC
Funding Acquisition: IR, GC
Investigation: IR, MA, GC
Methodology: IR, GC
Project Administration: IR, GC
Resources: IR, GC
Software: IR, GC
Supervision: GC
Validation: IR, SS, GC
Visualization: IR, SS, GC
Writing – Original Draft Preparation: IR, SS, GC
Writing – Review & Editing: IR, SS, GC

References

1. Pavez-Fox MA, Negron-Del Valle JE, Thompson IJ, Walker CS, Bauman SE, Gonzalez O, et al. Sociality predicts individual variation in the immunity of free-ranging rhesus macaques. Physiol Behav. 2021;241:113560.

2. Snyder-Mackler N, Burger JR, Gaydosh L, Belsky DW, Noppert GA, Campos FA, et al. Social determinants of health and survival in humans and other animals. Science. 2020;368(6493):eaax9553.

3. Snyder-Mackler N, Sanz J, Kohn JN, Brinkworth JF, Morrow S, Shaver AO, et al. Social status alters immune regulation and response to infection in macaques. Science. 2016;354(6315):1041–5.

4. Tung J, Barreiro LB, Johnson ZP, Hansen KD, Michopoulos V, Toufexis D, et al. Social environment is associated with gene regulatory variation in the rhesus macaque immune system. Proc Natl Acad Sci. 2012;109(17):6490–5.

5. Almberg ES, Cross PC, Dobson AP, Smith DW, Metz MC, Stahler DR, et al. Social living mitigates the costs of a chronic illness in a cooperative carnivore. Ecol Lett. 2015;18(7):660–7.

6. Loehle C. Social barriers to pathogen transmission in wild animal populations. Ecology. 1995;76(2):326–35.

7. Hart BL. Biological basis of the behavior of sick animals. Neurosci Biobehav Rev. 1988;12(2):123–37.

8. Lopes PC, French SS, Woodhams DC, Binning SA. Sickness behaviors across vertebrate taxa: proximate and ultimate mechanisms. J Exp Biol. 2021;224(9):jeb225847.

9. Stockmaier S, Stroeymeyt N, Shattuck EC, Hawley DM, Ancel Meyers L, Bolnick DI. Infectious
diseases and social distancing in nature. Science. 2021;371(6533):eabc8881.

10. Fishkin RJ, Winslow JT. Endotoxin-induced reduction of social investigation by mice: interaction
with amphetamine and anti-inflammatory drugs. Psychopharmacol Berl. 1997;132(4):335–41.

11. Moreno KR, Weinberg M, Harten L, Salinas Ramos VB, Herrera M. LG, Czirják GÁ, et al. Sick
bats stay home alone: fruit bats practice social distancing when faced with an immunological
challenge. Ann N Y Acad Sci. 2021;1505(1):178–90.

12. Stockmaier S, Bolnick DI, Page RA, Carter GG. An immune challenge reduces social grooming
in vampire bats. Anim Behav. 2018;140:141–9.

13. Lopes PC, Block P, Konig B. Infection-induced behavioural changes reduce connectivity and the
potential for disease spread in wild mice contact networks. Sci Rep. 2016;6:31790.

14. Fairbanks BM, Hawley DM, Alexander KA. The Impact of Health Status on Dispersal Behavior in
Banded Mongooses (Mungos mungo). EcoHealth. 2014;11(2):258–62.

15. Lopes PC, Woodhams DC, Binning SA. Infection avoidance behaviors across vertebrate taxa:
patterns, processes and future directions. In: Animal Behavior and Parasitism. Oxford University
Press; 2022.

16. Behringer DC, Butler MJ, Shields JD. Ecology: avoidance of disease by social lobsters. Nature.
2006;441(7092):421.

17. Kiesecker JM, Skelly DK, Beard KH, Preisser E. Behavioral reduction of infection risk. Proc Natl
Acad Sci U A. 1999;96(16):9165–8.

18. Poirotte C, Massol F, Herbert A, Willaume E, Bomo PM, Kappeler PM, et al. Mandrills use
olfaction to socially avoid parasitized conspecifics. Sci Adv. 2017;3(4).

19. Boillat M, Challet L, Rossier D, Kan C, Carleton A, Rodriguez I. The Vomeronasal System
Mediates Sick Conspecific Avoidance. Curr Biol. 2015;25(2):251–5.

20. Stockmaier S, Bolnick DI, Page RA, Carter GG. Sickness effects on social interactions depend
on the type of behaviour and relationship. J Anim Ecol. 2020; 89:1387-1394.

21. Poirotte C, Charpentier MJE. Unconditional care from close maternal kin in the face of parasites.
Biol Lett. 2020;16(2):20190869.

22. Stockmaier S, Ulrich Y, Albery GF, Cremer S, Lopes PC. Behavioural defences against
parasites across host social structures. Funct Ecol. 2023; 37, 809– 820.

23. Beltran-Bech S, Richard FJ. Impact of infection on mate choice. Anim Behav. 2014;90:159–70.

24. Love AC, Anthony AC, Nash A, Campos-Melara A, Kodali J, DuRant SE. Simulated infection
alters the behavior of pair bonded songbirds and their healthy neighbors. Behav Ecol.
2023;34(2):251–60.

25. Razik I, Brown BKG, Carter GG. Forced proximity promotes the formation of enduring
cooperative relationships in vampire bats. Biol Lett. 2022;18(4):20220056.

26. Carter GG, Farine DR, Crisp RJ, Vrtilek JK, Ripperger SP, Page RA. Development of New Food-
Sharing Relationships in Vampire Bats. Curr Biol. 2020;30(7):1275-1279.e3.

27. Razik I, Brown BKG, Page RA, Carter GG. Non-kin adoption in the common vampire bat. R Soc
Open Sci. 202;8(2):201927.

28. Canty A, Ripley B. boot: Bootstrap R (S-Plus) functions. R Package Version. 2017;1:3–20.

29. Douglas Bates MM. Fitting Linear Mixed-Effects Models Using lme4. J Stat Softw. 2015;67:1–48.

30. Razik I, Carter G, Stockmaier S, Abou-Elias M. Data and R code for Opportunistic evidence that
Staph infection affects social integration in vampire bats. 2023; Available from:
https://doi.org/10.6084/m9.figshare.22573018.v1

31. Wilkinson GS. Social grooming in the common vampire bat, Desmodus rotundus. Anim Behav.
1986;34(6):1880–9.

32. Carter GG, Wilkinson GS. Food sharing in vampire bats: reciprocal help predicts donations more
than relatedness or harassment. Proc Biol Sci. 2013;280(1753):20122573.

33. Cárdenas-Canales EM, Stockmaier S, Cronin E, Rocke TE, Osorio JE, Carter GG. Social effects
of rabies infection in male vampire bats (Desmodus rotundus). Biol Lett. 2022;18(9):20220298.

34. Fairbanks BM, Hawley DM, Alexander KA. No evidence for avoidance of visibly diseased
conspecifics in the highly social banded mongoose (Mungos mungo). Behav Ecol Sociobiol.
2015;69(3):371–81.

35. Bouwman KM, Hawley DM. Sickness behaviour acting as an evolutionary trap? Male house
finches preferentially feed near diseased conspecifics. Biol Lett. 2010;6(4):462–5.

36. Stephenson JF. Parasite-induced plasticity in host social behaviour depends on sex and
susceptibility. Biol Lett. 2019;15(11):20190557.

37. Stroeymeyt N, Grasse AV, Crespi A, Mersch DP, Cremer S, Keller L. Social network plasticity
decreases disease transmission in a eusocial insect. Science. 2018;362(6417):941.

38. Cremer S. Social immunity in insects. Curr Biol. 2019;29(11):R458–63.

39. Brownlee-Bouboulis SA, Reeder DM. White-nose syndrome-affected little brown myotis (Myotis
lucifugus) increase grooming and other active behaviors during arousals from hibernation. J
Wildl Dis. 2013;49(4):850–9.

40. Bos N, Lefevre T, Jensen AB, d'Ettorre P. Sick ants become unsociable. J Evol Biol.
2012;25(2):342–51.

41. Lopes PC, Adelman J, Wingfield JC, Bentley GE. Social context modulates sickness behavior.
Behav Ecol Sociobiol. 2012;66(10):1421–8.

42. Willette AA, Lubach GR, Coe CL. Environmental context differentially affects behavioral,
leukocyte, cortisol, and interleukin-6 responses to low doses of endotoxin in the rhesus monkey.
Brain Behav Immun. 2007;21(6):807–15.

43. Alciatore G, Ugelvig LV, Frank E, Bidaux J, Gal A, Schmitt T, et al. Immune challenges increase
network centrality in a queenless ant. Proc R Soc B. 2021;288(1958):20211456.

44. Aubert A, Goodall G, Dantzer R, Gheusi G. Differential Effects of Lipopolysaccharide on Pup
Retrieving and Nest Building in Lactating Mice. Brain Behav Immun. 1997;11(2):107–18.

45. Lopes PC. When is it socially acceptable to feel sick? Proc Biol Sci. 2014;281(1788):20140218.

46. Weil ZM, Bowers SL, Dow ER, Nelson RJ. Maternal aggression persists following
lipopolysaccharide-induced activation of the immune system. Physiol Behav. 2006;87(4):694–9.

47. Armour EM, Bruner TL, Hines JK, Butler MW. Low-dose immune challenges result in detectable
levels of oxidative damage. J Exp Biol. 2020;223(6).

399