DESpace: spatially variable gene detection via differential expression testing of spatial clusters

Peiying Cai[1], Mark D Robinson[1], and Simone Tiberi[1,2*]

[1]*Department of Molecular Life Sciences and Swiss Institute of Bioinformatics, University of Zurich, Zurich, Switzerland.*
[2]*Department of Statistical Sciences, University of Bologna, Bologna, Italy.*
[*] e-mail: Simone.Tiberi@unibo.it

Abstract

Spatially resolved transcriptomics (SRT) enables scientists to investigate spatial context of mRNA abundance. Here, we introduce *DESpace*, a novel approach to discover spatially variable genes (SVGs), i.e., genes whose expression varies across the tissue. Our framework inputs all types of SRT data, summarizes spatial information via spatial clusters, and identifies spatially variable genes by performing differential gene expression testing between clusters. Although several methods have been proposed to identify SVGs, our approach adds some unique features; in particular: it allows identifying (and testing) the specific areas of the tissue affected by spatial variability, and it enables joint modelling of multiple samples (i.e., biological replicates). Furthermore, in our benchmarks, *DESpace* displays a higher true positive rate than competitors, controls for false positive and false discovery rates, and is among the most computationally efficient SVG tools. *DESpace* is distributed as a Bioconductor R package.

keywords: Spatially resolved transcriptomics (SRT); Spatial omics; spatially variable genes; Differential gene expression; Spatial clustering.

1 Introduction

Spatially resolved transcriptomics (SRT) technologies allow the spatial characterization of gene expression profiles in a tissue. SRT techniques can be broadly grouped into two categories: sequencing-based methods (e.g., Slide-seq [29], Slide-seq-V2 [33], 10X Genomics Visium [1], spatial transriptomics [34], high definition spatial transcriptomics [37], STEREO-seq [6]), and in situ hybridization technologies (e.g., seqFISH [13], seqFISH2 [30], seqFISH+ [12], and MERFISH [17]). Sequencing-based approaches provide measurements across the entire transcriptome but the mRNA abundances (typically) refer to the aggregation of multiple cells; conversely, the imaging-based methods are targeted towards a (usually) limited number of genes but offer higher spatial resolution, e.g., molecular-level measurements. In both cases, computationally efficient methods are required to deal with an increasingly large number of genes and spatial measurements.

The emergence of SRT technologies has prompted the development of novel analysis frameworks that exploit the joint availability of gene expression and spatial information; in particular, below, we briefly introduce two such schemes: spatially resolved clustering and spatially variable gene identification. Other analyses include strategies to identify putative cell-cell communication via local ligand-receptor co-expression, and enrichment of cell type interactions [24].

Figure 1: Spatial clusters in three samples from *LIBD* [22] (Visium [1]), *melanoma* [35] (spatial transcriptomics [34]) and *mouse cerebellum* [5] (Slide-seqV2 [33]) datasets. Spatial clusters were obtained: via manual annotations form a pathologist (*LIBD*), *BayesSpace* [11] (*melanoma*), and *StLearn* [23] (*mouse cerebellum*).

While single-cell RNA-sequencing enables clustering cells based on transcriptional profiles, SRT allows the identification of spatial clusters (Figure 1). Such structures are typically estimated via spatially resolved clustering tools, such as *BayesSpace* [11], *StLearn* [23], *Giotto* [28], and *PRECAST* [18], which cluster spots based on both their gene expression and spatial localization; alternatively, spatial clusters can also be obtained via manual annotation, e.g., from histology [22]. Another popular analysis performed on SRT data consists in identifying genes whose expression patterns change across the tissue, also known as spatially variable genes (SVGs). In recent years, various methods have been proposed to discover SVGs; notably: *MERINGUE* [4], a density-independent approach built on the spatial autocorrelation of each gene; *nnSVG* [38], which fits nearest-neighbor Gaussian processes; *SpaGCN* [14], that identifies spatial domains based on a graph convolutional network, and performs a Wilcoxon rank-sum test on the mean-level expression changes between domains; *SPARK* [31], using generalized linear mixed models with overdispersed Poisson and Gaussian distributions; *SPARK-X* [32], from the authors of *SPARK*, based on a scalable non-parametric framework to test the dependence between gene expression covariance and distance covariance; *SpatialDE* [36], which infers the dependency between space and gene expression based on a non-parametric Gaussian process; *SpatialDE2* [16], that applies a Bayesian hidden Markov random field to segment spatial regions and a count-based likelihood to model the variance components of each gene within identified regions; and, *Trendsceek* [8], which evaluates significant spatial gene expression heterogeneity via marked point processes.

Despite the abundance of SVG tools, there are some limitations with the current approaches, including that: i) most methods are computationally demanding, which is particularly troublesome for recent SRT technologies, such as Slide-seq-V2, which provide hundreds of thousands of measurements per gene; ii) biological replicates are not allowed, and only individual samples can be processed, exposing results to a high degree of (biological and technical) variability; iii) few tools can incorporate information about spatial structures of interest, such as spatial clusters; and, iv) it is only possible to test the entire tissue for spatial variability (SV), while scientists cannot perform SV testing on specific regions of interest (e.g., white

matter in brain cortex).

In order to address these limitations, we propose *DESpace*, which is based on differential expression testing of spatial clusters. Our method requires pre-computed spatial clusters (Figure 1), which can be obtained via manual annotation or from spatially resolved clustering tools (e.g., *BayesSpace* [11], *StLearn* [23], Giotto [28], and PRECAST [18]); these spatial clusters are used as a proxy for the spatial information. We then fit a negative-binomial model, which was recently shown to accurately model SRT data [3, 40], via *edgeR* [7, 10, 21], a popular tool for differential gene expression; we use spatial clusters as covariates, and perform a differential gene expression test across these clusters. If the expression of a gene is significantly associated to the spatial clusters, its expression varies across the tissue, thus indicating a SVG. Clearly, our approach relies on spatial clusters being available, and accurately summarizing the spatial patterns of transcription, but we argue that these assumptions are usually fulfilled. In fact, even when pre-computed annotations are not available, spatially resolved clustering tools enable inferring clusters that summarize the spatial structure of gene expression. The method we propose has several advantages with respect to alternative approaches for SVG detection. In our benchmarks (see Results), *DESpace* displays good sensitivity and specificity, and is computationally efficient; furthermore, to the best of our knowledge, *DESpace* offers some unique features, compared to currently available SVG tools. In particular, our approach: i) can jointly model multiple samples, hence increasing statistical power, reducing the uncertainty that characterizes inference performed from individual samples, and identifying genes with coherent spatial patterns across biological replicates; ii) allows identifying the key areas of the tissue affected by SV, testing if the average expression in a particular region of interest (e.g., cancer tissue) is significantly higher or lower than the average expression in the remaining tissue (e.g., non-cancer tissue), hence enabling scientists to investigate changes in mRNA abundance in specific areas which may be of particular interest. Finally, our method is computationally efficient and flexible, and can input any type of SRT data.

Results

Inference with *DESpace*

Consider a SRT dataset, from a single sample, with N spots, and where a total of C spatial clusters have been identified; further assume that the c-th cluster corresponds to N_c spots, where $\sum_{c=1}^{C} N_c = N$. We define x_{gi} as the mRNA abundance for the g-th gene in the i-th spot belonging to the c-th cluster; we model this value with a negative binomial distribution:

$$x_{gi} \sim NB(\mu_{gi}, \phi_g), \tag{1}$$
$$log(\mu_{gi}) = log(M_i) + \beta_{gc}, \tag{2}$$

where $NB(\mu, \phi)$ denotes a negative binomial random variable with mean μ and variance $\mu(1 + \mu\phi)$; ϕ_g is the gene-specific dispersion parameter modeling the variability between spots; M_i is the effective library size (total count multiplied by TMM normalization factor [27]) for the i-th spot; and β_{gc} is the c-th spatial cluster coefficient for gene g, which represents the abundance of gene g in spatial cluster c.

In order to investigate if gene g is spatially variable, we verify if its gene expression varies across spatial clusters. In particular, we employ a likelihood ratio test (LRT) [39] to test whether the β_{gc} parameters differ across clusters, via the following system of hypotheses:

$$\begin{cases} H_0: & \beta_{g1} = \cdots = \beta_{gC} \\ H_1: & \text{otherwise.} \end{cases} \tag{3}$$

In order to identify the specific regions affected by spatial variability, for each gene, we also apply a LRT on each individual spatial cluster parameter, β_{gc}.

When biological replicates are available, we add a sample-specific parameter to the mean of the negative binomial model in (1). In particular, assuming that the observation comes from the

174 j-th replicate, equation (2) becomes:

$$log(\mu_{gi}) = log(M_i) + \beta_{gc} + \gamma_{gj}, \quad (4)$$

where γ_{gj} indicates the coefficient for the j-th
sample in gene g, with $\gamma_{g1} = 0$ for the first sam-
ple (taken as baseline) to ensure identifiability.
SVGs are identified, as in the individual-sample
case, by testing for the significance of β_{gc}'s pa-
rameters.
In all cases, parameter inference and differential
testing are performed via *edgeR* [7,10,21]. Note,
that we also tried alternative approaches for dif-
ferential gene expression, namely *DESeq2* [19]
and *limma* [26], but overall performance (in
terms of sensitivity, specificity, and runtime)
was better when using *edgeR*.

Simulation studies - anchor data

We have designed several benchmarks, on real
and semi-simulated data; all our analyses start
from one of three SRT datasets (collected from
distinct spatial technologies), denoted by *LIBD*
[22], *melanoma* [35] and *mouse cerebellum* [5]
(Figure 1).
The *LIBD* [22] data, collected via the Vi-
sium platform [1], contains 12 dorso-lateral pre-
frontal cortex samples from 3 independent hu-
man brain donors (i.e., biological replicates).
In all samples, the tissue was partitioned via
manual annotation from pathologists into white
matter and up to six layers (Figure 1). Overall,
this dataset contains, for each sample, on aver-
age, measurements for 33,538 genes across 3,973
spots; after filtering (see Methods), this reduces
to 14,628 genes and 3,936 spots (Supplementary
Table 1).
The *melanoma* [35] data consists of 8 melanoma
lymph node samples from 4 distinct individuals
(i.e., biological replicates), diagnosed with stage
IIIc melanoma. This data was obtained by ap-
plying spatial transcriptomics (ST) [34] tech-
nology to melanoma lymph node biopsies. In
the original study [35], the spots of each sam-
ple were clustered in four groups: melanoma,
stroma, lymphoid tissue, and remaining spots.
We clustered spots in four spatial clusters us-

ing *BayesSpace* (Figure 1), where cluster 4 indi-
cates the melanoma area, cluster 2 embeds both
stroma and lymphoid tissues, cluster 3 repre-
sents the border region between lymphoid and
tumor tissues, and cluster 1 includes lymphoid
tissue distant from the cancer. The dataset, on
average, consists of a total of 15,884 genes per
sample, measured on 294 spots; after filtering
(see Methods), we retained 8,173 genes and 290
spots (Supplementary Table 1).
The *mouse cerebellum* [5] data, obtained via
Slide-seqV2 [33] technology, consists of mea-
surements, from a single sample, for 20,141
genes across 11,626 beads. After filtering (see
Methods), these numbers reduced to 8,277 and
11,615, respectively (Supplementary Table 1).
Beads were clustered via *StLearn* (Figure 1),
obtaining similar clusters to those computed by
Zhu *et al.* (2021) [32], on the same dataset,
using RCTD software [5].
Note that for the semi-simulated data from the
LIBD and *melanoma* datasets, we generated
one simulation for each biological replicate (3
and 4 in the former and latter case, respec-
tively).

Simulation studies - SV profiles

We generated various semi-simulated datasets,
by using, as anchor data, the previously de-
scribed three datasets. In particular, we edited
the *runPatternSimulation* function from *Giotto*
[28], and re-arranged the real measurements to
partition the space into a highly and a lowly
abundant region (see Methods); by partition-
ing the tissue in different ways, we generated
five different SV patterns. We initially sepa-
rated the space in artificial regions: top *vs.* bot-
tom (or right *vs.* left) areas (Figure 2, *Bot-*
tom/Right), and inside *vs.* outside a circle
(Figure 2, *Circular*). Then, we followed real
data structures such as brain cortex layers and
melanoma regions (Figure 2, *Annotations*). In
these three scenarios, in each simulated dataset,
half of the SVGs are highly abundant in one re-
gion (e.g., bottom or centre of the circle), and
half are highly abundant in the complementary
region (e.g., top or outside the circle). However,

Figure 2: Three examples of simulated SVGs, from the *LIBD* data, following *bottom/right*, *circular*, and *annotations* patterns. Examples of SVGs from *mixture* and *inverted mixture* patterns are presented in Supplementary Figure 1.

in the cases described above, all SVGs follow the same spatial structures, which is unlikely to happen in real data. In order to improve the realism of the semi-simulated data, we designed two additional simulations, where SVGs follow multiple spatial patterns, all based on real data structures; we generated two such structures, referred to throughout as *mixture* and *inverted mixture*. In the former case, a small region is identified as highly abundant, while most of the tissue is lowly abundant; conversely, in the latter case, gene expression is uniform in most of the tissue, and a small region is characterized by low abundance (Supplementary Figure 1). In each simulation, we generated between 33% (*bottom/right*, *circular*, and *annotations* patterns) and 50% (*mixture*, and *inverted mixture* patterns) of SVGs, while the remaining genes are uniformly distributed; such uniform patterns were obtained by randomly permuting measurements across the tissue (see Methods). Note that, when running *DESpace*, we did not use the original spatial clusters that we simulated from. Instead, we inputted noisy spatial clusters inferred via *StLearn* and *BayesSpace*. We choose these two spatially resolved clustering tools, because they performed well in recent benchmarks [11].

Simulation studies - individual sample

In all simulations, we bechmarked *DESpace* against seven of the most popular tools for SVG detection: *SpaGCN* [14], *SpatialDE* [36], *SpatialDE2* [16], *SPARK* [31], *SPARK-X* [32], *MERINGUE* [4], and *nnSVG* [38]. Note that *Trendsceek* was excluded from our analyses, due to its extremely high computational cost [31, 32].

Figure 3 reports the true positive rate (TPR) *vs.* false discovery rate (FDR) of all methods for the three datasets and five spatial patterns. Rates for the *LIBD* and *melanoma* samples are aggregated across biological replicates; the results for the individual samples can be seen in Supplementary Figures 2-3. In all five SV patterns, *DESpace* controls for the false discovery rate, and leads to a higher statistical power than competitors; interestingly, this gap is smaller in simpler SV patterns (*bottom/right* and *circular*), and increases for the more complex ones (*annotations*, *mixture* and *inverted mixture*). This suggests that our framework may be particularly beneficial for detecting challenging spatial structures. Notably, results are highly consistent across the three datasets, and *DESpace* leads to analogous results when inputting spatial clusters estimated from *StLearn*

Figure 3: TPR vs. FDR for SVG detections in the individual sample simulations. Rows and columns refer to the anchor data used in the simulation, and to the SV profiles, respectively, *BayesSpace_DESpace* and *StLearn_DESpace* indicate *DESpace* based on spatial clusters computed via *BayesSpace* and *StLearn*, respectively.

or *BayesSpace*, which indicates that our approach is robust with respect to the spatial clusters provided.

Furthermore, in order to investigate false positive detections, in each simulation, we selected the subgroup of uniform genes (i.e., non-SVGs), and studied the distribution of raw p-values. Overall, most methods (including *DESpace*) display approximately uniform p-values, and none presents a significant inflation of false positives, with the only exception of *SPARK* on the *mouse cerebellum* data (Supplementary Figure 4). Notably, *SpatialDE* and *nnSVG* produce highly conservative results, with p-values significantly inflated towards 1, which is consistent with previous findings [31, 38].

Additionally, we investigated if methods are able to sort SVGs according to the strength of their spatial structure. In fact, it is desirable that genes displaying strong spatial patterns are ranked before (i.e., smaller p-value) genes with weaker spatial patterns, because the first ones are more likely to be biologically relevant. To this aim, we generated a simulation with SVGs only, where 50% of the genes display a strong spatial pattern, and 50% display a weak one (see Methods). Supplementary Figure 5 reports the TPRs for the strong and weak patterns; overall, *DESpace* ranks strong SV patterns better than competitors.

Figure 4: Expression plot, for four SVGs detected with *DESpace* individual cluster test, on the real *LIBD* dataset (sample 151673). SVGs were identifying by selecting high and low expression in white matter (genes MOBP and ENC1, respectively), and high and low abundance in layer 3 (genes HOPX and HS3ST4, respectively). Lines highlight the cluster being tested.

Simulation studies - individual cluster

We also investigated *DESpace*'s ability to identify the key spatial regions involved in SVGs. Note that, when only 2 spatial clusters are available, individual cluster testing corresponds to the global test described above; therefore, this analysis framework can only be applied when at least 3 clusters are available, and is particularly beneficial when there is a large number of clusters. To validate this approach, we considered the *mixture* and *inverted mixture* simulations in the *LIBD* dataset, which contain between 5 and 7 spatial clusters per biological replicate. In these simulations, it is possible to define a key spatial cluster that displays the biggest difference compared to the average signal; in the *mixture* (or *inverted mixture*) simulations this corresponds to the small region with higher (or lower) abundance than the majority of the tissue (Supplementary Figure 1).

On average, when using *BayesSpace* clusters, *DESpace* identified the main SV cluster in 99.6 and 88.9% of SVGs in *mixture* and *inverted mixture* patterns, respectively; using *StLearn* clusters led to similar numbers Table 1). Note that, in all samples, percentages are higher in the *mixture* simulations, because the change in abundance, between the key spatial cluster and the rest of the tissue, is higher compared to the *inverted mixture* patterns (Supplementary Figure 1). Furthermore, in non-SVGs, p-values were uniformly distributed in all spatial clusters, showing no inflation of false positive detections at the cluster level (Supplementary Figures 6-7). This testing feature may be particularly useful for computational biologists, as it enables them to directly target key regions of the tissue, for example by identifying SVGs characterized by high or low abundance in white matter or layer 3 (Figure 4). Finally, note that we also provide a faster implementation of the individual cluster testing, where we recycle the previously-computed dispersion estimates from the gene-level test (see Methods); this strategy performs similarly to the original one (analogous fraction of key spatial clusters identified), while the average runtime decreases significantly: from 32 to 2 minutes per sample (Supplementary Figures 8-9 and Supplementary Table 2).

Pattern	Sample	*BayesSpace*	*StLearn*
Mixture	151507	99.6	98.7
	151669	99.4	99.5
	151673	99.8	99.0
	Average	99.6	99.1
Inverted Mixture	151507	86.6	79.7
	151669	89.3	88.8
	151673	90.9	85.5
	Average	88.9	84.7

Table 1: Individual cluster results. Percentage of times that *DESpace* identified the main SV cluster in each *mixture* and *inverted mixture* simulations, using *BayesSpace* and *StLearn* clusters.

Figure 5: TPR vs. FDR for SVG detections in the multiple sample simulation. Rows and columns refer to the anchor data used in the simulation, and to the SV profiles, respectively. *Single_sample_DESpace* and *Multiple_samples_DESpace* indicate *DESpace* on single-sample and multi-sample modes, respectively. Note that TPRs are lower than in the individual simulation (Figure 3), because we have simulated slightly weaker spatial patterns here (see Methods).

Simulation studies - multiple samples

Using the *LIBD* and *melanoma* datasets, which contain three and four biological replicates, respectively, we designed a multi-sample simulation: we simulated the 5 SV profiles shown before, and ensured that SVGs (and their spatial profiles) are consistent across samples; we then fit *DESpace* on, both, single-sample and multi-sample modes. Here, we aim to compare the two *DESpace* approaches, based on identical input data; therefore, in this analysis, we used the same spatial clusters (i.e., the annotations we simulated from) for both single-sample and multi-sample modes. While both approaches control for the FDR, jointly modelling multiple samples leads, in all cases, to a significant increase of the TPR, which is expectable given that more information is available (Figure 5).

Applications to real data

We also applied all methods to the three real datasets considered. For the *LIBD* and *melanoma* datasets, we considered all 12 and 8 samples, respectively. In all datasets, we computed spatial clusters via *BayesSpace* and *StLearn*; in the *LIBD* dataset, we additionally considered pre-computed manually annotated clusters. In all cases, *DESpace* identifies significantly more SVGs (at 1% FDR threshold) than any other method, which is coherent with the higher statistical power displayed in simulations.

In *LIBD* and *melanoma* data, we also investigated the coherency of the top ranked genes across different replicates. In order to do that, we used the Jaccard index [15], which ranges between 0 and 1, and measures the similarity between two pairs of groups. In particular, for every method, we considered the top 1,000 (LIBD) or 200 (melanoma) ranked genes (i.e., smallest p-value) in each sample, and computed the Jaccard index on pairs of samples; we then averaged results across samples. In both datasets, *DESpace*, *SPARK* and textitSPARK-X display the highest Jaccard index, which indicates a greater coherency of top ranked genes across samples (Figure 6). Note that, the Jaccard index of all methods is significantly lower in the *melanoma* dataset, compared to the *LIBD* data, which is reasonable given the high degree of gene expression variability in cancer.

Figure 6: Jaccard index on the *LIBD* and *melanoma* datasets. For each method, the index is measured on the sets of top SVGs reported across replicates, and represents how coherent results are between samples: higher values indicate greater coherency.

Finally, we investigated the computational efficiency of methods, and computed each tool's runtime on the three real datasets (Figure 7). The ranking of methods is similar across datasets: *SPARK-X* and *SpaGCN* excel as the fastest methods, followed by *DESpace*, *SpatialDE* and *SpatialDE2*, while the slowest three methods appear to be *nnSVG*, *SpaGCN*, and *SPARK*. Note that *DESpace*'s time largely depends on the computational cost required to generate spatial clusters; in particular, in all datasets, *StLearn* is significantly faster than *BayesSpace*.

Discussion

In this manuscript, we have presented *DESpace*, an approach, based on differential testing across spatial structures, to identify spatially variable genes from SRT data. We have run extensive benchmarks on both semi-simulated and real datasets, where our method displays good control of false positives, a higher statistical power (i.e., TPR) than competitors, and is one of the most computationally efficient methods. Furthermore, to the best of our knowledge, our method is the only SVG tool that can i) iden-

tify the key SV regions, and ii) jointly model multiple samples. The former feature allows performing statistical testing on specific regions of the tissue, while the latter one provides increased TPR, and enables the identifications of SVGs with consistent spatial patterns across replicates.

Additionally, our approach is flexible and can work with SRT data from distinct technologies. Finally, *DESpace* is distributed, open-access, as a Bioconductor R package, which makes it easy to install and integrate with existing pipelines (such as the popular *SpatialExperiment* class [25]), is accompanied by an example vignettes that illustrate its usage, and includes plotting functions that facilitate visualization and interpretation of results.

We also acknowledge some limitations of our approach. In particular, our framework requires pre-computed spatial clusters and therefore is prone to detecting genes that follow such spatial structures. Nonetheless, we argue that spatial clusters can be successfully obtained in most datasets (e.g., via spatially resolved clustering tools), and that such clusters typically capture the expression patterns of SVGs. Furthermore, we are aware of the circularity (or double dip-

Figure 7: Runtime (in minutes), of each method, in the real datasets; for the *LIBD* and *melanoma* data, we report the average runtime across replicates. For *DESpace*, we colour differently the runtime to compute spatial clusters (brown top region), and the one of *DESpace*. Note that, for the *LIBD* dataset, *SPARK* successfully ran on Sample 151507 only, hence *SPARK*'s runtime refers to that sample only.

ping) issue of our framework; indeed, when relying on spatially resolved clustering tools, we use our data twice: i) to cluster spots, and ii) to perform differential analyses between such clusters. Nonetheless, each individual gene contributes marginally to spatial clustering results, which are (usually) based on the expression data from thousands of genes. Furthermore, in all our benchmarks, we have empirically shown that this issue does not impact performance results (i.e., good statistical power, and well calibrated false positive rates).

Finally, a consideration about the computational aspect: recent SRT technologies allow measurements at sub-cellular resolution; this leads to a significant increase in the size of datasets and, in turns, of the computational cost of downstream analyses. In order to mitigate the runtime of our tool, one may aggregate the signal from several spots that belong to the same spatial cluster, which would greatly diminish the number of observations available. We aim to explore this in the future to quantify the reduction in the computational burden, and assess how it affects performance.

Availability

DESpace is freely available as a Bioconductor R package at: https://bioconductor.org/ packages/DESpace. The scripts used to run all analyses are available on GitHub (https://github.com/peicai/DESpace_manuscript) and Zenodo (DOI: 10.5281/zenodo.7829817). The *LIBD* [22] data can be accessed via *spatialLIBD* ExperimentHub R Bioconductor packages; the *melanoma* [35] data set is available at https://www.spatialresearch.org/resources-published-datasets/doi-10-1158-0008-5472-can-18-0747/; and the *mouse cerebellum* [5] data can be reached through Broad Institute's single-cell repository (https://singlecell.broadinstitute.org/single_cell/study/SCP948).

Acknowledgements

We acknowledge all members of the Robinson lab for their feedback and suggestions. This work was supported by Swiss National Science Foundation grants 310030_175841 and 310030_204869 to MDR. MDR acknowledges support from the University Research Priority Program Evolution in Action at the University of Zurich.

Author contributions

PC implemented the method, performed all analyses, and contributed to writing the

manuscript. MDR contributed to designing the study and to writing the manuscript. ST conceived the method, designed the study and wrote the manuscript. All authors approve the article.

Competing interests

The authors declare no competing interests.

Methods

Adjusted p-values

In all methods, to control for the false discovery rate, we used adjusted p-values, obtained via Benjamini-Hochberg correction [2].

Quality control and filtering

Quality control followed the guidelines in the online book entitled "Orchestrating Spatially-Resolved Transcriptomics Analysis with Bioconductor" [20]. In particular, low quality spots were identified using *addPerCellQC* function from *Scuttle* R package [9]. Based on total UMI counts per spot, number of genes with non-zero UMI counts per spot (detected genes), proportion of read mapping to mitochondrial genes, and number of cells per spot, low quality spots were filtered with sample-specific thresholds. Furthermore, undetected genes and lowly expressed genes were removed to ensure that each gene had at least 20 non-zero spots of expression.

Simulating SV patterns

To generate our simulations, we initially partitioned the tissue into two regions, R_1 and R_2, with n_1 and n_2 spots, respectively; as described above, we generated a variety of spatial structures. Then, starting from these partitions, we took advantage of the *runPatternSimulation* function from *Giotto*'s R package [28]. In particular, *runPatternSimulation* considers the most highly abundant n_1 genes and assigns them to region R_1 with probability π. When $\pi = 1$, all the most highly abundant n_1 spots are assigned to region R_1, while $\pi = 0.5$ leads to uniform patterns.

Our simulations are composed of a mixture of uniform and SV genes: we obtained uniform patterns by setting $\pi = 0.5$, while SVGs were simulated using $\pi = 0.9$. Only in the multi-sample simulation, SVGs were simulated using $\pi = 0.8$; we choose to decrease π because the TPRs of both single-sample and multi-sample modes were close to 1: decreasing π allowed us to show a difference between the TPRs of the two modes. For the strong *vs.* weak patterns simulation, we set *pi* to 0.9 and 0.6 for the former and latter case, respectively.

In the *bottom/right*, *circular*, and *annotations* patterns, we simulated 1/3 of SVGs and 2/3 of uniform genes; in each one of these simulations, half of the SVGs are highly abundant one region (e.g., centre of the circle, or Layer 5; see Figure 2), and half are highly abundant in the complementary region (e.g., outside the circle, or outside Layer 5; see Figure 2). In *mixture* and *inverted mixture* simulations, we generate 50% of uniform genes and 50% of SVGs, in equal proportions from: 5, 3 and 4 distinct patterns in the *LIBD*, *melanoma*, and *mouse cerebellum* datasets, respectively.

Individual cluster testing

Computing dispersion estimates is the most computationally demanding task in *edgeR*. Since this step depends on the design matrix of the test performed, for each cluster being tested, *edgeR* requires re-computing such dispersion estimates. In order to accelerate calculations, we also provide an alternative implementation, where we use the dispersion estimates previously computed for the gene-level test. This results in a significant decrease in the computational burden (runtime decreased, on average, from 32 to 2 minutes per sample), resulting in analogous performance (Table 1, and Supplementary Table 2).

Software versions

Most analyses were performed in R (version 4.2.0, with Bioconductor packages from release 3.15); some methods (i.e., *StLearn*, *SpatialDE*, *SpatialDE2* and *SpaGCN*) were run in python (version 3.10.0).

References

[1] 10x Genomics Acquires Spatial Transcriptomics. https://www.10xgenomics.com/news/10x-genomics-acquires-spatial-transcriptomics., 2018. [Accessed November 2019].

[2] Y. Benjamini and Y. Hochberg. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal statistical society: series B (Methodological)*, 57(1):289–300, 1995.

[3] N. BinTayyash, S. Georgaka, S. John, S. Ahmed, A. Boukouvalas, J. Hensman, and M. Rattray. Non-parametric modelling of temporal and spatial counts data from RNA-seq experiments. *Bioinformatics*, 37(21):3788–3795, 2021.

[4] C. D. X. Z. Brendan F Miller, Dhananjay Bambah-Mukku and J. Fan. Characterizing spatial gene expression heterogeneity in spatially resolved single-cell transcriptomic data with nonuniform cellular densities. *Genome research*, May 2021.

[5] D. M. Cable, E. Murray, L. S. Zou, A. Goeva, E. Z. Macosko, F. Chen, and R. A. Irizarry. Robust decomposition of cell type mixtures in spatial transcriptomics. *Nature biotechnology*, 40:517–526, 2022.

[6] A. Chen, S. Liao, M. Cheng, K. Ma, L. Wu, Y. Lai, X. Qiu, J. Yang, J. Xu, S. Hao, X. Wang, H. Lu, X. Chen, X. Liu, X. Huang, Z. Li, Y. Hong, Y. Jiang, J. Peng, S. iu, and J. Wang. High-definition spatial transcriptomics for in situ tissue profiling. *Cell*, 185(10):1777–1792.e21, 2022.

[7] Y. Chen, A. T. Lun, and G. K. Smyth. From reads to genes to pathways: differential expression analysis of RNA-Seq experiments using Rsubread and the edgeR quasi-likelihood pipeline. *F1000Research*, 5, 2016.

[8] P. J. Daniel Edsgård and R. Sandberg. Identification of spatial expression trends in single-cell gene expression data. *Nature Methods*, 15:339–342, 2018.

[9] A. T. L. L. Q. F. W. Davis J McCarthy, Kieran R Campbell. Scater: pre-processing, quality control, normalization and visualization of single-cell RNA-seq data in R . *Bioinformatics*, 33(8):1179–1186, 2017.

[10] G. K. S. Davis J McCarthy, Yunshun Chen. Differential expression analysis of multifactor RNA-Seq experiments with respect to biological variation. *Nucleic acids research*, 40(10):4288–97, 2012.

[11] Z. Edward, M. R. Stone, R. Xing, G. Jamie, K. S. Smythe, P. Thomas, S. R. Williams, C. R. Uytingco, S. E. B. Taylor, N. Paul, J. H. Bielas, and G. Raphael. Spatial transcriptomics at subspot resolution with BayesSpace. *Nature Biotechnology*, 2021.

[12] C.-H. L. Eng, M. Lawson, Q. Zhu, R. Dries, N. Koulena, Y. Takei, J. Yun, C. Cronin, C. Karp, G.-C. Yuan, et al. Transcriptome-scale super-resolved imaging in tissues by RNA seqFISH+. *Nature*, 568(7751):235–239, 2019.

[13] T. Z. M. A. Eric Lubeck, Ahmet F Coskun and L. Cai. Single-cell in situ RNA profiling by sequential hybridization. *Nature methods*, 11(4):360–1, 2014.

[14] J. Hu, X. Li, K. Coleman, A. Schroeder, N. Ma, D. J. Irwin, E. B. Lee, R. T. Shinohara, and M. Li. SpaGCN: Integrating gene expression, spatial location and histology to identify spatial domains and spatially variable genes by graph convolutional network. *Nature Methods*, 18:1342 – 1351, Novemble 2021.

[15] P. Jaccard. Distribution de la flore alpine dans le bassin des dranses et dans quelques régions voisines. *Bull Soc Vaudoise Sci Nat*, 37:241–272, 1901.

[16] I. Kats, R. Vento-Tormo, and O. Stegle. SpatialDE2: Fast and localized variance component analysis of spatial transcriptomics . *bioRxiv*, 2021.

[17] J. R. M. S. W. X. Z. Kok Hao Chen, Alistair N Boettiger. RNA imaging. Spatially resolved, highly multiplexed RNA profiling in single cells. *Science (New York, N.Y.)*, 348(6233), 2015.

[18] W. Liu, X. Liao, Z. Luo, Y. Yang, M. C. Lau, Y. Jiao, X. Shi, W. Zhai, H. Ji, J. Yeong, et al. Probabilistic embedding, clustering, and alignment for integrating spatial transcriptomics data with PRECAST. *Nature Communications*, 14(1):296, 2023.

[19] M. I. Love, W. Huber, and S. Anders. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome biology*, 15(12):1–21, 2014.

[20] M. W. Lukas, T. Leonardo, Collado, and C. H. Stephanie. Orchestrating spatially-resolved transcriptomics analysis with bioconductor. https://lmweber.org/OSTA-book/quality-control.html. Accessed: 2022-08-07.

[21] G. K. S. Mark D Robinson, Davis J McCarthy. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics*, 26(1):139–140, 11 2009.

[22] B. Pardo, A. Spangler, L. M. Weber, S. C. Page, S. C. Hicks, A. E. Jaffe, K. Martinowich, K. R. Maynard, and L. Collado-Torres. spatialLIBD: an R/Bioconductor package to visualize spatially-resolved transcriptomics data. *BMC genomics*, 23(1):434, 2022.

[23] D. T. Pham, T. Xiao, X. Jun, L. F. Grice, P. Y. Lam, R. Arti, V. Jana, M. J. Ruitenberg, and Q. H. Nguyen. stLearn: integrating spatial location, tissue morphology and gene expression to find cell types, cell-cell interactions and spatial trajectories within undissociated tissues. *Bioinformatics*, 36(7):2293–2294, 2020.

[24] A. Rao, D. Barkley, G. S. França, and I. Yanai. Exploring tissue architecture using spatial transcriptomics. *Nature*, 596(7871):211–220, 2021.

[25] D. Righelli, L. M. Weber, H. L. Crowell, B. Pardo, L. Collado-Torres, S. Ghazanfar, A. T. Lun, S. C. Hicks, and D. Risso. SpatialExperiment: infrastructure for spatially-resolved transcriptomics data in R using Bioconductor. *Bioinformatics*, 38(11):3128–3131, 2022.

[26] M. E. Ritchie, B. Phipson, D. Wu, Y. Hu, C. W. Law, W. Shi, and G. K. Smyth. limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic acids research*, 43(7):e47–e47, 2015.

[27] M. D. Robinson and A. Oshlack. A scaling normalization method for differential expression analysis of RNA-seq data. *Genome biology*, 11(3):1–9, 2010.

[28] D. Ruben, Z. Qian, D. Rui, C.-H. L. Eng, L. Huipeng, L. Kan, F. Yuntian, Z. Tianxiao, S. Arpan, B. Feng, R. E. George, P. Nico, C. Long, and Y. Guo-Cheng. Giotto: a toolbox for integrative analysis and visualization of spatial expression data. *Genome Biol*, 22:78, 2021.

[29] A. G. C. A. M. E. M. C. R. V. J. W. L. M. C. F. C. . . E. Z. M. Samuel G Rodriques, Robert R Stickels. Slide-seq: A scalable technology for measuring genome-wide expression at high spatial resolution. *Science (New York, N.Y.)*, 363(6434):1463–1467, 2019.

[30] W. Z. L. C. Sheel Shah, Eric Lubeck. In situ transcription profiling of single cells reveals spatial organization of cells in the mouse hippocampus. *Neuron*, 92(2):342–357, 2016.

[31] J. Z. Shiquan Sun and X. Zhou. Statistical analysis of spatial expression patterns for spatially resolved transcriptomic studies. *Nature Methods*, 17:193–200, 2020.

[32] J. Z. Shiquan Sun and X. Zhou. SPARK-X: non-parametric modeling enables scalable and robust detection of spatial expression patterns for large spatial transcriptomic studies. *Genome Biology*, 22:184, 2021.

[33] R. R. Stickels, E. Murray, P. Kumar, J. Li, J. L. Marshall, D. J. Di Bella, P. Arlotta, E. Z. Macosko, and F. Chen. Highly sensitive spatial transcriptomics at near-cellular resolution with Slide-seqV2. *Nature Biotechnology*, 39:313 – 319, March 2021.

[34] P. L. Ståhl, F. Salmén, S. Vickovic, A. Lundmark, J. F. Navarro, J. Magnusson, S. Giacomello, M. Asp, J. O. Westholm, M. Huss, A. Mollbrink, S. Linnarsson, S. Codeluppi, Åke Borg, F. Pontén, P. I. Costea, P. Sahlén, J. Mulder, O. Bergmann, J. Lundeberg, and J. Frisén. Visualization and analysis of gene expression in tissue sections by spatial transcriptomics. *Science*, 353(6294):78–82, 2016.

[35] K. Thrane, H. Eriksson, J. Maaskola, J. Hansson, and J. Lundeberg. Spatially resolved transcriptomics enables dissection of genetic heterogeneity in stage iii cutaneous malignant melanoma. *Cancer Research*, 78(20):5970–5979, 2018.

[36] S. A. T. Valentine Svensson and O. Stegle. SpatialDE: identification of spatially variable genes. *Nature Methods*, 15:343–346, 2018.

[37] S. Vickovic, G. Eraslan, F. Salmén, J. Klughammer, L. Stenbeck, D. Schapiro, T. Äijö, R. onneau, L. Bergenstråhle, J. F. Navarro, J. Gould, G. K. Griffin, å. Borg, M. Ronaghi, J. Frisén, J. Lundeberg, A. Regev, and P. L. Ståhl. High-definition spatial transcriptomics for in situ tissue profiling. *Nature methods*, 16:987–990, 2019.

[38] L. M. Weber, A. Saha, A. Datta, K. D. Hansen, and S. C. Hicks. nnSVG: scalable identification of spatially variable genes using nearest-neighbor Gaussian processes. *bioRxiv*, May 2022.

[39] S. S. Wilks. The large-sample distribution of the likelihood ratio for testing composite hypotheses. *The annals of mathematical statistics*, 9(1):60–62, 1938.

[40] P. Zhao, J. Zhu, Y. Ma, and X. Zhou. Modeling zero inflation is not necessary for spatial transcriptomics. *Genome Biology*, 23(1):118, 2022.