The genomic landscape of adaptation to a new host plant

Rachel A. Steward[1]*, Kalle J. Nilsson[1]*, Jesús Ortega Giménez[1,2]*, Zachary J. Nolen[1], Chao Yan[1], Yajuan Huang[1], Julio Ayala López[1,3] and Anna Runemark[1]

*Equal contributions

Correspondence: rachel.steward@biol.lu.se

[1] Department of Biology, Lund University, Lund, Sweden

[2] Cavanilles Institute of Biodiversity and Evolutionary Biology, Universidad de Valencia, Paterna, Spain

[3] Division of Theoretical Ecology and Evolution, Universität Bern, Bern, Switzerland

Keywords: adaptive divergence, host plant adaptation, inversion, parallelism, genomic selection, introgression

1 Abstract

Adaptation to novel ecological niches is known to be rapid. However, how
the loci underlying ecological divergence are coupled to traits
reproductively isolating populations, ultimately enabling the formation of
persistent species, remains a consequential question in speciation research.
Here, we investigated the genomic differences underpinning colonization of
a new niche and formation of two partly sympatric host races of *Tephritis*
conura peacock flies. We took advantage of two independent sympatric
zones west and east of the Baltic Sea, where host plant specialists using the
thistle species *Cirsium heterophyllum* and *C. oleraceum* co-occur, and
address what regions of the genome maintain the host races in parallel.
Using genome-wide association, differentiation and divergence statistics,
we identified a large, highly divergent region associated with host use
among western and eastern populations. Within this region, we identified
unique haplotypes associated with each host race, indicative of a large
inversion, adding to the growing body of evidence that structural changes to
the genome are important for adaptations to persist in the face of gene flow.
We further showed strong signatures of selection in this region, especially
in populations of the derived *C. oleraceum* specialist host race. The region
also had reduced introgression, especially in western populations, while the
rest of the genome showed signs of extensive gene flow. Genes within
highly differentiated windows within the putative inversion were not only
enriched for functions involved in host adaptation, including phenology and
metabolic responses to different metabolites in the two host plants, but also
enriched for gametogenesis, fertilization and embryological development,
all of which suggest sequence divergence could have large consequences on
reproductive isolation between the host races. In conclusion, this study
suggests that structural changes in the genome may facilitate the formation
of persistent host races, and ultimately speciation, in face of gene flow.

Introduction

How biological variation and novel adaptation arise are fundamental and unresolved questions in our understanding of the origins of biodiversity. Ecological adaptation to novel environments and food sources are known to result in rapid phenotypic and genetic divergence (Bush, 1969; Boag and Grant, 1981; Feder *et al.*, 1988; Filchak *et al.*, 2000; Herrel *et al.*, 2008; Nosil, 2012), but to what extent this ecological adaptation leads to persistent speciation is contested (Schluter, 2009; Anderson and Weir, 2022; Anderson *et al.*, 2023). Both the ecological context, including the consistency (Hendry, 2009; Bolnick, 2011) and multidimensionality (Rice and Hostert 1993; Nosil and Sandoval 2008; Nosil et al. 2009; Chevin et al. 2014)(Rice and Hostert, 1993; Nosil and Sandoval, 2008; Nosil and Harmon, 2009; Chevin *et al.*, 2014) of the divergent niches, and the genomic underpinnings of adaptation to those niches (Chevin *et al.*, 2014) may influence the likelihood of divergent lineages becoming distinct species over time. Whether ecological adaptations and reproductive isolation are linked, at either the phenotypic or genomic scale, is important to the speciation process. Without reproductive isolation, ecological shifts can lead to collapses of formerly isolated populations, resulting in increased gene flow and breakdown of the speciation process (Gow *et al.*, 2006; Taylor *et al.*, 2006; Lackey and Boughman, 2017). Thus, a challenge for evolutionary biologists is to identify the genomic architecture that enables persistent species to form (Kulmuni *et al.*, 2020).

One way to uncover the genomic architecture that enables speciation is to assess sequence differences that are robust to gene flow between populations in sympatry (Dittmar *et al.*, 2016; Ferris *et al.*, 2017). Gene flow can be limited in regions of the genome harboring genes that are involved in reproductive isolation, and differences in these regions are likely contributing to the speciation process (Ellegren *et al.*, 2012; Coughlan and Matute, 2020; Schluter and Rieseberg, 2022). However, interpreting which of the differences between existing species *caused* speciation may be challenging, as additional genes involved in reproductive isolation may have accumulated since the species diverged (Cruickshank and Hahn, 2014; Ellegren and Galtier, 2016). Investigating the genomic differences between recently diverged taxa that are maintained in independent contact zones can help resolve this challenge (Schluter and Rieseberg, 2022). Regions that repeatedly differ among differentially adapted taxa are likely to play a role in speciation (Johannesson *et al.*, 2010; Meier *et al.*, 2018; Rivas *et al.*, 2018; Bohutínská *et al.*, 2021; Koch *et al.*, 2022). Extensive theoretical and empirical evidence has shown that genomic architecture coupling coadapted loci, including inversions, is

particularly important for persistence of differential adaptation in the face
of gene flow (Feder, Roethele, *et al.*, 2003; Kirkpatrick and Barton, 2006;
Wellenreuther and Bernatchez, 2018; Faria, Johannesson, *et al.*, 2019;
Berdan *et al.*, 2022; Schaal *et al.*, 2022; Lucek *et al.*, 2023). Yet, the extent
to which linkage of ecological adaptation and reproductive isolation loci is
facilitated by pleiotropic effects of the same genes, coupling of genes
through structural variation, or co-inheritance of uncoupled genomic
regions upheld by correlational selection remains an empirical challenge for
speciation researchers.

Herbivorous insects and the plants they interact with are excellent models
for studying the genomic basis of ecological adaptation and speciation.
These are two of the most speciose groups of eukaryotes (Mora *et al.*, 2011;
Christenhusz and Byng, 2016), and specialization of phytophagous insects
and their hosts is hypothesized to drive this divergence (Vidal and Murphy,
2018). The specificity and multidimensionality of the niches constituted by
the host plants (Hardy and Otto, 2014) and strong dependency on host
plants for reproduction and survival (e.g. assortative mating based on host
recognition; Bush, 1969) may increase the chance of ecological speciation
in herbivorous insects (Hardy and Otto, 2014). Changes in host repertoire,
either through host range expansions or host shifts, have previously been
associated with chemosensory genes underlying butterfly host choice (van
Schooten *et al.*, 2020), genes involved in oral secretions in aphids
(Nouhaud *et al.*, 2018; Boulain *et al.*, 2019; Shih *et al.*, 2023), genes coding
for enzymes in the digestive systems of both butterflies and aphids (Nallu *et
al.*, 2018; Singh *et al.*, 2020; Shih *et al.*, 2023), and genes underlying
phenological shifts in *Rhagoletis* flies (Feder *et al.*, 1988; Feder, Berlocher,
et al., 2003; Egan *et al.*, 2015; Hood *et al.*, 2020; Inskeep *et al.*, 2022).
However, how the genetic changes underlying ecological adaptation are
coupled to reproductive isolation at the genomic level (Butlin and Smadja,
2018), enabling co-existence and long-term persistence of differentially
adaptive ecotypes or host races (cf. Dres and Mallet, 2002), remains a
challenge to resolve. It is possible the time required for stable existence of
host races is determined by the accumulation of genomic architecture
coupling host adaptation and reproductive isolation loci. This is the case in
birds, where sister taxa with inversions co-exist at shorter divergence times
(Hooper *et al.*, 2019).

Here, we use the Tephritid fly *Tephritis conura* to uncover the genomic
basis of ecological divergence and reproductive isolation promoting stable
coexistence of two differentially adapted host races. These host races infest
the ancestral host thistle *Cirsium heterophyllum* and the derived host thistle
C. oleraceum, respectively (Romstock-Volkl, 1997). They differ in

phenology and host plant preference (Romstock-Volkl, 1997; Diegisser,
Johannesen*, et al.*, 2006), and have host-specific larval performance and
survival (Diegisser *et al.*, 2008) resulting in relatively strong reproductive
isolation (Seitz and Komma, 1984; Romstock-Volkl, 1997; Diegisser,
Johannesen*, et al.*, 2006; Diegisser, Seitz*, et al.*, 2006). Moreover,
mitochondrial networks and allozyme markers suggest they are genetically
differentiated (Diegisser, Johannesen*, et al.*, 2006; Diegisser, Seitz*, et al.*,
2006). Taken together, these findings suggest that there is strong divergent
ecological selection acting on these host races. This study system also
features independent contact zones where the host races are found in
sympatry both west and east of the Baltic Sea (Nilsson *et al.*, 2022),
enabling us to determine the genomic regions that differ between host races
in parallel, and hence are likely to confer ecological adaptation. We show
that a large, contiguous region of the ancestral dipteran X chromosome is
the basis of genomic differentiation between *T. conura* host races. This
region has unique haplotypes for each host race and is highly resistant to
introgression, consistent with an inversion. Not only does the putative
inversion exhibit signatures of positive selection, especially in the derived
host race, but it also couples genes that are enriched for functions critical to
host adaptation, mating and reproduction. Together, these results suggest
that structural variation has enabled colonization of a novel host and
facilitates host race persistence in sympatry.

Methods

Tephritis conura (Loew, 1844; Diptera) is a true fruit fly in the family
Tephritidae. Most species in this family are specialists, eating one or a few
plants within the same family or genus (Aluja and Norrbom, 2001). Larvae
feed on fruits, seeds, flowers, stems or plant roots (Christenson and Foote,
1960; Headrick and Goeden, 1998). Within Tephritidae, several examples
of host races occur in the genera *Eurosta* (Craig *et al.*, 1993), *Rhagoletis*
(Bush, 1969; Feder *et al.*, 1988) and *Tephritis* (Diegisser *et al.*, 2004;
Diegisser, Seitz*, et al.*, 2006). *Tephritis conura* host plant specialization has
been documented in populations in continental Europe, following a host
shift from *C. heterophyllum* (referred to as CH flies from here on) to the
derived host plant *C. oleraceum* (referred to as CO flies; Romstock-Volkl,
1997; Diegisser, Johannesen*, et al.*, 2006). The host races are nearly
indistinguishable morphologically, except for the ovipositor length, which
is shorter for CO females (Romstock-Volkl, 1997; Diegisser *et al.*, 2007;
Nilsson *et al.*, 2022). Sampling adult flies on their host plants or as larvae
in infested buds also facilitates discrimination.

To uncover the genomic regions involved in host plant adaptation in *T.*
conura, we assembled and annotated a reference genome and used whole
genome resequencing of four populations from each host race. For both
host races, *T. conura* larvae inside infested buds were collected during June
and July 2018. We sampled buds from eight populations, four for each host
race, including sites in allopatric populations of *C. oleraceum* in Germany
and Lithuania, allopatric populations of *C. heterophyllum* in northern
Sweden and Finland, and sympatric populations of both host races in
southern Sweden and Estonia (Fig. 1; Table S1). This design enables us to
identify the genomic regions that consistently differ between host races, as
the sympatric regions sampled on each side of the Baltic Sea constitute
independent contact zones. The pupa used for the reference genome was
sampled from a CH individual from the CHST population (Fig. 1D).

Thistle buds containing *T. conura* larvae were stored individually in netted
cups at 21^{o}C in a laboratory at Lund University, Sweden. We sampled adult
flies three days after eclosion. We considered all flies emerging from a
given bud as potential siblings and therefore sampled a single adult male
from each bud to avoid including related individuals. Adult flies and the
pupae were euthanized through snap freezing and stored at -80^{o}C.

Genome assembly and improvement

DNA from a single *T. conura* pupa was extracted and sequenced by
SciLifeLabs, Solna, Sweden following their in-house protocol for PacBio
sequencing. The pupa was from the allopatric CH population from
Grythyttan, Sweden (aka CHST). Briefly, the DNA was sequenced using
PacBio long read technology and sequences were error-corrected using
SMRT Link analysis. Sequencing resulted in 4,173,684 reads with a total of
74.3 Gbp and an N50 read length of 17.8 kb. PacBio sequences were
assembled with hifiasm 0.7-dirty-r255 (Cheng *et al.*, 2021), which was run
using default parameters, followed by purge_dups (v. 1.2.5; Guan *et al.*,
2020) to identify and remove heterozygous duplication (e.g. haplotigs) in
the resulting assemblies.

We used RepeatModeler (v. 2.0.3; Flynn *et al.*, 2020) and RepeatMasker
(v. 4.1.2; Smit *et al.*, 2015) to identify and soft mask repetitive elements in
the *T. conura* reference assembly. For RepeatModeler, we specified the
'rmblast' engine and ran the LTR structural pipeline. We ran the -gccalc
option in RepeatMasker to account for variable GC content. To put the *T.*
conura results in context we accessed repeat summaries for nine of the
Tephritid species with published genomes (Sproul *et al.*, 2023). While

Sproul *et al.* (2023) performed additional repeat screens, we used the output
from their first round of masking, in which they used RepeatModeler2.0
(search engine "ncbi") and RepeatMasker4.1.0 to generate custom repeat
libraries.

Contigs were arranged into chromosome-level scaffolds using HiC
sequencing data from a second *T. conura* individual. Following the Arima
HiC pipeline (v. 02; https://github.com/ArimaGenomics/mapping_pipeline;
accessed 06/2023), HiC reads were mapped to the hifiasm assembly with
bwa-mem (v.7.17-r1188; Li and Durbin, 2009), paired and filtered (mapQ >
10) using python scripts from the Arima pipeline, and duplicates were
marked and removed using Picard tools MarkDuplicates (v2.10.3; Picard
toolkit, 2019). The mapped reads were then used to scaffold existing
contigs using YaHS (Zhou *et al.*, 2022), specifying a minimum contig
length of 50 kb and requiring the use of existing contigs (i.e., hifi-asm
contigs were not broken by YAHS while scaffolding). The resulting contact
map was assessed visually using Juicebox
(https://github.com/aidenlab/juiceboxgui; Fig. S1A). Due to the low
coverage of our HiC data, we performed downstream analyses at the contig
level but used the scaffolding output to cluster contigs within hypothetical
linkage groups. We assessed the HiC scaffolds by comparing synteny with
the published genome of a close relative, *Rhagoletis pomonella*. We
hardmasked repeats and aligned the scaffolded assembly to the *R.*
pomonella assembly (GCF_013731165.1_Rhpom_1.0; accessed 11/2022;
length = 1223 Mbp, scaffolds = 32,060, scaffold N50 = 72 Mbp) using
minimap2 (asm20; Li, 2021) and visualized synteny using circlize (v.
0.4.15; Gu *et al.*, 2014). We used a similar approach to compare *T. conura*
scaffolds against a *Drosophila melanogaster* assembly (dmel-all-
chromosome-r6.48, accessed from FlyBase 01/2023;
https://ftp.flybase.org/releases/FB2022_05).

The reference assembly was scanned with Kraken2 (Wood *et al.*, 2019) to
identify bacterial, viral and fungal sequences. Contigs with a contamination
proportion of greater than 10% were removed (150 contigs; 47,038,732 bp),
a threshold that was chosen based on a large gap that occurred between 2
and 10%. The ancestral dipteran X chromosome was identified by finding
contigs with an excess of hits (at least one hit per million bp, approximately
top 5% of contigs) when blasted (tblastn; v.2.11.0; Camacho *et al.*, 2009)
with a set of *Drosophila melanogaster* X-linked proteins, as the the *D.*
*melanogaste*r X chromosome is homologous to the ancestral dipteran X
(Vicoso and Bachtrog, 2013, 2015). This resulted in 102 putative X-
enriched contigs, 80.3% of which were clustered on a single HiC scaffold:
scaffold_3. We used a similar approach to identify scaffold_7 as

homologous with *D. melanogaster* chromosome 4 (aka Muller element F),
which was previously identified as the Z chromosome in congener *T.*
californica (Vicoso and Bachtrog, 2015).

Gene prediction and annotation

We annotated the assembly using the BRAKER3 pipeline (Gabriel *et al.*,
2024). We generated expressed sequence predictions using RNAseq reads
from flash-frozen flies from various life stages, sexes and populations (see
Steward *et al.*, 2024 for details about extraction and sequencing). Reads
were mapped to the genome using the splice-aware aligner HISAT2 (Kim
et al., 2019). Protein hints were generated from the Arthropoda ODB_11
database (Kuznetsov *et al.*, 2022). We used the default BRAKER3 pipeline,
but a large proportion of single exon genes motivated us to re-merge the
protein-based and RNA-based annotations with TSEBRA, reducing the
support necessary for introns from 1 to 0.25. The result was a highly
complete annotation (BUSCO v. 5.3.1; Manni *et al.*, 2021; 98.0% complete
(S: 73.6%, D: 24.4%), 0.8% fragmented, 1.2% missing, diptera_ODB10, n
= 3285), with a total of 25,175 genes and 30,632 transcripts. We
functionally annotated the longest isoform of the predicted genes with
eggNOG emmaper (v. 2.1.5; Huerta-Cepas *et al.*, 2018) using default
settings. Functional annotations were returned for 17,123 genes, including
gene ontology (GO) terms for 10438 of these.

DNA extraction and sequencing

We extracted DNA for whole genome sequencing from entire flies using
Qiagen DNeasy Blood and Tissue Kit (Qiagen Corp., Valencia, CA),
following the standard protocol with some small modifications
(Supplementary Methods 1). Frozen flies were briefly thawed and ground
with plastic pestles in the tissue lysis buffer. Samples were incubated with
proteinase K (3.5h, 56°C), and then RNAse was added and the pellet eluted
using a 56μl of EB buffer. Library preparation and whole genome
resequencing were performed by SciLifeLab (NGI-Sweden, Solna,
Sweden) using Illumina TruSeq PCR-free prep kit with an insert size of 350
bp. Sequencing of 2x150 bp paired-end reads was carried out on 8 Illumina
HiSeqX lanes.

Mapping, variant calling and allele frequencies

We trimmed raw whole genome sequencing data using fastp (v. 0.23.2; Chen *et al.*, 2018), including removing poly-G tails common on patterned flow cells (-g). We aligned the trimmed reads to the reference genome using bwa mem (v. 0.7.17; Li, 2013), then merged parallel sequencing runs per sample, sorting and indexing the alignments using samtools (v. 1.10; Danecek *et al.*, 2021). PCR duplicates were identified and removed using MarkDuplicates in Picard tools (v. 2.23.4; Picard toolkit, 2019). Genome-wide mean coverage was calculated with samtools (Fig. S2). Raw sequence data processing and mapping, along with several latter genomic analyses were performed using a custom Snakemake (Mölder *et al.*, 2021).

To account for low coverage, we utilized genotype likelihood-based methods for analyses when possible. To accommodate both genotype likelihood and genotype call-based analyses, we produced both (1) a Beagle file containing genotype likelihoods for the 96 *T. conura* individuals produced using ANGSD (v. 0.940; Korneliussen *et al.*, 2014) and (2) a VCF containing genotype calls for the 96 individuals as well as three outgroup individuals produced using bcftools (Danecek *et al.*, 2021). The filters and steps to produce these two files are described in Supplementary Methods 2. The final Beagle file and VCF contained 10,421,531 and 32,991,264 SNPs, respectively.

Per population, we generated site allele frequency indices (SAF) with ANGSD using the *T. conura* reference to polarize the ancestral state and relying on the same filters as described for the Beagle file, excluding those used for SNP calling (Supplementary Methods 2). We produced folded 1D and 2D site frequency spectra (SFS) for each population and population pair using the SAF files in realSFS (Nielsen *et al.*, 2012).

Population clustering

To explore and visualize genetic clustering of the individuals, we used a principal component analysis performed in PCAngsd (v. 0.982; Skotte *et al.*, 2013) and then formally assessed clustering using NGSAdmix (v. 33; Skotte *et al.*, 2013) for 1-10 clusters (K). To prepare input files for these tools, we estimated pairwise linkage disequilibrium between all SNPs within 100 kb of each other using ngsLD (Fox *et al.*, 2019). We then used the included prune_ngsLD.py script to produce a list of linkage pruned SNPs, excluding edges from the initial graph between sites with a distance greater than 50 kb and with an r^2 of less than 0.1 (--max_dist 50000 --min_weight 0.1). We then subset our Beagle file to these linkage pruned

sites and used it as input for both PCAngsd and NGSAdmix. We also
performed PCAs using PCAngsd for each contig, and used PC1 to evaluate
patterns of genetic clustering across the genome. Contigs with fewer than
500 sites in the pruned Beagle file were excluded. For the clustering
analysis, we performed between 10-100 replicate optimizations for each K
with a maximum of 4000 iterations per replicate. Replication was halted
when the replicates converged or 100 replicates had completed, considering
replicates as converging when the three highest likelihood replicates came
within 2 log-likelihood units of each other, as in Pečnerová et al. (2021).
We selected the best K as the highest level of K achieving convergence.

Coalescent-based phylogenetic analyses

We reconstructed the phylogenetic relationship between samples using a
coalescent-based approach as implemented in ASTRAL-MP (Yin *et al.*,
2019), which takes into account the discordance between gene trees and the
true species tree expected in recently diverged taxa. For all *T. conura*
samples and the three outgroup samples we produced consensus FASTA
sequences for all 2,419 complete single-copy Dipteran BUSCO genes
present in our assembly using ANGSD, calling the consensus base with the
highest effective sequencing depth (-doFasta 3), excluding positions with
an individual sequencing depth less than 3 or greater than 100. For each
gene, we inferred the maximum likelihood gene tree using IQ-TREE2 (v.
2.2.6; Minh *et al.*, 2020), testing and selecting the best fitting substitution
model and performing 1000 ultrafast bootstrap replicates to assess node
confidence. We collapsed nodes with bootstrap support of less than 10%
into polytomies using Newick Utilities (v. 1.6; Junier and Zdobnov, 2010)
as collapsing low confidence nodes in gene trees can improve species tree
estimation accuracy (Zhang *et al.*, 2018). Finally, we inferred the
coalescent phylogenetic relationships from BUSCO gene trees with
sequencing data for all samples (2,418/2,419 genes) using ASTRAL-MP v.
5.15.5.

Demographic history

Demographic history was estimated for each population using MSMC2 (v.
2.1.2; Schiffels and Durbin, 2014). We generated the individual VCF and
mask bed file by using bcftools (v. 1.14; Li, 2011) and bamCaller.py script
from msmc-tools (https://github.com/stschiff/msmc-tools; accessed
09/2023). Sites with extremely low or high coverage, defined as 0.5x and
2x mean coverage respectively, were masked before using Samtools (v.

1.16) to compute the average sample coverage and generate a BED file per-
sample. We merged the VCF and mask files (msmc-tools,
generate_multihetsep.py script) and used the resulting file as input for
MSMC2. To scale the result, we used a mutation rate of 3.46×10^{-9}
mutations per site per generation, which is derived from estimates in *D.*
melanogaster (Keightley *et al.*, 2009) and a mean generational interval of
one year (Romstock-Volkl, 1997). Because this software is sensitive to read
coverage, neutrality, and repetitive content, we compared the MSMC2
output for three sets of samples (the full set of samples, the individuals with
highest coverage in each of the eight populations, and the individuals with
coverage closest to the global mean in each of the eight populations) at
three levels of VCF filtering (the VCF output from bcftools + bamCaller.py
script described above; the VCF excluding all repetitive content, and the
VCF excluding all outlier windows identified by population branch
structure analysis described below).

Genome-wide association analysis with BayPass

Genome-wide scans for association with ecotype were performed with
BayPass (v. 2.4; Gautier 2015). We converted the filtered VCF to allele
frequencies for each population with bcftools (v. 1.19; Danecek *et al.*,
2021) and removed invariant sites using an in-house script. The dataset was
further filtered to account for linkage by subsampling every 20th variant
across each contig into 20 unique sets of SNPs. We analyzed each set of
SNPs first under the BayPass core model using the default options, then
under the auxiliary model using a Markov Chain Monte Carlo (MCMC)
algorithm and host race as a covariate. Three independent runs (using the
option -seed) were performed for each of the 20 SNP sets. Support for host
race association was evaluated using the mean Bayes Factor of these three
runs. Results from the 20 independent runs were combined into a single
dataset for all sites, and reported in deciban units, with 10db corresponding
to 10:1 odds, 20db to 100:1 odds, etc. The mean Bayes Factor was further
summarized across 50kb windows. Because several windows were highly
repetitive, we removed windows containing 500 or fewer SNPs (1% of the
window size) from downstream analyses and visualizations.

Population differentiation and divergence

To measure population differentiation, F_{ST} was calculated for each site, first
by generating the site frequency spectrum (2D-SFS) for each population
pair, based on which we calculated per-site F_{ST} using a Bhatia estimator

(Bhatia *et al.*, 2013) as implemented in ANGSD (realSFS F_{ST}). Average
pairwise F_{ST} was estimated genome-wide and for nonoverlapping 50kb
windows (realSFS stats2). We used population branch statistics (PBS) to
assess differentiation between sympatric populations of the two host races
when corrected for divergence between populations within host races. This
accounts for heterogeneity in background differentiation between
populations, as differentiation within host races should reflect genetic drift
and the recombination rate landscape whereas differentiation between host
races should also reflect changes in nucleotide diversity resulting from
selection and introgression. We calculated PBS for each site (realSFS
stats2; Fig. S3) for four population triads composed of a focal sympatric
population, and the sympatric and allopatric populations of the other host
race in the same transect.

To measure population divergence, we first re-calculated the 2D-SFS for
every 50kb window, then calculated absolute nucleotide divergence (dxy)
using a modified script from
(https://github.com/marqueda/PopGenCode/blob/master/dxy_wsfs.py;
accessed 11/2023 and modified to run in R). We then calculated relative
node depth (RND; Fig S3), which quantifies genetic divergence between
two populations (here, two sympatric populations) relative to a third
population (usually an outgroup, but here an allopatric population). RND is
therefore analogous to PBS, as it controls for divergence within host races
while testing for divergence between host races. Because several windows
were highly repetitive, we removed PBS and RND windows with less than
20% coverage from downstream analyses.

Outlier analysis and gene ontology term enrichment

To identify genomic regions associated with host use we identified outliers
for three different metrics: BayPass (estimated for all eight populations),
PBS, and RND (both assessed for a focal sympatric population of each host
race in the western and eastern transects, for a total four combinations).
Windows that exceeded the genome-wide mean by three standard
deviations in at least two of these metrics were identified as outliers. We
identified outliers for the CH and CO host races in the western and eastern
transects separately, and defined highly differentiated windows as those
that were shared between comparisons in the western and eastern transects.

We performed gene set enrichment analysis for gene ontology (GO) terms
using the package TopGO (Alexa and Rahnenfuhrer, 2022), only
considering terms that were assigned to a minimum of five genes

(minimum node size = 5). We tested for enrichment of biological processes,
molecular functions and cellular components. Significant enrichment was
tested with one-sided Fisher's exact tests corrected with the parent-child
algorithm (Grossmann *et al.*, 2007). GO terms were considered
significantly enriched with a p-value < 0.05. We further generated a
STRING database from the *T. conura* genome annotation
(STRG0A87NSB; https://string-db.org/; updated 12/2023) and identified
networks within our highly differentiated genes genes using Markov
clustering implemented in STRING (inflation = 3). We report the 8 largest
networks and associated GO terms, KEGG pathways and enriched
reactomes.

Nucleotide diversity and divergence from neutrality

We evaluated nucleotide diversity (π) and divergence from neutrality, as
characterized by Tajima's D, to understand genome-wide and local
differences among populations. ANGSD was used to calculate pairwise
nucleotide diversity for each site (thetaD, realSFS saf2theta). These were
used to estimate Tajima's D and pairwise differences across the entire
genome and over 50kb non-overlapping windows (thetaStat do_stat). To
calculate π, pairwise differences were divided by the total number of sites
used in each window. For both metrics, we excluded windows with less
than 20% coverage from downstream analyses.

Selection statistics

To identify SNPs associated with host races and different biogeographic
scenarios, we used SelScan (v. 1.3; Szpiech and Hernandez, 2014) to
calculate the cross-population extended haplotype homozygosity (XP-
EHH). To prepare the data for XP-EHH analysis, SNPs unpruned for
linkage disequilibrium were phased with SHAPEIT2 v 2.r837 (Delaneau *et*
al., 2013) using default MCMC parameters (7 burn☐in MCMC iterations, 8
pruning iterations, and 20 main iterations), conditioning states for
haplotype estimation (K = 100), and window size set at 0.5 Mb as
recommended for whole genome sequence data. We also used an effective
population size (Ne) of 494, based on estimates calculated with SNeP (v.
1.1; Barbato *et al.*, 2015). Here, we used the mean across populations, as all
estimates were very similar. XP-EHH was calculated in windows of size
100 kb in each direction from core SNPs, allowing decay curves to extend
up to 1 Mb from the core, and SNPs with MAF < 0.05 were excluded from
consideration as a core SNP. As we lacked a fine☐scale genetic map for *T.*

conura, we assumed a constant recombination rate of 1.73 cM/Mb as found
for the related species *Bactrocera cucurbitae*. Scores were then normalized
within contigs with the norm version of Selscan v. 1.3.0.

The XP-EHH method identifies genomic regions that underwent a selective
sweep in one population but remained variable in the second population.
We calculated XP-EHH for six population pairs to test for differences in
selection between host races in sympatry and allopatry, and to test for
differences between western and eastern transects. For each pair, we
identified putative locality□specific sweeps by selecting those SNPs in the
top 1% of the genome-wide distribution. Genes under selection were
detected by comparing the SNPs detected by the XP-EHH test against a
gene annotation for *T. conura* with bedtools window 2 kb upstream and
downstream of these genes (v. 2.29.2; Quinlan and Hall, 2010). We also
used a sliding window approach, estimating the average normalized XP-
EHH score across all SNPs in 50 kb nonoverlapping windows. More
negative values indicate a stronger signature of positive selection in one
population in a pair while more positive values indicate a stronger signature
of positive selection in the other population.

Introgression statistics

We investigated evidence of introgression between populations by
estimating genome-wide Patterson's D (Green *et al.*, 2010; Durand *et al.*,
2011) and f_4-ratios (Patterson *et al.*, 2012) for all possible population trios,
as implemented in Dsuite's DTrios function (v0.5-r52; Malinsky *et al.*,
2021). To assess variation in introgression across the genome in sympatric
populations, we investigated f_{dM} in windows across the genome for eight
focal trios, as implemented in the script fourPopWindows.py
(https://github.com/simonhmartin/genomics_general; accessed 02/2024).
We selected this statistic as it is designed to detect introgression in
localized regions of the genome and is suitable for detecting introgression
between both P1 and P3, as well as P2 and P3 (Malinsky *et al.*, 2015).
Briefly, for each of the four sympatric populations as P2, we estimated f_{dM}
in non-overlapping 50 kb windows, assigning the closest allopatric
population of the same host plant to P1, with P3 being either the closest
allopatric or sympatric population of the alternate host plant. For all
analyses we utilized a *Tephritis hyoscyami* individual (sample ID
P18705_127) as the outgroup.

Results

Our final assembly of the *T. conura* genome was 1.99 Gbp in 2,713 contigs
(N50 = 1.75 Mbp). HiC scaffolding with YaHS arranged these contigs into
seven large (>100Mbp) linkage groups comprising over 1.77 Gbp of the
sequence genome (N50 = 323 Mbp). HiC scaffolds were highly syntenic
with *Rhagoletis pomonella* linkage groups (Fig. S1), and linkage group
(LG3) was syntenic with the *Drosophila melanogaster* X chromosome,
which is homologous to the ancestral dipteran X (Vicoso and Bachtrog,
2013, 2015). Benchmarking with Dipteran single copy orthologs found
97.0% completeness with low duplication (2.4%), making it not only the
longest but among the most complete Tephritidae genomes available to date
(Fig. S4). It was also the most repeat rich genome by more than 20%,
driven by an expansion of long terminal repeats that make up over 22% of
the genomic content (Fig. S5; Table S2). This history of genomic expansion
and high repetitive content, which increase the opportunity for both
structural change and chromatin state differences, suggests that *T. conura*
has the potential for extensive and rapid genomic divergence driven by
structural variants.

Genomically divergent host races

Using eight continental *T. conura* populations sampled from regions of host
plant allopatry and sympatry east and west of the Baltic Sea (Fig. 1D), we
found considerable evidence that the *C. heterophyllum* (CH) and *C.*
oleraceum (CO) host races represent distinct lineages. The two host races
separated under the best supported model in an Admixture clustering
analysis (K = 3; Fig. 1E; see Fig. S6A for remaining K). In a genome-wide
principal component (PC) analysis, individuals are generally separated by
host race on the first PC axis, explaining 2.13% of the variance (Fig. 1F;
see Fig. S6B for PC2 and PC3). The samples also split into two distinct
branches in a consensus tree constructed using ASTRAL from 3285 single
copy ortholog gene trees (posterior probability = 1, Fig. 1G, see Fig. S7 for
scaled branch lengths and posterior probabilities for all nodes). These
results were supported by pairwise genome-wide F_{ST} and dxy comparisons,
which showed the largest differentiation and divergence occurred between
populations of different host races (Fig. S8, S9). The western sympatric CH
population (CHSK) appears to be the most distinct lineage, separating from
both CO individuals and the remaining CH individuals along PC1, and
forming a separate cluster under the best supported model in the clustering
analysis (K = 3). This population also formed a distinct cluster in the

ASTRAL consensus tree constructed from 3285 single copy ortholog gene
trees (Fig. 1G).

Despite clear evidence of divergence between the two host races, we also
found signs of recent introgression or incomplete lineage sorting among the
sampled flies, especially in eastern populations. In the consensus tree, there
were five individuals (3 CO, 2 CH) that were not assigned to the same
major branch as the rest of their host race (Fig 1G). Combined with the
clustering analysis, which showed considerable admixture in some
individuals (Fig. 1E), and the PCA, in which several individuals of both
host races were highly divergent on the second PC axis (1.47% of the
variance; Fig 1F), these results suggest some level of admixture in these
individuals.

Although it was previously hypothesized that the derived host race arose
just before the onset of the last glacial maximum c. 18,000 years ago
(Diegisser, Seitz, *et al.*, 2006), an MSMC2 analysis suggests host races
diverged much earlier, between 0.5 - 1 million years ago (Fig. 1H). These
results broadly confirm findings from PCA and admixture analyses. In
particular, the highly divergent CHSK population again appears as a
distinct lineage, perhaps having diverged from the other CH populations
much earlier than the estimated divergence between host races. However,
similar patterns can arise when populations have been affected by recent
bottlenecks, and we are as yet unable to disentangle these alternative
explanations.

Genomic regions underlying host race formation

To identify genomic regions associated with host race divergence, we tested
for allele frequency associations with host race across all eight populations
using BayPass. We further evaluated genomic differentiation and genomic
divergence among population triads east and west of the Baltic Sea using
population branch statistics (PBS) and relative node depth (RND). PBS
leverages pairwise F_{ST} comparisons among three populations (triads) to
assess differentiation and signatures of selection, while RND uses dxy in an
analogous framework to PBS to compare divergence and introgression (Fig.
S3, S10-S12).

We found strong evidence for a large, contiguous region of linkage group 3
(LG3), spanning 104 Mb, 89 contigs and containing 1269 genes, being
highly genomically divergent between the host races (Fig. 2). This region
was consistently supported in tests for whole-genome association with host
race across all populations, as well as PBS and RND comparisons in all

triads (Fig. 2A-C, S13). In each case, we identified outlier windows for
each statistic as those falling more than 3 standard deviations outside the
mean, and considered outlier windows with support from at least two
metrics as highly differentiated between the two host races for CH and CO
triads (west: Fig. 2D, east: Fig. 2E). Overall, highly differentiated windows
were very consistent among host race comparisons and transects, with 214
windows shared across all four comparisons out of 289 total windows (Fig.
2F). All of these 214 windows were located within the contiguous region
on LG3.

This dense accumulation of highly differentiated windows on LG3 suggests
a large region with reduced recombination, possibly under strong selection
in one or both host races, as might be expected for an adaptive inversion.
To evaluate this hypothesis, we used PCAngsd to generate local PCAs for
each contig in our assembly (Fig. 3A). Within the highly differentiated
region, we found two distinct clusters on PC1 for each of the host races
(Fig. 3C), consistent with two inversion haplotypes. There was
considerably higher variance among CH individuals on PC1, but it is
unclear whether this is due to recent introgression into the CH haplotype or
ancestral variation among CH populations predating the formation of the
inversion and divergence of the CO host race. Interestingly, for contigs
outside of the putative inversion, there was little evidence of population
structure (Fig. 3B, D), suggesting that gene flow is high throughout the rest
of the genome. The local PCA revealed several individuals (4 CO, 3 CH
from all except the western sympatric populations CHSK and COSK) with
intermediate PC1 values throughout the entire putative inversion, which
may indicate that these are hybrid individuals. However, due to the high
variance among CH individuals, it is not clear whether these individuals are
truly heterozygous for the CH/CO haplotypes.

Strong selection on the derived host race (CO) within the inversion

To understand the selective pressures acting on the putative inversion, we
tested for evidence for selective sweeps, and whether these are stronger in
the derived CO host race. To do this, we assessed Tajima's D, nucleotide
diversity (π) and extended haplotype homozygosity (XP-EHH; Fig. 4).
When considering all windows across the genome, π within highly
differentiated windows was lower than in non-differentiated windows for
all populations (Fig. 4A-B, S14; Table S3). This effect was consistently
larger in CO compared to CH populations (Table S3), suggesting selection
has disproportionately decreased nucleotide diversity in the putative
inversion in the CO host race. Tajima's D was similar among populations

(Fig. 4C- D, S15; Table S4). The exception was CHSK, which had a much
higher Tajima's D, adding to the evidence from Admixture and MSMC
analyses suggesting that this population has experienced a recent
bottleneck. While Tajima's D tended to be lower in highly differentiated
windows compared to non-differentiated windows, especially in allopatric
populations, Tajima's D was higher in highly differentiated windows in two
sympatric populations (CHSK and COSK).

Extended haplotype homozygosity statistics revealed considerably stronger
positive selection in CO populations when contrasted with CH populations
(Fig. 4E-F, S16). In all comparisons between CH and CO population pairs,
the mean normalized XP-EHH was significantly more positive in highly
differentiated windows compared to non-differentiated regions, indicative
of stronger positive selection in CO populations (Fig. 4F, Table S5). We
found that the difference in XP-EHH was similar in all comparisons
between host races (Fig. 4F, Table S5), despite our expectation that positive
selection for host adaptation or reproductive isolation should be stronger in
sympatric than in allopatric populations (Kautt *et al.*, 2020). Meanwhile, in
comparisons within host races between contact zones, there was evidence
of stronger selection in sympatric populations in the western transect
(CHSK and COSK) compared to the eastern transect, although the effect
was much smaller than between host race comparisons (Fig. 4F, Table S5).

We further identified significantly selected SNPs in each population for
each pairwise contrast using the top 1% of SNPs. Between 42-48% of these
SNPs overlapped genes, including the 2,000 bp up and downstream of the
coding regions. We found more genes overlapping selected SNPs in the CH
host race (Table S6), as SNPs under selection in CO populations were more
densely clustered in a smaller number of annotated genes compared to CH
populations (Table S6). Furthermore, these selected genes in CO
populations were more likely than those in CH populations to overlap with
genes in highly differentiated windows or with genes in the putative
inversion (Fig. S17), adding evidence that selected SNPs in CO populations
fall in a more concentrated, more differentiated set of genes.

The putative inversion is a region of reduced recombination

Despite extensive evidence for prezygotic reproductive isolation between
the *T. conura* host races (Nilsson *et al. in prep*), we detected introgression
between the host races in both the eastern and western sympatric regions.
D-statistics suggest higher levels of introgression between host races in the

eastern than the western region, consistent with this being an older contact
zone for these host races (Table S7).

Introgression was not evenly distributed across the genome (Fig. 5A), as
illustrated by variation in f_{dM} in 50 kb windows. Values of f_{dM} are expected
to be positive when introgression occurred between P3 and P2, and
negative if it occurred between P3 and P1. f_{dM} values in highly
differentiated windows had a much narrower spread around the median
than did undifferentiated windows, suggesting these regions are more
resistant to introgression (Fig. 5B, S18; Table S8). Interestingly,
introgression in highly differentiated regions differed between western and
eastern transects (Fig. 5B, Table S9). In the west, f_{dM} in highly
differentiated regions was consistently lower than undifferentiated regions,
suggesting more introgression between allopatric populations of different
host races, rather than between sympatric populations of the different host
races. Negative values are consistent with greater introgression between the
more geographically distant populations, which may reflect historical gene
flow. These patterns suggest outlier regions have increased resistance to
introgression between sympatric populations. The pattern was opposite in
the east, with greater introgression between sympatric populations than
allopatric populations within the highly differentiated regions.

Highly differentiated windows enriched for host adaptation and
reproductive functions

To understand the possible functional consequences of the putative
inversion, we performed gene set enrichment analyses of genes overlapping
highly differentiated windows (278 genes; Fig. 6A-C; Table S10). Genes
within the putative inversion were enriched for numerous functions that
may be directly or indirectly related to adaptation to different host plants.
Genes involved in both renal development (including malpighian tubules in
flies; Bunt *et al.*, 2010; Fig. 6A; table S10) and gut development were
enriched. Both of these tissues are involved in processing and excreting
toxic substances and waste (Gautam *et al.*, 2017; Colombani and Andersen,
2020). Notably, response to toxic substances was also enriched, and was
associated with glutathione S-transferase and carboxylesterase genes within
the highly differentiated region. Both gene families are known to be
involved in generalized detoxification of plant metabolites (Low *et al.*,
2007; Yu *et al.*, 2009; Cruse *et al.*, 2023).

The two host plants differ significantly in their budding phenology, and
accordingly CH and CO have divergent flight times, even within sympatric

host ranges (Romstock-Volkl, 1997). Numerous terms involved in circadian
rhythmic processes and diapause were enriched among highly differentiated
genes (Table S10). In a network analysis of highly differentiated genes, the
second largest network was associated with both phototransduction and
circadian rhythm processes and pathways (Fig. 4D). While it has not been
thoroughly tested, it appears adult diapause termination is mediated by
temperature cues rather than photoperiod (Diegisser, 2005). Accordingly,
we also found that thermosensory behavior was significantly enriched in
highly differentiated genes.

While both ability to eat the alternate host plant and differing phenologies
serve as major reproductive barriers between the host races (Diegisser *et*
al., 2008), we also found highly differentiated genes enriched for functions
involved in potential intrinsic barriers. We recovered enrichment of
biological process terms involved in oogenesis, spermatogenesis,
fertilization, embryological development, embryological growth, and
mitosis (Table S10). Other processes broadly important to biological
development were also highly enriched, including cell signaling and intra-
and intercellular transport. Regulation, transcription, post transcriptional
modification, and degradation of RNAs, including mRNAs and ncRNAs,
were also enriched, suggesting large divergence in coding and noncoding
gene expression between the two host races. Several of the eight largest
networks recovered in the network analysis of highly differentiated genes
were associated with transcriptional regulation and post-transcriptional
modifications of RNAs (Fig. 6D). Together, these enriched processes and
networks suggest that the highly differentiated region contains many genes
that could be critical to reproductive isolation, either by limiting
gametogenesis, fertilization and embryogenesis, or by disrupting critical
biological processes like cell-signaling and gene expression.

We also found that genes encoding mating behavior and especially male
courtship behavior were highly enriched (Fig. 6A), especially conditioned
taste aversion, which was annotated to, among other genes, the bitter
gustatory receptor *Gr33a*. This receptor has previously been found to be
sensitive to pheromones in *Drosophila melanogaster*, and is also sensitive
to a range of antifeedants (Moon *et al.*, 2009) and may mediate either
identification of host plants, where peacock fly courtship occurs, and/or
courtship behavior itself. Calcineurin-mediated signaling and calmodulin
binding were the main functions of a network of six highly differentiated
genes (Fig. 6D). Calmodulin is involved in *D. melanogaster* courtship (Sato
et al., 2019), possibly mediated by its role in odorant binding receptor
sensitization (Jain *et al.*, 2021)

Discussion

While the genetic basis underlying ecological adaptation is increasingly
well understood (Bomblies and Peichel, 2022), it remains unclear how the
genomic regions associated with differential adaptation confer or are
coupled to reproductive isolation, enabling the formation of persistent
species (Butlin and Smadja, 2018; Kulmuni *et al.*, 2020). We document a
large inversion that differentiates the ecologically differentially adapted
host races of *T. conura*, reducing gene flow and enabling persistent
adaptation. The inversion is differentiating the host races in parallel in two
independent regions of secondary sympatric existence, as expected under
parallel selection (Johannesson, 2001; Stuart *et al.*, 2017), supporting its
role in ecological adaptation.

Inversions have been increasingly appreciated as key to enabling co-
existence of ecologically divergent lineages in sympatry (Hooper and Price,
2017; Faria, Chaube*, et al.*, 2019). And yet, how the inversions enabling the
formation of stable morphs, divergence or speciation originate remains a
consequential question (Faria, Johannesson*, et al.*, 2019; Lucek *et al.*,
2023). We document recent and extensive expansions of long terminal
repeats, constituting over 20% of the genome in *T. conura*, rendering
genome size in *T. conura* 55% bigger than that of the closely related
species *Rhagoletis pomonella*. While this expansion and its effects on
genomic architecture is biologically interesting, analyzing the genomic
content of the repetitive regions is intractable, and they have therefore been
excluded from the population genomic analyses. Importantly, this repeat
expansion has likely facilitated a number of structural changes, including
the inversion enabling host race divergence, and producing a more
contiguous assembly required for determining the exact nature of the break
points and thus origin of the inversion is an exciting venue for future
research.

The inversion spans 89 contigs (104 Mb) scaffolded to LG3, covering 33%
of the linkage group, and shows consistent and high divergence between the
host races. This region of LG3 is syntenic with contiguous regions in the
genome of *R. pomonella*. Individuals sampled from the different host races
consistently have native host plant specific inversion haplotypes, with only
7 out of 96 individuals potentially having heterozygous inversion
haplotype, and none being homozygous for the alternative inversion
haplotype. Jointly, the discrete nature of the divergent genomic region and
its strong association with host race suggest that this inversion is key for
enabling the formation of two host races of *T. conura*. Recent research has
shown that inversions are important for maintaining both discrete morphs

within species (Jones *et al.*, 2012; Küpper *et al.*, 2016; Tuttle *et al.*, 2016;
Faria, Chaube*, et al.*, 2019), facilitate speciation (Wellenreuther and
Bernatchez, 2018; Faria, Johannesson*, et al.*, 2019), and enable coexistence
with sister taxa after a shorter period of time (Hooper and Price, 2017).

Our findings suggest that the inversion may be important for coupling (cf.
Butlin and Smadja, 2018) ecologically co-adapted traits in *T. conura*
through physically linking genes involved in ecological adaptation and
reproductive isolation. Functional enrichment analyses suggest that this
inversion has linked genes important to multiple ecological traits associated
with divergent host plant use, including phenology (Romstock-Volkl,
1997), host preference (Diegisser *et al.*, 2008), or physiology including
digestion of plant defense chemicals (Diegisser *et al.*, 2008) and thus
facilitate coexistence between the host races. For instance, genes encoding
response to toxic substances and renal- and gut development are enriched,
which is expected as the host races suffer very strong extrinsic inviability as
larvae when feeding on the wrong host plants (Diegisser *et al.*, 2008).
Moreover, genes involved in male courtship behavior and conditioned taste
aversion, potentially key for host plant recognition and instrumental for
mating to take place on the native host plant as observed in *T. conura*
(Romstock-Volkl, 1997) are enriched in the inversion. Genes involved in
diapause, relevant to the divergent phenologies of the host races
(Romstock-Volkl, 1997) were also enriched. The combination of
enrichment of genes involved in performance on the host plants, preference
for the host plant and phenology which differs between host plants
(Romstock-Volkl, 1997; Diegisser *et al.*, 2008) adds to evidence that the
inversion has coupled (Butlin and Smadja, 2018; Kulmuni *et al.*, 2020;
Schluter and Rieseberg, 2022) genes involved in ecological adaptation and
those contributing to reproductive isolation.

Another set of functions that were enriched in the inverted outlier region
included regulation, transcription, post-transcriptional modification and
degradation of mRNA and ncRNA. Further studies assessing the extent to
which gene expression is differentiated between the host races are
necessary to assess the importance of altered regulation of gene expression
in host plant adaptation. Furthermore, while we had a prior expectation of
enrichment of genes involved in ecological adaptation to using the different
thistle species based on the strong documented selection pressures, the most
enriched gene functions were instead oogenesis, spermatogenesis, sperm
DNA decondensation and fertilization. The enrichment of these functions is
not easily interpreted, as it either could reflect selection for reproductive
isolation or potentially merely result from LG3 harboring the inversion
being syntenic to the ancestral dipteran X-chromosome and therefore

enriched for *Drosophila melanogaster* X-linked proteins. Although LG3
not the sex chromosome in *T.* conura, it has over evolutionary history been
subjected to sex chromosome- specific selection different to selection on
autosomes due to e.g. the lower effective population sizes of sex
chromosomes (Charlesworth *et al.*, 2018), their potential lower rates of
recombination (Charlesworth, 2017) and the faster X (Mank *et al.*, 2009),
all expected to increase divergence. An additional intriguing possibility is
that lower levels of gene flow could result in the typically more divergent
sex chromosomes, with a large role in reproductive isolation, representing
the ancestral phylogeny despite substantial introgression homogenizing
autosomal regions (Fontaine *et al.*, 2015). These processes could
complicate the interpretation of genomic regions or genes important for
ecological adaptation on these contigs, and disentangling the history of the
chromosome and the selection pressures shaping it in detail remains a topic
for future research.

Instead, the Z chromosome in Tephritis species is likely homologous to *D.*
melanogaster chromosome 4 (aka Muller element F), as previously
described by Vicoso and Bachtrog (2015). Using BLAST, we found that
LG7 in our assembly is the most homologous to *D. melanogaster*
chromosome 4. However, LG7 was highly repetitive in *T. conura*,
averaging 91-97% repetitive content in 50 Kb windows (Fig. S5). As a
result, windows within this chromosome were excluded from most
analyses. Future comparisons of coverage across the genomes of males and
females and gene expression data from reproductive tissues will shed
further light on the sex chromosome in *T. conura* and its importance for
host race formation.

Other major model systems where strong, divergent ecological selection
has resulted in the formation of host races or ecotypes include *R. pomonella*
apple maggot flies infesting hawthorn and apple, respectively (Feder *et al.*,
1988; Filchak *et al.*, 2000; Feder, Berlocher*, et al.*, 2003), the parapatric
ecomorphs of *Littorina* snails adapted to either wave-exposure or crab
predation (Johannesson and Johannesson, 1996; Hollander *et al.*, 2005;
Faria, Chaube*, et al.*, 2019; Perini *et al.*, 2020), and *Timema* walking sticks
adapted to be cryptic on different host plants (Nosil *et al.*, 2002; Nosil,
2007; Nosil and Sandoval, 2008). In contrast to our findings in *T. conura*,
strong selection pressures generally maintain ecologically divergent
populations in sympatry despite more incomplete reproductive isolation in
these systems. *Rhagoletis* flies are isolated by phenological difference and
host preference (Feder, Roethele*, et al.*, 2003; Inskeep *et al.*, 2022), but the
genomic differences that underlie the differently adapted ecotypes is
polygenic and can rapidly be recovered through selection on standing

genetic variation within the ancestral hawthorn host race (Egan *et al.*,
2015). In contrast, the genomic basis conferring mimicry on different host
plants in *Timema* walking consists of few loci of large effects, and the
background genomic divergence is lower (Villoutreix *et al.*, 2020; Nosil *et*
al., 2023). The genomic basis of the *Littorina* ecotypes is similar to that in
T. conura in that variants within inversions contribute to approximately half
of the ecotype divergence (Koch *et al.*, 2022). In contrast to *T. conura*,
several smaller inversions segregate in *Littorina*, enabling a gradual change
in both phenotype and genotype in intermediate selection environments
(Morales *et al.*, 2019; Stankowski *et al.*, 2024).

Compared to those systems, the *T. conura* host races have relatively high
levels of genetic divergence and strong reproductive isolation, arising from
the combination of preference, phenology and performance in the
alternative niche (Diegisser, Johannesen*, et al.*, 2006; Diegisser, Seitz*, et*
al., 2006; Diegisser *et al.*, 2007; Diegisser *et al.*, 2008; Nilsson *et al.*,
2022). Potentially, the adaptation is more multifarious, the genomic
architecture has been permissive for coupling i.e. by physical linkage
(Butlin and Smadja, 2018; Faria, Johannesson*, et al.*, 2019), or adaptation
may have taken place during a longer period of time. In contrast to previous
divergence time estimates based on mitochondria, MSMC analyses suggest
that host race divergence may be old. *Tephritis conura* CH and CO
populations could be older than 500 kya as there is no convergence between
the host races in the MSMC plots even at this time scale. Further
exploration of the dependency of the MSMC inference of which fractions
of the genome are included would be necessary to draw any firm
conclusions though. Given the extreme morphological similarity of the host
races (Nilsson *et al.*, 2022), it may seem surprising that divergence times
exceed previous estimates of just before the last glacial maximum.
However, our new estimates are consistent with the high level of genomic
divergence and limited introgression on LG3 observed in this study. While
divergence is limited across the rest of the genome, this could reflect gene
flow rather than evolutionary divergence time (Fontaine *et al.*, 2015).
Furthermore, we believe this timeframe is also in accordance with strong
reproductive isolation in this system, evidenced by phenological differences
(Romstock-Volkl, 1997), oviposition preference and larval mortality on the
new host plant (Diegisser *et al.*, 2008). Hence, *T. conura* provides a
window into a later stage of ecological speciation continuum (Stankowski
and Ravinet, 2021) with the potential to offer interesting insights into the
development of reproductive barriers following ecological specialization.

Conclusions and future directions

In spite of mounting support for inversions having a role in speciation (Jones *et al.*, 2012; Küpper *et al.*, 2016; Tuttle *et al.*, 2016; Hooper and Price, 2017; Wellenreuther and Bernatchez, 2018; Faria, Chaube*, et al.*, 2019; Faria, Johannesson*, et al.*, 2019; Lucek *et al.*, 2023), studies that uncover how they contribute to coupling of genes underlying ecological and reproductively isolating traits, enabling the formation of persistent species are scarce (but see Meyer *et al.*, 2023; Moan *et al.*, 2023). We find compelling evidence for a large inversion enabling host race specific adaptation and progress towards speciation in *T. conura*. The enrichment of both genes underlying the ability to metabolize host race defense chemicals and gene functions putatively important for mate choice, host plant recognition and phenology which constitutes an important reproductive barrier between host races could imply that this is an empirical example of such coupling through strengthening of physical linkage between genes within the inverted region. Future studies should focus on determining the evolutionary origin of the inversion, including if recruitment of genes to the inverted region has taken place, and experimentally assess if host race specific inversion karyotype genes are associated with ecological and reproductive isolation.

The extensive and recent repeat expansion also means that *T. conura* has the potential to become a study system for investigating the role of transposable elements to contribute in reproductive isolation or to serve as a source for novel variation through altering regulation of gene expression and remodeling chromosomes, has recently been recognized (Serrato-Capuchina and Matute, 2018). To which extent recent TE releases has contributed to the potential to adapt to novel niches, and to reproductively isolate the host races of *T. conura* is an interesting venue for future research that could help shed light on the role of TE releases in ecological speciation. Improved long-read assemblies of the host races would be an exciting future venue, enabling new insights into the chromosomal structural variation underlying host race formation and persistence. *Tephritis conura* exemplifies ongoing ecological speciation of two host races and future research on this study system will offer unique insights into changes in the genome enabling the formation of persistent species to arise during adaptation to novel niches.

Data availability

All code used will be deposited on GitHub upon acceptance. The reference genome will be available from NCBI and all resequencing data will be available at NCBI/ENA upon manuscript acceptance.

Author contributions

AR conceptualized and funded the study and performed field work. KJN and JOG performed field work, and JOG performed all molecular lab work. Bioinformatic analyses were performed by RAS and ZJN, with contributions from JOG, YH, CY and YH, and oversight from AR. The first draft of the manuscript was written by RAS and AR, with contributions from KJN, ZJN, and JOG. RAS, AR, and KJN made significant revisions to the text for subsequent versions.

Acknowledgements

This work was funded by a fellowship from the Wenner-Gren foundations, a Swedish Research Council starting grant, and additional grants from the Crafoord Foundation, Erik Philip Sörensens Stiftelse and Carl Tryggers Stiftelse to AR covering field-, sequencing- and salary costs. Grants from The Royal Physiographic Society of Lund to JO contributed to field work (Young Researchers Endowment) and RNAseq (Nilsson-Ehle Endowment) costs. A Lund Djurskyddsfond grant to JO covered additional field work expenses. Authors acknowledge support from the National Genomics Infrastructure in Stockholm funded by Science for Life Laboratory, the Knut and Alice Wallenberg Foundation and the Swedish Research Council, and SNIC/Uppsala Multidisciplinary Center for Advanced Computational Science for assistance with massively parallel sequencing and access to the UPPMAX computational infrastructure. We thank J. Johanneson for support when starting the work on this study system, H. Vogel for help with long-read extractions, E. Kärrnäs and M. Schnuriger for assistance during field work, and S. Jacobsen Ellerstrand for exploration of the sex chromosome.

References

Alexa, A. and J. Rahnenfuhrer. 2022. topGO: enrichment analysis for gene ontology. R package version 2.48.0.

Aluja, M. and A. L. Norrbom. 2001. *Fruit flies (Tephritidae): phylogeny and evolution of behaviour*. CRC Press, Boca Raton, USA.

Anderson, S. A. S., H. López-Fernández, and J. T. Weir. 2023. Ecology and the origin of nonephemeral species. *The American Naturalist* 201:619-638.

Anderson, S. A. S. and J. T. Weir. 2022. The role of divergent ecological adaptation during allopatric speciation in vertebrates. *Science* 378:1214-1218.

Barbato, M., P. Orozco-terWengel, M. Tapio, and M. W. Bruford. 2015. SNeP: a tool to estimate trends in recent effective population size trajectories using genome-wide SNP data. *Frontiers in Genetics* 6:109.

Berdan, E. L., T. Flatt, G. M. Kozak, K. E. Lotterhos, and B. Wielstra, eds. 2022. Genomic architecture of supergenes: causes and evolutionary consequences [Theme Issue]. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences* 377:20210192

Bhatia, G., N. Patterson, S. Sankararaman, and A. L. Price. 2013. Estimating and interpreting FST: the impact of rare variants. *Genome research* 23:1514-1521.

Boag, P. T. and P. R. Grant. 1981. Intense natural selection in a population of Darwin's finches (*Geospizinae*) in the Galápagos. *Science* 214:82-85.

Bohutínská, M., J. Vlček, S. Yair, B. Laenen, V. Konečná, M. Fracassetti, T. Slotte, and F. Kolář. 2021. Genomic basis of parallel adaptation varies with divergence in *Arabidopsis* and its relatives. *Proceedings of the National Academy of Sciences* 118:e2022713118.

Bolnick, D. I. 2011. Sympatric speciation in threespine stickleback: why not? *International Journal of Ecology* 2011:e942847.

Bomblies, K. and C. L. Peichel. 2022. Genetics of adaptation. *Proceedings of the National Academy of Sciences* 119:e2122152119.

Boulain, H., F. Legeai, J. Jaquiéry, E. Guy, S. Morlière, J.-C. Simon, and A. Sugio. 2019. Differential expression of candidate salivary effector genes in pea aphid biotypes with distinct host plant specificity. *Frontiers in Plant Science* 10:1301.

Bunt, S., C. Hooley, N. Hu, C. Scahill, H. Weavers, and H. Skaer. 2010.
Hemocyte-secreted type IV collagen enhances BMP signaling to guide
renal tubule morphogenesis in *Drosophila*. *Developmental Cell*
19:296-306.

Bush, G. L. 1969. Sympatric host race formation and speciation in
frugivorious flies of genus *Rhagoletis* (Diptera, Tephritidae).
Evolution 23:237-251.

Butlin, R. K. and C. M. Smadja. 2018. Coupling, reinforcement, and
speciation. *The American Naturalist* 191:155-172.

Camacho, C., G. Coulouris, V. Avagyan, N. Ma, J. Papadopoulos, K.
Bealer, and T. L. Madden. 2009. BLAST+: architecture and
applications. *BMC Bioinformatics* 10:421.

Charlesworth, B., J. L. Campos, and B. C. Jackson. 2018. Faster-X
evolution: theory and evidence from *Drosophila*. *Molecular Ecology*
27:3753-3771.

Charlesworth, D. 2017. Evolution of recombination rates between sex
chromosomes. *Philosophical Transactions of the Royal Society of
London. Series B: Biological Sciences* 372:20160456.

Chen, S., Y. Zhou, Y. Chen, and J. Gu. 2018. fastp: an ultra-fast all-in-one
FASTQ preprocessor. *Bioinformatics* 34:i884-i890.

Cheng, H., G. T. Concepcion, X. Feng, H. Zhang, and H. Li. 2021.
Haplotype-resolved de novo assembly using phased assembly graphs
with hifiasm. *Nature Methods* 18:170-175.

Chevin, L.-M., G. Decorzent, and T. Lenormand. 2014. Niche
dimensionality and the genetics of ecological speciation. *Evolution*
68:1244-1256.

Christenhusz, M. J. M. and J. W. Byng. 2016. The number of known plants
species in the world and its annual increase. *Phytotaxa* 261:201-217.

Christenson, L. D. and R. H. Foote. 1960. Biology of fruit flies. *Annual
Review of Entomology* 5:171-192.

Colombani, J. and D. S. Andersen. 2020. The *Drosophila* gut: a gatekeeper
and coordinator of organism fitness and physiology. *WIREs
Developmental Biology* 9:e378.

Coughlan, J. M. and D. R. Matute. 2020. The importance of intrinsic
postzygotic barriers throughout the speciation process. *Philosophical
Transactions of the Royal Society of London. Series B: Biological
Sciences* 375:20190533.

Craig, T. P., J. K. Itami, W. G. Abrahamson, and J. D. Horner. 1993.
Behavioral evidence for host-race formation in *Eurosta solidaginis*.
Evolution 47:1696-1710.

Cruickshank, T. E. and M. W. Hahn. 2014. Reanalysis suggests that
genomic islands of speciation are due to reduced diversity, not
reduced gene flow. *Molecular Ecology* 23:3133-3157.

Cruse, C., T. W. Moural, and F. Zhu. 2023. Dynamic roles of insect
carboxyl/cholinesterases in chemical adaptation. *Insects* 14:194.

Danecek, P., J. K. Bonfield, J. Liddle, J. Marshall, V. Ohan, M. O. Pollard,
A. Whitwham, T. Keane, S. A. McCarthy, R. M. Davies, and H. Li.
2021. Twelve years of SAMtools and BCFtools. *Gigascience*
10:giab008.

Delaneau, O., B. Howie, Anthony J. Cox, J.-F. Zagury, and J. Marchini.
2013. Haplotype estimation using sequencing reads. *The American*
Journal of Human Genetics 93:687-696.

Diegisser, T. 2005. *Artbildung via Wirtsrassen bei Tephritis conura*
(Diptera: Tephritidae). Johannes Gutenberg-Universität Mainz.

Diegisser, T., J. Johannesen, C. Lehr, and A. Seitz. 2004. Genetic and
morphological differentiation in *Tephritis bardanae* (Diptera:
Tephritidae): evidence for host-race formation. *Journal of*
Evolutionary Biology 17:83-93.

Diegisser, T., J. Johannesen, and A. Seitz. 2006. The role of geographic
setting on the diversification process among *Tephritis conura*
(Tephritidae) host races. *Heredity* 96:410-418.

Diegisser, T., J. Johannesen, and A. Seitz. 2008. Performance of host-races
of the fruit fly, *Tephritis conura* on a derived host plant, the cabbage
thistle *Cirsium oleraceum*: Implications for the original host shift.
Journal of Insect Science 8:66.

Diegisser, T., A. Seitz, and J. Johannesen. 2006. Phylogeographic patterns
of host-race evolution in *Tephritis conura* (Diptera: Tephritidae).
Molecular Ecology 15:681-694.

Diegisser, T., A. Seitz, and J. Johannesen. 2007. Morphological adaptation
in host races of *Tephritis conura*. *Entomologia Experimentalis Et*
Applicata 122:155-164.

Dittmar, E. L., C. G. Oakley, J. K. Conner, B. A. Gould, and D. W.
Schemske. 2016. Factors influencing the effect size distribution of
adaptive substitutions. *Proceedings of the Royal Society B: Biological*
Sciences 283:20153065.

Dres, M. and J. Mallet. 2002. Host races in plant-feeding insects and their
importance in sympatric speciation. *Philosophical Transactions of the*
Royal Society of London. Series B: Biological Sciences 357:471-492.

Durand, E. Y., N. Patterson, D. Reich, and M. Slatkin. 2011. Testing for
ancient admixture between closely related populations. *Molecular*
Biology and Evolution 28:2239-2252.

Egan, S. P., G. J. Ragland, L. Assour, T. H. Q. Powell, G. R. Hood, S.
Emrich, P. Nosil, and J. L. Feder. 2015. Experimental evidence of
genome-wide impact of ecological selection during early stages of
speciation-with-gene-flow. *Ecology Letters* 18:817-825.

Ellegren, H. and N. Galtier. 2016. Determinants of genetic diversity. *Nature*
Reviews Genetics 17:422-433.

Ellegren, H., L. Smeds, R. Burri, P. I. Olason, N. Backström, T. Kawakami,
A. Künstner, H. Mäkinen, K. Nadachowska-Brzyska, A. Qvarnström,
S. Uebbing, and J. B. W. Wolf. 2012. The genomic landscape of
species divergence in *Ficedula* flycatchers. *Nature* 491:756-760.

Faria, R., P. Chaube, H. E. Morales, T. Larsson, A. R. Lemmon, E. M.
Lemmon, M. Rafajlović, M. Panova, M. Ravinet, K. Johannesson, A.
M. Westram, and R. K. Butlin. 2019. Multiple chromosomal
rearrangements in a hybrid zone between *Littorina saxatilis* ecotypes.
Molecular Ecology 28:1375-1393.

Faria, R., K. Johannesson, R. K. Butlin, and A. M. Westram. 2019.
Evolving inversions. *Trends in Ecology & Evolution* 34:239-248.

Feder, J. L., S. H. Berlocher, J. B. Roethele, H. Dambroski, J. J. Smith, W.
L. Perry, V. Gavrilovic, K. E. Filchak, J. Rull, and M. Aluja. 2003.
Allopatric genetic origins for sympatric host-plant shifts and race
formation in *Rhagoletis*. *Proceedings of the National Academy of*
Sciences 100:10314-10319.

Feder, J. L., C. A. Chilcote, and G. L. Bush. 1988. Genetic differentiation
between sympatric host races of the apple maggot fly *Rhagoletis*
pomonella. *Nature* 336:61-64.

Feder, J. L., J. B. Roethele, K. Filchak, J. Niedbalski, and J. Romero-
Severson. 2003. Evidence for inversion polymorphism related to
sympatric host race formation in the apple maggot fly, *Rhagoletis*
pomonella. *Genetics* 163:939-953.

Ferris, K. G., L. L. Barnett, B. K. Blackman, and J. H. Willis. 2017. The
genetic architecture of local adaptation and reproductive isolation in
sympatry within the Mimulus guttatus species complex. *Molecular*
Ecology 26:208-224.

Filchak, K. E., J. B. Roethele, and J. L. Feder. 2000. Natural selection and
sympatric divergence in the apple maggot *Rhagoletis pomonella*.
Nature 407:739-742.

Flynn, J. M., R. Hubley, C. Goubert, J. Rosen, A. G. Clark, C. Feschotte,
and A. F. Smit. 2020. RepeatModeler2 for automated genomic
discovery of transposable element families. *Proceedings of the*
National Academy of Sciences 117:9451-9457.

Fontaine, M. C., J. B. Pease, A. Steele, R. M. Waterhouse, D. E. Neafsey, I.
V. Sharakhov, X. Jiang, A. B. Hall, F. Catteruccia, E. Kakani, S. N.
Mitchell, Y.-C. Wu, H. A. Smith, R. R. Love, M. K. Lawniczak, M. A.
Slotman, S. J. Emrich, M. W. Hahn, and N. J. Besansky. 2015.
Extensive introgression in a malaria vector species complex revealed
by phylogenomics. *Science* 347:1258524.

Fox, E. A., A. E. Wright, M. Fumagalli, and F. G. Vieira. 2019. ngsLD:
evaluating linkage disequilibrium using genotype likelihoods.
Bioinformatics 35:3855-3856.

Gabriel, L., T. Brůna, K. J. Hoff, M. Ebel, A. Lomsadze, M. Borodovsky,
and M. Stanke. 2024. BRAKER3: Fully automated genome annotation
using RNA-seq and protein evidence with GeneMark-ETP,
AUGUSTUS and TSEBRA. *bioRxiv* 2023.2006.2010.544449.

Gautam, N. K., P. Verma, and M. G. Tapadia. 2017. *Drosophila*
malpighian tubules: a model for understanding kidney development,
function, and disease. Pp. 3-25 *in* R. K. Miller, ed. *Kidney*
Development and Disease. Springer, Cham.

Gow, J. L., C. L. Peichel, and E. B. Taylor. 2006. Contrasting hybridization
rates between sympatric three-spined sticklebacks highlight the
fragility of reproductive barriers between evolutionarily young
species. *Molecular Ecology* 15:739-752.

Green, R. E., J. Krause, A. W. Briggs, T. Maricic, U. Stenzel, M. Kircher,
N. Patterson, H. Li, W. Zhai, M. H.-Y. Fritz, N. F. Hansen, E. Y.
Durand, A.-S. Malaspinas, J. D. Jensen, T. Marques-Bonet, C. Alkan,
K. Prüfer, M. Meyer, H. A. Burbano, J. M. Good, R. Schultz, A.
Aximu-Petri, A. Butthof, B. Höber, B. Höffner, M. Siegemund, A.
Weihmann, C. Nusbaum, E. S. Lander, C. Russ, N. Novod, J.
Affourtit, M. Egholm, C. Verna, P. Rudan, D. Brajkovic, Ž. Kucan, I.
Gušic, V. B. Doronichev, L. V. Golovanova, C. Lalueza-Fox, M. de la
Rasilla, J. Fortea, A. Rosas, R. W. Schmitz, P. L. F. Johnson, E. E.
Eichler, D. Falush, E. Birney, J. C. Mullikin, M. Slatkin, R. Nielsen, J.
Kelso, M. Lachmann, D. Reich, and S. Pääbo. 2010. A draft sequence
of the Neandertal genome. *Science* 328:710-722.

Grossmann, S., S. Bauer, P. N. Robinson, and M. Vingron. 2007. Improved
detection of overrepresentation of gene-ontology annotations with
parent child analysis. *Bioinformatics* 23:3024-3031.

Gu, Z., L. Gu, R. Eils, M. Schlesner, and B. Brors. 2014. circlize
implements and enhances circular visualization in R. *Bioinformatics*
30:2811-2812.

Guan, D., S. A. McCarthy, J. Wood, K. Howe, Y. Wang, and R. Durbin.
2020. Identifying and removing haplotypic duplication in primary
genome assemblies. *Bioinformatics* 36:2896-2898.

Hardy, N. B. and S. P. Otto. 2014. Specialization and generalization in the
diversification of phytophagous insects: tests of the musical chairs and
oscillation hypotheses. *Proceedings of the Royal Society B: Biological*
Sciences 281:20132960.

Headrick, D. H. and R. D. Goeden. 1998. The biology of nonfrugivorous
tephritid fruit flies. *Annual Review of Entomology* 43:217-241.

Hendry, A. P. 2009. Ecological speciation! Or the lack thereof? *Canadian*
Journal of Fisheries and Aquatic Sciences 66:1383-1398.

Herrel, A., K. Huyghe, B. Vanhooydonck, T. Backeljau, K. Breugelmans, I.
Grbac, R. Van Damme, and D. J. Irschick. 2008. Rapid large-scale
evolutionary divergence in morphology and performance associated
with exploitation of a different dietary resource. *Proceedings of the*
National Academy of Sciences 105:4792-4795.

Hollander, J., M. Lindegarth, and K. Johannesson. 2005. Local adaptation
but not geographical separation promotes assortative mating in a snail.
Animal Behaviour 70:1209-1219.

Hood, G. R., T. H. Q. Powell, M. M. Doellman, S. B. Sim, M. Glover, W.
L. Yee, R. B. Goughnour, M. Mattsson, D. Schwarz, and J. L. Feder.
2020. Rapid and repeatable host plant shifts drive reproductive
isolation following a recent human-mediated introduction of the apple
maggot fly, *Rhagoletis pomonella*. *Evolution* 74:156-168.

Hooper, D. M., S. C. Griffith, and T. D. Price. 2019. Sex chromosome
inversions enforce reproductive isolation across an avian hybrid zone.
Molecular Ecology 28:1246-1262.

Hooper, D. M. and T. D. Price. 2017. Chromosomal inversion differences
correlate with range overlap in passerine birds. *Nature Ecology &*
Evolution 1:1526-1534.

Huerta-Cepas, J., D. Szklarczyk, D. Heller, A. Hernández-Plaza, S. K.
Forslund, H. Cook, D. R. Mende, I. Letunic, T. Rattei, Lars J. Jensen,
C. von Mering, and P. Bork. 2018. eggNOG 5.0: a hierarchical,
functionally and phylogenetically annotated orthology resource based
on 5090 organisms and 2502 viruses. *Nucleic Acids Research*
47:D309-D314.

Inskeep, K. A., M. M. Doellman, T. H. Q. Powell, S. H. Berlocher, N. R.
Seifert, G. R. Hood, G. J. Ragland, P. J. Meyers, and J. L. Feder. 2022.
Divergent diapause life history timing drives both allochronic
speciation and reticulate hybridization in an adaptive radiation of
Rhagoletis flies. *Molecular Ecology* 31:4031-4049.

Jain, K., S. Lavista-Llanos, V. Grabe, B. S. Hansson, and D. Wicher. 2021.
Calmodulin regulates the olfactory performance in Drosophila
melanogaster. *Scientific Reports* 11:3747.

Johannesson, B. and K. Johannesson. 1996. Population differences in
behaviour and morphology in the snail *Littorina saxatilis*: Phenotypic
plasticity or genetic differentiation? *Journal of Zoology* 240:475-493.

Johannesson, K. 2001. Parallel speciation: a key to sympatric divergence.
Trends in Ecology & Evolution 16:148-153.

Johannesson, K., M. Panova, P. Kemppainen, C. André, E. Rolán-Alvarez,
and R. K. Butlin. 2010. Repeated evolution of reproductive isolation
in a marine snail: unveiling mechanisms of speciation. *Philosophical*
Transactions of the Royal Society of London. Series B: Biological
Sciences 365:1735-1747.

Jones, F. C., M. G. Grabherr, Y. F. Chan, P. Russell, E. Mauceli, J.
Johnson, R. Swofford, M. Pirun, M. C. Zody, S. White, E. Birney, S.
Searle, J. Schmutz, J. Grimwood, M. C. Dickson, R. M. Myers, C. T.
Miller, B. R. Summers, A. K. Knecht, S. D. Brady, H. Zhang, A. A.
Pollen, T. Howes, C. Amemiya, J. Baldwin, T. Bloom, D. B. Jaffe, R.
Nicol, J. Wilkinson, E. S. Lander, F. Di Palma, K. Lindblad-Toh, and
D. M. Kingsley. 2012. The genomic basis of adaptive evolution in
threespine sticklebacks. *Nature* 484:55-61.

Junier, T. and E. M. Zdobnov. 2010. The Newick utilities: high-throughput
phylogenetic tree processing in the Unix shell. *Bioinformatics*
26:1669-1670.

Kautt, A. F., C. F. Kratochwil, A. Nater, G. Machado-Schiaffino, M. Olave,
F. Henning, J. Torres-Dowdall, A. Härer, C. D. Hulsey, P. Franchini,
M. Pippel, E. W. Myers, and A. Meyer. 2020. Contrasting signatures
of genomic divergence during sympatric speciation. *Nature* 588:106-
111.

Keightley, P. D., U. Trivedi, M. Thomson, F. Oliver, S. Kumar, and M. L.
Blaxter. 2009. Analysis of the genome sequences of three *Drosophila*
melanogaster spontaneous mutation accumulation lines. *Genome*
research 19:1195-1201.

Kim, D., J. M. Paggi, C. Park, C. Bennett, and S. L. Salzberg. 2019. Graph-
based genome alignment and genotyping with HISAT2 and HISAT-
genotype. *Nature Biotechnology* 37:907-915.

Kirkpatrick, M. and N. Barton. 2006. Chromosome inversions, local
adaptation and speciation. *Genetics* 173:419-434.

Koch, E. L., M. Ravinet, A. M. Westram, K. Johannesson, and R. K.
Butlin. 2022. Genetic architecture of repeated phenotypic divergence
in *Littorina saxatilis* ecotype evolution. *Evolution* 76:2332-2346.

Korneliussen, T. S., A. Albrechtsen, and R. Nielsen. 2014. ANGSD:
analysis of next generation sequencing data. *BMC Bioinformatics*
15:356.

Kriventseva, E. V., D. Kuznetsov, F. Tegenfeldt, M. Manni, R. Dias, F. A.
Simão, and E. M. Zdobnov. 2019. OrthoDB v10: sampling the
diversity of animal, plant, fungal, protist, bacterial and viral genomes
for evolutionary and functional annotations of orthologs. *Nucleic*
Acids Res 47:D807-d811.

Kulmuni, J., R. K. Butlin, K. Lucek, V. Savolainen, and A. M. Westram,
eds. 2020. Towards the completion of speciation: the evolution of
reproductive isolation beyond the first barriers [Theme issue].
Philosophical Transactions of the Royal Society of London. Series B:
Biological Sciences 375:20190528

Küpper, C., M. Stocks, J. E. Risse, N. dos Remedios, L. L. Farrell, S. B.
McRae, T. C. Morgan, N. Karlionova, P. Pinchuk, Y. I. Verkuil, A. S.
Kitaysky, J. C. Wingfield, T. Piersma, K. Zeng, J. Slate, M. Blaxter,
D. B. Lank, and T. Burke. 2016. A supergene determines highly
divergent male reproductive morphs in the ruff. *Nature Genetics*
48:79-83.

Kuznetsov, D., F. Tegenfeldt, M. Manni, M. Seppey, M. Berkeley,
Evgenia V. Kriventseva, and E. M. Zdobnov. 2022. OrthoDB v11:
annotation of orthologs in the widest sampling of organismal diversity.
Nucleic Acids Research 51:D445-D451.

Lackey, A. C. R. and J. W. Boughman. 2017. Evolution of reproductive
isolation in stickleback fish. *Evolution* 71:357-372.

Li, H. 2011. A statistical framework for SNP calling, mutation discovery,
association mapping and population genetical parameter estimation
from sequencing data. *Bioinformatics* 27:2987-2993.

Li, H. 2013. Aligning sequence reads, clone sequences and assembly
contigs with BWA-MEM. *arXiv preprint*

Li, H. 2021. New strategies to improve minimap2 alignment accuracy.
Bioinformatics 37:4572-4574.

Li, H. and R. Durbin. 2009. Fast and accurate short read alignment with
Burrows–Wheeler transform. *Bioinformatics* 25:1754-1760.

Low, W. Y., H. L. Ng, C. J. Morton, M. W. Parker, P. Batterham, and C.
Robin. 2007. Molecular evolution of glutathione S-transferases in the
genus *Drosophila*. *Genetics* 177:1363-1375.

Lucek, K., M. D. Giménez, M. Joron, M. Rafajlović, J. B. Searle, N.
Walden, A. M. Westram, and R. Faria. 2023. The impact of
chromosomal rearrangements in speciation: from micro- to
macroevolution. *Cold Spring Harbor Perspectives in Biology*
15:a041447.

Malinsky, M., R. J. Challis, A. M. Tyers, S. Schiffels, Y. Terai, B. P.
Ngatunga, E. A. Miska, R. Durbin, M. J. Genner, and G. F. Turner.
2015. Genomic islands of speciation separate cichlid ecomorphs in an
East African crater lake. *Science* 350:1493-1498.

Malinsky, M., M. Matschiner, and H. Svardal. 2021. Dsuite - fast D-
statistics and related admixture evidence from VCF files. *Molecular
Ecology Resources* 21:584-595.

Mank, J. E., K. Nam, and H. Ellegren. 2009. Faster-Z evolution is
predominantly due to genetic drift. *Molecular Biology and Evolution*
27:661-670.

Manni, M., M. R. Berkeley, M. Seppey, and E. M. Zdobnov. 2021.
BUSCO: assessing genomic data quality and beyond. *Current
Protocols* 1:e323.

Meier, J. I., D. A. Marques, C. E. Wagner, L. Excoffier, and O. Seehausen.
2018. Genomics of parallel ecological speciation in Lake Victoria
cichlids. *Molecular Biology and Evolution* 35:1489-1506.

Meyer, L., P. Barry, F. Riquet, A. Foote, C. D. Sarkissian, R. Cunha, C.
Arbiol, F. Cerqueira, E. Desmarais, A. Bordes, N. Bierne, B. Guinand,
and P.-A. Gagnaire. 2023. Divergence and gene flow history at two
large chromosomal inversions involved in long-snouted seahorse
ecotype formation. *bioRxiv* 2023.2007.2004.547634.

Minh, B. Q., H. A. Schmidt, O. Chernomor, D. Schrempf, M. D.
Woodhams, A. von Haeseler, and R. Lanfear. 2020. IQ-TREE 2: new
models and efficient methods for phylogenetic inference in the
genomic era. *Molecular Biology and Evolution* 37:1530-1534.

Moan, A. L., S. Stankowski, M. Rafajlovic, O. Ortega-Martinez, R. Faria,
R. Butlin, and K. Johannesson. 2023. Coupling of 12 chromosomal
inversions maintains a strong barrier to gene flow between ecotypes.
bioRxiv 2023.2009.2018.558209.

Mölder, F., K. Jablonski, B. Letcher, M. Hall, C. Tomkins-Tinch, V.
Sochat, J. Forster, S. Lee, S. Twardziok, A. Kanitz, A. Wilm, M.
Holtgrewe, S. Rahmann, S. Nahnsen, and J. Köster. 2021. Sustainable
data analysis with Snakemake *F1000Research* 10:33.

Moon, S. J., Y. Lee, Y. Jiao, and C. Montell. 2009. A *Drosophila* gustatory
receptor essential for aversive taste and inhibiting male-to-male
courtship. *Current Biology* 19:1623-1627.

Mora, C., D. P. Tittensor, S. Adl, A. G. B. Simpson, and B. Worm. 2011.
How many species are there on earth and in the ocean? *Plos Biology*
9:8.

Morales, H. E., R. Faria, K. Johannesson, T. Larsson, M. Panova, A. M.
Westram, and R. K. Butlin. 2019. Genomic architecture of parallel
ecological divergence: beyond a single environmental contrast.
Science Advances 5:eaav9963.

Nallu, S., J. A. Hill, K. Don, C. Sahagun, W. Zhang, C. Meslin, E. Snell-
Rood, N. L. Clark, N. I. Morehouse, J. Bergelson, C. W. Wheat, and
M. R. Kronforst. 2018. The molecular genetic basis of herbivory
between butterflies and their host plants. *Nature Ecology & Evolution*
2:1418-1427.

Nielsen, R., T. Korneliussen, A. Albrechtsen, Y. Li, and J. Wang. 2012.
SNP calling, genotype calling, and sample allele frequency estimation
from new-generation sequencing data. *Plos One* 7:e37558.

Nilsson, K. J., J. Ortega, M. Friberg, and A. Runemark. 2022. Non-parallel
morphological divergence following colonization of a new host plant.
Evolutionary Ecology 36:859-877.

Nosil, P. 2007. Divergent host plant adaptation and reproductive isolation
between ecotypes of *Timema cristinae* walking sticks. *The American*
Naturalist 169:151-162.

Nosil, P. 2012. *Ecological speciation*. Oxford University Press, Oxford,
UK.

Nosil, P., B. J. Crespi, and C. P. Sandoval. 2002. Host-plant adaptation
drives the parallel evolution of reproductive isolation. *Nature*
417:440-443.

Nosil, P. and L. Harmon. 2009. Niche dimensionality and ecological
speciation. Pp. 127-154 *in* R. Butlin, J. Bridle, and D. Schluter, eds.
Speciation and Patterns of Diversity. Cambridge University Press,
Cambridge, UK.

Nosil, P. and C. P. Sandoval. 2008. Ecological niche dimensionality and the
evolutionary diversification of stick insects. *Plos One* 3:e1907.

Nosil, P., V. Soria-Carrasco, R. Villoutreix, M. De-la-Mora, C. F. de
Carvalho, T. Parchman, J. L. Feder, and Z. Gompert. 2023. Complex
evolutionary processes maintain an ancient chromosomal inversion.
Proceedings of the National Academy of Sciences 120:e2300673120.

Nouhaud, P., M. Gautier, A. Gouin, J. Jaquiéry, J. Peccoud, F. Legeai, L.
Mieuzet, C. M. Smadja, C. Lemaitre, R. Vitalis, and J.-C. Simon.
2018. Identifying genomic hotspots of differentiation and candidate
genes involved in the adaptive divergence of pea aphid host races.
Molecular Ecology 27:3287-3300.

Patterson, N., P. Moorjani, Y. Luo, S. Mallick, N. Rohland, Y. Zhan, T.
Genschoreck, T. Webster, and D. Reich. 2012. Ancient admixture in
human history. *Genetics* 192:1065-1093.

Pečnerová, P., G. Garcia-Erill, X. Liu, C. Nursyifa, R. K. Waples, C. G.
Santander, L. Quinn, P. Frandsen, J. Meisner, F. F. Stæger, M. S.
Rasmussen, A. Brüniche-Olsen, C. Hviid Friis Jørgensen, R. R. da
Fonseca, H. R. Siegismund, A. Albrechtsen, R. Heller, I. Moltke, and
K. Hanghøj. 2021. High genetic diversity and low differentiation
reflect the ecological versatility of the African leopard. *Current
Biology* 31:1862-1871.e1865.

Perini, S., M. Rafajlovic, A. M. Westram, K. Johannesson, and R. K.
Butlin. 2020. Assortative mating, sexual selection, and their
consequences for gene flow in *Littorina*. *Evolution* 74:1482-1497.

Picard toolkit. 2019. Broad Institute. http://broadinstitute.github.io/picard/.

Quinlan, A. R. and I. M. Hall. 2010. BEDTools: a flexible suite of utilities
for comparing genomic features. *Bioinformatics* 26:841-842.

Rice, W. R. and E. E. Hostert. 1993. Laboratory experiments on speciation:
what have we learned in 40 years? *Evolution* 47:1637-1653.

Rivas, M. J., M. Saura, A. Pérez-Figueroa, M. Panova, T. Johansson, C. André, A. Caballero, E. Rolán-Alvarez, K. Johannesson, and H. Quesada. 2018. Population genomics of parallel evolution in gene expression and gene sequence during ecological adaptation. *Scientific Reports* 8:16147.

Romstock-Volkl, M. 1997. Host race formation in *Tephritis conura*: determinants from three trophic levels. Pp. 21-38 *in* K. Dettner, G. Bauer, and W. Völkl, eds. *Vertical Food Web Interactions*. Springer, Berlin, Germany.

Sato, K., M. T. Ahsan, M. Ote, M. Koganezawa, and D. Yamamoto. 2019. Calmodulin-binding transcription factor shapes the male courtship song in Drosophila. *PLOS Genetics* 15:e1008309.

Schaal, S. M., B. C. Haller, and K. E. Lotterhos. 2022. Inversion invasions: when the genetic basis of local adaptation is concentrated within inversions in the face of gene flow. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences* 377:20210200.

Schiffels, S. and R. Durbin. 2014. Inferring human population size and separation history from multiple genome sequences. *Nature Genetics* 46:919-925.

Schluter, D. 2009. Evidence for ecological speciation and its alternative. *Science* 323:737-741.

Schluter, D. and L. H. Rieseberg. 2022. Three problems in the genetics of speciation by selection. *Proceedings of the National Academy of Sciences* 119:e2122153119.

Seitz, A. and M. Komma. 1984. Genetic polymorphism and its ecological background in tephritid populations (*Diptera: Tephritidae*). Pp. 143-158 *in* K. Wöhrmann, and V. Loeschcke, eds. *Population Biology and Evolution*. Springer Berlin Heidelberg, Berlin, Heidelberg.

Serrato-Capuchina, A. and D. R. Matute. 2018. The role of transposable elements in speciation. *Genes* 9:254.

Shih, P.-Y., A. Sugio, and J.-C. Simon. 2023. Molecular mechanisms underlying host plant specificity in aphids. *Annual Review of Entomology* 68:431-450.

Singh, K. S., B. J. Troczka, A. Duarte, V. Balabanidou, N. Trissi, L. Z.
Carabajal Paladino, P. Nguyen, C. T. Zimmer, K. M. Papapostolou, E.
Randall, B. Lueke, F. Marec, E. Mazzoni, M. S. Williamson, A.
Hayward, R. Nauen, J. Vontas, and C. Bass. 2020. The genetic
architecture of a host shift: an adaptive walk protected an aphid and its
endosymbiont from plant chemical defenses. *Science Advances*
6:eaba1070.

Skotte, L., T. S. Korneliussen, and A. Albrechtsen. 2013. Estimating
individual admixture proportions from next generation sequencing
data. *Genetics* 195:693-702.

Smit, A. F., R. Hubley, and P. Green. 2015. RepeatMaster Open-4.0.
http://www.repeatmasker.org.

Sproul, J. S., S. Hotaling, J. Heckenhauer, A. Powell, D. Marshall, A. M.
Larracuente, J. L. Kelley, S. U. Pauls, and P. B. Frandsen. 2023.
Analyses of 600+ insect genomes reveal repetitive element dynamics
and highlight biodiversity-scale repeat annotation challenges. *Genome*
research 33:gr.277387.277122.

Stankowski, S. and M. Ravinet. 2021. Defining the speciation continuum.
Evolution 75:1256-1273.

Stankowski, S., Z. B. Zagrodzka, M. D. Garlovsky, A. Pal, D. Shipilina, D.
G. Castillo, H. Lifchitz, A. Le Moan, E. Leder, J. Reeve, K.
Johannesson, A. M. Westram, and R. K. Butlin. 2024. The genetic
basis of a recent transition to live-bearing in marine snails. *Science*
383:114-119.

Steward, R. A., J. O. Giménez, S. Choudhary, S. Yi, O. V. Aken, and A.
Runemark. 2024. Evolved and plastic gene expression in adaptation to
a novel niche. *bioRxiv* 2024.2003.2002.583104.

Stuart, Y. E., T. Veen, J. N. Weber, D. Hanson, M. Ravinet, B. K. Lohman,
C. J. Thompson, T. Tasneem, A. Doggett, R. Izen, N. Ahmed, R. D.
H. Barrett, A. P. Hendry, C. L. Peichel, and D. I. Bolnick. 2017.
Contrasting effects of environment and genetics generate a continuum
of parallel evolution. *Nature Ecology & Evolution* 1:0158.

Szpiech, Z. A. and R. D. Hernandez. 2014. selscan: an efficient
multithreaded program to perform EHH-based scans for positive
selection. *Molecular Biology and Evolution* 31:2824-2827.

Taylor, E. B., J. W. Boughman, M. Groenenboom, M. Sniatynski, D.
Schluter, and J. L. Gow. 2006. Speciation in reverse: morphological
and genetic evidence of the collapse of a three-spined stickleback
(Gasterosteus aculeatus) species pair. *Molecular Ecology* 15:343-355.

Tuttle, Elaina M., Alan O. Bergland, Marisa L. Korody, Michael S. Brewer,
Daniel J. Newhouse, P. Minx, M. Stager, A. Betuel, Zachary A.
Cheviron, Wesley C. Warren, Rusty A. Gonser, and Christopher N.
Balakrishnan. 2016. Divergence and functional degradation of a sex
chromosome-like supergene. *Current Biology* 26:344-350.

van Schooten, B., J. Meléndez-Rosa, S. M. Van Belleghem, C. D. Jiggins,
J. D. Tan, W. O. McMillan, and R. Papa. 2020. Divergence of
chemosensing during the early stages of speciation. *Proceedings of the*
National Academy of Sciences 117:16438-16447.

Vicoso, B. and D. Bachtrog. 2013. Reversal of an ancient sex chromosome
to an autosome in *Drosophila*. *Nature* 499:332-335.

Vicoso, B. and D. Bachtrog. 2015. Numerous transitions of sex
chromosomes in Diptera. *Plos Biology* 13:e1002078.

Vidal, M. C. and S. M. Murphy. 2018. Bottom-up vs. top-down effects on
terrestrial insect herbivores: a meta-analysis. *Ecology Letters* 21:138-
150.

Villoutreix, R., C. F. de Carvalho, V. Soria-Carrasco, D. Lindtke, M. De-la-
Mora, M. Muschick, J. L. Feder, T. L. Parchman, Z. Gompert, and P.
Nosil. 2020. Large-scale mutation in the evolution of a gene complex
for cryptic coloration. *Science* 369:460-466.

Wellenreuther, M. and L. Bernatchez. 2018. Eco-evolutionary genomics of
chromosomal inversions. *Trends in Ecology & Evolution* 33:427-440.

Wood, D. E., J. Lu, and B. Langmead. 2019. Improved metagenomic
analysis with Kraken 2. *Genome Biology* 20:257.

Yin, J., C. Zhang, and S. Mirarab. 2019. ASTRAL-MP: scaling ASTRAL
to very large datasets using randomization and parallelization.
Bioinformatics 35:3961-3969.

Yu, Q.-Y., C. Lu, W.-L. Li, Z.-H. Xiang, and Z. Zhang. 2009. Annotation
and expression of carboxylesterases in the silkworm, *Bombyx mori*.
BMC Genomics 10:553.

Zhang, C., M. Rabiee, E. Sayyari, and S. Mirarab. 2018. ASTRAL-III:
polynomial time species tree reconstruction from partially resolved
gene trees. *BMC Bioinformatics* 19:153.

Zhou, C., S. A. McCarthy, and R. Durbin. 2022. YaHS: yet another Hi-C
scaffolding tool. *Bioinformatics* 39:btac808.

1467

1468
1469

Figure Captions

Figure 1. Genomic divergence of Tephritis conura (A; female pictured)
host races specializing on Cirsium heterophyllum (CH; B) or C.
oleraceum (CO; C) thistle buds. (D) Flies were sampled from populations
of the CH (purple points) or CO (green points) host race along transects
east (diamonds) and west (circles) of the Baltic Sea. Transects traversed
regionds of northern Europe where *C. heterophyllum* exists in allopatry
(purple background), *C. oleraceum* exists in allopatry (green background),
or both host plants are found in sympatry (striped background). (**E**)
Admixture proportions estimated for K=2 and K=3 (see Fig. S6A for K4-
9). (**F**) Principal component (PC) analysis of genetic differences among
populations separates CH (purple) and CO (green) individuals on the first
PC axis (see Fig. S6B for comparison of the second and third PC axes). (**G**)
ASTRAL consensus cladogram of flies used in the study. Terminal nodes
are colored by population and internal nodes with strong support (posterior
probability > 0.8) are colored in red. For branch lengths and posterior
probabilities of all nodes, see Fig. S7. (**H**) MSMC analysis of demographic
history using eight individuals with the most similar coverage from each
population and excluding highly divergent or repetitive regions.

Figure 2. Highly divergent genomic windows between host races based
on (**A**) BayPass analysis for association with host race using all 8
populations (**B**) population branch statistics (PBS), and (**C**) relative node
depth (RND) for COSK, the sympatric CO population in the west, as a
representative (see Fig. S13 for remaining triads). Statistics were evaluated
over 50 kb non-overlapping windows. Windows were considered outliers
when greater than three standard deviations away from the mean (dashed
line). (**D**) Overlap between outlier sets within CH and CO comparisons
within western and eastern transects. Windows were considered highly
differentiated if they were outliers in at least two of the three metrics tested.
(**H**) Overlap between all highly differentiated windows.

Figure 3. Host race-specific haplotypes within the highly differentiated
region suggest a large inversion. (**A**) PC1 across ordered contigs in the
genome for all 96 individuals. Zoom panel highlights the putative
inversion, with potential break points indicated by black lines. Contigs were
included if more than 200 filtered variants were used to calculate the PCA
and are ordered according to HiC scaffolds (Fig. S1). Hypothetical linkage
groups are shown with alternating gray and white bands. The right-most
white band represents unplaced scaffolds. The convergence of haplotypes at

760 Mbp is likely a scaffolding error. (**B-D**) PCA of two contigs outside
and one inside the putative inversion. Locations of contigs in panels B-D
are indicated in A.

Figure 4. Signatures of selection across the genome. (**A**) Nucleotide
diversity (π) in 50 kb windows across the genome for two representative
populations (CHSK and COSK, see Fig. S14 for remaining populations).
Windows with lower than 20% coverage were excluded. Highly
differentiated windows are colored by population. (**B**) π in highly
differentiated windows and undifferentiated windows for all populations.
Horizontal lines in violin plots show 25th, 50th, and 75th quantiles and
points indicate the mean. (**C**) Tajima's D in 50 kb windows across the
genome for two representative populations (CHSK and COSK, see Fig. S15
for remaining populations). (**D**) Tajima's D in highly differentiated and
undifferentiated windows for all populations. (**E**) Average XP-EHH in 50
kb windows across the genome for a representative comparison (COSK vs.
CHSK, see Fig. S16 for remaining populations). (**F**) Average XP-EHH in
highly differentiated and undifferentiated windows for contrasts between
sympatric populations and between allopatric populations of different host
races in the west and east, and between western and eastern sympatric
populations of the same host race ('Between'). Medians were compared
between highly differentiated windows and undifferentiated windows using
Mann-Whitney tests ($p < 0.05$ *, $p < 0.01$ **, $p < 0.001$ ***, $p < 0.0001$
****, Table S3-S5).

Figure 5. Introgression in across the genome. (**A**) Introgression was
compared between triads of *T. conura* populations using f_{dM}. Triads were
composed of two populations of the same host race (P1 = allopatric, P2 =
sympatric) and a third population of the other host race (P3). Pictured are
f_{dM} in 50 kb windows for two representative P3 populations, CHSK and
COSK. Values close to zero indicate little to no introgression, negative
values indicate greater relative introgression between P1 and P3, while
positive values indicate greater relative introgression between P2 and P3.
(**B**) Introgression in highly differentiated windows (colored according to
P3) and undifferentiated windows (gray) for all eight populations. Medians
were significantly higher in the west and lower in the east in highly
differentiated windows compared to undifferentiated windows (Mann-
Whitney, $p < 0.0001$ ****, Table S9), while variance around the median
was consistently significantly lower in highly differentiated windows
(Levene's test; Table S8).

1546

A *T. conura*

B *C. heterophyllum*

C *C. oleraceum*

D

CHFI
CHST
CHES
COES
COSK
CHSK
COLI
COGE

E

○ CHSK ● CHST ◆ CHFI ◇ CHES ◇ COES ◆ COLI ● COGE ● COSK

Admixture proportion

K = 2

K = 3

Individual

F

PC2 (1.47%)

West
● CHST
● CHSK
● COSK
● COGE

East
◆ CHFI
◇ CHES
◇ COES
◆ COLI

Sympatric host ranges

PC1 (2.13%)

G

outgroups

H

Effective population size

CHST   CHFI
CHSK   CHES
COSK   COES
COGE   COLI

Time (years ago)