Do glucocorticoids indicate stress? Meta-analysis reveals a tight association between glucocorticoids and metabolic rate

Blanca Jimeno[1, 2*] & Simon Verhulst[3]

[1] Instituto de Investigación en Recursos Cinegéticos (IREC), CSIC-UCLM-JCCM, Ronda de Toledo 12, 13005 Ciudad Real, Spain

[2] Instituto Pirenaico de Ecologia (IPE), CSIC, Avda. Nuestra Señora de la Victoria, 16, Jaca, Spain

[3] University of Groningen, the Netherlands

* Instituto Pirenaico de Ecologia (IPE), CSIC, Avda. Nuestra Señora de la Victoria, 16, Jaca, Spain

bjimenorev@gmail.com

ABSTRACT

Glucocorticoid (GC) variation has long been thought to reflect variation in organismal 'stress', but associations between GCs and Darwinian fitness components are diverse in magnitude, direction, and highly context-dependent. This paradox reveals our poor understanding of the causes of GC variation, contrasting with the detailed knowledge of the functional consequences of GC variation. Amongst other things, GCs orchestrate energy availability, to anticipate and recover from predictable and unpredictable environmental fluctuations and challenges. Although this is mechanistically well-known, the extent to which GC levels are quantitatively explained by energy metabolism is unresolved. We investigated this association through meta-analysis, selecting studies of endotherms in which (1) an experiment was performed that affected metabolic rate and (2) metabolic rate and GC-levels were measured simultaneously. We found that an increase in metabolic rate was associated with an increase in GC levels in 20 out of 21 studies (32 out of 35 effect sizes). More importantly, there was a strong positive correlation between the increases in metabolic rate and GCs (p=0.003). This pattern was similar in birds and mammals, and independent of the nature of the experimental treatment. We conclude that metabolic rate is a major driver of GC variation within individuals. Stressors often affect metabolic rate, leading us to question whether GC levels provide information on 'stress' beyond the stressor's effect on metabolic rate.

KEYWORDS

Glucocorticoid, cortisol, corticosterone, metabolism, energy expenditure.

INTRODUCTION

The accepted - but often overlooked - metabolic role of glucocorticoids

Glucocorticoid hormones (GCs; e.g. cortisol, corticosterone) were identified by Hans Selye (1907-1982) as the key molecular mediators of the 'stress reaction', and named in reference to their capacity to increase glucose in blood. Selye highlighted the fundamental role of GCs in the 'general adaptation syndrome' - i.e. *"the physiological mechanisms that help to raise resistance to damage as such, irrespective of the specific nature of the damaging agents"* (Selye 1950). In this view, GCs play a role in 'adapting' to the challenge, by triggering mechanisms that help the organism return to, or maintain, homeostasis after an environmental challenge. Indeed, towards the later stages of his career, Selye's definition of stress was *"the nonspecific response of the body on any demand on it"* (Selye 1976), making the point that very different stimuli (i.e. 'stressors') triggered the same response. However, in the literature this perspective has changed over time, and 'stress' and, by extension, GCs, have been widely linked to negative outcomes (but see Koolhaas *et al.* 2011; Herman 2022). Consequently, researchers in fields from biomedicine to conservation physiology and animal husbandry have focused on GCs to find proxies of 'physiological and/or psychological stress' to evaluate physical and/or welfare status. Indeed, GCs have predominated over other traits throughout the stress physiology literature, to the point of being referred to as vertebrate 'stress hormones' (Madliger *et al.* 2015; Madliger & Love 2014; McCormick & Romero 2017). While several authors have argued against this simplified view of GC regulation during the past decades (e.g. Koolhaas *et al.*, 2011; MacDougall-Shackleton *et al.* 2019; Herman 2022; Romero 2004; Landys *et al.* 2006; Bonier *et al.* 2009), such association prevails, in the sense that publications still abound in which GC levels are assumed to provide information on organismal stress.

The focus on GCs to measure organismal stress can be understood from their role as key mediators of organismal responses to challenges, triggering a cascade of effects on many physiological systems (Koolhaas *et al.*, 2011; Sapolsky, Romero & Munck, 2000; Deviche *et al.* 2017; Zimmer *et al.* 2019; Zimmer *et al.* 2020). Furthermore, acute increases in circulating GCs are a defining aspect of the so-called "stress response" (Koolhaas *et al.*, 2011; Sapolsky *et al.* 2000). Consequently, higher GC levels circulating in plasma or deposited in keratinized tissues (i.e. hair or feathers) have traditionally been interpreted as an indication of homeostatic unbalance, poor condition, implicating low fitness prospects (reviewed in Schoenle *et al.* 2021, Romero & Fairhurst 2016). However, the latter assumption is at best poorly supported by the literature, with associations between GCs and fitness (survival / reproduction) being diverse in direction and often non-existent (Schoenle *et al.* 2021; Zimmer *et al.* 2022; Busch & Hayward 2009; Bonier *et al.* 2009; Petrullo *et al.* 2022). This inconsistency raises the question what alternative inference can be made from GC-variation. The urgency of this question is further underlined by the observation that GCs also increase in response to experiences we would not normally qualify as stress; for example, sexual activity induces a GC increase in humans, horses and rodents (Siciliani 2000; Colborn *et al.* 1991; Buwalda *et al.* 2012). We suggest that this question can potentially be resolved through a better understanding of physiological mechanisms and environmental factors driving GC variation.

Selye (1976) emphasised that homeostatic challenges of any kind lead to increases in body demands, loosely defined as *"the rate at which we live at any one moment"*. Activation of the Hypothalamus-Pituitary-Adrenal

(HPA) axis, represents a primary hormonal response to homeostatic challenges that, through the release of
GCs, mobilises the resources needed to fuel the current or anticipated rise in energy expenditure (current
and anticipatory responses) or recover from an immediate threat that induced an unanticipated increase in
energy expenditure (reactive response) (Herman *et al.* 2016, McEwen & Wingfield 2003). In this context, GCs
are involved in the metabolism of most types of energy reserves, modulating glucose, fat and protein
metabolism in liver, skeletal muscle, and other target tissues (Box 1). Allusions to the energetic role of GCs
and their tight link to energy expenditure are common in physiological and ecological studies, especially
those using approaches which underline the adaptive function of GC responses (McEwen & Wingfield 2003;
Landys et al. 2006; Romero *et al.* 2009; Deviche *et al.* 2017), but the extent to which GC variation can be
quantitatively explained as facilitator of variation in energy expenditure has rarely been addressed (but see
Jimeno *et al.* 2018, Malkoc *et al.* 2021 supplementary information). We here schematically review the role of
GCs in energy metabolism (Box 1), and investigate this link quantitatively through meta-analysis.

Revisiting the associations between GCs and metabolic rate. A meta-analytic approach

Although previous evidence supports the link between energy expenditure and GC secretion (e.g. Koolhaas
et al. 2011; Sapolsky *et al.* 2000; Beerling *et al.* 2011; Buwalda *et al.* 2012; Jimeno *et al.* 2018; Malkoc *et al.*
2021 supplementary information), the qualitative importance of both this association and the underlying
processes to explain GC variation remains unexplored; this is however fundamental towards accurately
interpreting GC variation. We here test whether changes in energetic demands are associated with variation
in GC levels. Specifically, we (i) use a meta-analytic approach to test whether experimental manipulations
leading to increases in metabolic rate in endotherms also lead to an increase in GCs (qualitative approach).
We included only experimentally-induced increases of energy expenditure to avoid potential masking effects
of anticipatory responses or delayed effects of GCs. Because metabolic rate and GCs can fluctuate rapidly,
we targeted metabolic rate and GC measurements taken simultaneously or when animals could be assumed
to be in the same physiological state (e.g. within the same day and experimental treatment). We further
investigated (ii) whether the magnitude of the experimentally induced changes in metabolic rate and GCs
were correlated (quantitative approach) through meta-regression. Our predictions are that (i) increases in
metabolic rates are associated with increases in plasma GC concentrations, (ii) that changes in GCs are
proportional to induced changes in metabolic rate, and (iii) that the association between increases in
metabolic rate and GCs is independent of the treatment used to increase the metabolic rate.

BOX 1: Glucocorticoids and energy metabolism

We here consider GC regulation from the perspective of their role in fuelling metabolic rate. When metabolic rate is low, for example during periods of inactivity, circulating GCs are maintained at low levels and glucose release from fuel stores is released in the blood stream at a low rate matching the modest metabolic needs. (*permissive* actions; Sapolsky *et al.* 2000; Fig. 1). An increase in metabolic rate can be anticipated or unanticipated (GCs will exert *preparative* or *stimulating* actions, respectively; Sapolsky *et al.* 2000; Fig. 1), and acute or gradual. Unanticipated but gradual increases in metabolic rate will occur for example when thermoregulatory costs unexpectedly increase.

In both gradual and acute increases in metabolic rate, a main role of GCs is to increase circulating glucose at a rate matching the metabolic requirements through diverse mechanisms. Decreasing plasma glucose levels trigger a series of hormonal changes that promote a switch in energy usage. Together with a decrease of insulin, GCs are released into the circulation (Andrews & Walker 1999; Rosmond & Bjorntorp 2000) reducing anabolic insulin actions (Vegiopoulos & Hertzig 2007). Blood glucose level then increases, both by mobilization from existing stores, and by inhibition of further storage. GCs also inhibit glucose uptake and glycogen synthesis in the liver, redirecting resources to gluconeogenesis and glycogenolysis, along with glucagon and catecholamines as part of the most immediate acute response. Catecholamines act quickly and increase within seconds to induce the release of energy needed to fuel the response (Herman *et al.* 2016; Romero & Beattie 2021; Sapolsky *et al.* 2000). The GC response lags in time -as GCs are produced de novo at the adrenal and take minutes to be secreted— and lasts substantially longer, depending on active (feedback signalling) and passive (GC degradation) processes (Herman *et al.*, 2016; Fig. 1), enhancing and prolonging the increase in blood glucose (Nonogaki 2000; Romero & Beattie 2021), or recovering energy stores after a brief burst of activity. In addition, inhibition of peripheral glucose transport and utilization in response to GCs increases the availability for other tissues, such as the brain (reviewed in Sapolsky *et al.* 2000; Herman *et al.* 2016). GCs also act in other substrates, further increasing lipolysis by inducing hormone-sensitive lipase (Slavin *et al.* 1994), and reducing lipoprotein lipase activity in peripheral fat depots. They also promote pre-adipocyte differentiation, pro-lipogenic pathway activity, and cellular hypertrophy in central fat (Vegiopoulos & Hertzig 2007) as well as decreased thermogenesis in brown adipose tissue (Soumano *et al.* 2000). In various muscle types, GCs suppress protein synthesis while promoting protein degradation and amino acid export. When the energetic and substrate requirements of the organism are further increased (e.g. during fasting or illness), muscle tissue (40% of total body mass) becomes a rich source of amino acids, which can be mobilized as substrates for energy generation, gluconeogenesis and protein synthesis (Kuo *et al.* 2013).

Given the existing evidence on the metabolic role of GCs, and the different time scales associated to the kind of response (anticipatory vs perceived) as well as to the hormones and substrates' physiological actions (see above), GC concentrations cannot be expected to always reflect the 'immediate' energy expenditure. However, we would expect changes in energetic demands to always require a GC response/input to meet the derived metabolic needs (Fig. 1). This prediction of a strong association between GCs and metabolic rate, however, is not well researched, and does not necessarily imply that one trait necessarily affects the other per se, as their interplay is likely to be shaped by the environmental or physiological context. Note further that we make no distinction between baseline and stress-induced GC-levels, and thereby in effect assume these to be points in a continuum from a metabolic perspective; a perspective supported by the monotonic effects of GCs on glucose uptake and fat depletion (Kattwinkel & Munck 1966; Dallman *et al.* 1993). Additionally, although we consider GCs to be regulated to meet energetic demands, we are aware GCs have many complex downstream effects at both baseline and stress-induced levels, besides energy-mobilization (Fig. 1).

Fig. 1: Schematic representation of the association between metabolic rate and plasma levels of glucocorticoids and glucose. Green

METHODS

Literature search

We reviewed the literature to identify empirical studies reporting measurements of both metabolic rate and plasma GCs. We compiled studies that met all following criteria: (1) Including an experimental manipulation of any kind leading to increases in metabolic rate which was quantified (i.e. being or not significant). Among these, we also included those studies reporting heart rate as a metabolic measure, as heart rate and metabolic rate are strongly correlated (Bevan *et al.* 1994; Bevan *et al.* 1995; Butler *et al.* 2004, Word *et al.* 2022). (2) Including measurements of natural GC concentrations in plasma (i.e. not exogenous or chemically induced –e.g. with ACTH or CRH). (3) measurements of GCs and metabolism had to be on the same individuals and measured in the same physiological state. The latter condition excludes, for example, studies with daily energy expenditure measurements combined with GCs measured at one time-point. Finally, we only included studies on endotherms (birds and mammals), because metabolic regulation differs strongly between endotherms and ectotherms.

We conducted a database search (Web of Science, 20/07/2021) to identify candidate studies, using the following two combinations of search terms: *"energy expenditure" AND (glucocorticoid OR cortisol OR corticosterone)* and *"metabolic rate" AND (glucocorticoid OR cortisol OR corticosterone)*. After the search, we consecutively selected articles after a) abstract review, b) full text review and c) data availability for effect size calculations. Using this approach, we identified a total of 14 studies that met all our criteria (see Table S1 for additional information on the number of studies obtained on each of the search steps). We also systematically checked the reference list of these 14 papers, which yielded an additional 7 papers. Thus, we included a total of 21 papers in our analyses, of which 12 were on birds, and 9 on mammals. Nine of the 22 papers included more than one experimental treatment, yielding a total of 35 effect sizes. For each of these studies, we extracted information on study species or metabolic and GC variables reported, among others (Table S2). Additionally, we recorded variation related to the experimental design, the variables that were quantified, and the type of treatment used: a) Before / after design: whether the experimental manipulation included a time effect (i.e. individuals served as their own control, being measured before and after the experimental manipulation); b) Experiment / control design: Whether the experiment accounted for within-individual variation (i.e. all individuals went through all experimental treatments); c) Whether metabolic rate or heart rate was the metabolic variable; and d) The type of treatment that induced an increase in metabolic rate (see below) (Table S2).

Effect size calculations

To estimate effect sizes of metabolism and GCs, we used the web-based effect size calculator Practical Meta-Analysis Effect Size Calculator (www.campbellcollaboration.org/escalc/html/EffectSizeCalculator-Home.php), following Lipsey & Wilson 2000 and Nakagawa & Cuthill 2007. We calculated standardized mean-difference effect sizes (Cohen's D), which we computed from means and standard deviations (19 studies) or t-test (2 studies). When metrics were presented graphically only, we extracted data from the figure(s) using the GetData Graph Digitizer software (http://getdata-graph-digitizer.com/). See Table S3 for details on data extraction and effect size calculations.

For each study, we compared the mean metabolic rate and level of plasma GCs of individuals in the
treatment group(s) to that of individuals in the control group. For studies in which treatment was
confounded with time, because pre-treatment measurements were used as control and compared with
measurements during treatment, the pre-treatment measure was used as control when calculating effect
sizes in studies where there was a single treatment. When studies with a before-after design included more
than one experimental treatment, the treatment yielding the lowest metabolism was taken as control for
the effect size calculations. Thus, confounding time with treatment was avoided whenever possible.

Statistical analyses

We conducted all meta-analyses using the *rma.mv* function from the *metafor* package (Viechtbauer 2010),
implemented in R (version 4.0.1, R Core Team 2020). Standard errors were used for the weigh factor. All
models contained a random intercept for study identity to account for inclusion of multiple experimental
treatments or groups from the same study. Most species were used in a single study, and we therefore did
not include species as a random effect in addition to study identity. The number of species was however
insufficient to reliably estimate phylogenetic effects, we therefore limited the analysis in this respect with a
comparison between birds and mammals (see below). The dependent variable was either the metabolic rate
or the GC effect size. One model was fitted with the metabolic rate effect size as a dependent variable, to
estimate the average effect on metabolic rate across all studies in the analyses. All other models had the GC
effect size as dependent variable, and metabolic rate effect size as a moderator. Distribution of metabolic
rate effect sizes was skewed which was resolved by ln-transformation, which yielded a better fit when
compared to a model using the linear term (evaluated using AIC, see results for details). Our first GC model
contained only the metabolic rate effect size as a fixed independent variable. This model provides a
qualitative test of whether GC levels increase when metabolic rate is increased and tests prediction (i) by
providing an estimate of the intercept, which represents the average GC effect size because we mean
centered the ln-transformed metabolic rate effect size (Schielzeth 2010). The same model tests prediction
(ii) whether the GC effect increases with an increasing metabolic rate effect size, which will be expressed in a
significant regression coefficient of the metabolic rate effect size.

Following the model with which we tested our main predictions, we ran additional models to test for the
effects on GC effect sizes of (a) taxa (birds vs. mammals), two design effects, namely (b) before / after effect,
and (c) experiment / control effect, (d) metabolic variable measured (metabolic rate or heart rate), (e) type
of treatment (categorized in climate, psychological, or other). This last factor tests our prediction iii. We
included these variables as modulators in the analysis, as well as the two-way interactions of these factors
with the metabolic rate effect size. All factors were coded as 1 (bird / no before − after effect / no
experiment-control effect / metabolic rate) or 2 (mammal / before-after effect / experiment-control effect /
heart rate), and then mean-centered. Treatment type was categorized as 1 (climate), 2 (psychological), or 3
(others). We compared models with vs. without these additional variables using Akaike's Information
Criterion, with correction for small sample sizes (AICc, Akaike, 1974)) for which a change in AICc of 2 is
considered significant (Burnham, Anderson & Huyvaert 2011). Models within ⊠AICc < 2 were considered
best fitting models, and we further explored the effects of the main predictors when present in these top
models. To rule out publication bias effects (i.e. regression test for funnel plot asymmetry), we included a

weighing variable (square root of the sample size) as moderator in the models, as Egger's test is not a
reliable test of funnel plot asymmetry in multilevel models. Variable effects and results remained
quantitatively very similar and qualitatively unchanged.

RESULTS

Among the studies selected for inclusion in the analysis, the treatment effect size on metabolic rate (MR)
was on average 1.85 ± 0.87 (Fig. S1). In accordance with prediction, effects on GCs were positive in the
majority of cases (32/35; Fig. 2), and consequently the overall average effect size deviated significantly from
zero, with the average GC effect size estimated at 0.73 ± 0.11 (Table 1). There was a strong association
between MR effect sizes and GC effect sizes (Table 1, Fig. 3), thus confirming prediction ii. It is further worth
noting that the residual heterogeneity did not exceed the level expected by chance (Table 1). MR Cohen's D
was Ln-transformed (see methods) to normalize the distribution (Fig. S2), and AICc of models including Ln
MR were significantly lower when compared to models including untransformed MR Cohen's D (AICc = 70.58
vs 74.97, respectively).

The association between MR and GC effect sizes remained statistically significant when adding Taxa, before /
after, experiment / control effect, metabolic variable or treatment type one by one to the model.
Furthermore, none of these variables had a significant effect on GC effect size, nor did the association
between MR and GC effect sizes depend on those factors (i.e. interactions between these variables and MR
effect sizes were always non-significant; Table 2, S4). The latter result confirms prediction iii. Given that none
of these effects significantly improved the model, the final model when removing all factors was the one
including MR effect size as only predictor of GC effect size (Table 1). Despite these modulators being non-
significant, the associations were in the expected directions, with studies including within-individual
variation (i.e. experiment / control effect), and not including a before / after effect reporting higher GC
effect sizes (Table S4, Fig. S3).

Table 1. Meta regression model testing the association between metabolic rate (MR) effect sizes and
glucocorticoid effect sizes.

	Estimate	s.e.	Z	P	95% C.I..
Intercept	0.72	0.11	6.40	<0.0001	0.50-0.95
MR effect size (ln)	0.31	0.10	2.94	0.003	0.10-0.51

Variance components: Study.ID (Sigma^2)– Estimate = 0.00, sqrt = 0.00, n= 21

Residual heterogeneity: QE(df = 33) = 28.40, p = 0.70

Test of moderators: QM(df = 1) = 8.64, p= 0.003

212   **Table 2.** Table showing the main effects of all variables considered (Metabolic Rate, Taxa, Time effect,
213   Within-individual variation, Metabolic variable, and Treatment Type) to modulate glucocorticoid effect sizes
214   across studies. Full models are shown in Table S4.

Variable	P
MR effect size (ln)	**0.003**
Taxa	0.63
Before / after	0.49
Experiment / control	0.68
Metabolic variable	0.94
Treatment type	
Treat. 2	0.93
Treat. 3	0.60

219   **Fig. 2**: Forest plot showing the glucocorticoid (GC) effect sizes (Cohen's D ± 95% CI) associated to
220   experimental manipulations of metabolic rate, grouped by treatment group and study. Area of squares is
221   proportional to the experiment sample size (1/s.e.).

Fig 3. Glucocorticoid effect size (Cohen´s D) increases with increasing metabolic rate effect size similarly in studies of mammals (open circles) and birds (closed circles). Area of dots is proportional to the experiment sample size (i.e. square root of the number of individuals in which GCs were measured).

DISCUSSION

Finding a consistent interpretation of GC variation has proven challenging, and to this end we presented a simplified framework focusing on the interplay between energy metabolism and GCs (Box 1). Based on this framework, we made three predictions that we tested through a meta-analysis of studies in endotherms in which metabolic rate was manipulated and GCs were measured at the same time. The analysis confirmed our predictions, showing that experimental manipulations that increased metabolic rate induced a proportional increase in GCs (Fig.3), and our interpretation of this effect is that GC secretion facilitated increases in MR. This association indicates that fluctuations in energy turnover are a key factor driving variation in GC levels. From this perspective, the many downstream effects of GCs (e.g. down regulation of immune function and reproduction, enhanced learning; McEwen & Wingfield 2003; Sapolsky *et al.* 2000) may be interpreted as allocation adjustments to the metabolic level at which organisms operate.

The effect of metabolic rate on GC levels was independent of the type of manipulation used to increase metabolic rate, confirming our third prediction. Note however that confirmation of this prediction relied on the absence of a significant effect, and absence of evidence is not evidence of absence. However, the residual heterogeneity of our final model did not deviate from a level expected due to sampling variance, providing additional support for our third prediction.

We restricted the meta-analysis to experimental studies, and expect the association between MR and GCs to
be less evident in a more natural context. Associations between GCs and MR will be most evident when
animals are maintained at different but stable levels of metabolic rate, because then the rate at which
tissues are fuelled is likely to be in equilibrium with the metabolic needs. While equilibrium conditions can
be created in laboratory studies, conditions will usually be more variable in the wild. When metabolic rate
fluctuates, e.g. due to short term variation in activity, GC variation will track metabolic rate fluctuations, but
with a time lag (Box 1), thus adding complexity to the MR / GC association and its detectability. Furthermore,
experiments yield estimates of associations within the average individual in the study, while data collected in
a natural context usually rely on variation between individuals (but see Malkoc *et al.* 2022). Associations
between individuals will be less strong than associations within individuals due to individual variation in GC
levels and GC reactivity (e.g. Liu *et al.* 1997; Weaver *et al.* 2004; Yehuda *et al.* 2014; Taff et al 2018, 2022).
Between subject variation is likely to be even larger on an interspecific level, and in line with this expectation
the comparative evidence on the MR / GC association is mixed. While a strong positive MR / GC correlation
was reported for mammals (Haase et al 2016), Francis et al (2018) found no consistent GC / MR association
in birds and other tetrapod taxa. However, recent comparative analyses showed avian GC variation to be
positively correlated with estimated thermoregulatory costs (Rubalcaba & Jimeno 2022), and GC variation in
lizards to be positively correlated with body temperature, which directly influences metabolic rate in
ectotherms (Rubalcaba & Jimeno *in press*). The contrast between the findings may be due to the MR and GC
data not always being collected on individuals in a comparable state. We emphasize therefore the
importance of measuring metabolic rate and GCs when animals are in the same state, preferably by
measuring both variables at the same time.

GCs increased in the studies included in our meta-analysis in response to an induced increase in MR, but GCs
can also increase in response to an *anticipated* increase in MR (Box 1). For example, the early morning GC
increase in humans, known as the 'cortisol awakening response' (Fries et al 2009) can be interpreted as
preparation for an increase in MR – indeed "early birds" show higher GC-levels than "night owls" in the hour
after awakening (Kudielka et al 2006). Likewise, GC levels increase in athletes preceding competition (van
Paridon et al 2017), although separating effects of psychological stress from anticipated metabolic needs is
difficult in this context. Experiments in which animals are trained to anticipate an increase in MR to
investigate whether this generates an anticipatory increase in GCs would be an interesting additional test of
the framework laid out in Box 1.

Given that GC levels are often assumed to provide information on organismal 'stress' and welfare, the
question arises whether the observed pattern can be the consequence of effects on psychological stress
instead of metabolic rate. This question arises because manipulations of energy expenditure are always
'indirect', in that an external treatment is used to induce an increase in metabolic rate, as opposed to a
direct manipulation of metabolic rate, and this leaves room for other factors to cause the observed effects.
While we acknowledge that it is not possible to demonstrate conclusively that a process is not happening,
we consider it unlikely that 'stress effects' explain our findings. Firstly, because the way metabolic rate was
manipulated varied widely between studies but manipulation type had no discernible effect on the MR / GC
association. This was also the conclusion of one of the studies included in our meta-analysis, designed
specifically to compare MR / GC associations between two MR increasing treatments, ambient temperature

and noise as psychological stressor (Jimeno et al 2018). Secondly, the finding that the GC increase was
proportional to the increase in MR can only be explained by psychological stress when the induced
psychological stress was proportional to the induced MR. Thirdly, the pattern is consistent with what is
known of the functional consequences of GC variation (Box 1). Lastly, diverse non-injurious psychological
stressors increase metabolic rate in humans (Sawai *et al.* 2007, Balanos *et al.* 2010, Carroll *et al.* 2009),
mammals (Harris *et al.* 2006; mild and unpredictable chronic stress; García-Díaz *et al.* 2007) and birds
(Jimeno *et al.* 2018), explaining why GCs generally increase in response to a stressor. We conclude therefore
that while a causal link between MR and GCs is not the only possible explanation of our findings, we argue it
to be the most parsimonious explanation. Direct manipulations of MR are challenging, but could confirm or
reject this explanation, and may for example be achieved using thyroid hormones, which have been shown
to affect MR (Moreno *et al.* 2002, Kim 2008). Additionally, it would be informative to investigate whether
energy turnover immediately before blood sampling is a predictor of GC levels, as we would predict on the
basis of the interpretation of our findings. Increasing the use of devices and techniques that monitor energy
expenditure or its proxies (e.g. accelerometers) may be a way to increase our understanding of the
generality of the GC-MR association.
298
Authors that assumed GC levels to be a proxy of physiological stress have struggled with the interpretation
of findings such as the mixed results with respect to fitness consequences of GC-variation. Our findings offer
a way to interpret such variation: GCs are regulated with respect to their role in facilitating energy
metabolism, and we encourage researchers to approach and interpret findings from this perspective. For
example, a positive association between GCs and reproductive success may indicate that individuals that are
able to sustain high metabolic rates attain higher fitness (e.g. Bauch et al, 2016), while a negative association
indicates the opposite effect (e.g. Ouyang et al 2013; see Atema *et al.*, 2021, for a more general discussion of
this specific contrast). Given that GCs have many other downstream effects (Sapolsky *et al.* 2000), for
example supressing immune function (Cain & Cidlowski 2017), and growth (Allen 1996), this may seem an
overly simplistic approach. However, in this framework, downstream effects of GCs can be understood as
responses to a system level readout of the current level of energy metabolism, with high levels affecting the
allocation of energy to different energy demanding processes. In this view, the link between GCs and energy
demanding processes is asymmetric, in the sense that GCs affect energy allocation to e.g. growth, but there
is no direct feedback from growth to GC levels. In conclusion, whereas GCs are widely seen as 'stress
hormones', we offer a different interpretation and question whether GC variation reveals any physiological
stress beyond fluctuations in energy expenditure.

FUNDING

BJ was funded by grant FJC2019-039748-I of MCIN/AEI /10.130 39/501100011033.

REFERENCES

References marked with an asterisk were included in the meta-analysis but not otherwise cited in the paper.

Akaike, H. (1974). A New Look at the Statistical Model Identification. *IEEE Transactions on Automatic Control*,
19(6), 716–723. doi: 10.1109/TAC.1974.1100705

Allen, D. B. (1996). Growth suppression by glucocorticoid therapy. *Endocrinology and metabolism clinics of*
North America, *25*(3), 699-717.

Andrews, R. C., & Walker, B. R. (1999). Glucocorticoids and insulin resistance: old hormones, new targets.
Clinical Science, *96*(5), 513–523. doi: 10.1042/CS0960513

Atema, E., van Noordwijk, A. J., & Verhulst, S. (2021). Telomere dynamics in relation to experimentally
increased locomotion costs and fitness in great tits. *Molecular Ecology,* doi.org/10.1111/mec.16162.

Balanos, G. M., Phillips, A. C., Frenneaux, M. P., McIntyre, D., Lykidis, C., Griffin, H. S., & Carroll, D. (2010).
Metabolically exaggerated cardiac reactions to acute psychological stress: the effects of resting blood
pressure status and possible underlying mechanisms. *Biological Psychology*, *85*(1), 104-111.

Bauch, C., Riechert, J., Verhulst, S., & Becker, P. H. (2016). Telomere length reflects reproductive effort
indicated by corticosterone levels in a long-lived seabird. *Molecular Ecology*, *25*(22), 5785-5794.

Beerling, W., Koolhaas, J. M., Ahnaou, A., Bouwknecht, J. A., de Boer, S. F., Meerlo, P., & Drinkenburg, W. H.
I. M. (2011). Physiological and hormonal responses to novelty exposure in rats are mainly related to
ongoing behavioral activity. Physiology & behavior, 103(3-4), 412-420.

Bevan, R. M., Woakes, A. J., Butler, P. J., & Boyd, I. L. (1994). The use of heart rate to estimate oxygen
consumption of free-ranging black-browed albatrosses Diomedea melanophrys. *The Journal of*
experimental biology, *193*(1), 119-137.

Bevan, R. M., Speakman, J. R., & Butler, P. J. (1995). Daily energy expenditure of tufted ducks: a comparison
between indirect calorimetry, doubly labelled water and heart rate. *Functional Ecology*, 40-47.

Bonier, F., Martin, P. R., Moore, I. T., & Wingfield, J. C. (2009). Do baseline glucocorticoids predict
fitness?. *Trends in ecology & Evolution*, *24*(11), 634-642.

* Booth-Mclean, M. E., Schwartzkopf-Genswein, K. S., Brown, F. A., Holmes, C. L., Schaefer, A. L., McAllister,
T. A., & Mears, G. J. (2007). Physiological and behavioural responses to short-haul transport by stock
trailer in finished steers. Canadian journal of animal science, 87(3), 291-297.

* de Bruijn, R., & Romero, L. M. (2011). Behavioral and physiological responses of wild-caught European
starlings (Sturnus vulgaris) to a minor, rapid change in ambient temperature. Comparative Biochemistry
and Physiology Part A: Molecular & Integrative Physiology, 160(2), 260-266.

* de Bruijn, R., & Romero, L. M. (2013). Artificial rain and cold wind act as stressors to captive molting and
non-molting European starlings (Sturnus vulgaris). Comparative Biochemistry and Physiology Part A:
Molecular & Integrative Physiology, 164(3), 512-519.

Burnham, K. P., Anderson, D. R., & Huyvaert, K. P. (2011). AIC model selection and multimodel inference in
behavioral ecology: Some background, observations, and comparisons. *Behavioral Ecology and*
Sociobiology, *65*(1), 23–35. doi: 10.1007/S00265-010-1029-6/FIGURES/3

Busch, D. S., & Hayward, L. S. (2009). Stress in a conservation context: a discussion of glucocorticoid actions
and how levels change with conservation-relevant variables. *Biological Conservation*, *142*(12), 2844-

2853.

Burton, T., Killen, S. S., Armstrong, J. D., & Metcalfe, N. B. (2011). What causes intraspecific variation in
359   resting metabolic rate and what are its ecological consequences?. *Proceedings of the Royal Society B:*
360   *Biological Sciences, 278*(1724), 3465-3473.

361   Butler, P. J., Green, J. A., Boyd, I. L., & Speakman, J. R. (2004). Measuring metabolic rate in the field: the pros
362   and cons of the doubly labelled water and heart rate methods. *Functional ecology, 18*(2), 168-183.

363   Buwalda, B., Scholte, J., de Boer, S. F., Coppens, C. M., & Koolhaas, J. M. (2012). The acute glucocorticoid
364   stress response does not differentiate between rewarding and aversive social stimuli in rats. Hormones
365   and behavior, 61(2), 218-226.

366   Cain, D. W., & Cidlowski, J. A. (2017). Immune regulation by corticosteroids. *Nat Rev Immunol, 17*, 233-247.

367   Carroll, D., Phillips, A. C., & Balanos, G. M. (2009). Metabolically exaggerated cardiac reactions to acute
368   psychological stress revisited. *Psychophysiology, 46*(2), 270-275.

369   * Celi, F. S., Brychta, R. J., Linderman, J. D., Butler, P. W., Alberobello, A. T., Smith, S., ... & Chen, K. Y. (2010).
370   Minimal changes in environmental temperature result in a significant increase in energy expenditure
371   and changes in the hormonal homeostasis in healthy adults. European journal of
372   endocrinology/European Federation of Endocrine Societies, 163(6), 863.

373   * Cohen, A., Hau, M., & Wikelski, M. (2008). Stress, metabolism, and antioxidants in two wild passerine bird
374   species. Physiological and Biochemical Zoology, 81(4), 463-472.

375   Colborn, D. R., Thompson Jr, D. L., Roth, T. L., Capehart, J. S., & White, K. L. (1991). Responses of cortisol and
376   prolactin to sexual excitement and stress in stallions and geldings. *Journal of Animal Science, 69*(6),
377   2556-2562.

378   * Cyr, N. E., Wikelski, M., & Romero, L. M. (2008). Increased energy expenditure but decreased stress
379   responsiveness during molt. Physiological and Biochemical Zoology, 81(4), 452-462.

380   Dallman, M. F., Strack, A. M., Akana, S. F., Bradbury, M. J., Hanson, E. S., Scribner, K. A., & Smith, M. (1993).
381   Feast and famine: critical role of glucocorticoids with insulin in daily energy flow. *Frontiers in*
382   *neuroendocrinology, 14*(4), 303-347.

383   Deviche, P., Bittner, S., Gao, S., & Valle, S. (2017). Roles and mechanistic bases of glucocorticoid regulation of
384   avian reproduction. *Integrative and comparative biology, 57*(6), 1184-1193.

385   Francis, C. D., Donald, J. W., Fuxjager, M. J., Goymann, W., Hau, M., Husak, J. F., ... & Downs, C. J. (2018).
386   Metabolic scaling of stress hormones in vertebrates. *Integrative and comparative biology, 58*(4), 729-
387   738.

388   * Frank, S. M., Higgins, M. S., Fleisher, L. A., Sitzmann, J. V., Raff, H., & Breslow, M. J. (1997). Adrenergic,
389   respiratory, and cardiovascular effects of core cooling in humans. American Journal of Physiology-
390   Regulatory, Integrative and Comparative Physiology, 272(2), R557-R562.

391   Fries, E., Dettenborn, L., & Kirschbaum, C. (2009). The cortisol awakening response (CAR): facts and future
392   directions. *International journal of Psychophysiology, 72*(1), 67-73.

393   García-Díaz, D. F., Campion, J., Milagro, F. I., Lomba, A., Marzo, F., & Martínez, J. A. (2007). Chronic mild
394   stress induces variations in locomotive behavior and metabolic rates in high fat fed rats. *Journal of*
395   *physiology and biochemistry, 63*(4), 337-346.

Haase, C. G., Long, A. K., & Gillooly, J. F. (2016). Energetics of stress: linking plasma cortisol levels to

metabolic rate in mammals. *Biology letters*, *12*(1), 20150867.

Harris, R. B., Palmondon, J., Leshin, S., Flatt, W. P., & Richard, D. (2006). Chronic disruption of body weight
but not of stress peptides or receptors in rats exposed to repeated restraint stress. *Hormones and*
behavior, *49*(5), 615-625.

* Harlow, H. J., Thorne, E. T., Williams, E. S., Belden, E. L., & Gern, W. A. (1987). Adrenal responsiveness in
domestic sheep (Ovis aries) to acute and chronic stressors as predicted by remote monitoring of
cardiac frequency. Canadian journal of zoology, 65(8), 2021-2027.

Herman, J. P., McKlveen, J. M., Ghosal, S., Kopp, B., Wulsin, A., Makinson, R., … Myers, B. (2016). Regulation
of the Hypothalamic-Pituitary-Adrenocortical Stress Response. In *Comprehensive Physiology* (Vol. 6, pp.
603–621). doi: 10.1002/cphy.c150015

Herman, J. P. (2022). The neuroendocrinology of stress: Glucocorticoid signaling mechanisms.
Psychoneuroendocrinology, 105641.

* Hipolide, D. C., Suchecki, D., de Carvalho Pinto, A. P., Chiconelli Faria, E., Tufik, S., & Luz, J. (2006).
Paradoxical sleep deprivation and sleep recovery: effects on the hypothalamic–pituitary–adrenal axis
activity, energy balance and body composition of rats. Journal of neuroendocrinology, 18(4), 231-238.

Jessop, T. S., Lane, M. L., Teasdale, L., Stuart-Fox, D., Wilson, R. S., Careau, V., & Moore, I. T. (2016).
Multiscale evaluation of thermal dependence in the glucocorticoid response of vertebrates. *The*
American Naturalist, *188*(3), 342-356.

Jimeno, B., Hau, M., & Verhulst, S. (2018). Corticosterone levels reflect variation in metabolic rate,
independent of 'stress.' *Scientific Reports*, *8*(1). doi: 10.1038/s41598-018-31258-z

Kattwinkel, J., & Munck, A. (1966). Activities in vitro of glucocorticoids and related steroids on glucose
uptake by rat thymus cell suspensions. *Endocrinology*, *79*(2), 387-390.

Kim, B. (2008). Thyroid hormone as a determinant of energy expenditure and the basal metabolic
rate. *Thyroid*, *18*(2), 141-144.

Kudielka, B. M., Federenko, I. S., Hellhammer, D. H., & Wüst, S. (2006). Morningness and eveningness: the
free cortisol rise after awakening in "early birds" and "night owls". *Biological psychology*, *72*(2), 141-
146.

* Kaciuba-Uscilko, H., Kruk, B., Szczypaczewska, M., Opaszowski, B., Stupnicka, E., Bicz, B., & Nazar, K. (1992).
Metabolic, body temperature and hormonal responses to repeated periods of prolonged cycle-
ergometer exercise in men. *European journal of applied physiology and occupational physiology*, 64(1),
26-31.

* Keselman, B., Vergara, M., Nyberg, S., & Nystrom, F. H. (2017). A randomized cross-over study of the acute
effects of running 5 km on glucose, insulin, metabolic rate, cortisol and Troponin T. PloS one, 12(6),
e0179401.

* Kleist, B., Wahrburg, U., Stehle, P., Schomaker, R., Greiwing, A., Stoffel-Wagner, B., & Egert, S. (2017).
Moderate walking enhances the effects of an energy-restricted diet on fat mass loss and serum insulin
in overweight and obese adults in a 12-week randomized controlled trial. The Journal of
nutrition, 147(10), 1875-1884.

Koolhaas, J. M., Bartolomucci, A., Buwalda, B., de Boer, S. F., Flügge, G., Korte, S. M., … & Fuchs, E. (2011).
Stress revisited: a critical evaluation of the stress concept. *Neuroscience & Biobehavioral*
Reviews, *35*(5), 1291-1301.

Kuo, T., Harris, C. A., & Wang, J. C. (2013). Metabolic functions of glucocorticoid receptor in skeletal
muscle. *Molecular and cellular endocrinology*, *380*(1-2), 79-88.

Landys, M. M., Ramenofsky, M., & Wingfield, J. C. (2006). Actions of glucocorticoids at a seasonal baseline as
compared to stress-related levels in the regulation of periodic life processes. *General and comparative*
endocrinology, *148*(2), 132-149.

Lipsey, M. W., & Wilson, D. B. (2001). *Practical meta-analysis*. SAGE publications, Inc.

Liu, D., Diorio, J., Tannenbaum, B., Caldji, C., Francis, D., Freedman, A., Meaney, M.J., 1997. Maternal care,
hippocampal glucocorticoid receptors, and hypothalamic-pituitary-adrenal responses to stress. Science
277 (5332), 1659–1662. doi: 10.1126/science.277.5332.1659.

MacDougall-Shackleton, S. A., Bonier, F., Romero, L. M., & Moore, I. T. (2019). Glucocorticoids and "Stress"
Are Not Synonymous. *Integrative Organismal Biology*, *1*(1). doi: 10.1093/IOB/OBZ017

Madliger, C. L., & Love, O. P. (2014). The need for a predictive, context-dependent approach to the
application of stress hormones in conservation. *Conservation Biology*, *28*(1), 283-287.

Madliger, C. L., Semeniuk, C. A. D., Harris, C. M., & Love, O. P. (2015). Assessing baseline stress physiology as
an integrator of environmental quality in a wild avian population: Implications for use as a conservation
biomarker. *Biological Conservation*, *192*, 409–417. doi: 10.1016/j.biocon.2015.10.021

Malkoc, K., Casagrande, S., & Hau, M. (2021). Inferring whole-organism metabolic rate from red blood cells
in birds. *Frontiers in Physiology*, *12*.

Malkoc, K., Mentesana, L., Casagrande, S., & Hau, M. (2022). Quantifying glucocorticoid plasticity using
reaction norm approaches: There still is so much to discover!. *Integrative and Comparative*
Biology, *62*(1), 58-70.

McCormick, S. D., & Romero, L. M. (2017). Conservation Endocrinology. *BioScience*, *67*(5), 429–442. doi:
10.1093/BIOSCI/BIX026

McEwen, B. S., & Wingfield, J. C. (2003). The concept of allostasis in biology and biomedicine. *Hormones and*
behavior, *43*(1), 2-15.

Moreno, M., Lombardi, A., Beneduce, L., Silvestri, E., Pinna, G., Goglia, F., & Lanni, A. (2002). Are the effects
of T3 on resting metabolic rate in euthyroid rats entirely caused by T3 itself?. *Endocrinology*, *143*(2),
504-510.

Nakagawa, S., & Cuthill, I. C. (2007). Effect size, confidence interval and statistical significance: a practical
guide for biologists. *Biological Reviews*, *82*(4), 591–605. doi: 10.1111/J.1469-185X.2007.00027.X

* Nephew, B. C., & Romero, L. M. (2003). Behavioral, physiological, and endocrine responses of starlings to
acute increases in density. Hormones and Behavior, 44(3), 222-232.

* Nephew, B. C., Kahn, S. A., & Romero, L. M. (2003). Heart rate and behavior are regulated independently of
corticosterone following diverse acute stressors. General and comparative endocrinology, 133(2), 173-
180.

Nonogaki, K. (2000). New insights into sympathetic regulation of glucose and fat
metabolism. *Diabetologia*, *43*(5), 533-549.

Ouyang, J. Q., Sharp, P., Quetting, M., & Hau, M. (2013). Endocrine phenotype, reproductive success and
survival in the great tit, Parus major. *Journal of Evolutionary Biology*, *26*(9), 1988-1998.

van Paridon, K. N., Timmis, M. A., Nevison, C. M., & Bristow, M. (2017). The anticipatory stress response to
sport competition; a systematic review with meta-analysis of cortisol reactivity. *BMJ open sport &*
exercise medicine, *3*(1), e000261.

Patterson, S. H., Hahn, T. P., Cornelius, J. M., & Breuner, C. W. (2014). Natural selection and glucocorticoid
physiology. *Journal of Evolutionary Biology*, *27*(2), 259–274. doi: 10.1111/jeb.12286

* Peake, J. M., Tan, S. J., Markworth, J. F., Broadbent, J. A., Skinner, T. L., & Cameron-Smith, D. (2014).
Metabolic and hormonal responses to isoenergetic high-intensity interval exercise and continuous
moderate-intensity exercise. American Journal of Physiology-Endocrinology and Metabolism, 307(7),
E539-E552.

Petrullo, L., Delaney, D., Boutin, S., McAdam, A. G., Lane, J. E., Boonstra, R., ... & Dantzer, B. (2022). The
glucocorticoid response to environmental change is not specific to agents of natural selection in wild
red squirrels. *Hormones and Behavior*, *146*, 105262.

R Core Team (2020). R: A language and environment for statistical computing. R Foundation for Statistical
Computing, Vienna, Austria. URL https://www.R-project.org/.

Romero, L. M. (2004). Physiological stress in ecology: lessons from biomedical research. *Trends in ecology &*
evolution, *19*(5), 249-255.

Romero, L. M., Dickens, M. J., & Cyr, N. E. (2009). The reactive scope model—a new model integrating
homeostasis, allostasis, and stress. *Hormones and behavior*, *55*(3), 375-389.

Romero, L. M., & Beattie, U. K. (2021). Common myths of glucocorticoid function in ecology and
conservation. *Journal of Experimental Zoology Part A: Ecological and Integrative Physiology*. doi:
10.1002/JEZ.2459

Romero, L. M., & Fairhurst, G. D. (2016). Measuring corticosterone in feathers: strengths, limitations, and
suggestions for the future. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative*
Physiology, *202*, 112-122.

Rosmond, R. A., & Björntorp, P. (2000). The hypothalamic–pituitary–adrenal axis activity as a predictor of
cardiovascular disease, type 2 diabetes and stroke. *Journal of internal medicine*, *247*(2), 188-197.

Rubalcaba, J. G., & Jimeno, B. Body temperature and activity patterns modulate glucocorticoid levels across
lizard species. A macrophysiological approach. Frontiers in Ecology and Evolution, *in press*.

Rubalcaba, J. G., & Jimeno, B. (2022). Biophysical models unravel associations between glucocorticoids and
thermoregulatory costs across avian species. *Functional Ecology*, *36*(1), 64-72.

Samra, J. S., Clark, M. L., Humphreys, S. M., MacDonald, I. A., Bannister, P. A., & Frayn, K. N. (1998). Effects of
physiological hypercortisolemia on the regulation of lipolysis in subcutaneous adipose tissue. *The*
Journal of Clinical Endocrinology & Metabolism, *83*(2), 626-631.

Sapolsky, R. M., Romero, L. M., & Munck, A. U. (2000). How do glucocorticoids influence stress responses?
Integrating permissive, suppressive, stimulatory, and preparative actions. *Endocrine reviews*, *21*(1), 55-
89.

Sawai, A., Ohshige, K., Yamasue, K., Hayashi, T., & Tochikubo, O. (2007). Influence of mental stress on
cardiovascular function as evaluated by changes in energy expenditure. *Hypertension Research*, *30*(11),
1019-1027.

Schielzeth, H. (2010). Simple means to improve the interpretability of regression coefficients. *Methods in*

Ecology and Evolution, *1*(2), 103–113. doi: 10.1111/J.2041-210X.2010.00012.X

Schoenle, L. A., Zimmer, C., Miller, E. T., & Vitousek, M. N. (2021). Does variation in glucocorticoid
concentrations predict fitness? A phylogenetic meta-analysis. *General and Comparative Endocrinology*,
Vol. 300, p. 113611. doi: 10.1016/j.ygcen.2020.113611

Selye, H. (1950). Stress and the general adaptation syndrome. *British medical journal*, *1*(4667), 1383.

Selye, H. (1976). Stress without distress. In *Psychopathology of human adaptation* (pp. 137-146). Springer,
Boston, MA.

Siciliani, J. L. 2000. "The Effect of Sexual Activity on Secretory Immunoglobulin a and Cortisol Levels in
Married Couples." Dissertation. University of Tennessee Knoxville.

Slavin, B. G., Ong, J. M., & Kern, P. A. (1994). Hormonal regulation of hormone-sensitive lipase activity and
mRNA levels in isolated rat adipocytes. *Journal of lipid research*, *35*(9), 1535-1541.

Soumano, K., Desbiens, S., Rabelo, R., Bakopanos, E., Camirand, A., & Silva, J. E. (2000). Glucocorticoids
inhibit the transcriptional response of the uncoupling protein-1 gene to adrenergic stimulation in a
brown adipose cell line. *Molecular and cellular endocrinology*, *165*(1-2), 7-15.

* Šrámek, P., Šimecková, M., Janský, L., Šavlíková, J., & Vybíral, S. (2000). Human physiological responses to
immersion into water of different temperatures. European journal of applied physiology, 81(5), 436-
442.

Taff, C. C., Zimmer, C., Ryan, T. A., van Oordt, D. C., Aborn, D. A., Ardia, D. R., … & Vitousek, M. N. (2022).
Individual variation in natural or manipulated corticosterone does not covary with circulating glucose in
a wild bird. *Journal of Experimental Biology*, *225*(4), jeb243262.

Taff, C. C., Schoenle, L. A., & Vitousek, M. N. (2018). The repeatability of glucocorticoids: a review and meta-
analysis. *General and Comparative Endocrinology*, *260*, 136-145.

Vegiopoulos, A., & Herzig, S. (2007). Glucocorticoids, metabolism and metabolic diseases. *Molecular and*
cellular endocrinology, *275*(1-2), 43-61.

Versteegh, M. A., Helm, B., Dingemanse, N. J., & Tieleman, B. I. (2008). Repeatability and individual
correlates of basal metabolic rate and total evaporative water loss in birds: a case study in European
stonechats. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative*
Physiology, *150*(4), 452-457.

Viechtbauer, W. (2010). Conducting Meta-Analyses in R with the metafor Package. *Journal of Statistical*
Software, *36*(3), 1–48. doi: 10.18637/JSS.V036.I03

Weaver, I.C.G., Cervoni, N., Champagne, F.A., D'Alessio, A.C., Sharma, S., Seckl, J.R., Meaney, M.J., 2004.
Epigenetic programming by maternal behavior. Nat. Neurosci. 7 (8), 847–854.

* Wikelski, M., Lynn, S., Breuner, J. C., Wingfield, J. C., & Kenagy, G. J. (1999). Energy metabolism,
testosterone and corticosterone in white-crowned sparrows. *Journal of Comparative Physiology*
A, 185(5), 463-470.

Word, K.R., Austin, S.H. & Wingfield JC (2022) Allostatic Load in Gambel's White Crowned Sparrow,
Zonotrichia leucophrys gambelii: Relationships With Glucocorticoids. *Frontiers in Ecology and Evolution*
10:855152

* Xu, X. M., Chi, Q. S., Cao, J., & Zhao, Z. J. (2018). The effect of aggression I: the increases of metabolic cost
and mobilization of fat reserves in male striped hamsters. Hormones and Behavior, 98, 55-62.

Yehuda, R., Daskalakis, N.P., Lehrner, A., Desarnaud, F., Bader, H.N., Makotkine, I., Meaney, M.J., 2014.
Influences of maternal and paternal PTSD on epigenetic regulation of the glucocorticoid receptor gene
in holocaust survivor offspring. *American Journal of Psychiatry* 171 (8), 872–880.

Zimmer, C., Taff, C. C., Ardia, D. R., Ryan, T. A., Winkler, D. W., & Vitousek, M. N. (2019). On again, off again:
acute stress response and negative feedback together predict resilience to experimental
challenges. *Functional Ecology*, *33*(4), 619-628.

Zimmer, C., Taff, C. C., Ardia, D. R., Rose, A. P., Aborn, D. A., Johnson, L. S., & Vitousek, M. N. (2020).
Environmental unpredictability shapes glucocorticoid regulation across populations of tree
swallows. *Scientific reports*, *10*(1), 1-13.

Zimmer, C., Woods, H. A., & Martin, L. B. (2022). Information theory in vertebrate stress physiology. *Trends*
in Endocrinology & Metabolism, *33*(1), 8-17.

SUPPLEMENTARY INFORMATION

569
Do glucocorticoids indicate stress? Meta-analysis reveals a tight association
570
between glucocorticoids and metabolic rate

571 **Blanca Jimeno & Simon Verhulst**

573 **Table S1.** Study selection steps and number of studies found

"energy expenditure" AND (glucocorticoid OR cortisol OR corticosterone) (20/07/2021)			
Total	After abstract	After text	After effect size calculations
707	108	43	9
"metabolic rate" AND (glucocorticoid OR cortisol OR corticosterone) (20/07/2021)			
Total	After abstract	After text	After effect size calculations
447	71 (64*)	28	5
Selected papers' reference check			
Total (excl. previously included)	After abstract	After text	After effect size calculations
20	15	9	7

Text step requires paper selection criteria to be met. Effect size calculation step requires enough data reported to calculate effect sizes by experimental treatment.

Main reasons for not including articles after abstract / text step: i) Experimental treatment had no effect on metabolic rate, ii) chemical or drug induction of metabolic rate increase, iii) GCs were measured in other tissues than plasma, iv) no experimental manipulation of metabolic rate.

*Number of studies excluding those overlapping with previous search, which were not further considered.

Table S2: Table with the information extracted from each study included in the meta analysis (excel sheet available).

Table S3: Effect size calculations. The document Includes one sheet per study with the data extracted, the part of the
article it was extracted from, and the effect size calculations and results further included in the meta analysis and Table
S2 (excel sheets available).

Table S4. Meta regression model (quantitative approach) testing the effect of (a) Taxa, (b) Before / after effect, (c)
Experiment / control effect, (d) Use of Metabolic Rate or Heart Rate as metabolic variable and (e) Treatment type, on
the association between metabolic rate (MR) and glucocorticoid effect sizes across studies.

(a)	Estimate	s.e.	Z	P	95% C.I.
Intercept	0.70	0.12	6.00	<0.0001	0.47-0.93
MR effect size (ln)	**0.29**	**0.11**	**2.70**	**0.007**	**0.08-0.50**
Taxa (mammal)	0.12	0.24	0.49	0.63	-0.35-0.58
MR : Taxa	0.17	0.22	0.80	0.42	-0.25-0.60

Variance components: Study.ID (Sigma^2)– Estimate = 0.00, sqrt = 0.00, n= 21

Residual heterogeneity: QE(df = 31) = 27.44, p = 0.65

Test of moderators: QM(df = 3) = 9.60, p=0.022

(b)	Estimate	s.e.	Z	P	95% C.I.
Intercept	0.73	0.11	6.42	<0.0001	0.51-0.95
MR effect size (ln)	**0.31**	**0.11**	**2.95**	**0.003**	**0.10-0.52**
Before / after effect (yes)	-0.26	0.38	-0.70	0.49	-0.99-0.48
MR : Time effect	0.02	0.31	0.05	0.96	-0.59-0.62

Variance components: Study.ID (Sigma^2)– Estimate = 0.00, sqrt = 0.00, n= 21

Residual heterogeneity: QE(df = 31) = 27.91, p = 0.62

Test of moderators: QM(df = 3) = 9.13, p=0.028

(c)	Estimate	s.e.	Z	P	95% C.I.
Intercept	0.73	0.11	6.51	<0.0001	0.51-0.96
MR effect size (ln)	**0.31**	**0.10**	**2.97**	**0.003**	**0.11-0.51**
Exp./ control effect (yes)	0.05	0.31	0.16	0.87	-0.56-0.65
MR : Exp. / control effect	-0.06	0.30	-0.22	0.83	-0.64-0.51

Variance components: Study.ID (Sigma^2)– Estimate = 0.00, sqrt = 0.00, n= 21

Residual heterogeneity: QE(df = 31) = 28.18, p = 0.61

Test of moderators: QM(df = 3) = 8.94, p= 0.03

(d)	Estimate	s.e.	Z	P	95% C.I.
Intercept	0.73	0.12	6.11	<0.0001	0.50-0.97
MR effect size (ln)	**0.33**	**0.11**	**2.97**	**0.003**	**0.11-0.54**
Met. variable (HR)	0.14	0.24	0.57	0.57	-0.33-0.60
MR : Met. variable	-0.02	0.22	-0.07	0.94	-0.45-0.42

Variance components: Study.ID (Sigma^2)– Estimate = 0.01, sqrt = 0.09, n= 21

Residual heterogeneity: QE(df = 31) = 28.09, p = 0.62

Test of moderators: QM(df = 3) = 8.67, p= 0.022

e)	Estimate	s.e.	Z	P	95% C.I.
Intercept	0.78	0.21	3.79	0.0002	0.38-1.19
MR effect size (ln)	**0.27**	**0.19**	**1.39**	**0.166**	**-0.11-0.64**
Treat. Type 2	-0.03	0.27	-0.09	0.93	-0.56-0.51
Treat. Type 3	-0.16	0.31	-0.52	0.60	-0.78-0.45
MR : Treat. Type 2	0.05	0.25	0.22	0.83	-0.44-0.55
MR : Treat. Type 3	0.08	0.29	0.27	0.79	-0.48-0.64

Variance components: Study.ID (Sigma^2)– Estimate = 0.02, sqrt = 0.14, n= 21

Residual heterogeneity: QE(df = 29) = 27.99, p = 0.52

Test of moderators: QM(df = 5) = 8.86, p= 0.115

Fig S1: Forest plot showing the metabolic rate (MR) effect sizes (Cohen's D ± 95% CI) associated to experimental manipulations of metabolic rate, grouped by treatment group and study. Area of squares is proportional to the experiment sample size (1/s.e.).

587
588
589

Fig S2. Relationship between metabolic rate and glucocorticoid effect sizes (Cohen's D) across studies. Panels show the association without ln-transforming metabolic rate effect sizes (left panel) and when ln-transforming them (right panel). Size of dots is proportional to the experiment sample size (i.e. square root of the number of individuals in which glucocorticoids were measured). Shaded areas represent 95% C.I. Note that the number of data points in the graph is higher than the number of studies, as some of the studies included multiple experimental treatments (Study ID was included as random factor in statistical analyses).

Fig S3. Relationship between metabolic rate and glucocorticoid effect sizes (Cohen's D) across studies as a function of a) Before / after effect (open circles and dashed line for studies including a time effect; closed circles and continuous line for studies not including a time effect; see methods) and b) Experiment / control effect (open circles and dashed line for studies including within-individual variation; closed circles and continuous line for studies not including within-individual variation). Size of dots is proportional to the experiment sample size (i.e. square root of the number of individuals in which glucocorticoids were measured). Shaded areas represent 95% C.I. Note that the number of data points in the graph is higher than the number of studies, as some of the studies included multiple experimental treatments (Study ID was included as random factor in statistical analyses). Before / after effect and Experiment / control effect results should be interpreted with caution, as most studies do include within individual variation and do not include time effect.